Table of Contents

Cover 1
About this book 2
Copyright 3
Front of Book 5
- Abbreviations 5
- Acknowledgments 16
- Dedication 17
- Foreword 18
- Normal values: worth knowing by heart 19
- Preface 21
- Reviewers 22
- The author 25
- What is new? 26

The basis of general practice 32
- Chapter 1 - The Nature and Content of General Practice 32
- Chapter 2 - The Family 39
- Chapter 3 - Consulting Skills 48
- Chapter 4 - Communication Skills 59
- Chapter 5 - Counselling Skills 73
- Chapter 6 - Difficult, Demanding and Angry Patients 87
- Chapter 7 - Palliative Care 93
- Chapter 8 - The Elderly Patient 105
- Chapter 9 - Prevention in General Practice 125
- Chapter 10 - Nutrition in Health and Illness 140
- Chapter 11 - Health Promotion and Patient Education 151
- Chapter 12 - Pain and Its Management 160
- Chapter 13 - Whole Person Approach to Management 177
- Chapter 14 - Travel and Tropical Medicine 184
- Chapter 15 - Research and Evidence-Based Medicine 209
- Chapter 16 - Laboratory Investigations 220

Diagnostic perspective in general practice 235
- Chapter 17 - Inspection as A Clinical Skill 235
- Chapter 18 - A Safe Diagnostic Strategy 242
- Chapter 19 - Genetic Conditions 254
- Chapter 20 - Depression 275
- Chapter 21 - Diabetes Mellitus: Diagnosis 286
- Chapter 22 - Drug Problems 295
- Chapter 23 - Anaemia 309
- Chapter 24 - Thyroid and Other Endocrine Disorders 319
- Chapter 25 - Spinal Dysfunction 333
- Chapter 26 - Urinary Tract Infection 336
- Chapter 27 - Malignant Disease 346
- Chapter 28 - HIV/AIDS - Could it be HIV? 357
- Chapter 29 - Baffling Viral and Protozoal Infections 370
- Chapter 30 - Baffling Bacterial Infections 382
- Chapter 31 - Chronic Kidney Disease 398
- Chapter 32 - Connective Tissue Disease and The Vasculitides 407
- Chapter 33 - Neurological Dilemmas 421

Problem solving in general practice 441
Chapter 34 - Abdominal Pain 441
Chapter 35 - Arthritis 478
Chapter 36 - Anorectal Disorders 512
Chapter 37 - Low Back Pain 524
Chapter 38 - Thoracic Back Pain 558
Chapter 39 - Bruising and Bleeding 580
Chapter 40 - Chest Pain 593
Chapter 41 - Constipation 626
Chapter 42 - Cough 641
Chapter 43 - Deafness and Hearing Loss 664
Chapter 44 - Diarrhoea 679
Chapter 45 - The Disturbed Patient 702
Chapter 46 - Dizziness 725
Chapter 47 - Dyspepsia (Indigestion) 738
Chapter 48 - Dysphagia 753
Chapter 49 - Dyspnoea 758
Chapter 50 - The Painful Ear 778
Chapter 51 - The Red and Tender Eye 800
Chapter 52 - Pain in the Face 822
Chapter 53 - Fever and Chills 840
Chapter 54 - Faints, Fits and Funny Turns 854
Chapter 55 - Haematemesis and Melaena 866
Chapter 56 - Headache 870
Chapter 57 - Hoarseness 900
Chapter 58 - Jaundice 904
Chapter 59 - Nasal Disorders 927
Chapter 60 - Nausea and vomiting 940
Chapter 61 - Neck Lumps 948
Chapter 62 - Neck Pain 954
Chapter 63 - Shoulder Pain 975
Chapter 64 - Pain in the Arm and Hand 994
Chapter 65 - Hip, Buttock and Groin Pain 1021
Chapter 66 - Pain in the Leg 1039
Chapter 67 - The Painful Knee 1066
Chapter 68 - Pain in the Foot and Ankle 1098
Chapter 69 - Walking Difficulty 1127
Chapter 70 - Palpitations 1137
Chapter 71 - Sleep Disorders 1154
Chapter 72 - Sore Mouth and Tongue 1167
Chapter 73 - Sore Throat 1184
Chapter 74 - Tiredness 1198
Chapter 75 - The Unconscious Patient 1209
Chapter 76 - Urinary Disorders 1220
Chapter 77 - Visual Failure 1237
Chapter 78 - Weight Gain 1256
Chapter 79 - Weight Loss 1270
Child and adolescent health 1279
Chapter 80 - An Approach to The Child 1279
Chapter 81 - Specific Problems of Children 1290
Chapter 82 - Surgical Problems in Children 1316

Chapter 83 - Common Childhood Infectious Diseases (Including Skin Eruptions) 1331
Chapter 84 - Behaviour Disorders in Children 1354
Chapter 85 - Child Abuse 1366
Chapter 86 - Emergencies in Children 1377
Chapter 87 - Adolescent Health 1401
Women's health 1409
Chapter 100 - Infections in Pregnancy 1409
Chapter 101 - High-Risk Pregnancy 1415
Chapter 102 - Postnatal Care 1436
Chapter 88 - Cervical Cancer and Pap Smears 1447
Chapter 89 - Family Planning 1457
Chapter 90 - Breast Pain (Mastalgia) 1468
Chapter 91 - Lumps in the Breast 1477
Chapter 92 - Abnormal Uterine Bleeding 1495
Chapter 93 - Lower Abdominal and Pelvic Pain in Women 1506
Chapter 94 - Premenstrual Syndrome 1524
Chapter 95 - The Menopause and Osteoporosis 1530
Chapter 96 - Vaginal Discharge 1544
Chapter 97 - Vulvular Disorders 1557
Chapter 98 - Violence Against Women 1569
Chapter 99 - Basic Antenatal Care 1576
Men's health 1587
Chapter 103 - Men's Health: An Overview 1587
Chapter 104 - Scrotal Pain 1591
Chapter 105 - Inguinoscrotal Lumps 1598
Chapter 106 - Disorders of The Penis 1613
Chapter 107 - Disorders of the Prostate 1623
Sex-related problems 1632
Chapter 108 - The Subfertile Couple 1632
Chapter 109 - Sexual Health 1642
Chapter 110 - Sexually Transmitted Infections 1659
Problems of the skin 1676
Chapter 111 - A Diagnostic and Management Approach to Skin Problems 1676
Chapter 112 -Pruritus 1692
Chapter 113 - Common Skin Problems 1708
Chapter 114 - Acute Skin Eruptions 1745
Chapter 115 - Skin Ulcers 1766
Chapter 116 - Common Lumps and Bumps 1778
Chapter 117 - Pigmented Skin Lesions 1807
Chapter 118 - Hair Disorders 1822
Chapter 119 - Nail Disorders 1835
Chronic disorders: continuing management 1853
Chapter 120 - Alcohol Problems 1853
Chapter 121 - Allergic Disorders Including Hay Fever 1864
Chapter 122 - Anxiety 1874
Chapter 123 - Asthma 1888
Chapter 124 - Chronic Obstructive Pulmonary Disease 1905
Chapter 125 - Epilepsy 1916
Chapter 126 - Hypertension 1927
Chapter 127 - Dyslipidaemia 1953
Chapter 128 - Diabetes Mellitus: Management 1959

Chapter 129 - Chronic Heart Failure 1975

Accident and emergency medicine 1984

Chapter 130 - Emergency Care 1984

Chapter 131 - The Doctor's Bag and Other Emergency Equipment 2006

Chapter 132 - Stroke and Transient Ischaemic Attacks 2014

Chapter 133 - Thrombosis and Thromboembolism 2022

Chapter 134 - Common Skin Wounds and Foreign Bodies 2031

Chapter 135 - Common Fractures and Dislocations 2052

Chapter 136 - Common Sporting Injuries 2089

Indigenous health 2115

Chapter 137 - The Health of Indigenous Peoples 2115

Appendices 2127

Appendix I. Percentile charts: infant girls 2127

Appendix II. Percentile charts: infant boys 2128

Appendix III. Percentile charts: girls 2129

Appendix IV. Percentile charts: boys 2130

Appendix V. Respiratory function tables 2131

Appendix VI. The Australian Nutrition Foundation: weight for height charts 2133


John MURTAGH'S
general practice
FOURTH EDITION

John Murtagh's General Practice, 4th Edition

Compiled by KhCh

Courtesy of Z

ABOUT THIS BOOK



John Murtagh's general practice, Fourth Edition
John Murtagh
© 2007 McGraw-Hill Australia Pty Ltd
ISBN13: 978-0-074717-79-0
ISBN10: 007-4-717-79-0

Murtagh's ***General Practice*** is widely recognised as the most influential publication for general practice and primary health care. Drawn from John Murtagh's extensive experience as a rural GP, it attempts to address the issue of the broad knowledge and skills required in modern general practice.

Murtagh's ***General Practice*** adopts an approach typical of Western medicine: strategies for solving common problems, continuing care, basic medical skills and 'tricks of the trade'. The approach is to promote relatively fail-safe strategies to reduce the margin of error. Guidelines for early referral are included and, importantly, information and advice aimed at developing counselling and therapeutic skills in the general practitioner are also included.

Murtagh's ***General Practice*** has selected those problems that are common, significant, preventable and treatable.

This edition includes six new chapters for access to current guidelines on nasal disorders, surgical problems in children, infections in pregnancy, chronic obstructive pulmonary disease, chronic heart failure and thrombosis and thromboembolism.

"...in an extraordinary way the book became known as the 'Bible of General Practice' in Australia. In addition to being widely used by practising doctors, it has become a popular and standard textbook in several medical schools and also in the teaching institutions for alternative health practitioners such as chiropractic, naturopathy and osteopathy. "

GC Schofield,
Dean of Medicine, Monash University, Melbourne 1977-88

COPYRIGHT

NOTICE

Medicine is an ever-changing science. As new research and clinical experience broaden our knowledge, changes in treatment and drug therapy are required. The editors and the publisher of this work have checked with sources believed to be reliable in their efforts to provide information that is complete and generally in accord with the standards accepted at the time of publication. However, in view of the possibility of human error or changes in medical sciences, neither the editors, nor the publisher, nor any other party who has been involved in the preparation or publication of this work warrants that the information contained herein is in every respect accurate or complete. Readers are encouraged to confirm the information contained herein with other sources. For example, and in particular, readers are advised to check the product information sheet included in the package of each drug they plan to administer to be certain that the information contained in this book is accurate and that changes have not been made in the recommended dose or in the contraindications for administration. This recommendation is of particular importance in connection with new or infrequently used drugs.

This fourth edition published 2007. Reprinted 2008, Reprinted 2009
First edition published 1992
Second edition published 1998
Third edition published 2003
Text copyright © 2007 John E. Murtagh
Illustrations copyright © 2007 John E. Murtagh and McGraw-Hill Australia Pty Ltd
Design copyright © 2007 McGraw-Hill Australia Pty Ltd
Additional owners of copyright are named in on-page credits and on the acknowledgments page. Every effort has been made to trace and acknowledge copyright material.
The authors and publishers tender their apologies should any infringement have occurred.

Reproduction and communication for educational purposes
The *Australian Copyright Act 1968* (the Act) allows a maximum of one chapter or 10% of the pages of this work, whichever is the greater, to be reproduced and/or communicated by any educational institution for its educational purposes provided that the institution (or the body that administers it) has sent a Statutory Educational notice to Copyright Agency Limited (CAL) and been granted a licence. For details of statutory educational and other copyright licences contact: Copyright Agency Limited, Level 15, 233 Castlereagh Street, Sydney NSW 2000. Telephone: (02) 9394 7600. Website: www.copyright.com.au

Reproduction and communication for other purposes
Apart from any fair dealing for the purposes of study, research, criticism or review, as permitted under the Act, no part of this publication may be reproduced, distributed or transmitted in any form or by any means, or stored in a database or retrieval system, without the written permission of McGraw-Hill Australia including, but not limited to, any network or other electronic storage.

Enquiries should be made to the publisher via www.mcgraw-hill.com.au or marked for the attention of the Rights and Permissions Manager at the address below.

National Library of Australia Cataloguing-in-Publication data

Murtagh, John
John Murtagh's general practice.
4th ed.
Bibliography.
Includes index.
ISBN 9780074717790.
ISBN 9780070135963 (RACGP edition)
1.Family medicine 2.Physicians (General Practice)
I.Title
610

Published in Australia by
McGraw-Hill Australia Pty Ltd
Level 2, 82 Waterloo Road, North Ryde NSW 2113

Publisher: Nicole Meehan
Project Editor: Jane Riley

Editorial Assistant: Hollie Zondanos
Editor: Joy Window
Proofreader: Pauline O'Carolan
Indexer: Max McMaster
Art Director: Steve Randles
Cover design: Jennifer Pace Walter
Internal design: Jan Schmoeger/Designpoint
Illustrations: Alan Laver/Shelley Communications, Chris Sorell, John Murtagh, and Trade Wind Creations
Cover image: Masterfile
Typeset in 10/11 pt Minton Pro by Aptara Inc., New Delhi, India
Printed on 70 gsm matt art by 1010 Printing International Ltd, China

ABBREVIATIONS

AAA	abdominal aortic aneurysm
AAFP	American Academy of Family Physicians
ABC	airway, breathing, circulation
ABCD	airway, breathing, circulation, dextrose
ABFP	American Board of Family Practice
ABI	ankle brachial index
AC	air conduction
AC	acromioclavicular
ACAH	autoimmune chronic active hepatitis
ACE	angiotensin-converting enzyme
ACL	anterior cruciate ligament
ACR	albumin creatine ratio
ACTH	adrenocorticotrophic hormone
AD	aortic dissection
AD	autosomal dominant
ADHD	attention deficit hyperactivity disorder
ADT	adult diphtheria vaccine
AFI	amniotic fluid index
AFP	alpha-fetoprotein
AI	aortic incompetence
AICD	automatic implantable cardiac defibrillator
AIDS	acquired immunodeficiency syndrome
AIIRA	angiotension II(2) reuptake antagonist
AKF	acute kidney failure
ALE	average life expectancy
ALL	acute lymphocytic leukaemia
ALP	alkaline phosphatase
ALT	alanine aminotransferase
ALTE	apparent life-threatening episode
AMI	acute myocardial infarction
AML	acute myeloid leukaemia
ANA	antinuclear antibody
ANCI	antineutrophil cytoplasmic antibody
ANF	antinuclear factor
a/n/v	anorexia/nausea/vomiting
AP	anterior - posterior
APF	Australian pharmaceutical formulary
APH	ante-partum haemorrhage
APTT	activated partial thromboplastin time
AR	autosomal recessive
ARC	AIDS-related complex
ARR	absolute risk reduction
ASD	atrial septal defect
ASIS	anterior superior iliac spine
ASOT	antistreptolysin 0 titre
AST	aspartate aminotransferase
ATFL	anterior talofibular ligament
AV	atrioventricular
AVM	arteriovenous malformation
AZT	azidothymidine

BC	bone conduction
BCC	basal cell carcinoma
BCG	bacille Calmette-Guérin
bDMARDs	biological disease modifying antirheumatic drugs
BMI	body mass index
BOO	bladder outlet obstruction
BP	blood pressure
BPH	benign prostatic hyperplasia
BPPV	benign paroxysmal positional vertigo
BSE	breast self-examination

Ca	carcinoma
CABG	coronary artery bypass grafting
CAD	coronary artery disease
CAP	community acquired pneumonia
CBE	clinical breast examination
CBT	cognitive behaviour therapy
CCF	congestive cardiac failure
CCP	cyclic citrinullated peptide
CCT	controlled clinical trial
CCU	coronary care unit
CD $_4$	T helper cell
CD $_8$	T suppressor cell
CDT	combined diphtheria/tetanus vaccine
CEA	carcinoembryonic antigen
CFL	calcaneofibular ligament
CFS	chronic fatigue syndrome
cfu	colony forming unit
CHD	coronary heart disease
CHF	chronic heart failure
CI	confidence interval
CIN	cervical intraepithelial neoplasia
CJD	Creutzfeldt-Jakob disease
CK	creatinine kinase
CK - MB	creatinine kinase - myocardial bound fraction
CKD	chronic kidney disease
CKF	chronic kidney failure
CMC	carpometacarpal
CML	chronic myeloid leukaemia
CMV	cytomegalovirus
CNS	central nervous system
co	compound
COAD	chronic obstructive airways disease
COC	combined oral contraceptive
COMT	catechol-O-methyl transferase
COPD	chronic obstructive pulmonary disease
COX	cyclooxygenase
CPA	cardiopulmonary arrest
CPAP	continuous positive airways pressure
CPK	creatine phosphokinase
CPPD	calcium pyrophosphate dihydrate

CPR	cardiopulmonary resuscitation
CPS	complex partial seizures
CR	controlled release
CRD	computerised reference database system
CREST	calcinosis cutis; Raynaud ' s phenomenon; oesophageal involvement; sclerodactyly; telangiectasia
CRF	chronic renal failure
CRFM	chloroquine-resistant falciparum malaria
CRH	corticotrophin-releasing hormone
CR(K)F	chronic renal (kidney) failure
CRP	C-reactive protein
CSF	cerebrospinal fluid
CSFM	chloroquine-sensitive falciparum malaria
CSIs	COX-2 specific inhibitors
CSU	catheter specimen of urine
CT	computerised tomography
CTD	connective tissue disorder
CTG	cardiotocograph
CTS	carpal tunnel syndrome
CVA	cerebrovascular accident
CVS	cardiovascular system
CXR	chest X-ray
DBP	diastolic blood pressure
DC	direct current
DDAVP	desmopressin acetate
DDH	developmental dysplasia of the hip
DEXA	dual energy X-ray absorptiometry
DHA	docosahexaenoic acid
DI	diabetes insipidus
DIC	disseminated intravascular coagulation
DIDA	di-imino diacetic acid
DIMS	disorders of initiating and maintaining sleep
DIP	distal interphalangeal
dL	decilitre
DMARDs	disease modifying antirheumatic drugs
DNA	deoxyribose-nucleic acid
DOM	direction of movement
DRE	digital rectal examination
DRABC	defibrillation, resuscitation, airway, breathing, circulation
drug	bd - twice daily
dosage	tid, tds - three times dailyqid, qds - four times daily
ds	double strand
DS	double strength
DSM	diagnostic and statistical manual (of mental disorders)
DU	duodenal ulcer
DUB	dysfunctional uterine bleeding
DVT	deep venous thrombosis
DxT	diagnostic triad
EAR	expired air resuscitation
EBM	Epstein-Barr mononucleosis (glandular fever)
EBNA	Epstein-Barr nuclear antigen

EBV	Epstein-Barr virus
ECC	external chest compression
ECG	electrocardiogram
ECT	electroconvulsive therapy
ED	emergency department
EDD	expected due date
EEG	electroencephalogram
ELISA	enzyme linked immunosorbent assay
EMG	electromyogram
ENA	extractable nuclear antigen
EO	ethinyloestradiol
EPA	eicosapentaenoic acid
EPL	extensor pollicis longus
EPS	expressed prostatic secretions
ER	external rotation
ESRF	end-stage renal failure
ESR(K)F	end stage renal (kidney) failure
ERCP	endoscopic retrograde cholangiopancreatography
esp.	especially
ESR	erythrocyte sedimentation rate
ET	embryo transfer
ETT	endotracheal tube

FAD	familial Alzheimer disease
FAP	familial adenomatous polyposis
FB	foreign body
FBE	full blood count
FDL	flexor digitorum longus
FEV_1	forced expiratory volume in 1 second
FHL	flexor hallucis longus
fL	femto-litre (10^{-15})
FRC	functional residual capacity
FSH	follicle stimulating hormone
FTA- ABS	fluorescent treponemal antibody absorption test
FTT	failure to thrive
FUO	fever of undetermined origin
FVC	forced vital capacity
FXS	fragile X syndrome

g	gram
GA	general anaesthetic
GABHS	group A beta-haemolytic streptococcus
GBS	Guillain-Barr é syndrome
GCA	giant cell arteritis
GESA	Gastroenterological Society of Australia
GFR	glomerular filtration rate
GGT	gamma-glutamyl transferase
GI	glycaemic index
GIFT	gamete intrafallopian transfer
GIT	gastrointestinal tract
GnRH	gonadotrophin-releasing hormone
GO	gastro-oesophageal

GORD	gastro-oesophageal reflux
GP	general practitioner
G-6-PD	glucose-6-phosphate
GSI	genuine stress incontinence
GU	gastric ulcer
GV	growth velocity

HAV	hepatitis A virus
anti-HAV	hepatitis A antibody
Hb	haemoglobin
HbA	haemoglobin A
anti-HBc	hepatitis B core antibody
HBeAg	hepatitis Be antigen
anti-HBs	hepatits B surface antibody
HBsAg	hepatitis B surface antigen
HBV	hepatitis B virus
HCG	human chorionic gonadotropin
HCV	hepatitis C virus
anti-HCV	hepatitis C virus antibody
HDL	high-density lipoprotein
HDV	hepatitis D (Delta) virus
HEV	hepatitis E virus
HFA	hydrofluoro alkane
HFM	hand, foot and mouth
HFV	hepatitis F virus
HGV	hepatitis G virus
HHC	hereditary haemochromatosis
HIDA	hydroxy iminodiacetic acid
HIV	human immunodeficiency virus
HLA-B $_{27}$	human leucocyte antigen
HMGCoA	hydroxymethylglutaryl CoA
HNPCC	hereditary nonpolyposis colorectal cancer
HPV	human papilloma virus
HRT	hormone replacement therapy
HSIL	high grade squamous intraepithelial lesion
HSV	herpes simplex viral infection
H	hypertension

IBS	irritable bowel syndrome
ICE	ice, compression, elevation
ICHPPC	International Classification of Health Problems in Primary Care
ICS	inhaled corticosteroid
ICS	intercondylar separation
ICSI	intracytoplasmic sperm injection
IDDM	insulin dependent diabetes mellitus
IDU	injecting drug user
IgE	immunoglobulin E
IgG	immunoglobulin G
IgM	immunoglobulin M
IHD	ischaemic heart disease
IHS	International Headache Society
IM, IMI	intramuscular injection

IMS	intermalleolar separation
inc.	including
INR	international normalised ratio
IOC	International Olympic Committee
IOFB	intraocular foreign body
IP	interphalangeal
IPPV	intermittent positive pressure variation
IR	internal rotation
ITP	idiopathic (or immune) thrombocytopenia purpura
IUCD	intrauterine contraceptive device
IUGR	intrauterine growth retardation
IV	intravenous
IVF	in-vitro fertilisation
IVI	intravenous injection
IVP	intravenous pyelogram
IVU	intravenous urogram
JCA	juvenile chronic arthritis
JVP	jugular venous pulse
KA	keratoacanthoma
kg	kilogram
KOH	potassium hydroxide
LA	local anaesthetic
LABA	long acting beta agonist
LBBB	left branch bundle block
LBO	large bowel obstruction
LBP	low back pain
LCR	ligase chain reaction
LDH/LH	lactic dehydrogenase
LDL	low-density lipoprotein
LFTs	liver function tests
LH	luteinising hormone
LHRH	luteinising hormone releasing hormone
LIF	left iliac fossa
LMN	lower motor neurone
LNG	levonorgestrel
LPC	liquor picis carbonis
LRTI	lower respiratory tract infection
LSD	lysergic acid
LSIL	low grade squamous intraepithelial lesion
LUQ	left upper quadrant
LUTS	lower urinary tract symptoms
LV	left ventricular
LVH	left ventricular hypertrophy
MAIS	*Mycobacterium avium intracellulare* or *M. sacrofulaceum*
mane	in morning
MAOI	monoamine oxidase inhibitor
MAST	medical anti-shock trousers
MB	myocardial base

MCL	medial collateral ligament
MCP	metacarpal phalangeal
MCU	microscopy and culture of urine
MCV	mean corpuscular volume
MDI	metered dose inhaler
MG	myaesthenia gravis
MI	myocardial infarction
MIC	mitral incompetence
MID	minor intervertebral derangement
MND	motor neurone disease
MRCP	magnetic resonance cholangiography
MRI	magnetic resonance imaging
MRSA	methicillin-resistant *staphylococcus aureus*
MS	multiple sclerosis
MSU	midstream urine
MTP	metatarsophalangeal
MVA	motor vehicle accident
N	normal
N saline	normal saline
NAD	no abnormality detected
NET	norethisterone
NF	neurofibromatosis
NGU	non-gonococcal urethritis
NHL	non-Hodgkin ' s lymphoma
NH & MRC	National Health and Medical Research Council
NIDDM	non-insulin dependent diabetes mellitus
NNT	numbers needed to treat
nocte	at night
NR	normal range
NRT	nicotine replacement therapy
NSAIDs	non-steroidal anti-inflammatory drugs
NSCLC	non-small cell lung cancer
NSU	non-specific urethritis
(o)	taken orally
OA	osteoarthritis
OCP	oral contraceptive pill
OGTT	oral glucose tolerance test
OSA	obstructive sleep apnoea
OSD	Osgood-Schlatter disorder
OTC	over the counter
PA	posterior - anterior
PAN	polyarteritis nodosa
Pap	Papanicolaou
PBG	porphobilinogen
PBS	Pharmaceutical Benefits Scheme
pc	after meals
PCA	percutaneous continuous analgesia
PCB	post coital bleeding
PCL	posterior cruciate ligament

PCOS	polycystic ovarian syndrome
PCP	pneumocystitis pneumonia
PCR	polymerase chain reaction
PCV	packed cell volume
PD	Parkinson ' s disease
PDA	patent ductus arteriosus
PDD	pervasive development disorders
PEF	peak expiratory flow
PEFR	peak expiratory flow rate
PET	pre-eclamptic toxaemia
PET	positron emission tomography
PFO	patent foramen ovale
PFT	pulmonary function test
PGL	persistent generalised lymphadenopathy
PH	past history
PHR	personal health record
PID	pelvic inflammatory disease
PIP	proximal interphalangeal
PKU	phenylketonuria
PLISSIT	permission: limited information: specific suggestion: intensive therapy
PLMs	periodic limb movements
PMDD	premenstrual dysphoric disorder
PMS	premenstrual syndrome
POP	plaster of Paris
POP	progestogen-only pill
PPI	proton-pump inhibitor
PPROM	preterm premature rupture of membranes
PR	per rectum
prn	as and when needed
PRNG	penicillin-resistant gonococci
PROM	premature rupture of membranes
PSA	prostate specific antigen
PSGN	post streptococcal glomerulonephritis
PSIS	posterior superior iliac spine
PSVT	paroxysmal supraventricular tachycardia
PT	prothrombin time
PTC	percutaneous transhepatic cholangiography
PTFL	posterior talofibular ligament
PU	peptic ulcer
PUO	pyrexia of undetermined origin
PUVA	psoralen + UVA
pv	per vagina
PVC	polyvinyl chloride
PVD	peripheral vascular disease
qds, qid	four times daily
RA	rheumatoid arthritis
RACGP	Royal Australian College of General Practitioners
RAP	recurrent abdominal pain
RBC	red blood cell
RCT	randomised controlled trial

RF	rheumatic fever
Rh	rhesus
RIB	rest in bed
RICE	rest, ice, compression, elevation
RIF	right iliac fossa
RPR	rapid plasma reagin
RR	relative risk
RRR	relative risk reduction
RSD	reflex sympathetic dystrophy
RSI	repetition strain injury
RSV	respiratory syncytial virus
RT	reverse transcriptase
rtPA	recombinant tissue plasminogen activator
RUQ	right upper quadrant
s	serum
SABA	short acting beta agonist
SAH	subarachnoid haemorrhage
SARS	severe acute respiratory distress syndrome
SBE	subacute bacterial endocarditis
SBO	small bowel obstruction
SBP	systolic blood pressure
SC/SCI	subcutaneous/subcutaneous injection
SCC	squamous cell carcinoma
SCFE	slipped capital femoral epiphysis
SCG	sodium cromoglycate
SCLC	small cell lung cancer
SIADH	syndrome of secretion of inappropriate antidiuretic hormone
SIDS	sudden infant death syndrome
SIJ	sacroiliac joint
SL	sublingual
SLD	specific learning disability
SLE	systemic lupus erthematosus
SLR	straight leg raising
SND	sensorineural deafness
SNHL	sensorineural hearing loss
SNPs	single nuceotide polymorphisms
SNRI	serotonin noradrenaline reuptake inhibitor
SOB	shortness of breath
sp	species
SPA	suprapubic aspirate of urine
SPECT	single photon emission computerised tomography
SPF	sun penetration factor
SR	sustained release
SSRI	selective serotonin reuptake inhibitor
SSS	sick sinus syndrome
statim	at once
STI	sexually transmitted infection
STD	sodium tetradecyl sulfate
SUFE	slipped upper femoral epiphysis
SVC	superior vena cava
SVT	supraventricular tachycardia

T_3	tri-iodothyronine
T_4	thyroxine
TA	temporal arteritis
TB	tuberculosis
tds, tid	three times daily
TENS	transcutaneous electrical nerve stimulation
TFTs	thyroid function tests
TG	triglyceride
TIA	transient ischaemic attack
TIBC	total iron binding capacity
TM	tympanic membrane
TMJ	temporomandibular joint
TOE	transoesophageal echocardiography
TOF	tracheo-oesophageal fistula
TORCH	toxoplasmosis, rubella, cytomegalovirus, herpes virus
TPHA	Treponema pallidum haemoglutination test
TSE	testicular self-examination
TSH	thyroid-stimulating hormone
TUIP	transurethral incision of prostate
TURP	transurethral resection of prostate
TV	tidal volume
U	units
UC	ulcerative colitis
U & E	urea and electrolytes
μ g	microgram
UMN	upper motor neurone
URTI	upper respiratory tract infection
US	ultrasound
UTI	urinary tract infection
U	ultraviolet
VAS	visual analogue scale
VBI	vertebrobasilar insuffiency
VC	vital capacity
VDRL	Venereal Disease Reference Laboratory
VF	ventricular fibrillation
VMA	vanillylmandelic acid
VPG	venous plasma glucose
VRE	vancomycin-resistant enterococci
VSD	ventricular septal defect
VT	ventricular tachycardia
VUR	vesicoureteric reflux
VVS	vulvar vestibular syndrome
VWD	von Willebrand ' s disease
WBC	white blood cells
WBR	white → blue → red
WCC	white cell count
WHO	World Health Organization
WPW	Wolff-Parkinson-White

XL	sex linked

ACKNOWLEDGMENTS

The author would like to thank the Publication Division of the Royal Australian College of General Practitioners for supporting my past role as Medical Editor of *Australian Family Physician* , which has provided an excellent opportunity to gather material for this book. Acknowledgment is also due to those medical organisations that have given permission to use selected information from their publications. They include the Preventive and Community Medicine committee of the RACGP (Guidelines for Preventive Activities in General Practice), Therapeutic Guidelines Limited (*Therapeutic Guidelines* series), the Hypertension Guideline Committee: Research Unit RACGP (South Australian Faculty), and the *Medical Observer* , publishers of *A Manual for Primary Health Care* , for permitting reproduction of Appendices I - IV.

Special thanks to Chris Sorrell, graphic designer, for his art illustration, and to Nicki Cooper, Jenny Green and Caroline Menara for their skill and patience in typing the manuscript.

Figures 65.6, 113.9c, 113.9d were provided by Dr Levent Efe.

Many of the quotations at the beginning of chapters appear in either Robert Wilkins (ed), *The Doctor ' s Quotation Book* , Robert Hale Ltd, London, 1991 or Maurice B. Strauss (ed), *Familiar Medical Quotations* , Little, Brown & Co., New York, 1958.

Thanks are also due to Dr Bruce Mugford, Dr Lucie Stanford and to Lesley Rowe, Deputy Director of Pharmacy, Nepean Hospital, Penrith, NSW, for reviewing the manuscript, and to the publishing and production team at McGraw-Hill Australia for their patience and assistance in so many ways.

Finally, thanks to Dr Ndidi Victor Ikealumba for his expert review of *General Practice fourth edition* and his subsequent contribution to this reprint.

DEDICATION

To my wife, Jill, and our children, Paul, Julie, Caroline, Luke and Clare, for their understanding, patience and support

and

to all my medical colleagues, past and present, who have provided the vast reservoir of knowledge from which the content of this book was made possible.

FOREWORD

In 1960 a young schoolmaster, then teaching biology and chemistry in a secondary school in rural Victoria, decided to become a country doctor. He was admitted to the first intake of students into the Medical School of the newly established Monash University and at the end of the six-year undergraduate medical course and subsequent intern and resident appointments his resolve to practise community medicine remained firm.

During his years of undergraduate and early postgraduate study Dr Murtagh continued to gather and record data relating to the diagnostic and therapeutic procedures and clinical skills he would require in solo country practice. These records, subsequently greatly expanded, were to provide at least the foundation of this book. Happily, after graduation, he married Dr Jill Rosenblatt, a young graduate from Melbourne University, who shared his vocational interests. Subsequently they also shared the fulfilment of family life and the intellectual and emotional satisfaction of serving as doctors in a rural setting.

In the meantime the Royal Australian College of General Practitioners had established postgraduate training programs that had a significant influence on standards of professional practice. At the same time Monash University established a Department of Community Medicine at one of its suburban teaching hospitals, under the Chairmanship of Professor Neil Carson and staffed by practitioners in the local community.

While in practice Dr Murtagh gained a Fellowship of the College through examination. The College recognised his unique clinical, educational and communication skills and immediately commissioned him to prepare educational programs, especially the CHECK programs. His outstanding expertise as a primary care physician led to his appointment as a senior lecturer in the University Department of Community Medicine.

The success of the initial academic development in Community Medicine at Monash University, and its influence on the clinical skills of its graduates as they relate to primary care, led to a University decision to establish a further Department of Community Medicine at another suburban teaching hospital in Melbourne. It was considered by the University to be entirely appropriate that Dr Murtagh be invited to accept appointment as Professor and Head of that Department. Four years later Professor Murtagh was appointed Head of the extended Department and the first Professor of General Practice at Monash University.

John Murtagh has now become a national and international authority on the content and teaching of primary care medicine. As Medical Editor of *Australian Family Physician* from 1986 to 1995 he took that journal to the stage where it was the most widely read medical journal in Australia.

This textbook provides a distillate of the vast experience gained by a once-upon-a-time rural doctor whose career has embraced teaching from first to last, whose interest is ensuring that disease, whether minor or life-threatening, is recognised quickly, and whose concern is that strategies to match each contingency are well understood.

General Practice is the outcome of the vision of a schoolteacher of great talent who made a firm decision to become a country doctor; through this book his dream has become a reality for all who are privileged to practise medicine in a community setting.

The first edition of this book, published in 1994, achieved remarkable success on both the national and international scene. The second and third editions built on this initial success and in an extraordinary way the book became known as the ‘ Bible of General Practice ’ in Australia. In addition to being widely used by practising doctors, it has become a popular and standard textbook in several medical schools and also in the teaching institutions for alternative health practitioners, such as chiropractic, naturopathy and osteopathy. In particular, medical undergraduates and graduates struggling to learn English have found the book relatively comprehensible. This fourth edition has been updated and expanded, and has retained the successful format of the previous three editions but with a more attractive and user-friendly format including clinical photographs and illustrations in colour.

John Murtagh ' s works have been translated into Italian by McGraw-Hill Libri Italia s.r.l., Portuguese by McGraw-Hill Nova Iorque and Spanish by McGraw-Hill Interamericana Mexico, and into Chinese, Greek, Polish and Russian.

GC SCHOFIELD
OBE, MD, ChB(NZ), DPhil(Oxon), FRACP,
FRACMA, FAMA
Professor of Anatomy,
Monash University , 1961 - 77,
Dean of Medicine,
Monash University , 1977 - 88

NORMAL VALUES: WORTH KNOWING BY HEART

The following is a checklist that one can use as a template to memorise normal quantitative values for basic medical conditions and management.

Hypertension	
BP	> 140/90 mmHg
Alcohol excessive drinking	
Males	> 4 standard drinks/day
Females	> 2 standard drinks/day
Anaemia - haemoglobin	
Males	< 130 g/L
Females	< 115 g/L
Body mass index Wt (kg)/Ht (m^2)	
Normal	20 - 25
Overweight	> 25
Obesity	> 30
Jaundice	
Serum bilirubin	> 19 μ mol/L
Fever - temperature (morning) (a)	
Oral	> 37.2 ° C
Rectal	> 37.7 ° C
Diabetes mellitus - blood sugar	
Random (1 reading if symptomatic, 2 readings if asymptomatic)	> 11.1 mmol/L
Fasting	> 7.0 mmol/L
or	the 2 values from an oral GTT
Hypokalaemia	
Serum potassium	< 3.5 mmol/L
Hyperkalaemia	
Serum potassium	> 5.0 mmol/L

Vital signs (average)	**< 6 months**	**6 months - 3 years**	**3 - 12 years**	**Adult**
Pulse	120 - 140	110	80 - 100	60 - 100
Respiratory rate	45	30	20	14
BP (mmHg)	90/60	90/60	100/70	≤ 130/85

Children's weight Rule of thumb:	1 - 10 year old: Wt = (age + 4) × 2 kg

(a) There is considerable diurnal variation in temperature so that it is higher in the evening (0.5 - 1°C). I would recommend the definition given by Yung et al. in *Infectious Diseases: a Clinical Approach* : ‘ Fever can be defined as an early morning oral temperature > 37.2 ° C or a temperature > 37.8 ° C at other times of the day’ .

PREFACE

The discipline of general practice has become complex, diffuse and challenging. With the ever burgeoning phenomenon of specialisation in medicine, with its associated complex technology and hospital basis, the need for competent, caring generalists is more important than ever before.

This textbook attempts to address the issue of the broad knowledge and skills required in modern general practice. Some of the basics of primary health care remain the same. In fact, there is an everlasting quality about many of the medical problems that affect the human being, be it a splinter under a nail, a stye of the eyelid, a terminal illness or simply stress-related anxiety. Many of the treatments and approaches to caring management are universal and timeless. Unfortunately, the modern general practitioner has lost a considerable amount of counselling and therapeutic skills compared with his or her counterpart of 50 years ago and this book encourages GPs to develop these basic skills.

General practice remains the most cost-effective health service and there is evidence that the consumer and government agencies are focusing more on general practitioners as a solution to the high-cost, high-technology, depersonalised services in the health care system. They are searching for value, satisfaction, early diagnosis and holistic management without inappropriate invasive investigations and potent medication.

This textbook of general practice attempts to take these issues into account and adopts an approach typical of Western medicine: strategies for solving common problems, continuing care, basic medical skills and ‘ tricks of the trade ’ .

The discipline of general practice can at times be overwhelming in its breadth and depth, especially with patients presenting with undifferentiated problems with an overlap of organic and psychosocial aspects. There is the constant challenge to make an appropriate early diagnosis and detect the ever-lurking life-threatening illness. The approach in this book is to promote relatively fail-safe strategies to reduce the margin of error. This diagnostic model, which pervades all the chapters on problem solving, is based on my own experience, but readers can draw on their own experience to make the model work effectively for themselves.

Guidelines for early referral are included.

Another feature is the prominence given to ‘ masquerades ’ in general practice so that an increased awareness of these baffling problems will lead to improved diagnostic skills and more interesting practice. The fourth edition includes an important diagnostic initiative in the form of diagnostic triads, which are a brief *aide-m é moire* to assist in identifying a disorder from three key symptoms or signs.

A particular challenge in the preparation of this book was to identify as much appropriate evidence-based material as possible. This information, which still has its limitations, has been combined with considerable collective wisdom from experts, especially from the *Therapeutic Guidelines* series. For the everyday nitty-gritty problems of general practice, recommended treatments include those that have worked effectively for myself and my colleagues.

This fourth edition also addresses the concern from many colleagues and patients in particular about the emphasis of our profession on ‘ disease ’ . This term for a particular condition has been replaced where appropriate by the term syndrome or disorder.

In this edition, each chapter covering clinical problems includes a section on those problems affecting children and the elderly. If problems affecting those age groups are not found in ‘ child and adolescent health ’ they will be found in the clinical chapters such as arthritis, dyspnoea and hypertension.

Such a book cannot possibly present all the medical problems likely to be encountered, but an attempt has been made to select those problems that are common, significant, preventable and treatable. I am confident that my general practice colleagues will identify with the book's content and methodology.

General Practice is written with the recent graduate, the international medical graduate and the medical student in mind. It is a comprehensive textbook that focuses on the very basics of medical principles and management. However, it is hoped that all practitioners will gain useful information from the book's content.

REVIEWERS

The fourth edition underwent a rigorous peer review process to ensure that *General Practice* remains the gold standard reference for general practitioners around the world.

To that end, the author and the publishers extend their sincere gratitude to the following people who generously gave their time, knowledge and expertise.

Content consultants

The author is indebted to the many consultants for their help and advice after reviewing various parts of the manuscript that covered material in their particular area of expertise.

Dr Rob Baird	laboratory investigation
Dr Roy Beran	epilepsy; fits and funny turns; neurological dilemmas
Dr Peter Berger	a diagnostic and management approach to skin problems
Professor Geoff Bishop	basic antenatal care
Dr John Boxall	palpitations
Dr Jill Cargnello	hair disorders
Dr Paul Coughlin and Professor Hatem Salem	bruising and bleeding; thrombosis and thromboembolism
Mr Rod Dalziel	shoulder pain
Dr David Dunn and Dr Hung The Nguyen	the health of Indigenous peoples
Dr Robert Dunne	common skin wounds and foreign bodies
Genetic Health Services, Victoria	genetic disorders
Dr Lindsay Grayson	medical advice for travellers
Dr Michael Gribble	anaemia
Mr John Griffiths	pain in the hip and buttock
Professor Michael Grigg	pain in the leg
Dr Gary Grossbard	the painful knee
Dr Peter Hardy-Smith	the red and tender eye; visual failure
Professor David Healy	abnormal uterine bleeding
Assoc Professor Peter Holmes	cough; dyspnoea; asthma; COPD
Professor Michael Kidd, Dr Ron McCoy and Dr Alex Welborn	human immunodeficiency virus infection
Professor Gab Kovacs	abnormal uterine bleeding; the infertile couple
Professor Even Laerum	research in general practice
Dr Barry Lauritz	common skin problems; pigmented skin lesions
Mr Peter Lawson (deceased) and Dr Sanjiva Wijesinha	disorders of the penis; prostatic disorders
Dr Peter Lowthian	arthritis
Mr Frank Lyons	common fractures and dislocations
Professor Barry McGrath	hypertension
Dr Joe McKendrick	malignant disease
Professor Robyn O ' Hehir	allergic disorders, including hayfever
Dr Michael Oldmeadow	tiredness
Dr Frank Panetta	chest pain

Professor Roger Pepperell	high risk pregnancy
Dr Geoff Quail	pain in the face, sore mouth and tongue
Mr Ronald Quirk	pain in the foot and ankle
Dr Ian Rogers	emergency care
Dr Jill Rosenblatt	the menopause; cervical cancer and Pap smears
Professor Avni Sali	abdominal pain; lumps in the breast; jaundice; constipation; dyspepsia; nutrition
Dr Hugo Standish	urinary tract infection; chronic kidney failure
Dr Richard Stark	neurological diagnostic triads
Dr Paul Tallman	stroke and transient ischaemic attacks
Professor Greg Whelan	alcohol problems
Dr Sanjiva Wijesinha	men's health, scrotal pain, inguinoscrotal lumps
Dr Alan Yung	fever and chills; sore throat
Dr Ronnie Yuen	diabetes mellitus; thyroid and other endocrine disorders

Reviewing consultants

A substantial number of people were involved in the reviewing of this book and their invaluable contribution is acknowledged below. We also take the opportunity to thank the other participants who preferred not to be named in this collective.

Academics

Associate Professor Sue Smith
Dr Elena Ghergori
Dr Fred De Looze
Dr George Kostalas
Dr Sue Hookey
Professor Jon Emery
Professor Mark Nelson
Professor Wes Fabb

General practitioners, educators and registrars

Ashraf Aboud
Mohammed Al Kamil
Anne Balcomb
Jill Benson
Kathy Brotchie
Shane Brun
Daniel Byrne
Paul Carroll
Peter Charlton
Rudolph W. M. Chow
Patrick Clancy
Jennifer Cook-Foxwell
Alice Cunningham
Gabrielle Dellit
Michael Desouza
Matthew Dwyer
Judith Ellis
Say Poh Eng
Iain Esslemont
Marian Evans
Cyril Fernandez
Oliver Frank
Brett Garrett
Tarek Gergis
Naomi Ginges
Jim Griffin
Ranjan Gupta
Hadia Haikal-Mukhtar
Pedita Hall
Mark Henschke
Edward C. Herman
Seyed Ebrahim Hosseini
Brett Hunt
Farhana Hussein
Robyn Hüttenmeister
John inkwater
Aravinda Jawali
Les Jenshel
Fiona Joske
Meredith Joslin
Inas Abdul Karim
Sophia Kennelly
Jim Kourdoulos
Christine Lonergan
Justin Madden
Meredith Makeham
Shahid Malick
Linda Mann
Muhammad Mannan
Cameron Martin
Ronald Mccoy
Robert Meehan
Brad Murphy
Keshwan Nadan
Harry Nespolon
Ching-Luen Ng
Christopher Oh
John Padgett
Peter Parkes
W. J. Patterson
Anoula Pavli
Matthew Penn
Satish Prasad
Tereza Rada
Muhammad Raza
Kate Roe
Daniel Rouhead
Safwat Saba
Amin Sauddin
Kelly Seach
Leslie Segal
Isaac Seidl
Rubini Selvaratnam
Pravesh Shah
Russell Shute
G. Sivasambu
Jane Smith
Lucie Stanford
Sean Stevens
Hui Tai Tan
Judy Toman
Khai Tran
Anthony Wickins
Belinda Woo

Overseas trained doctors and international medical graduates

Ibrahim K. Botros
Gordana Cuk
Nazih Hamzeh
Erfanul Haque
Dac Luu
Hemant Mahagaonkar
Charles Mutandwa
Mitra Babazadeh Shahri

Yock Seck Ding
Diosdado Javellana
Patrick Mulhern
Heinz Tilenius

Medical students

Barrie Coulson
Therese Cox
Anthony Fok
Rosalyn Hunt
Gloria Jove
Megha Mulchandani
Brent O'Carrigan
Jamie Sharples
Joseph V. Turner

THE AUTHOR

John Murtagh AM

MBBS, MD, BSc, BEd, FRACGP, DipObstRCOG

Professor in General Practice, School of Primary Health,
Monash University, Melbourne

Professorial Fellow, Department of General Practice,
University of Melbourne

Adjunct Clinical Professor, Graduate School of Medicine,
University of Notre Dame, Fremantle, Western Australia

JMurtagh

John Murtagh was a science master teaching chemistry, biology and physics in Victorian secondary schools when he was admitted to the first intake of the newly established Medical School at Monash University, graduating in 1966. Following a comprehensive postgraduate training program, which included surgical registrarship, he practised in partnership with his medical wife, Dr Jill Rosenblatt, for ten years in the rural community of Neerim South, Victoria.

He was appointed Senior Lecturer (part-time) in the Department of Community Medicine at Monash University and eventually returned to Melbourne as a full-time Senior Lecturer. He was appointed to a professorial chair in Community Medicine at Box Hill Hospital in 1988 and subsequently as chairman of the extended department and Professor of General Practice in 1993 until retirement from this position in 2000. He now holds teaching positions as Professor in General Practice at Monash University, Adjunct Clinical Professor, University of Notre Dame and Professorial Fellow, University of Melbourne. He combines these positions with part-time general practice, including a special interest in musculoskeletal medicine. He achieved the Doctor of Medicine degree in 1988 for his thesis ‘ The management of back pain in general practice ’.

He was appointed Associate Medical Editor of *Australian Family Physician* in 1980 and Medical Editor in 1986, a position held until 1995. In 1995 he was awarded the Member of the Order of Australia for services to medicine, particularly in the areas of medical education, research and publishing.

One of his numerous publications, *Practice Tips*, was named as the British Medical Association's Best Primary Care Book Award in 2005. In the same year he was named as one of the most influential people in general practice by the publication ‘ Australian Doctor ’ . John Murtagh was awarded the inaugural David de Krester medal from Monash University for his exceptional contribution to the Faculty of Medicine, Nursing and Health Sciences over a significant period of time. Members of the Royal Australian College of General Practitioners may know that he was recently bestowed the honour of the namesake of the College library.

Today John Murtagh continues to enjoy active participation with the diverse spectrum of general practitioners - whether they are students or experienced practitioners, rural or urban based, local or international medical graduates, clinicians or researchers. His vast experience with all of these groups has provided him with tremendous insights into their needs, which is reflected in the culminated experience and wisdom of Murtagh's *General Practice* .

WHAT IS NEW?

The fourth edition is a substantial content revision with updating throughout all of the book. Highlights of the revision include:

- **Six new chapters:**
  - Nasal disorders
  - Surgical problems in children
  - Infections in pregnancy
  - Chronic obstructive pulmonary disease
  - Chronic heart failure
  - Thrombosis and thromboembolism
- **Diabetes has been expanded** to two chapters of coverage:
  - Diabetes mellitus - diagnosis
  - Diabetes mellitus - management
- **A greater emphasis on chronic disease management** throughout the book reflects the changing nature of general practice.
- **Rarer conditions have now been included** so that *General Practice* is the one essential source where you can find everything you need. All content has been thoroughly reviewed and revised.
- **Selected content has been moved** to the *General Practice Companion Handbook* , such as emergency medicine, rational antibiotic prescribing, and special community disasters - allowing access to such vital information either inside or outside of the practice.
- **Full colour presentation** with an outstanding design allows you to find what you need more quickly and gain more from illustrations and photos.

A summary of the major new inclusions is provided in tabular form.

Chapter	Chapter title	Major updates - new or expanded coverage from the 3e
Part 1	**The basis of general practice**	
1	The nature and content of general practice	Symptoms and conditions related to litigation; chronic disease management
2	The family	
3	Consulting skills	
4	Communication skills	Use of analogy; ‘ road blocks ’ to good communication
5	Counselling skills	Problem gambling
6	Difficult, demanding and angry patients	
7	Palliative care	Gold Standards Framework (UK); opioid rotation
8	The elderly patient	Loneliness in the elderly; rules of ‘ 7 ’ for the non-coping elderly patient; mental state examination tests; driving; later-life depression and suicide
9	Prevention in general practice	Updated immunisation; colorectal cancer and prostate cancer
10	Nutrition in health and illness	
11	Health promotion and patient education	Promotion of healthy lifestyle; introduction of the SNAP guide
12	Pain and its management	Coxibs; current use of paracetamol; somatoform disorders
13	Whole person approach to management	
14	Travel and tropical medicine	General updates throughout
15	Research and evidence-based medicine	More on qualitative approaches including phenomenology, ethnography, grounded theory
16	Laboratory investigations	

Part 2	**Diagnostic perspective in general practice**	
17	Inspection as a clinical skill	Adenoid facies; choleric facies; smoker ' s facies; uraemic facies
18	A safe diagnostic strategy	Acknowledgment of bullying in workplace and stress
19	Genetic conditions	Expanded terminology and definitions; inherited adult onset neurological disorders; hereditary haemoglobinopathies; haemolytic disorders, bleeding and clotting disorders
20	Depression	Postpartum depression; current information on antidepressants and pharmacological management; rule of ‘ 7 ’
21	Diabetes mellitus: diagnosis	**90 % NEW coverage, now with two chapters on diabetes diagnosis and management**
22	Drug problems	
23	Anaemia	Anaemia and bone marrow; Vitamin B12 deficiency
24	Thyroid and other endocrine disorders	
25	Spinal dysfunction	
26	Urinary tract infection	Cranberry juice and UTI prevention; prostatitis; vulvovaginitis in children
27	Malignant disease	Common cancers and 5 year survival rate
28	HIV/AIDS - could it be HIV?	Pneumocystitis carimi
29	Baffling viral and protozoal infections	SARS and avian bird flu
30	Baffling bacterial infections	Antibiotic prophylaxis; Hansen ' s disease (leprosy)
31	Chronic kidney disease	Chronic kidney disease classification; ACR guidelines drug prescribing; goals of management
32	Connective tissue disorders and the vasculitides	Autoimmune diseases; connective tissue disorders and vasculitides; CREST syndrome; Sj ö gren ' s syndrome; Raynaud ' s phenomenon; Takayasu ' s arteritis; Bechet ' s syndrome
33	Neurological dilemmas	Motor neurone disease; Parkinson ' s; cognitive impairment with Parkinson ' s
Part 3	**Problem solving in general practice**	
34	Abdominal pain	Updating of abdominal pain
35	Arthritis	Features of viral arthritis; pharmacological management; new therapies in DMARDs; management of gout and pharmacological treatment of gout
36	Anorectal disorders	
37	Low back pain	Yellow flag pointers in low back pain
38	Thoracic back pain	Red flag pointers in thoracic back pain
39	Bruising and bleeding	Henoch sch ö nlein purpura (HSP); acute thrombocytopema of childhood; splenectomy
40	Chest pain	Intra-coronary stents; hospital management; fibrinolytic therapy; oral anticoagulation
41	Constipation	
42	Cough	
43	Deafness and hearing loss	

44	Diarrhoea	Whipple disease; blastocystitis hominus
45	The disturbed patient	Somatisation; dysmorphophobia; conversion; dissociation; the acutely psychotic patient; cardiac dysfunction; body dysmorphic disorder
46	Dizziness	
47	Dyspepsia	
48	Dysphagia	Odynophagia
49	Dyspnoea	Pleural effusion; pulmonary fibrosis; acute respiratory distress syndrome (ARDS); severe acute respiratory syndrome (SARS)
50	The painful ear	Necrotising otitis externa
51	The red and tender eye	The painful red eye; key symptoms; features of episcleritis, scleritis and uveitis (iritis); corneal disorders; endophthalmitis
52	Pain in the face	Ludwig ' s angina; chronic sinusitis; temporal arteritis
53	Fever and chills	
54	Faints, fits and funny turns	
55	Haematemesis and melaena	Drugs associated with gastrointestinal haemorrhage
56	Headache	Post-lumbar puncture headache
57	Hoarseness	
58	Jaundice	Guidelines on jaundice
59	Nasal disorders	**ALL NEW**
60	Nausea and vomiting	Gastroparesis
61	Neck lumps	Thyroid nodule
62	Neck pain	Acute neck pain; evidence of benefit
63	Shoulder pain	Common shoulder conditions; rotator cuff tears; shoulder instability; osteoarthritis of the glenohumeral joint; glenoid labrum
64	Pain in the arm and hand	Intersection syndrome; ischaemic necrosis; ganglia; Raynaud ' s phenomenon
65	Hip and buttock pain	Avulsion body injuries; avascular necrosis; groin pain; fascia latia syndrome; ischial bursitis
66	Pain in the leg	
67	The painful knee	Anterior, lateral and medial knee pain; the Ottowa knee rules; complex regional pain syndrome 1; Baker ' s cyst
68	Pain in the foot and ankle	Fat pad disorder treatment
69	Walking difficulty	Flat feet, claw feet; hammer toes, claw toes
70	Palpitations	Atrial flutter; atrial fibrillation
71	Sleep disorders	
72	Sore mouth and tongue	Halitosis
73	Sore throat	Recurrent tonsilitis
74	Tiredness	
75	The unconscious patient	
76	Urinary disorders	Overactive bladder; uterovaginal prolapse
77	Visual failure	Referral; floaters and flashes

78	Weight gain	
79	Weight loss	
Part 4	**Child and adolescent health**	
80	An approach to the child	Guidelines for feeding infants; guidelines for toilet training; blocked nasocacrimal duct; growing pains; constipation; DSM IV criteria for enuresis; Mongolian blue spot
81	Specific problems of children	
82	Surgical problems in children	**ALL NEW**
83	Common childhood infectious diseases (including skin eruptions)	
84	Behaviour disorders in children	Habit cough; Asperger ' s disorder; childhood bullying
85	Child abuse	
86	Emergencies in children	Paediatric advanced life support; button and disc battery ingestion; grading system for croup; red flags bile
87	Adolescent health	Aged and informed consent; major depression
Part 5	**Women ' s health**	
88	Cervical cancer and Pap smears	New lesion classification; medico-legal issues; vaccination
89	Family planning	Delaying a period
90	Breast pain (mastalgia)	
91	Lumps in the breast	
92	Abnormal uterine bleeding	Uterine fybroids; carcinoma of the cervix; endometrial cancer
93	Lower abdominal and pelvic pain in women	Features of chronic pain; pelvic congestion syndrome
94	Premenstrual syndrome	
95	The menopause and osteoporosis	
96	Vaginal discharge	
97	Vulvar disorders	Vulvovaginitis in pre-pubertal girls; mild vulvovaginitis; moderate/persisent vulvovaginitis; labial adhesions; Bartholin ' s cyst
98	Violence against women	Management issues of sexual assault
99	Basic antenatal care	Hypotension; pruritis; obesity; breathlessness in pregnancy
100	Infections in pregnancy	**ALL NEW**
101	High-risk pregnancy	**90 % NEW Maternal mortality; perinatal mortality; guidelines for specialist obstetric consultation; hypertensive disorders; medical conditions in pregnancy; multiple pregnancy; preterm labour; trauma; drugs**
102	Postnatal care	Tiredness; hair loss; back pain and coccygodynia
Part 6	**Men's health**	
103	Men ' s health: an overview	
104	Scrotal pain	
105	Inguinoscrotal lumps	Sperm granulomas; comparison of common testicular cancers; vasectomy
106	Disorders of the penis	
107	Disorders of the prostate	

Part 7	**Sex-related problems**	
108	The subfertile couple	
109	Sexual health	Oral medication; gender identity concerns
110	Sexually transmitted infections	
Part 8	**Problems of the skin**	
111	A diagnostic and management approach to skin problems	Erythema, miium, papilloma; zallus, exfoliation, keratoderma; topical corticosteroids for chronic dermatoses
112	Pruritis	
113	Common skin problems	Golfer ' s vasculitis; flea bites, bed bug bites; differences between chilblains and Raynaud ' s phenomenon
114	Acute skin eruptions	
115	Skin ulcers	
116	Common lumps and bumps	Stucco keratoses; sebaceous hyperplasia; lumps on ears; chondrodermatitis nodularis helicus; bursae
117	Pigmented skin lesions	The early nodular melanoma problem; pitfalls in diagnosis of melanoma; guidelines for excision margins
118	Hair disorders	Causes of virilisation
119	Nail disorders	Myxoid pseudocyst
Part 9	**Chronic disorders: continuing management**	
120	Alcohol problems	Risk of harm in short and long term
121	Allergic disorders including hay fever	Allergic rhinitis
122	Anxiety	Body dysmorphic disorder; somatisation disorder
123	Asthma	Asthma severity classification; Omalizumab; starting treatment; stepwise interval management plan; plan for children
124	Chronic obstructive pulmonary disease	**ALL NEW**
125	Epilepsy	Current anti-epileptic drugs
126	Hypertension	Risk stratification and calculation of cardiovascular risk; pharmacological management
127	Dyslipidaemia	Current lipid-lowering drugs
128	Diabetes mellitus: management	**80 % NEW with more on Type 2 diabetes; starting insulin in Type 2 diabetes; metabolic complications of diabetes; management in summary; diabetes control estimation of cardiovascular risk in the future**
129	Chronic heart failure	**ALL NEW**
Part 10	**Accident and emergency medicine**	
130	Emergency care	Advanced life support
131	The doctor ' s bag	
132	Stroke and transient ischaemic attacks	Differential diagnosis ' stroke mimics '
133	Thrombosis and thromboembolism	**ALL NEW**
134	Common skin wounds and foreign bodies	Burns to the hand
135	Common fractures and dislocations	Patella dislocation and subluxations; ankle/talus/subtular joint dislocations
136	Common sporting injuries	Abductor muscle or tendon injury; talar dome injury; tibiofibular

		syndesmosis injury
Part 11	**Indigenous health**	
137	The health of Indigenous peoples	CARPA Standard Treatment Manual; cardiovascualr disease; GABHS infections; Hansen's disease (leprosy)
Appendices		

CHAPTER 1. THE NATURE AND CONTENT OF GENERAL PRACTICE

Medical practice is not knitting and weaving and the labour of the hands, but it must be inspired with soul and be filled with understanding and equipped with the gift of keen observation; these together with accurate scientific knowledge are the indispensable requisites for proficient medical practice.

MOSES BEN MAIMON (1135 - 1204)

THE NATURE AND CONTENT OF GENERAL PRACTICE

General practice is a traditional method of bringing primary health care to the community. It is a medical discipline in its own right, linking the vast amount of accumulated medical knowledge with the art of communication.

DEFINITIONS

General practice can be defined as that medical discipline which provides ‘community-based, continuing, comprehensive, preventive primary care’, sometimes referred to as the CCCP model.

The Royal Australian College of General Practitioners (RACGP) uses the following definitions of general practice and primary care:

> General practice is that component of the health care system which provides initial, continuing, comprehensive and co-ordinated medical care for all individuals, families and communities and which integrates current biomedical, psychological and social understandings of health.
>
> General practitioner is a medical practitioner with recognised generalist training, experience and skills, who provides and co-ordinates comprehensive medical care for individuals, families and communities.
>
> Primary care involves the ability to take responsible action on any problem the patient presents, whether or not it forms part of an ongoing doctor - patient relationship. In managing the patient, the general/family practitioner may make appropriate referral to other doctors, health care professionals and community services. General/family practice is the point of first contact for the majority of people seeking health care.
>
> In the provision of primary care, much ill-defined illness is seen; the general/family practitioner often deals with problem complexes rather than with established diseases.
>
> The practitioner must be able to make a total assessment of the person's condition without subjecting the person to unnecessary investigations, procedure and treatment.

The American Academy of Family Physicians (AAFP)[1] and the American Board of Family Practice (ABFP) have defined family practice as:

> … the medical specialty that provides continuing and comprehensive health care for the individual and the family. It is the specialty in breadth that integrates the biological, clinical and behavioural sciences. The scope of family practice encompasses all ages, both sexes, each organ system and disease entity.

The AAFP has emphasised that the primary objective of the specialty of family practice is to deliver primary health care, described as:

> … a form of medical care delivery that emphasises first contact care and assumes ongoing responsibility for the patient in both health maintenance and therapy of illness. It is personal care involving a unique interaction and communication between the patient and the physician. It is comprehensive in scope and includes the overall co-ordination of care of the patient's health problems, be they biological, behavioural or social. The appropriate use of consultants and community resources is an important part of effective primary care.

The AAFP has expanded on the function of delivery of primary health care.[1, 2]

Primary care is a form of delivery of medical care that encompasses the following functions:

1. It is ‘first-contact’ care, serving as a point-of-entry for patients into the health care system.
2. It includes continuity by virtue of caring for patients over a period of time, both in sickness and in health.

3. It is comprehensive care, drawing from all the traditional major disciplines for its functional content.
4. It serves a co-ordinative function for all the health care needs of the patient.
5. It assumes continuing responsibility for individual patient follow-up and community health problems; and
6. It is a highly personalised type of care.

Pereira Gray[3] identifies six principles - primary care, family care, domiciliary care and continuing care all designed to achieve preventive and personal care. ‘We see the patient as a whole person and this involves breadth of knowledge about each person, not just depth of disease.’

General practice is not the summation of specialties practised at a superficial level and we must avoid the temptation to become ‘specialoids’. In the current climate where medicine is often fragmented there is a greater than ever need for the generalist. The patient requires a trusted focal point in the often bewildering health service jungle. Who is to do this better than the caring family doctor taking full responsibility for the welfare of the patient and intervening on his or her behalf? Specialists also need highly competent generalists to whom they can entrust ongoing care.

UNIQUE FEATURES OF GENERAL PRACTICE

Anderson, Bridges-Webb and Chancellor[4] emphasise that ‘the unique and important work of the general practitioner is to provide availability and continuity of care, competence in the realm of diagnosis, care of acute and chronic illness, prompt treatment of emergencies and a preventive approach to health care’.

The features that make general practice different from hospital- or specialist-based medical practices include:

- first contact
- diagnostic methodology
- early diagnosis of life-threatening and serious disease
- continuity and availability of care
- personalised care
- care of acute and chronic illness
- domiciliary care
- emergency care (prompt treatment at home or in the community)
- family care
- palliative care (at home)
- preventive care
- scope for health promotion
- holistic approach
- health care co-ordination

Apart from these processes the GP has to manage very common problems including a whole variety of problems not normally taught in medical school or in postgraduate programs. Many of these problems are unusual yet common and can be regarded as the ‘nitty gritty’ or ‘bread and butter’ problems of primary health care.

CONTINUING CARE

The essence of general practice is continuity of care. The doctor - patient relationship is unique in general practice in the sense that it covers a span of time which is not restricted to a specific major illness. The continuing relationship involving many separate episodes of illness provides an opportunity for the doctor to develop considerable knowledge and understanding of the patient, the family and its stresses, and the patient's work and recreational environment.

Strategies to enhance continuing care

A philosophical commitment

Underlying appropriate patient care is the attitude of the provider. A caring, responsible practitioner who is competent, available and a trusted friend is ‘like gold’ to his or her patients.

Medical records

An efficient medical record system is fundamental. Ideally, it should include a patient profile, a database, problem lists, special investigation lists, medication lists, adverse drug reactions and ‘at risk’ details.

Checklists

The use of checklists or questionnaires to assemble information on presenting problems may enhance knowledge as well as assist earlier diagnosis.

Home visits

Home visits are a goldmine of information about intrafamily dynamics. They should cement the doctor - patient relationship if used appropriately and discreetly. We are the only doctors who practise domiciliary care. We must treasure it. Sitting in the office chair practising ‘conveyor belt’ medicine is contrary to the ideals of general practice.

Anticipatory guidance

Unfortunately patients do not usually perceive the family doctor as a counsellor, but opportunities should be taken to offer advice about anticipated problems in situations such as premarital visits, antenatal care and preadolescent contact.

Patient education

Whenever possible, patients should be given insight into the nature of their illness, and reasons for the treatment and prognosis. Patient education leaflets, such as those published in journals, can be used as a starting point, although there is no substitute for careful personal explanation. This should lead to better compliance and an improved relationship between doctor and patient.

Personal health records

These excellent wallets, which are handed to parents of newborn babies, have a very important place in the ongoing care of children. Their purpose is to supply an outline of preventive health care, beginning from birth. They provide an inbuilt recall list directed at a most compliant source - mothers. In fact, they provide a complete record of health care throughout a person's lifetime.

Patient register

An age-and-sex register of all patients in the practice is a very useful acquisition. The main strategy is to find out who are the patients, what are their basic characteristics and who suffers from chronic diseases, such as cancer, diabetes and emphysema.

Recall lists

Use of recall lists based on the patient register should significantly improve health care delivery. Dentists have been using this technique successfully for some time. In the United States, Canada and many other countries doctors use recall lists regularly to remind patients that preventive items, such as immunisation schedules and cancer smear tests, are due.

Computers

Computers have simplified and streamlined the design and use of practice registers and patient-recall systems in addition to their use for accounting purposes. Their potential for patient education and doctor education is considerable.

COMMON PRESENTING SYMPTOMS

Common presenting symptoms in Australian practices are presented in Table 1.1,[5] where they are compared with those in the United States.[6] The similarity is noticed but the different classification system does not permit an accurate comparison. In the third national survey of morbidity in general practice in Australia[5] the most common symptoms described by patients were cough (6.2 per 100 encounters), throat complaints (3.8 per 100), back complaints (3.6 per 100) and upper respiratory tract infection (URTI) (3.2 per 100). In addition, very common presentations included a check-up (13.7 per 100) and a request for prescription (8.2 per 100). McWhinney lists the 10 most common presenting symptoms from representative Canadian and British practices but they are divided between males and females.[7]

Table 1.1 Most frequent presenting problems/symptoms (excluding pregnancy, hypertension, immunisation and routine check-up)

	Australia	**United States**
Cough	1	1
Throat complaint	2	2
Back pain	3	4
URTI	4	11
Rash	5	5
Abdominal pain	6	6
Depression	7	
Ear pain	8	3
Headache	9	10
Fever	10	7
Weakness/tiredness	11	
Diarrhoea	12	
Asthma	13	
Nasal congestion/sneeze	14	12
Chest pain	15	13
Knee complaint	16	8
Visual dysfunction		9

Source: Australian figures: Britt et al.[5]; United States figures (all specialties): De Lozier & Gagnon[6]

For males in the Canadian study these symptoms are (in order, starting from the most common) cough, sore throat, colds, abdominal/pelvic pain, rash, fever/chills, earache, back problems, skin inflammation and chest pain.

For females the five other symptoms that are included are menstrual disorders, depression, vaginal discharge, anxiety and headache.

In the British study the most common symptoms are virtually identical between males and females and include cough, rash, sore throat, abdominal pain, bowel symptoms, chest pain, back pain, spots, sores and ulcers, headache, muscular aches and nasal congestion.[8]

Most frequent presenting symptoms in the author's practice

The most common presenting symptoms in the author's practice[9] were identified, with the emphasis being on pain syndromes:

- cough
- disturbance of bowel function
- pain in abdomen

- pain in back
- pain in chest
- pain in head
- pain in neck
- pain in ear
- pain in throat
- pain in joints/limbs
- rashes
- sleep problems
- tiredness/fatigue
- vaginal discomfort

These symptoms should accurately reflect Australian general practice since the rural practice would represent an appropriate cross-section of the community's morbidity, and the recording and classification of data from the one practitioner would be consistent.

SYMPTOMS AND CONDITIONS RELATED TO LITIGATION

Medical defence organisations have highlighted the following areas as being those most vulnerable for management mishaps:

- acute abdominal pain
- acute chest pain
- breast lumps
- children's problems, especially the sick febrile child <2 years, groin pain and lumps
- dyspnoea ±cough (? heart failure, cancer, TB)
- headache

COMMON MANAGED DISORDERS

Excluding a general medical examination, hypertension and upper respiratory tract infection (URTI) were the two most common problems encountered in both the Australian and US[10] studies. The 23 most frequent individual disorders are listed in Table 1.2 and accounted for over 40% of all problems managed.[5, 11]

Table 1.2 Most frequently managed disorders/diagnoses (rank order) excluding prescriptions

	Australia	United States
General medical examination	1	1
Hypertension	2	3
URTI	3	2
Immunisation	4	[*]
Depression	5	6[†]
Acute bronchitis/bronchiolitis	6	13
Asthma	7	29
Back complaint	8	
Diabetes mellitus	9	8
Lipid metabolism disorder	10	[*]

Osteoarthritis	11	10
Sprain/strain	12	5
Contact dermatitis	13	9
Acute otitis media	14	18
Anxiety	15	6[†]
Sleep disorders/insomnia	16	-
Urinary tract infection	17	11
Female genital check-up, Pap smear	18	(under 1)
Sinusitis	19	25
Occupational check-up	20	27
Oesophageal disease	21	27
Menopause complaint	22	27
Viral disease	23	-
	Soft tissue injuries	4
	Ischaemic heart disease	7

[*] not listed

[†] combined

Source: Australian figures: Britt et al.[5]; United States figures: Rosenblatt et al.[10]

The content of this textbook reflects what is fundamental to the nature and content of general practice - that which is common but is significant, relevant, preventable and treatable.

CHRONIC DISEASE MANAGEMENT

A study of international target conditions[12] in chronic disease management have highlighted the importance of the following (as common themes)

- coronary heart disease
- chronic heart failure
- stroke
- hypertension
- diabetes mellitus type 2
- chronic obstructive pulmonary disease
- asthma
- epilepsy
- hypothyroidism
- chronic mental illness
- medication monitoring

REFERENCES

1. American Academy of Family Physicians. Official definition of Family Practice and Family Physician (AAFP Publication No. 303). Kansas City, Mo, AAFP, 1986.
2. Rakel RE. *Essentials of Family Practice*. Philadelphia: WB Saunders Company, 1993: 2 - 3.

3. Pereira Gray DJ. Just a GP. JR Coll Gen Pract, 1980; 30: 231 - 9.
4. Anderson NA, Bridges-Webb C, Chancellor AHB. *General Practice in Australia*. Sydney: Sydney University Press, 1986: 3 - 4.
5. Britt H, Sayer GP et al. *Bettering the Evaluation and Care of Health: General Practice in Australia* 1998 - 9. Sydney: University of Sydney & the Australian Institute of Health & Welfare, 1998 - 99.
6. De Lozier JE, Gagnon RO. *1989 Summary: National Ambulatory Medical Care Survey*. Hyattsville, Md, National Centre for Health Statistics, 1991.
7. McWhinney IR. *A Textbook of Family Medicine* (2nd edn). New York: Oxford University Press, 1997: 40 - 4.
8. Wilkin D, Hallam L et al. *Anatomy of Urban General Practice*. London: Tavistok, 1987.
9. Murtagh JE. *The Anatomy of a Rural Practice*. Melbourne: Monash University, Department of Community Practice Publication, 1980: 8 - 13.
10. Rosenblatt RA, Cherkin DC, Schneeweiss R et al. The structure and content of family practice: Current status and future trends. J Fam Pract, 1982; 15(4): 681 - 722.
11. Bridges-Webb C, Britt H, Miles D et al. Morbidity and treatment in general practice in Australia. Aust Fam Physician, 1993; 22: 336 - 46.
12. Piterman L *Chronic Disease Management OSP Report*. Melbourne: Monash University, 2004.

CHAPTER 2. THE FAMILY

The family only represents one aspect, however important an aspect, of a human being's functions and activities - A life is beautiful and ideal, or the reverse, only when we have taken into our consideration the social as well as the family relationship.

HAVELOCK ELLIS 1922, *LITTLE ESSAYS OF LOVE AND VIRTUE*

THE FAMILY

Working with families is the basis of family practice. Families living in relative harmony provide the basis for the good mental health of their members and also for social stability.

However, the traditional concept of the nuclear family where the wife stays at home to care for the children occurs in only about 15% of Australian families, and approximately 46% of Australian marriages end in separation. This results in many psychosocial problems that family doctors will have to address.

Family therapy is ideally undertaken by general practitioners, who are in a unique position as providers of continuing care and family care. It is important for them to work together with families in the counselling process and to avoid the common pitfalls of working in isolation and assuming personal responsibility for changing the family.

Bader1 summarises working with families succinctly:

> From the perspective of family therapy, working with families means avoiding the trap of being too directive, too responsible for the family's welfare, with the result that the family becomes overly dependent on the general practitioner for its health and development. From the perspective of family education, working with families means developing the skills of anticipating guidance, helping families to prepare, not only for the normal changes occurring as the family develops, but also for the impact of illness on the family system.

CHARACTERISTICS OF HEALTHY FAMILIES

Successful families have certain characteristics, an understanding of which can give the family doctor a basis for assessing the health of the family and a goal to help set targets for change in disrupted families. Such characteristics are:

- *Healthy communication*. In this situation family members have freedom of expression for their feelings and emotions.
- *Personal autonomy*. This includes appropriate use of power sharing between spouses.
- *Flexibility*. This leads to appropriate ‘give and take’ with adaptation to individual needs and changing circumstances.
- *Appreciation*. This involves encouragement and praise so that members develop a healthy sense of self-esteem.
- *Support networks*. Adequate support from within and without the family engenders security, resistance to stress and a healthy environment in general. The family doctor is part of this network.
- *Family time and involvement*. Studies have shown that the most satisfying hallmark of a happy family is ‘doing things together’.
- *Spouse bonding*. The importance of a sound marital relationship becomes obvious when family therapy is undertaken.
- *Growth*. There needs to be appropriate opportunities for growth of individual family members in an encouraging atmosphere.
- *Spiritual and religious values*. An attachment to spiritual beliefs and values is known to be associated with positive family health, supporting the saying ‘The family that prays together stays together’.

FAMILIES IN CRISIS

Doctors are closely involved with families who experience unexpected crises, which include illnesses, accidents, divorce, separation, unemployment, death of a family member and financial disasters.

The effect of illness

Serious illness often precipitates crises in individual members of the family, crises which have not previously surfaced in the apparently balanced family system. It is recognised, for example, that bereavement over the unexpected loss of a child may lead to marital breakdown, separation or divorce.

In the long term, other family members may be affected more than the patient. This may apply particularly to children and manifest as school underachievement and behaviour disturbances.

During the crisis the obvious priority of the doctor is to the patient but the less obvious needs of the family should not be ignored.

Guidelines for the doctor

- Include the family as much as possible, starting early in the acute phase of the illness. It may necessitate family conferences.
- Include the family on a continuing basis, especially if a long-term illness is anticipated. It is helpful to be alert for changes in attitudes, such as anger and resentment towards the sick member.
- Include the family in hospital discharge planning.
- If a serious change in family dynamics is observed, the use of experts may be needed.

SIGNIFICANT PRESENTATIONS OF FAMILY DYSFUNCTION

The following presentations may be indicators that all is not well in the family, and so the doctor needs to ‘think family’:

- marital or sexual difficulties
- multiple presentations of a family member - ‘the thick file syndrome’
- multiple presentations by multiple family members
- abnormal behaviour in a child
- the ‘difficult patient’
- inappropriate behaviour in the antenatal and/or postpartum period
- drug or alcohol abuse in a family member
- evidence of physical or sexual abuse in wife or child
- psychiatric disorders
- susceptibility to illness
- increased stress/anxiety
- complaints of chronic fatigue or insomnia

It is important that the family doctor remains alert to the diversity of presentations and takes the responsibility for identifying an underlying familybased problem.

THE PATIENT AND FAMILY DYNAMICS

Family doctors see many patients who present with physical symptoms that have primarily an emotional or psychosocial basis with either little or no organic pathology. As many as 50 - 75% of patients utilising primary care clinics have a psychosocial precipitant as opposed to biomedical problems as the main cause of their visit.[2]

In order to understand the clinical manifestations of the sick role of patients, family doctors should first understand the individual's response to stress stimuli, which may come from external (family, work or sexual behaviour) or internal (personality trait or psychosocial) sources (Figure 2.1 and Table 2.1).


External sources
Dysfunction of
• family
• work
• sexuality
somatic symptoms
behaviour (sick role)
Threshold
values
Growing environment
feelings
expectations
personality trait
Internal sources

Figure 2.1 Family dynamics and psychosomatic illness iceberg

Table 2.1 Areas of possible biopsychosocial dysfunction

Work	Family	Sex
• Type of work	• Present family (change of structure and function)	• Sexual dysfunction
	• Workload	• Disharmony
• Work environment	• Extended family (parents and relatives)	• Deprivation
	• Goals	• Guilt
• Work satisfaction	• Growing environment (family tree)	

HOW TO EVALUATE THE FAMILY DYNAMICS

- Carefully observe family members interacting.
- Invite the whole family to a counselling session (if possible).
- Visit the home: an impromptu home visit (with some pretext such as a concern about a blood test result) on the way home from work may be very revealing.
- Prepare a genogram (Figure 19.1): family dynamics and behaviour can be understood by drawing a family map or genogram (a diagrammatic representation of family structure and relationships).[3, 4]

The genogram

The genogram is a very valuable pedigree chart that usually covers three generations of a family tree.[3] Genograms are a useful strategy for involving family members who may have been reluctant to be involved in discussions on family matters.[4] An example, including the use of symbols, is shown in Figure 19.1.

THE FAMILY LIFE CYCLE

Helpful in understanding the dynamics of the family is the concept of the family life cycle,[5] which identifies several clearly defined stages of development (Table 2.2). Such an understanding can help the doctor form appropriate hypotheses about the problems patients are experiencing at a particular stage. Each stage brings its own tasks, happiness, crises and difficulties. This cycle is also well represented in Figure 2.2 overleaf (after McWhinney[6] and Duvall[7]), which indicates the approximate length of time on each of the stages.

Table 2.2 The family life cycle[1]

Stage	Tasks to be achieved
1 Leaving home	Establishing personal independence. Beginning the emotional separation from parents.
2 Getting married	Establishing an intimate relationship with spouse. Developing further the emotional separation from parents.
3 Learning to live together	Dividing the various marital roles in an equitable way. Establishing a new, more independent relationship with family.
4 Parenting the first child	Opening the family to include a new member. Dividing the parenting roles.
5 Living with the adolescent	Increasing the flexibility of the family boundaries to allow the adolescent(s) to move in and out of the family system.
6 Launching children: the empty-nest phase	Accepting the multitude of exits from and entries into the family system. Adjusting to the ending of parenting roles.
7 Retirement	Adjusting to the ending of the wage-earning roles. Developing new relationships with children, grandchildren and each other.
8 Old age	Dealing with lessening abilities and greater dependence on others. Dealing with losses of friends, family members and, eventually, each other.


2 years
2.5 years
3.5 years
married couples (no children)
childbearing families
families with preschool children
families with school children
ageing family members
10 to 15 ± years
1
2
3
8
4
7 years
retirement
5
7 years
families with teenagers
middle-aged parents
7
6
15 ± years
families with young adults leaving home
8 years

Figure 2.2 The family life cycle (approximate time in each stage). After McWhinney[6] and Duvall

Family assessment

The assessment of families with problems can be formalised through a questionnaire which allows the collection of information in a systematic way in order to give an understanding of the functioning of the family in question.

The questionnaire[1]

1 Family of origin

- Could each of you tell us something about the families you grew up in?
- Where do you come in the family?
- Were you particularly close to anyone else in the family?
- Were there any severe conflicts between family members?
- Did anyone abuse you in any way?
- Do you have much contact with any of your family now?
- Have you tried to model (or avoid) any features for your own family?

2 History of the couple's relationship

- How did you two meet?
- What attracted you to each other?
- Why did you marry this person rather than someone else?
- How did your families react to your choice?
- How did the birth of your children affect your relationship?
- When was your relationship at its best? Why?

3 Experience in counselling and enrichment

- Have any of you been to ‘marriage encounter’ or similar programs?
- Have any of you been to any form of counselling?
- Did you go alone or with another family member?
- What did you like or dislike about the experience?
- In what way was it helpful or unhelpful?

4 Expectations and goals

- Whose idea was it to come here?
- What was the reaction of other family members?
- Why did you come now?
- Was there any particular event that triggered the decision?
- What do each of you hope to gain by coming for an assessment?

5 Family function[1]

- What is it like for each of you to live in this family? (If children are present, they should be asked first.)
- Do you have any difficulty in talking to other members of the family? (Again, children first.)
- Do you have any difficulty in expressing appreciation to each other? (Mention here that studies on healthy families show that both communication and appreciation rank in the top qualities.)
- How do you show appreciation in this family?
- How do you show affection in this family? (Again, children first.)
- How satisfied are you with the present arrangement? Are there any changes you would like to see?
- What ways have you used to resolve disagreements or change the way the family functions?

Assessment based on the questionnaire

- family members present in interview (names and ages)
- missing members (names and ages)
- presenting problems or reasons for family interview identified by whom? Any attempted solutions?
- roles - structure, organisation (who is dominant and so on)
- affect - predominant emotional tone and expressed emotions
- communication - who dominates? Who talks? who listens to whom?

- stage in the family life cycle
- illness and sickness roles
- coping mechanisms

FAMILY-BASED MEDICAL COUNSELLING

There are several brief counselling models to assist the family doctor in probing and counselling, using a simple infrastructure such as the BATHE model.

The BATHE technique[8]

This really represents a diagnostic technique to identify sources of disharmony, which can act as a springboard for counselling.

The acronym BATHE stands for background, affect, trouble, handling and empathy, and can be summarised as follows.

Background

Enquire about possible areas of psychosocial problems to help elicit the context of the patient's visit.

- What is happening in your life?
- Is there anything different since before you got sick?
- How are things at home?

Affect

Affect is the ‘feeling state’ and includes anxiety, so it is wise to probe potentially sensitive areas.

- How do you feel about what is going on in your life?
- How do you feel about your home life?
- How do you feel about work/school?
- How do you feel about your (husband or wife or daughter or …)?
- What is your mood like? Do you feel sad or happy?

Trouble

Enquire about how the patient's problems are troubling the patient.

- What about the situation troubles you most?
- What troubles or worries you most in your life?
- What worries you most at home?
- How stressed and upset are you about this problem?
- How do you think this problem affects you?

Handling

- How are you handling this problem?
- Do you think that you have mishandled anything?
- Do you get support at home to help handle the problem?
- Where does your support come from?
- How do you feel that you are coping?

Empathy

Indicate an understanding of the patient's distress and legitimise his or her feelings.

- That must be very difficult for you.
- That sounds really tough on you.

STEPS TO BRING ABOUT BEHAVIOUR CHANGE

Fabb and Fleming have introduced the model of change, which is fundamental to initiating therapy. The five steps are:

1. *Dissatisfaction*. There must be dissatisfaction with the present pattern of behaviour.
2. *Alternative*. There must be an acceptable alternative behaviour pattern available.
3. *Emotional commitment*. There must be an emotional commitment to the new pattern of behaviour over the old.
4. *Practice with feedback*. There must be practice of the new behaviour, with feedback, to establish the new pattern as an available behaviour.
5. *Habituation with support*. There must be installation of the new behaviour in the normal work/living situation *with support*.

All of these must be present for change to occur. Steps 4 and 5 are often neglected, with the result that change does not occur or is less successful.

MARITAL DISHARMONY

Family doctors often have to provide marital counselling for one or both partners. The problems may be resolved quite simply or be so complex that marital breakdown is inevitable despite optimal opportunities for counselling.

Opportunities for prevention, including anticipatory guidance about marital problems, do exist and the wise practitioner will offer appropriate advice and counselling. Examples include an accident to a child attributable to neglect by a parent, or similar situation in which that parent may be the focus of blame, leading to resentment and tension. The practitioner could intervene from the outset to alleviate possible feelings of guilt and anger in that marriage.

Some common causes of marital disharmony are:

- selfishness
- unrealistic expectations
- financial problems/meanness
- not listening to each other
- sickness (e.g. depression)
- drug or alcohol excess
- jealousy, especially in men
- fault finding
- ‘playing games’ with each other
- driving ambition
- immaturity
- poor communication

BASIC COUNSELLING OF COUPLES

The following text on basic counselling of couples,[9] which should be regarded as a patient education sheet, includes useful advice for couples:

> The two big secrets of marital success are *caring* and *responsibility*.

Some important facts

- Research has shown that we tend to choose partners who are similar to our parents and that we may take our childish and selfish attitudes into our marriage.
- The trouble spots listed above reflect this childishness; we often expect our partners to change and meet our needs.
- If we take proper care and responsibility, we can keep these problems to a minimum.
- Physical passion is not enough to hold a marriage together - ‘when it burns out, only ashes will be left ’.
- While a good sexual relationship is great, most experts agree that what goes on *out* of bed counts for more.
- When we do something wrong, it is most important that we feel forgiven by our partner.

Positive guidelines for success

1. *Know yourself*. The better you know yourself, the better you will know your mate. Learn about sex and reproduction.
2. *Share interests and goals*. Do not become too independent of each other. Develop mutual friends, interests and hobbies. Tell your partner ‘I love you’ regularly at the right moments.

3. *Continue courtship after marriage*. Spouses should continue to court and desire each other. Going out regularly for romantic evenings and giving unexpected gifts (such as flowers) are ways to help this love relationship. Engage in some high-energy fun activities, such as massaging and dancing.
4. *Make love, not war*. A good sexual relationship can take years to develop, so work at making it better. Explore the techniques of lovemaking without feeling shy or inhibited. This can be helped by books such as *The Joy of Sex* and videos on lovemaking. Good grooming and a clean body are important.
5. *Cherish your mate*. Be proud of each other, not competitive or ambitious at the other's expense. Talk kindly about your spouse to others - do not put him or her down.
6. *Prepare yourself for parenthood*. Plan your family wisely and learn about child-bearing and -rearing. Learn about family-planning methods and avoid the anxieties of an unplanned pregnancy. The best environment for a child is a happy marriage.
7. *Seek proper help when necessary*. If difficulties arise and are causing problems, seek help. Your general practitioner will be able to help. Stressrelated problems and depression in particular can be lethal in a marriage - they must be ‘nipped in the bud’.
8. *Do unto your mate as you would have your mate do unto you*. This gets back to the unconscious childhood needs. Be aware of each other's feelings and be sensitive to each other's needs. Any marriage based on this rule has an excellent chance of success.

The **BE** Attitudes (virtues to help achieve success)

BE honest	**BE** loyal
BE loving	**BE** desiring
BE patient	**BE** fun to live with
BE forgiving	**BE** one
BE generous	**BE** caring

Making lists - a practical task

Make lists for each other to compare and discuss.

- List qualities (desirable and undesirable) of your parents.
- List qualities of each other.
- List examples of behaviour each would like the other to change.
- List things you would like the other to do for you.

Put aside special quiet times each week to share these things.

Pitfalls[1]

The general practitioner who is too closely attached to one or more members of the family can easily become trapped in the role of the ‘rescuer’ or ‘saviour’ of those members. The best defence against this trap is to respect the family's autonomy and work with the family to achieve the goals the family sets for itself, thus avoiding three major pitfalls for the general practitioner in treating families:

1. assuming personal responsibility for changing the family
2. working alone, neglecting the assistance of the family
3. becoming a ‘rescuer’ or ‘saviour’

Other pitfalls

- conducting therapy in the absence of a significant member
- breaching confidentiality of individuals within the family
- failing to recognise the ‘ganging-up effect’
- taking sides
- failing to use available resources
- overrelating to your own experiences

Possible solutions to avoid pitfalls[1]

- Let the patients do the work.
- Share the burden with a colleague or other resources.
- Ensure that the goals for therapy are realistic.
- Point out that all family members have to work together and that therapy works best when there is openness on all sides.
- Identify any tendency to look for scapegoats within the family.
- Avoid trying to achieve quick solutions.
- Obtain clear-cut agreements on confidential matters and record this in the history.
- Keep an open mind and avoid forcing your own values on to the family.

REFERENCES

1. Bader E. Working with families. Aust Fam Physician, 1990; 19: 522 - 8.
2. Fabb W. *Handbook for Medical Students*. Hong Kong: Chinese University of Hong Kong, 1995: 31.
3. McGoldrick M, Gerson R. *Genograms in Family Assessment*. New York: WW Norton, 1985: chs 1 - 4.
4. Jackson L, Tisher M. Family therapy skills for general practitioners. Aust Fam Physician, 1996; 25: 1265 - 9.
5. Van Doorn H. *Common Problems Checklist for General Practice*. Melbourne: Royal Australian College of General Practitioners, 1989: 19.
6. McWhinney IR. *A Textbook of Family Medicine* (2nd edn). Oxford: Oxford University Press, 1997: 240 - 56.
7. Duvall EM. *Family Development* (5th edn). Philadelphia: Lippincott, 1977.
8. Stuart MR, Leiberman JA III. *The 15-Minute Hour: Applied Psychotherapy for the Primary Care Physician*. New York: Praeger, 1986.
9. Murtagh JE. *Patient Education* (3rd edn). Sydney: McGrawHill, 2000: 3.

CHAPTER 3. CONSULTING SKILLS

The consultation is a formalised interaction between doctor and patient in settings that may vary from a clearly defined task such as suturing a simple wound to the complexities of vague undifferentiated illness with profound psychological issues.

> The essential unit of medical practice is the occasion when in the intimacy of the consulting room the person who is ill or believes himself (or herself) to be ill, seeks the advice of a doctor whom he (she) trusts. This is the consultation and all else in the practice of medicine derives from it.
>
> SIR JAMES SPENCE 1960

CONSULTING SKILLS

The *objectives of the consultation* are to:

- determine the exact reason for the presentation
- achieve a good therapeutic outcome for the patient
- develop a strong doctor-patient relationship

THE SKILLS OF GENERAL PRACTICE

A successful outcome to the medical consultation depends on a whole array of skills required by the GP. Although interrelated, these skills, which can be collectively termed ‘consulting skills’, include clinical skills, diagnostic skills, management skills, communication skills, educative skills, therapeutic skills, manual skills and counselling skills.

Communication skills, which are fundamental to consulting skills, are the key to the effectiveness of the doctor as a professional, and expertise with these skills is fundamental to the doctor-patient relationship. Communication skills are essential in obtaining a good history and constitute one of the cornerstones of therapy.

A skilled interviewer will succeed in transmitting his or her findings to the patient so that they are clearly understood, are not unduly disturbing, and inspire trust and confidence in the physician.

MODELS OF THE CONSULTATION

Several models that formalise the general practice consultation can be very useful for developing an understanding of the process of the consultation. Two classic models are those by Pendleton and his colleagues,[1] and by Stott and Davis.[2] Pendleton and colleagues, in their landmark book *The Consultation: An Approach to Learning and Teaching*,[1] defined seven key tasks to the consultation, which serve as helpful guidelines:

1. To define the reason for the patient's attendance, including:
    - the nature and history of problems
    - their aetiology
    - the patient's ideas, concerns and expectations
    - the effect of the problems
2. To consider other issues:
    - continuing problems
    - risk factors
3. To choose, with the patient, an appropriate action for each problem
4. To achieve a shared understanding of the problems with the patient
5. To involve the patient in the management and encourage him or her to accept appropriate responsibility
6. To use time and resources efficiently and appropriately:

  - in the consultation
  - in the long term

7. To establish or maintain a relationship withthe patient that helps to achieve the other tasks.

The exceptional potential in each primary care consultation described by Stott and Davis,[2] which is presented in Table 3.1, also acts as an excellent aide-memoire to achieve maximal benefit from the consultation.

Table 3.1 The potential in each primary care consultation

A	**B**
Management of presenting problems	Modification of healthseeking behaviour
C	**D**
Management of continuing problems	Opportunistic health promotion

Source: Stott and Davis[2]

PHASES OF THE CONSULTATION

The consultation can be considered in three phases, as follows:

1. Establishment of rapport
2. Diagnostic phase
    - the history
    - the physical and mental examination
    - investigations
3. Management phase
    - explanation and education
    - prescribing medication
    - procedural - therapeutic or extended diagnostic
    - referral
    - follow-up

ESTABLISHING RAPPORT

Although rapport building occurs throughout all phases of the consultation, the initial encounter with the patient sets the foundation for the relationship during the consultation. It is good policy to walk into the waiting room and call the patient by the most appropriate name. Valuable clinical information can be gleaned by observing the patient's affect, movements and walking. It is also most appropriate to quickly familiarise oneself with the patient's notes from wellkept records, preferably before seeing the patient.

Practice tip

Remembering the patients preferred name and their basic past history is powerful rapport.

Rapport-establishing techniques include:

- Greet the patient with a friendly interested manner.
- Treat the patient with respect and courtesy.
- Greet the patient by his or her preferred name.

- Shake hands if appropriate.

Figure 3.1 The consultation: establishment of good rapport is the foundation to successful consulting skills

- Make the patient feel comfortable.
- Be 'unhurried' and relaxed.
- Be well briefed about prior consultations.
- Focus firmly on the patient.
- Listen carefully and appropriately.
- Make appropriate reassuring gestures.
- Start with: 'What would you like to tell me?' or 'How can I help you?'

THE HISTORY

The doctor has four basic tasks to perform during the history-taking phase of the consultation. These are to determine:

1. the patient's stated reason for attending
2. why the patient is attending today, or at this particular time in the course of this illness
3. a list of problems or supplementary symptoms
4. any other initially unspoken or hidden reason for attending (e.g. the fear of cancer)

The old medical cliché that 'a good history is the basis of the clinical examination' is as relevant as always. The art of history taking, which is based on good communication, is the most fundamental skill in general practice and requires a disciplined approach.

A very good approach is that used by Professor Rita Charon of Columbia University: 'I'm going to be your doctor, so I need to know a great deal about your body, health and life. Please tell me what you think I should know about yourself and your situation.'

Guidelines include:[3]

- Commence by eliciting the presenting complaint.
- Permit an uninterrupted history.
- Use appropriate language - keep the questions simple.
- Use specific questions to clarify the presenting complaint.
- Write notes or use the keyboard to record information but maintain as much eye contact as possible.

- Enquire about general symptoms, such as fatigue, weight changes, fever, headache, sleep and coping ability.
- Undertake a relevant systems review.
- An historical checklist includes past medical history, complete medication history, drug habits and sensitivities, family history, psychosocial history and preventive care history.
- Give feedback to the patient about your understanding of the problems and agenda, and correct any misconceptions.

Good questions

In order to determine any underlying agenda or significant psychosocial problems, it is very helpful to use analytical questions. Such questions and inviting statements could include:

- Why have you come to see me today?
- Do you have any particular concern about your health?
- That really interests me - tell me more - it seems important.
- Where would you put your real feelings between 0 and 100%?
- What is it that's really upsetting or bothering you?
- What do you really think deep down is the cause of your problem?
- Are you basically satisfied with your life?
- Is there anything that I haven't asked you and that you should tell me about?
- Tell me about things at home.
- Tell me about things at work.
- Do you experience any bullying?
- Are you afraid that something bad is going to happen to you?
- Is your relationship with any particular loved one/person causing you stress? (This may lead to information about sensitive issues such as domestic violence or sexual problems.)
- Is there anything in your life that you would like to change?
- I'm concerned about what you are not telling me.

Basic interviewing techniques

There are a number of basic interviewing techniques[4] that encourage communication. It is important to use the least controlling interview techniques before embarking on direct questioning.

Questions

When the patient is asked a question the doctor tends to take control of the interview, and so directs it along the lines of his or her own thinking or hypothesis generation. The problem is that if questions are used too early in the interview, the amount of desirable information is restricted and may disrupt the true priorities of the patient's concerns.

Open-ended questions and direct questions are very useful at appropriate times, while other questions are very restrictive. Examples, using pain as the ‘problem’, are:

- Open-ended question: ‘Tell me about the pain.’
- Direct question: ‘Where is the pain?’
- Closed question: ‘Is the pain severe?’
- Leading question: ‘The pain is severe?’
- Reflective question: ‘You want to know the cause of the pain?’

The open-ended question

The open-ended question is essential in initiating the interview. A question such as ‘What kind of troubles have you been having?’ says to the patient ‘I'm interested in anything you may feel is important enough for you to tell me’.

The open-ended question gives the patient an opportunity to take temporary control of the consultation and to outline problems and concerns.

Listening and silence

Silence is a means of encouraging communication. While the patient is communicating freely, the doctor's behaviour of choice is an interested, attentive and relaxed silence. An attentive facial expression and posture tells the patient non-verbally that he or she has an interested listener. Silence can also encourage communication but one has to be careful that the person does not feel uncomfortable with the process. There is one time when it is mandatory for the doctor to use silence - this is when the patient has stopped speaking from being overwhelmed with emotion.

Facilitation

Facilitation encourages communication by using manner, gesture or words that do not specify the kind of information that is being sought. It suggests that the doctor is interested and encourages the patient to continue. Silence and facilitation go hand in hand.

A common mode of facilitation is the nod of the head, conveying ‘I'm listening', ‘I understand what you are saying’ or ‘Go on’. A similar message is conveyed to the patient with an occasional ‘mmm-mmm’ or by postural shifts towards the patient or into a position of greater alertness. The doctor may also interject short words or phrases, such as ‘Yes’ or ‘I see’, without interrupting the flow of the patient's narrative, or following a pause saying, ‘Yes, I understand - please continue‘.

Confrontation

When one senses that the patient is not speaking freely or clearly, the technique of confrontation may be used whereby the interviewer describes to the patient something striking about his or her verbal or non-verbal behaviour. Examples are: ‘You look sad’, ‘You seem frightened’, ‘You sound angry’ or ‘I notice that you have been rubbing the back of your neck’. Confrontation has to be used with tact and skill, and should reflect sympathetic interest in the patient. It is appropriate too to confront a patient when his or her voice, posture, facial expression or bodily movements betray emotions, for example, ‘You seem tense’ or ‘You're trembling’.

Support and reassurance

The doctor's ability to be appropriately supportive and reassuring helps to create an atmosphere in which the patient is encouraged to communicate. Examples of supportive statements are ‘I understand’ or ‘That must be very upsetting’. Reassurance includes words or actions which tend to restore the patient's sense of well-being, worthiness or confidence.

Summarising

Summarising what the patient has said can keep the patient on track and check the accuracy of the information by providing the patient with the opportunity to revise any misunderstandings, for example, ‘If I've understood you correctly you have told me… ’.

Information from other sources

Sometimes it is important to obtain information from other sources, especially friends or relatives. Off -handed comments from others may be loaded with ‘cues’ and one should be listening intently.

Problem definition

Part of the diagnostic process is defining the patient's problem or problems. The more complex the presentation, the more necessary it is to have an orderly approach. It is clearly important to list the problems in a priority order. These problems may have been ‘offered’ by the patient, ‘observed’ by the doctor, ‘derived’ during the interview or ‘known’ from the past history. Problems can be conveniently considered as organic or physiological, and intrapersonal or social.[5]

Touching the patient

Sometimes a natural response is to touch the distressed patient as a reassuring gesture. It is best to adopt a caring-and-support gesture, such as offering a box of tissues to the weeping patient, but it may be quite acceptable in most patients to give a reassuring momentary touch somewhere between the shoulder and wrist on the arm nearest to you. Touching should be a natural gesture that is comfortable for both the doctor and patient. Touch elsewhere should generally be avoided.

THE PHYSICAL AND MENTAL EXAMINATION

If a diagnostic hypothesis based on the history is being tested, the examination may be confined to one system or to one anatomical region. However, other regions, systems or a general examination may be undertaken for medicolegal or preventive reasons. Patients tend to feel vulnerable during the physical examination, so their sensitivity and modesty have to be respected. Generally, the examination is conducted in relative silence, with the doctor instructing the patient what to do.

Patients need to be warned of possible discomfort or pain that may accompany certain examinations, of the reason for the examination, and of its immediate results, particularly if normal. Continued silence on the doctor's part is often interpreted by patients as being indicative of something serious or unusual being found. For the same reason the doctor's nonverbal behaviour is important.

MEDICOLEGAL GUIDELINES FOR EXAMINATIONS[6, 7]

The following guidelines have been recommended by the NSW Medical Board for consultations and physical examinations:

- Carefully explain the nature and purpose of the physical examination before you start. Take particular care with explanations before rectal, vaginal, breast and genital examination.
- Indicate when an examination may be uncomfortable and ask the patient to advise if you are causing pain.
- If a patient is required to disrobe, explain to what extent undressing is required and why.
- A patient's modesty should be preserved when undressing and dressing before and after a physical examination. Privacy screens, sheets and gowns should be provided as a matter of course. Clinic staff should not interrupt physical examinations.
- If the patient requests the presence of a chaperone or a friend, this should be respected.
- Do not lock the door of the consultation room. The setting should allow the patient confidence to terminate the consultation at any time if he or she is uncomfortable.

INVESTIGATIONS

It is often necessary to arrange for special tests to assist in the diagnostic process or to monitor the progress of certain illnesses or response to treatment. The informed consent of patients must be obtained. A collaborative decision for or against certain tests may be negotiated.

GPs have a responsibility (clinical and economic) to be very discerning and selective in the investigations that they choose. The questions that should be asked in decision making include:

- Is this investigation necessary?
- Will it change my management?

Richard Asher (1954) listed the questions a clinician should ask before requesting an investigation:[8]

- Why am I ordering this test?
- What am I going to look for in the result?

- If I find it, will it affect my diagnosis?
- How will this affect my management of the case?
- Will this ultimately benefit the patient?

In general, investigations should be performed only when the following criteria are satisfied:[8]

- The consequence of the result of the investigation could not be obtained by a cheaper, less intrusive method (e.g. taking a better history or using time).
- The risks of the investigation should relate to the value of the information likely to be gained.
- The result will directly assist in the diagnosis or have an effect on subsequent management.

The three strikes and you're out rule

A very useful rule is to bail out of the diagnosis and refer to a colleague if you have failed to make a diagnosis after three consultations.

MANAGEMENT PHASE OF THE CONSULTATION

The management phase of the consultation may immediately follow the information-gathering interview, or it may take place on review, after diagnostic tests or referral. It should be remembered that there are at least two people concerned in management: the doctor *and* the patient. Poor patient compliance with any proposed therapy can be a result of a poorly conducted management phase. It is necessary not only for the doctor to make statements concerning therapy and the reasons for the chosen therapy, but also for the information to be conveyed in a language appropriate to each patient's understanding.

Management includes immediate care, prevention and long-term care. Doctors generally tend to be authoritarian in their management proposals. Whole-person management, however, implies that the patient's views are listened to, explanations are offered where necessary by the doctor, and an educative approach is adopted to encourage the patient to actively participate in management and preventive behaviour, where possible.

No longer can the patient be expected to act as the passive receiver of advice or to submit without question to procedures, as in the past. There is evidence that the patient's compliance with management plans is improved if the patient has been involved in the decision making.

The objectives of the management phase of the consultation are summarised in Table 3.2.

Table 3.2 Objectives of the management phase of the consultation

• To make use of the doctor-patient relationship in therapy
• To involve the patient as far as possible in the management of his or her own problem
• To educate the patient about the illness
• To promote rational prescribing
• To achieve compliance in therapy
• To emphasise preventive opportunities
• To provide appropriate reassurance
• To encourage continuity of ongoing care

The sequence of the management interview

The following is a suggested *10-point plan* or sequence for conducting a management interview. These guidelines will not always need to be applied in their entirety, and may need to be staged over a number of consultations. The use of this sequence should ensure identification of all the patient's problems by the doctor (including fears, feelings and expectations); adequate patient understanding of their problems; an acceptable and appropriate treatment plan being defined for each problem; preventive opportunities being addressed; and the patient being satisfied with the consultation and being clear about follow-up arrangements.

The sequence is as follows.

1 Tell the patient the diagnosis

If a diagnosis is not possible, describe the problem as it relates to the presenting symptoms.

2 Establish the patient's knowledge of the diagnosis

This information provides a clear-cut baseline of information from which to launch the management phase of the consultation.

3 Establish the patient's attitude to the diagnosis and management

Unless this is done the doctor may already have begun to enter a conflicting relationship with the patient without knowing why and be unaware of underlying fears.

4 Educate the patient about diagnosis

- Correct any incorrect health beliefs recognised in point 2.
- Supplement the patient's existing knowledge to a level appropriate to the needs of the patient and the doctor.

Such illness education will be facilitated by the use of appropriate language, special charts and diagrams, models, investigation reports and other relevant aids (e.g. X-rays and ECGs).

5 Develop a management plan for the presenting problem

With precise instructions using three headings:

- Immediate: always included, even if no action is proposed
- Long term: for chronic, long-term or recurrent illnesses
- Preventive: sometimes specific measures apply - often patient education is the method required

The patient should be encouraged at this stage to participate in decision making regarding management and to make a commitment to the plans.

6 Explore other preventive opportunities

Common examples of preventive opportunities include immunisation, screening status (e.g. Pap smear), and advice about smoking and alcohol problems, and safe sex.

7 Reinforce the information

Emphasise information already given about the diagnosis and management by the use of other techniques, for example:

- Use the patient's own results (e.g. X-rays and ECGs).
- Encourage the patient to participate in the decision making and in accepting some degree of responsibility for his or her own management.

This process may be facilitated by having patients learn drug names and dosages, record body weight and urine tests, and monitor temperatures and blood pressure - when relevant.

8 Provide takeaway information

Examples of this important strategy include patient instruction leaflets and resource contacts.

9 Evaluate the consultation

When time permits the doctor should encourage feedback regarding the patient's reaction to the way the consultation has been conducted, and establish whether the objectives of both have been met and the patient is happy with the outcome.

10 Arrange follow-up

Clear instructions for review need to be made, preferably by providing appointments or stating that no further review is needed. Follow-up not only shows patient response to management, but also enables the reinforcement and clarification of preventive measures and information given. It also allows involvement of others, particularly family members where appropriate.

Closing the session

Good closure is an important strategy; ask ‘Has this visit helped you and your problems - is there anything more I can do?’

Discussion

The first three points should be attended to clearly right at the start of the management phase to avoid any conflict or misunderstanding between doctor and patient. The doctor can then commence education about the diagnosis, correcting any incorrect health beliefs recognised in point 2, and supplement the patient's existing knowledge to a level appropriate to the needs of the patient and doctor.

The management plan for the presenting problem must be clear and specific - if complicated, the steps involved should be written down for the patient to take home. Immediate management should always be included even if no action is proposed. Longterm management strategies are useful for chronic or recurrent illness to help the patient envisage what the future holds. Preventive measures may be a specific part of the long-term management or just require patient education. The patient encouraged to participate in decision making should indicate commitment to the plans, which reinforces the essential information and encourages the patient to accept some degree of responsibility for the management of his or her own illness.

The final, yet essential, step in the management consultation is precise follow-up. Follow-up strengthens an ongoing doctor - patient relationship where the doctor indicates genuine concern and interest in that patient's long-term health needs.

A patient management strategy

Brian McAvoy, writing in Fraser's excellent book *Clinical Method: A General Practice Approach* , presents a helpful aide-memoire in the approach to patient management:[8]

1. reassurance and/or explanation
2. advice
3. prescription
4. referral
5. investigation
6. observation (follow-up)
7. prevention

Prescriptions

It is worth emphasising that prescribing medicine is a relatively complex skill that requires considerable knowledge of the disease, patient's expectations, the drugs prescribed, their interactions and their adverse reactions. Part of this skill is making a decision not to prescribe medication when it is not absolutely necessary and then explaining the reasons and including non-pharmacological measures. This decision may be made in the context of a patient expecting a biochemical solution for his or her problem. As McAvoy points out, ‘If in doubt whether or not to give a drug - don't’.[8]

Referral

The decision to refer a patient is also another important skill. It is often difficult to find the right balance. Some practitioners refer excessively - others cling to the patients inappropriately. It is a mistake not to refer a patient with a serious chronic or life-threatening disease. Apart from consultants and hospitals, referral should be considered to GP colleagues or partners with special interests or expertise, support groups and other members of the primary health care team, such as physiotherapists, dietitians, chiropodists and social workers. At all times the GP should act as the focal reference point and maintain control of patient management.

The ‘gatekeeper’ role of the GP

The patient's general practitioner is the obvious and ideal linchpin in the health care system to take responsibility for the patient's health concerns and management. The patient may become confused with the system, especially if his or her problems are many and complex. The patient's GP has a vital role in acting as a ‘gatekeeper’ between primary and secondary care, and between paramedical services. He or she should always act in the patient's best interests and intervene, if necessary, to ensure that the patient is getting the best possible care.

The healing art of the doctor

The counselling process in general practice is based on the therapeutic effect of the doctor. This well-recorded feature is reinforced if the doctor has a certain professional charisma, and is caring and competent. We cannot underestimate the dependency of our patients on this healing factor, especially where significant psychic factors are involved.

MAKING THE PATIENT FEEL GOOD ABOUT THE CONSULTATION

Many patients do not particularly enjoy visiting the general practitioner and some, for a variety of reasons, visit as a last resort. It is important to make patients feel good about their visit. They should feel accepted, believed and important in the sense that they have reached out to a friend in whom they can trust and be confident with. This is where appropriate explanation, reassurance and basic counselling are so important in patient management. Even seemingly trivial complaints should be managed with understanding and explanation. Patient education handouts are appreciated. Attention to even simple problems with simple therapies, such as the palpation of a tender trigger spot on the back followed by the application of an ‘analgesic’ spray or rub with gentle massage is much more powerful than a disinterested, albeit reassuring, dismissal that the problem is ‘nothing much to worry about’.

KEY POINTS ON PATIENT MANAGEMENT

- It is difficult, perhaps impossible, to reassure patients in the absence of an appropriate physical examination and certain investigations.
- Reassurance must always be appropriate and therefore based on a substantial foundation: inappropriate reassurance damages the credibility of both the doctor and his or her profession.
- The two key characteristics of the doctor in establishing the basis of a successful outcome for the doctor-patient relationship are caring and responsibility.
- Vital factors included in this relationship are good communication, genuine interest and trust.

REFERENCES

1. Pendleton D et al. *The Consultation: An Approach to Learning and Teaching*. Oxford: Oxford University Press, 1984.
2. Stott N, Davis R. The exceptional potential in each primary care consultation. JR Coll Gen Pract, 1979; 29: 201 - 5.
3. Nyman KC. *Successful Consulting*. Melbourne: Royal Australian College of General Practitioners, 1996: 11 - 32.
4. Rose AT. Basic interview techniques. In Kidd M, Rose A. *An Introduction to Consulting Skills*. Community Medicine

Student Handbook. Melbourne: Monash University, 1991: 32 - 40.

5. Barrard J. The consultation in general practice: patients, problems and resources. Med J Aust, 1991: 154: 671 - 6.
6. Johnson P. Bedside manners: advice for doctors in training. UMP Journal, 1998; 2: 2.
7. Guidelines for Medico-Legal Consultation and Examinations. Sydney: NSW Medical Board, 1997.
8. Fraser RC. *Clinical Method: A General Practice Approach* (2nd edn.) Oxford: Butterworth-heinemann, 1992: 59 - 76.

CHAPTER 4. COMMUNICATION SKILLS

Most people have a furious itch to talk about themselves and are restrained only by the disinclination of others to listen. Reserve is an artificial quality that is developed in most of us as a result of innumerable rebuffs. The doctor is discreet. It is his business to listen and no details are too intimate for his ears.

W. SOMERSET MAUGHAM (1874 - 1965), *Summing Up*

COMMUNICATION SKILLS

Hippocrates wrote:

> In the art of medicine there are three factors - the disease, the patient and the doctor … It is not easy for the ordinary people to understand why they are ill or why they get better or worse, but if it is explained by someone else, it can seem quite a simple matter - if the doctor fails to make himself understood he may miss the truth of the illness.[1]

Francis Macnab, Doctor of Divinity and patient, wrote: 'The style of the doctor, the communication of the doctor and the person of the doctor at the level of primary contact and primary care can be crucial in a person's life'.[2]

Much of the art of general practice lies in the ability to communicate.

Research continues to focus the 'blame' for communication breakdown on the doctor, ignoring the role of the patient.[3]

COMMUNICATION

Communication can be defined as 'the successful passing of a message from one person to another'.

There are five basic elements in the communication process:

- the communicator
- the message
- the method of communicating
- the recipient
- the response

Important principles facilitating the communication process are:

- the rapport between the people involved
- the time factor, facilitated by devoting more time
- the message, which needs to be clear, correct, concise, unambiguous and in context
- the attitudes of both the communicator and the recipient

Communication in the consultation

The doctor requires appropriate communication skills for complete diagnosis (physical, emotional and social) and competent management. The majority of interaction between doctor and patient occurs in the traditional consultation. Table 4.1 shows where the communication pattern swings between being 'patient focused' and 'doctor focused'.[4]

Important positive doctor behaviour

At first contact:

- address the patient by his or her preferred name
- make the patient feel comfortable
- be 'unhurried' and relaxed
- focus firmly on the patient
- use open-ended questions where possible
- make appropriate reassuring gestures

Table 4.1 Phases of doctor-patient communication[4]

Phase 1 Patient focus	Phase 2 Doctor focus	Phase 3 Mutual focus

Introduction Present complaint Other medical history Family history Social history	Examination Investigation	Management discussion Follow-up Sign-off

Active listening

Listening is the single most important skill.[4] Listening is an active process, described by Egan as follows:

> One does not listen with just his ears: he listens with his eyes and with his sense of touch. He listens by becoming aware of the feelings and emotions that arise within himself because of his contact with others (that is, his own emotional resonance is another 'ear'), he listens with his mind, his heart, and his imagination. He listens to the words of others, but he also listens to the messages that are buried in the words or encoded in all the cues that surround the words. He listens to the voice, the demeanour, the vocabulary, and the gestures of the other. He listens to the context, verbal messages and linguistic pattern, and the bodily movements of others. He listens to the sounds, and to the silences.[5]

Listening includes four essential elements:

- checking facts
- checking feelings
- encouragement
- reflection

Listen with understanding, in a relaxed, attentive silence. Use reflective questions, such as:

- You seem very sad today.
- You seem upset about your husband.
- It seems you're having trouble coping.
- You seem to be telling me that …
- Your main concern seems to me to …

Attitudes

- caring
- empathy
- respect
- interest
- concern
- confidence
- competence
- responsibility
- trust
- sensitivity
- perceptiveness
- diligence

Communicating strategies

- Modify language.
- Avoid jargon.
- Provide clear explanations.
- Give clear treatment instructions.
- Evaluate the patient's understanding.
- Summarise and repeat.
- Avoid uncertainty.
- Avoid inappropriate reassurance.
- Arrange appropriate referral (if necessary).
- Ensure patient is satisfied.
- Obtain informed consent.

Follow-up

- Be available for telephone calls.
- Ensure patients obtain results of investigations ordered, including Pap smears.

- Ensure any promised follow-up is carried out.
- Phone the patient if you have any lingering concerns (this could be handled by the receptionist).
- Arrange referral if inadequate response to treatment.
- Act as an advocate if necessary (e.g. pressing for hospital admission).

Use of analogy

When communicating concepts/problems to people such as in counselling, it is worth being creative in the use of comprehensive common analigues; for example, in explaining cardiac arrhythmias refer to the similarity of a car motor misfiring because of electrical problems (distributor, spark plugs) and point out how the human engine (the heart) can be 'fixed' by medication or pacemakers.

DIFFICULTIES IN COMMUNICATION

The Victorian Medical Board lists poor communication as the most important factor causing complaints from patients and relatives against doctors.[6]

Effective communication depends on four interrelated factors concerning the message - the doctor (the sender), the patient (the recipient), the message itself and the environment in which the message is sent (Figure 4.1).[7]



Figure 4.1 The four key factors affecting communication

Environment

The physical environment is important (Table 4.2). The appearance, size and layout of consulting rooms, waiting rooms and patients' rooms will affect communication, sometimes adversely, especially if privacy is threatened by, say, leaving the consulting room door open. The doctor can create an obstacle simply by a physical 'barrier', for example, a large desk distancing the doctor from the patient (Figure 4.2).

The hospital environment will encourage the 'sick' role and generally is not conducive to good communication because of a low level of privacy.

A busy practice affected by time constraints on doctor or patient will influence communications seriously. A doctor in Wales has a notice in his waiting room: 'If the doctor is a long time with a patient don't get mad: it might be you'.

The message

The nature and content of the message may be uncomfortable for the doctor or the patient or both (Table 4.3). This applies to emotionally charged, complex or subtle content, such as sexual problems, malignant disease, drug abuse, bereavement, malingering and psychiatric disorders.

The patient may find the message difficult to comprehend because of inappropriate delivery or explanation by the doctor. Failure to use good follow-up strategies, including appointment times and appropriate patient education material, will aggravate communication breakdown. Language difficulties can distort the message and generate frustration in both parties. Good interpreters often help.

Table 4.2 Summary of environmental factors that can adversely influence communication

Waiting room	Poor physical layout Length of waiting time
Time pressure	'Traffic' level
	? busy
	? noisy
	? sense of urgency
Physical factors	Desk - barriers
	Layout inappropriate
	Poor record system
	Substandard examination couch
Privacy	Dressing/undressing Sound
	Interruptions - telephone



Figure 4.2 The physical barrier

The doctor may also fail to appreciate that certain symptoms, such as chronic pain or the presence of a lump, mean 'cancer' to the patient. Failure to reassure the patient (where appropriate) distracts the patient.

Doctor - patient interaction

There are several general characteristics that affect communication between doctor and patient. These include:

- poor past relationships and experiences leading to unresolved interpersonal conflict (e.g. an incorrect diagnosis or poor treatment outcome and indifferent compliance in following treatment or paying accounts)
- personal differences, openly expressed, which may create subtle barriers, including differences in age, sex, religion, culture, social status and doctor/patient roles (occasionally influenced by political factors)
- the communication skills of doctor and patient, both as the sender and receiver of messages
- the personal honesty and integrity of both parties in dealing with difficult messages
- psychosocial problems that will establish barriers (e.g. psychiatric illness or speech impediments)
- familiarity between patient and doctor (e.g. friends or relatives)

Table 4.3 Negative communication related to the message

Language difficulties
Complex problems
Emotional problems
Uncertainty and doubt
Examples:
• sexuality, such as incest, STDs
• malignancy
• multiple complaints: 'the shopping list'
• infertility
• unwanted pregnancy
• abortion

The doctor

Although we believe that most doctors satisfactorily meet professional standards, there are times when the communication factor is adversely affected by inbuilt negative forces, including chronic tiredness, stress, domestic problems and poor health (Table 4.4).

Furthermore, there are many strategies, roles, 'games' or 'hobbyhorses' that some of us appear to rely on, especially when confronted with difficult or threatening circumstances, such as the management of the terminally ill.

Dare we recognise in ourselves some of the following unkind caricatures of doctors (i.e. personality types who may generate unfavourable communication)?[8, 9]

Table 4.4 The doctor's personal factors that influence communication

Age	Elderly, young
Sex	Opposite
Senses	Deafness, speech idiosyncrasy
Handicap	Relative immobility
Competence	Health understanding Professional training Social awareness Empathy
Attitudes	Bias - patient attending other doctors or alternative practitioners
Communication style differences	Religion, sexual practices Social class Ethnic group Political group

Dress

Eccentricities

Familiarity



Figure 4.3 Dr Al Oof

Dr Al Oof (Figure 4.3). The prima donna doctor (not necessarily a surgeon); aloof; omnipotent; dark suit with matching Mercedes; club tie or bow tie; feared by medical students; partial to Scotch; pronounces certain cures; powerfully dispels doubts; no faith in the healing process before surgery but unshakeable faith after surgery; unavailable in the patient's decline.

Dr N. Zyme (Figure 4.4). The scientific doctor; machine-like; cool; assured; obsessive; drives an Italian car; orders a new test and drug at every visit; conversant with the cellular biochemistry of the disease process but ignorant of its host.

Dr G. Rumble. The gruff doctor; grunts in mono-syllables; brilliant but appears tough and unapproachable; actually quite shy, soft and kind behind the facade; drives a Ford.



Figure 4.4 Dr N. Zyme

Figure 4.5 Dr I. Knowall

Dr No Komento. The secretive doctor; strong and silent, or is he weak and silent, threatened? In another world! Drives a BMW; a computer buff.

Dr I. Knowall (Figure 4.5). Glib; assured; garrulous; drives latest red sports car; drapes stethoscope around neck; accepts invitations to lecture on all subjects; rarely available on the phone; keeps patients waiting for hours.

Dr S. Winger. Modern, swinging and trendy; superficial; on first name terms with patients; drives beaten-up Renault held together by political stickers; works only 35 hours a week; cavalier; undiplomatically blunt.

Dr X. Cytabull. Fanatic; madly enthusiastic about rarities; overreacts to physical abnormalities; compulsive writer to medical editors; refers patients ad nauseam; drives yellow Porsche.

Dr Genghis M. Pyre. Longs for a mega-practice, assistants (not partners) and a pathology service; addicted to conferences and cocktail parties; also yearns for a Daimler, a halo and New Year's honours.

Dr Buzz Bee. Ever busy; flits from one consulting room to another; frequent telephone user during consultation; creates a sense of urgency everywhere; charming to patients but intimidates them; overservices; holds pilot's licence; drives Landcruiser when licence not suspended.

Dr Go Along Cassidy. Feels comfortable when he is giving patients what they ask for; has a ‘conveyor belt’ type practice; rarely leaves his chair and doesn’t examine his patients; drives a Mustang.

Dr I. Kling. Protective and possessive; hangs on to patients; refers only under pressure; overconfident; likes to be liked; indifferent medical record system; compulsive drug prescriber; still drives 1982 Volvo.

Dr Nat Ure. Strong on ‘alternatives’; pleasant chap; keen on Blackmore's publications and remedies; health shop (run by spouse) next door for fibre, sprouts and vitamin pills; attracts an attractive clientele; mutters audibly while writing the rare script; into massage, yoga and transcendental meditation; wears a knitted tie; rides a bicycle.

Dr FiMayle. The invisible doctor; juggles patients, children and the PC with one hand while cooking dinner with the other; earns less, pays more; shuns cocktail parties in preference to continuing medical education (CME) meetings with child care; prefers to be really achieving something through her division rather than waiting for the power boys to do it for her; finds continuing care and collegiate relationships difficult; drives whatever will take her reliably from A to B many times a day.[10] (reprinted with permission)

Dr Amy Preschool. Ever late to start surgery; smartly dressed in three-year-old fashions (bought before the baby was born) bearing tell-tale infant food stains; babysitting problems; caring of mums and kids; constant attender of paediatric continuing education programs to find the cause of her child's continual diarrhoea; drives a late model Japanese-built station wagon with a recommended car seat in the back.[10]

Dr Family Practice. The conservative doctor; married to a university sweetheart; practising from home for many years with her husband; prescribes mist magnesium trisilicate for peptic ulceration, Relaxa-Tabs for panic states and the ‘Red tonic’ for depression; children grown up and at university; both left at home with the ageing parents, two dogs and two cars; she drives the Mini Minor.[11]

Dr Susie Nirvana. Pap smear queen; always working, never in the same place twice; takes her entourage of similar searching patients with her. Drives someone else's car.[11]

Dr Magoo. Always in court; popular with solicitors; never examines patients; great conversationalist; rarely looks; misses obvious signs; can’t afford a car.

Dr Ann Osmia. Socialite doctor with ‘special’ clientele; senseless to sensory signs, such as abnormal smells, sights and sounds; has difficulty with diagnosing alcohol abuse, gastrointestinal disorders and diabetes; harbours secret bad experiences with neurology tutor; drives a Saab cabriolet with matching poodle.

Dr Otto Sclerosis. The clinic's unpopular doctor; doesn’t listen; doesn’t hear; preoccupied; has gambling and drinking problem; drives Range Rover when driver's licence not suspended.

These caricatures mirror something of ourselves, so that, it is to be hoped, we can understand our own attitudes and behaviour. The stereotypes portrayed may well adversely affect our relationship with our patients and colleagues.

The patient

Do we recognise, with significant emotion, these patients in our practice?

- ‘Smith speaking - I insist on speaking to you directly and not to the “iron curtain” out front.’
- ‘Doctor, I’ve lost my script again - be a good fellow and … ’
- ‘Those pills you prescribed yesterday are doing nothing for me.’
- ‘Doctor, you’re the only one who can help me.’

Yes, doctors are human and can harbour hostility towards the difficult patient, including the demanding patient, the seductive patient, the ‘compo’ patient, the difficult ‘ethnic’ patient, the hypochondriac, the bad debtor or the manipulative patient.

Some patients appear to have the irrepressible ability to create conflict, so often heralded by an upset receptionist, thus setting the scene for a potentially difficult consultation (Table 4.5).

However, doctors have a professional responsibility to transcend interpersonal conflict and facilitate productive communication by establishing a caring and responsible relationship, even with ‘difficult’ patients. Not surprisingly, such patients can also be found to be warm and pleasantly human beneath their ‘shoulder-chip’ facade and so be helped immeasurably by an empathetic doctor.

It is important to bear in mind that medical communication often occurs in an emotional environment because ‘disease’ has important emotional connotations for patients and their relatives and friends. Inappropriate communication and management can generate hostility.

Keep in mind also the variation in patients’ cultural and educational background and tailor your communication to the person in front of you. Consider an interpreter.

The doctor-become-patient in the hands of his colleagues learns fast, but possibly too late. Illness plus defective communication can bring confusion, anxiety and pain; suspicion and confinement add new dimensions to suffering.

Sooner or later we come to see ourselves as persons, both as doctor and as patient (‘wearing his moccasins’). The patient in us longs for the ideal doctor who is truly professional with sound knowledge and sane judgment, who is available, unhurried, caring and responsible.

Table 4.5 Patient characteristics that can influence communication

Age	Adolescent, elderly
Sex	Opposite
Senses	Deaf, blind, speech impairment
Handicapped	Speech disorders, visual impairment
Illness	Acutely ill/injured
Psychological	
Attitudes	Aggressive, hostile Demanding Aggrieved (e.g. fees, mistakes) Perception of doctor's authority
Anxiety/depression	
Dementia	
Fears and phobias (e.g. AIDS)	
Health understanding	
Hysteria	
Hypochondriasis	
Personality disorders	
Sensitive issues (e.g. sexuality, bereavement, malignancy)	
Social	
Social class	
Ethnic group	
Education	
Dress	
Political group	
Familiarity	

‘ROAD BLOCKS’ TO GOOD COMMUNICATION

In the communication process it is important to eliminate blocking approaches such as highlighted in the ‘dirty dozen’.

The dirty dozen[12]

The following list, referred to as ‘the dirty dozen’, represents well-identified blocking techniques to effective patient communication.

Judging

1. Criticising: ‘You didn’t bother to follow up that test’
2. Name-calling: ‘You are becoming a worrisome drug addict’
3. Diagnosing: ‘I can read you like a book’
4. Praising evaluative: ‘You’re a good patient - I know you can manage this … ’

Sending solutions

1. Ordering: ‘You must stop smoking’
2. Threatening: ‘If you don’t change, you will be dead meat in 12 months’
3. Moralising: ‘I cannot condone that sort of behaviour - it's wrong and you will pay the penalty’
4. Excessive/inappropriate questioning
5. Advising/patronising: ‘When you’re in Thailand, be perfectly good’

Avoiding the other's concerns

1. Diverting/changing the subject: ‘What did you think of the election result?’

2. Logical argument: ‘This wouldn’t have happened if you … ’
3. Reassuring: ‘What are you worrying about? Hundreds of people have to face up to that … ’

The seven sins of medicine

It is interesting to reflect on the editorial comments of Sir Richard Asher in *The Lancet* in 1949 outlining the ‘Seven Sins of Medicine’:[13]

- sloth
- spanophilia (love of the rare)
- obscurity
- bad manners
- over-specialisation
- cruelty
- common stupidity

NON-VERBAL COMMUNICATION

Non-verbal communication or body language is a most important feature of the communication process. Birdwhistle has shown that more human communication takes place by the use of gestures, postures, position and distances (non-verbal communication or *body language*) than by any other method.[14] Albert Mehrabian showed that non-verbal cues comprise the majority of the impact of any communicated message (Table 4.6)[15]

Recognition of non-verbal cues in our communication is important, especially in a doctor-patient relationship. The ability to recognise non-verbal cues improves communication, rapport and understanding of the patient's fears and concerns. Recognising body language can allow doctors to modify their behaviour, thus promoting optimum communication.[16]

Table 4.6 Impact of the message

	%
Words alone	7
Tone of voice	38
Non-verbal communication	55

Interpreting body language[14, 16]

The interpretation of body language, which differs between cultures, is a special study in its own right but there are certain cues and gestures that can be readily interpreted. Examples illustrated include the depressed patient (Figure 4.6), barrier-type signals often used as a defensive mechanism to provide comfort or indicate a negative attitude (Figure 4.7) and a readiness gesture indicating a desire to terminate the communication (Figure 4.8).

Having noted the non-verbal communication the doctor must then deal with it. This may require confrontation, that is, diplomatically bringing these cues to the patient's attention and exploring the associated feeling further.

It is not difficult to appreciate the importance of body language in the doctor-patient relationship. A hunch or gut feeling can be better understood, reinforced or corrected by skilled observation and interpretation of body language. A doctor can recognise a patient's non-verbal cues and explore the issues raised. By improving one's skills, and modifying one's behaviour (and consulting room configuration) the doctor can encourage communication and a better understanding of the patient.

The skill to interpret non-verbal cues can be achieved by conscious observation of people's interaction, including our own. A technique suggested by Pease[14] is to watch television without sound for 15 minutes each day and check your interpretation each 5 minutes. By the end of three weeks, he suggests, you will have become a more skilled body language observer.



Figure 4.6 Posture of a depressed person - head down, slumped, inanimate; position of desk and people correct



Figure 4.7 Body language - barrier signals: **(a)** arms folded; **(b)** legs crossed; **(c)**‘ankle lock’ pose



Figure 4.8 Body language - ‘readiness to go’ gestures

RAPPORT-BUILDING TECHNIQUES

A person can develop a rapport with another by mimicking their body language, speech, posture, pace and other characteristics. This method is a type of neuro-linguistic programming based on the work of Bandler and Grinder.[17] Such techniques can be used to help the doctor communicate better with the patient and also to improve a patient's attitude by changing the patient's body language position. It will be difficult for the patient to maintain a negative attitude if the body language position is not congruent.[16]

Mirroring

Mirroring is a useful technique whereby the limb positions and body angles of the person you are talking to can be copied. A mirror image is formed of their position so that when they look at you they see themselves as in a mirror. It is not necessary to copy uncomfortable gestures or unusual limb positions, such as hands behind the head. A partial mirror is often sufficient.

Pacing

People exhibit a certain rhythm or pace that can be revealed through their breathing, talking, and movements of the head, hands or feet. If you can copy the pace of another person, it will establish a sense of oneness or rapport with them. Once this pace is established you can change their pace by changing yours. This is called *leading*.

Practice tips

- A fundamental prerequisite for effective communication is listening; this includes not only hearing the words but also understanding their meaning, in addition to being sensitive to the feelings accompanying the words.[19]
- Undertake the strategies of paraphrasing and summarising during the consultation to emphasise that listening is occurring and to provide a basis for defining the problems.
- Associated with listening is the observation of the non-verbal language, which may in many instances be the most significant part of the communication process.
- Good communication between doctors and patients decreases the chance of dissatisfaction with professional services, even with failed therapy, and the likelihood of litigation.

Vocal copying

Vocal copying is a rapid and effective way to develop rapport with people. It involves copying intonation, pitch, volume, pace, rhythm, breathing and length of the sentence before pausing.

Engaging in these strategies will bring you into such close rapport that you can intuitively pick up all kinds of things about people that were not obvious beforehand. It may also have the unfortunate effect of making you feel that you are 'drowning' in their problems. If you feel overwhelmed, then break the rapport and diplomatically go into a leading phase.[18]

Key features of good communication

- Active listening
- Appropriate address (i.e. name of patient)
- Empathy
- Open-ended questions
- Summarising
- Checking understanding and feelings
- Good closure

REFERENCES

1. Elliott-Binns E. *Medicine: The Forgotten Art*. Tunbridge Wells, Kent: Pitman Books, 1978: 35.
2. Macnab F. Changing levels of susceptibility in sickness and in health. Aust Fam Physician, 1986; 15: 1370.
3. Dunn S, Allard B. Communication breakdown. Australian Doctor, 30 May 2003; I - VIII.
4. Mansfield F. Basic communicating skills. Aust Fam Physician, 1987; 16: 216 - 22.
5. Kidd M, Rose A. *An Introduction to Consulting Skills*. Department of Community Medicine, Student Handbook. Melbourne: Monash University, 1991: 15.
6. Medical Board of Victoria. Third Annual Report, 1982/3. Melbourne: FD Atkinson, Government Printers, 1983: 12.
7. Carson N, Findlay D. *Communication Skills*. Student Handbook. Melbourne: Monash University, Department of Community Medicine, 1986: 31.
8. Elliott CE. 'How am I doing?' Med J Aust, 1979; 2: 644 - 5.
9. Murtagh JE, Elliott CE. Barriers to communication. Aust Fam Physician, 1987; 16: 223 - 6.
10. Ivory K. The invisible doctor. Med Observer, 1996; 12 April: 19.
11. Saltman D. Rectifying a sexual bias (Letter to the editor). Aust Fam Physician, 1987; 16: 545.
12. Baker L, Ghanen A, Morton A, Mendel S. *Communication Skills*. Proceedings: Centwest/R*hed West* Training Program Workshop 2003: 2 - 3.
13. Asher R 'The Seven Sins of Medicine' The Lancet, 1949; 27 August.
14. Pease A. *Body Language*. London: Camel Publishing, 1985: 1 - 63.
15. Mehrabian A. *Silent Messages*. Belmont, CA: Wadsworth, 1971.
16. Findlay D. Body language. Aust Fam Physician, 1987; 16: 229.
17. Bandler R, Grinder J. *Re-framing: Neuro-linguistic Programming and the Transformation of Meaning*. Moab, UT: Real People Press, 1982: 1 - 203.
18. Oldham J. Neuro-linguistic programming. Aust Fam Physician, 1987; 16: 237 - 40.
19. Lloyd M, Bor R. *Communication Skills for Medicine*. London: Churchill Livingstone, 1996: 17 - 25.

CHAPTER 5. COUNSELLING SKILLS

> The doctor should have a kind disposition, great patience, self-possession, meticulous freedom from prejudice, an understanding of human nature resulting from an abundant knowledge of the world, adroitness in conversation and a special love of his calling.
>
> G. GRIESINGER 1840

COUNSELLING SKILLS

The Macquarie Dictionary says that counselling is 'giving advice': that it is 'opinion or instruction given in directing the judgment or conduct of another'. In the clinical context counselling can be defined as 'the therapeutic process of helping a patient to explore the nature of his or her problem in such a way that he or she determines his or her decisions about what to do, without direct advice or reassurance from the counsellor'.

The counselling process in general practice is based on the therapeutic effect of the doctor. There is an enormous and ever-increasing need for people in the community to have many of their emotional and social problems addressed by the health profession. Modern medicine has acquired a much more scientific face over recent years at the expense of its once respected humanistic one. Medicine is primarily a humanitarian pursuit, not an economic or scientific one, and uses science as a tool. Many feel that medicine is losing sight of this, at the considerable expense of its standing in the community.[1]

The public perceives that GPs can and do counsel people because more people go to their GP for counselling than to any other group of health workers, including psychologists, psychiatrists, social workers, marriage guidance counsellors and clergy.[1] People do not generally tell the doctor or even realise that counselling is exactly what led them to come to the doctor in the first place. The GP is, therefore, ideally placed in the community to make the most significant contribution to fill the community's needs in this area.

THE GP AS AN EFFECTIVE COUNSELLOR

GPs can be effective counsellors for the following reasons:[2]

- They have the opportunity to observe and understand the patients and their environment
- GPs are ideally placed to treat the whole patient
- Their generalist skills and holistic approach permit them to have a broad grasp of the patient's problems and a multifaceted approach to treatment
- They can provide treatment in comfortable and familiar surroundings, including the doctor's rooms and the patient's home
- They are skilled at working as a member of a professional team and directing patients to more expert members of the team as necessary
- They can readily organise 'contracts' with the patient
- They have an intimate knowledge of the family and the family dynamics
- They fit comfortably into continuing patient care with appropriate follow-up treatment programs

To be an effective counsellor the GP must first prepare for this role. Following a commitment to its importance the GP can acquire the knowledge and skills for basic counselling by reading, by attending workshops and by discussing cases with colleagues who are skilled in counselling.[2] Well-developed interviewing skills are essential, as is self-discipline to appreciate one's strengths and limitations.

FEATURES OF COUNSELLING

Doctors can respond to patients' problems and distress by a spectrum of behaviours from doctorcentred directive behaviour or advice at one end, to patient-centred non-directive behaviour at the other. In handling psychosocial



problems, advice giving is at one end of the spectrum and psychotherapy at the other.

Counselling, as an activity in general practice, can be represented by a moving point between these two extremes.[1]

Counselling can be seen as having the following features:[1]

- It is a clear-cut treatment option, like a course of antibiotics
- It is a co-operative problem-solving process
- It is an educational venture where patients learn new information and new activities
- It is a developmental process for patients
- It is a change process - often moving a patient from a ‘stuck state’
- It is a goal-directed activity
- It is a process of energising patients and lifting their morale
- It is a sensitive response to problems within a caring relationship

A PROBLEM-SOLVING APPROACH

Defining the problem (what the matter really is) is the most important step in the process of patient care. The following outline is one approach to counselling that is applicable to a general practice context.[1]

1. Listen to the problem of first presentation: this involves listening not only to issues, events and experiences, but also to the patient's feelings and distress. The emphasis here is more on the communication skills of facilitation, silence, clarification, reflection, paraphrasing, confrontation and summary, than on questioning. In many cases this phase of the counselling constitutes the major part of the therapy; in grief or bereavement counselling, where the doctor supports the patient through a natural but distressing process.
2. Define a problem, if possible in behavioural terms:

   > Beneath the feeling is the experience, beneath the experience is the event, the event is related to a problem.[3]

3. Establish a contract for counselling, with an agreed number of visits initially (e.g. weekly half hour or hour appointments for 4 to 6 weeks).
4. Define short-term and long-term goals for action.
5. Decide on one option - ’experimental action’.
6. Build an action program with the patient - negotiate ‘homework’ for the patient between visits.
7. Evaluate progress.
8. Continue action or select another option.
9. Evaluate progress.
10. Terminate or refer.

COUNSELLING MODELS

The PLISSIT model

The PLISSIT model, developed by Annon (1974)[4] as an aid in therapy for sexual problems, is a very useful model for problems presented as feelings where there is limited scope for intervention by the therapist.

The mnemonic PLISSIT stands for the following:

- P is for permission giving
- LI is for limited information
- SS is for specific suggestion

- IT is for intensive therapy

Annon emphasises that every primary care practitioner should be competent to offer ‘permission giving’ and ‘limited information’.

The Colagiuri and Craig model

The medical counselling model was developed by Colagiuri and Craig (Figure 5.1 on p. 34)[5] as a useful tool for teaching contraceptive, abortion and sterilisation counselling. It can be applied in most situations as it empowers the patients to make their own decisions through facilitation as opposed to the directive and advisory learning model.


Information giving
+
Facilitation
Decision making by client/s
Making it easy
Accurate
Unbiased
Appropriate
Comprehensive
Active listening
Questioning—
• open-ended
• circular
• reflective
Drawing comparisons
Challenging attitudes
Clarifying needs and desires
Self-determination
Independent
• In control
• Empowered even if wrong decision

Figure 5.1 Medical counselling model. After Colagiuri and Craig, reprinted with permission.

THE VALUE OF PATIENT-CENTRED COUNSELLING

There is evidence that the use of non-directive counselling techniques leads to more accurate diagnosis and therefore to more appropriate management and an improved outcome.[6]

Jerome Frank wrote in 1967: ‘The field of counselling and psychotherapy has for years presented the puzzling spectacle of unabating enthusiasm for forms of treatment where effectiveness could not be objectively demonstrated’.[7] Traux and Carkhuff [8] measured important aspects of the psychotherapeutic relationship and demonstrated what had long been recognised: the outcome was enhanced if practitioners had such qualities as accurate and sensitive awareness of the patient's feelings, deep concern for the patient's welfare (without attempting to dominate) and openness about their own reactions.

The essential feature of the patient-centred approach is that the counsellor is more like a facilitator; that is, by the asking of well-directed questions it is hoped that patients can realise their own solutions for their problems.[1] This encourages patients to attain understanding and personal growth themselves rather than just put their personal affairs in the hands of someone else. This does not mean to say that the facilitator is passive in the process of assessing the relative merit of various solutions produced by the patient. The doctor-centred approach is most applicable for patients who are so confused or distraught that their ability to reflect usefully is temporarily or permanently inaccessible. Here, taking a more active and authoritarian role may be just what is required. It is therefore important to be flexible and move between the two ends of the spectrum as needed.

BASICS OF COUNSELLING OR PSYCHOTHERAPY

- Listening and empathy are the beginning of counselling
- Good communication is the basis of counselling
- The therapist must really care about the patient
- Always be aware of the family context
- It is important for therapists to handle and monitor their own feelings and emotions
- Maintain eye contact
- The therapist must tolerate and be comfortable with what the patient says

- Confidentiality is essential
- Counselling is easier if there is a good rapport with the patient, especially if a long-standing relationship exists
- Counselling is difficult if a social relationship is present
- Don't say to the patient ‘I'm counselling you’ or ‘I'm giving you psychotherapy’ - make it a natural communication process
- The therapist must be versatile and adapt a counselling style to the clinical occasion
- Characteristics of the effective counsellor have been demonstrated to be genuineness, non-possessive warmth for the patient, and accurate and empathic understanding

Some useful interviewing skills used in counselling are summarised in Table 5.1 overleaf.

Table 5.1 Interview skills used in counselling

• Use reflective statements.
• Use silence.
• Allow expressions of emotion.
• Offer supportive comments.
• Paraphrase and summarise.
• Allow patients to correct your interpretations of their feelings.
• Observe lack of congruence.
• Try to understand what the patient is feeling:
- anger
- hostility
- fear
- manipulation
- seduction
- insecurity
• Make intelligent guesses to prompt patient to continue.
• Don't reassure too soon.

Counselling strategies[3, 6]

- The therapy should be patient-centred
- Use gentle, clever, probing questions
- Facilitate the discussion to draw out relevant areas
- It is important to be non-judgmental
- Counsel through intuition and base it on commonsense
- Do not tell the patient what to do
- Do not try to rush patients into achieving a happy ending
- Provide guidance to allow the patient to gain insight
- Wherever possible, make therapy nonauthoritarian and non-directional
- Use appropriate ‘gentle’ confrontation to allow self-examination
- Help patients to explore their own situation and express emotions, such as anxiety, guilt, fear, anger, hope, sadness, self-hate, hostility to others and hurt feelings
- Explore possible feelings of insecurity and allow free expression of such feelings
- Explore patients' belief systems and consider and respect their spiritual aspirations and conflicts
- Ask key searching questions, such as:

  - ‘What would be different in your life if you were well?’
  - ‘Who are you mad at?’
  - ‘If I understand you correctly you are telling me that … ’
  - ‘You seem to be telling me that … ’
  - ‘Correct me if I'm on the wrong track, but you are saying that … ’
  - ‘What do you think deep down is the cause of your problem?’
  - ‘What does your illness do to you?’
  - ‘Do you really worry about any things in particular?’
  - ‘How do you think your problem should be treated?’
  - ‘If you could change anything in your life what would it be?’

Avoid:

- telling patients what they must do/offering solutions
- giving advice based on your own personal experiences and beliefs
- bringing up problems that the patient does not produce voluntarily

What counselling is not

- giving information
- giving advice
- being judgmental
- imposing one's own values, behaviour and practices
- the same as interviewing
- handing out patient education material

Cautions[1]

- Individual doctors cannot be useful to all patients, so be selective
- We cannot solve patients' problems for them
- Patients' problems belong to them and not to their counsellors
- Patients often have to change by only an inch in order to move a mile
- If a counselling relationship is no longer productive, then terminate and refer
- Most patients in primary care need information, support and a lift in morale, not long-term psychotherapy

PATIENTS UNLIKELY TO BENEFIT

The following groups of patients are not likely to benefit from counselling therapy[1] (i.e. relative contraindications):

- psychotic patients
- patients who have had an unrewarding experience with psychiatrists and other psychotherapists
- people who are antagonistic to the notion of a psychosocial diagnosis, subsequently found to be organic
- patients with little awareness or language to express emotional difficulty
- patients who do not believe doctors can treat psychosocial problems
- patients who are dependent on contact with the doctor and are willing to do almost anything to maintain the relationship
- patients with a vested interest in remaining unwell who are therefore resistant to change (e.g. patients with work-related disabilities awaiting legal settlement)
- patients with chronic psychosomatic tendencies who are willing to do almost anything to maintain the relationship
- those in an intractable life situation who are unable or unwilling to change
- patients who are unwilling to examine and work on painful or uncomfortable areas of their life

COGNITIVE BEHAVIOUR THERAPY

Cognitions are thoughts, beliefs or perceptions. Cognitive behaviour therapy (CBT) involves the process of knowing or identifying, understanding or having insight into these thought processes. The therapy then aims to change behaviour. CBT is based on an underlying theoretical rationale that an individual's behaviour, attitude and thinking is based on the way in which he or she perceives the world. It can be applied in any area of medical practice as a form of psychotherapy and is suitable in general practice for the treatment of depression, insomnia, eating disorders, delusions and hallucinations in psychotic disorders and anxiety in all forms, especially social anxiety disorder and phobias - in all of which CBT has proved to be better than placebo.[9]

It is a relatively brief, active, directive and practical form of therapy. However, not all therapists or patients are suited to CBT.

The basic processes of CBT are to:

- educate the patient
- teach basic skills for symptom control, relaxation and breathing control (especially for hyperventilation)
- identify, challenge and change maladaptive thoughts, feelings, perceptions and behaviour

Some basic principles and objectives of CBT are:

- aim to bring about a desired change in patients' lives
- assess, monitor and attempt to modify thoughts and behaviour
- reinforce positive behaviour, discourage negative behaviour
- educate about any misconceptions about a patient's illness
- encourage the patient to be an active participant (not a passive recipient)
- get patients to establish a problem list and hierarchy of problems
- aim for more realistic thinking and more adaptive responses

SPECIFIC AREAS OF COUNSELLING

Opportunities for basic counselling by the general practitioner are ubiquitous in medical practice. Complex problems require referral but, irrespective of that situation, the general practitioner still has an important role in continuing management.

Areas demanding counselling include:

- any crisis situation - breaking bad news
- bereavement or grief
- terminal illness/palliative care (Chapter 7)
- marital problems (Chapter 2)
- family problems (Chapter 2)
- sexual dysfunction (Chapter 109)
- chronic pain
- anxiety and stress (Chapter 122)
- depression (Chapter 20)
- intellectual handicap in a child
- infertility (Chapter 108)
- any disease or illness, especially severe illness
- sexual abuse/child abuse (Chapter 85)
- domestic violence (Chapter 98)
- insomnia and other sleep disturbances (Chapter 71)

Crisis management

Crisis situations are not uncommon in general practice and people in crisis are usually highly aroused and demanding. Examples include tragic deaths, such as children drowning or sudden infant death syndrome (SIDS), unexpected marital break-up and breaking bad news.

Figure 5.2 Counselling skills: these skills include good eye contact, listening, empathy and appropriate communication skills

Aims of crisis intervention

- Resolve the crisis and restore psychological equilibrium as quickly and constructively as possible
- Encourage the person in crisis to regain control and take appropriate action

Principles of management

- Intervene early - actively and directly
- Establish an empathic alliance
- Be accessible
- Attend to family and social supports
- Be prepared for the difficult phase of 24 - 48 hours
- Do not carry the burden of crisis
- Aim for brief time-limited intervention (no more than six interviews over 6 weeks)
- When necessary, be prepared to provide short-term use of psychotropic drugs (e.g. a hypnotic), for two or three nights of good sleep

Ten rules to help those in distress

The following rules are given to those in crisis (personal explanation followed by a take-home handout):

1. *Give expression to your emotions*. You simply must accept your reactions as normal and not be afraid to cry or call out. Do not bottle up feelings.
2. *Talk things over with your friends*. Do not overburden them but seek their advice and listen to them. Do not avoid talking about what has happened.
3. *Focus on things as they are now- at this moment*. Do not brood on the past and your misfortune. Concentrate on the future in a positive way.
4. *Consider your problems one at a time*. Do not allow your mind to race wildly over a wide range of problems. You can cope with one problem at a time.

5. *Act firmly and promptly to solve a problem*. Once you have worked out a way to tackle a problem, go for it. Taking positive action is a step in allowing you to get on with life.
6. *Occupy yourself and your mind as much as possible*.
   Any social activity - sports, theatre, cards, discussion groups, club activity - is better than sitting around alone. Many people find benefit from a holiday visit to an understanding friend or relative. Religious people usually find their faith and prayer life a great source of strength at this time.
7. *Do not nurse grudges or blame other people*. This is not easy but you must avoid getting hostile. In particular, do not get angry with yourself and your family, especially your spouse.
8. *Set aside some time every day for physical relaxation*. Make a point of doing something physical, such as going for a walk, swimming or enjoying an easy exercise routine.
9. *Stick to your daily routine as much as possible*. At times of crisis a familiar pattern of regular meals and chores can bring a sense of order and security. Avoid taking your problems to bed and thus ensuring sleepless nights. Try to ‘switch off ’ after 8 pm. Taking sleeping tablets for those few bad nights will help.
10. *Consult your family doctor when you need help*. Your doctor will clearly understand your problem because stress and crisis problems are probably the commonest he or she handles. Consult your doctor sooner rather than later.
    - Remember that there are many community resources to help you cope (e.g. ministers, social workers, community nurses, crisis centres and church organisers)
    - Take care: do drive carefully and avoid accidents, which are more common at this time

Bereavement

Bereavement or grief may be defined as deep or intense sorrow or distress following loss.[10] Raphael uses the term to connote ‘the emotional response to loss: the complex amalgam of painful affects including sadness, anger, helplessness, guilt, despair’.[11]

The GP will see grief in all its forms over a wide variety of losses. Although the nature of loss and patient reaction to it varies enormously, the principles of management are similar.

Stages of normal bereavement

1. *Shock or disbelief*. Feelings include numbness and emptiness, searching, anxiety, fear and suicidal ideation, ‘I don't believe it’. Concentration is difficult and spontaneous emotions, such as crying, screaming or laughing, tend to occur. There may be a sense of the deceased's presence, and hallucinations (visual and auditory) may occur.
2. *Grief and despair*. Feelings include anger, ‘Why me?’, guilt and self-blame, and yearning. Social withdrawal and memory impairment may occur. The feeling of intense grief usually lasts about 6 weeks and the overall stage of grief and despair for about 6 months, but it can resurface occasionally for a few years. The last few months involve feelings of sadness and helplessness.
3. *Adaptation and acceptance*. Features of the third stage include significant feelings of apathy and depression. This phase takes a year or more. Physical illness is common and includes problems such as insomnia, asthma, bowel dysfunction, headache and appetite disturbances.

Pathological bereavement

Pathological bereavement can occur and may manifest as intense emotion, particularly anger, and multiple visits with somatic complaints; the patient often gets around to long dissertations about the deceased and the circumstances surrounding death. Extreme anger is likely when the sense of rejection is great, as with divorce or sudden death. Guilt can also be intense.[10]

Raphael's classification of the patterns of pathological grief and its various resolutions are presented in Table 5.2.[11]

Table 5.2 Patterns and resolution of pathological grief

Morbid or pathological patterns • Absence, inhibition or delay of bereavement • Distorted bereavement • Chronic grief (intense anguish continues unabated)
Outcome • Normal resolution, satisfactory adjustment; reintegration in life, satisfying attachments • General symptomatology (leading to increased care-eliciting behaviour) • Depression, suicidal behaviour • Other psychiatric disorders (anxiety state, phobia, mania, alcoholism, criminal activity, such as shoplifting) • Altered relationship patterns • Vulnerability to loss • Anniversary phenomenon • Death (more likely in the first 12 months)

The GP as counsellor[1]

Important rules to bear in mind:

- The bereaved may be feeling very guilty
- They may be angry towards their doctor or the medical profession in general
- They need a clear explanation as to the exact cause and manner of death. Autopsy reports should be obtained and discussed
- The bereaved tend to view an apparent lack of concern and support as disinterest or guilt[10]
- Early intervention averts pathological grief

The GP probably had a close relationship with the deceased and the family. The GP will have a special awareness of those at risk and the nature of the relationships within the family. The family is likely to maintain the relationship with the GP, expressing the physical and psychological effects of grief and consulting about intercurrent problems.[10]

Working through the stages of grief with patients will allow GPs to reach some acceptance of their own emotions, as well as ensure that patients feel supported and cared for, rather than distanced by embarrassment.

Help from religious sources is highly valued as it can meet both spiritual and personal needs. Other resources include funeral directors, hospice (and other) counsellors and support groups, such as those for SIDS.[10]

At least 30 minutes should be allowed for consultations.

Long-term counselling

Normal bereavement can persist for years. Ongoing counselling is indicated if it continues unabated or psychiatric referral sought if grief is extreme. Regular enquiries during routine consultations or meetings are important if the patient appears to be coping.

Breaking bad news

Good communication skills are fundamental to giving bad news appropriately. When bad news is broken insensitively or inadequately the impact can be distressing for both giver and recipient, leaving lasting scars for the latter. For the doctor it may represent professional failure, fear of people's reaction and feelings of guilt. Doctors should have a plan for this difficult process and learn how to cope with the recipient's reaction. Most of the circumstances described apply to unexpected death or anticipated death.

Optimal approach

Some basic initial rules:[12]

- If relatives have to be contacted it is preferable for the doctor (if at all possible) or a sympathetic police officer to make the contact personally, rather than a relatively matter-of-fact telephone call from the hospital or elsewhere
- If a telephone message is necessary it should be given by an experienced person
- The relatives or close friends should not drive to the clinic alone

The setting for the interview:

- Use a suitable quiet private room if possible
- See the recipients of the news alone in the room
- Advise that the meeting should be undisturbed

Guidelines for the doctor

- Be well prepared: check the facts and plan your approach
- Always ask those involved if they have heard any news or know the reasons for the consultation
- Always assess their understanding
- Give information in an unhurried, honest, balanced, empathic manner[13]
- Look directly at the person you are talking to, be honest and direct, and keep information simple (avoid technical language)
- The sad news must be accompanied by positive support, understanding and encouragement
- Give recipients time to react (offer time and moments of silence to allow the facts to sink in) and opportunities to ask questions
- Avoid false reassurance
- Remember that relatives appreciate the truth and genuine empathy
- In the event of death, relatives should be given a clear explanation of the cause of death

A list of guidelines for the interview is summarised in Table 5.3.[12]

Table 5.3 Breaking bad news: recommended actions during the interview

Allow: • time • opportunities to react • silence • touching • free expression of emotions • questions • viewing of a dead or injured body
Avoid: • rushing • bluntness • withholding the truth • platitudes • protecting own inadequacies • euphemisms • the notion ‘nothing more can be done’ • using medical jargon

- meeting anger with anger
- leaving the patient or loved one without a follow-on contact

Source: After McLauchlan[12]

Coping with patient responses

- The responses cover a wide range - stunned silence, disbelief, acute distress, anger, extreme guilt
- Be prepared for any of these responses
- Appropriate training using simulated patients, video replays and skilled feedback improves communication skills
- Give permission and encouragement for reactions, such as crying and screaming
- Have a box of tissues available
- A comforting hand on the shoulder or arm or holding a hand is an acceptable comfort zone
- Offer a cup of tea or a cool drink if available
- Ask the patients or relatives how they feel, what they would like to do and if they want you to contact anyone
- Arrange follow-up
- Give appropriate patient education material
- Provide information about support services

The depressed patient

Studies have emphasised the importance and therapeutic efficacy of counselling in the management of the depressed patient.[14] The most practical approach by the GP to the depressed patient is empathy, support and a logical explanation of their malaise. The author gives the following explanation to the patient.

> Depression is a very real illness that affects the entire mind and body. It seriously dampens the five basic activities of humans, namely their energy for activity, sex drive, sleep, appetite and ability to cope with life. They cannot seem to lift themselves out of their misery or fight it themselves. Superficial advice to ‘snap out of it’ is unhelpful because the person has no control over it.
>
> The cause is somewhat mysterious but it has been found that an important chemical is present in smaller amounts than usual in the nervous system. It is rather like a person low in iron becoming anaemic.
>
> Depression can follow a severe loss such as the death of a loved one, a marital separation or financial loss. On the other hand it can develop for no apparent reason although it may follow an illness such as glandular fever or influenza, an operation or childbirth.

Emphasising the ‘missing chemical’ theory really helps patients and family come to terms with an illness that tends to have socially embarrassing connotations. It also helps compliance with therapy when antidepressant medication is prescribed.

Ongoing contact, support and availability are an important component of counselling with appropriate referral to someone with more expertise, should that be required. CBT is a most effective and important approach to the management of depressive illness.

Chronic pain

Patients suffering from long-term pain are a special problem, especially those with back pain who seem to be on a merry-go-round of failed multiple treatments and complex psychosocial problems. These patients are frequently treated in pain clinics. As family doctors we often observe an apparently normal, pleasant person transformed into a person who seems neurotic, pain-driven and doctor-dependent. The problem is very frustrating to the practitioner, often provoking feelings of suspicion, uncertainty and discomfort.

De Vaul et al.[15] list five subgroups of patients where perplexing pain presents as the major symptom:

1. pain as a symptom of depression
2. pain as a delusional symptom of psychosis
3. Pain as a conversion symptom of hysterical neurosis

4. pain as a symptom of an unresolved bereavement reaction
5. pain as a symptom of a ‘need to suffer’

Patients who somatise their symptoms present one of the most difficult challenges to our skills and usually require a multidisciplinary team approach.

Management involves:

- thorough medical assessment
- psychological assessment
- detailed explanations to the patient and family about treatment
- rational explanations about the cause of the pain
- management of associated problems (e.g. depression, sexual dysfunction)
- behavioural modification to encourage increased activity and a gradual return to normality

A useful explanation

The author finds the following account a most useful method of explaining perplexing continuing back pain or neuralgia to patients (where there is no evidence of a persisting organic lesion).

> Part of the problem is that psychological factors continue to aggravate and maintain the problem even though the reason for the pain in the first place may have disappeared. It is a similar problem to a person who has had a painful leg amputated. Even though it has been removed, the patient can still feel the leg and maybe even the pain. The patient has a ‘phantom limb’. The nervous system, especially the brain, can play funny tricks on us in this way.
>
> This means that even though the original disc injury has settled after several weeks, the body can still register the pain. This is more likely to occur in people who have become anxious and depressed about their problem. The pain continues. Someone once described it as a ‘tension headache that has slipped down to the back’.

Problem gambling

Problem or pathological gambling is persistent and recurrent maladaptive gambling behaviour despite its detrimental effect (disruption of personal, family or work life). It is undoubtedly a dependence disorder similar to alcohol and other drugs with a similar approach to management. Refer to DSM-IV criteria for pathological gambling. Prevalence: 0.5 - 1.5% adult population.

Dangers

- suicide risk (high)
- major depression (up to 75%)
- stress-related problems
- domestic violence

Key warning signs

- gambling > $100 week
- chasing losses

Other telltale signs

- Spending many hours gambling
- Placing larger, more frequent bets
- Lying about behaviour
- Being secretive
- Promising cutting back, but not doing it
- Impulsive activity
- Mood swings

- Gambling at the expense of other pleasant social activities
- Growing debts
- Excessive drinking

First-line management

- Ask (as part of social history)
- Consider South Oaks Gambling screen to support provisional diagnosis[16]
- Firm confrontation if suspected
- Consider using the Prochaska and Di Clemente model of change (p. 1249)
- Provide education material
- Look into the family (? domestic violence) and provide support
- Advise the family not to provide ‘rescue money’
- There is no recommended pharmacological treatment

Counselling approach[17]

Problem gambling is a treatable condition and GPs can provide a central role in management. As for smoking and alcohol dependence, a brief intervention and education consultation session about the impact of excessive behaviour can be most effective. CBT is a very effective treatment for gambling. It combines systematic discussion and carefully structured behavioural assignments to help patients modify problematic thinking patterns and behaviours. ‘CBT directed towards correcting erroneous perceptions, irrationals beliefs and misunderstanding of concepts of randomness and independence of chance events is a fundamental element of any therapeutic approach.’ [17]

It is appropriate to use specialist gambling counsellors when one's intervention is not proving effective or where there is evidence of disturbing gambling problems. (Refer < http://www.gamblersanonymous.org >.)

SUMMARY: COUNSELLING SKILLS STRATEGIES

- Provide guidance and facilitation to allow the patient to gain insight
- Use appropriate ‘gentle’ confrontation to allow self-examination
- Help patients to explore their own situation and express emotions such as anxiety, guilt, fear, anger, hostility and hurt feelings
- Explore possible feelings of insecurity and allow free expression of such feelings
- Ask key searching questions, e.g:
    - What do you think deep down is the cause of your problem?
    - How do you think your problem should be treated?
- Provide ‘okay’ specific suggestions, e.g:
    - I wonder if your basic problem is that you are a perfectionist?
    - Many people in your situation feel guilty about something which may even be trivial and need to feel forgiven.

Effective counselling comes from commitment, experience and a genuine caring compassionate feeling for patients and their ethos.

If one feels out of one's depth, then immediate referral to an expert is important. CBT is a most appropriate therapy for most conditions.

KEY RULES TO COUNSELLING

1. The patient must leave feeling better.

2. Provide *insight* into their illness and/or behaviour.
3. Address any feelings of *guilt* (people must feel okay or forgiven about any perceived transgression).

REFERENCES

1. Hassed C. Counselling. In: *Final Year Handbook*. Melbourne: Monash University, Department of Community Medicine, 1992: 97 - 104.
2. Ramsay AT. The general practitioner as an effective counsellor. Aust Fam Physician, 1990; 19: 473 - 9.
3. Harris RD, Ramsay AT. *Health Care Counselling*. Sydney: Williams & Wilkins, 1988: 68 - 95.
4. Annon JS. *The Behavioural Treatment of Sexual Problems*. Vols 1 and 2. Honolulu: Enabling Systems Inc, 1974.
5. Craig S. A medical model for infertility counselling. Aust Fam Physician, 1990; 19: 491 - 500.
6. Cook H. Counselling in general practice: Principles and strategies. Aust Fam Physician, 1986, 15: 979 - 81.
7. Frank JF. Foreword. In: Traux CB, Carkhuff RR. *Towards Effective Counselling and Psychotherapy: Training in Practice*. New York: Aldine, 1967: ix.
8. Traux CB, Carkhuff RR. *Towards Effective Counselling and Psychotherapy*. New York: Aldine, 1967.
9. Tiller JWG. Cognitive Behaviour Therapy in Medical Practice. Aust Prescriber, 2001; 24(2): 33 - 7.
10. Williams AS. Grief counselling. Aust Fam Physician, 1986; 15: 995 - 1002.
11. Raphael B. *The Anatomy of Bereavement: A Handbook for the Caring Professions* . London: Hutchinson, 1984: 33 - 62.
12. McLauchlan CAJ. Handling distressed relatives and breaking bad news. In: Skinner D. *ABC of Major Trauma*. London: British Medical Association, 1991: 102 - 6.
13. Cunningham C, Morgan P, McGucken R. Down syndrome: is dissatisfaction with disclosure of diagnosis inevitable? Devel Med Child Neurol, 1984; 26: 33 - 9.
14. Jackson HJ, Moss JD, Solinski S. Social skills training: an effective treatment for unipolar non-psychotic depression? Aust NZ J Psychiatry, 1985; 19: 342 - 53.
15. De Vaul RA, Zisook S, Stuart HJ. Patients with psychogenic pain. J Fam Pract 1977; 4(1): 53 - 5.
16. Lesieur HR, Blume S. The South Oaks Gambling Screen (SOGS): a new instrument for the identification of pathological gamblers. American J of Psychiatry, 1987; 144: 1184 - 8.
17. Blaszczynski A. How to treat: Problem gambling. Australian Doctor, 2005; 12 August: 37 - 44.

CHAPTER 6. DIFFICULT, DEMANDING AND ANGRY PATIENTS

There are patients in every practice who give the doctor and staff a feeling of ‘heartsink’ every time they consult.

THOMAS O'DOWD 1988

DIFFICULT, DEMANDING AND ANGRY PATIENTS

Weston defines a ‘difficult patient’ as one with whom the physician has trouble forming an effective working relationship.[1] However, it is more appropriate to refer to difficult problems rather than difficult patients - it is the patients who have the problems while doctors have the difficulties.

Some characteristics of problematic patients, from the doctor's perspective, include:

- frequent attenders with trivial illness
- multiple symptomatology
- undifferentiated illness
- chronic tiredness
- negative investigations
- dissatisfaction with treatment, especially procedures
- non-compliant
- hostile or angry
- attending multiple therapists
- demanding on staff
- inconsiderate of the doctor's time
- taking multiple drugs
- seductive, then demanding
- garrulous
- manipulative
- taciturn and uncommunicative
- all knowing
- personality disorders in general, borderline personality disorders in particular

Such patients are often referred to as the ‘heartsink’ patients,[2] referring to that certain sinking feeling on seeing them in the waiting room or their name on the booking list. They can provoke negative feelings in us and we have to discipline ourselves to be patient, responsible and professional.

An inevitably poor consultation will follow if we allow feelings of hostility to affect our communication with the difficult patient, especially the demanding, angry or ‘compo’ patient.

However, it is important not to misdiagnose organic disease and also to consider the possibilities of the following disorders, which may be masked:

- anxiety
- depression
- obsessive compulsive disorder
- bipolar disorder (manic depression)
- drug dependency
- alcohol abuse
- schizophrenia

It is therefore appropriate to maintain traditional standards by continually updating the database, integrating psychosocial aspects, carefully evaluating new symptoms, conducting an appropriate physical examination and being discriminating with investigations.

Management of the violent and dangerous patient is presented in Chapter 45 on the disturbed patient (p. 497).

MANAGEMENT STRATEGIES

Our professional responsibility is to rise above interpersonal conflict and facilitate productive communication by establishing a caring and responsible relationship with such patients. An appropriate strategy is to follow Professor Aldrich's precepts for the ‘difficult’ patients who do not have an organic disorder or a psychiatric illness.[3]

1. Give up trying to cure them - they are using their symptoms to maintain their relationship with you: accept them as they are.
2. Accept their symptoms as expressions of their neurosis. Make a primary positive diagnosis - only test if you have to.
3. Structure a program for them, for example, ‘Mrs Jones, I have decided that we should meet for 15 minutes every second Wednesday at 10 am.’
4. During the consultation, demonstrate your genuine interest in the person's life, garden, work and so on; show less interest, even boredom, for the litany of complaints.

Other management guidelines include:

- Use reassurance with caution - it is insufficient by itself and should be soundly based.
- Be honest and maintain trust.
- Allow the patient a fair share of your time - this is your part of the contract. At the same time indicate that there are limits to your time (set rules).
- Be polite yet assertive.
- Avoid using labels of convenience and placebo therapy.
- Be honest about your understanding (or lack of understanding) of the problems.
- Remember that the consultation is often the therapy, without a prescription.
- Do not undermine other doctors. Avoid collusion.
- Have limited objectives - zealous attempts to cure may be inappropriate.
- Do not abandon the patient, however frustrating the relationship. Accept this as a legitimate role.
- Remain available if alternative therapies are sought by the patient.
- Take extra care with the ‘familiar’ patient and sometimes the patient who brings gifts. Maintain your professional role.
- If you are uncomfortable with counselling, consider early referral to a counsellor while maintaining contact in the future.
- You may have to accept that there are some people no one can help.

A ‘heartsink’ survival kit

A pilot workshop of managing ‘heartsink’ patients described by Mathers and Gask[4] led to the formulation of a ‘heartsink survival’ model for the management of patients with somatic symptoms of emotional distress.

The first part of the three-part model, which is called ‘feeling understood’, includes a full history of symptoms, exploration of psychosocial cues and health beliefs, and a brief focused physical examination. In the second stage, termed ‘broadening the agenda’, the basic aim is to involve discussion of both emotional and physical aspects during the consultation. It includes reframing the patient’s symptoms and complaints to provide insight into the link between physical, psychological and life events.

In the third stage, ‘making the link’, simple patient education methods are used to explain the causation of somatic symptoms, such as the way in which stress, anxiety or depression can exaggerate symptoms. It also includes projection or identification techniques using other sufferers as examples.

THE ANGRY PATIENT

Anger in patients and their relatives is a common reaction in the emotive area of sickness and healing. The anger, which may be concealed or overt, might be a communication of fear and insecurity. It is important to bear in mind that many apparently calm patients may be harbouring controlled anger. The practice of our healing art is highly emotive and can provoke feelings of frustration and anger in our patients, their friends and their relatives.

Anger is a normal and powerful emotion, common to every human being, yet with an enormous variety of expression. The many circumstances in medicine that provoke feelings of anger include:[5]

- disappointment at unmet expectations
- crisis situations, including grief
- any illness, especially an unexpected one
- the development of a fatal illness
- iatrogenic illness
- chronic illness, such as asthma
- financial transactions, such as high cost for services
- referral to colleagues, which is often perceived as failure
- poor service, such as long waits for an appointment
- problems with medical certificates
- poor response to treatment
- inappropriate doctor behaviour (e.g. brusqueness, sarcasm, moralistic comments, aloofness, superiority)

The patient's anger may manifest as a direct confrontation with the doctor or perhaps with the receptionist, with litigation or with public condemnation.

In an extreme example, a Melbourne doctor was shot and killed by an angry patient who had been denied a worker's compensation certificate for a claim considered unjustified.

When a patient expresses anger about the medical profession or our colleagues it may be directed at us personally and, conversely, if directed to us it may be displaced from someone else, such as a spouse, employer or other figure of authority.

What is anger?

Anger is a person's emotional response to provocation or to a threat to his or her equilibrium. If inappropriate, it is almost always the manifestation of a deeper fear and of hidden insecurity. Angry abusive behaviour may be a veiled expression of frustration, fear, self-rejection or even guilt.

On the other hand, its expression may be a defence against the threat of feeling too close to the doctor, who could have an overfamiliar, patronising or overly friendly attitude towards the patient. Some patients cannot handle this threatening feeling.

Basically, anger may be a communication of fear and insecurity. The patient could be saying, ‘I am afraid there is something seriously wrong with me. Are you doing everything to help me?’.

Consulting strategies[6]

When one feels attacked unfairly, to react with anger is a natural human response. This response, however, must be avoided since it will damage the doctor-patient relationship and possibly aggravate the problem.

- The initial response should be to remain calm, keep still and establish eye contact.
- ‘Step back’ from the emotionally charged situation and try to analyse what is happening.
- Ask the patient to sit down and try to adopt a similar position (the mirroring strategy) without any aggressive pose.
- Address the patient (or relative) by the appropriate name, be it Mr or Mrs Jones or a first name.
- Appear comfortable and controlled.
- Be interested and concerned about the patient and the problem.

- Use clear, firm, non-emotive language.
- Listen intently.
- Allow patients to ventilate their feelings and help to relieve their burdens.
- Allow patients to ‘be themselves’.
- Give appropriate reassurance (do not go overboard to appease the patient).
- Allow time (at least 20 minutes).

Analysing the responses

- Search for any ‘hidden agenda’.
- Recognise the relationship between anger and fear.

Recognising distress signals

It is important to recognise signs of deteriorating emotional distress:[7]

- body language (demonstrative agitated movements or closing in)
- speech (either becoming quiet or more rapid and louder)
- colour (either becoming flushed or pale)
- facial expression (as above, tense, tightening of muscles of eye and mouth, loss of eye contact)
- manner (impatient, threatening)

Skilful consulting strategies should then be employed. It is worthwhile having a contingency plan, such as memorising a telephone number to summon ecurity help.

Questions to uncover the true source of anger

The following represent some typical questions or responses that could be used during the interview.

Rapport building

- ‘I can appreciate how you feel.’
- ‘It concerns me that you feel so strongly about this.’
- ‘Tell me how I can make it easier for you.’

Confrontation

- ‘You seem very angry.’
- ‘It's unlike you to be like this.’
- ‘I get the feeling that you are upset with … ’
- ‘What is it that's upsetting you?’
- ‘What really makes you feel this way?’

Facilitation, clarification

- ‘I find it puzzling that you are angry with me.’
- ‘So you feel that … ’
- ‘You seem to be telling me … ’
- ‘If I understand you correctly … ’
- ‘Tell me more about this … ’
- ‘I would like you to enlarge on this point - it seems important.’

Searching

- ‘Do you have any special concerns about your health?’
- ‘Tell me about things at home.’

- ‘How are things at work?’
- ‘How are you sleeping?’
- ‘Do you have any special dreams?’
- ‘Do you relate to anyone who has a problem like yours?’
- ‘If there's any one thing in your life that you would like to change, what would it be?’

Some important guidelines are summarised in Table 6.1.

Table 6.1 Guidelines for handling the angry patient

Do	Don't
Listen	Touch the patient
Be calm	Meet anger with anger
Be comfortable	Reject the patient
Show interest and concern	Be a ‘wimp’
Be conciliatory	Evade the situation
Be genuine	Be overfamiliar
Allay any guilt	Talk too much
Be sincere	Be judgmental
Give time	Be patronising
Arrange follow-up	
Act as a catalyst and guide	

Management

When confronted with an angry patient the practitioner should be prepared to remain calm, interested and concerned. It is important to listen intently and allow time for the patient to ventilate his or her feelings.

A skilful consultation should provide both doctor and patient with insight into the cause of the anger and result in a contract in which both parties agree to work in a therapeutic relationship. The objective should be to come to amicable terms which, of course, may not be possible, depending on the nature of the patient's grievance.

If the problem cannot be resolved in the time available a further appointment should be made to continue the interview.

Sometimes it may be appropriate to advise the patient to seek another opinion. If the angry patient does have problems with relationships and seeks help, it would be appropriate to arrange counselling so that the patient acquires a more realistic selfimage, thus leading to improved self-esteem and effectiveness in dealing with people. In addition, it should lead to the ability to withstand frustration and cope with the many vicissitudes of life - a most rewarding outcome for a consultation that began with confrontation.

REFERENCES

1. McWhinney I. *A Textbook of Family Medicine*. New York: Oxford, 1989: 96 - 8.
2. O'Dowd TC. Five years of ‘heartsink’ patients in general practice. Br Med J, 1988; 297: 528 - 30.
3. Elliott CE. ‘How am I doing?’ Med J Aust, 1979; 2: 644 - 5.
4. Mathers NJ, Gask L. Surviving the ‘heartsink’ experience. Fam Practice, 1995; 12: 176 - 83.
5. Murtagh JE. The angry patient. Aust Fam Physician, 1991; 20: 388 - 9.
6. Mongtomery B, Morris L. *Surviving: Coping with a Crisis*. Melbourne: Lothian, 1989: 179 - 86.
7. Lloyd M, Bor R. *Communication Skills for Medicine*. London: Churchill Livingstone, 1996: 135 - 7.

CHAPTER 7. PALLIATIVE CARE

When the cancer that later took his life was first diagnosed, Senator Richard L. Neuberger remarked upon his 'new appreciation of things I once took for granted - eating lunch with a friend, scratching my cat Muffet's ears and listening to his purrs, the company of my wife, reading a book or magazine in the quiet of my bed lamp at night, raiding the refrigerator for a glass of orange juice or a slice of toast. For the first time, I think I actually am savouring life'.

BETTER HOMES AND GARDENS MAGAZINE

Palliative care is the active total (holistic) care of patients whose disease is not responsive to curative treatment … The goal of palliative care is achievement of the best quality of life for patients and their families.

WHO CANCER PAIN RELIEF & PALLIATIVE CARE. GENEVA, WHO, 1990

To enable a person to live in dignity, peace and comfort throughout their illnesses means responding to physical, psychological, emotional, social and spiritual needs.[1]

PALLIATIVE CARE: THE NITTY GRITTY HANDBOOK. E FAIRBANK, T BANK

PALLIATIVE CARE

Palliative care is comprehensive, continuing, multidisciplinary patient care that involves the patients and their carers, consultants, domiciliary nurses, social workers, clergy and other health professionals who are able to contribute to optimal team care.

The fundamental principles of palliative care are:[2]

- good communication
- management planning
- symptom control
- emotional, social and spiritual support
- medical counselling and education
- patient involvement in decision making
- support for carers

THE DISEASES

Palliative care applies not only to incurable malignant disease and HIV/AIDS but also to several other diseases, such as end-stage organ failure (heart failure, kidney failure, respiratory failure and hepatic failure) and degenerative neuromuscular diseases.

THE SPECIAL ROLE OF THE FAMILY DOCTOR

The GP is the ideal person to manage palliative care for a variety of reasons - availability, knowledge of the patient and family, and the relevant psychosocial influences. A key feature is the ability to provide the patient with independence and dignity by managing palliative care at home. Someone has to take the responsibility for leadership of the team and the most appropriate professional is a trusted family doctor.

Most patients and their families require answers to six questions:[3]

- What is wrong?
- What can medical science offer?
- Will I suffer?
- Will you look after me?
- How long will I live?
- Can I be looked after at home?

Caring honesty is the best policy when discussing the answers to these questions with the patient and family. Never lie to a patient and always avoid thoughtless candour.

SUPPORT FOR PATIENTS AND CARERS

Studies have indicated that the most common complaints of patients are boredom and fear of the unknown. This highlights the importance for the attending doctor of the following points:

- Give emotional support.
- Listen and be receptive to unexpressed 'messages'.
- Treat the sufferer normally, openly,
- enthusiastically and confidently.
- Show empathy and compassion.
- Employ good communication skills.
- Give honest answers without labouring the point or giving false hope.
- Provide opportunities for questions and clarification.
- Show an understanding of the patient's needs and culture.
- Adopt a whole person approach: attend to physical, psychosocial and spiritual needs.
- Anticipate and be prepared for likely problems. Special points worth emphasising are:
- The patient needs a feeling of security.
- Provide reassurance that the patient will not suffer unnecessarily.
- Be prepared to take the initiative and call in others who could help (e.g. clergy, cancer support group, massage therapists).
- Patients must not be made to feel isolated or be victims of the so-called 'conspiracy of silence' in which families collude with doctors to with hold information from the patient.
- The worst feeling a dying patient can sense is one of rejection and discomfort on the part of the doctor.
- Always be prepared to refer to an oncologist or appropriate therapist for another opinion about further management. The family and patient appreciate the feeling that every possible avenue is being explored.

Note: Always establish what the patient knows and wants to know.

THE GOLD STANDARDS FRAMEWORK (UK)

This framework, which provides an optimal model for palliative care by the primary care team, focuses on seven key tasks:

1. optimal quality of care
2. advanced planning (including out of hours)
3. teamwork
4. symptom control
5. patient support
6. carer support
7. staff support

The strategy has been shown to increase the number of patients dying in their preferred place with improved quality of care.[4]

SYMPTOM CONTROL

Common symptoms

- boredom (the commonest symptom)
- loneliness/isolation
- fear/anxiety
- pain
    - physical
    - emotional
    - spiritual
    - social
- anorexia
- nausea and vomiting
- constipation

The grief reaction

This follows five stages, as identified by Kübler-Ross:[5]

1. denial and isolation
2. anger
3. bargaining
4. depression
5. acceptance

This model provides a useful guideline in understanding the stages a patient and family will be experiencing.

The principles of symptom management are summarised in Table 7.1, and the goals of treatment according to the different stages of cancer are presented in Figure 7.1.

Table 7.1 Principles of symptom management[2]

Determine the cause.
Treat simply.
Provide appropriate explanation of symptoms and treatment.
Provide regular review.
Give medication regularly around the clock, not ad hoc.
Plan ‘breakthrough’ pain-relieving doses.
Provide physical treatment as necessary (e.g. paracentesis, pleural tap, nerve block).
Provide complementary conservative therapy (e.g. massage, physiotherapy, occupational therapy, dietary advice, relaxation therapy).
Provide close supervision.


Time of diagnosis
Progression of time
Time of death
At first
Then
Eventually
treatment is potentially curative
treatment is palliative
radiotherapy
chemotherapy
surgery
hormonal
symptom control
symptom control at any time

Figure 7.1 Stages of cancer management: the goals of treatment differ according to the different stages After J. Buchanan et al.

6

PAIN CONTROL IN CANCER

Achieving pain relief is one of the most important functions of palliative care and patients need reassurance that they can expect such relief. The principles of relief of cancer pain are:[1]

1. Treat the cancer.
2. Raise the pain threshold:
    - Provide appropriate explanation.
    - Allow the patient to ventilate feelings and concepts.
    - Give good psychosocial support.
    - Use antidepressants or hypnotics.
3. Add analgesics according to level of pain, for example, opioids (if necessary).
4. Use specific drugs for specific pain - not all pain responds to analgesics (refer Table 7.2 on p. 50).

Table 7.2 Treatment options for cancer pain[6] (based on aetiology)

Aetiology	1st line treatment	2nd line treatment	Other treatment modalities to consider
Nociceptive pain: stimulation of sensory nerves	Aspirin Antidepressants	Opioids Corticosteroids NSAIDs	Radiotherapy Neurosurgery
Neuropathic pain: direct nerve involvement (e.g. brachialgia, sciatica)	Opioids Antidepressants (e.g. amitriptyline) Antiepileptics (e.g. carbamazepine, gabapentin)		Spinal morphine Local anaesthetic Ketamine

Dysaesthesia: superficial burning pain	Antidepressants	Opioids	Local anaesthetic TENS
Pressure pain: tumour-associated oedema (e.g. raised intracranial pressure)	Corticosteroids (e.g. dexamethasone)	Opioids	Radiotherapy Neurosurgery
Bony metastases and other tissue destruction	NSAIDs Aspirin	Opioids	Radiotherapy (the most effective) Hormones Orthopaedic surgery
Muscle spasm pain	Diazepam Clonazepam Baclofen	Opioids Dantrolene	
Viscus (hollow organ) obstruction (e.g. colic, tenesmus)	Antispasmodics (e.g. hyoscine)	Opioids Chlorpromazine Corticosteroids	Palliative surgery Radiotherapy
Metabolic effects: hypercalcaemia	Biphosphonates (APD)		
Skin infiltration/ulceration	Aspirin Opioids	Corticosteroids	Treat infection Dressings Palliative surgery Radiotherapy

5. Set realistic goals.
6. Organise supervision of pain control.

Note: The right drug, in the right dose, given at the right time relieves 80 - 90% of the pain.[1]

Use of analgesics[7]

These should be given by the clock and administered according to the three-step method.

Step 1: Mild pain

Start with basic non-opioid analgesics: aspirin 600 - 900 mg (o) 4 hourly (preferred) *or*

paracetamol 1 g (o) 4 hourly ± NSAID

Step 2: Moderate pain

Use low dose or weak opioids (according to age and condition) or in combination with non-opioid analgesics (consider NSAIDs): add morphine 5 to 10 mg (o) 4 hourly increase in increments of 30 - 50% up to 15 - 20 mg *or* oxycodone up to 10 mg (o) 4 hourly or CR 10 mg (o) 12 hourly *or*

oxycodone 30 mg, rectally, 8 hourly

Step 3: Severe pain

Maintain non-opioid analgesics. Larger doses of opioids should be used and morphine is the drug of choice:

- morphine 10 - 15 mg (o) 4 hourly, increasing to 30 mg if necessary *or*
- morphine CR/SR tabs or caps (o) 12 hourly or once daily
  - Give dosage according to individual needs (morphine CR/SR comes in 5, 10, 15, 20, 30, 50, 60, 90, 100, 120, 200 mg tablets or capsules).
  - The proper dosage is that which is sufficient to alleviate pain.
  - The usual starting dose is 20 - 30 mg bd.
  - Give usual morphine 10 mg with first dose of morphine CR/SR and then as necessary for 'rescue dosing'.
  - Gauge the probable dose of the long-acting morphine from the standard dosage.
  - To convert to morphine CR/SR, calculate the daily oral dose of regular morphine and divide by 2 to get the 12 hourly dose.
  - Do not crush or chew the tablets or capsules.

Guidelines

- Ensure that pain is likely to be opioid-sensitive.
- Give morphine orally (if possible) either by mixture (preferred) or tablets.
- Starting doses are usually in the range of 5 - 20 mg (average 10 mg).
- If analgesia is inadequate, the next dose should be increased by 50% until pain control is achieved.
- Give it regularly, usually 4 hourly, before the return of the pain (see Figure 7.2 on p. 51).



Figure 7.2 Appropriate schedule of analgesia to achieve optimal pain control

- Many patients find a mixture easier to swallow than tablets (e.g. 10 mg/10 ml solution).
- Constipation is a problem, so treat prophylactically with regular laxatives and carefully monitor bowel function.
- Order a 'rescue dose' (usually 5 - 10 mg) for breakthrough pain or anticipated pain (e.g. going to toilet).
- Order antiemetics (e.g. haloperidol prn at first; usually can discontinue in 1 - 2 weeks as tolerance develops).
- Reassure the patient and family about the safety and efficacy of morphine (see Table 7.3 on p. 51). (Beware of opiophobia.)

Table 7.3 Common myths about morphine[7]

• **Morphine is a last resort.** This is not so, and there is no maximum dose.
• **The patient will become addicted.** This is rare and probably irrelevant in the context of palliative care.
• **The patient will need ever-increasing amounts.** The drug does not lose its effect but is usually increased according to disease progression.
• **Morphine will cause respiratory depression.** This is rarely a problem and may even help those with dyspnoea. An overdose can be reversed with an injection of naloxone.
• **Morphine will shorten life.** The reverse may in fact apply. It is not being used for euthanasia but to control pain.

- Using morphine as a mixture with other substances (e.g. Brompton's cocktail) has no particular advantage.
- Pethidine is not recommended (short half - life, toxic metabolites) and codeine and IM morphine should be avoided.
- Other opioids are sometimes used instead of morphine (Table 7.4).

Table 7.4 Non-morphine opioids used in pain control1, [8]

Opioid (oral unless indicated other)	Duration of action hours)	Dose equivalent to oral morphine 10 mg
Codeine	3 - 5	60 mg
Dextromoramide (Palfium)	2	20 mg
Diamorphine (heroin)	3 - 5	7 mg
Fentanyl (transdermal) (Durogesic)	72	100 μg
Hydromorphone (Dilaudid)	3 - 4	2.5 mg (oral)
Methadone	variable 4 - 12	7 mg
Oxycodone		
Endone/OxyNorm (oral)	3 - 5	10 mg
Oxycontin (CR)	3 - 4	5 mg
Proladone (rectal)	6 - 12	10 mg
Tramadol (Tramal)	4 - 6	40 mg

- Fentanyl is a potent synthetic opioid which is available as a transdermal system. Effective and good for compliance. It is the least constipating opioid.
- Hydromorphone is a potent analgesic available as oral liquid, tablets and injection and is now widely used in palliative care. It facilitates oral dosing when a high opioid dose is required and because of its short half-life (2 - 3 h) may reduce the incidence of side-effects in the frail and elderly.

Opioid rotation

This practice involves changing from one strong opioid to another in patients with dose limiting sideeffects. Different opioids have differences in opioid receptor binding.

Morphine can be alternated with oxycodone, hydromorphone, methadone, fentanyl and others. Transdermal patches of fentanyl are an alternative to parenteral morphine.

Parenteral morphine

This is generally given subcutaneously (not IV or IM). Indications are:[6]

1. unable to swallow (e.g. severe oral mucositis; dysphagia; oesophageal obstruction)
2. bowel obstruction
3. severe nausea and vomiting
4. at high oral dose (i.e. above 100 - 200 mg dose) there appears to be no additional benefit from further dose increments.

Adjuvant therapy

Refer to Table 7.2. In all three steps of pain control consider the use of adjuvant ‘analgesics’; while not strictly speaking analgesics, they contribute to pain relief. Examples are corticosteroids, antidepressants, psychotropic agents and anticonvulsants.

Pain control

Bone pain:

- aspirin, paracetamol, NSAIDs are helpful co-analgesics

Neuropathic pain (direct involvement of nerves):

- antidepressants (e.g. amitriptyline)
- anticonvulsants (e.g. carbamazepine, gabapentin)
- ketamine, an anaesthetic agent, is valuable for difficult pain but requires expertise in administration.

Neurological pressure:

- corticosteroids for spinal cord compression and raised intracranial pressure
  e.g. dexamethasone 4 - 16 mg (o, sc or im) daily in morning *or* prednisolone 25 - 100 mg (o) daily

Continuous subcutaneous infusion of morphine

When the oral and/or rectal routes are not possible or are ineffective a subcutaneous infusion with a syringe pump can be used.

It is also useful for symptom control when there is a need for a combination of drugs (e.g. for pain, nausea and agitation). It may avoid bolus peak effects (sedation, nausea or vomiting) or trough effects (breakthrough pain) with intermittent parenteral morphine injections.

Practical aspects

- Access to the subcutaneous space is via a 21 gauge butterfly needle, which is replaced regularly (1, 2, 3 or 4 days).
- Most regions are suitable. The more convenient are the abdomen, the anterior thigh, and the anterior upper arm (usually the anterior abdominal wall is used).
- The infusion can be managed at home.
- About one-half to two-thirds of the 24 hour oral morphine requirement is placed in the syringe.
- The syringe is placed into the pump driver, which is set for 24 hour delivery.
- Areas of oedema are not suitable.

Spinal morphine

Epidural or intrathecal morphine is sometimes indicated for pain below the head and neck, where oral or parenteral opioids have been ineffective. It is necessary to insert an epidural or intrathecal catheter (anaesthetist or neurosurgeon).[7]

COMMON SYMPTOM CONTROL

Common symptoms can be controlled as follows:[1]

Anorexia

metoclopramide 10 mg tds *or*

corticosteroids (e.g. dexamethasone 2 - 8 mg tds) high-energy drink supplements

Constipation[6]

If opioids need to be maintained, the laxatives need to be peristaltic stimulants, not bulkforming agents. Aim for firm faeces with bowels open about every third day.

e.g. docusate (Coloxyl) with senna, 2 daily *or* bd bisacodyl (Durolax) 5 - 10 mg bd *or*

Movicol, one sachet in 125 ml water 1, 2 or 3 times daily.

Rectal suppositories, microenemas or enemas may be required (e.g. Microlax).

Shaw's (or PCU) cocktail is useful for severe constipation. With a small quantity of water melt one tablespoon of Senokot granules in a microwave oven. Add 20 ml Agarol and constitute to 100 ml with cold or warm milk or icecream.

Terminal respiratory congestion (death rattles)[7,8]

- Hyoscine hydrobromide 0.4 mg SC, 4 hourly or 0.8 - 1.6 mg daily by SC infusion
- Hyoscine butylbromide or atropine can also be given by SC injection.

These agents dry secretions and stop the ‘death rattle’.

Dyspnoea

Identify the cause, such as a pleural effusion, and treat as appropriate. Pleural taps can be performed readily in the home. Corticosteroids can be given for lung metastases. Oxygen may be necessary to help respiratory distress in the terminal stages and bedside oxygen can be readily obtained. Morphine can be used for intractable dyspnoea, together with haloperidol or a phenothiazine for nausea.

Terminal distress/restlessness[7, 8, 9, 10]

(Exclude reversal causes, e.g. drugs, fear, faecal impaction, urinary retention.)

1st choice:

clonazepam	0.5 mg SC bolus or 0.25 - 0.5 mg (o) 12 hourly (drops SL) or tabs[7]
or	1 - 4 mg over 24 hours in SC syringe driver
midazolam (very effective but expensive)	2.5 - 5 mg SC 1 - 3 hourly prn or 15 mg/day by SC infusion

If very severe: add phenobarbitone as SC infusion or (with care because of fitting) haloperidol

Nausea and vomiting

If due to morphine: haloperidol 1.5 - 5 mg daily[1] (can be reduced after 10 days) *or*

prochlorperazine (Stemetil) 5 - 10 mg (o) qid *or* 25 mg rectally bd

If due to poor gastric emptying, use a prokinetic agent: metoclopramide or cisapride or domperidone

Consider ondansetron (Zofran) for nausea and vomiting induced by cytotoxic chemo - therapy and radiotherapy.

Cerebral metastases

Common symptoms are headache and nausea. Consider corticosteroid therapy (e.g. dexamethasone 4 - 16 mg daily). Analgesics and antiemetics such as haloperidol are effective.

Paraplegia

Paraplegia is especially prone to occur with carcinoma of the prostate, even when treated with LHRH analogues. The warning signs are the development of new back pain, paraesthesia in limbs or the recent development of urinary retention.[1] The objective is to prevent paraplegia developing. High-dose corticosteroids are given while arranging urgent hospital admission.

Hiccoughs[7, 8]

Try a starting dose of chlorpromazine 25 mg tds or 25 mg IM as bolus *or* haloperidol 2.5 mg bd

Swallowing granulated sugar with or without vinegar does not appear to be effective. Other drugs reported to be beneficial include baclofen, midazolam, clonazepam, nifedipine and metoclopramide.

Weakness and weight loss

This problem may be assisted by a high-calorie and high-protein diet. A list of high-energy drink supplements is provided in Palliative Care: *The Nitty Gritty Handbook*.[1]

Examples

Banana Sustagen milk

• milk	2 cups
• banana	one
• egg	one
• Sustagen powder	3 dessertspoons
• skim milk powder	1 dessertspoon
• Glucodin	1 dessertspoon
• Ice	crushed
(Vitamise all together.)	

Egg flip

• egg	one
• milk	one cup
• vanilla syrup or essence	to taste (13 drops)
• sugar	1 teaspoon
• brandy	(optional)

(Vitamise all together, strain, sprinkle with nutmeg.)

High-energy cordial

• cordial	1 tablespoon
• Glucodin	1 teaspoon
• water	1 cup
• ice	crushed

(Blend cordial and Glucodin till smooth, stir in water.)

High-energy juice

• juice	1 cup
• Glucodin	1 dessertspoon

(Blend Glucodin with a little juice till smooth. Stir in remaining juice.)

The AIDS patient

The same principles of management apply to the person suffering from the many manifestations of terminal AIDS. Many of these patients wish to die at home and there are excellent caring support groups to help. It is important to become acquainted with the service network. Because of opportunistic infections there are many challenges facing the palliative care of such patients and some management guidelines are included in Chapter 8. A very helpful guide to symptom control is presented in *Therapeutic Guidelines by* Mashford et al. (pp. 239 - 46).[7]

DYING AND GRIEVING

The stages of the grieving process as described by Kübler-Ross may be experienced by both the patient and the family, albeit not exactly according to the five stages. The grieving process following the death of a loved one can vary enormously but many people are devastated. The principles of care and counselling include:[1]

- Be available and be patient.
- Allow them to talk while you listen.
- Reassure them that their feelings are normal.
- Accept any show of anger passively.
- Avoid inappropriate reassurance.
- Encourage as much companionship as possible, if desired.

(See guidelines for crisis counselling given in Chapter 5.)

Communicating with the dying patient

Good communication is essential between the doctor and patient in order to inform, explain, encourage and show empathy. However, it can be very difficult, especially with the cancer patient.

Good communication is dependent on honesty and integrity in the relationship. Telling the truth can be painful and requires sensitivity, but it builds trust that enables optimal sharing of other difficult concerns and decisions, such as abandoning curative treatment, explaining the dying process and perhaps addressing thoughts on euthanasia.

Improved communication will lead not only to better ‘spiritual’ care but also to better symptom control.[1] Give patients every opportunity to talk about their illness and future expectations, and be available and patient in offering help and support.

Spiritual issues

Spirituality is an important issue for all people, especially when faced with inevitable death. Many people are innately spiritual or religious and those with deep faith and a belief in ‘paradise’ appear to cope better with the dying process. Others begin to reflect seriously about spirituality and search for a meaning for life in this situation. Carers, including the attending doctor, should be sensitive to their needs and turmoil and reach out a helping hand, which may simply involve contacting a minister of religion.

Spiritual care builds on patients' existing resources to enable them to rise above the physical, emotional and social effects of their terminal illness.[1]

The question of euthanasia

It should be an uncommon experience to be confronted with a request for the use of euthanasia,[3] especially as the media clichés of ‘extreme suffering’ and ‘agonising death’ are rarely encountered in the context of attentive whole person continuing care. The non-use of life support systems, the use of ‘round the clock’ morphine, cessation of cytotoxic drugs, the use of ancillary drugs such as antidepressants and antiemetics, various nerve blocks and loving attention almost always help the patient cope without undue pain and suffering.

Practice tips

- Consider prescribing antidepressants routinely for patients in pain.
- Remember the ‘sit down rule’ whereby the home visit is treated as a social visit - sitting down with the patient and family, having a ‘cuppa’ and sharing medical and social talk.[3]
- Early referral of terminal patients with difficult-to-control problems, especially pain, to a hospice or multidisciplinary team can enhance the quality of care. However, the patient's family doctor must still be the focus of the team.

REFERENCES

1. Fairbank E, Banks T. *Palliative Care: The Nitty Gritty Handbook*. Melbourne: RACGP Services Division, 1993: 1 - 18.
2. McGuckin R, Currow D, Redelman P. *Palliative care: your role*. Med Observer, 1992; 27 November: 41 - 2.
3. Carson NE, Miller C. *Care of the Terminally Ill*. Melbourne: Monash University, Department of Community Medicine Handbook, 1993: 107 - 15.
4. NICE: > http://www.nice.org.uk <
5. Kübler-Ross E. *On Death and Dying*. London: Tavistock, 1970.
6. Buchanan J et al. *Management of Pain in Cancer*. Melbourne: Sigma Clinical Review, 1991; 18: 8 - 10.
7. Mashford ML (Chair) et al. *Therapeutic Guidelines*: Palliative Care (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2005.
8. Woodruff R. *Palliative Medicine* (3rd edn). Melbourne: Oxford University Press, 1999.
9. Burke AL. Palliative care: an update on ‘terminal restlessness’. Med J Aust, 1997; 166: 39 - 42.
10. Twycross R. *Introducing Palliative Care*. Oxford: Radcliffe Medical Press, 1996: 147.

CHAPTER 8. THE ELDERLY PATIENT

Last scene of all, That ends this strange eventful history, Is second childishness, and mere
oblivion, Sans teeth, sans eyes, sans taste, sans everything

WILLIAM SHAKESPEARE (1564 - 1616), AS YOU LIKE IT

THE ELDERLY PATIENT

The ageing (over 65 years) are the fastest growing section of the Australian population. The number of 'old-old' (over 85 years) is increasing at an even faster rate.[1]

The over 65s in 2001 made up 12.7% of the Australian population (13.4% in the United States). It is expected that this group will make up 13.4% in 2011 and 20% in the year 2031. A similar trend is expected in the United States with 18% in 2040.[2]

The over 65s use twice the number of health services per head of population. They account for 25% of all hospital costs and 75% of all nursing home costs. They represent 25% of all general practice consultations.[1] Many are affected by multisystem disease. All are affected to a greater or lesser extent by the normal physiological changes of organ ageing.

Figure 8.1 Establishing rapport and support through the home visit to the elderly patient is an important security gesture

Ageing is characterised by the following:[1]

- decrease in metabolic mass
- reduction in the functional capacity of organs
- reduced capacity to adapt to stress
- increased vulnerability to disease
- increased probability of death

Age-associated deterioration occurs with hearing, vision, glucose tolerance, systolic blood pressure, kidney function, pulmonary function, immune function, bone density, cognitive function, mastication and bladder function.

One of the main contributing factors is the problem of disuse. Encourage exercise, especially walking and water aerobics.

AGEING AND DISEASE

Degenerative cardiovascular disease emerges with ageing according to the following approximate guidelines:

- aged 40 - obesity
- 50 - diabetes
- 55 - ischaemic heart disease
- 65 - myocardial infarction
- 70 - cardiac arrhythmias
- 75 - heart failure
- 80 - cerebrovascular accidents

Deterioration in health and the 'masquerades'

Unexpected illness, including mental confusion (one of the major hallmarks of disease in the elderly), can be caused commonly by any of the so-called masquerades outlined in Chapter 18:

- depression
- drugs, including alcohol, anticholinergics
- diabetes mellitus
- anaemia
- thyroid disease
- urinary tract infection
- neurological dilemmas
    - Parkinson's disease
    - cerebrovascular accident
- infections (e.g. bronchopneumonia)
- neoplasia
- giant cell arteritis/polymyalgia rheumatica

Common significant management disorders encountered in the elderly include:

- hypertension
- ischaemic heart disease and heart failure
- depression
- diabetes (type 2)
- dementia
- social and physical isolation
- osteoarthritis
- disorders of the prostate
- urinary incontinence
- locomotive (lower limb) disorders
    - neurological
        - peripheral neuropathy
        - ataxia
    - claudication due to vascular insufficiency
    - other peripheral vascular disease
    - claudication due to spinal canal stenosis
    - sciatica/nerve root paresis
    - osteoarthritis: hips, knees, feet
    - foot disorders (e.g. ingrown toenails)
    - leg ulceration

Important problems affecting the elderly are presented in Figure 8.2.


CVA
• Parkinson's disease
• dementia
• depression
Eye disorders
including
• loss of visual acuity
• cataracts
• glaucoma
• macular degeneration
• xerophthalmia
Xerostomia
Polymyalgia rheumatica
Lung
• asthma
• chronic bronchitis
• COPD
• infection
Gastrointestinal
• constipation
• incontinence
• faecal impaction
Osteoarthritis
Hip fracture
Locomotive (lower limb) disorders
Ear disorders
• deafness
• tinnitus
Temporal arteritis
Thyroid
• hypothyroidism
• hyperthyroidism
Cardiovascular
• ischaemic heart disease
• arrhythmias
• heart failure
• hypertension
• hypotension
Kidney failure
Diabetes mellitus
especially type 2
Spinal problems
• spondylosis
• spinal canal stenosis
• sciatica
• osteoporosis
Genitourinary
• urinary dysfunction
• urinary tract infection
• prostate disorders
• incontinence of urine
• impotence
General
• drugs/medication
• anaemia
• isolation

Figure 8.2 Significant problems affecting the elderly

The 'classic' triad[3]

Be mindful of the classic triad:


confusion
falls
incontinence

This can represent non-specific signs of acute illness, particularly infection. Aggressive antibiotic treatment is worth considering while awaiting the results of culture or clinical developments.

CHANGES IN SENSORY THRESHOLDS AND HOMEOSTASIS

A clinically significant feature in some elderly patients is the raising of the pain threshold and changes in homeostatic mechanisms, such as temperature control. Consequently, these patients may have an abnormal response to diseases such as appendicitis, pyelonephritis, internal abscess, pneumonia and septicaemia. There may be no complaints of pain and no significant fever but simply general malaise and abnormal behaviour, such as delirium, agitation and restlessness.

ESTABLISHING RAPPORT WITH THE ELDERLY PATIENT

The elderly patient especially requires considerable support, understanding, caring and attention from a GP who can instil confidence and security in a patient who is likely to be lonely, insecure and fragile. This means taking time, showing a genuine interest and a modicum of humour, and always leaving detailed instructions.

One of the best ways to generate a good relationship is through home visitation. The value of home visits can be considered under the concepts of the RACGP.[4]

1. Assessment, both initial and continuing: 'You don't know your patient until you've visited their home'.
2. Continuing care:
    - security to the patient
    - support for 'caring' family
    - effective monitoring/intervention role
    - effective liaison with patient/family
    - checking medication

    Home visits can be considered in three categories:

    a. an 'unexpected' visit (especially to a new patient)
    b. a patient-initiated but routine request for a 'check-up and tablets'
    c. the regular call - usually 2 to 4 weeks

These home visits are a 'security gesture' to the patient - evidence that they are supported in their desire to remain independent for as long as possible in their own home. They strengthen the patient - doctor relationship as a position of trust, which is of special importance to frail, elderly people feeling increasingly insecure and threatened.

If the patient is being supported by a spouse or relative, the doctor can provide continuous reassurance and support to all concerned as well as continual assessment, both physical and mental. Finally, the home visit may become part of the terminal care of a dying patient, something that is very important to the elderly patient. Home visits can enhance the quality of life, both physical and mental, of an ageing person.

LONELINESS IN THE ELDERLY

Forbes points out that at least one in three elderly people feel lonely.5 It is more likely to affect the 'oldold', widows and widowers, and those affected by disability. They may tend to stay indoors, troubled by depression, agarophobia, social phobia, sensory impairment or incontinence of urine or faeces.

Possible signs of loneliness include:

- verbal outpouring
- drab clothing
- dependence on television
- body language with a 'defeated' demeanor
- prolongation of visit including holding on to one's hand

DOCTOR BEHAVIOUR THAT CAN IRRITATE AND CONFUSE ELDERLY PATIENTS

Ellard[6] has written a most interesting paper, based on a compendium of complaints by the elderly about their doctors, on behaviour that upsets the elderly:

- having a consulting room with slippery steps, poor lighting and inadequate handrails
- non-attention to simple courtesies by receptionist staff
- keeping them waiting
- having low soft chairs in the waiting room and surgery
- being overfamiliar, with addresses such as 'Pop' or first names for elderly females

- shouting at them on the assumption that they are deaf
- appearing rushed and keen to get the consultation over quickly
- forgetting their psychosocial problems and concentrating only on their physical problems (i.e. not treating the whole person)
- forgetting that they have several things wrong with them and using a different priority list from theirs
- being unaware that they may have seen other practitioners or may be taking additional medication
- failing to ask patients to give their understanding of what is wrong
- omitting to give printed patient education handouts about their problems and medications
- omitting to explain how the medication will work
- treating them as though they would have little comprehension of their health and treatment
- failing to respect their privacy, such as not knocking before entering the examination room
- failing to provide appropriate advice on various social services such as meals on wheels and other support groups
- failing to reevaluate carefully their health and medication
- failing to take steps to reverse any deterioration in their health, including reluctance to refer.

ASSESSMENT OF THE ELDERLY PATIENT

The initial consultation should include a thorough clinical examination on the traditional lines of history, physical examination and selective laboratory investigation. At regular intervals during continuing care this careful assessment may need to be repeated.

A golden rule

You don't know your patient until you've visited them at home.

History

The medical history may be difficult to obtain and the help of a family member is recommended. The use of questionnaires, which can be completed at leisure at home with the help of family members, is most helpful as complementary to the medical interview.

Important specific areas to focus on are:

- previous medical history and hospitalisation
- immunisation status
- medications and OTC drugs
- alcohol intake, smoking
- problem list of complaints
- dependence on others
- members of household
- household problems
- comforts: heating, cooling, bedding, etc.
- ambulation/mobility
- meals: diet
- hygiene: bathing
- toileting: continence
- teeth: condition, ? dentures
- vision
- hearing (always ask about this)
- systems review, especially:
  - genitourinary function

  - gastrointestinal
  - cardiorespiratory
- locomotion, including feet
- nervous system, ? falls, giddiness, faints
- emotional and mental health
- evidence of depression
- history of bereavement
- history of abuse by carers, family members
- financial/insurance status

A thorough family history and psychosocial history is of prime importance. At all times concentrate on a general assessment of the patients' ability to communicate by evaluating mental status, comprehension, hearing, vision, mood and speech.

Physical examination

The routine for the physical examination is similar to that of the younger adult but certain areas require more attention. The elderly patient expects to be examined adequately (especially having the blood pressure measured) but requires appropriate dignity. It is recommended that the practice nurse supervise dressing and undressing and prepares the patient for examination.

The following areas should be examined.

Practice nurse

- Prepares for examination.
- Helps with questionnaire.
- Records weight and height.
- Takes temperature, pulse and respiration.
- Checks audiometry (if hearing problem).
- Checks ocular tension (if appropriate).
- Prepares cervical smear tray for female patient (if relevant).

Doctor

- general appearance, including skin, hair and face (evaluate nutritional status)
- mental status examination (Figure 8.3)[7]



Mini mental state examination

ASK the patient	*INSTRUCTIONS for assessor*	*Score*
ORIENTATION		
What is the year, month, day and date? What is the season?	Ask specifically for omitted details, e.g. month, year Score 1 point for each correct answer.	/5
Can you tell me where we are now?	Expect street no., road name, suburb, city, state. Ask in turn for each place (if necessary). Score 1 point for each correct answer.	/5
REGISTRATION		
'I am going to test your memory—I want you to repeat these 3 items after me, and hold them in your mind for when I ask you later.'	Name three unrelated objects (e.g. orange, camel, table) speaking clearly and slowly; repeat up to three times, subtracting 1 point each time. Repeat until all three are learnt.	/3
ATTENTION AND CALCULATION		
'Beginning at 100, count backwards by 7', or, if patient is unable to perform this task, ask: 'Spell the word "WORLD" backwards'	Stop after five answers; each correct answer scores one. Score equals number of letters before first mistake.	/5
RECALL		
'What are the three words I asked you before?'	Score 1 point for each correct answers.	/3
LANGUAGE		
'What is the name of this object?'	Show two objects. Point to a watch, and then a pen.	/2
'Please repeat this sentence "NO IFS, ANDS, OR BUTS".'		/1
Giving the patient a blank piece of paper, say: 'Take this paper in your right hand, fold it in half, and put it on the floor.'	Score 1 point for each step.	/3
PRAXIS (3 TASKS)		
1. 'Read and obey this task.'	**CLOSE YOUR EYES** Score only if patient closes eyes.	/1
2. WRITE A SENTENCE …………………………………… …………………………………… …………………………………… …………………………………… ……………………………………	Sentence must be sensible and contain a verb and subject.	/1
3. COPY THIS DESIGN	All 10 angles must be present with two intersecting (ignore tremor, scale and rotation).	/1
		/30
Scoring guide only: 18–24 (probable mild dementia); 10–17 (probable moderate impairment) < 10 (severe impairment). A decline in score of > 2 points over 1 year may be significant. *Note:* These scores are used as a baseline to determine the criteria for prescribing anti-dementia drugs		Max score

Figure 8.3 A practical adaptation of the mini-mental state examination. Adapted from M.F. Folstein, S.E. Folstein and P.R. McHugh, Mini-Mental State, J Psych Res, 1975: 12: 189

- eyes: visual acuity
- ears: simple hearing test; auroscopic examination
- oral cavity, including teeth and gums
- neck especially thyroid
- lungs: consider peak flow meter
- pulse and blood pressure (repeat)
- heart; breasts
- abdomen; hernial orifices
- spine
- lower limbs: joints; circulation; feet including nails
- gait
- men: rectal examination; scrotum and testes
- women: cervical smear if appropriate

'Rules of 7' for assessment of the non-coping elderly patient

If your elderly patient presents with non-specific symptoms, unexpected deterioration in health and/or an inability to cope with the activities of daily living, consider the following checklist in your assessment. Apart from confusion, other non-specific symptoms include drowsiness, poor concentration, apathy, fatigue/weakness/tiredness/lethargy, anorexia, nausea, weight loss, dyspnoea, immobility, 'stuck in bed or chair', stumbles or falls. It is also important to consider infections including pneumonia and the masquerades (see p. 161).

1 MENTAL STATE	? Confusion/dementia Depression

	Bereavement, incl. pets Elderly abuse/bullying
2 EYES	? Visual acuity Cataracts/glaucoma
3 EARS	? Deafness, e.g. wax Tinnitus
4 MOUTH	? Dentition Xerostomia Malnutrition
5 MEDICATION	? Polypharmacy Adverse reactions
6 BLADDER and BOWELS	? Incontinence Retention Urinary infection
7 LOCOMOTION	? Gait - antalgic - movement disorder, esp. Parkinson's Arthritis - hips/knees Back/sciatica Feet - nails - neuropathy Circulation Leg ulcers

The mini-mental status examination

Evidence of memory difficulty remains the best single indicator of dementia and should always be evaluated by formal memory testing. However, memory problems may be due to factors other than dementia, and demonstrating failure in other areas of cognitive functioning (language, spatial ability, reasoning) is necessary to confirm the diagnosis of dementia.[7] A number of screening tests are available but the minimental status examination (MMSE), particularly the Folstein MMSE depicted in Figure 8.3 is commonly used and is the recommended test to use.

Another somewhat simpler test is the ‘Quick 10-step cognitive impairment test’ (Table 8.1 at p. 62).[8, 9]

Table 8.1 The quick 10-step cognitive impairment test

Scoring: questions 1 to 8: correct - 0, incorrect - 2 questions 9 and 10: correct - 0, 1 error - 2, > 1 error - 4
1 When were you born?
2 What year is it?
3 What month is it?
4 What is the date today?
Remember the following address: *25 Main Street, Newcastle*
5 What is your telephone number? (or if no telephone) What is your street address?
6 What time is it (to nearest hour)?
7 Who is the prime minister of Australia?
8 What year did World War II end?
9 Count backwards from 20 to 1.
10 Repeat the memory test I gave you.
Evaluation: 0 - 8 not significant 9 - 12 probably significant 13 - 24 significant

Adapted from Hodkinson (1972)[8] and Kingshill (2000)[9]

The clock-face drawing test

This relatively simple test provides a ready qualitative screening test to differentiate normal elderly from patients with cognitive impairment, particularly dementia.[10]

Method

- Give the patient a blank A4-sized piece of paper.
- Ask the patient to draw the face of a round clock and put the numbers in their correct positions on the face of the clock to represent the hours. (Figure 8.4).



Figure 8.4 The clock-face drawing test

- Ask the patient to draw in the hands of the clock, indicating 10 minutes after 11 (or some other suitable time). (Figure 8.4).
- Repeat the instructions if the patient does not understand them.

Scoring system

- A closed circle is drawn - 3 points
- Numbers are in the correct position - 2 points
- All correct numbers are included - 2 points
- Clock hands are placed in the correct position - 2 points

The maximum score is 9. A low score indicates the need for further evaluation and does not establish the criteria for dementia, being a pointer only.

Laboratory investigations

The laboratory tests should be selected according to the evaluation of the patient and to costs versus potential benefits.

Recommended investigations for suspected dementia include:[11]

- kidney function
- hepatic function
- thyroid function
- full blood screen
- blood glucose
- serum electrolytes (especially if on diuretics)
- urinalysis
- serum vitamin B_{12} and folate
- syphilis serology
- chest X-ray
- neuroimaging: CT or MRI (preferably)
- Positron Emission Tomography (PET) scan or SPECT scan - for further information

Behavioural changes in the elderly

As GPs we are often called to assess abnormal behaviour in the elderly patient, with the question being asked, ‘Is it dementia?’ or ‘Is it Alzheimer's, doctor?’.

There are many other causes of behavioural changes in people over the age of 65 years and dementia must be regarded as a diagnosis of exclusion. The clinical presentation of some of these conditions can be virtually identical to early dementia.

The clinical features of early dementia include:

- poor recent memory
- impaired acquisition of new information
- mild anomia (cannot remember names)
- personality change (e.g. withdrawn, irritable)
- minimal visuospatial impairment (e.g. tripping easily)
- inability to perform sequential tasks

The differential diagnosis for behavioural changes apart from dementia include several other common and important problems (which must be excluded) and can be considered under a mnemonic for dementia.[11]

All these conditions should be considered with the onset of deterioration in health of the elderly person. Even apparently minor problems, such as the onset of deafness (e.g. wax in ears), visual deterioration (e.g. cataracts), diuretic therapy, poor mastication and diet, urinary tract intercurrent infection, boredom and anxiety can precipitate abnormal behaviour.

- **D** drugs and alcohol depression
- **E** ears
  eyes
- **M** metabolic, e.g. hyponatraemia diabetes mellitus hypothyroidism
- **E** emotional problems (e.g. loneliness)
- **N** nutrition: diet (e.g. vitamin B group deficiency, teeth problems)
- **T** tumours of CNS
  trauma of CNS
- **I** infection
- **A** arteriovascular disease → cerebral insufficiency

ELDER ABUSE

It is important to keep in mind the possibility of abuse of the elderly, especially where there is a family history of abuse of members. The issue is as important as child and spouse abuse. Over 1 million elderly people are estimated to be the victims of physical or psychological abuse each year in the United States.[3] We should keep in mind the occasional possibility of Munchausen's syndrome by proxy.

DEPRESSION AND DEMENTIA

The main differential diagnosis of dementia is depression, especially major depression, which is termed pseudodementia. The mode of onset is one way in which it may be possible to distinguish between depression and dementia. Dementia has a slow and surreptitious onset that is not clear-cut, while depression has a more definable and clear-cut onset that may be precipitated by a specific incident. Patients often have a past history of depression. Those with dementia have no insight while those with depression have insight, readily give up tasks, complain bitterly and become distressed by their inability to perform their normal enjoyable tasks.

In response to cognitive testing, the typical response of the depressed patient is ‘don't know’, while making an attempt with a near miss typifies the patient with dementia (see Table 8.2).

Table 8.2 Comparison of dementia and pseudodementia (commonly severe depression)

	Dementia	Pseudodementia
Onset	Insidious	Clear-cut, often acute
Course over 24 hours	Worse in evening or night	Worse in morning
Insight	Nil	Present
Orientation	Poor	Reasonable
Memory loss	Recent > remote	Recent ° remote
Responses to mistakes	Agitated	Gives up easily
Response to cognitive testing (questions)	Near miss! Difficulty understanding	‘Don't know’ Slow and reluctant but understands words (if co-operative)

It is vital to detect depression in the elderly as they are prone to suicide. ‘Nothing to look back on with pride and nothing to look forward to.’ The middleaged and elderly may not complain of depression, which can be masked. They may present with somatic symptoms or delusions.

Note that depression occurs commonly in people with dementia, especially in the early stages.

Dementia (chronic organic brain syndrome)

The incidence of dementia increases with age, affecting about 1 person in 10 over 65 years and 1 in 5 over 80 years. The important causes of dementia are:

- degenerative cerebral diseases, including
    - Alzheimer's disease (about 60%)
    - dementia of frontal type (up to 10%)
    - dementia with Lewy bodies (up to 10%)
- vascular (15%)
- alcohol excess (5%)

Note: Mixed dementia should be considered

Other causes of dementia:

- AIDS dementia
- cerebral tumours
- Cruetzfeldt-Jakob disease

- Pick's disease
- neurosyphilis

In Alzheimer's, there is insidious onset with initial forgetfulness progressing to severe memory loss. In frontal dementias the earliest manifestations are personality change and alteration of behaviour, including social dysfunction. Dementia with Lewy bodies is characterised by any two of visual hallucinations, spontaneous motor parkinsonism and fluctuations in the mental state. Vascular dementia usually starts suddenly and is accompanied by focal neurological signs with evidence of cerebrovascular disease.

The characteristic feature of dementia is impairment of memory. Abstract thinking, judgment, verbal fluency and the ability to perform complex tasks also become impaired. Personality may change, impulse control may be lost and personal care deteriorates.

Risk factors for dementia include:

- family history
- late onset depression
- hypothyroidism
- Down syndrome
- history of head injury
- HIV/AIDS
- generalised atherosclerosis
- Parkinson's disease

Differential diagnosis of dementia includes:

- normal cognitive impairment of ageing
- delirium
- major depression
- drug abuse
- amnestic disorder
- various medical conditions, e.g. anaemia, thyroid/endocrine disorders

The DSM-IV (TR) criteria for dementia are presented in Table 8.3 and clinical clues suggesting dementia in Table 8.4.

Table 8.3 DSM-IV (TR) criteria for dementia of Alzheimer's type (modified)

Diagnosis of dementia requires evidence of:
A1 Memory impairment
A2 At least one of the following cognitive disturbances:
• Language ° aphasia
• Motor actions ° apraxia
• Recognition ° agnosia
• Executive functioning (e.g. organising)
B Disturbance significantly interferes with social and work functions
C Gradual onset and continuing cognitive decline
D Not due to known organic causes (e.g. drugs, cerebrovascular disorders)
E Not due to delirium
F Not due to another axis 1 disorder (e.g. major depression)

(*a*) Classified with or without behavioural disturbance (*b*) Also with early onset < 65 years or late onset > 65 years

Source: Diagnostic and Statistical Manual (4th edn, revised). Washington, DC: American Psychiatric Association, 2000.

Table 8.4 Clinical clues suggesting dementia

1 Patient presentations
• new psychological problems in old age
• ill-defined and muddled complaints
• uncharacteristic behaviour
• relapse of physical disorders
• recurrent episodes of confusion
2 Problems noted by carers
• 'not themselves' - change in personality (e.g. humourless)
• domestic accidents, especially with cooking and heating
• unsafe driving
• false accusations
• emotional, irritable outbursts
• tendency to wander
• misplacing or losing items (e.g. keys, money, tablets, glasses)
• muddled on awakening at night
3 Mental state observations
• vague, rambling or disorganised conversation
• difficulty dating or sequencing past events
• repeating stock phrases or comments
• playing down obvious, perhaps serious, problems
• deflecting or evading memory testing

Source: After McLean[7]

The many guises of dementia can be considered in terms of four major symptom groups:[7]

1. Deficit presentations - due to loss of cognitive abilities, including:
    - forgetfulness
    - confusion and restlessness
    - apathy (usually a late change)
    - self-neglect with no insight
    - poor powers of reasoning and understanding
2. Unsociable presentations - based on personality change, including:
    - uninhibited behaviour
    - risk taking and impulsive behaviour
    - suspicious manner
    - withdrawn behaviour
3. Dysphoric presentations - based on disturbed mood and personal distress, including:

    - depression (hopeless and helpless)
    - irritability with emotional outbursts
    - lack of co-operation
    - insecurity

4. Disruptive presentations - causing distress and disturbance to others, including:
    - aggressive, sometimes violent, behaviour
    - agitation with restlessness

The problem occasionally results in marked emotional and physical instability. It is sad and difficult for relatives to watch their loved ones develop aggressive and antisocial behaviour, such as poor table manners, poor personal cleanliness, rudeness and a lack of interest in others. Sometimes severe problems, such as violent behaviour, sexual promiscuity and incontinence, will eventuate.

There is always the likelihood of accidents with household items such as fire, gas, kitchen knives and hot water. Accidents at the toilet, in the bath and crossing roads may be a problem, especially if combined with failing sight and hearing. Such people should not drive motor vehicles.

Without proper supervision they are likely to eat poorly, neglect their bodies and develop medical problems, such as skin ulcers and infections. They can also suffer from malnutrition and incontinence of urine or faeces.

Management of suspected dementia

Exclude reversible or arrestable causes of dementia:

- full medical history (including drug and alcohol intake)
- mental state examination
- physical examination
- investigations (see p. 488)

Management of dementia

There is currently no cure for dementia - the best that can be offered to the patient is tender loving care.

Education, support and advice should be given to both patient and family. Multidisciplinary evaluation and assistance are needed. Regular home visits by caring sympathetic people are important. Such people include relatives, friends, general practitioners, district nurses, home help, members of a dementia self-help group, religious ministers and meals on wheels. The sufferers tend to manage much better in the familiar surroundings of their own home and this assists in preventing behaviour disturbance.

Special attention should be paid to organising memory aids, such as lists, routines and medication, and to hygiene, diet and warmth. Adequate nutrition, including vitamin supplements if necessary, has been shown to help.

Driving

Driving is a problem especially as many are reluctant to give up their licence. Those with mild dementia are more likely to cause road accidents. In some states it is compulsory for doctors to report patients who are unfit to drive. If uncertainties arise or a patient is recalcitrant refer to the local Road Traffic Authority. In Sweden it is recommended that those with moderate to severe dementia should not drive.

Co-morbidity/associated problems[12]

Depression can occur early in dementia and requires intervention. Demented patients are vulnerable to superimposed delirium which is often due to:

- urinary tract infection
- other febrile illness
- prescribed medication

- drug withdrawal

Delirium should be suspected if a stable patient becomes acutely disturbed.

Dementia and Parkinson's disease

This is a very difficult yet common problem. One problem is that medication affects the mental processes. The choice of drugs is critical to care, so referral to an experienced team which can provide a good neuro psychiatric assessment is advisable. The best option appears to be the administration of:

- Levo-dopa to maximum dose
- Quetiapine at night

Medication[12]

Demented patients often do not require any psychotropic medication. Antidepressant drugs can be prescribed for depression. The cholinesterase inhibitors donepezil, galantamine, and rivastigmine appear to delay progression of dementia to a modest extent only.

Available drugs for Alzheimer's disease

Cholinesterase inhibitors

- donepezil 5 mg (o) nocte for 4 weeks, increase to 10 mg nocte as tolerated
- galantamine 4 mg (o) bd for 4 weeks, increase to 8 mg bd.
- rivastigmine 1.5 mg (o) bd for 2 weeks, increase gradually up to 6 mg bd

Aspartate (NMDA) antagonist

- memantine 5 mg (o) mane for 1 week → 5 mg bd week 2 → 10 mg bd

Based on double-blind randomised trials of the two drugs donepezil and rivastigmine and using Cochrane data,[12, 13] the following points emerge:

- only modest improvement overall
- greatest improvement with higher doses
- higher doses less well tolerated
- long-term efficacy unknown
- clinical effectiveness in severe disease has not been demonstrated.[14]

The newer agent memantine appears to have similar outcomes and can be used in combination.

Using evidence-based medicine criteria on numbers needed to treat (p. 137), the evidence shows that 13 patients must be treated with rivastigmine 6 - 12 mg/day for 6 months for one patient to display clinically meaningful improvement.[14, 15]

To control psychotic symptoms or disturbed behaviour probably due to psychosis:[12]

- olanzapine 2.5 - 10 mg (o) daily *or*
- risperidone 0.5 - 2 mg (o) daily *or*
- haloperidol 0.5 mg (o) nocte up to 2 mg bd

To control symptoms of anxiety and agitation use:

> oxazepam 15 mg (o) 1 to 4 times daily but benzodiazepines should be used only for short periods (maximum 2 weeks) as they tend to exacerbate cognitive impairment in dementia.

Complementary therapy

As yet there is insufficient evidence for the efficacy ofcomplementary medicines such as *Ginkgo biloba*,[16] vitamin E[17] and other antioxidants in treating (alleviating symptoms in) dementia, despite some epidemiological evidence, especially in the case of vitamin E. However, we should encourage our patients in preventive healthy lifestyle strategies, such as optimal nutrition rich in essential vitamins and exercise.

LATE LIFE DEPRESSION AND SUICIDE

The risk of suicide increases with age in males (Figure 20.1 on p. 191). The risk factors for late life suicide are:[18]

- male
- single
- recent bereavement
- social isolation
- recent relocation
- poor pain control
- feeling helpless/hopeless
- anhedonia
- indicating a wish to die
- recent alcohol abuse

The principles of treatment include supportive care, regular visits, CAT team, interpersonal psychotherapy, cognitive behavioural therapy and family support/interventions.

PARAPHRENIA

Paraphrenia is that condition where the symptoms and signs of paranoid psychosis appear for the first time in the elderly. In this non-psychotic mental illness, the patient, who is usually an elderly female, presents with paranoid delusions, such as a feeling of being watched or persecuted. These are usually associated with visual and hearing problems.

FALLS IN THE ELDERLY

Falls in the elderly are a major problem as 30% of people over the age of 65 experience at least one fall per year, with 1 in 4 of these having significant injury. About 5% of falls result in fracture.[19] In 2002 (in Australia) there were approximately 1200 deaths in people over 75 years following falls.

The main causes are:

- neurological (e.g. cerebrovascular disease)
- sensory impairment (e.g. visual, vestibular)
- cardiovascular (e.g. postural hypotension)
- musculoskeletal (e.g. arthritis, foot disorders)
- cognitive and psychological conditions (e.g. dementia, delirium)
- medication/drug related (e.g. sedatives, alcohol)
- physiological changes (e.g. gait disorders)
- environmental factors (e.g. slipping or tripping)
- combinations of the above

The most significant clinical risk factors for falls have been shown to be visual impairment, impaired general function, postural hypotension, hearing impairment, low morale/depression, drug usage, especially sedatives, decreased lower limb strength including arthritis, and impaired balance and gait.

Assessment

The history should embrace the above causes and risk factors. In particular, a description of the fall by a witness, the perceived dysfunction at the time of the fall and whether there was loss of consiousness is particularly important.

The physical examination should include cardiac function, neurological status, including the minimental function test, and the musculoskeletal system, including assessment of gait. The 'get up and go' test (Table 8.5) is useful.

Table 8.5 The 'get up and go' test: a brief test of postural competence

1 Get up from chair without use of arms.
2 Observe normal gait and 360° turn.
3 Carry out the Romberg test (slight push with eyes closed).
4 Observe tandem walking (heel toe, straight line).

As special investigations (especially for those proving difficult to evaluate) consider full blood examination and ESR, blood sugar, urea and electrolytes, thyroid function tests, cardiac investigations (e.g. ambulatory ECG monitoring, ambulatory blood pressure monitoring), vestibular function testing, and CT scans or MRI scans.[20]

Management and prevention

Steps should be taken to correct any medical disorders and risk factors. It is appropriate to refer to a multidisciplinary team, including occupational therapists and physiotherapists. Assessment of circumstances in the home is very helpful. This may lead to reducing environmental hazards and providing walking aids. Exercise training and medication reduction are also important strategies.

PRESCRIBING AND ADVERSE DRUG REACTIONS

Ageing is associated with increased rates of adversedrug reactions.[1] The rate of adverse drug reactions for a single medication rises from about 6% at age 20 years to about 20% at age 70 years.

For less than six medications taken concurrently the rate of adverse drug reactions is about 6%. For greater than six medications taken concurrently the rate of adverse drug reactions jumps to 20%.[1] Approximately 15% of elderly patients admitted to hospital are suffering adverse drug reactions. Most adverse drug reactions are type A (dose related) rather than type B (idiosyncratic).[21]

Factors predisposing to adverse drug reactions in the elderly[1]

Most adverse drug reactions in the elderly are entirely predictable. Most are merely an extension of the pharmacological action of the drug (e.g. all antihypertensives will reduce blood pressure and have the capacity to cause hypotension and falls in a person with impaired baroreceptor function or poor homeostasis in their vascular tree). Very few adverse reactions are idiosyncratic or unexpected.

The five mechanisms of adverse drug reactions in the elderly are:

1. *Drug - drug interaction*
   For example, beta-blockers given concomitantly with digoxin increases the risk of heart block and bradycardia. Alcohol used in combination with antidepressants increases the risk of sedation.
2. *Drug - disease interaction*
   For example, in the presence of kidney impairment, tetracyclines carry an increased risk of kidney deterioration.
3. *Age-related changes leading to increases in drug plasma concentration*
   Decreased kidney excretion can extend the halflife of medication, leading to accumulation and toxicity.
4. *Age-related changes leading to increased drug sensitivity*
   For example, there is some suggestion that the pharmacological response to warfarin, narcotics and

benzodiazepines is increased in the elderly.
Conversely, the pharmacological response to insulin, theophylline and beta-adrenergic blocking agents is thought to be decreased.

5. *Patient error*
   Multiple medications can lead to patient error. The incidence and prevalence of dementia also increases with age. Other problems include failing eyesight and reduced manual dexterity.

Risk factors predisposing to medicine-related problems

Adverse drug effects in older people are influenced by multiple factors, with many exposed to more than one of these factors at a given time. Furthermore, their recovery from serious incidents such as a hip fracture is jeopardised, possibly ending in death.

Increasing the number of simultaneous medications increases the risk for all five mechanisms of adverse drug reactions.

In a study on adverse drug reaction in elderly patients the drugs most frequently causing admission to hospital were:[22]

- digoxin
- diuretics
- antihypertensives (including beta-blockers)
- psychotropics and hypnotics
- analgesics and NSAIDs

The same study showed that drugs regularly prescribed without revision were:

- barbiturates
- benzodiazepines
- antidepressants
- antihypertensives
- beta-blockers
- digoxin
- diuretics

Drug regimens should be kept as simple as possible to aid compliance and avoid or minimise drug interactions.

The elderly may need much lower doses of anxiolytics and hypnotics than younger patients to produce the same effect, thus rendering them more susceptible to adverse effects and toxicity. The elderly are especially liable to accumulate the longer-acting benzodiazepines.

In particular, any drug or combination of drugs with anticholinergic properties (e.g. tricyclic antidepressants, antiparkinsonian agents, antihistamines, phenothiazines and some cold remedies) can precipitate a central anticholinergic syndrome.[7]

The elderly are very prone to adverse effects to most of the more potent drugs, especially those for cardiac dysfunction and hypertension (Table 8.6). Both ACE inhibitors and calcium-channel blockers have been shown to produce a greater fall in blood pressure in elderly compared with younger subjects, presumably related also to a reduced homeostatic response.[21, 23]

Table 8.6 Risk factors predisposing to medicine related problems[21]

• 5 or more regular medicines
• >12 doses of medicine per day
• attending several different doctors
• significant recent treatment changes

- drugs with narrow therapeutic window
- drugs requiring monitoring e.g. warfarin
- non-compliance
- confusion/dementia
- language/literacy problems
- recent discharge from hospital or facility

Starting medications[12]

The starting dose of a drug in the aged[7] should be at the lower end of recommended ranges. Dosage increments should be gradual and reviewed regularly.

That is, start low, go slow and monitor frequently. It is important to individualise doses for the elderly with the simplest dosage regimen.

Minimising medication problems

- Write simple instructions on all prescriptions.
- Individualise doses for the elderly

Table 8.7 Drug side-effects in the elderly: common presentations[22]

Drug	Side-effects
Benzodiazepines	Confusion, falls, psychomotor impairment
β-blockers	Confusion, falls, asthma, insomnia
Cimetidine	Confusion
Digoxin	Nausea, confusion
Diuretics	Incontinence, falls, hyponatraemia, hypokalaemia
Levodopa	Confusion, falls, dystonia, hallucinations, agitation, postural hypotension
Metoclopramide	Confusion, extrapyramidal symptoms
Narcotic analgesics	Constipation, confusion
NSAIDs	Confusion, GIT bleeding, oedema, kidney dysfunction, headache
Phenothiazines	Confusion, postural hypotension, falls, constipation
Phenytoin	Confusion, falls, ataxia, parkinsonism, urinary problems
Prazosin	Postural hypotension, incontinence
SSRI antidepressants	Nausea, agitation, insomnia
Theophylline	Nausea, tremor, confusion
Tricyclic antidepressants	Confusion, falls, postural hypotension, constipation, urinary problems, eye problems
Verapamil	Constipation

- Give patients a list of their medications.
- Ask them to bring the list and their medications at each visit.
- Update this list as necessary.
- Keep medication regimen as simple as possible.

- Consider a Dosette box or Webster pack for polypharmacy.
- At home visits use the opportunity to inspect medications.
- Observe for drug interactions and toxicity.
- Keep detailed records of all medications prescribed.
- Carefully review medication after discharge from hospital.

Other tips for medication in the elderly

- Obtain relatives/carers' consent for medication, especially in the confused.

REFERENCES

1. Harris E. *Prescribing for the Ageing Population*. Update Course Proceedings Handbook. Melbourne: Monash University Medical School, 1992.
2. Australian Institute of Health and Welfare Canberra: Australian Government, Australia's Health 2004.
3. Mold JW. Principles of geriatric care. American Health Consultants. Primary Care Rep, 1996: 2(1): 2 - 9.
4. Lang D. Home visits to the elderly. Aust Fam Physician, 1993; 22: 264.
5. Forbes A. Caring for older people: loneliness. BMJ, 1996; 313: 352 - 4.
6. Ellard J. How to irritate and confuse your elderly patients - 20 simple rules. Mod Med Aust, 1990; 7: 66 - 8.
7. McLean S. Is it dementia? Aust Fam Physician, 1992; 21: 1762 - 76.
8. Hodkinson HM. Evaluation of a mental test score for assessment of mental impairment in the elderly. BMJ, 1972; 1: 233 - 8.
9. Kingshill 2000 < www.Kingshill-research.org >
10. Fredman M, Leach L, Kaplan E et al. *Clock Drawing - a Neuropsychological Analyses*. New York: Oxford University Press, 1994.
11. Turnbull JM. *Dementia and Depression in the Elderly*. California Medical Association: Audio-Digest Family Practice, 1993; 41: 10.
12. Bochner F (Chair). *Therapeutic Guidelines: Psychotropic* (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2000: 94 - 7.
13. Birks JS et al. Donepezil for mild and moderate Alzheimer's disease (Cochrane Review). In: The Cochrane Library. Issue 1, 2001. Oxford: Update Soft ware.
14. Birks JS et al. Rivastigmine for Alzheimer's disease (Cochrane Review). In: The Cochrane Library. Issue 1, 2001. Oxford: Update Soft ware.
15. New Alzheimer's drugs show only modest benefit. NPS News, 2001; 16: 1 - 6.
16. Le Bars PL et al. *Ginkgo biloba* and dementia. JAMA, 1997; 278: 1327 - 32.
17. Tabet N et al. Vitamin E for Alzheimer's disease (Cochrane Review). In: The Cochrane Library. Issue 1, 2001. Oxford: Update Soft ware.
18. Jeffreys D. Late-life depression. Medical Observer, 18 July 2003; 36 - 7.
19. Hindmarsh JJ, Estes H. Falls in older persons: causes and intervention. Arch Intern Med, 1989; 149: 2217.
20. Quail GG. An approach to the assessment of falls in the elderly. Aust Fam Physician, 1994; 23: 873 - 83.
21. NPS News. Medicines and older people: an accident waiting to happen? NPS, 2004; 34: 1 - 4.
22. Briant RH. Medication problems of old age. Patient Management, 1988; 5: 27 - 31.
23. Scharf S, Christophidis N. Drug treatment in the elderly. In: *MIMS Disease Index* (2nd edition). Sydney: IMS Publishing, 1996; 159 - 60.

CHAPTER 9. PREVENTION IN GENERAL PRACTICE

When meditating over a disease, I never think of finding a remedy for it, but, instead, a means of preventing it.

LOUIS PASTEUR 1884

DEFINITIONS[1]

Prevention may be defined as the means of promoting and maintaining health or averting illness.

It is concerned with removal or reduction of risks, early diagnosis, early treatment, limitation of complications, including those of iatrogenic origin, and maximum adaptation to disability.

The promotion of health concerns helping well people to learn healthy behaviours and to accept responsibility for their own well-being.

A preventive attitude implies that the doctor understands and can utilise the preventive potential in each primary care consultation by an ‘opportunistic approach’. In addition to the traditional management of both presenting and continuing problems, the doctor takes the opportunity to modify the patient's healthseeking behaviour, to provide education about the illness, and to promote health by relating the patient's present condition to previous unhealthy behaviour.

Primary prevention

Primary prevention includes action taken to avert the occurrence of disease. As a result there is no disease. Primary preventive strategies include:

1. education to bring about changes in lifestyle factors known to be associated with diseases (e.g. smoking cessation, healthy balanced diets, reduction in alcohol intake, exercise)
2. sterilisation of surgical instruments and other medical equipment
3. eradication, as with vector control of mosquitoes to prevent malaria
4. immunisation against infective diseases
5. sanitation, keeping our water supplies clean and disposing efficiently of sewage and industrial wastes
6. legislation to ensure that some of these primary preventive measures are carried out

Secondary prevention

Secondary prevention includes actions taken to stop or delay the progression of disease.

The term is usually applied to measures for the detection of disease at its earliest stage, i.e. in the presymptomatic phase, so that treatment can be started before irreversible pathology is present. The early recognition of hypertension through routine testing (screening) of patients allows treatment during the presymptomatic phase of the illness process. Screening for cervical cancer allows the treatment of cervical dysplasia, a premalignant condition. Other examples include mammography and endoscopy for polyps of the large bowel.

Tertiary prevention

Tertiary prevention includes the management of established disease so as to minimise disability.

The term is usually applied to the rehabilitation process necessary to restore the patient to the best level of adaptation possible when there has been damage of an irreversible nature. A patient who has suffered a stroke because of hypertension may be restored to a useful lifestyle with appropriate rehabilitation.

Relationship between types of prevention

It can be seen that there is a clearer demarcation between primary and secondary prevention than between secondary and tertiary prevention, although the latter term is particularly useful in dealing with the elderly and the handicapped.

Conceptually, curative medicine falls within the definitions of secondary and tertiary prevention while public health measures are mainly concerned with primary prevention. Prevention is really wider than medical practice but because of the success of public health practices in the past, more attention is now being focused on prevention by doctors (see Figure 9.1).[2]


PRIMARY PREVENTION
SECONDARY PREVENTION
TERTIARY PREVENTION
Healthy individual (no disease)
Biological onset of disease
Clinical onset
cure
disability
death
Chronic disease
Risk factors
Asymptomatic signs
Symptoms and signs
Functional status
Education Health promotion Immunisation Prophylaxis
Screening Surveillance
Diagnosis and treatment
Rehabilitation Support Prophylaxis

Figure 9.1 The phases of prevention in relation to the natural history of disease

As GPs our role in prevention is twofold:

1. First, we can recognise the preventable factors that are involved in an illness process and determine appropriate interventions.
2. Second, we can act to implement the preventive measure. In cases where the responsibility rests with the individual or the community, doctors can support prevention through education, applying political pressure or working with community agencies.

THE PRACTICE OF PREVENTIVE MEDICINE BY THE DOCTOR

What is preventable?

The first step in the implementation of prevention is to define which specific diseases can be prevented and to what extent, given certain restraints such as human resources, technology and the cost to the community. All diseases have a potential pre ventability but it may be unrealistic to try to achieve this.

Diseases that can be prevented can be grouped according to their aetiology. They fall into the following broad categories:

- genetic disorders
- conditions occurring during pregnancy and the puerperium
- developmental disorders
- accidents
- infections
- addictions
- behavioural disorders
- occupational disorders
- premature vascular disease
- neoplasms
- handicap in the disabled
- certain ‘other’ diseases (e.g. diverticular disease)

Mortality is the only reliable index by which the outcome of preventive activities can be judged. Conditions can be ranked in importance as causes of premature death according to the ‘person years of life lost before 70 years’ as follows:[1]

Accidents, poisoning and violence	29%
Neoplasms	19%
Circulatory diseases	17%
Perinatal conditions	10%
Congenital conditions	7%

This gives quite a different perspective to prevention and explains why the efforts of public health authorities and practising doctors do not always coincide.

The interventions available to us in medical practice are as follows:

1. educational - health promotion, health education and illness education
2. screening
3. surveillance
4. interventional care - immunisation, behaviour modification and drug prophylaxis
5. rehabilitation

OPTIMAL OPPORTUNITIES FOR PREVENTION

Primary prevention par excellence can be practised in general practice under the opportunities provided by the following clinical circumstances:

- antenatal care
- postnatal care
- advising people travelling overseas
- visits by infants with their parents
- times of crisis or potential crisis
- the premarital check-up

The Royal College of General Practitioners (UK) has identified the seven most important opportunities for prevention as:

1. family planning
2. antenatal care
3. immunisation
4. fostering the bonds between mother and child
5. discouragement of smoking
6. detection and management of raised blood pressure
7. helping the bereaved

MORTALITY AND MORBIDITY CONSIDERATIONS

An understanding of the mortality and morbidity patterns in the modern human being is essential to the planning of preventive programs. The great infectious diseases of the past, such as tuberculosis, syphilis, smallpox, influenza, diphtheria and streptococcal infections, have been largely contained but other diseases have become prominent as life expectancy increases. The great modern diseases are atherosclerosis (hardening of the arteries), malignant disease (cancer), HIV infection and iatrogenesis (doctor-induced illness).

These diseases and the common causes of mortality (Table 9.1) act as a focus for our energies in addressing preventive programs.

Table 9.1 Common causes of deaths in Australia in 2004

Circulatory disease Ischaemic heart disease 18.5% Cerebrovascular disease 9% Other vascular 3%	30.5%
Malignant disease	29%
Accidents, poisoning and violence	6%
Respiratory disease	5%
Suicide	2%

Source: Australian Bureau of Statistics, *Year Book Australia, 2004*.

It is worth focusing on the changes in disease indices during the past generation in order to evaluate the effect of preventive and health promotion programs during this period (Table 9.2).[3] The messages are to harness and promote with renewed vigour those strategies that are working, such as prevention of death from coronary artery disease and motor vehicle accidents, and to reevaluate those important areas, such as Aboriginal mortality, HIV infection, cancer, suicide and asthma, which are bad news!

Table 9.2 Apparent changes in disease indices 1960 - 1988

Improvements	No change or deterioration
Overall mortality	Aboriginal mortality (minor improvement)
Heart attack	
Stroke	Cancers
Unintentional injuries/poisonings	- pancreas
	- breast
- motor vehicle	- skin
- drownings	- lung (women)
- falls	- prostate
- occupational injury	AIDS
Most infectious diseases	Alcohol-related diseases
Cancer of lung (men)	Drug abuse
Cervical cancer (women)	Asthma
Cancer of stomach	Senile dementia
Unintentional pregnancies	STIs
Violence/homicide	Suicide in the young
Pregnancy complications	Cirrhosis
Congenital abnormalities	Musculoskeletal disorders
Dental health	
Perinatal complications	
Chronic bronchitis	

Source: G. Egger, R. Spark and J. Lawson, *Health Promotion Strategies and Methods*, Sydney, McGraw-Hill, 1990: 2 - 3.

A GLOBAL STRATEGY FOR GOOD HEALTH

The WHO defines good health as ‘a state of dynamic harmony between the body, mind and spirit of a person and the social and cultural influences which make up his or her environment’.

A considerable amount of epidemiological information has emerged to support what GPs have known for a long time - that a commonsense healthy lifestyle not only promotes good health but also reduces the risk of the main causes of mortality and morbidity in this country, including cardiovascular disease and cancer.

The common theme for virtually all disease is to follow the nutrition and lifestyle guidelines presented in Chapter 10.

Behaviour modification

Lifestyle habits that have developed over many years can be very difficult to change even when the individual is well motivated to change. There are a variety of instructional, motivational and behavioural techniques that can be used to initiate a lifestyle change program; GPs should be aware of these and use the resources of a multidisciplinary team to give support to motivated people who as a rule find behaviour modification difficult.

VASCULAR DISEASE

Risk factors for vascular disease (atherosclerosis) are:

- hypertension
- smoking
- high cholesterol
- diabetes
- obesity
- sedentary lifestyle
- stress
- alcohol excess
- poor diet
- family history

The guidelines for good health given on page 86, if followed, will help prevent the development of cardiovascular and cerebrovascular disease.

It is worth noting that the death rate from coronary heart disease is about 70% higher for smokers than for non-smokers and for very heavy smokers the risk is almost 200% higher. It has been shown that the incidence of heart disease falls in those who have ceased smoking.

Of particular interest is the UK study that showed high-dose vitamin E (400 - 800 IU daily) was cardioprotective and reduced the incidence of myocardial infarction in angina patients.[4] This has not been verified by further studies.

GPs can estimate the absolute 5 year risks of cardiovascular events in their patients by referring to the New Zealand Guidelines Group Cardiovascular Risk Tables (http://www.nzgg.org.nz)

The parameters used are:

- gender and age
- smoking status
- diabetes status
- blood pressure
- total cholesterol/HDL ratio

MALIGNANT DISEASE

Primary prevention of cancer is an important objective and there is a need to focus on this vital factor as much as on secondary prevention. The interesting statistics on the 5-year survival rate for specific cancers are shown in Table 9.3.

Table 9.3 Cancer prognosis: 5-year survival

	USA (whites)		Australia	
	1960 - 63	1983 - 88	1960	1982 - 97
Cancer	**%**	**%**	**%**	**%**
Lung	8	13	10	11
Bowel	40	58	35	55
Breast	63	79	60	82
Pancreas	1	3	2	5
Stomach	11	16	10	21

Source: Statistics provided by Dr Graham Giles, Anti Cancer Council of Victoria.

That environmental factors are involved in the aetiology of colorectal cancer and other cancers is indicated by wide variations in incidence between different countries.

Suspicion falls on diet and there is epidemiological evidence implicating diets high in animal fats and low in insoluble fibre, fruits and vegetables, and also high alcohol consumption. It is noted that there are higher incidence rates in people migrating from low-to high-risk countries, such as Japanese to Hawaii[5] and Greeks and Italians to Australia.[6]

Studies in the United States indicate that at least 35% of all cancer deaths are related to diet. Obese individuals have an increased risk of colon, breast and uterine cancers. High-fat diets are a risk factor for prostate, breast and colon cancers. Salt-cured, smoked and nitrate-cured foods increase the risk of upper GIT cancers. Foods rich in vitamin A and folate (dark green and deep yellow vegetables and fruits) and vitamin C and cruciferous vegetables (cabbage, brussels sprouts, broccoli and cauliflower) are all considered to have protective effects for various cancers.7 Photochemicals (plant chemicals) exist in these foods and in other vegetables and fruit that have a cancer-protective effect.[8]

Overall, diet, smoking, alcohol and occupational exposures (5%) appear to account for over 73% of all cancer mortality.[7]

Doll and Peto[9] consider that environmental factors are responsible for 80 - 90% of cancers and estimate that diet is a major factor in the cause of cancer in 40% of men and 60% of women.

The role of immunity in cancer

The development of a number of cancers appears to be related to a depression of the individual's immune system, particularly in relation to cellular immunity, in a similar way (albeit on a different scale) to the effect of HIV infection. Studies have shown that the immune system is adversely influenced by:[10]

- stress, especially bereavement
- depression
- ageing
- drugs
- pollutants
- cigarette smoke
- inappropriate diet
- alcohol
- radiation

On the other hand, a protective effect on the immune system may be provided by:

- food antioxidants (Table 10.2 on p. 84)
- tranquillity
- meditation

In some instances malignancies appear to undergo unpredictable remissions with patients following an optimal diet, taking antioxidants, changing their lifestyle and practising meditation. However, an Australian study indicated that the enthusiasm for the value of antioxidants may be unjustified (p. 84).

Diet certainly appears to be a most important factor in the primary prevention of disease. If immune deficient diseases can respond in such a way, imagine what a powerful primary preventive force such a lifestyle represents for all disease.

ASTHMA AND OTHER RESPIRATORY DISEASES

The death rate and morbidity rate for asthma and other respiratory diseases is unacceptable and much of it can be prevented.[11] A report on the cost of asthma claimed that there is evidence that a significant proportion of diagnosed asthmatics are currently receiving treatment that does not provide the best possible control of the disease.[11]

Prevention means being better informed and treating such an ‘irritable’ disease as bronchial asthma aggressively. It means focusing on better assessment and monitoring (e.g. home use of the mini peak flow meter), better delivery of medication to the airways (e.g. use of spacers attached to inhalers and/or use of pumps and nebulisers) and appropriate management of the cause (inflammation of the bronchial tree) by the use of inhaled corticosteroids or sodium cromoglycate as the first-line treatment for significant asthma. Following the six-step asthma management plan (Table 9.4) of the National Asthma Campaign would certainly promote prevention in this fickle disease.

Table 9.4 The six-step asthma management plan (National Asthma Campaign: Australia)

1 Establish the severity of the asthma.
2 Achieve best lung function.
3 Maintain best lung function - identify and avoid trigger factors.
4 Maintain best lung function - follow an optimal medication program.
5 Develop an action plan.
6 Educate and review regularly.

The protective effect for asthma and COPD of vitamin C, fish oils, a low-salt diet and other natural antioxidants is highlighted by Sridhar.[12]

PERIODIC HEALTHEXAMINATION

Since 86% of the population visit a general practitio ner at some stage of the year,3 and these people visit about five times each year (on average), GPs are in an ideal position to develop strategies for a periodic health examination. An emphasis should be placed on the history in addition to the physical examination and related basic investigations.

As for any smooth-running quality professional program, it is important to be organised with prepared practice staff, checklists and record systems. The RACGP has developed a College Record System, which has several leaflets covering all approaches to the patient ‘check-up’.[13]

The following guidelines for the periodic health examination are adapted from those recommended by the Preventive and Community Medicine Committee of the RACGP.13 This represents appropriate screening at the front line of primary health care.

Aims of screening

In practice, screening is not only to detect disease at its earliest stage, but also to find individuals at risk or those with established disease who are not receiving adequate care. There are three levels at which screening can be applied in general practice:

1. 'well' individuals with risk factors that predispose to disease (e.g. obesity, uncomplicated essential hypertension, hyperlipidaemia)
2. asymptomatic individuals with signs of early disease or illness (e.g. developmental dysplasia of the hip, ectopic testis, glaucoma, bacteriuria of pregnancy, carcinoma in situ of cervix)
3. symptomatic individuals whose irreversible abnormalities are unreported but the effects can be controlled or assisted (e.g. visual defects, deafness, mental handicap).

THE HISTORY[13]

An appropriate history will allow the recognition of certain risk factors that may foreshadow future disease. Though established patients will have a previously acquired database, their history should be reviewed and updated. It is recommended that the following items be included in history taking in the appropriate age groups.

Family history. In particular, cardiovascular disease, some cancers (breast, bowel, melanoma with dysplastic naevi), diabetes, asthma, genetic disorders and bowel disease will alert the doctor to specific risk factors (and psychological factors) for these patients.

Suicide and accidents. Consider the risk factors predisposing to suicide and accidents, which are the major preventable causes of death in children and young adults.

Substance abuse. Tobacco and alcohol are the major causes of preventable death in adults, although other drugs contribute to a lesser extent. Counselling by GPs, about smoking in particular, has been shown to be effective.

Exercise and nutrition. These factors have a role to play in preventing cardiovascular disease and to a lesser extent in blood pressure control, cancer, diabetes and constipation. They have an even greater role to play in improving general well-being and preventing morbidity.

Occupational health hazards. Consider these in working adults, as occupational health hazards can significantly contribute to morbidity and mortality (e.g. exposure to toxic substances, unsafe work practices). Specific examples include:

- coal miners - pneumoconiosis
- gold, copper and tin miners - silicosis
- asbestos workers and builders - asbestosis, mesothelioma
- veterinarians, farmers, abattoir workers - zoonoses
- aniline dye workers - bladder cancer
- health care providers - hepatitis B

Physical functioning, home conditions and social supports. Consider these in elderly people, as physical function and social supports are of crucial importance in determining whether they can care for themselves - intervention can prevent accidents and death.

Sexuality/contraception. Sexually transmitted infections are all preventable, as are unwanted pregnancies. Opportunities should be sought to ask young people, in particular, about their sexuality, and to counsel them. The question 'Do you have any concerns about sex?' is very useful in this context.

Osteoporosis. Osteoporosis affects nearly a third of all postmenopausal women, most of whom suffer osteoporotic fractures. Fractures of the femoral neck have a particularly poor prognosis, with up to a third of these women dying within 6 months, and many more requiring continuing nursing home care. Bone loss accelerates at the time of the menopause, and can be reduced by hormone replacement therapy.

Women at risk of osteoporosis are short, slim, Caucasian; they drink coffee and alcohol, smoke, eat a high-protein and high-salt diet, and don't exercise.

Masquerades in general practice. It is worth considering the ‘masquerades’ (Chapter 18, Tables 18.4 and 18.5 on pp. 161 and 162), which may present as undifferentiated illness, as a means of following the important medical principle of early detection of disease: engendering a certain awareness.

Primary masquerades to consider are:

- depression
- diabetes mellitus
- drug problems
- anaemia
- thyroid disorders, especially hypothyroidism
- urinary tract infection
- vertebral (spinal) dysfunction

Hypothyroidism has been estimated to exist in up to 15% of women aged 60 and above, and searching for clues may elicit subtle symptoms and signs previously attributed to ageing.

Relationships and psychosocial health. Consider the mental health of patients, particularly the elderly, by enquiring about how they are coping with life, how they are coping financially, about their peace of mind and how things are at home. Focus on the quality of their close relationships (e.g. husband - wife, father - son, mother - daughter, employer - employee). Enquire about losses in their life, especially family bereavements.

EXAMINATION SCREENING SPECIFIC TO CHILDREN[13]

Childhood health record books provide an excellent opportunity for communication between different health care givers; parents should be provided with the record books and encouraged to bring them to every visit. Various recommendations for screening are made under the following headings.

Height/weight/head circumference. Record height from age 3 and weight at regular intervals to age 5 years. Record head circumference at birth and then up to 6 months. The adequacy of a child's growth cannot be assessed on one measurement and serial recordings on growth charts are recommended. Head circumference recordings may provide further data about a child's growth.

Hips. Screen for congenital dislocation at birth, 6 - 8 weeks, 6 - 9 months and 12 - 24 months.

The flexed hips are abducted, checking for movement and a ‘clunk’ of the femoral head forwards (the test is most likely to be positive at 3 - 6 weeks and usually negative after 8 weeks). Shortening or limited abduction is also abnormal. Ultrasound examination is more sensitive than the clinical examination especially up to 3 to 4 months. Observe gait when starting to walk.

Strabismus. Strabismus should be sought in all infants and toddlers by occlusion testing (not very sensitive), examining light reflexes and questioning parents, which must be taken very seriously. Amblyopia can be prevented by early recognition and treatment of strabismus by occlusion and surgery. Early referral is essential.

Visual acuity. At birth and 2 months, eyes should be inspected and examined with an ophthalmoscope with a 3+ lens at a distance of 20 - 30 cm to detect cataracts and red reflexes. At 9 months gross vision should be determined by assessing ability to see common objects. Visual acuity should be formally assessed at school entry using Sheridan Gardiner charts.

Hearing. Hearing should be tested by distraction at 9 months or earlier; also by pure tone audiometry at 1000 and 4000 hertz when a child is 4 years (preschool entry) and 12 years.

Note. Formal audiological evaluation should be carried out at any time if there is clinical suspicion or parental concern. No simple screening test is very reliable for sensorineural or conductive deafness.

Testes. Screen at birth, and 6 - 8 weeks, 6 - 9 months and 3 years for absence or maldescent. Those who have been treated for maldescent have a higher risk of neoplastic development in adolescence.

Dental assessment/fluoride. Advise daily fluoride drops or tablets, if water supply is not fluoridated. Children's teeth should be checked regularly, particularly if a school dental service is not available. Advice should be given on sugar consumption, especially night-time bottles, and tooth cleaning with fluoride toothpaste to prevent plaque.

Scoliosis. Screening of females by the forward flexion test, which is carried out around 12 years of age, is of questionable value because of poor sensitivity and specificity.

Congenital heart disease. The heart should be auscultated at birth, in the first few days, at 6 - 8 weeks and on school entry.

Femoral pulses. Testing for absence of femoral pulses or delay between brachial/femoral pulses at birth and 8 weeks will exclude coarctation of the aorta. Refer the child immediately if concerned.

Speech and language. A child's speech should be intelligible to strangers by 3 years. It is related to hearing.

SCREENING FOR ADULTS[13]

The following recommendations apply for adults.

Weight. Weight should be recorded at least every few years. Obesity is a major reversible health risk for adults contributing to many diseases (e.g. heart disease, diabetes, arthritis). Body mass index (BMI) should ideally be between 20 and 25.

BMI ° (Weight (kg)) + (Height (m)^2)

Abdominal obesity is a major risk factor for adults. The waist:hip circumference ratio is regarded as a useful predictor of cardiac disease. Recommended waist:hip ratios are:

- males <0.9
- females <0.8

Blood pressure. Blood pressure should be recorded at least every 1 - 2 years on all people 16 years and over. There is no dispute that control of blood pressure results in reduced mortality from cerebrovascular accidents and, to a lesser extent, heart disease, kidney failure and retinopathy.

Cholesterol. All adults aged 18 and over should have a 5-yearly estimation of serum cholesterol. Total cholesterol is adequate for screening purposes. HDL levels give additional information. The National Heart Foundation recommends keeping cholesterol levels below 4.0 mmol/L. For most, dietary modification is sufficient to achieve these levels; some may require drug treatment.

Carcinoma of the cervix. Women aged 18 - 70 who have ever been sexually active should have a Pap smear every 2 years. Those over 70 who have never been screened should have two successive tests before screening is ceased. after consideration of the relative risks to the individual woman, she and her physician may choose to increase the interval between smears, but the interval should not be greater than 3 years.

Risk factors include:

- all women who are or ever have been sexually active
- early age at first sexual intercourse
- multiple sexual partners
- genital wart virus infection
- cigarette smoking

- those with LSIL and HSIL on Pap smears

Carcinoma of the breast. Mammography should be performed at least every 2 years on women aged 50 - 70 years. It is not useful for screening prior to age 40 years due to difficulty in discriminating malignant lesions from dense tissue. There is insufficient evidence of long-term benefit to recommend screening in the 40 - 50 age group. Mammography must not be used alone to exclude cancer if a lump is palpable. Such lesions require a complete appraisal since, even in the best hands, mammography still has a false negative rate of at least 10%. Genetic testing should be considered in those at risk.

Colorectal cancer (CRC). A history should be taken, with specific enquiry as to family history of adenomas or colorectal cancer, past history of inflammatory bowel disease and rectal bleeding. Rectal examination should be performed as part of an examination. Faecal occult blood testing (FOBT) every 2 years is now recommended for screening for people over 50 years without symptoms and with average or slightly above average risk.

Should a positive history be elicited, then the following are recommended:

- past history or large bowel cancer or colonic adenomas - colonoscopy
- past or present history of ulcerative colitis - colonoscopy with biopsies
- familial polyposis, Gardner's syndrome - sigmoidoscopy or colonoscopy

Prophylactic colectomy needs consideration in some individuals.

Apart from FOBT screening the NH&MRC currently recommends 2-yearly colonoscopy for people from 25 - 30 years of age if there is a family history on the same side of the family of:

- three or more first or second degree relatives with CRC at any age
- two or more first or second degree relatives diagnosed as CRC < 50 years of age
- a family member where genetic studies identify a high risk (Refer to RACGP *Guidelines for Preventive Activities in General Practice*13 for further information.)

Genetic testing should be considered in those at risk.

Prostate cancer. Screening is controversial. The RACGP guidelines do not recommend routine screening with DRE, PSA or transabdominal ultrasound. Patients should make their own decision after being fully informed of the potential benefits, risks and uncertainties of testing. Doctors should also use their clinical judgement for their individual male patient.

Skin cancer. All patients should be informed regularly about the need for protection of the skin from solar radiation, using clothing and sunscreens, and avoiding exposure during midday hours.

Skin cancer, which is increasing in incidence, is common in Australia, particularly in more northern areas. Squamous cell carcinoma, and melanoma in particular, may be lethal. Detection and treatment of early lesions prevents mortality and morbidity. Prevention of skin cancer by reduction of sun exposure should be taught to all patients.

Oral hygiene/cancer. Patients should be counselled about cessation of smoking and alcohol consumption, and dental hygiene should be taught. The oral cavity should be inspected annually in patients over the age of 40 years.

Although oral cancer has a relatively low incidence, premalignant lesions may be detected by inspection of the oral cavity. Its incidence is highest in elderly people with a history of heavy smoking or drinking. Poor dental hygiene may result in poor nutrition, particularly among the elderly.

Cancer screening in summary[13]

- Screen for breast, cervical and colorectal cancer.
- Routine population-based screening is at this stage not recommended for lung, melanoma, ovarian, prostate and testicular cancers.

IMMUNISATION

Basic diseases (diphtheria, tetanus, polio, whooping cough, measles, mumps, rubella) should be covered. Children and adolescents should be immunised according to the current NH&MRC recommended standard vaccination schedule.

Age	Immunisation
Birth	Hepatitis B
2 months	DTP, Hib, hepatitis B, polio, pneumococcus
4 months	DTP, Hib, hepatitis B, polio, pneumococcus
6 months	DTP, polio, Hib pneumococcus and hepatitis B (or at 12 months)
12 months	Measles/mumps/rubella, Hib, meningococcus C and hepatitis B (or at 6 months)
18 months	Varicella, pneumococcus (ATSIP)
4 years	DTP, measles/mumps/rubella, polio
15 - 17 years (prior to leaving school)	Adult diphtheria/tetanus, polio

Hib ° *Haemophilas influenzae* b.

DTP (triple antigen) ° diphtheria, tetanus, pertussis

ATSIP ° Aboriginal and Torres Strait Islander People

Special considerations. The NH&MRC endorses hepatitis B vaccine for all infants and preadolescents (3 doses). It also advises against injections into the buttock (challenged by many), preferring the anterolateral thigh for children under 12 months with the deltoid region being an alternative in older children and adults. The recommended needle is 23 gauge, 25 mm long. Do not postpone immunisation for minor illnesses such as mild URTI. Acellular pertussis vaccine, which reduces the risk of reactions has become a standard component of triple antigen.

Figure 9.2 Immunisation of an older child: important continuing preventive care

All adults should receive an adult diphtheria and tetanus (ADT) booster *each* 10 years.

All women of child-bearing years should have their rubella antibody status reviewed.

The general immunisation schedule is presented in Table 9.5.

Table 9.5 NH&MRC standard vaccination schedule (2006)

Initial courses			
Diphtheria, tetanus and pertussis	Intramuscular	DTP (triple antigen)	2, 4, 6 months
Poliomyelitis	Intramuscular	IPV (with DTP)	2, 4, 6 months
Measles, mumps and rubella	Subcutaneous	MMR	12 months
Haemophilus influenzae b (Hib)	Intramuscular	various	2, 4, 6, 12 months
Hepatitis B	Intramuscular	HBsAg subunit	0, 2, 4, then either at 6 or 12 months
Pneumococcus	Intramuscular	7 v PCV	2, 4, 6 months
	Intramuscular	23 v PPV	18 months (ATSIP)
Meningococcus C	Subcutaneous	various	12 months
Varicella	Subcutaneous	various	18 months
Booster doses			
Diphtheria, tetanus and pertussis	Intramuscular	DTP (Triple antigen)	4 years or prior to school entry
Poliomyelitis	Oral	Sabin	4 years and 15 - 19 years
Measles, mumps and rubella	Subcutaneous	MMR	4 years
Diphtheria and tetanus	Intramuscular	Adult diphtheria/tetanus (ADT)	15 - 17 years or prior to school leaving, then every 10 years

Fever and illness. Children with minor illness (providing the temperature is <38.5°C) may be vaccinated safely. Otherwise it should be delayed. A simple past febrile convulsion or pre-existing neurological disease is not a contraindication to pertussis vaccination. Absolute contraindications include encephalopathy within 7 days of a previous DTP or an immediate severe or anaphylactic reaction to DTP.

Paracetamol prophylaxis. Consider giving 15 mg/kg paracetamol liquid to children within a 30 minute period before the injection and then a further dose in 2 to 4 hours, to reduce the side effects of the vaccines.

influenza. influenza immunisation is recommended on an annual basis for persons of all ages with chronic debilitating diseases, especially chronic cardiac, pulmonary, kidney and metabolic diseases, persons over 65 years of age, Indigenous Australian adults over 50 years of age and persons receiving immunosuppressant therapy. Health care personnel may wish to consider it for their own use.

Pneumococcal disease. This should be considered for the same risk groups as influenza vaccine. Those at higher risk of fatal pneumococcal infection (e.g. post-splenectomy or Hodgkin's lymphoma), should receive a booster every 5 years. This is currently provided for all children.

Hepatitis A. Immunisation is recommended for:

- certain occupational groups at risk (e.g. health workers, child care workers, sewage workers)
- non-immune homosexual men
- those with chronic liver disease
- recipients of blood products
- travellers to hepatitis A endemic areas

Hepatitis B. Immunisation is recommended routinely for all children at birth, 2 months, 4 months and at either 6 or 12 months, and for individuals of all ages who, through work or lifestyle, may be exposed to hepatitis B and have been shown to be susceptible. Such groups would include health care personnel, personnel and residents of institutions, prisoners and prison staff, persons with frequent and/or close contact with high-risk groups, and persons at increased risk due to their sexual practices. Household contacts of any of the above groups should be considered for immunisation. Booster doses are not recommended for immunocompetent people but are recommended for immunosuppressed individuals. Universal vaccination represents a preventive step against hepatocellular cancer.

Haemophilus influenza type b. Hib immunisation is recommended for all children, especially those in child care. It is ideal to achieve immunity by the age of 18 months and preferably commencing at 2 months. Risk factors for Hib disease include day care attendance, presence of ill siblings under 6 years of age in the home and household crowding.

Q fever. People at reasonable risk from Q fever, particularly abattoir workers, should be given this vaccine, which is virtually 100% effective.

Measles-mumps-rubella. Both females and males should be immunised against measles, mumps and rubella at the age of 12 months and 4 - 5 years using the trivalent vaccine. All non-immune women who are postpartum or of child-bearing age should be immunised.

Varicella vaccine. This is available and one dose is given at 18 months. Those over 12 years have a course of two injections.

Meningococcal vaccine. Meningococcal disease is caused by *Neisseria meningitides,* which has thirteen serogroups of which A, B and C account for over 90% of isolated cases, with serogroup B responsible for most cases. A vaccine against serogroup B is not yet available. The main vaccine that is available is a quadrivalent polysaccharide vaccine against serogroups A, C, Y and W125 for use in individuals over 2 years as a single injection. Universal prevention by immunisation remains unsatisfactory. It is most useful when a community outbreak due to proven serogroup C occurs.

REFERENCES

1. Piterman L, Sommer SJ. *Preventive Care.* Melbourne: Monash University, Department of Community Medicine, Final Year Handbook, 1993: 75 - 85.
2. Silagy C. Prevention in general practice. In: McNeil J et al., eds, *A Textbook of Preventive Medicine,* Melbourne: Edward Arnold, 1990: 269 - 77.
3. National Health Strategy. *The Future of General Practice.* Issues paper No. 3. Canberra: AGPS, 1992: 54 - 169.
4. Stephens NG, Parsons A, Schofield P et al. Randomised controlled trial of vitamin E in patients with coronary disease: Cambridge Heart Antioxidant Study (CHAOS). Lancet, 1996; 347: 781 - 6.
5. Locke FB, King H. Cancer mortality risk among Japanese in the United States. National Cancer Institute, 1980; 65: 1149.
6. Potter JD, McMichael AJ. Diet and cancer of the colon and rectum: a case control study. National Cancer Institute, 1986; 76: 557 - 69.
7. Rakel RE. *Essentials of Family Practice.* Philadelphia: Saunders, 1993: 126 - 7.
8. Editorial. Position of the American Dietetic Association: photochemicals and functional foods. J Am Diet Assoc, 1993; 93: 493 - 6.
9. Doll R, Peto R. *The Causes of Cancer.* New York: Oxford University Press, 1981: 1197 - 219.
10. Sali A. Strategies for cancer prevention. Aust Fam Physician, 1987; 16: 1603 - 13.
11. Antic R. *Report on the Cost of Asthma in Australia*. Melbourne: National Asthma Campaign, 1992: 14 - 33.
12. Sridhar MK. Editorial. Nutrition and lung health. Br Med J, 1995; 310: 75 - 6.
13. Royal Australian College of General Practitioners. *Guidelines for Preventive Activities in General Practice* (6th edn). Melbourne: RACGP, 2005.

CHAPTER 10. NUTRITION IN HEALTH AND ILLNESS

On the 20th of May 1747 on board the *Salisbury* at sea I took 12 patients in the scurvy with putrid gums, the spots and lassitude and weakness of their knees. They had a common diet - two were ordered a quart of cyder a day - two, elixir vitriol three times a day - two, 2 spoonfuls of vinegar three times a day - two were put on a course of sea water - two had 2 oranges and one lemon each day - two, nutmeg, garlic and other additives to barley water. The consequence was, that the most sudden and visible good effects were perceived from the use of the oranges and lemons with one being at the end of 6 days fit for duty.

JAMES LIND, A (SUMMARISED) ACCOUNT OF SCURVY, 1747

NUTRITION IN HEALTH AND ILLNESS

Good nutrition is fundamental to good health. It influences management in all branches of medicine. Modern people's health varies from the excesses of inappropriate nutrition, resulting in obesity and various degenerative disorders, to malnutrition and other deficiency states seen in those unfortunates deprived of nutrients.

Nutritional factors may play a vital role in the causation of several of the major diseases, such as coronary artery disease, hypertension, diabetes and cancer.

Special diets are important in the management of many hereditary metabolic disorders, such as phenylketonuria and galactosaemia, and several other disorders such as coeliac disease.

NUTRITIONAL ASSESSMENT

The first step in nutritional assessment is to identify the high-risk patient.[1] Those at high risk of nutritional insufficiency include those with a history of obesity, eating disorders, chronic illness, psychological disorders, the elderly, the institutionalised, trauma victims and those with long periods of hospitalisation, including major surgery. Of particular interest is the rate of growth and development in infants and children and the body composition in children and adults.

When taking the history it is appropriate to include a 24-hour recall of foods eaten and ideally get the patient to complete a symptom questionnaire that can then be linked to a computerised nutritional evaluation program, such as Nutricheck.[1] Evaluate sunlight exposure.

A nutritionally focused physical examination should be performed on each patient at risk, with the emphasis on body weight, waist size, muscle wasting, fat stores and signs of micronutrient deficiencies.

Examples of the latter include zinc deficiency, which affects taste and smell and the skin. Deficiencies of vitamins B_6 and B_{12} cause neurological disorders, such as peripheral neuropathy.[1] Alcoholism and malnutrition affect many systems, including the gastrointestinal system. The oral cavity, especially the gums, teeth and buccal mucosa, are affected by vitamin B complex and vitamin C deficiencies. Bones and joints are affected in scurvy, rickets, osteomalacia and osteoporosis. The important anthropometric measurements include height and weight, skin-fold thickness and waist - hip circumference ratio (refer p. 850). Laboratory investigations depend on the clinical examination and should be selective.

THE GENERAL PRINCIPLES OF OPTIMAL NUTRITION

In order to help people make healthy choices, the Health Foundations of several countries have developed recommendations for eating a healthy diet in the form of a pyramid (Figure 10.1) although its effectiveness is being reviewed.[1, 3]


Fat, oil, and sweets group
Use sparingly
OIL
Milk, yoghurt, and cheese group
2–3 servings
MILK
Yogu
Meat, poultry, fish, dry beans, eggs and nuts group
2–3 servings
Vegetable group
3–5 servings
Fruit group
2–4 servings
Bread, cereal, rice, and pasta group
6–11 servings

Figure 10.1 Daily food guide pyramid

The CSIRO of Australia has formulated a 12345+ food and nutrition plan.

Daily recommendations include:

- breads and cereals 5+ servings
- vegetables 4 servings
- fruit 3 servings
- milk and milk products 2 servings
- meats and alternatives 1 serving
- indulgences or extras no more than 2 servings

The corresponding recommendations from the US Department of Agriculture were breads and cereals (6 - 11 servings), vegetables (3 - 5), fruit (2 - 4), milk and dairy products (2 - 3), meat, poultry and fish (2 - 3), and a group of fats, oils and sweets to be used sparingly.[1] The number of recommended servings varies depending on an individual's energy needs, which can vary from 6700 kJ to more than 10 050 kJ (1600 - 2400 calories) a day.

The heart-healthy eating pyramid of the Australian Nutrition Foundation (1997)[4] has a simplified system, namely:

- Eat most - vegetables, dried peas, beans and lentils, cereals, bread, fruit and nuts
- Eat moderately - lean meat, eggs, fish, chicken (without skin), milk, yoghurt, cheese
- Eat in small amounts - oil, margarine, reducedfat spreads, butter, sugar

What is a serving?[4]

The following guides represent one serve:

Milk and milk products group:	1 cup (300 mL) of milk 200 g yoghurt 40 g cheese
Meat group:	60 - 100 g cooked lean meat, poultry or fish 2 eggs 2/3 cup of cooked legumes
Vegetable group:	1 cup raw leafy vegetables 1/2 cup other vegetables (cooked or raw) 3/4 cup vegetable juice
Fruit group:	1 medium apple, banana, orange 1/2 cup of chopped, cooked or canned fruit 3/4 cup fruit juice 1/4 melon

Tips for the overweight patient

With patients who are overweight, it is helpful to encourage them to reduce meal sizes by one-third and to avoid second helpings (see p. 853). Thus, people are able to eat what they enjoy, they just eat a little less. It is also important to eat slowly. Other suggestions include:

- Select fish, poultry and lean meats.
- Trim excess fat from meat and skin from poultry.
- Limit the amount of butter or margarine on vegetables and bread.
- Use a minimum of cooking fats.
- Limit the intake of full-cream products, fried foods and fatty takeaway foods.
- Eat less sugar - avoid lollies, soft drinks, biscuits, syrups and cakes.
- Increase intake of complex carbohydrates that contain starch and fibre.
- Eat more breads, cereals, fruit and vegetables.
- Drink more water.
- Plant food is good for you, have it as part of breakfast.
- What you usually eat matters most, not what you occasionally eat.

PROTEIN[4]

Proteins are composed of carbon, hydrogen, oxygen, nitrogen, phosphorus, sulphur and iron. They make up the greater part of plant and animal tissue and provide the amino acids essential for the growth and repair of tissue. Protein in the body in muscle, connective tissue and enzymes is constantly being broken down, while dietary protein is hydrolysed to amino acids that are both essential and non-essential. A complete protein is one that contains all the nine indispensable amino acids, namely, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan and valine.

Protein in animal products (fish, meat and milk) is of high quality and that in vegetable products is lower because of a limited supply of lysine (in cereals) and methionine and cysteine (in legumes).4 Vegetarian diets are usually adequate in protein, especially if the combining vegetable groups complement each other in basic amino acid groups. Diets that exclude all animal products may be inadequate, especially in children. Infants and children require 2 - 2.2 g protein/kg/day.

- High protein content foods - lean beef and lamb, chicken, fish, eggs, milk, cheese, soy beans
- Medium protein content foods - bread, spaghetti, corn, potatoes (cooked), rice (cooked), cabbage, cauliflower

Protein - energy malnutrition

This is a deficiency syndrome with a reduction in all macronutrients, energy (kilojoules) and many micronutrients due to an inadequate intake of protein and energy food stuffs.

It is commonly found in infants and children in developing countries but can occur in persons of any age in any country.

Clinically, protein-energy malnutrition has three forms:

1. dry (thin, desiccated) - *marasmus*
2. wet (oedematous, swollen) - *kwashiorkor*
3. combined - *marasmic kwashiorkor*

Marasmus

Features:

- grossly underweight
- gross muscle wasting
- no fat
- hungry
- 'old man's' face
- no oedema
- normal hair

Caused by diet low in protein and calories.

Kwashiorkor

Features:

- oedema
- 'moon' face
- anorexic

- hair pale and thinned
- apathetic
- skin changes

Caused by a diet low in protein with some carbohydrate, leading to hypoalbuminaemia.

CARBOHYDRATES

Dietary carbohydrates include simple sugars, complex carbohydrates (starches) and indigestible carbohydrate (dietary fibre). Carbohydrates are the main source of dietary energy. The two most important crops feeding the world are rice and wheat, which are rich in starch. Starch and sucrose account for the majority of carbohydrates consumed in all diets. Carbohydrates that are available in food are:

- sugars - sucrose, lactose, maltose, glucose, fructose
- polyols - sorbitol, xylitol, maltilol, lactilol
- starch - amylose, amylopectin
- dextrose

As long as adequate energy and protein are provided in the diet, there is no specific requirement for dietary carbohydrate. A small amount - 100 g/day - is necessary to prevent ketosis.[4]

The glycaemic index (GI)

The GI, which applies to carbohydrate foods, is a numerical index based on a reference point of 100. It is a measure of the capacity to increase post- prandial glucose levels compared to a glucose load. The standard food is glucose, which is given an arbitrary level of 100.

The higher the GI, then the higher the rise in blood glucose level and thus the greater the insulin response.

A list of GI values of various foods is shown in Table 10.1.

Table 10.1 Glycaemic index values[3]

Low GI (<60)	Moderate GI (60 - 85)	High GI (>85)
Apple/peach/pear/orange/banana (medium)	Noodles	Raisins
	Wholemeal/white bread (1 slice)	Dried dates
Pasta	Sweet corn	Nutri-Grain/Weetbix/Rice Bubbles (50 g)
Baked beans	Chocolate (other)	Cornflakes
All Bran cereal	Orange juice	Jasmine white rice
Porridge/Special K	White rice	Most Australian rices
Yoghurt Ice-cream low-fat	Potato chips	Corn chips
	Can of soft drink	Sweet biscuits
Milk	Rye breads	Sugar
Brown rice	Rockmelon/pineapple/apricot	Cordial/Gatorade/other sports drinks
Peanuts	Most biscuits and cakes	Glucose
Muesli and oats		Potato (1 medium - 120 g)
Dark chocolate		French fries
Mixed grain bread		White bread
Sweet potato/peas		Baked or boiled potatoes/pumpkin
		Watermelon

FAT

Dietary fat, which is composed mainly of fatty acids and dietary cholesterol, is the most concentrated source of food energy.[4]

Fatty acids are classified according to the number of unsaturated double bonds:

- nil - saturated (e.g. butyric and stearic acids)
- one - monounsaturated (e.g. oleic acid)
- more than one - polyunsaturated (e.g. linoleic acid, eicosapentanoic acid [EPA], docosahexanoic acid [DHA])

Polyunsaturated fatty acids (two or more unsaturated bonds) can be subdivided into:

- n-6 (e.g. linoleic acid, 2 unsaturated bonds; arachidonic acid, 4 unsaturated bonds)
- n-3 - omega-3 fatty acids (e.g. ?-linolenic acid, 3 unsaturated bonds; EPA, 5 unsaturated bonds; DHA, 6 unsaturated bonds)

The n-3 and n-6 polyunsaturated fatty acids with chain lengths of 18 or more are called essential fatty acids because they are required for vital body functions and animals, including humans, are unable to synthesise them.[4]

The proportions of saturated, monounsaturated and polyunsaturated fatty acids in the diet are important determinants of health and disease.[4] The current strategy is to reduce total fat intake and reduce saturated fats and increase unsaturated fats, especially n-3 polyunsaturated fats.

Fish oil contains omega-3 fatty acids, which are considered more potent than the omega-3 fatty acids found in plants. The value of omega-3 fatty acids in preventing cardiovascular mortality has been well proven. They have no effect on cholesterol levels but have a well-documented potent hypotriglyceridaemic effect.[5]

Cholesterol, which is a major constituent of cell membranes, is synthesised by the body and is not an essential nutrient. The plasma cholesterol level, and hence the amount of cholesterol in the diet, has been related to the development of atherosclerosis. However, other factors are related to the development and prevention of atherosclerosis. In the past decade, specific nutrients, especially plant sterols (phyto sterols), soy protein and soluble fibre, have been shown to improve the plasma cholesterol profile, which has led to their appropriate inclusion in 'designer' foods targeting the health-conscious consumer market.[5]

The Lyon Heart Study [6]

This very impressive prospective, randomised, singleblinded, secondary prevention trial investigated the effects of a 'Mediterranean'-type diet containing a plant oil rich in omega-3 fatty acids on 605 subjects who had survived their first heart attack. The control group were given a low-cholesterol diet with a mix of unsaturated fats. The intervention group had a diet higher in oleic acid, omega-3 fatty acid, fibre and vitamin C (olive and canola oils were used in food preparation). In this group, the cardiovascular mortality was reduced by 73%. This was far greater than the control group and also for groups taking lipid-lowering agents, and the benefit was independent of any reduction in cholesterol or triglycerides. One explanation is that the antioxidants and phytochemicals in the plant-based diet stabilise the endothelium of arteries.

Some consequent lessons for people with a highlipid profile are:[3]

- Eat more fish - at least twice a week.
- Switch to skim-based dairy products - low-fat milk and yoghurts.
- Eat plenty of fruit, vegetables and grain food.
- Use monounsaturated (olive) oils and perhaps margarine instead of butter. However, use these cooking oils lightly.
- Use alternative toast spreads, e.g. hommus, baked beans, lentils, salmon.
- Limit intake of dietary cholesterol (less than 300 mg a day). Eat less animal-based food, dairy products, eggs (limit two a week), saturated fats, bakery food, fast food.

Ornish nutritional program

Dean Ornish, a US cardiologist, was one of the first doctors to bring out a nutritional program to treat heart disease. The program involves drinking a lot of water, cutting down consumption of dairy produce and eating plenty of complex carbohydrates. Patients also exercise, undergo stress reduction and achieve ideal weight. The results showed that intensive changes in diet lead to regression of coronary heart disease.[7]

Antioxidants

> They may be much more important than doctors thought in warding off cancer, heart disease and the ravages of ageing. And no, you may not be getting enough of these crucial nutrients in your diet.

T ime magazine [8]

The antioxidant issue is still controversial and unclear. Empirical observation of healthy communities over the years indicates good health outcomes, especially with cardiovascular status, in people having an optimal diet containing high levels of vitamins and minerals (especially from fruit and vegetables).

Food antioxidants (Table 10.2) appear to protect against free radicals, which can suppress immunity. Free radicals, which are usually a toxic form of oxygen containing an odd number of electrons, are produced by a variety of toxins.[9] Apart from the possible adverse effect to immunity from free radicals, they may also damage body tissues, such as the liver in alcoholics, as well as increasing susceptibility to degenerative diseases.[10]

Table 10.2 Food antioxidants

Vitamin A, especially beta-carotene
Vitamin C
Vitamin E
Ubidecarenone (coenzyme Q10)
Selenium
Zinc
Manganese nutrient cofactors
Copper

Source: After Sali[9]

However, Bury, in a review of the literature of antioxidant nutrients, concludes that 'high intake of antioxidant nutrients from food sources appears to offer some health advantage but claims for any therapeutic benefit of antioxidant supplements are premature and scientifically unjustified at present'.[11] On the other hand, studies have proved that antioxidants have a preventive role in macular degeneration (see p. 844). The jury appears to be still out on the true value of antioxidant supplements.

Prime sources of antioxidants in food[12]

- Vitamin C - citrus fruits, berries, papaya, green leafy vegetables
- Vitamin E - seed-like cereal grains, nuts and oils (plants), eggs
- Beta-carotene - orange-coloured and darkgreen leafy vegetables
- Selenium - grains, meats, Brazil nuts, fish
- Copper - cocoa, wheat bran, yeast
- Ubiquinone - meats, fish, peanuts
- Phytochemicals - soy, tea, green tea, herbs, apples, onions, cocoa

Folate-containing foods[8]

- Green leafy vegetables - broccoli, spinach
- Wheat grain
- Wholegrain cereals
- Starchy beans - kidney and butter
- Peas, corn, cauliflower
- Nuts
- Avocado
- Liver
- Folic acid fortified foods (e.g. breakfast cereals)

NUTRITION AND GOUT

Certain foods can aggravate gout. These include:

- tinned fish (e.g. sardines, anchovies etc.)
- organ meats (e.g. liver, pancreas, brain, kidney etc.)
- alcohol (the major one)

Gout often occurs in males after they binge drink in their late 20s or 30s. Beer is particularly problematic. It is useful to recommend an increase in water consumption. Since gout and hyperuricaemia is a proven association with coronary artery disease a healthy heart prevention diet is advisable.

THE MAIN PRINCIPLES OF THE DIETARY CONTROL OF DIABETES

Diabetes is an endemic problem in many communities, particularly in Indigenous communities. It is thought that for every two people diagnosed with diabetes, there may be one who is not diagnosed. Prevention of diabetes is very important and diet can play a significant role in this.

As many as 30% of people who suffer from diabetes are now taking insulin and oral hypoglycaemic medications. However, if sufferers can maintain an ideal weight and control symptoms through diet, the need for medications is reduced. Exercise is also very important.

Dietary principles for preventing and control of diabetes include:

- Follow the guidelines of the healthy eating pyramid.
- Keep to an ideal weight.
- Eat low GI foods.
- Limit the consumption of fat and fatty products.
- Eat complex carbohydrates, such as starchy fibre foods, cereals, wholemeal bread.
- Avoid simple carbohydrates (e.g. sugar).
- Spread the consumption of carbohydrates throughout the day.

In summary, the diet for both type 1 and type 2 diabetes is based on achieving ideal weight and following a diet of high-fibre carbohydrates with a low GI and low fat.

IRON THERAPY

Iron-deficiency anaemia is a common problem in our society, particularly in children from 6 months to 2 years old who have been given a lot of cow's milk. In such cases it is important to educate people about iron-rich foods and the quantities they need.

GUIDELINES FOR SAFE CONSUMPTION OF ALCOHOL

Males

- Problematic drinking occurs with more than four standard drinks per day.
- Dangerous drinking is six or more standard drinks per day.

Females

- It is preferable to limit consumption to two standard drinks per day
- Dangerous drinking is four or more standard drinks per day.

Refer to p. 1251 for alcohol guidelines.

COELIAC DISEASE

Coeliac disease occurs as a result of sensitivity to gluten. It is quite common and often undiagnosed. In some cases, it can occur after a bout of gastroenteritis. It can be treated with a gluten-free diet.

- For breakfast, exclude wheat, barley, oats and rye.
- Any food containing gluten (e.g. flour and bread) should be avoided.
- Any food containing hidden ingredients (e.g. stock cubes) should be avoided.

FOODS ASSOCIATED WITH MIGRAINE

Common ones include:

- wine, particularly red wine
- cheese
- oranges
- tomatoes
- caffeine in some people

Other culprit foods might include amine-rich foods, such as bananas and avocados. It is helpful to ask patients to keep a diary and see if they can find links between what they eat and their migraine attacks. Some migraines are not linked to diet.

NUTRITION AND CHRONIC SIMPLE CONSTIPATION

Important advice for treatment of constipation includes:

- Drink a lot of fluids, especially water and fruit juices.
- Eat foods with bulk and roughage (e.g. bran, fresh and dried fruits, wholemeal bread).

Foods that are high in roughage include (in order):

- bran
- carrots
- apples
- lettuce
- cabbage
- peas
- cauliflower
- bananas
- potatoes

Fruits are high in fibre and some are natural laxatives, such as:

- prunes
- figs
- rhubarb
- apricots
- pears

Studies show that most people do not eat enough fruit. According to a recent survey of Victorian schoolchildren, one-third did not have fruit during the day, one-third did have fruit and one-third only had it spasmodically.

RECURRENT URINARY CALCULI

The dietary advice for recurrent urinary calculi includes:

1. Drink at least 2 L of water every day, or more if there is increased fluid loss: this is the most important step.
2. Minimise consumption of foods that contain oxalate or uric acid. Foods that contain oxalate include:
    - chocolate
    - coffee
    - cola drinks
    - rhubarb
    - tea
3. Avoid milk in tea - calcium precipitates oxolate. Foods that contain uric acid include:
    - beer
    - red wine
    - red meat
    - organ meats
4. Avoid processed meats, organ meats (e.g. brain, kidney, liver, sweetbread), yeast spreads and other high salt foods. Restrict salt intake.
5. Reduce animal protein consumption: restriction to one major meat meal a day (includes chicken and fish).
6. Add citrate-containing fruit juices to the diet, including grapefruit, apple and orange juice.
7. Eat a healthy diet of vegetables and fruit with a high fibre content.

DIETARY GUIDELINES FOR CHILDREN AND ADOLESCENTS (NH&MRC)

1. Encourage and support breastfeeding.
2. Children need appropriate food and physical activity to grow and develop normally. Growth should be checked regularly.
3. Enjoy a wide variety of nutritious foods.
4. Eat plenty of breads (least processed), cereals, vegetables (including legumes) and fruits.
5. Low-fat diets are not suitable for young children. For older children, a diet low in fat and, in particular, low in saturated fat, is appropriate.
6. Encourage water as a drink. Alcohol is not recommended for children.
7. Eat only a moderate amount of sugars and foods containing added sugars.
8. Choose low salt foods.
9. Eat foods containing calcium.

10. Eat foods containing iron.

DIETARY GUIDELINES FOR OLDER PEOPLE (NH&MRC)

1. Enjoy a wide variety of nutritious foods.
2. Keep active to maintain muscle strength and a healthy body weight.
3. Eat at least three meals every day.
4. Care for your food - prepare and store it correctly.
5. Eat plenty of vegetables (including legumes) and fruit.
6. Eat plenty of cereals, breads and pastas.
7. Eat a diet low in saturated fat.
8. Drink adequate amounts of water and/or other fluids.
9. If you drink alcohol, limit your intake.
10. Choose foods low in salt and use salt sparingly.
11. Include foods high in calcium.
12. Use added sugars in moderation.

GENERAL DIETARY GUIDELINES FOR GOOD HEALTH - A SUMMARY

- Keep to an ideal weight.
- Eat a high fibre diet.
- Eat more fruits and vegetables, least processed breads and cereals, preferably wholegrain.
- Eat fish at least twice a week (daily if possible).
- Choose a nutritious diet.
- Eat less saturated fat, refined sugar and salt.
- Use low-fat dairy products - milk and yoghurt.
- Avoid fast foods and deep-fried foods.
- Do not eat animal meat every day, and then only in small portions. Note that processed meats, such as sausages, have a very high fat content.
- Use monounsaturated (olive) oils and perhaps margarine instead of butter.
- Use olive oil for cooking rather than polyunsaturated oils.
- Always trim fat off meat.
- Drink more water.
- Limit salt intake; pepper is okay.
- Limit caffeine intake (0 - 3 drinks a day maximum).
- Check plasma cholesterol level and, if it is elevated, aim to reduce it with diet.

A very appropriate food pyramid is the Kiat model (Figure 10.2).[13]


Fats
olive oil, nuts
Proteins
egg whites, fish, nuts, legumes, white meats, tofu, soy drinks, non-fat milk
Carbohydrates
fruit, cereals, rice, pasta, noodles, pumpkin, potatoes
Greens + water
fruit and vegetables juices, soya, non-fat milk

Figure 10.2 The Kiat simplified good health pyramid

These nutritional recommendations need to be supplemented with other lifestyle and mental health ones, such as:

- Do not smoke.
- Limit alcohol intake to two standard drinks per day.
- Reserve alcohol for special occasions and to only one occasion in the day.
- Get sufficient sunlight to gently ‘pink’ your arms and face.
- Take regular exercise (e.g. 30 minutes per day, 3 - 4 days per week).
- Practise relaxation.
- Partake in ample recreational activities.
- Encourage a circle of friends who can offer emotional support.
- Give expression to feelings rather than suppressing them.
- Discuss problems regularly with someone with a listening ear. Everyone needs one close friend to talk to.

REFERENCES

1. Sydney-Smith M. Nutritional assessment. Curr Ther, 2000; September: 13 - 22.
2. Beers MH, Berkow R. *The Merck Manual of Diagnoses and Therapy* (17th edn). New Jersey: Merck Research Laboratories, 1999: 12 - 16.
3. Crimmins B. Nutrition. CHECK program 391: Melbourne: RACGP; 2004; 391: 1 - 30.
4. Wahlqvist ML. *Food and Nutrition*. Sydney: Allen & Unwin, 1997.
5. Howe P. Nutrition and cardiovascular risk. Med Observer, 2001; 16 November: 36 - 7.
6. De Lorgeril M, Renaud S, Mamelle S et al. Mediterranean alpha-linoleic acid-rich diet in secondary prevention of coronary heart disease. Lancet, 1994; 343: 1001 - 29.
7. Ornish D, Scherwitz LW, Billings JH et al. Intensive lifestyle changes for reversal of coronary heart disease. JAMA, 1998; 280: 2001 - 7.
8. Toufexis A. The new scope on vitamins. Time, 1992; April 6: 50 - 5.

9. Sali A. Strategies for cancer prevention. Aust Fam Physician, 1987; 16: 1603 - 13.
10. Dormandy TL. An approach to free radicals. Lancet, 1983; 2: 1010 - 14.
11. Bury R. *Clinical Applications of Antioxidant Nutrients* . Melbourne: Department of Human Services, 1996: 26.
12. Wahlqvist ML, Wattanapenpaiboon N. Antioxidant nutrients. Aust Prescriber, 1999; 22(6): 142 - 4.
13. Kiat H. *East-West Medical Makeover*. Sydney: Image of Distinction, 2002.

CHAPTER 11. HEALTH PROMOTION AND PATIENT EDUCATION

Never believe what a patient tells you his doctor said.

SIR WILLIAM JENNER (1815 - 98)

HEALTH PROMOTION[1]

Health promotion is the motivation and encouragement of individuals and the community to see good health as a desirable state that should be maintained by the adoption of healthy practices. It is also the process of helping people obtain their optimal health.

For those who feel healthy, the message may have little meaning, but it is reinforced by contact with others who become ill, particularly within the family.

HEALTH EDUCATION

Health education is the provision of information about how to maintain or attain good health.

There are many methods, including the advertising of health practices, the provision of written information (e.g. about diet and exercise, immunisation, accident prevention and the symptoms of disease); and methods to avoid disease (e.g. sexually transmitted infection).

ILLNESS EDUCATION

A lot of so-called 'health' education is, in reality, information about the cause of particular illnesses. Clearly, the medical practitioner is in a pre-eminent position to provide his or her patients with specific information about the cause of an illness at the time, either individually or to the family. This educative strategy has a preventive objective that is often the modification of help-seeking behaviour.

Every consultation is an opportunity to provide information about the condition under care and this can be reinforced in written, diagrammatic or printed form. Patients' own X-rays can be similarly used to illustrate the nature of the problem.

HEALTH PROMOTION IN GENERAL PRACTICE

GPs are ideally placed to undertake health promotion and prevention, mainly due to opportunity.

There are several reasons for this health promotion role:

- Population access: over 80% of the population visit a GP at least once a year.[2]
- On average, people visit a GP about five times each year.
- GPs have a knowledge of the patient's personal and family health history.
- The GP can act as leader or co-ordinator of preventive health services in his or her local area.
- The GP can participate in community education programs.
- GPs should undertake opportunistic health promotion - the ordinary consultation can be used not just to treat the presenting problem, but also to manage ongoing problems, co-ordinate care with other health professionals, check whether health services are being used appropriately and undertake preventive health activities.[2]

OPPORTUNISTIC HEALTH PROMOTION

The classic model by Stott and Davis (Table 3.1 on p. 16) highlights the opportunities for health promotion in each consultation.[3] Since the consultation is patient-initiated, it is the doctor who needs to be the initiator of preventive health care. The potential in the consultation involves reactive and proactive behaviour by the doctor (Figure 11.1).[4]


Reactive
Presenting complaint overt/covert
Continuing care
Prevention, early detection, health promotion
Co-ordination of care
Modification of abnormal help-seeking behaviour
Proactive
Professional behaviours

Figure 11.1 The potential in every general practice consultation. Reproduced with permission from M. Sales [4]

Reactive professional behaviour deals only with the presenting complaint. It may be performed with skill but if the practitioner is only trained to perform reactively then the opportunity for preventive and promotive health care will be lost.

Proactive behaviour is defined as professional behaviour that is necessary for the patient's well-being, but it is performed not merely as a response to the presenting problem and it is initiated by the doctor.[4] It includes health promotion, preventive care and screening and the early detection of disease before it becomes symptomatic. Other aspects of proactive care are seen in Figure 11.1.

Proactive behaviour also includes:[4]

- continuing care of a previously treated problem (e.g. rechecking blood pressure, checking diabetic control, follow-up bereavement counselling)
- co-ordination of care by organising referral to appropriate agencies or specialists and maintaining adequate medical records
- the modification of abnormal or inappropriate help-seeking behaviour (e.g. the person who never attends is at risk from ‘silent disease’; the too frequent attender wastes resources and serious illness may be overlooked)

This mix of reactive and proactive behaviour is not appropriate in every consultation. It requires counselling skills and training in the delivery of quality general practice.

Methods

- Being informed and updated by maintaining continuing medical education, especially in preventive roles.
- Using health promotional material for patient education:
    - handouts
    - waiting room posters
    - waiting room video systems.
- Having an efficient medical record system.
- Operating a patient register and recall system.

- Encouraging regular health checks for at-risk groups.
- Providing regular advice on:
  - nutrition
  - exercise
  - stress management
  - weight control.
- Providing personal health records to the parents of newborn babies.

HEALTH GOALS AND TARGETS

Health goals and targets as determined by the Health Targets and Implementation Committee[5] were set in three areas - population groups, major causes of illness and death, and risk factors (Table 11.1).

Table 11.1 Health promotion areas in which goals and targets have been set

Population groups
The socioeconomically disadvantaged, Indigenous Australians, migrants, women, men, older people, children and adolescents.
Major causes of illness and death Heart disease and stroke, cancers (including lung, breast, cervical and skin cancer), injury, communicable diseases, musculoskeletal disease, diabetes, disability, dental disease, mental illness, asthma.
Risk factors Drugs (including tobacco smoking, alcohol misuse, pharmaceutical misuse or abuse, illicit drugs and substance abuse), nutrition, physical inactivity, high blood pressure, high blood cholesterol, occupational health hazards, unprotected sexual activity, environmental health hazards.

Source: Health Targets and Implementation Committee[5]

The six priority health areas for Australians are:

- asthma
- cancer control
- cardiovascular health
- diabetes mellitus
- injury, prevention and control
- mental health

PROMOTING HEALTHY LIFE STYLE IN GENERAL PRACTICE

General practitioners can provide a simple framework to encourage patients to adopt a healthy lifestyle whether they have a particular disorder or not. The acronyms act as a good aide-mémoire for practitioners for opportunistic health promotion.

The 'SNAP' guide[6]

The 'SNAP' guide has been developed by the RACGP to address important risk factors with patients with a view to encouraging change if appropriate. The guide comes as a comprehensive booklet and includes the 'Estimation of absolute 5-year risk of cardiovascular events' tables. The SNAP guide can be summarised by the following risk factors.

S ° smoking

N ° nutrition

A ° alcohol

P ° physical activity

The guide emphasises that there are health inequalities in the community because the risk factors are far more prevalent in people from low socioeconomicstatus backgrounds and Indigenous Australians.

The guide focuses on the 5 As as stages of change theory to promote change on lifestyle where appropriate.

The 5 As

Ask (1)	• identify patients with risk factors
Assess (2)	• level of risk factor and its relevance to the individual in terms of health • readiness to change/motivation
Advise (3)	• provide written information • provide a lifestyle prescription • brief advice and motivational interviewing
Assist (4)	• pharmacotherapies • support for self-monitoring
Arrange (5)	• referral to special services • social support groups • phone information/counselling services • follow up with the GP

The information can be accessed at < http://www.racgp.org.au/guidelines/snap >

The NEAT guide

The NEAT guide is basically similar to the SNAP guide, but with a greater emphasis on counselling the patient about lifestyle and the importance of stress management.

N ° Nutrition: optimal diet

E ° Exercise/physical activity

A ° Avoidance or moderation of potential

harmful substances (CATS):

- Caffeine
- Alcohol
- Tobacco
- Sugar, salt and social drugs

T ° Tranquility and promotion of recreation, relaxation techniques, meditation

PSYCHOSOCIAL HEALTH PROMOTION

There is a tendency for health goals and targets to focus mainly on physical illness and not emphasise mental health. However, this area represents an enormous opportunity for anticipatory guidance. It includes the important problems of stress and anxiety, chronic pain, depression, crisis and bereavement, sexual problems, adolescent problems, child behavioural problems, psychotic disorders and several other psychosocial problems.

Time spent in counselling, giving advice and stressing ways of coping with potential problems such as suicide and deterioration in relationships is rewarding. GPs need to pay more attention to promoting health in this area, which at times can be quite complex.

PATIENT EDUCATION

Evidence has shown that intervention by GPs can have a significant effect on patients' attitudes to a change to a healthier lifestyle. If we are to have an impact on improving the health of the community, we must encourage our patients to take responsibility for their own health and thus change to a healthier lifestyle. They must be supported, however, by a caring doctor who follows the same guidelines and maintains a continuing interest. Examples include modifying diet, cessation of smoking, reduction of alcohol intake and undertaking exercise.

In an American survey of 360 patients, 90% reported wanting a pamphlet at some or all of their office visits. Overall, 67% reported reading or looking through and saving pamphlets received, 30% read or looked through them and then threw them away, and only 2% threw them away without review. Only 11% of males and 26% of females reported ever asking a doctor for pamphlets. More patients desire pamphlets than are receiving them.[7]

Patient educational materials have been shown to have a beneficial effect. Giving patients a handout about tetanus increased the rate of immunisation against tetanus among adults threefold.[8] An ed ucational booklet on back pain for patients reduced the number of consultations made by patients over the following year and 84% said that they found it useful.[9] The provision of systematic patient education on cough significantly changed the behaviour of patients to follow practice guidelines and did not result in patients delaying consultation when they had a cough lasting longer than 3 weeks or one with ‘serious’ symptoms.[10]

There is no evidence that patient education has a harmful effect. Patient education about drug sideeffects has been shown not to have any detectable adverse effects.[11]

One form of patient education is giving handouts (either prepared or printed from a computer at the time of the consultation) to the patient as an adjunct to the verbal explanation which, it must be emphasised, is more important than the printed handout.

The patient education leaflets should be in nontechnical language and focus on the key points of the illness or problem. The objectives are to improve the quality of care, reduce costs and encourage a greater input by patients in the management of their own illness. In modern society where informed consent and better education about health and disease is expected, this information is very helpful from a medicolegal viewpoint.

The author has produced a book called Patient Education, which has a one-page summary of each of 252 common medical conditions.[12] The concept is to photocopy the relevant problem or preventive advice and hand it to the patient or relative. Over the years the greatest demand (following a survey of requests for prints of the sheets) has been for the following (in order):

- exercises for your back (Figure 11.2)



Figure 11.2 Patient education leaflet (diagrammatic part only): exercises for your lower back

- backache (Figure 11.3)



Figure 11.3 Patient education leaflet on backache (diagrammatic part only): rules of care for sitting, lying and bending

- exercises for your neck
- your painful neck
- exercises for your knee
- breastfeeding and milk supply
- how to lower cholesterol
- breast self-examination
- testicular self-examination
- vaginal thrush
- menopause
- anxiety
- coping with stress
- depression
- bereavement

SUMMARY

Recommended target areas for health promotion in general practice include:

- nutrition
- weight control
- substance abuse and control
    - smoking
    - alcohol
    - other drugs
- exercise practices
- appropriate sleep, rest and recreation
- safe sexual practices
- promotion of self-esteem and personal growth
- stress management

Important health promotion recommendations are to encourage patients to:[13]

- cease smoking
- reduce alcohol intake to safe levels
    - women no more than 2 standard drinks per day
    - men no more than 4 standard drinks per day
    - 3 alcohol-free days per week
- limit caffeine intake to 3 drinks per day
- increase regular physical activity
    - 30 minutes per day for three days per week, sufficient to produce a sweat
- reduce fasting plasma cholesterol to 4.0 mmol/L or less
- have a diastolic BP of less than 85 mmHg
- have a BMI of between 20 and 25 (see p. 76)
- reduce fat, refined sugar and salt intake in all food
- increase dietary fibre to 30 g per day
- build up a circle of friends who offer emotional support
- express their feelings rather than suppress them
- discuss their problems regularly with some other person
- work continuously to improve their relationships with people
- not drive a car when angry, upset or after drinking
- have a 2-yearly Pap smear
- avoid casual sex
- practise safe sex
- have an HIV antibody check before entering a relationship

REFERENCES

1. Piterman L, Sommer SJ. *Preventive Care*. Melbourne: Monash University, Department of Community Medicine, Final Year Handbook, 1993: 75 - 85.
2. National Health Strategy. *The Future of General Practice*. Issues paper No 3. Canberra: AGPS, 1992: 54 - 169.
3. Stott N, Davis R. The exceptional potential in each primary care consultation. JR Coll Gen Pract, 1979; 29: 201 - 5.
4. Sales M. Health promotion and prevention. Aust Fam Physician, 1989; 18: 18 - 21.
5. Health Targets and Implementation (Health for All) Committee. Health for All Australians. Canberra: AGPS, 1988.
6. Harris M. SNAP. *A Population Guide to Behavioural Risk Factors in General Practice*. South Melbourne: RACGP, 2004.
7. Shank JC, Murphy M, Schulte-Mowry L. Patient preferences regarding educational pamphlets in the family practice center. Fam Med, 1991; 23(6): 429 - 32.
8. Cates CJ. A handout about tetanus immunisation: influence on immunisation rate in general practice. Br Med J, 1990; 300(6727): 789 - 90.
9. Roland M, Dixon M. Randomised controlled trial of an educational booklet for patients presenting with back pain in general practice. JR Coll Gen Pract, 1989; 39(323): 244 - 6.
10. Rutten G, Van Eijk J, Beek M, Van der Velden H. Patient education about cough: effect on the consulting behaviour of general practice patients. Br J Gen Pract, 1991; 41(348): 289 - 92.
11. Howland JS, Baker MG, Poe T. Does patient education cause side effects? A controlled trial. J Fam Pract, 1990; 31(1): 62 - 4.

12. Murtagh J. *Patient Education* (4th edn). Sydney: McGraw-Hill, 2004.
13. Fisher E. The botch of Egypt: Prevention better than cure. Aust Fam Physician, 1987; 16: 187.

CHAPTER 12. PAIN AND ITS MANAGEMENT

> It is as much of the business of a physician to alleviate pain, and to smooth the avenues of death,
> when unavoidable, as to cure diseases.
>
> JOHN GREGORY 1725 - 1773, LECTURES ON THE DUTIES AND QUALIFICATIONS OF A PHYSICIAN

PAIN AND ITS MANAGEMENT

Pain is the capital symptom of humans - the great hallmark of disease - the signal par excellence to the patient and doctor that all is not well.

In modern medicine the successful management of chronic pain, in particular, still poses a great challenge. Its management or mismanagement is a yardstick of the excellence of that important bond - the doctor - patient relationship.

Pain is a multifactorial problem. The suffering patient must not only deal with the painful sensation itself but also cope with its possible serious significance.

- Does chest pain imply a pending heart attack?
- Does chronic ache or acute pain signify cancer?
- Does ‘whiplash’ imply neck pain to the grave?

Chronic pain is the most challenging problem. It must be emphasised that it always starts as an acute episode and that back pain accounts for the majority of cases of chronic pain encountered in general practice.

Golden rules:

- Acute pain ° acute anxiety.
- Chronic pain ° chronic depression.

DEFINITIONS

Pain is defined as ‘an unpleasant sensory and motional experience associated with actual or potential tissue damage or described in terms of such damage’. The box below defines the variety of types of pain.

Glossary of terms[1]

Allodynia

Pain due to a stimulus that does not normally provoke pain.

Mechanical - light touch feels painful.

Temperature - hot/cold stimulus (normally not painful) is painful.

Analgesia

Absence of pain in response to stimulation that would normally be painful.

Anaesthesia dolorosa

Pain in an area or region that is anaesthetic.

Causalgia

A syndrome of sustained burning pain, allodynia and hyperpathia after a traumatic nerve lesion, often combined with vasomotor and sudomotor dysfunction and later trophic changes (now known as complex regional pain syndrome II).

Central pain

Pain associated with a lesion of the central nervous system.

Dysaesthesia

An unpleasant abnormal sensation, whether spontaneous or evoked (e.g. formication - a feeling like ants crawling on the skin).

Hyperaesthesia

Increased sensitivity to stimulation, excluding the special senses.

Hyperalgesia

An increased response to a stimulus that is normally painful (i.e. painful stimulus feels much more painful than expected, such as firm finger pressure).

Hyperpathia

A painful syndrome, characterised by an increased reaction to a stimulus, especially a repetitive stimulus, as wellas an increased threshold for sensory detection.

Hypoaesthesia

Decreased sensitivity to stimulation, excluding the special senses.

Hypoalgesia

Diminished pain in response to a normally painful stimulus.

Neuralgia

Pain in the distribution of a nerve or nerves.

Neuritis

Inflammation of a nerve or nerves.

Neuropathy

A disturbance of function or pathological change in a nerve.

Paraesthesia

Abnormal sensation, whether spontaneous or evoked.

Phantom pain

The sensation of the presence of a missing body part.

Somatoform pain

Pain that has the qualities of pain arising from a physical (somatic) cause but not attributable to any objectively demonstrable organic causation (i.e. the expression of psychological distress as physical symptoms).

Origins of pain[1, 2]

In general terms, the origin of conscious pain can be subdivided into three broad types - nociceptive, neurogenic or psychogenic.

1. *Nociceptive pain* is pain arising from stimulation of superficial or deep tissue pain receptors (nociceptors) from tissue injury or inflammation. It requires an intact nervous system. *Nociception* is stimulation of peripheral nociceptors (i.e. nerve endings sensitive to a noxious stimulus).
2. *Neurogenic pain* is pain caused or initiated by a primary lesion or dysfunction (i.e. damage) in the peripheral or central nervous system. It can be subdivided into central pain, when the primary lesion is in the central nervous system, or peripheral pain. Neuropathic pain is a form of neurogenic pain in which there is actual nerve cell or axonal damage due to inflammation, trauma or degenerative disease.
3. *Psychogenic pain* is pain arising in the absence of any discernible injury and where the predominant aetiology is psychological or psychiatric. However, it is a realand distressing entity to the patient.

It should be emphasised that these types may overlap and patients may have more than one type of pain.

The features of the main types of pain are summarised in Table 12.1.

Table 12.1 Features of main types of pain[2]

Type of pain	Origin of pain	Symptoms and signs	Examples
Nociceptive			
1 Somatic	Skin mucosa	Localised stinging or burning	Skin ulcers/abscess
			Aphthous ulcers
	Bones and joints	Dullache	Arthritis
	Pleura and peritoneum	± Pain on movement	Minor fractures

			Pleurisy
2 Visceral	Solid or hollow organs	Deep, diffuse pain	Abdominal tumours
		Poorly localised	Bowel obstruction
		± Colic	Major surgery
		± Nausea and vomiting	Ureteric/biliary colic
3 Muscle spasm	Skeletal muscle	Pain worse on movement	Acute low back pain (some)
	Smooth muscle	Severe colic	Ureteric/biliary colic
			Bowel obstruction
		Tenesmus	Rectal dysfunction
Neurogenic			
1 Peripheral nerve	Nerve compression	Pain in nerve distribution	Prolapse IV disc
		± Motor deficit	Early brachial/lumbar plexus tumour
		± Sensory deficit	
			Infiltration
2 Neuropathic pain	Nerve damage	Various pain syndromes (see text)	Post-herpetic neuralgia
		± Trophic changes	Peripheral neuropathy
			Late plexus infiltration

Components of pain[2]

A conceptual approach to the components of pain is illustrated in Figure 12.1.



Figure 12.1 A conceptual approach to the components of pain: a biopsychosocial model

Developing an understanding of pain depends on the knowledge of relationships among nociception, pain, suffering and pain behaviour. The first three components cannot be measured or completely under stood in individuals and only the pain behaviour can be observed and measured by parties other than the patient.

CATEGORIES OF PAIN

Pain can be described simply as:

- acute pain
- cancer pain
- chronic non-cancer pain

Acute pain

Acute pain is pain of rapid onset and short duration that usually has an obvious cause and a predictable duration. Early diagnosis and treatment is usually successful. It is usually opioid sensitive. It is almost always due to nociception.

Pain behaviour may be modified by socioenvironmental factors (e.g. appropriate prepa ration for an operation results in lower postoperative analgesic requirements). Psychological factors are not usually a significant factor.

Early and appropriate interventions by GPs can have a major impact in improving the assessment and treatment of pain and helping to prevent the development of chronic pain, as in the chronic pain cycle. Th is applies particularly to acute musculoskeletal injury, such as acute neck and back pain and sporting injuries.

Problem areas in acute pain management include:

- acute back pain (Chapter 37)
- acute neck pain (Chapter 62)
- acute headache (Chapter 56)
- acute orofacial pain (Chapter 52)
- acute abdominal pain (Chapter 34)
- acute haemarthrosis (e.g. knee in haemophilia/sports injuries)
- acute chest pain (Chapter 40)
- sports injuries (Chapter 136)
- trauma
- burns
- neuropathic pain (e.g. herpes zoster)

Chronic pain

Chronic pain may be defined as pain present for a period greater than 3 months or pain present for 4 weeks more than the expected time of recovery.

Chronic pain is the classic 'heartsink' problem of practitioners as it is far more difficult to manage and is often unresponsive to agents effective in acute pain relief, such as opioids. It is, in itself, a type of disease state.

The management of chronic pain may require a multidisciplinary pain clinic strategy in which nociceptive, neuropathic, psychological and environmental contributions to pain behaviour can be evaluated and managed.

The most demanding chronic pain problems include:

- chronic back pain
- osteoarthritis
- rheumatoid arthritis
- chronic recurrent headache

ASSESSING AND MEASURING PAIN

Pain is a subjective symptom and generates considerable emotion and frustration, and thus can be difficult to assess.

The history

The classic historical approach is still the most important approach. Patients can use the PQRST approach to describe their pain,[2] namely, P - palliative and provocative factors, Q - quality, R - radiation, S - severity and T - timing. Most practitioners use the SROT-SARA approach to pain, namely:

- site
- radiation
- onset and off set
- type (quality)
- severity
- aggravating factors
- relieving factors
- associations

Use of body charts

To assist with diagnosis and ongoing management of pain, it is helpful to get patients to chart the site and radiation of their pain on body charts - either total body charts or regional charts (e.g. the head). This is particularly helpful for spinal pain with referral patterns and fibromyalgia.

Measurement of pain

Despite its subjective nature, it is good to record some type of repeatable measurement, especially for chronic pain requiring treatment outcomes. This involvement also helps the patient.

Unidimensional scales

Visual analogue scales (VAS), which are used as a research tool and well validated, can be useful in recording both acute and chronic pain levels. An example of a simple VAS linear scale on which the pain indicates the severity of their current pain is shown in Figure 12.2. A pictorial (faces) scale applicable to children is shown in Figure 12.3.


Mark the areas on your body where you feel the various sensations
pain
numbness
pins and needles
intolerable pain
10
9
8
7
6
5
moderate pain
4
3
2
1
0
no pain
Mark your level of pain on tthis scale
Back
Front
Left
Right
Right
Left

Figure 12.2 Assessing pain using a visual analogue scale and body chart: ideal for lumbosacral pain


0
No hurt
1
Hurts little bit
2
Hurts little more
3
Hurts even more
4
Hurts whole lot
5
Hurts worst

Figure 12.3 Visual analogue scale. A faces pain rating scale ideal for children

Multidimensional scale

These scales, which are usually employed for chronic pain, take into account several aspects of pain perception in addition to assessing functional effects and levels of disability. Examples include the:

- McGill Pain Questionnaire
- Pain Disability Index
- Form 36 Health Survey (SF-36)

- Oswestry low-back pain questionnaire

THE HOLISTIC APPROACH TO PAIN MANAGEMENT

The principles of this approach are outlined in Chapter 13, in which a case history approach for the perennial problem of back pain is considered.

Doctors should assess not only the somatic basis of the pain but also the patient's psyche, mood, coping ability, attitude and family.

The pain threshold can be lowered by factors such as fatigue, anger, depression, loneliness, home or work environment. On the other hand, the factors that raise the pain threshold include a happy or contented disposition, empathy, companionship, sleep, rest, pleasant diversions and good home support.[3]

It is useful to have a therapeutic model on which to base an orderly approach to management and the following set of guidelines on the management of arthritic disorders is one such example.

Management of arthritic disorders

General principles

1. Explanation/patient education and appropriate reassurance.
2. Attention to lifestyle - the NEAT approach:
    - Nutrition - optimal diet, ideal weight, low fat, low sugar etc.
    - Exercise - a balanced lifestyle program
    - Avoidance of toxins - avoid or limit alcohol, nicotine, other drugs
    - Tranquillity - relaxation techniques
3. Shared care - multidisciplinary approach.
   Examples include:
    - Rheumatologist, especially for chronic problems (e.g. systemic lupus erythematosus [SLE], rheumatoid arthritis [RA])
    - Physiotherapist - important for most disorders
    - Occupational therapy
    - Pain clinic
4. Pharmacological management, for example:
    a. simple analgesics (e.g. paracetamol for pain; consider aspirin)
    b. non-steroidal anti-inflammatory drugs (NSAIDs) for arthritic symptoms
    c. disease-modifying agents:
        - sulphasalazine for RA, spondyloarthropathies
        - hydroxychloroquine - SLE, RA
        - injectable/oral gold - RA
        - D-penicillamine - RA
        - methotrexate - RA, spondyloarthropathies
        - cyclosporin - RA
        - cyclophosphamide - RA
        - azathioprine - RA, SLE
        - leflunomide - RA
        - biological DMARDS - RA
        - fish oil (long-chain omega-3 fatty acids) - RA
        - glucosamine and chondroitin - OA
    d. disease-suppressant agents:
        - corticosteroids
5. Intra-articular injections, for example:

    - corticosteroids
    - hylan

6. Surgery (e.g. synovectomy, joint replacement, arthrodesis)

ANALGESIC PRESCRIBING

The general practitioner has an important responsibility not only to relieve patients' pain skilfully and completely (if possible) but to ensure that the appropriate drugs are prescribed without leading to dependence.

The three categories of pain - nociceptive pain, neurogenic pain and psychogenic pain - each require different therapeutic agents.[4] For nociceptive pain the traditional analgesics are used: paracetamol, aspirin, other NSAIDs and opioids. Neurogenic pain is treated with antidepressants, antiepileptics and membrane stabilisers. Agents used to treat muscle pain are muscle relaxants and baclofen.

Medications used to treat pain can be considered as:

- ‘simple’ analgesics (aspirin, paracetamol)
- NSAIDS, cyclo-oxygenase-2 (COX-2) specific inhibitors
- opioids
- antidepressants (amitriptyline)
- membrane stabilisers (anticonvulsants, antiarrhythmics)
- NMDA blockers (ketamine)

Chronic pain, such as chronic low back pain, represents a challenge but the challenge is to manage acute pain skilfully and prevent chronicity. O'Reilly[5] has provided some useful guidelines for the role of the general practitioner in the management of chronic pain:

- Emphasise that cure is unlikely.
- Encourage the patient to accept responsibility for pain management.
- Provide ongoing support and interest.
- Promote activity rather than rest.
- Educate patients about their condition: review X-rays and discuss drugs, especially the longterm disadvantages of opioids and sedatives.
- Encourage time-contingent versus paincontingent use of medication.
- Withdraw unsuitable drugs (e.g. opioids and sedatives).
- Consider use of undiagnosed depression or anxiety as amplifiers of pain perception and treat where necessary.
- Promote the use of relaxation, distraction and heat or ice packs rather than increased medication.
- Discuss strategy for management of flare-ups in advance.
- Refer for psychological support: cognitive behavioural psychologist or psychiatrist recommended.

Analgesics in common use[2, 4]

Paracetamol

Paracetamol has minimal anti-inflammatory acti vity but moderate analgesic (equipotent with aspirin) and antipyretic properties. It is metabolised by the liver and has an excretion half-life of approximately 4 hours. Adverse effects are uncommon but gastrointestinal discomfort such as dyspepsia and nausea can occur occasionally. It is well tolerated by patients with peptic ulcers and has no effect on platelet function. Use of 4 g or more per day over 12 months has been associated with chronic liver disease. It should be administered with caution in patients with kidney or hepatic dysfunction. Although less effective than NSAIDs for pain relief, its excellent safety in therapeutic doses makes it the first-line analgesic for mild to moderate pain.

Usual dosage: 1 g (o) 4 hourly (max. 4 g/day)

Formulations of paracetamol include:

- immediate release tablets and capsules
- oral solution and suspension
- extended-action formulation - 665 mg (may provide up to 8 hours pain relief)
- suppositories (125 mg, 250 mg, 500 mg forms)
- injectable formulation

Aspirin

Aspirin has both analgesic and anti-inflammatory activity and is a very effective drug in adults for mild to moderate acute pain. It has an extremely short half-life and is metabolised to salicylic acid, which shares the above properties. The major problems with aspirin are gastrointestinal discomfort, ulceration and bleeding.

Usual dosage: 600 mg (o) 4 hourly (max. 4 g/day)

Opioid analgesics

These agents are usually reserved for the treatment of severe pain. Commonly used agents are the weaker opioids - codeine, oxycodone, dextropropoxyphene - and the stronger ones - morphine, pethidine and methadone. Other types are dextromoramide, pentazocine, paraveretum and fentanyl.

Adverse effects, which are common, dose dependent and quite variable, include nausea and vomiting, constipation, respiratory depression and dysphoria.

Codeine

Codeine, which is methylmorphine, is metabolised to morphine. Controlled trials have shown codeine 32 mg to be no more effective than aspirin 600 mg.[6] Disturbing side-effects include nausea and constipation.

Usual dosage: 30 - 60 mg (o) 4 hourly (max. 180 mg/day)

Oxycodone

Oxycodone is a synthetic opioid that is very effective orally. It is useful for moderate pain in bridging the gap between the simple analgesics and strong opioids.[6] The oral form has a duration of action of 4 hours. It is well absorbed through the rectal mucosa and is useful as a night-time suppository with a duration of 7 - 8 hours.[2]

Adverse side-effects include nausea, constipation, confusion and itch.

Usual dosage: 10 mg (o) 4 hourly (max. 60 mg/day) 30 mg rectally 12 hourly

Dextropropoxyphene

This controversial drug is structurally related to methadone. Adverse effects include dysphoria, confusion, lightheadness and constipation. It has been associated with relatively sudden death when taken in overdose, especially with alcohol.[2] Continuous use should be discouraged, particularly in elderly patients and those with cardiac disease.[2]

Morphine

Morphine is the most effective opioid for the relief of cancer pain (Chapter 7). It is worth noting that injections are not more effective than oral administration in achieving pain relief. It should be titrated according to patients' needs. Dose reduction may need to be made in patients with impaired liver or kidney function and in elderly or debilitated patients.

Note: Dose requirements vary considerably and patient response should be monitored frequently. There may be no upper limit to dosage in cancer patients.

Usual dosage:[2] 10 mg (o) 4 times daily IM or SC injection: 7.5 - 15 mg intravenous: bolus - 2.5 - 5 mg followed by 1 - 2 mg increments at 5 minute intervals slow - 2.5 - 5 mg over 5 - 10 minutes, then increments infusion - 2 - 5 mg per hour

Methadone

Methadone is an effective oral analgesic with a long but variable half-life; it is given, preferably, once or at most twice a day. Unsuitable for acute pain, it is valuable for chronic pain, although longacting morphine preparations have displaced it. It should not be used in elderly patients or those with kidney dysfunction. Its place is in management of opioid dependency but it needs to be used with care because of the risk of respiratory depression and accumulation.

Pethidine

Pethidine is a synthetic opioid with a short duration of action. A problematic adverse effect is accumulation of its toxic metabolite (norpethidine) which can cause myoclonic and general seizures.[6] It has no place in the management of chronic pain, whether cancer or non-cancer and possibly no place in acute pain.

Fentanyl[2]

Fentanyl is a very potent synthetic opioid which can be administered IV, IM, SC, transdermal or by the epidural route. Its efficacy is similar to morphine but it has fewer side-effects. The conversion factor is: 10 mg (SC) morphine ° 150 µg (SC) fentanyl.

Usual dosage: preoperation/postoperation pain, 50 - 100 µg IM or slow IV patches (various strengths, deliver over 72 hours), initial dose 25 µg/hour

Hydromorphone

This is structurally similar to morphine but 5 to 7 times more potent. It is available for oral, parenteral and intraspinal use in moderate to severe pain. The risk of drug dependence is high.

Usual dosage: 2 - 4 mg (o) every 4 hours 1 - 2 mg IM, SC or IV (slow) every 4 - 6 hours

Tramadol

Tramadol is an atypical analgesic with both opioid and non-opioid features. Its use has increased dramatically. Its advantages include analgesia with minimal sedation or respiratory depression and low abuse/dependence potential. Side-effects include dizziness, vertigo, nausea, vomiting, constipation, headache, somnolence, tremor, confusion and hypersensitivity reactions. Seizures have been reported.[7]

Usual dosage: (as 50 mg caps or SR tabs - 100, 150 or 300 mg) 50 - 100 mg (o) 2 - 4 times daily or 100 - 200 mg (SR) twice daily injection 100 mg/2 mL: 50 - 100 mg 4 - 6 hourly IM or slow IV

Buprenorphine

This is a mixed agonist - antagonist opioid derived from the opium alkaloid thebaine. It has very limited use in pain management. Sublingual preparations (0.2 mg, 0.4 mg, 2 mg, 8 mg) are used for acute, chronic and cancer pain. It is also available as a transdermal patch, e.g. Norspan and injection formulation. It is used in the treatment of opioid dependence.

Pentazocine

This is also a mixed agonist - antagonist opioid and not structurally related to the other opioids. It is usually given orally or by SC/IM injection. Its use is not recommended.[2]

Combined analgesics

Combined analgesics are freely available both as prescribed and over-the-counter (OTC) preparations for moderately severe pain. They usually consist of a simple analgesic such as paracetamol or aspirin combined with an opioid analgesic (usually codeine). The analgesics have an additive effect because they act at different receptor sites. The use of such preparations is generally not recommended but if both agents are needed it is advisable to prescribe them separately, thus enabling individual dose adjustments.[2]

Methoxyflurane

This is administered as inhalational analgesia with the Penthrox Inhaler in emergency situations such as at the roadside. It can be used with oxygen or air. It provides pain relief after 8 - 10 breaths and it continues for several minutes.

PAIN IN CHILDREN

The management of pain in children requires considerable sensitivity and wisdom. There are many clinical myths about this issue, especially that children have a high pain tolerance and analgesia is relatively dangerous. The inappropriate pain management of circumcision is a classic example.

Guiding principles

- All children feel pain - as much if not more than adults.
- Neonates do feel and remember pain - even the fetus feels pain.
- Children can localise and describe their pain.
- Pain in children can be masked and they often do not report pain.
- If a child complains of pain, they are serious and it is organic until proved otherwise.

- Consider other confusing factors (e.g. developmental delay).
- Consider children's fear of analgesic injections.
- Anxiety from anticipation of a procedure is a big issue - explanation, distraction and parental involvement help.
- Adverse effects of opioids, such as addiction, do not necessarily pose serious problems in children.[2]
- Analgesic medication should be dosed on a per-weight basis.
- Consider analgesic combinations with more severe pain (e.g. NSAID plus opioid).

Assessment of pain

This is very important in children of allages and should involve a careful history and examination. Selfreporting of pain is reliable in children over 4 years of age. Close observation of non-verbal cues and behaviour is important (e.g. severe pain may be indicated by quiet withdrawal and whimpering; tachycardia).

Scaling strategies can be used - the modified faces pain scale (Figure 12.3 on p. 99) is useful in younger children, while older children and teenagers can use a visual analogue scale.

Analgesia in children[2]

The three most commonly used analgesics in children are paracetamol, NSAIDs and opioids. Table 12.2 lists the hierarchy of analgesics to be used in children.

Table 12.2 The analgesic hierarchy[8]

- Paracetamol
- NSAID (ibuprofen, naproxen)
- Combination oral therapy:
    - paracetamol/codeine mixes
    - alternate: NSAID/paracetamol
- Parenteral opioid
    - bolus IM, IV; infusion; continuous PCA
- Combination parenteral therapy:
    - NSAID/opioid/ketamine
    - adjuvant clonidine

Paracetamol

This is generally safe and effective even in asthmatics in therapeutic doses. It is rapidly absorbed orally within 30 minutes and well absorbed rectally.

Usual dosage: 15 mg/kg (o) every 4 hours (max. 6 doses/day) 20 mg/kg rectally, every 6 hours

Rules: maximum daily dose:[7]

- 90 mg/kg to max. 4 g (for 2 days only)
- 60 mg/kg/day if aged < 6 months a dose of 10 mg/kg is no more effective than placebo

- Don't exceed recommended daily doses.
- Don't use continuously > 48 hours without review.
- Don't double up, e.g. use cold and flutablets with paracetamol.
- Don't use in a trivial fever.

Cautions:

- Jaundice
- Repeated doses over several days when febrile and/or dehydrated

Hepatotoxicity is rare and does not usually occur in doses below 150 mg/kg/day but acute paracetamol overdose (single doses of > 100 mg/kg) is a potentially life-threatening event.

Aspirin

Aspirin is not in common use in children and should not be used < 18 years since it was associated with Reye's syndrome.

For antiplatelet effects use low doses 2 - 5 mg/kg/day.

NSAIDs

NSAIDs have a proven safety and efficacy in children for mild to moderate pain and can be used in conjunction with paracetamol and opioids such as codeine and morphine. The advantage is their opioid-sparing effect. Contraindications include known hypersensitivity, severe asthma (especially if aspirin sensitive), bleeding diatheses, nasal polyposis and peptic ulcer disease.

Those commonly used for analgesia are:

- ibuprofen: 5 - 10 mg/kg (o) 6 - 8 hourly (max. 40 mg/kg/day)
- naproxen: 5 - 10 mg/kg (o) 12 - 24 hourly (max. 1 g/day)
- indomethacin: 0.5 - 1 mg/kg (o) 8 hourly (max. 200 mg/day)
- diclofenac: 1 mg/kg (o) 8 hourly (max. 150 mg/day)

The rectal dose is double the oral dose (e.g. indomethacin 2 mg/kg) but only administered twice a day.

Opioid analgesics

Oral opioids

These have relatively low bioavailability but can be used for moderate to severe pain when weaning from parenteral opioids, for ongoing severe pain (e.g. burns) and where the IV route is unavailable.

Codeine

Usual dosage:

- 0.5 - 1 mg/kg (o), 4 - 6 hourly prn (max. 3 mg/kg/day)

More effective if used combined with paracetamol.

Morphine

Immediate release:

- 0.3 mg/kg (o) 4 hourly prn

Sustained release:

- 0.6 - 0.9 mg/kg, 12 hourly

Tramadol

Usual dosage:

- 1 - 2 mg/kg (o) 4 hourly

Oxycodone

Immediate release:

- 0.2 - 0.3 mg/kg (o) 4 hourly (max. 10 mg)

Sustained release:

- 0.6 mg/kg (o) 12 hourly

Hydromorphone

Usual dosage:

- 0.04 mg/kg (o) 4 hourly

Fentanyl

Fentanyl citrate can be administered orally (transmucosal) as ‘lollipops’ or transcutaneous as ‘patches’.

Parenteral opioids[9]

These are the most powerful parenteral analgesics for children in severe pain and can be administered in intermittent boluses (IM, IV or SC) or by continuous infusion (IV or SC). Infants under 6 months are more sensitive and need careful monitoring (e.g. pulse oximetry). This management is invariably in the hospital. Administration of parenteral opioid should not be undertaken without the availability of oxygen, resuscitation equipment and naloxone to reverse overdose.

Maximum dosage of IM opioids:

- morphine: 0.2 mg/kg (max. 10 - 15 mg), 4 hourly prn
- pethidine: 2 mg/kg (max. 25 - 100 mg), 3 hourly prn

ANALGESICS IN THE ELDERLY

As a general rule, most elderly patients are more sensitive to opioid analgesics and to aspirin and other NSAIDs but there may be considerable individual differences in tolerance between patients. Patients over 65 years should receive lower initial doses of opioid analgesics with subsequent doses being titrated according to the patient's needs.[2]

Some general rules and tips[2]

- Give analgesics at fixed times by the clock rather than ‘prn’ for ongoing pain.
- Regularly monitor your patient's analgesic requirements and modify according to needs and adverse effects.
- Start with a dose towards the lower end of the dose range and then titrate upwards depending on response.
- Provide ongoing interest and support. This will magnify any placebo effect.
- Avoid using compound analgesics and prescribe simple and opioid analgesics separately.
- Never cut suppositories in half with the intention of halving the dose.
- Reserve the use of antiemetics for the development of nausea and vomiting with opioids. Extrapyramidal reactions (dystonia and oculogyric crises) can be a problem with antiemetics.
- Advise patients about the benefits of high-fibre foods if on analgesics. Prescribe a bulking agent or lactulose if necessary.

NON-STEROIDAL ANTI-INFLAMMATORY DRUG PRESCRIBING

The prescribing of aspirin and other NSAIDs is an area of increasing concern to all general practitioners, especially in the elderly with their increasing propensity to the pain of arthritic conditions and demands for relief. About 15% of the patients on a general practitioner's list in the United Kingdom will present with a locomotor problem.[10] A common response to the symptoms of rheumatic complaints is to prescribe NSAIDs; in the United States 55% of all visits by patients with arthritis will lead to a prescription for an NSAID.[11]

Unfortunately, the use of NSAIDs involves a high incidence of side-effects, ranging from the trivial to the lethal, with many deaths from bleeding ulcers, especially in the elderly. The NSAIDs ‘epidemic’ prevails because the agents continue to be perceived as the best available for the relief of arthritic symptoms. Of particular concern, however, is the widespread use of NSAIDs for common problems such as back pain when the main cause is dysfunctional or mechanical without evidence of inflammation. Surprisingly, many of these patients appear to achieve a good response to NSAIDs.

The choice of NSAID can be controversial. Some are claimed to have more adverse effects than others but there is little evidence of substantial differences.[4, 12] There is no doubt that particular NSAIDs are of great value in specific rheumatic conditions. Both indomethacin and phenylbutazone are more effective for the spondyloarthropathies and gout.[12] It is worthwhile being mindful of their half-lives. Those with short half-lives include aspirin, diclo fenac, tiaprofenic acid, keto-profen, ibuprofen and indomethacin. NSAIDs with long half-lives include diflunisal, naproxen, sulindac, piroxicam and tenoxicam, and these are useful in the treatment of chronic pain such as bony metastases in cancer.

Table 12.3 shows the classification of NSAIDs.

Table 12.3 Classification of NSAIDs[2]

Action	Example
Non-selective inhibitors of COX-1 and COX-2, mainly in CNS	Paracetamol
Non-selective inhibitors of COX-1 and COX-2, acting in both CNS and periphery	Aspirin Other NSAIDs
Specific inhibitors of COX-2	Celecoxib Lumiracoxib

Preferential inhibitors of COX 2 over COX-1	Meloxicam
COX ° cyclo-oxygenase	

Studies have identified certain patients who appear to have significant risk factors for the likely development of NSAID lesions (gastropathy). NSAID ulceration and its complications (Table 12.4) are more prevalent in the elderly, particularly in females.

Table 12.4 Persons at higher risk from NSAID-induced side-effects (after Ryan)[12]

Definite
Age > 65 years
Prior ulcer disease or complication
High-dose, multiple NSAIDs
Concomitant corticosteroid therapy
Duration of therapy (< 3 months)
Possible
Condition necessitating NSAID treatment (e.g. RA)
Female sex
Smoking
Alcohol excess
Helicobacter pylori

It is likely that NSAIDs have been prescribed too readily in Australia, particularly in general practice.[12] In a review,[13] a reduction of NSAID prescribing was recommended, particularly for conditions where inflammation is not a major feature (e.g. osteoarthritis and 'mechanical' back pain). Consumption has declined, probably in part because of media publicity. Haslock emphasises that those who both stop and start NSAIDs must recognise some clinical responsibilities.[14]

Those who start the drug have obligations to:

- ensure there is an appropriate indication for it
- provide education about the drug to the patient - its purpose, side-effects and dosage regimen
- monitor its continuing need, effectiveness and safety
- give the patient permission to telephone about any concerns
- be prepared to deal with side-effects

Those who stop medication have an equal obligation to:

- ascertain whether there is still a need for its therapeutic action
- take the necessary steps, if the need exists, to provide an alternative

There is evidence that peptic ulcers that develop in patients taking NSAIDs heal faster if the NSAID is dropped.[15]

There is no evidence that using an enteric-coated NSAID preparation or using the rectal route for administration reduces gastric damage as it is mediated almost entirely systemically after the NSAID has been absorbed.[12] Furthermore, trials have indicated that the efficacy of using H_2-receptor antagonists for preventing NSAID gastrointestinal complacations is low to absent.

An anti-inflammatory dose of fish oil should be considered as part of long-term treatment to minimise NSAID use.

The coxibs are a group of NSAIDs synthesised to inhibit COX-2 specifically. They are on a par as an analgesic with the COX-1 inhibitors. Their gastrointestinal adverse reactions are less, but they have all the other adverse effects and drug interactions of COX inhibitors. The cardiovascular problems, including increased blood pressure, thrombosis (fatal myocardial infarction and stroke) and impairment of kidney function, experienced with rofecoxib indicates the potential problems of these agents. If prescribed for patients at risk of thrombosis aspirin should be added.[16]

Parecoxib is a COX-2 agent that is very effective for postoperative pain as an IM or IV injection. Newer agents include lumiracoxib and etoricoxib.

NSAIDs and the elderly

NSAIDs with short half-lives (e.g. ibuprofen and diclofenac), may be safer in the elderly and all NSAIDs should be used in reduced dosage. Patients should be monitored for fluid retention and hypertensive control.[2, 17]

Prescribing recommendations

- Recommend intermittent use of NSAIDs based on appropriate patient education. Intermittent courses for 14 days can work well in chronic conditions, remembering that it takes about 10 days for NSAIDs to achieve maximal effectiveness.
- Use paracetamol as an alternative, especially for osteoarthritis conditions.
- Consider protective combinations (e.g. diclofenac + misoprostol).

NEUROPATHIC PAIN

Neuropathic pain can be defined as pain associated with injury, disease or surgical section of the peripheral or central nervous system. It is a common (affects about 1% of the population) although underidentified condition.[18]

Considerations regarding neuropathic pain include:

- essentially due to nerve damage
- frequently involves both peripheral and central sensitisation
- has a multiplicity of clinical pain features
- many responsible mechanisms (e.g. neuroma, demyelination, loss of normal sensory input, enhanced sympathetic activity)

Clinical features[1]

- Burning, shooting, pulsating or stabbing pain
- Paroxysmal or spontaneous pain
- Pain in the absence of ongoing tissue damage
- Pain in an area of sensory loss
- Allodynia (e.g. pain on light touch)
- Hyperpathia
- ± Hyperalgesia
- ± Dysaesthesia (e.g. ‘ants crawling on skin’)
- ± Delay in onset of pain after nerve injury
- Radiating electric shock like pains in nerve distribution (e.g. Tinel's sign - tapping of neuroma or nerve)
- Often refractory to simple analgesics and NSAIDs
- Poor response to opioids
- ± Autonomic nervous system instability (e.g. pallor or cyanosis, excessive warmth or cold, sweating changes)

Examples of neuropathic pain states include:

- post-herpetic neuralgia
- trigeminal neuralgia (‘tic douloureux’)
- atypical facial pain
- phantom limb pain
- complex regional pain syndrome I (reflex sympathetic dystrophy)
- complex regional pain syndrome II (causalgia)
- other neuralgias
- diabetic/alcoholic peripheral neuropathy
- spinal cord injury pain
- late brachial/lumbar sacral plexus tumour infiltration
- post-stroke pain
- arachnoiditis

Treatment

For initial pain relief, use aspirin or paracetamol or a NSAID. However, it is usual to depend on the use of analgesic adjuvants, either orally (mainly tricyclic antidepressants [TCAs] and antiepileptics) or parenterally, such as lignocaine or ketamine. Adjuvants should be started in low doses and increased as necessary to the maximum tolerated level.

amitriptyline 10 - 25 mg (o) nocte increasing every 7 days to 75 - 100 mg max.

Consider another TCA (e.g. nortriptyline or doxepin)[2] *and/or* carbamazepine 50 - 100 mg (o) bd initially increasing to 400 mg bd max. *or* gabapentin 300 mg (o) daily (nocte) initially, increasing as tolerated to 3 times daily (max. 2400 mg)

Note:

- It is renally excreted, therefore caution in elderly and renally impaired.
- Try a test dose of 100 mg (o) at night in the elderly.
- Side-effects include drowsiness, dizziness and generalised fatigue. *or*
  sodium valproate 200 - 600 mg (o) bd *or*
  pregabalin 150 mg (o) daily initially
  If refractory consider the cardiac drugs.
  mexiletine 50 - 200 mg (o) tds
  flecainide 50 mg (o) bd increasing to 300 mg max. per day

Some guiding rules[2]

- TCAs may be better for constant burning pain.
- Antiepileptics may be better for sharp shooting pain.
- Carbamazepine is the drug of choice for trigeminal neuralgia.
- Doses of TCAs are much smaller than those for treating depression.

Evidence base

- Level 1 evidence shows that about 70% of patients have a significant response to carbamezepine.[18, 19]
- The newer antiepileptic gabapentin has proven efficacy for diabetic neuropathy[20] and post-herpetic neuralgia.[21]

Painful diabetic neuropathy[22]

It is important to realise that painful diabetic neuropathy tends to occur in mild or even unrecognised cases of diabetes and sometimes may be a presenting symptom. Pain in the feet and legs is found in 11.6% of people with type 1 diabetes and in 31.2% of people with type 2 diabetes.

The classic complaint is burning in the feet with possible associated aching, cramping and tingling sensations. The symptoms are often worse during the night. Non-diabetic causes of painful neuropathy can include deficiency states associated with alcoholism and vitamin B_{12} deficiency, uraemia and ischaemic neuropathy associated with peripheral vascular disease. First-line treatment is TCAs and then gabapentin.

SOMATOFORM PAIN DISORDER

In somatoform pain, which is sometimes referred to as psychogenic pain, the patient complains of severe and distressing pain that has the qualities of pain arising from a physical (somatic) cause, but which cannot be attributed to objectively demonstrable organic pathology.[2] Pain occurs in association with emotional conflict or psychosocial problems deep enough to point to their possible causation.

Differential diagnoses include occult organic pain, depression, substance abuse, malingering and rare disorders such as sickle cell anaemia and porphyria. Management is difficult and is based on caring support devoid of promoting the ‘sick role’. Psychological treatments are directed towards helping the patient to cope and ‘live with the pain’. The possibility of comorbid depression should be addressed and if it is possible an appropriate trial of antidepressants can be tried. Referral for CBT or similar psychotherapies or to a pain clinic are options.

REFERENCES

1. Gray M, Cousins M. Pain. CHECK, 1994; issue 264: 3 - 18.
2. Mashford ML (Chair). *Therapeutic Guidelines: Analgesic* (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2006.
3. Turk DC, Meichenbaum D. *Textbook of Pain* (2nd edn). London: Churchill Livingstone, 1989: 1001 - 9.
4. Moulds RFW (Chair). *Analgesic Guidelines* (3rd edn). Melbourne: Victorian Medical Postgraduate Foundation, 1997 - 98.
5. O'Reilly S. Treatment of chronic non-malignant pain in general practice. Aust Fam Physician, 1994; 23: 281.
6. Crammond T. Pain relief. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 380 - 5.

7. NPS. Minimising the risks of using analgesics for musculoskeletal pain. NPS News 28, 2003; 5 - 12.
8. Moloney G. Paediatric analgesia. Curr Ther, 2001; August: 67 - 71.
9. Smart J. *Paediatric Handbook* (6th edn). Melbourne: Blackwell Science, 2000: 55 - 6.
10. Khalig N, Wood PNH. *Arthritis and Rheumatism in the Eighties*. London: Arthritis & Rheumatism Council, 1986.
11. Baum C, Kennedy DL, Forbes MB. Utilisation of NSAIDs. Arthritis Rheum, 1985; 28: 686 - 92.
12. Ryan P. Pharmacology of musculoskeletal treatment agents. Monash Distance Education: Graduate Diploma in Family Medicine. Musculoskeletal Medicine Notes, 2001; 6(1): 10 - 13.
13. Brooks PM, Yeomans ND. NSAIDs gastropathy: is it preventable? Aust NZ Med, 1992; 22: 685 - 91.
14. Haslock I. Should we use NSAIDs? Ailment Pharmacol Therap, 1988; 25: 1 - 8.
15. Lancaster-Smith MJ, Jaderberg ME, Jackson DA. Ranitidine in the treatment of NSAIDs-associated gastric and duodenal ulcers. GUT, 1991; 32: 252 - 5.
16. Day RO, Graham G. The vascular effects of COX - 2 selective inhibitors. Aust Prescriber, 2004; 27: 142 - 5.
17. Scharf S, Christophidis N. Drug treatment in the elderly. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 159 - 60.
18. Bashford GM. The use of anticonvulsants for neuropathic pain. Aust Prescriber, 1999; 22(6): 140 - 1.
19. Nicol CF. A four year double-blind study of Tegretol in facial pain. Headache, 1969; 9: 54 - 7.
20. Backonja M, Beydoun A et al. Gabapentin for the symptomatic treatment of painful neuropathy in patients with diabetes mellitus. JAMA, 1998; 280: 1831 - 6.
21. Rowbottom M, Harden N et al. Gabapentin for the treatment of postherpetic neuralgia. JAMA, 1998; 280: 1837 - 42.
22. Gouche R. Neuropathic pain. Part 1. Med Observer, 2001; 22 June: 44 - 5.

CHAPTER 13. WHOLE PERSON APPROACH TO MANAGEMENT

> Never forget that it is not a pneumonia, but a pneumonic man who is your patient.
>
> Sir WILLIAM GULL (1816 - 90)

WHOLE PERSON APPROACH TO MANAGEMENT

The management of the whole person, or the holistic approach, is an important approach to patient care in modern medicine. Whole person diagnosis is based on two components:[1]

1. the disease-centred diagnosis
2. the patient-centred diagnosis

The disease-centred consultation is the traditional medical model based on the history, examination and special investigations, with the emphasis on making a diagnosis and treating the disease. The diseasecentred diagnosis, which is typical of hospital-based medicine, is defined in terms of pathology and does not focus significantly on the feelings of the person suffering from the disease.

The patient-centred consultation not only takes into account the diagnosed disease and its management but also adds another dimension - that of the psychosocial hallmarks of the patient (Table 13.1), including details about:[1]

- the patient as a person
- emotional reactions to the illness
- the family
- the effect on relationships
- work and leisure
- lifestyle
- the environment

Table 13.1 Whole person diagnosis and management

Disease-centred diagnosis	Patient-centred diagnosis
• aetiology of disease	• significance of illness to patient • effect on family and relationships • effect on work and income • psychological effects - stress and anxiety - abnormal illness/behaviour - sleep - depression • effect on sexuality • effect on attitudes and spirituality
Disease-centred management	**Patient-centred management**
• rest	• psychological support
• drugs	• appropriate reassurance
• intervention	• patient education
• surgery	• empowering self-responsibility
• other invasive techniques	• anticipatory guidance/special hazards

- prevention
- health promotion
- lifestyle recommendations/modifications
  - diet/nutrition
  - exercise
  - alcohol
  - smoking
  - stress management
- family and social supports
- self-help groups alternative options
- consultation and referral
- follow-up

DIMENSIONS TO WHOLE PERSON MANAGEMENT

In the diagnostic model presented in Chapter 18 the fifth and final question is: ‘Is this patient trying to tell me something else?’ and this self-posed question should be scrutinised and answered in most instances. This presupposes being tuned in to the patient, watching for cues and listening.

An efficient medical record system also helps the process, since following a set routine generally ensures that important facets of the patient's psychosocial history are not omitted.

The answer to the above question takes into account the patient's:

- feelings
- fears or concerns
- expectations of the doctor
- future aspirations

Such an approach may determine whether there is a ‘hidden agenda’ in the presentation and whether various stressors, including interpersonal conflicts, are significant factors in the illness.

ILLUSTRATIVE CASE HISTORY

The patient. JR, aged 47, bus driver.

Disease-centred problem. Persistent, moderateto-severe low back pain for 5 months, after sudden onset of left sciatica (now settled) and low back pain following lifting a suitcase at work.

JR (a real patient) is representative of many patients encountered in general practice with the common problem of chronic back pain. His case serves to illustrate the importance of managing the whole person.

After taking a detailed history according to the pain analysis guidelines (p. 97), a physical examination focusing on his lumbosacral spine was performed. He had localised tenderness over the L4 - 5 interspinous space with restriction (stiff ness and discomfort) of forward flexion and extension. There was no neurological abnormality. His previous practitioner had recommended spinal surgery and prescribed piroxicam 20 mg (o) daily and OTC analgesics. He smoked 20 cigarettes a day and averaged four standard drinks of alcohol a day.

Investigations

- Plain X-ray (past) - slight narrowing L4 - 5 disc, minimal degenerative changes in facet joints.

- CT scan - mild spondylosis (facet joints) and L4 - 5 disc degeneration: nerve roots unaffected.

Disease-centred diagnosis

- L4-5 disc degeneration

Disease-centred management

- Back brace
- Piroxicam 20 mg daily
- Codeine and paracetamol compound tablets prn

Patient-centred problems

A detailed history about the effect of the back disorder on the patient revealed the following:

- humiliation about being on workers' compensation
- boredom and frustration leading to irritable behaviour (‘not like me’)
- anxious about his future, particularly occupation
- fears of being a cripple ‘in a wheelchair’
- conflicts with family members
- concern about reduction of sexual performance
- anger with lack of response from medical management
- anger with lack of interest in his problem
- concerns about something more serious (he admitted to a fear of cancer)
- feelings of depression: unable to cope, sleep disturbance
- concern about taking drugs, especially NSAIDs

Patient-centred physical examination

- General: tense and anxious man looking older than his years, senile arcus
- Cardiovascular examination: BP 165/100 mmHg
- Weight: 85 kg
- Height: 1.67 m
- BMI: 30.5

Problem list

- Chronic back pain
- Hypertension
- Obesity
- Anxiety
- Depression
- Concerns about:
    - serious disease
    - future employment
    - finances
    - sexual performance
    - drug taking
- Family disruption
- Problem drinking
- Nicotine excess
- Type 2 diabetes risk

- Ischaemic heart disease risk

Approaches to management

Psychological support

Factors that make patients feel supported and generate their confidence in the doctor include:

- showing an interest in all aspects
- examining the problematic area
- performing a more thorough examination
- picking up ‘cues’ and drawing attention to them

It is noteworthy just how impressed some patients are when they are physically examined. It is appreciated particularly by those who say: ‘The other doctor did not get me to take off my clothes nor examine me’.

Support includes acting as the patient's advocate during the complex process of coping with the red tape of workers' compensation formalities, examinations by other doctors and any medicolegal issues.

Appropriate reassurance

Reassurance should be given from an appropriate knowledge base. It should be emphasised to the patient that on the law of averages the chronic back pain should eventually resolve and the patient can look forward to resuming a relatively normal life.

The patient should also be reassured that he does not have malignancy or a physically crippling disorder.

Explanation, basic counselling and patient education

One of the most important features of management is careful and appropriate explanation, including the reasons for the pain and why it is self-perpetuating. A useful approach is that given on page 40, whereby the explanation is that the main triggering factor (the prolapsed disc) has basically resolved but the central nervous system is still registering the pain, analogous to a ‘phantom limb’. The development of anxiety and depression has caused continuing overactivity of the pain pathways, which need to be ‘switched off ’ by the various treatments that will be given.

Explore various concerns, including the effect of his illness on his family relationships and sexual activity, and provide counselling as necessary. The patient should then be handed suitable patient education material as presented in Figure 11.3 (at p. 93) and encouraged to undertake an exercise program (Figure 11.2 at p. 92). Education about the problems of anxiety, depression and lifestyle factors should also be addressed.

Empowering self-responsibility

Emphasise to the patient that he must take the responsibility for his own health and rehabilitation. This involves an active program of exercises and swimming, taking preventive measures and following recommendations about health promotion. It also involves dedication to restoration of ideal weight since his obesity (BMI >30) would be contributing to his back pain.

Medication

Explain tactfully that since there is no clinical evidence of inflammation, the NSAIDs prescribed by another practitioner can be ceased.

Because he is depressed with an anxiety component and insomnia, an antidepressant can be prescribed to be taken at night (see explanation for the patient on page 186 - 7). He can be assured that not only will this help his psychological state but it should help ‘shut the pain pathway gate’ and reverse the overactivity of the nervous system.

Patient education, including patient education handouts about possible side-effects and how to cope with them, would be appropriate.

Prevention and health promotion

The patient should be informed in a supportive way that he has the following risk factors:

- family history of cardiovascular disease
- family history of diabetes
- hypertension (related to obesity, NSAIDs, alcohol excess and smoking)
- smoking
- obesity
- possible hypercholesterolaemia (senile arcus noted)
- stress and anxiety

It would be important to stress (despite the fact that he might not be receptive during a depressed state) that for the sake of his future health it is time to change his lifestyle to try to reverse the risk factor process.

Advice on the prevention of recurrence of his back problem is important, with advice on daily activities, lifting, bending and so on. The use of a brace is not generally recommended as dependency on the supportive device tends to lead to aggravation of the chronicity of the back.

Lifestyle recommendation

In JR's case, and indeed for many patients, he should receive advice, education, guidelines and support in the basic lifestyle factors, namely:

- diet control:
    - weight reduction
    - low fat/cholesterol diet
    - high fibre
    - alcohol restriction
- smoking
- exercise and suitable physical activities
- interesting hobbies
- relaxation techniques

Family support and counselling

It would be ideal to include Mrs JR in the next consultation or, better still, arrange a home visit to discuss the role the family can take in his rehabilitation. Explanations and seeking family support would be very helpful.

Other positive suggestions would be to encourage Mrs JR to perform massage (as instructed) on the aching back and co-operate with his diet and other lifestyle issues, and to assess the state of the bed and the chairs as a possible aggravation factor for the back pain.

Anticipatory guidance and special hazards

Possible problems for JR include aggravation of his back pain by inappropriate physical and emotional stressors, marital break-up, loss of job and even suicide. These issues should be diplomatically discussed with the patient in a positive way. It is to be hoped that the interest, encouragement and support he receives can help avert these serious problems.

Recommended treatment for JR

- Attention to lifestyle factors
- Antidepressant medication
- Referral for therapy:
    - hydrotherapy
    - mobilisation

    - active exercises
- Therapeutic massage

Alternative options

There are many alternative therapies that can be used to help alleviate his painful condition. It is important for GPs to keep abreast of these therapies and their relative success rate based on scientific evidence. (Where known, the relative success rate, based on research by the author, is indicated.2) If the response is slow, one or more of these treatment modalities can be selected and the patient referred to a reputable therapist experienced in the therapy.

- Spinal mobilisation and manipulation (64%)
- Manipulation under general anaesthesia (34%)
- Transcutaneous electrical nerve stimulation (TENS)
- Acupuncture
- Biofeedback
- Meditation
- Facet joint injections (20%)
- Nerve blockage (48% after 6 months)
- Epidural injections
- Neurofasciotomy (42%)
- Radiofrequency denervation (65%)
- Electrotherapy
    - for superficial heating:
        - radiant heat lamps
        - infra-red lamps
    - for deep heating:
        - short-wave diathermy
        - ultrasound

Consultation and referral

One of the skills of a responsible GP is to sum up his or her own limitations with a particular problem and consult with and/or refer to a colleague if it is in the patient's best interests. Effective communication and advice is only a telephone call away. Patients with a chronic illness should be referred to an expert in the discipline to satisfy patient, family and GP that everything possible is being attempted to help the patient. Patients and their families often judge us by the quality of the referral process and its outcome so considerable thought has to be given to the most outstanding consultant for that particular problem at a given time. The GP should remain in charge of the team and direct management.

JR could be referred to medical consultants such as an orthopaedic surgeon, an orthopaedic physician or a rheumatologist. It would be inappropriate to refer him to a psychiatrist in the early stage of his pain but if a psychological problem intervened and complicated management (as in this case) referral should be strongly considered. As a rule depression and anxiety can be managed well by a caring family doctor.

Probably the best option for a patient like JR is referral to a pain clinic that is staffed by a multidisciplinary team of specialists including psychiatrists, social workers, physiotherapists, occupational therapists and other medical specialists. They look at the total problem and the whole person. As a patient similar to JR said after attending a pain clinic: ‘For the first time all my problems - physical, psychological, drug dependency and other disturbed fragments of my life - have been dealt with by the whole team’.

Follow-up

Careful instructions to ensure follow-up visits by the patient is important as there is a tendency for some dissatisfied and emotional patients to seek attention elsewhere and thus get on the chronic disorder ‘merry-go-round’.

Outcome

JR gradually improved, presumably because depression had complicated the problem. The aphorism 'acute pain ° acute anxiety and chronic pain ° depression' applied to JR and relief of his depression, combined with supportive measures, such as physiotherapy, hydrotherapy, physical therapy and strong psychological support from his family and GP, provided the 'holistic' approach to healing.

THE HOLISTIC APPROACH TO MANAGEMENT[3]

In a healing profession obsessed with interventionism, invasive technology and drug management, the general practitioner has an obligation to his or her patients to use natural healing methods wherever possible and be very discerning and conservative with investigatory medicine.

Patients appreciate natural remedies and taking responsibility for their own management wherever possible and appropriate. Examples include relative rest, exercise, swimming, stress management, meditation, spiritual awareness, antioxidant therapy (e.g. vitamin C, vitamin E, selenium), weight control, optimal healthy nutrition, avoidance of toxins (e.g. illicit drugs, nicotine, caffeine and alcohol) and sexual fulfilment.

Underlying a successful outcome is motivation, and the healing factor of the physician in being the motivator, teacher and facilitator should never be underestimated.

REFERENCES

1. Carson N. The diagnostic process. In: *Department of Community Medicine Final Year Handbook*. Melbourne: Monash University, 1990: 9 - 25.
2. Murtagh J. *Back pain in general practice*. Melbourne: Monash University, 1988: (MD thesis). 160 - 5.
3. Hetzel R. *The New Physician: Tapping the Potential for True Health*. Melbourne: Houghton Mifflin, 1991.

CHAPTER 14. TRAVEL AND TROPICAL MEDICINE

If you can't peel it, boil it or cook it - don't eat it.

ANON

EMPORIATICS - THE SCIENCE OF TRAVEL MEDICINE

With over 400 million international trips being taken annually the health problems faced by travellers are considerable and variable depending on the countries visited and the lifestyle adopted by the traveller.[1] There is evidence that many travellers are receiving inaccurate predeparture travel advice.[2] Travellers to North America, Europe and Australasia are usually at no greater risk of getting an infectious disease than they would be at home, but those visiting the less developed tropical and subtropical countries of Africa, Central and South America and South-East Asia are at significant risk of contracting infectious diseases.

Problems range in complexity from the most frequent and usually benign problems, such as traveller's diarrhoea, to more exotic and potentially fatal infections such as malaria, Japanese encephalitis and HIV. It must also be remembered that in some countries with volatile political changes there is the possibility of injury, incarceration or being left stranded. Travel means transport and thus the potential for accidents and crippled body and bank balance. Insurance for such contingencies is as important as preventive health measures.

KEY FACTS AND CHECKPOINTS

- The main diseases facing the international traveller are traveller's diarrhoea (relatively mild) and malaria, especially the potentially lethal chloroquine-resistant falciparum malaria (CRFM).
- Most cases (up to 75%) of traveller's diarrhoea are caused by enterotoxigenic *Escherichia coli*.
- Enteroinvasive *E. coli* (a different serotype) produces a dysentery-like illness similar to Shigella.
- Traveller's diarrhoea is contracted mainly from contaminated water and ice used for beverages, washing food or utensils or cleaning teeth.
- Poliomyelitis is endemic in at least 20 countries and thus immunisation for polio is still important.
- One bite from an infected mosquito during a single overnight stop in a malaria area can result in a possible lethal infection.
- Infections transmitted by mosquitoes include malaria, yellow fever, Rift Valley fever, Japanese encephalitis, chikungunya and dengue fever. Avoiding their bites is excellent prevention.
- Every year over 500 Australians catch malaria while travelling overseas.
- Malaria is a dusk-till-dawn risk only, but bites from daytime mosquitoes can cause dengue.
- CRFM is steadily increasing, as is resistance to newer antimalarials.
- It is important for general practitioners to consult a travel medicine database to obtain specific information about 'at risk' countries.
- Avoid tattooing, ear-piercing, acupuncture or any skin puncturing while overseas.
- The commonest cause of death in travellers overseas is trauma, particularly traffic accidents and homicides.

PRINCIPLES OF PRE-TRAVEL HEALTH CARE

- Advise the patient to plan early - at least 8 weeks beforehand.
- Advise a dental check.
- Allow adequate time for consultation (e.g. 30 - 45 minutes).
- Individualise advice.
- Provide current information.
- Provide written as well as verbal advice.
- Provide a letter concerning existing medical illness and treatment.
- Encourage personal responsibility.

GASTROINTESTINAL INFECTIONS

The commonest problem facing travellers is traveller's diarrhoea but other important diseases caused by poor sanitation include hepatitis A, and worm infestations such as hookworm and schistosomiasis.

Contamination of food and water is a major problem, especially in third world countries.

Reputable soft drinks, such as Coca-Cola, should be recommended for drinking. Indian-style tea, in which the milk is boiled with tea, is usually safe, but tea with added milk is not. The food handlers can be infected and the water used to wash food may be contaminated.

Traveller's diarrhoea

Traveller's diarrhoea is a special problem in Mexico, Nepal, India, Pakistan, Latin America, the Middle East and central Africa and its many colourful l abels include 'Bali Belly', 'Gippy Tummy', 'Rangoon Runs', 'Tokyo Trots' and 'Montezuma's Revenge'. It occurs about 6 - 12 hours after taking infected food or water.

The illness is usually mild and lasts only 2 or 3 days. It is unusual for it to last longer than 5 days. Symptoms include abdominal cramps, frequent diarrhoea with loose watery bowel motions and possible vomiting. Very severe diarrhoea, especially if associated with the passing of blood or mucus, may be a feature of a more serious bowel infection such as amoebiasis.

Most traveller's diarrhoea is caused by an enterotoxigenic *E. coli*. Travellers are infected because they are exposed to slightly different types or strains of *E. coli* from the ones they are used to at home.[3]

The possible causes of diarrhoeal illness are listed in Table 14.1.

Table 14.1 Causes of diarrhoea in travellers

	Causative organism	Type of illness
Bacteria	*Escherichia coli*	Traveller's diarrhoea
	Shigella species	Dysentery
	Salmonella species	Typhoid fever, food poisoning
	Campylobacter jejuni	Traveller's diarrhoea, dysentery
	Vibrio cholerae	Cholera
	Yersinia enterocolitica	Traveller's diarrhoea
	Aeromonas hydrophilia	Traveller's diarrhoea
	Staphylococcus aureus (toxin)	Food poisoning
	Clostridium perfringens	Food poisoning
	Bacillus cereus	Food poisoning
Viruses	Rotavirus	Children's diarrhoea
	Norwalk virus	Traveller's diarrhoea
Protozoa parasites	Cryptosporidium	
	Entamoeba histolytica	Amoebiasis
	Giardia lamblia	Giardiasis
	Strongyloides stercoralis	Strongyloidiasis
	Schistosomiasis	
Chemicals	Capsicum (chilli)	

Treatment

Refer to Figure 14.1.[3,4]


• Nausea
• Watery diarrhoea
• Abdominal cramps
Fever?
yes
no
Bloody stools
no
Rx • fluids +
* Imodium or Lomotil
Still unwell at 48 hours
yes
no
Continue trip
yes
Rx • fluids
• glucose electrolyte solution
+
• antibiotics: norfloxacin or azithromycin
Still unwell at 48 hours?
no
yes
Seek medical advice
Specimens:
? amoebiasis
? giardiasis
? *Salmonella*
? drug-resistant organism
? other cause

Figure 14.1 Algorithm for adult travellers with acute diarrhoea. After Locke [3]

The key factor in treatment is rehydration.

Mild diarrhoea

- Maintain fluid intake - cordial or diluted soft drink.
- Antimotility agents (judicious use: if no blood in stools) - avoid in children. loperamide (Imodium) 2 caps statim then 1 after each unformed stool (max. 8 caps/day) *or*
- diphenoxylate with atropine (Lomotil) 2 tablets statim then 1 - 2 (o) 8 hourly. Imodium is the preferred agent.

Moderate diarrhoea[4]

- Attend to hydration.
- Patient can self-administer antibiotic - e.g. norfloxacin 400 mg bd for 3 days or 800 mg (o) statim; in children, (20 mg/kg) up to 800 mg (o) as single dose.
- Avoid Lomotil or Imodium.

Severe diarrhoea (patient toxic and febrile)

- ? Admit to hospital.
- Attend to hydration - use an oral hydrate solution (e.g. Gastrolyte or WHO formulation).
- Avoid Lomotil and Imodium.
- Antibiotic: norfloxacin, ciprofloxacin or azithromycin.

Note: There is increasing resistance to doxycycline and cotrimoxazole, especially in South-East Asia.

Persistent diarrhoea

Any travellers with persistent diarrhoea after visiting less developed countries, especially India and China, may have a protozoal infection such as amoebiasis or giardiasis. If the patient has a fever and mucus or blood in the stools, suspect amoebiasis. Giardiasis is characterised by abdominal cramps, flatulence, and bubbly, foul-smelling diarrhoea persisting beyond 2 to 4 days. Take three specimens of faeces for analysis. In some cases serology may be helpful, e.g. amoebiasis.

Treatment

- Giardiasis: tinidazole or metronidazole
- Amoebiasis: metronidazole or tinidazole

Patient can self-administer these drugs and carry them if visiting areas at risk, but they can have a severe adverse reaction with alcohol.

Preventive advice

The following advice will help prevent diseases caused by contaminated food and water. These ‘rules’ need only be followed in areas of risk such as Africa, South America, India and other parts of Asia.

- Purify all water by boiling for 10 minutes. Adding purifying tablets is not so reliable, but if the water cannot be boiled some protection is provided by adding Puratabs (chlorine) or iodine (2% tincture of iodine), which is more effective than chlorine - use 4 drops iodine to 1 litre of water and let it stand for 30 minutes.
- Do not use ice. Drink only boiled water (supplied in some hotels) or well-known bottled beverages (mineral water, 7-Up, Coca-Cola, beer).
- Avoid fresh salads or raw vegetables (including watercress). Salads or uncooked vegetables are often washed in contaminated water. Bananas and fruit with skins are safe once you have peeled and thrown away the skin but care should be taken with fruit that may possibly be injected with water.
- Be wary of dairy products such as milk, cream, ice-cream and cheese.
- Avoid eating raw shellfish and cold cooked meats.
- Avoid food, including citrus fruits, from street vendors.
- Drink hot liquids wherever possible.
- Use disposable moist towels for hand washing.

The golden rule is: *If you can't peel it, boil it or cook it - don't eat it.*

MALARIA

General aspects:

- Travellers to all tropical countries are at some risk.
- Malaria is endemic in 102 countries;[5] 2.3 billion people are at risk.
- The risk is very low in the major cities of Central and Southern America and South-East Asia but can be high in some African cities.
- Malaria is either benign (vivax, ovale) or malignant (falciparum).
- Resistance to many drugs is increasing:
  - The lethal **Plasmodium falciparum** is developing resistance to chloroquine and the antifolate malarials (Fansidar and Maloprim).[5]
  - Resistance is now reported to mefloquine.
  - CRFM is common in South-East Asia, Papua New Guinea, northern South America and parts of Africa.
- Chloroquine is still effective against *P. ovale* and *P. vivax* (the most common forms).
- The long-awaited vaccine will make all the complex drug management much simpler. However, it still appears to be many years away despite considerable research.
- Patients who have had splenectomies are at grave risk from P. falciparum malaria.
- People die from malaria because of delayed diagnosis, delayed therapy, inappropriate therapy and parasite - host factors.
- Practitioners should follow updated recommended guidelines (e.g. NH&MRC, WHO).

DxT: fever + chills + headache ° malaria

Malaria risk assessment

The risk of catching malaria is increased by:

- being in a malaria area, especially during and after the wet season
- a prolonged stay in a malaria area, especially rural areas, small towns and city fringes
- sleeping in unscreened rooms without mosquito nets over the bed
- wearing dark clothing with short-sleeved shirts and shorts
- taking inappropriate drug prophylaxis

Malarial prevention

Travellers should be advised that malaria may be prevented by following two simple rules:

- avoid mosquito bites; and
- take antimalarial medicines regularly

In order to avoid mosquito bites, travellers are advised to:

- keep away from rural areas after dusk
- sleep in air-conditioned or properly screened rooms
- use insecticide sprays to kill any mosquitoes in the room or use mosquito coils at night
- smear an insect repellent on exposed parts of the body; an effective repellent is diethyl-mtoluamide (Muskol, Repellem, Rid)
- use mosquito nets (tuck under mattress; check for tears)
- impregnate nets with permethrin (Ambush) or deltamethrin
- wear sufficient light-coloured clothing, long sleeves and long trousers, to protect whole body and arms and legs when in the open after sunset
- avoid using perfumes, cologne and after-shave lotion (also attracts insects)

Important considerations in malaria prophylaxis

1. Minimise exposure to mosquitoes and avoid bites.
2. Know areas of risk:
    - tropical South America (southern Mexico to northern half South America)
    - tropical Africa (sub-Sahara to northern South Africa)
    - Nile region, including Egypt
    - Southern Asia, especially tropical areas
3. Know areas of widespread chloroquine resistance:
    - Asia, tropical South America (rare north of Panama Canal), sub-Sahara, East Africa
4. Consider several factors:
    - intensity of transmission
    - season and length of stay
    - itinerary
        - urban - hotel
        - urban - non-hotel
        - rural - housing
        - rural - backpacking
    - resistance patterns
    - host factors
        - age
        - pregnancy
        - associated illness
        - compliance
5. Know the antimalarial drugs (Table 14.2).

Table 14.2 Common drugs used for malarial prophylaxis[4]

	Adult dosage	Children's dose	Comments
Chloroquine (Chlorquin)	300 mg base (2 tabs) same day each week 1 week before, during, 4 weeks after exposure	5 mg base/kg up to maximum adult dose	Only antimalarial approved for pregnancy Aggravates psoriasis Beware of retinopathy
Doxycycline	100 mg each day, 2 days before, during, 4 weeks after	> 8 years only 2 mg/kg/day up to 100 mg	Photosensitivity reactions
Mefloquine (Lariam)	250 mg (1 tab) same day each week, 1 week before, during, 4 weeks after	Not recommended < 45 kg; > 45 kg as for adults	Side-effects: dizziness, 'fuzzy' head, blurred vision Beware of beta-blockers
Proguanil (Paludrine)	200 mg (2 tabs) same day each week 1 day before, during, 4 weeks after	> 1 year: ¼ tablet 1 - 4 years: ½ tablet 5 - 8 years: 1 tablet 9 - 14 years: 1½ tablett > 14 years: adult	Safe in lactation and pregnancy (give folic acid) Side-effects: GIT disturbances, headache, dizziness, rash
Atovaquone+ proguanil Malarone)	250 mg/100 mg (1 tab) with food same day each week 2 days before, during, 7 days later	Junior tablets 62.5 mg/25 mg 11 - 20 kg: 1 tablet/day 21 - 30 kg: 2 tablet/day 31 - 40 kg: 3 tablet/day > 40 kg: 1 adult tab/day	Avoid in pregnant women or women breastfeeding infants < 11 kg Avoid in severe kidney impairment Side-effects: GIT upset, headache, dizziness, myalgia - others

6. Balance risk benefit of drug prophylaxis: drug side-effects versus risk of CRFM.
7. Visiting areas of CRFM does not automatically require the use of potentially harmful drugs.[1]
8. Those at special risk are pregnant women, young children and the immunocompromised. Advise against travel.
9. No drugs give complete protection.

Drug prophylaxis

Refer to WHO guidelines. < www.who.int/ith/en >.

Guidelines

- Accommodation in large air-conditioned hotels in most cities of South-East Asia (dusk - dawn) for < 2 weeks: no prophylaxis required.
- For low-risk travel (urban: dusk - dawn) in areas of high resistance for < 2 weeks: chloroquine adequate; use a treatment course of mefloquine if necessary (Table 14.3 on p. 118).

Table 14.3 Drugs used for chloroquine-resistant malaria (presumptive breakthrough where professional medical care unavailable)[4]

	Adult dose	Children's dose
Artemether/lumefantrine (Riamet)	4 tablets at 0, 8, 24, 36, 48, 60 hours	Only if > 12 years, > 35 kg
Mefloquine (Lariam)	750 mg (3 tablets) statim, then 500 mg after 6 - 8 hours or 1000 mg statim	Only recommended if weight > 45 kg (as for adults)
Atovaquone/proguanil (Malarone)	4 tablets daily for 3 days	11 - 20 kg: 1 tablet 21 - 30 kg: 2 tablets 31 - 40 kg: 3 tablets

*Unsuitable for Thailand.[4]

- For short-and long-term travel to rural areas of high resistance (e.g. South-East Asia including Thailand, Kenya, Tanzania, Ecuador, Venezuela, Brazil): doxycycline daily alone or mefloquine (once a week).

Summary of recommendations[4]

1. CSFM area: chloroquine
2. CRFM area: mefloquine 250 mg/week *or*
   doxycycline 100 mg/day or atovaquone + proguanil
3. Multidrug-resistant area: doxycycline 100 mg/day chloroquine for stays > 8 weeks
   +
   doxycycline 50 - 100 mg/day for stays > 8 weeks

Standby treatment: mefloquine or artemether + lumefantrine or atovaquone + proguanil

Note: CSFM ° chloroquine-sensitive falciparum malaria CRFM ° chloroquine-resistant falciparum malaria

SPECIFIC INFECTIOUS DISEASES AND IMMUNISATION

Protection from many types of infection is available through immunisation. All travellers should be immunised against tetanus, polio and diphtheria and measles. Protection against tetanus requires an initial course of three injections followed by a booster every 10 years.

Vaccinations are required for special circumstances. Yellow fever vaccination is a legal requirement for any travellers returning from a yellow fever endemic area. Cholera is not usually required.

Some travellers may be exposed to tuberculosis, hepatitis, plague, rabies, typhoid, typhus, and meningococcal infection. Immunisation against these is available and recommended for those at risk. Smallpox has now been eradicated from the world and therefore smallpox vaccination is no longer required for any traveller.

Japanese Bencephalitis presents as a special problem to the traveller.

Table 14.4 outlines a summary of recommen dations to consider.[7]

Table 14.4 Summary of preventive measures and vaccinations

All travellers, all destinations
Tetanus toxoid and diphtheria booster
if > 10 years since last dose
if > 5 years for third world travel
give CDT < 8 years ADT > 8 years
All travellers to developing countries free of malaria

Tetanus toxoid booster

Polio immunisation if > 10 years

Measles immunisation (consider MMR)

Influenza and pneumococcus vaccination (for those at risk)

Yellow fever (if compulsory)

Preventive measures against:

- gastrointestinal infections
- sexually transmitted infections
- mosquito bites

Travellers to developing and other countries at high risk of infection

As above plus:

- malaria prophylaxis
- hepatitis A-vaccine or immunoglobulin
- hepatitis B
- typhoid
- tuberculosis (BCG if Mantoux - ve)

Other vaccinations - consider:

- meningococcus (required in some countries)
- Japanese Bencephalitis
- rabies
- typhus
- plague
- anthrax
- cholera

COMPULSORY IMMUNISATIONS

The two vaccinations that may be required before visiting ‘at risk’ areas are meningococcus and yellow fever.

Yellow fever

Yellow fever is a serious viral infection spread by mosquitoes and, like malaria, is a tropical disease. Yellow fever vaccination is essential for travel to or through equatorial Africa and northern parts of South America, and for reentry to Australia from those countries.

DxT: fever + bradycardia + jaundice ° yellow fever

One injection only is required and the immunisation is valid for 10 years. Children aged less than 9 months should not be given this vaccine. It should not be given within 3 weeks of cholera vaccine.

Note: It is important to check specific country requirements in the WHO book on vaccination requirements.[8]

According to WHO a certificate against yellow fever is the only certificate that should be required for international travel. The requirements of some countries are in excess of International Health Regulations. However, vaccination against yellow fever is strongly recommended to all travellers who intend to visit places other than the major cities in the countries where the disease occurs in humans.

Meningococcal infection

Meningitis due to this organism is a contagious lethal disease. It is common in Nepal, Mongolia, Vietnam and parts of Africa and

Asia, especially in the dry season. Travellers trekking through the Kathmandu valley of Nepal and those attending the Haj pilgrimage to Saudi Arabia are at special risk and should have the vaccine. However, some countries require immunisation for entry.

VOLUNTARY IMMUNISATION

Precautions against the following diseases are recommended for those travellers who may be at special risk.

Hepatitis A, B, E

Hepatitis A is a common problem in rural areas of developing countries. There is a declining level of antibodies to hepatitis A in developed countries and adults are at special risk so one or two doses of hepatitis A vaccine should be given. If there is insufficient time a single injection of human immunoglobulin (IG) can give protection for 3 to 6 months. It is safe for all age groups but children under 8 years should not need it. A blood test for hepatitis A antibodies can be carried out to determine a person's immunity.

Prevention of hepatitis A

The rules of avoiding contaminated food and water apply (as for traveller's diarrhoea). Hepatitis A vaccine is given as a course of two injections.

Hepatitis B is endemic in South-East Asia, South America and other developing countries. Vaccination is recommended, especially for people working in such countries, particularly those in the health care area or those who may expect to have sexual or drug contact. If patients have a ‘negative’ HBV core IgG titre, then vaccination would be worthwhile (three doses: 0, 1 and 6 months). Hepatitis E has a high mortality rate in pregnant women.

The usual approach for non-immunised people is to give the combined hepatitis A and B vaccine (Twinrix) as a course of three injections.

Typhoid

Typhoid immunisation is not required for entry into any country but is recommended for travel to third world countries where the standards of sanitation are low. It should be considered for travellers to smaller cities, and village and rural areas in Africa, Asia, Central and South America and Southern Europe.

The parenteral (subcutaneous) vaccine can be used but the new single dose typhi Vi vaccine or the oral vaccine, which have fewer side-effects, are generally preferred. The oral vaccine, which is given as a series of three or four capsules, appears to afford protect ion for about 5 years but is contraindicated in the immunocompromised.

Cholera

Cholera vaccination is not officially recommended by the WHO because it has only limited effectiveness. It is advisable for health care workers or others at risk entering an endemic area. Cholera is given in two injections 7 to 28 days apart or usually as an oral vaccine (Dukoral) over 1 week prior to exposure. It is not recommended in children under 5 years or pregnant women.

Japanese Bencephalitis

This mosquito-borne flavivirus infection presents a real dilemma to the traveller and doctor because it is a very severe infection (mortality rate 20 - 40%) with high infectivity and high prevalence in endemic countries. The vaccine is prone to give allergic react ions and anaphylaxis. It may be obtained only in very restricted circumstances but can be obtained more readily abroad.

The disease is prevalent during the wet season in the region bound in the west by Nepal and Siberian Russia and in the east by Japan and Singapore, especially in Nepal, Burma, Korea, Vietnam, Thailand, China, eastern Russia and the lowlands of India. Rice paddies and pig farms are areas of risk. The usual preventive measures against mosquito bites are important.

DxT: febrile illness + vomiting + stupor ° Japanese Bencephalitis

Rabies

Rabies vaccination is recommended for some international aid workers or travellers going to rabies-prone areas for long periods. The vaccination can be effective after the bite of a rabid animal, so routine vaccination is not recommended for the traveller. Affected animals include dogs, cats, monkeys and feral (wild) animals. A traveller who sustains a bite or scratch or even is licked by an animal in countries at risk should wash the site immediately with soap or a detergent, and then seek medical help. The prebite vaccination does not remove the need for postexposure vaccination.

DxT: painful bite + paraesthesia + hydrophobia pain with drinking) ° rabies

Typhus

Typhus is transmitted to humans by bites from lice, fleas or ticks. Immunisation against typhus is desirable for doctors, nurses, and agricultural and technical advisers whose work takes them to remote areas of Bolivia, Burundi, Ethiopia, Mexico, Rwanda and mountainous regions of Asia. A booster injection should be given at 12 months if the traveller is still in an area of risk.

Plague

Plague is still prevalent in rodents in several countries, such as Vietnam, Brazil, Peru, Ecuador, Kenya and Malagasy Republic. Although not compulsory, vaccination is recommended for those engaged in field operations in plague areas and rural health workers who may be exposed to infected patients. Two doses are given to adults (three to children < 12 years) and a booster every 6 months.

SPECIAL PROBLEMS

Prevention of sexually transmitted infections

Casual sexual contacts place the traveller at risk of contracting a serious, perhaps fatal, STI. The common STIs especially prevalent in South-East Asia and Africa are non-specific urethritis (NSU), gonorrhoea (especially penicillin-resistant strains), hepatitis B and syphilis. HIV infection is a rapidly increasing problem, with heterosexual transmission common in Africa and in South-East Asia. Unusual STIs such as lymphogranuloma venereum, chancroid and donovanosis are encountered more commonly in tropical developing countries. A practical rule is to assume that all ‘at risk’ travellers are both ignorant and irresponsible and advise accordingly.

Rule: never permit blood, semen or vaginal fluids from a sexual partner to enter your body, unless you are absolutely certain they are not STD carriers.

Prevention

Abstinence or take your partner (condoms and diaphragms do not give absolute protection).

Drugs

Possession of and trafficking in drugs is very hazardous and many people are held in foreign prisons for various drug offences. The penalty for carrying drugs can be death!

Countries that currently may enforce the death penalty are Burma, Indonesia, Malaysia, Singapore, Thailand and Turkey. Travellers should be warned about taking cannabis while in a foreign country, as it can cause profound personality changes in the user.

Drug addicts should under no circumstances travel. Young travellers should be wary about accepting lift s or hitchhiking in countries ‘at risk’.

PROBLEMS IN THE RETURNED TRAVELLER

- Most will present within 2 weeks except HIV infection.
- Common infections encountered are giardiasis, amoebiasis, hepatitis A, B or E, gonorrhoea or *Chlamydia trachomatis*, malaria and helminthic infestations.
- An important non-infected problem requiring vigilance is deep venous thrombosis (DVT) and thromboembolism.
- The asymptomatic traveller may present for advice about exposure (without illness) or about an illness acquired such as rabies, malaria, schistosomiasis and STIs.

Exposure to STIs

If a patient has had unprotected intercourse and is at definite risk of acquiring an STI, such as penicillin - resistant gonorrhoea or NSU, the following may be appropriate:[1]

- ceftriaxone 250 mg IM (as a single dose)
- doxycycline 100 mg (o) for 10 days or azithromycin 1 g (o) statim

Gastrointestinal symptoms

Mild diarrhoea

- Stool microscopy and culture
- Look for and treat associated helminthic infestation (e.g. roundworms, hookworms)

Moderate or prolonged (> 3 weeks) diarrhoea

Usually due to *Giardia lamblia*, *Entamoeba histolytica*, *Campylobacter jejuni*, *Salmonella*, *Yersinia enterocolitica* or *Cryptosporidium*.

- Stool examination (three fresh specimens):
  - microscopy
  - wet preparation
  - culture
- Treat pathogen (see guidelines under diarrhoea, pp. 114 - 116)

Non-pathogens such as *E. coli* and *Endolimax nana* are often reported but do not treat specifically.

Note: Consider exotic causes such as schistosomiasis, strongyloidiasis and ciguatera in unusual chronic post-travel ‘gastroenteritis’.

Persistent abdominal discomfort

This common syndrome includes bloating, intestinal hurry and borborygmi, and often follows an episode of diarrhoea. Usually no pathogens are found on stool examination. However, giardiasis can be difficult to detect and an empirical course of tinidazole (2 g statim) is worthwhile. Any persistent problem then is a type of postinfective bowel dys-function or irritable bowel. Reassurance is important.

Fever

- Causes range from mild viral infections to potentially fatal cerebral malaria (Table 14.5) and meningococcal septicaemia.

Table 14.5 Fever and malaise in the returned traveller: diagnostic strategy model

Note: All fever in a returned traveller is malaria until proved otherwise!
Q. Probability diagnosis **A.** Viral respiratory illness (e.g. influenza) Hepatitis (may be subclinical) Gastroenteritis **Q. Serious disorders not to be missed** **A.** Malaria Typhoid Japanese Bencephalitis Meningococcal meningitis Melioidosis Amoebiasis (liver abscess) HIV seroconversion illness
Q. Pitfalls (often missed) **A.** Ascending cholangitis

Infective endocarditis

Cytomegalovirus

Epstein-Barr virus

Dengue fever

Lyme disease

Bronchopneumonia

Ross River fever

Rarities

Chikungunya

Legionnaire's disease

Schistosomiasis

African trypanosomiasis

Yellow fever

Rift Valley fever

Spotted fever

Lassa fever

Note: Three causes of a dry cough (in absence of chest signs) are malaria, typhoid, amoebic liver abscess.

Q. Seven masquerades checklist

A. Drugs (reaction to antimalarials)

Urinary infection

Investigations (if no obvious cause)

- Full blood examination
- Thick and thin blood films
- Blood culture
- Liver function tests
- Urine - micro and culture
- Stool - micro and culture
- ESR
- New malaria tests

- The common serious causes are malaria, typhoid, hepatitis (especially A and B), dengue fever and amoebiasis.
- Most deaths from malaria have occurred after at least 3 or 4 days of symptoms that may be mild. Death can occur within 24 hours. Factors responsible for death from malaria include delayed presentation, missed or delayed diagnosis (most cases), no chemoprophylaxis and old age.
- Refer immediately to a specialist unit if the patient is unwell.
- Be vigilant for meningitis and encephalitis.
- Be vigilant for amoebiasis - can present with a toxic megacolon, especially if antimotility drugs given.
- If well but febrile, *first-line screening tests*:
  - full blood examination and ESR
  - thick and thin films
  - liver function tests
  - urine for micro and culture
- Refer immediately if malaria is proven or if fever persists after a further 24 hours.

Malaria

See Figure 14.2.


Typical features
headache
malaise
fever/chills
prostration
sweating
myalgia
Other possible features
cerebral problems
• delirium
• convulsions
• coma
anaemia
jaundice
vomiting
T ↑↑
P ↑
BP N-↓
R N-↑
postural hypotension
cough
splenomegaly
hepatomegaly
abdominal pain
diarrhoea
arrhythmia
dark urine (blackwater fever)

Figure 14.2 Clinical features of malaria

- Incubation period: *P. falciparum* 7 - 14 days; others 12 - 40 days
- Most present within 2 months of return
- Can present up to 2 or more years
- Can masquerade as several other illnesses

Symptoms

- High fever, chills, rigor, sweating, headache
- Usually abrupt onset
- Can have atypical presentations (e.g. diarrhoea, abdominal pain, cough)

Other features

- Beware of modified infection.
- Must treat within 4 days.
- Typical relapsing patterns often absent.
- Thick smear allows detection of parasites (some laboratories are poorly skilled with thick films).

- Thin smear helps diagnose malaria type.
- If index of suspicion is high, repeat the smear (‘No evidence of malaria’ ° 3 negative daily thick films). Monocytosis and thrombocytopenia is a helpful diagnostic clue. Newer tests (e.g. PCR tests and ICT card tests) show promise. Cerebral malaria and blackwater fever are severe and dramatic. The Para check V test (a desktop test) is accurate and needs to be positive before prescribing artemether in some areas.

Treatment [4, 6]

- Admit to hospital with infectious disease expertise
- Supportive measures, including fluid replacement
- *P. vivax*, *P. ovale*, *P. malariae* [5] (Check G-6-PD activity first) chloroquine 4 tabs (o) statim, then 2 tabs in 6 hours, then 2 tabs on day 2 and day 3 + primaquine 15 mg (o) daily for 14 days
- *P. falciparum* [5] uncomplicated: artemether + lumefantrine (20 mg + 120 mg) 4 tablets with food at 0, 8, 24, 36, 48, 60 hours (i.e. 24 tablets) in 60 hours *or* quinine sulphate 600 mg (o) 8 hourly, 7 days + doxycycline 100 mg (o) 12 hourly, 7 days *or* clindamycin 300 mg (o) tds, 7 days *or* (alone as alternative to above) mefloquine *or* atovaquone + proguanil (Malarone) 4 tabs (o) daily for 3 days (if not used for prophylaxis)
  complicated (severe): quinine dihydrochloride 20 mg/kg up to 1.4 g IV (over 4 hours) *then* after 4-hour gap 7 mg/kg IV 8 hourly until improved then quinine (o) 7 days + Fansidar statim

Note: Check for hypoglycaemia. Beware if antimalarial use in previous 48 hours.

Typhoid fever

Incubation period 10 - 14 days.

Clinical presentation

DxT: ‘stepladder’ fever + abdominal pain + relative bradycardia ° typhoid (early)

- Insidious onset
- Headache prominent
- Dry cough
- Fever gradually increases in ‘stepladder’ manner over 4 days or so
- Abdominal pain and constipation (early)
- Diarrhoea (pea soup) and rash - rose spots (late)

Diagnosis

- On suspicion - blood culture
- Serology not very helpful

Treatment

- ciprofloxacin 500 mg (o) bd for 14 days

Dengue fever

Also known as ‘breakbone’ fever, it is wide-spread in the south-east Pacific and endemic in Queensland. A returned traveller with myalgia and fever <39°C is more likely to have dengue rather than malaria.

Features

- Mosquito-borne (*Aedes aegyptii*) viral infection
- Incubation period 5-6 days
- Abrupt onset fever, malaise, headache, nausea, pain behind eyes, severe backache, prostration
- Sore throat
- Severe aching of muscles and joints
- Fever subsides for about 2 days, then returns
- Maculopapular rubelliform rash on limbs → trunk
- Generalised erythema with ‘islands of sparing’

- ± Diarrhoea
- The rare haemorrhagic form is very severe; may present with shock which is usually fatal
- Later severe fatigue and depression (prone to suicide)

Note: A large-scale survey of dengue patients showed the symptoms as fever 100%, myalgia 79%, rash 74%, headache 68%, nausea 37%.

DxT: fever + severe aching + rash ° dengue fever

Diagnosis

- Dengue-specific IgM serology
- FBE: leukopaenia; thrombocytopenia in haemorrhagic form

Treatment

- Symptomatic with rest, fluids and analgesics

Prevention

- Avoid mosquito bites - no vaccine available

Chikungunya

This is an alpha-viral mosquito-borne infection with a similar clinical picture to dengue fever and can cause haemorrhagic fever. It is encountered in tropical South-East Asia, Indian Ocean Islands and parts of Africa. Diagnosis - positive serology.

Japanese Bencephalitis and meningococcal meningitis

Consider these serious infections in a patient presenting with headache, fever and malaise before neurological symptoms such as delirium, convulsions and coma develop.

Other tropical infections

Travellers to tropical or subtropical areas are at risk of more unusual infections. Most of these infections are contracted through contaminated food and water, insect bites and walking barefoot on contaminated soil. The risk of such infections is highest in rural areas of countries other than Europe, North America and Australasia.

African trypanosomiasis (sleeping sickness)

Clinical features

- Incubation period about 3 weeks
- Fever, headache and a skin chancre or nodule
- Lymphadenopathy, hepatosplenomegaly

Diagnosis

- Demonstrating trypomastigotes in peripheral blood smear or chancre aspirate

Treatment

- Suramin IV
- Infectious disease consultation essential

Prevention

- Avoid bites of the tsetse fly. If visiting areas of East, Central and West Africa, especially the ‘safari game parks’, travellers should use insect repellent and wear protective light-coloured clothing, including long sleeves and trousers.

Cutaneous leishmaniasis

This may be encountered in travellers and servicemen and servicewomen returning from the Middle East, especially the Persian

Gulf. The protozoa is transmitted by a sandfly and has an average incubation period of 9 weeks. The key clinical finding is an erythematous papuleFigure 14.3. Diagnosis is made by performing a punch biopsy and culturing tissue in a special medium. Treatment for extensive lesions is with high dosage ketoconazole for 1 month. Smaller lesions should be treated topically with 15% paromomycin and 12% methyl benzethonium chloride ointment applied bd for 10 days.[9] A special vaccine is available in some Middle Eastern countries (e.g. Israel).



Figure 14.3 Cutaneous leishmaniasis in a serviceman after returning from the Middle East.

Schistosomiasis (bilharzia)

The first clinical sign is a local skin reaction at the site of penetration of the parasite (it then invades liver, bowel and bladder). This site is known as ‘swimmer's itch’. Within a week or so there is a generalised allergic response, usually with fever, malaise, myalgia and urticaria. A gastroenteritis-like syndrome can occur (nausea, vomiting, diarrhoea) and respiratory symptoms, particularly cough. Clinical findings, such as in trypanosomiasis, include lymphadenopathy and hepatosplenomegaly.

The infestation is caused by parasite organisms (schistosomes) whose eggs are passed in human excreta, which contaminates watercourses (notably stagnant water) and irrigation channels in Egypt, other parts of Africa, South America, some parts of South-East Asia and China. Freshwater snails are the carriers (vectors).

Diagnosis

- Detecting eggs in the stools, the urine or in a rectal biopsy

Treatment

- Praziquantel

Prevention

- Travellers should be warned against drinking from, or swimming and wading in, dams, watercourses or irrigation channels, especially in Egypt and other parts of Africa.

Hookworm and strongyloidiasis

Hookworm and Strongyloides are parasites that are acquired by walking barefoot (or wearing thongs or sandals) on earth contaminated by faeces. The larvae penetrate the skin, travel through the lungs and settle in the small intestine. The first sign is local irritation or ‘creeping eruption’ at the point of entry, known as ‘ground itch’, which is often unnoticed. This subsides within 2 days or so and is followed 1 - 2 weeks later by respiratory symptoms, which may be associated with bronchitis and bronchopneumonia. Eventually a severe and chronic anaemia may develop. Hookworm infection is the commonest cause of iron deficiency anaemia in the world. Strongyloidiasis, which usually does not present with anaemia, is an important pathogen since it remains undetected for decades and presents as a severe infection, such as septicaemia, when the host becomes immunocompromised for any reason.

Diagnosis

- Detecting larvae or ova in the stool

Treatment (adults)

- Hookworm - mebendazole 100 mg bd for 3 days
- Strongyloidiasis - thiabendazole 1.5 g bd for 3 days (or more) or albendazole
- Adverse effects are common. Beware of these drugs in pregnancy and children.

Prevention

- Travellers should be warned to wear shoes and socks in endemic areas to prevent entry of the larvae into the skin of the feet.

Cutaneous larva migrans

Cutaneous larva migrans (creeping eruption) should be suspected in any pruritic, erythematous lesion with a serpiginous eruption on the skin, especially the hands and feet of a person from a subtropical or tropical area. It is caused by the larvae of dog or cat hookworms penetrating and migrating throughout human skin, the larva always being just ahead of the lesion it causes. The diagnosis is based on the classic clinical appearance and by eosinophilia. Biopsy is usually not indicated. The problem is usually self-limiting.

Diagnosis

- Clinical (characteristic appearance), eosinophilia (biopsy usually not indicated)

Treatment

- Albendazole or thiabendazole[4]
- Antihistamines for pruritus

Note: This is usually a self-limiting problem.

Prevention

- As for hookworm. Moist sandy soil contaminated with dog or cat faeces is a common source.

Cutaneous myiasis

Myiasis, which refers to the infestation of body tissues by the larvae (maggots) of flies, often presents as itchy 'boils'. Primary myiasis invariably occurs in travellers to tropical areas such as Africa and Central America, whereby the fly can introduce the larvae into the skin, or it can be due to secondary invasion of pre-existing wounds. Close inspection of lesions may reveal part or all of the larva. The simplest treatment is lateral pressure and tweezer extraction.

Trichuriasis (whipworm)

This common tropical parasite is sometimes seen in northern Australia. When the eggs are ingested with food, the larvae hatch in the small intestine and develop into adult worms, 3 to 5 cm long, which attach by means of their whiplike end to the mucosa of the large intestine. The majority of infected persons are asymptomatic but in heavy infection, abdominal cramps, diarrhoea, distension, nausea and vomiting and even rectal prolapse may develop.

Diagnosis

- Identifying characteristic eggs in faecal smears

Treatment

- Oral mebendazole or albendazole

Scrub typhus

- Scrub typhus is found in South-East Asia, northern Australia and the western Pacific. It is caused by *Rickettsia tsutsugamushi*, which is transmitted by mites.

Features

- Abrupt onset febrile illness with headache and myalgia

- A black eschar at the site of the bite with regional and generalised lymphadenopathy
- Short-lived macular rash
- Can develop severe complications (e.g. pneumonitis, encephalitis)

Diagnosis

- Serological assays and microimmunofluorescence

Treatment

- Doxycycline 100 mg bd for 7 - 10 days

Queensland tick typhus

Queensland tick typhus, which is caused by *Rickettsia australis*, is directly related to a tick bite. The symptoms are almost identical to scrub typhus, although less severe, and the treatment is identical.

Melioidosis

This serious disease with a high mortality is caused by the Gram-negative bacillus, Burkholderia pseudomallei, a soil saprophyte that infects humans mainly by penetrating through skin wounds, especially abrasions. It is mostly acquired while wading in rice paddies. It is mainly a disease of third world countries and occurs between 20° North and 20° South of the equator, mainly in South-East Asia and including northern Australia. It may manifest as a focal infection or as septicaemia with abscesses in the lung, kidney, liver or spleen. It presents with fever, headache, cough, pleuritic pain and generalised myalgia. It is called the 'Vietnamese time bomb' because it can present years after the initial infection in Vietnamese war veterans.

DxT: fever + pneumonia + myalgia ° melioidosis

Diagnosis

- Blood culture, swabs from focal lesions, haemagglutination test

Treatment (adults)[4, 6]

Cotrimoxazole 320/1600 mg (o) or IV, 12 hourly + ceft azidime 2g IV, 6 - 8 hourly *or* meropenem *or* imipenem all for at least 14 days followed by oral cotrimoxazole or doxycycline bd for 3 months

Prevention

- Traumatised people with open wounds (especially diabetics) in endemic areas (tropical South-East Asia) should be carefully nursed.

Ciguatera

This is a type of fish food poisoning caused by eating tropical fish, especially large coral trout and large cod, in tropical waters (e.g. the Caribbean and tropical Pacific). The problem is caused by a type of poison that concentrates in the fish after they feed on certain microorganisms around reefs. Ciguatera poisoning presents as a bout of 'gastroenteritis' (vomiting, diarrhoea and stomach pains) and then symptoms affecting the nervous system, such as muscle aching and weakness, paraesthesia and burning sensations of the skin, particularly of the fingers and lips. There is no cure for the problem but it can be treated with IV fluids and possibly mannitol or gammaglobulin. It is unwise to eat large predatory reef fish, especially their offal (mainly the liver).

PREGNANCY AND TRAVEL

Most international airlines do not allow passengers to travel after the thirty-sixth week of pregnancy and may require a doctor's certificate after twentyeight weeks. Air travel is contraindicated in the last month of pregnancy and until the seventh day after delivery. The past obstetric history should be taken into account. The same health risks apply except that most antimalarial tablets and vaccinations are not recommended. Only chloroquine is definitely safe. Live vaccinations (measles, rubella, influenza) are generally contraindicated[8] but the WHO considers it safe to have polio vaccine. Administration of killed or inactivated vaccines, toxoids and polysaccharides is permitted during pregnancy. Yellow fever vaccine is considered safe after the sixth month. As a general rule pregnancy and travel to third world countries do not mix and pregnant women should be advised not to travel to these countries.

Tetanus immunisation is important as protection is passed on to the child during early infancy. Immunoglobin can be safely given as prevention against hepatitis.

The antimalarial drugs chloroquine, quinine and proguanil may be given to pregnant women but all others mentioned in Table 14.2 (on p. 117) are contraindicated.

CHILDREN AND TRAVEL

Although children, including infants, are good travellers and adapt well, their resistance, especially to heat and infections, is lower. A child can suffer from acute dehydration very rapidly.[8] Air travel is not recommended for infants of less than 7 days or premature infants.

The change in atmospheric pressure on landing can cause distressing ear pain, so taking a bottle during descent is recommended.

In tropical areas it is important to keep children well hydrated and they should wear loose cotton clothing. A good guide to the health of children is the amount and colour of their urine. If it is scanty and concentrated they are not getting sufficient fluid.

Most vaccines (diphtheria, tetanus, poliomyelitis, BCG) can safely be given in the first few weeks of life. Measles is common overseas and it is worthwhile considering it even under 12 months. Yellow fever vaccine should not be given under 12 months. Hence the importance of protection against mosquito bites. Malaria prophylaxis is important. Chloroquine, proguanil and quinine may be given safely to infants. However, as a rule young children should be discouraged from travel.

AIR TRAVEL

Air travel is safe and comfortable, but jet lag and air sickness are problems that face many travellers.

Jet lag

This is the uncomfortable aftermath of a long flight in which the person feels exhausted and disoriented, and has poor concentration, insomnia and anxiety. The problem on arrival is poor concentration and judgment during the daytime.

Other symptoms that may occur include anorexia, weakness, headache, blurred vision and dizziness.

Jet lag is a feature of flying long distances east - west or west - east through several time zones, causing the person's routine daily rhythm of activity and sleep to get out of phase. The worst cases appear to be in those travelling eastbound from England to Australia. It can occur with travel in any direction, but the north - south flights are not so bothersome.

Factors influencing jet lag

Personal factors. These include age, state of health, tolerance to change, preparation for the long trip and, very importantly, the emotional and mental state.

General factors. Noise, vibration, air humidity and sitting still for long periods can influence jet lag.

Specific factors. Duration of the flight, time of departure, and changes in climate and culture at the destination affect the severity of jet lag. The problem is aggravated by:

- stress of the pretrip planning
- last minute rushing and anxiety
- lack of sleep during the trip
- overeating and excessive alcohol during the flight
- smoking

How to minimise the problem (advice to patients)

Before the flight

- Allow plenty of time for planning.
- Plan a stopover if possible.
- If possible, arrange the itinerary so that you are flying into the night.

- Ensure a good sleep the night before flying.
- Ensure a relaxed trip to the airport.
- Take along earplugs if noise (75 - 100 decibels) is bothersome.

During the flight

- *Fluids*. Avoid alcohol and coffee. Drink plenty of non-alcoholic drinks such as orange juice and mineral water.
- *Food*. Eat only when hungry and even skip a meal or two. Eat the lighter, more digestible parts of your meals and avoid fatty foods and rich carbohydrate foods.
- *Dress*. Women should wear loose clothes (e.g. long skirts, comfortable jeans, light jumpers) and avoid restrictive clothing. Wear comfortable (not tight) shoes and take them offduring the flight.
- *Sleep*. Try to sleep on longer sections of the flight (give the movies a miss). Close the blinds, wear special eye masks and ask for a pillow. Sedatives such as temazepam (Euhypnos or Normison) or antihistamines can help sleep.
- *Activity*. Try to take regular walks around the aircraft and exercise at airport stops. Keep feet up when resting, and exercise by flexing the major muscles of the legs. Avoid resting the calves of legs against the seat for long periods. Rest without napping during daylight sectors.
- *Special body care*. Continually wet the face and eyes. A wetting agent such as hydromellose 0.5% eye drops can help those with a tendency to sore eyes.

At the destination

Take a nap for 1 - 2 hours if possible.

Wander around until you are tired and go to bed at the usual time. It is good to have a full day's convalescence and avoid big decision making soon after arrival. Allow about three days for adjustment after the London to Australia flight.

Role of melatonin[10]

The importance of melatonin, a hormone secreted by the pineal gland, is somewhat controversial. However, a major review of trials by the Cochrane library pointed to its efficacy for alleviating jet lag. Oral medication in a 5 mg dose close to the desired or usual bedtime decreased jet lag. One possible explanation is that it has a low-grade hypnotic effect.

Who is fit to fly?

Patients with these problems should avoid flying[8] or be assessed for fitness:

- upper airways congested by infection, including influenza, e.g. within 6 weeks of severe acute respiratory illness
- acute gastroenteritis
- severe respiratory disease (COPD, chronic bronchitis, pneumothorax)
- recent thoracic surgery
- cystic fibrosis
- pulmonary tuberculosis (people should not fly until rendered non-infective)
- past history of respiratory problems while flying (dyspnoea, chest pain, confusion)
- unstable heart failure
- severe anaemia (below 7.5 g/dL)
- pregnancy beyond 200 days (28 weeks) (up to 36 weeks if necessary)
- previous violent or unpredictable behaviour
- within 7 days of a myocardial infarction
- within 3 days of a cerebrovascular accident
- within 5 - 10 days of major surgery
- brain tumour or recent skull fracture
- recent eye surgery
- severe or poorly controlled hypertension
- poorly controlled epilepsy

Special precautions are required by travellers with the following problems:

- *Colostomy*. Patients should wear a large colostomy bag and take extra bags.
- *Varicose veins*. Such patients should wear supportive stockings and exercise frequently.
- *Plaster casts*. Those with broken limbs in plaster should be careful of swelling.

- *Pacemakers*. Those with pacemakers may have a problem with X-rays at some overseas airports. Mention it to security officials before passing through security equipment.
- *Epilepsy*. Medication should be increased on the day of travel.
- *Diabetics*. Diabetics should discuss their therapy and control with their doctor. They should carry sweets.

Prevention of DVT

There is a risk of DVT in any person flying on longinternational flights. Risk factors include: increasingage, clotting tendency, i.e. thrombophilia, past historyof DVT, family history of DVT, smoking, obesity, varicose veins, dehydration, significant illness, recentmajor surgery and oestrogen therapy.

Prevention is by:

- keeping hydrated - drink ample fluids but avoid alcohol and caffeine drinks
- in-flight exercises, such as foot pumps, ankle circles, knee lift s
- compression stockings
- medication for those at risk, e.g. injections of LMW heparin for those with thrombophilia

TRAVEL SICKNESS

Almost everyone is sick when sailing on rough seas. However, some people, especially children, suffer sickness from the effect of motion on a boat, in a car or in a plane. The larger the boat, plane or car, the less the likelihood of sickness; travel by train rarely causes sickness. Nearly all children grow out of the tendency to have travel sickness, but many adults remain ‘bad’ sailors.

The problem is caused by sensitivity of the semicircular canals of the inner ear. They are affected by the movement and vibration of travel. Some people have sensitive inner ear canals and are prone to sickness, especially on certain types of journeys (e.g. winding roads through hills) and in certain vehicles.

The main symptoms of travel sickness are nausea, vomiting, dizziness, weakness and lethargy. Early signs are pallor and drowsiness, and sudden silence from an active, talkative child.

How to minimise the problem

1. Keep calm and relaxed before and during travel. With children avoid excitement and apprehension about the travelling. Encourage activities such as looking at distant objects; discourage activities such as reading and games that require close visual concentration.
2. Lie down, if possible, because this rests the inner ear canals and reduces the urge to vomit. If travelling by car, stop regularly for breaks. Passengers should use the front seat if possible.
3. Do not have a large meal a few hours before the journey or during it; avoid milk and fried or greasy foods. Do not travel with an empty stomach: have a light simple meal about an hour before and do not drink too much. Glucose drinks such as lemonade are suitable, as are glucose sweets and biscuits while travelling.

Medication for travel sickness

Many medicines are available for travel sickness. They include hyoscine, various antihistamines and other phenothiazine derivatives, all of which can cause drowsiness; although a problem to drivers, this sedative effect may be helpful for children or for those travelling long distances by plane.

Phenothiazine derivatives that provide appropriate anti-labyrinthine activity include prochlorperazine (Stemetil), promethazine hydrochloride (Phenergan) and promethazine theoclate (Avomine).

Combination antihistamine and hyoscine preparations for travel sickness include Travacalm and Benacine (Table 14.6).

Table 14.6 Medication to consider for motion sickness

Drug (genre)	Brand names and formulations	Dosage	
		Adults	*Children*
Antihistamines			

Dimenhydrinate	Andrumin 25 mg, 50 mg	50 mg statim then 4 hourly prn (max. 300 mg/24 hours)	avoid < 2 years
			2 - 6 years: 6.25 mg
			6 - 8 years: 12.5 mg
			8 - 12 years: 25 mg
	Dramamine 25 mg, 50 mg: Syrup 12.5 mg/5 mL		> 12 years: 50 mg tds (max. 3 - 4 doses/24 hours)
Pheniramine	Avil, 10 mg, 50 mg	25 - 50 mg tds	infants 10 mg bd
	Syrup 3 mg/mL		< 10 years: 10 mg tds
			> 10 years: 10 - 20 mg tds
Promethazine theoclate	Avomine 25 mg	25 mg statim or nocte for long journeys	> 5 years: ¼ tab
			5 - 10 years: ½ tab
			> 10 years: 1 tab
Promethazine hydrochloride	Phenergan 10 mg, 25 mg	25 mg bd	1 - 5 years: 5 mg bd
	Syrup 1 mg/mL		5 - 12 years: 10 mg bd
Related phenothiazines			
Prochlorperazine	Stemetil 5 mg suppositories 5 mg, 25 mg	5 - 15 mg tds	0.2 mg/kg bd or tds
			> 10 kg: avoid
Hyoscine			
Hyoscine hydrobromide	Kwells 0.3 mg tab Travacalm HO	1 - 2 tab statim then 1 tab 4 - 6 hour prn (max. 4/24 hours)	2 - 7 years: ¼ tab > 7 years: ½ tab (max. 4 doses/24 hours)
	Scop 1.5 mg transdermal	1 patch per 72 hours	avoid < 10 years
Combinations			
Hyoscine (0.2 mg) + Dimenhydrinate (50 mg) + Caffeine (20 mg)	Travacalm original	1 - 2 tabs statim (max. 4/24 hours)	> 2 years: avoid
			2 - 3 years: ¼ tab
			4 - 7 years: ¼ - ½ tab
			8 - 13 years: ½ - 1 tab (max. 4 doses/24 hours)

General rules: All tablets should be taken 30 - 60 minutes before departure and repeated 4 - 6 hourly as necessary (aim for maximum of 4 doses per 24 hours). Antihistamines should be used less frequently and some may be used once a day. Take care with drowsiness, pregnancy, the elderly and prostatic problems. Common adverse effects are drowsiness, irritability, dry mouth, dizziness and blurred vision, which are compounded by alcohol, antidepressants and tranquillisers. Hyoscine overdosage (from skin discs) can include confusion, memory loss, giddiness and hallucinations.

Hyoscine comes in tablet form, either alone or in combination and in the now popular adhesive patches.

Recommended medications

Car travel: adult passengers and children

- Dimenhydrinate (Andrumin, Dramamine) *or* Promethazine theoclate (Avomine) *or* Hyoscine (Kwells) These preventive oral preparations should ideally be taken 30 - 60 minutes before the trip and can be repeated 4 - 6 hourly during the trip (maximum 4 tablets in 24 hours).
- Hyoscine dermal discs (Scop) One of these adhesive patches should be applied to dry unbroken hairless skin behind the ear, 5 - 6 hours before travel and left on for 3 days. Wash the hands thoroughly after applying and removing the disc - be

careful of accidental finger-to-eye contact.

Sea travel

Sea travel generally poses no special problems apart from motion sickness and the possibility of injuries in the aged. The larger the ship, the less likely the problem. Those prone to sea sickness are advised to take antiemetics 60 minutes before sailing and for the first 2 days at sea until they obtain their ‘sea legs’. However, the use of hyoscine transdermal delivery systems is recommended for convenience.

Experienced seamen's ‘tricks’:

- always keep looking to the horizon
- plug one ear with cotton wool or ‘Blu-Tack’
- drink ginger drinks, e.g. ‘dry ginger’, ginger beer

Severe sea sickness. The standard treatment is promethazine (Phenergan) 25 mg IM injection. If injections are not possible, prochlorperazine (Stemetil) suppositories can be used.

The aged. Generally the elderly travel well but should take safeguards to avoid falls. The Chief Surgeon on P & O's flagship recommends that elderly people should bring the following:

- a letter from their doctor stating diagnosis and medication
- a spare set of spectacles
- a spare set of dentures
- a walking stick (if appropriate)

ALTITUDE SICKNESS[11]

High altitudes pose special problems for people who live at low altitude, especially if they have heart and lung disease. The severity depends on altitude, the speed of ascent, the temperature and level of activity. The high altitudes of Africa (Kilimanjaro, Kenya), India, Nepal (Himalayas), the Rockies of Canada and the United States, and South America provide such problems. It is usually safe to trek under 2500 m altitude but problems may occur over 3000 m.

Forms

1. Acute mountain sickness (mild → severe)
2. High-altitude pulmonary oedema
3. High-altitude cerebral oedema

Clinical features

- Usually within 8 - 24 hours of exposure
- Frontal headache (worse in morning and when supine)
- Malaise, fatigue, anorexia, nausea, insomnia

More severe: fluid retention, dyspnoea, vomiting, dry cough, dizziness

Serious: marked dyspnoea, neurological symptoms and signs

Management

Prevention

- Careful acclimatisation with gradual ascent[11]
- Spend 2 - 3 days at intermediate altitudes
- Ascent rate less than 300 m per day above 3000 m
- Ample fluid intake
- Acetazolamide (Diamox) 250 mg 8 hourly the day before ascent; continue 3 - 6 days (deaths from mountain sickness have still occurred while on this drug)

Treatment

- Immediate (urgent and rapid) descent to below 2000 m
- Oxygen
- Dexamethasone (e.g. 4 mg, 6 hourly)

TRAVELLERS' MEDICAL KIT

If a person intends to travel for a long time the following represents a comprehensive medical kit. It should not be regarded as an alternative to seeking appropriate medical help if it is available. Typical examples of general items are included in brackets.

Materials

- Bandaids and elastoplast dressing strip
- Bandages (2 cotton gauze, 2 crepe × 10 cm)
- Pocket torch
- Steristrips or 'butterfly strips' (to patch small cuts)
- Sterile gauze and cotton wool
- Thermometer
- Scissors and tweezers
- Safety pins
- Water purification tablets

Topical items

- Antifungal cream
- Chlorhexidine/cetrimide antiseptic cream (Savlon)
- Condoms
- Corticosteroid cream (e.g. hydrocortisone)
- Insect repellent containing diethyl-m-toluamide (Muskol, Repellem or Rid)
- Insecticide spray
- Mosquito net repellent solution: permethrin (Ambush - ICI)
- Nasal spray or drops
- Stingose spray (for bites and stings)
- Strepsils
- UV antisunburn cream (factor 15+)

Medication checklist

The medications below marked with * usually require a prescription.

- Antibiotics *
  - norfloxacin 400 mg (6 tablets for 3 days)
  - cotrimoxazole (for children)
- Antacid tablets - for heartburn or indigestion
- Antimalarials * - where appropriate
- Diamox tablets * for acute mountain sickness
- Fasigyn * 2 g or Flagyl * 2.4 g - for amoebiasis or giardiasis
- Laxative (Senokot)
- Imodium * or Lomotil * - for diarrhoea
- Motion sickness tablets (Avomine, Kwells or Phenergan)
- Paracetamol tablets - for fever or pain
- Sleeping tablets * (temazepam, promethazine)
- Rehydration mixture (Gastrolyte)
- Throat lozenges

GENERAL TIPS FOR THE TRAVELLER

Checklist for 'at risk' countries

- 'If you can't peel, boil or cook it, don't eat it.'
- Boil or purify water, avoid dairy products, ice-cream, shellfish, food left in open, salads, watercress, ice and recooked or reheated food.
- Never walk around barefoot at night in snake areas (and use a torch).
- Always shake your shoes before putting them on.
- Never wear nylon items in hot tropical areas.
- Never bathe, wade or drink in rivers, lakes or harbours unless you know they are bilharzia free.
- Keep yourself well covered after dark and use a mosquito net.
- Use insect repellent on skin frequently.
- Use an insecticide spray in your bedroom.
- Seek medical help if bitten by an excited dog, after washing bite.

Other tips

- Organise a dental check before departure.
- Arrange stopovers on a long flight (if possible).
- Take along a spare pair of spectacles and adequate medication.
- Arrange health and travel insurance.
- Check out your nearest embassy/consulate when visiting remote areas or politically unstable countries.
- Consider a traveller's medical kit.
- Never carry a parcel or luggage through Customs to oblige a stranger or recent acquaintance.
- Abstain from sex with a stranger.
- Have a credit card that allows a quick cash advance or an airline ticket purchase (for many countries a policy of 'if you get sick, then get out' is necessary).
- Most death and injury among travellers is caused by motor accidents. Avoid buses in India (and elsewhere) - trains are safer.

REFERENCES

1. Oldmeadow M. *Travel Medicine*. Melbourne: Monash University. Proceedings of Update Course for General Practitioners, 1991: 4.
2. Grayson L, McNeill J. Preventive health advice for Australian travellers to Bali. Med J Aust, 1988: 149: 462 - 6.
3. Locke DM. Traveller's diarrhoea. Aust Fam Physician, 1990; 19: 194 - 203.
4. Spicer J (Chairman). *Therapeutic Guidelines*: Antibiotic (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 85 - 49.
5. Currie B. Malaria. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 293 - 6.
6. Bochner F. (Chairman). Australian Medicines Handbook. Adelaide, 2006; 202 - 215.
7. Munro R, Macleod C. Recommendations for international travellers. Mod Med Aust, 1991: August: 50 - 7.
8. World Health Organization. International Health for Travellers, 2001. In: The Cochrane Library, Issue 1, 2001. Oxford: Update soft ware.
9. Amichai B, Finkelstein E et al. Think cutaneous leishmaniasis. Aust Fam Physician, 1993; 22: 1213 - 17.
10. Herxheimer A. Melatonin and jet lag. In: The Cochrane Library, Issue 1, 2001. Oxford: Update soft ware.
11. McDonnell L. Altitude sickness. Aust Fam Physician, 1990; 19: 205 - 8.

CHAPTER 15. RESEARCH AND EVIDENCE-BASED MEDICINE

Not the possession of truth, but the effort of struggling to attain it brings joy to the researcher.

GOFFHOLD LASSING (1729 - 81)

RESEARCH AND EVIDENCE-BASED MEDICINE

Effective research is the trademark of the medical profession. When confronted with the great responsibility of understanding and treating human beings we need as much scientific evidence aspossible to render our decision making valid, credible and justifiable.

Research can be defined as ‘a systematic method in which the truth ofevidence is based on observing and testing the soundness of conclusionsaccording to consistentrules’ [1] or, to put it more simply, ‘research is organised curiosity’,[2] the end point being new and improved knowledge.

In the medical context the term ‘research’ tends to conjecturebench-type laboratory research. However, the discipline of general practice provides a fertile research area in which toevaluate the morbidity patterns and the nature of common problems in addition to the processes specific to primary health care.

There has been an excellent tradition of research conducted by GPs. Tim Murrell in his paper ‘Nineteenth century masters of general practice’ [3] describes the contributions of Edward Jenner, Caleb Parry, John Snow, Robert Koch and James MacKenzie, and notes that ‘amongthe characteristics they shared was their capacity to observe and record natural phenomena, breaking new frontiers of discovery in medicine using an ecological paradigm’.

This tradition was carried into the 20th century by GPs such as William Pickles, the first president of the Royal College of General Practitioners, Keith Hodgkin and John Fry, all of whom meticulously recorded data that helped to establish patterns for the nature of primary health care. In Australia the challenge was taken up by such people as CliffordJungfer, Alan Chancellor, Charles Bridges-Webb, Kevin Cullen and Trevor Beard in the 1960s,[4] and now the research activities of the new generation of GPs, academic-basedor practice-based, have been taken to a higher level with the development of evidence-based medicine (EBM).

Based on the work of the Cochrane Collaboration and the initiatives of Chris Silapy in particular it has developed in the context of Australian general practice andnow beyond that. The focus of EBM has been to improve health care and health economics. Its development has gone hand in hand with improved information technology. EBM is inextricably linkedto research.

The aim of this chapter is to present a brief overview of research and EBM and, in particular, to encourage GPs, either singly or collectively, to undertake research - simple or sophisticated - and also to publish their work. The benefits of such are well outlined in John Howie's classic text Research in General Practice.[5]

WHY DO RESEARCH?

The basic objective of research is to acquire new knowledge and justification for decision making in medical practice. Research provides a basis for the acquisitionof many skills, particularly those of critical thinking and scientific methodology. The discipline of general practice is special to us with its core content of continuing, comprehensive, community-based primary care, family care, domiciliary care, whole person care and preventive care. To achieve credibility and parity with our specialist colleagues we need to research this area with appropriate meth-odology and to define the discipline clearly. There is no area of medicine that involves such a diverse range and quantity of decisions each day as general practice, and therefore patient management needs as much evidence-based rigour as possible.

Our own patch, be it an isolated rural practice or an industrial suburban practice, has its own microepidemiological fascination. Thus, it provides a unique opportunity to find answers to questions and make observations about that particular community.

There are also personal reasons to undertake research. The process assists professional development, encouraging clear and critical thinking, improvement of knowledge and the satisfaction ofdeveloping new skills and opening horizons.

The author undertook many small studies on common everyday problems during 10 years in country practice to determine the most effective treatments for which no or minimal evidence in the literature could be found. Many of these recommendations for problems, such as tennis elbow, cold sores, aphthous ulcers, ingrown toenails, hiccoughs, back pain, nightmares, temporomandibular dysfunction and warts, appear in this text. Although the numbers were relatively small it was a useful study to compare treatments for about 10 or 20 cases to test hypotheses and allow trends to emerge. The results from a large controlled trial would, of course, take precedence over these recommendations if they differed. However, the exercise, albeit limited, added immense interest to one's practice, which at times can be tedious without such scholarly challenges.

An important reason to undertake research is to conform with quality assurance processes that are now being expected of practitioners. The significant processes evaluating our accountability for quality control include audits of our own records, studies of critical incidents and morbidity studies.

WHO SHOULD DO RESEARCH?

Any GP searching for answers to questions and who has the opportunities should undertake research. Research is largely opportunistic and for some may be an impulsive reaction to a fascinating observation and in others a carefully conceived plan.

It can be undertaken in solo practice where the ability for personalised supervision of outcomes in patients is unique but where double-blind controlled trials are nigh impossible.

Research can be collaborative, and in fact this is an excellent way to get started. This can occur in a group practice.

Practitioners with computer skills and information technology at their fingertips are ideally placed to undertake research. Many GPs who have started ‘small’ have progressed to great heights of research activity, especially using their computer skills. In the process of posing questions and eventually finding the answers they frequently refer to the experience as ‘good fun’.

Asking questions

We often ask questions during the course of managing patients and such questions can form the basis of a research project, however simple.

Typical questions might be:

- Is all night-time cough in children due to asthma?
- Is suicide or attempted suicide in adolescent males precipitated by sexual problems?
- Is cancer precipitated by stress or unhappiness?
- Is recurrent migraine caused by cervical dysfunction?
- Is there any significant difference in response to various antibiotics to treat otitis media inchildren presenting in general practice?
- Does the distribution of leaflets by the receptionist in the waiting room lead to increased immunisation rates or Pap smears?
- Are my patients satisfied with the services they receive?
- Does the provision of patient information leaflets for the management of hypertension (or diabetes) lead to better compliance?

RESEARCH ON WHAT?

General practice has its own unique characteristics including illnesscontent, processes, epidemiology, health services, quality assurance and doctor-patient relationships. The special contact with patients provides opportunities to evaluate the patient's perspective on health service delivery, psychosocial issues and communication skills. The old saying ‘dig where you are’ is relevant to all of us. GPs invariably develop their ownspecial interests and this is a logical area in which to conduct research. Conducting a morbidity and prescribing survey in a practice is a simple and fascinating study. If the results are added to a wider study invaluable information about the nature of general practice is obtained.[5, 6, 7]

The development of the International Classification of Health Problems in Primary Care and the International Classification of Process in Primary Care by the World Organization of National Colleges and Assemblies of General Practice and the WHO has greatly assisted the process of morbidity studies. This information is now presented in the publication International Classification of Primary Care (ICPC).[8]

Research in general practice obviously covers many clinical areas studied by other groups but we may ask different types of questions, study different populations and use different methodologies, especially qualitative methods.

It would be logical to conduct research on those common problems requiring continuing care by the GP. These include:

- alcohol problems
- allergic disorders (e.g. asthma, allergic rhinitis)
- anxiety and depression
- arthritis
- chronic back pain and neck pain

- cancer
- cardiovascular disorders
- diabetes mellitus
- epilepsy
- hypertension
- migraine and other headache

Special opportunities, such as the observation that certain diseases or conditions are linked with specific circumstances, present frequently in primary care. An example is the observation that a group of farmers who presented to theirrural practitioner overa period of time with lymphosarcoma were all exposed to a specific herbicide to control blackberry growth on their farms. This led to further statewide investigations of this association, which indicated a significant link between the agent and the disease.

UNDERSTANDING TERMINOLOGY

Validity and reliability

- An ideal method of collecting research material is one that is valid.
- A valid method is one that measures what itclaims to measure.
- A reliable method is one that produces repeatable results.

Validity refers to the 'true' answer, which must be relevant, complete and accurate. Three significant questions that evaluate validity are:[9]

- Is the study useful or is the result inconclusive?
- Do you accept the results of this study as applied to the source population?
- Do the results apply to the population in which you would be interested?

Internal validity refers to the adequacy of the study methods in reference to the study population, while external validity refers to the generalisability ofthe results to the general population.

Reliability refers to the stability of question-and-answer response and is most successfully measured by testing and then retesting (repeatedly). The most frequently used method of testing for repeatability is to repeat application of the test.

Sensitivity, specificity and predictive values

Sensitivity and specificity, which are integral to validity, are important considerations in decision making in medicine, particularly in choosing appropriate investigations for disease diagnoses. The method of calculation of sensitivity, specificity and predictive values is summarised in Figure 15.1.



	Test positive	**Test negative**		
Condition present	A True positive	C False negative	A + C	Sensitivity $\frac{A}{A + C}$ %
Condition absent	B False positive	D True negative	B + D	Specificity $\frac{D}{B + D}$
	Positive predictive value $\frac{A}{A + B}$ %	Negative predictive value $\frac{C}{C + D}$ %		

Sensitivity:	How often a test shows pathology when it is present
Specificity:	How often a test is normal when no pathology present
Positive predictive value:	Indicates the likelihood of the patient having disease when the test is positive
Negative predictive value:	Indicates the likelihood of the patient not having disease when the test is negative

Figure 15.1 Definitions of sensitivity, specificity and predictive values

The *sensitivity* of a test depends on the proportion of people with the characteristic (disease) in whom the test is positive (i.e. percentage positive with disease). The ultimate sensitive test is one that detects all true positive cases.

The *specificity* of a test depends on the proportion of people without the characteristic (disease) in whom the test is negative (i.e. percentage negative of healthy people). The ultimate specific test is one that detects all the truly negative (disease-free) cases. A *gold standard* test is one that is as close to 100% specificity and 100% sensitivity as possible.

A clinical example of sensitivity and specificity is presented in Table 15.1.

Table 15.1 The predictability of signs and symptoms for carpal tunnel syndrome[9]

	Sensitivity (%)	**Specificity (%)**
Paraesthesia	97	4
Waking at night	91	14
Anaesthesia	57	61
Phalen's test	58	54
Tinel's test	42	63
Two point discrimination test	6	98

Predictive values that are useful indices of validity can be expressed as positive and negative values. Consider the example of a patient presenting with haematuria. In general practice the positive predictive value for carcinoma as the cause would be less than 5% but is about 50% in the inpatient hospital setting.

Incidence and prevalence

It is easy to confuse the meanings of these two terms:

- *Incidence* refers to the number of new cases of a disease (or factor of interest) occurring in a defined population within a specified period of time.
- *Prevalence* refers to the total number of individuals who have the disease (or factor of interest) at a particular time in a population.

  - Example: the prevalence of multiple sclerosis in temperate climates is 1 in 1000 - 2000 compared with 1 in 10 000 in the tropics. The incidence of multiple sclerosis in the Australian state of Victoria (population 4 400 000) is 8 per 100 000 per year.

Bias

This is any effect occurring during the investigation that tends to produce results that depart systematically from the true values. Varieties of bias include *measurement bias*(e.g. fault with a sphygmomanometer recording blood pressure), *confounding bias* (e.g. influence of alcohol on a study investigating the association between stress and hypertension) and *selection bias*(e.g. using hospital outpatients in a community-based study).

Confounding

This is a situation in which a measure of the effect of exposure on risk is distorted by the association of exposure with other (known or unknown) factors that influence the outcome.[1] A confounder is a factor that distorts the apparent magnitude of the effectof a study on risk.

Chance

One must question the probability that the results favouring the experimental intervention could have occurred by chance; therefore, we resort to statistical help in the form of a probability statement or significance level.

HOW IS THE RESEARCH UNDERTAKEN?

'Getting started' can be quite difficult for the beginner. However, assistance that should be accessed is available from several sources including individualGPs with researchexperience, university departments of general practice and the RACGP research committee. It is appropriate to seek out a suitable supervisor for the study. A chronological method follows.

1. *The idea*. Start with an idea or question, which needs to be interesting, relevant, significant and answerable.[10] It may be appropriate to develop a hypothesis at this stage.
2. *Floating the idea*. Next, discuss theidea with colleagues or an appropriate accessible authority.
3. *Literature search*. Review the literature, for example, a Medline search or checking witha central research 'bank'.
4. *Preparing a plan*. This can be a short written plan outlining the methodology for the study.
5. *Evaluation of the plan*. Then, contact a supervisor or appropriate authority to evaluate the study plan, which may be referred to a reference group or research committee.
6. *Development of a protocol*:
   - Prepare background; outline objectives and develop a hypothesis.
   - Select target population using clear criteria and appropriate numbers.
   - Design the research:
     - qualitative or quantitative?
     - questionnaire/s
   - Assess internal validity.
   - Consider statistical implications early:
     - number of patients
     - method for data analysis
   - Recruit subjects and assistants.
   - Assess the time frame.
   - Assess the ethical considerations →ethics approval committee.
7. *Pilot study and timetable*. Consider a preliminary pilot study and project timetable.
8. *Seek funding*. Solicit advice for appropriate funding bodies.
9. *The study*. Conduct the study.
10. *Publication*. Prepare for publication.

RESEARCH DESIGN

Hypothesis development

The reasoning process of the researcher is based on the null hypothesis - that is, an experimental group does not differ from a control 'normal' group in outcome. The question toconsider is: 'What is the probability that results from the experimental intervention would have occurred by chance?'. The answer is based on a probability statement: 'The probabilitythat the positive results occurred by chance is less than 5% ($P < 0.05$)'.[11]

Selecting a representative sample of appropriate size

Two basic components of subject selection are sample size and sample representativeness. The latter should be selected in a well-controlled manner.

A common question is: 'What is the ideal size of the sample?'. There is no fixed answer but it must be adequate to produce statistically meaningful results.

Recruitment of patients is a particular skill and often hard work, but it is easier if the researcher has a large pool of patients with whom he or she enjoys a good relationship. A useful rule is to aim to approach $3n$patients if you wish to work with a sample size of n.

Some guidelines for choosing the sample size are:[11]

- the more the individuals in the population differ, the larger the required sample
- the more planned comparisons, the larger the size
- larger sample sizes allow detection of smaller differences

TYPES OF RESEARCH [1]

The two broad categories of research in general practice are qualitative research, which is based on observation and talking with people, and quantitative research, which is based on measurement and analysis of data collection.

Qualitative research

This research is basically concerned with evaluating human behaviour from the subjects' perspective. It is based on close observation and is expressed in a descriptive way.

Common qualitative approaches

- Phenomenology
- Ethnography
- Grounded theory
- Biography (life story, narrative enquiry)
- Case study

The methods used are:

- interviews (open ended, semi-structured)
- focus groups
- participation observation
- document analysis

Qualitative research is an excellent method for generating hypothesesand can lead to quantitative research.

Phenomenology

The central focus of philosophy/method is the lived experience of theworld of everyday life. It describes events, situations, experiences and concepts.

It provides:

- detailed descriptions of an experience or event as it is lived
- deeper understandings and sensitivities

- improved thoughtful provision of care

Examples:

- effects of Viagra (and other agents) on marital/sexual relationships
- experience of carers in Alzheimer's disease
- effects of workplace bullying on absenteeism

Ethnography (ethnos ° a nation)

This examines cultures, peoples and societies including subgroups, e.g. adolescents. It is the basis of anthropology. The investigator usually identifies one or more key witnessess (informants) and interviews them to clarify observations.

Grounded theory

This is the development of new theory through the collection and analysis of data. It seeks to identify the core social processes within a given context in order tobuild theory that is grounded in the reality of those being studied.

Quantitative research

Quantitative research is research based on the collection of data in numerical quantities. It is concerned with hypothesis testing, reliability and validity. It canbe classified broadly asobservational, which includes case control, cross-sectional and cohortstudies, and experimental, which includes the classic controlled trial.[12]

- *Case control (or retrospective) study*isan observational study inwhich people with a disease (cases) are compared with those without i(control group).
  - Examples: Patients with mesotheliomas were investigated for exposure to asbestosor other agents; the mothers of children born with birth defects were investigated for an association with drug intake during pregnancy.
- *Cross-sectional or prevalence study* follows a correlation approach using existing databases. It is a survey of the frequency of disease, risk factors or other characteristics in a defined population at one particular time.
  - Example: The prevalence of diabetes mellitus (diagnosed and undiagnosed) was investigated in an Aboriginal community living in a particular area of metropolitan Sydney.
- *Cohort (or prospective) study*is also referred to as ‘follow-up’. The study follows a group (cohort) of individuals with a specified characteristic or disease over a periodof time. Comparisons may be made with a control group.
  - Example: 120 patients with chronic sciaticawere followed over 10 years to determine the outcome of their pain and neurological deficit. These were compared with a matched group who had undergone laminectomy.
- *Clinical controlled trial*is an experimental study that tests for hypothesised outcomes. An intervention is conducted on a randomly selected group of people and compared with a matched control group not subject to the specific intervention. The objective is to establish a causal relationship between the intervention and thehypothesised outcome. The ideal scientific trial is a **double-blind trial** where neither staff nor the participating patient are aware the participant is in the intervention or control group. This is the typical study when assessing the outcome of a drug trial as compared with placebo.

Meta-analysis

Meta-analysis is the process that systematically assesses compatible randomised controlled trials by merging the data (from usually smaller and inconclusive trials)to draw a ‘firmer’ conclusion from larger numbers of subjects.

EVIDENCE-BASED MEDICINE

EBM is a process of basing clinical practice on validated information. According to one of its modern architects, David Sackett, it is ‘the explicit, judicious and conscientious use of current best available evidence in making decisions about the care of individual patients’.[13]According to Silagy ‘EBM is the integration of the best available scientific evidence with yourclinical expertise andknowledge, your intuition, your wisdom’.[14]

The process of using EBM should be very comfortable for GPs because scientific methodology and evidence is second nature to us and has been the basis of our clinical decision making prior toand subsequent to graduation.

The proposed five steps of EBM are similar to basic research methodology:[14]

1. Construct a clinical question or define the problem.
2. Search for the evidence.

3. Appraise the quality and relevance of the evidence.
4. Apply it to the care of an individual patient.
5. Evaluate how effective it is.

The statistical methodologies used in EBM cover the traditional research methods but there is an emphasis on the methods of risk reduction, absolute and relative risk reduction and numbers needed to treat. These definitions are included in the glossary of terminology on pages 138 - 9.

GPs have a responsibility to their patients to be well versed with the best evidence when making decisions about management (Table 15.2), whether it be for a minor surgical procedure, selection of drugs, selection of an investigation or referral to the most appropriate consultant. If the best evidence reveals that a certain practice we are using is of no value or there may be more efficaciousmethods to our favoured method, then we should be prepared to change. On the other hand, if we find that a certain method works for us and there is no current evidence that it is the most appropriate one or the evidence is equivocal, then there is no compelling reason to change.

Table 15.2 Levels of evidence

1	Evidence obtained from a systematic review of all relevant randomised trials.
2	Evidence obtained from at least one properly designed randomised controlled trial.
3	Evidence from well-controlled trials that are not randomised, or well-designed cohort or case - control studies, or multiple time series (with and without the intervention).
4	Opinions of respected authorities; based on clinical experience; descriptive studies; or reports of expert committees.

Source: Modified from the NH & MRC.

GPs need a healthy scepticism about what is best evidence and claims for treatment in addition to the skill of critical appraisal of research/evidence. We tend to be impressed by the perception that evidence is a numbers game. However, the great work of JamesLind (see p. 80) shows that facts do not necessarily involve large numbers.

For EBM to be accepted by GPs the information needs to be readily accessible, user friendly, significant, relevant and, perhaps, believable.

The strength of EBM is that it can provide the answers to very important everyday decisions, especially in screening and preventive medicine where guidelines have fluctuated over the decades. The recent RACGP guidelines for preventive activities in general practice[15] highlight the value of current evidence.

GPs are currently faced with important decisions about the effectiveness of complementary therapies, which are very tempting to embrace or trial when searching forways to manage difficult problems, such as chronic fatigue syndrome, fibromyalgia, chronic asthma, chronic pain syndromes and other chronic diseases. We are hopeful that EBM can provide theanswers to best practice in addition to evaluating individual therapies.

To counterbalance the strengths (potential or real) of EBM there are concerns that it will be seized by bureaucrats to develop ‘cookbook’ guidelines, Holy Writ or economic rationalisation. Others are concerned about the perceived lack of flexibility.A very interesting critical review, especially affecting psychiatry, has been presented by John Ellard in his paper ‘What exactly is evidence-based medicine?’.[16] He questions the validity of the evidence underpinning EBM and the biases of both the proponents of ‘science’ and ‘art’ with the caution of Louis Pasteur: ‘The greatest derangement of the mind is to believe in something because one wishes it tobe so’.[17]

Glossary of terms[18,19]

Apart from the terms and definitions used in preceding pages it is important to highlight the following terms used in EBM/research.

Experimental event rate (EER)

The percentage of subjects in the intervention group that experiencedthe event of interest.

Control event rate (CER)

The percentage of subjects in the control group that experienced the event of interest.

Risk (R)

The probability that an event (death or disease) will occur.

Relative risk (RR)

The ratio of the risk of the outcome (e.g. disease or death) in the treatment/exposure group compared with the control/unexposed group.

*RR*informs us how many times more likely an event will occur inthe treatment group compared with the control group.

Calculation: *RR* ° *EER*/*CER*

RR ° 1 means no difference, so treatment has no effect

RR > 1 means the treatment increases the risk disease/death

RR < 1 means the treatment decreases the risk

Example: If the risk of death from people exposed to inhalation of anthrax spores is reduced from 10 in 100 cases to 2 in 100 cases with 60 days of prophylactic ciprofloxacin, the *RR*of death in this group is 0.20 or 20%.

Absolute risk reduction (ARR)

The absolute difference in event rates between two intervention or treatment groups. It gives an indication of the baseline risk and treatment effect. An *ARR*of 0 means no difference and thus the treatment has no effect.

Example: The *ARR*for prophylactic ciprofloxacin in the case cited is 10-2 ° 8 per 100 (0.08) or 8%.

Relative risk reduction (RRR)

The proportional reduction of adverse events between the treatment/experimental and the control groups in a trial (i.e. *RRR*is the ratio of the absolute risk reduction to the risk of the outcome in the control group). An alternative way to calculate the *RRR*is to subtract the *RR*from 1 (i.e. *RRR* ° 1 -*RR*).

In the example it is: 1-0.2 ° 0.80 or 80% or: RRR ° ARR/10 ° 8/10 ° 0.80 or 80%

*RRR*is probably the most commonly reported measure of treatment effects but the *ARR* gives a more realistic picture.

Number needed to treat (NNT)

The number of people who must be treated over a given period of time with the experimental therapy (specific intervention) to achieve one good outcome or prevent one adverse outcome. This incorporates the duration of treatment. It is a measure of the absoluterelative risk. Obviously the lower the *NNT*the better the treatment. It is calculated as 100/*ARR*(%); that is, the reciprocal of the *ARR*.

In the above example the number needed to treat would be 12 to 13 for60 days.

Note: The NNT will be different for different patient populations depending on their baseline risk for developing the outcome of interest.

Analysis of variance

This allows comparisons between the means of two samples of similar populations with a normal distribution. The contribution to variance for each variable can be determined and tested for statistical significance.

Clinical significance

The benefit to people receiving an intervention compared to the control group being great enough to warrant the intervention. It is based on measure of effect.

Statistical significance

The likelihood of a difference between two groups being real. It is the possibility that the difference occurred by chance alone. It is based on confidence intervals and *P*values. A 0.5 confidence level suggests that 19 times out of 20 the finding of the difference between the two groups is valid.

P value

The probability that an observed difference occurred by chance. The standard convention is that there is only a 5% (1 in 20) probability that the difference would fall outside this range by chance alone (i.e. a *P*value of 0.05 or 5%).

Confidence interval

The statistically derived range of values around a trial result in which the probability is that the true result will be within the range.

A 95% (standard) confidence interval for a sample indicates that there is a 95% chance that the interval includes the true population proportion whose care complies with the evidence. It isa measure of the certainty that the figures are correct.

Odds ratio

The probability of the occurrence of an event to its non-occurrence.

Type I error

A type I error occurs when a study concludes that there is a difference between two groups when there is no difference.

Type II error

A type II error occurs when a study concludes that no difference exists between groups when there is a true difference

Kappa

Cohen's kappa measures the agreement between the evaluations of two raters when both are rating the same object. A value of 1 indicates perfect agreement. A value of 0 indicates that agreement is no better than chance. It is an appropriate statistic for tables that have the same categories in the columns as in the rows (e.g. when measuring agreement between two raters).

Accessing the evidence

By computer/CD-ROM:

- The Cochrane library includes:
  - The Database of Systemic Reviews
  - Database of Abstracts of Reviews of Effectiveness
- Search tool: http://www.update-software.com/publications/cochrane
- Browse: http://www.nicsl.com.au (useful links - Cochrane library)
- Free online: http://www.update-software.com/publications/cochrane
- Bandolier: http://www.jr2.ox.ac.uk/bandolier/ A UK journal on EBM with excellent summaries on NNT for a host of common interventions

Publications

- Clinical evidence: BMJ Publishing Group, refer to < http://www.clinicalevidence.org >
- Evidence-based medicine. BMJ Publishing Group

CRITICAL APPRAISAL OF PUBLISHED RESEARCH

The objective of critically appraising a paper is to determine if themethods and results of the research have significant validity to produce useful information. The appraisal starts with a careful review of the abstract, which ideally should be presented in a structured format.

The critical questions are as follows:

1. What were the objectives of the study?
2. Were the ethical aspects properly followed?
3. What was the study design?
4. Were there any potential problems associated with the design?
5. Were all the patients who entered the studyproperly accounted for at its conclusion?
6. What were the important results?
7. How would you interpret and explain these results?

FURTHER READING

Armstrong D et al. *Research Methods for General Practice*. Oxford: Oxford Medical Publications, 1990.

Howie JGR. *Research in General Practice*(2nd edn). London: Chapman & Hall, 1992.

Sackett DL et al. *Evidence Based Medicine: Howto Practice and Teach EBM*. London: Churchill Livingstone, 1997.

Sackett DL, Haynes RB, Guyatt GH, Tugwell P. *Clinical Epidemiology: A Basic Science for Clinical Medicine* (2nd edn). Boston: Little Brown & Co., 1991.

Silagy C, Haines A. *Evidence Based Practice in Primary Care*. London: BMJ Books, 1998.

REFERENCES

1. Schattner P. Introduction to research and research designs. In: Piterman L ed. *Introduction to Research in Family Medicine MFM 2006*, 1996: 2 - 7.
2. Eimerl T. Organised curiosity. JR Coll Gen Pract, 1960; 3(1): 246 - 52.
3. Murrell TGC. Nineteenth century masters of general practice. Med J Aust, 1991; 155: 785 - 92.
4. Anderson NA, Bridges-Webb C, Chancellor A. *General Practice in Australia*. Sydney: Sydney University Press, 1986: 124 - 30.
5. Howie JGR. *Research in General Practice*(2nd edn). London. Chapman & Hall, 1992: 12 - 14.
6. Bridges-Webb C. The Australian general practice morbidity and prescribing survey, 1969 - 1976. Med J Aust (Suppl.) 1976; 2: 5 - 20.
7. Bridges-Webb C, Britt H, Miles DA et al. Morbidity and treatment in general practice inAustralia 1990 - 1991. Med J Aust, 1992; 157: 51 - 7.
8. Lamberts H, Wood M. *ICPC: International Classification of Primary Care*. Oxford: Oxford University Press, 1987.
9. Buch-Jaeger N, Foucher G. Carpal tunnel syndrome: validity of clinical signs. J Hand Surg, 1994; 19B: 72 - 4.
10. Silagy CA, Schattner P eds. *An Introduction to GeneralPractice Research*. Melbourne: Monash University, Department of Community Medicine, 1990.
11. Rakel RE. *Essentials of Family Practice*. Philadelphia: WB Saunders Company, 1993: 182 - 91.
12. Fowkes FGR, Fulton PM. Critical appraisal of published research: introductory guidelines. Br Med J, 1991; 302: 1136 - 40.
13. Sackett DL et al. Evidence Based Medicine: what it is and what it isn't. Br Med J, 1996; 312: 71 - 2.
14. Silagy C, Haines A. *Evidence Based Practice in Primary Care*. London: BMJ Books, 1998.
15. RACGP. *Guidelines for Preventive Activities in General Practice*(5th edn). Melbourne RACGP: 2001.
16. Ellard J. What exactly is evidence based medicine? Mod Med Aust, 1997; (September): 22 - 5.
17. Pasteur L. Speech to the French Academy of Medicine, July18, 1876. In: Strauss MB, ed. *Familiar Medical Quotations*. Boston: Little Brown & Co., 1968: 502.
18. Rosser M, Shafir MS. *Evidence-based Family Medicine*. Hamilton: BC Decker Inc., 1998.
19. NPS News. Drug promotion: distinguishing the good oil from snake oil. NPS News, 2002; 25: 1 - 4.

CHAPTER 16. LABORATORY INVESTIGATIONS

As is your pathology, so is your medicine!

SIR WILLIAM OSLER (1849 - 1919)

LABORATORY INVESTIGATIONS

Appropriate use of the laboratory, particularly the judicious selection of investigations, is an important skill for the general practitioner to perfect. It is wise to remember that the laboratory staff includes clinical pathologists, microbiologists and haematologists, who can offer invaluable assistance and advice. Hence, it is important to provide a properly collected specimen accompanied by a succinct and relevant clinical history.

This section discusses useful investigations and their clinical interpretation, including some that tend to mystify. A summary of reference values appears at the end of the chapter.

It is advisable for practitioners to be conversant with the specificity and sensitivity of the various tests in order to make rational decisions about their interpretation and to provide appropriate counselling to their patients.

POLYMERASE CHAIN REACTION (PCR)

PCR is one of the newer wonder tests that is linked to the genetic material DNA and RNA. It is a type of nucleic acid amplification technology that has opened the frontiers of improved diagnosis in virology, and slow-growing and fastidious organisms. More than 60 of these tests are now available and the scope is growing.

Polymerase is an enzyme that catalyses formation of nucleotides into DNA molecules before cell division, or RNA molecules before protein synthesis.

PCR is a process that permits making, in vitro, exponential numbers of copies of genes. This is initiated with a single molecule of DNA, leading to the generation of billions of similar molecules within a few hours. This has huge practical importance as a method of investigating genetic material. Thus, the technique of PCR can be used in investigating bacterial infections, parasites, viruses associated with cancer, human immunodeficiency virus (HIV), genetic disorders such as diabetes and breast cancer, and various disorders of the blood such as thalassaemia and of muscles.

ERYTHROCYTE SEDIMENTATION RATE (ESR)[1]

The ESR relies on the principle that blood components separate faster in illness. It is mainly determined by the effect of serum proteins on the negative electric charge on the erythrocyte surface. The ESR is a marker of inflammation and malignant disease (Table 16.1). It reflects the presence of all acute-phase proteins (especially fibrinogen) as well as the immunoglobulins. It should be used to screen asymptomatic patients for the presence of disease.

Table 16.1 Relative values (mm/h) of typical examples of ESR readings

Very high (up to 100+ mm/h)	High (40 - 80 mm/h)	Moderate to low elevation (20 - 40 mm/h)	Low (<1 mm/h)
Giant cell arteritis/polymyalgia rheumatica/temporal arteritis	Rheumatic fever	Most acute and chronic infections (e.g. recent viral)	Idiopathic - normal
	Pyelonephritis		Sickle-cell anaemia
	Other bacterial infections	Severe other illness	Polycythaemia
Multiple myeloma	Viral infections with cold agglutinins	Anaemia	NSAIDs

Tuberculosis		Pregnancy	Old specimen
Deep abscess	Collagen disorders (e.g. rheumatoid arthritis, SLE)	Drugs, especially contraceptives	
Bacterial endocarditis			
Acute osteomyelitis	Solid tumours, especially metastases	Elevated serum cholesterol level	
	Leukaemia/lymphomas	Laboratory error (e.g. tilted tube)	
	Myocardial infarction Inflammation of healing		
		Idiopathic - normal	

There is a lag phase of 24 - 48 hours between the onset of inflammatory stimulation and the production of inflammatory proteins that increase in the ESR. There is also a delay in the fall of the ESR after resolution of the inflammation because the fibrinogen levels can remain elevated for 6 days or so after acute tissue damage - this can take up to 4 - 8 weeks to return to normal.

A normal value of <20 mm/h generally excludes inflammation. The OCP can push the level to 20 - 25 mm/h.

Normal values of ESR - reference interval

• Child:	2 - 15 mm/h
• Adult male	
17 - 50 years:	1 - 10 mm/h
>50 years:	2 - 15 mm/h
• Adult female	
17 - 50 years:	3 - 12 mm/h
>50 years:	5 - 20 mm/h

C-REACTIVE PROTEIN (CRP)[1]

C-reactive protein was discovered in the blood by Tillet and Francis in 1930 and named because of the manner it reacted with the C polysaccharide of *Streptococcus pneumoniae.*

It is an important product of the acute-phase response and is generally accepted as the most accurate measure of the acute-phase response and, hence, of tissue inflammation. Like the ESR it is a non-specific marker of inflammation and neoplastic disease.

CRP levels rise within 6 hours and may double every 8 hours, reaching peak levels at 50 hours. Levels can fall very rapidly but resolve with a 24-hour halflife following tissue injury.

- A CRP level > 100 mg/L has an 80% sensitivity and 88% specificity for bacterial infection.
- A CRP level of 10 - 40 mg/L has a 69% sensitivity and 54% specificity for viral infection.[2]

Rules for inflammation

4 - 10 ° mild inflammation

10 - 20 ° moderate inflammation

>40 ° marked inflammation

The CRP can be used to follow the response to therapy (e.g. antibiotic treatment) or activity of disease (e.g. Crohn's, spondyloarthropathy).

Levels above 100 mg/L are more likely to be associated with bacterial infection. Refer to Table 16.2.

Table 16.2 A guide to C-reactive protein levels

Marked elevation >40 mg/L	Normal to mild elevation
Bacterial infection	Viral infection
Abscess	Ulcerative colitis
Crohn's disease	SLE
Active rheumatic disease:	Atherosclerosis
- rheumatic fever	
Connective tissue disorders:	Steroid/oestrogen therapy
- rheumatoid arthritis	
- vasculidities	
Malignant disease	
Trauma/tissue injury	

Comparison between ESR and CRP

- There tends to be a broad correlation between them.
- Both are markers of inflammation.
- The CRP levels rise faster than the ESR.
- The levels are similar after 24 hours or so.
- CRP levels fall faster than the ESR.
- CRP levels (unlike ESR) are not affected by pregnancy.
- ESR may be very high with a normal CRP in giant cell arteritis/polymyalgia rheumatica.
- CRP costs more. Normal value of CRP: <10 mg/L

RECOMMENDED TESTS FOR INFECTIOUS DISEASES[3]

Adenovirus

- Serum for antibody levels.

Amoebiasis

- Stool examination for trophozoites and cysts.
- Serum for antibody levels (positive titre is 1:128 or more) usually in extraintestinal amoebiasis only (e.g. hepatic). Note that the test remains positive for as long as 10 years after treatment.

Bordetella pertussis

- Nasopharyngeal swabs or aspirate (preferred) for culture and/or PCR studies.
- Serum for IgA detection - may take several weeks to rise, especially in infants and children. Hence, repeat testing may be necessary. Not affected by immunisation (no antibody response).

Brucella

- Serum for *Brucella* antibodies. Acute and convalescent (3 - 4 weeks) samples. Blood culture if febrile.

Cat-scratch disorder

- Acute and convalescent sera to detect a fourfold rise in *Bartonella henselae* infection.

Chickenpox/varicella zoster virus

- Usually a clinical diagnosis, but where it is unclear or atypical or to determine susceptibility to varicella zoster infection, take blood sample for viral antibody detection. A fourfold increase over 2 - 4 weeks supports diagnosis of acute varicella infection. Also, viral culture of vesicle fluid and smears of base of lesion for viral inclusions, fluorescent antibody tests or PCR.

Herpes simplex virus (HSV)

- The above methods can be used for detection of HSV but blood tests for antibodies are of limited use in differentiating type I or II herpes. If needed, culture and smears from the lesions for PCR testing is available.

Chlamydia

- *Chlamydia pneumoniae*: for atypical pneumonia - acute and convalescent blood samples for antibodies.
- *Chlamydia trachomatis*: for diagnosis of STIs, conjunctivitis and pneumonia in neonates - swabs or endotracheal aspirate for culture and PCR tests.

The PCR test does not necessarily require cervical or urethral swabs as it allows prompt diagnosis of genitourinary chlamydia infection in both males and females using the first 20 - 30 ml of the stream.

Patients should not have urinated for the previous 2 hours specimens should be stored in a yellow top urine container at 4°C and sent to the laboratory ASAP.

Chlamydia antibodies are non-specific in the diagnosis of genital tract infection or trachoma.

Clostridium difficile

- Fresh faeces to detect *Clostridium difficile* toxin and culture where antibiotic diarrhoea/colitis is suspected.

Cryptococcal infection

- *Cryptococcusr* antigens in specimens of blood and CSF (antibodies not so diagnostic).
- Positive in >95% of patients with cryptococcal meningitis.

Cytomegalovirus (CMV)

- Acute and convalescent (2 weeks) blood samples for CMV antibodies. A fourfold increase indicates recent infection. The presence of specific IgM in a neonate may represent intrauterine infection. This can be supplemented by PCR tests.

Epstein-Barr mononucleosis (EBM)

- EBM screening tests: white cell count, blood film, Monospot and Paul Bunnell.
- Antibody tests: blood for immunoassay for IgM and IgG antibodies to viral capsule antigen and Epstein-Barr nuclear antigen (EBNA) - used in those with the mononucleosis syndrome when EBM screening tests are negative.

Fungal (topical) infections

- Topical antifungals should be stopped at least 3 days before a specimen is taken.
- Collect specimen into a sterile yellow-topped jar. The more material submitted the greater the odds of positive findings.

- Skin lesions: take a scraping of the advancing edge.
- Nails: collect clippings and scrapings, include necrotic debris from beneath the nail.
- Hair: include hair roots (plucked hairs).
- Microscopy is usually reported immediately while cultures are maintained and examined on a regular basis for 2 - 4 weeks.

Hepatitis A, B, C, D, E

- Blood for immunoassay for respective antibodies and hepatitis B virus antigens.
- PCR tests for viral load with hepatitis B and C. Genotyping is available and important for determining those who benefit best from treatment. A PCR test on blood can be performed for hepatitis G virus.

HIV/AIDS

- Blood for HIV-1 and HIV-2 antibodies are routinely tested. Positive results indicate HIV infection. Negative results do not exclude infection if serum has been taken within 3 - 4 weeks of a risk exposure. Repeat testing is recommended after that time. The sensitivity of this screening test is almost 100%.
- HIV antigen may assist in the diagnosis of early HIV infection and neonatal HIV infection.
- Detuned ELISA testing can date the time of primary infection.

Markers

- CD_4 lymphocyte counts.
- HIV viral load.

Influenza

- Blood for antibodies: requires acute and convalescent (2 - 4 weeks) sera. Has low sensitivity and unhelpful for treatment as diagnosis is retrospective.
- Viral culture: can be isolated from nose and throat swabs and nasopharyngeal aspirates but the result can take some days.
- Direct antigen detection: a rapid test with high sensitivity and specificity but can only be performed on a nasopharyngeal aspirate.
- PCR test: this developing test is rapid and has good specificity and sensitivity.

Legionella

- Blood for antibodies: require acute and convalescent (4 - 6 weeks) sera. Requires a fourfold increase in titre to >128. A single titre of >256 is suggestive of infection.
- Sputum microscopy and culture is the quickest and most reliable test if *Legionella* sp. are identified.
- Urine antigen tests for *Legionella pneumophila* type I.

Leptospirosis

- Blood for antibodies will give a diagnosis according to levels matched with clinical features.

Malaria

- Thick and thin blood films for microscopic examination - usually require repeat examination (at least 3 at separate times).
- Serological tests (ELISA) are not commonly used but field-based card assays using agglutination tests are useful for travellers e.g. Paracheck V test.
- PCA methods are highly specific and sensitive but not widely practical at present because specialised laboratory methods are required.

Mumps

- Blood for antibody testing: diagnoses immune status and mumps infection (acute and convalescent sera) with IgM assays.
- CSF: the presence of IgG in CSF confirms the diagnosis of mumps meningitis although levels may be low for 2 - 3 days after the onset of the illness.

Mycobacterium tuberculosis

- Sputum (3 separate samples)/bronchial brushing or washings: acid-fast staining and microscopy.
- PCR testing on sputum.

Mycoplasma pneumoniae

- Blood for antibodies (acute and convalescent samples). Presence of IgM antibodies and a rise in titre indicates infection. High titres can persist for more than 12 months.

Q fever

- Blood in acute phase and 2 - 3 weeks after onset of illness for antibody levels.

Parvovirus B19

- Blood for antibody detection. Suspected fifth disease (erythema infectiosum) and other clinical conditions, such as maternal infection with hydrops foetalis, aplastic crisis in chronic haemolysis, polyarthritis and rash in adults. PCR for confirmation.

Rubella

- Blood for antibody tests - acute phase and convalescent (after 10 - 14 days).
- A fourfold increase in IgG indicates recent infection.
- IgM antibody becomes positive about 7 days after onset of the illness but will become undetectable after 8 weeks. IgM antibodies in maternal serum indicates high fetal risk, while in cord blood suggests congenital infection.

Toxoplasmosis

- Blood for acute and convalescent (2+ weeks) antibody testing. A fourfold rise in IgG titre is diagnostic for toxoplasmosis. IgM antibodies at a level >16 indicate recent infection. The diagnosis of congenital toxoplasmosis is supported by the presence of IgM antibodies. Antibody testing will date the infection.

Viral skin rashes[4]

When a patient presents with a fine maculopapular skin rash, serological tests can be performed for all of the following causative agents. Invariably, a rising antibody level between acute phase and convalescent sera (2 - 4 weeks) is required for diagnosis.

- Measles - rising IgM titre diagnostic (raised IgM ° previous infection or immunisation)
- Rubella - rising IgM or IgG ° recent infection
- Parvovirus B19
- Echovirus
- EBV
- CMV
- Ross River virus
- Barmah Forest virus
- Dengue fever
- Other arboviruses

Sexually transmitted infections[4]

- *Neisseria gonorrhoea*
  - Culture (urethral, cervical, rectal, pharyngeal)
  - PCR is excellent on cervical or urethral swabs or first-stream urine
- *Chlamydia trachomatis*
  - Antigen detection (PCR recommended) on cervical or urethral swabs or first-stream urine (preferably first 20 - 30 ml)
  - Culture available on request Serology not recommended
- Syphilis
  - Serology (RPR, TPHA, FTA - ABS, EIA)
- Hepatitis B
  - Serology
- HIV
  - Serology
- *Trichomonas vaginalis*
  - Microscopy from vaginal swab
  - PCR
- Herpes simplex
  - Viral culture
  - Antigen detection (PCR best)
  - Serology
- Lymphogranuloma venereum
  - *Chlamydia* serology
  - Lymph node biopsy
- Chancroid
  - Microscopy/culture for *Haemophilus ducreyi*
- Granuloma inguinale
  - Biopsy

Urinary tract infection (UTI)[4]

Microscopy

White blood cell count >10 per µL is abnormal and reflects an immune response to local infection.

- Higher counts have greater significance.
- Epithelial cells suggests the possibility of genital contamination in females (i.e. poor sample).

Culture

Counts are expressed as organisms per mL.

- Counts >10[5] organisms per mL are most significant.
- UTI can occur at lower counts, especially in pure growth and supported by the clinical picture and significant pyuria.
- Significant organisms are usually in pure growth (not mixed).
- Significant UTI are usually associated with a pyuria but may occur in its absence.

Fever in returning travellers

- Serology
  - Dengue fever (acute and convalescent)

  - Typhoid fever (acute and convalescent) - limited use
  - Viral hepatitis A, B
  - Amoebiasis - liver abscess
  - Typhus
- Blood culture
  - Typhoid fever (best to detect typhoid)
  - Meningococcal infection
- Full blood count (FBE), thick and thin film
  - Malaria
- Spot agglutination tests
  - Dengue Fever
  - Malaria
- Stool examination
  Culture
  - *Campylobacter, Salmonella, Shigella*,
    Typhoid
    Microscopy
  - *Amoeba, Giardia, others*
- Liver function tests
  - Hepatitis
- ESR
  - Screening

Lymphadenopathy

- FBE, ESR, mononucleosis screen (e.g. Paul Bunnell)
- Serology
  - EBV, CMV, HIV
  - Toxoplasma
  - Rubella, syphilis, cat-scratch disorder

INTERPRETATION OF IRON STUDIES[5]

A sound knowledge of the metabolism of iron and its transportation helps in the interpretation of iron studies (see Table 16.3). The serum (or plasma) level of *iron* falls gradually below the normal range (about 14 - 30 μmol/L) when the amount of iron in the body decreases after the iron reserves become exhausted. The level in the serum of *transferrin*, the major iron-transporting protein in the circulation, rises under these circumstances to perhaps above normal levels. A subnormal level of iron plus a high or normal level of *transferrin* is strong evidence of iron deficiency.

Table 16.3 The interpretation of iron studies[4]

Condition	Serum Fe	TIBC	% Transferrin Saturation	Ferritin
Iron deficiency	↓	Nor ↑	↓	↓↓
Thalassaemia	Nor ↑	N	Nor ↑	↑
Anaemia of chronic disease	↓	Nor ↓	↓	Nor ↑
Sideroblastic anaemia	Nor ↑	N	Nor ↑	↑
Haemochromatosis	↑	↓	↑↑	↑↑

Transferrin, as the carrier protein, binds most of the iron in the serum. The capacity of transferrin is represented by the total amount of iron that can be bound to serum protein, meaning that the *total iron-binding capacity* (TIBC) provides an alternative estimation of the concentration of transferrin.

Transferrin saturation is the extent to which the iron-binding sites on transferrin are occupied by iron. This is calculated by dividing the iron level by the serum iron-binding capacity. The percentage saturation is normally 20 - 55%. It is markedly elevated in haemochromatosis - above 50% - and is the key marker for that disorder.

The *serum ferritin* level bears a direct relationship to the amount of iron stores in the body and subnormal values can be detected when iron stores are exhausted even before the serum iron level has significantly declined. The normal range varies between sexes and age groups: 20 - 250 μ/L in males, 10 - 150 μg/L in females and lower again in children.

LIVER FUNCTION TESTS (LFTS)[4]

A comprehensive summary of liver function tests is presented in Chapter 58. The following parameters are tested.

Plasma bilirubin

- Unconjugated - from breakdown of red blood cells.
- Conjugated - after metabolism in the liver.

Albumin

- Synthesised in the liver with a half-life of 20 days (a good indicator of chronic liver disease, not acute).

Plasma transferases

- Alanine aminotransferase (ALT) - specific to liver.
- Aspartate aminotransferase (AST).
  Both are indicators of hepatocellular damage.

Plasma alkaline phosphatase (ALP)

- Present on the sinusoidal surface of hepatocytes and in bile canaliculi and ducts. Not specific to liver but an indicator of cholestasis (e.g. obstruction, infiltration, cirrhosis).

Gamma-glutamyl transferase (GGT)

- Present in bile canaliculi.
  Raised levels with cholestasis, other liver diseases and with drug and alcohol intake.

Differential diagnosis of jaundice[4]

Differentiating jaundice due to acute hepatocellular damage from extrahepatic obstruction on routine LFTs can only be suggested in the early stages according to the following guideline.

	Acute hepatitis	**Obstruction**
ALP	Normal to <3 times normal	ɰ times normal
ALT/AST	10 - 100 times normal	<10 times normal

Alcohol abuse

The following test indicators point to the diagnosis of alcohol excess:

- GGT - limited sensitivity and specificity.
- Mean corpuscular volume - macrocytosis, also limited sensitivity and specificity.
- Carbohydrate deficient transferrin

THYROID FUNCTION TESTS (TFTS)

A summary of thyroid function tests is given in Chapter 24, which includes a table (Table 24.2) of summarised tests. The key first-line TFT is the serum thyroid stimulating hormone (TSH) level, which has to be interpreted with care and thought. Because of its high sensitivity it can miss the occasional case of thyroid disorder, especially in the presence of underlying pituitary disorder, treated thyrotoxicosis and non-thyroidal illness. The next screening tests are serum free tri-iodothyronine (T_3) and thyroxine (T_4) tests. If there are discrepancies, then anti-thyroid antibody tests can be valuable in finding the cause of thyroid disorders. These include anti-thyroid peroxidase antibodies (especially), anti-thyroglobulin antibodies and anti-TSH receptor antibodies. Other tests include thyroxine-binding globulin and thyroglobulin. Follow-on tests may include nuclear medicine scanning and ultrasound. It is advisable to seek expert help in the interpretation of these tests, especially in patients with systemic illness.

SERUM ELECTROLYTE LEVELS[6]

An understanding of serum electrolyte levels is very important for the clinical implications of very ill patients, disorders of fluid loss and retention, and the use of cardiovascular drugs, especially diuretics. The key ions are potassium (K^+), sodium (Na^+), chloride (Cl^-), and bicarbonate (HCO_3^-). Normal intracellular and extracellular levels of sodium and potassium are fundamental for good health.

The anion gap

The anion gap is a useful clinical calculation to assess in general acid - base problems. It is calculated from the serum electrolyte values as the difference between the cation Na^+ and the sum of the two main anions Cl^- and HCO_3^-. The charges on the other cations (e.g. K^+) and anions (e.g. phosphate, PO_4^-) tend to balance out. Negatively charged plasma proteins account for most of the anion gap.

Anion gap ° $(Na^+) - (Cl^- + HCO_3^-)$

In a healthy person the anion gap is around 8 - 16.

An increased anion gap infers metabolic acidosis. Metabolic acidosis with a normal anion gap is called *hyperchloraemic acidosis* because the reduction in HCO_3^- is balanced by an increased Cl^- (e.g. chronic diarrhoea, kidney tubular acidosis).

Hypernatraemia

$Na^+ > 145$ mmol/L

Causes

- Water depletion (e.g. diabetes insipidus)
- Water and sodium depletion (e.g. diarrhoea)
- Corticosteroid excess (e.g. Cushing's syndrome, Conn's syndrome)

- Excess IV hypertonic Na solutions

Clinical features

- Thirst, confusion, oliguria
- Orthostatic hypotension
- Muscle twitching or cramps
- Severe: seizures, delirium, hyperthermia, coma

Hyponatraemia

$Na^+ < 135$ mmol/L

Causes

- Water retention (e.g. CCF, hypoalbuminaemia)
- Kidney failure to conserve salt (e.g. nephritis, diabetes mellitus)
- Gastrointestinal losses of Na^+ (e.g. diarrhoea, vomiting)
- Drugs (e.g. diuretics, ACE inhibitors)

Clinical features

- Lethargy, confusion, mental changes (e.g. in personality)
- Severe: convulsions, coma, death

Hyperkalaemia

$K^+ > 5$ mmol/L

The first sign of hyperkalaemia (e.g. > 6) may be a cardiac arrest.

Causes

- Kidney failure
- Acidosis
- Mineralocorticoid deficiency: Addison's disease (p. 224), aldosterone antagonists
- Excessive intake of K^+ (e.g. ↑ IV fluids with K)
- Drugs (e.g. spironolactone, ACE inhibitors, NSAIDs)

Clinical features

- Muscle weakness, flaccid paralysis (rare)
- May be asymptomatic until cardiac toxicity
- May cause cardiac arrest - asystole versus fibrillation
- ECG: peaked T waves, ↓ QT, ↑ PR interval→ arrhythmias

Hypokalaemia

$K^+ < 3.5$ mmol/L

Causes

- Kidney disease

- Gastrointestinal loss: vomiting, diarrhoea
- Alkalosis
- Mineralocorticoid excess: Cushing's syndrome, ↑ aldosterone
- Loss in extracellular fluid to intracellular (e.g. burns, other trauma)
- ↓ Intake of K^+
- Drugs (e.g. diuretics - frusemide, thiazides)

Clinical features

- Lethargy, muscle weakness and cramps, mental lethargy and confusion
- Severe flaccid paralysis, tetany, coma
- ECG: prominent U waves, depressed ST segment, ↓ T waves, arrhythmias

LABORATORY REFERENCE VALUES

The reference values and ranges for these blood tests are given in the system of international units (SI) and may vary from laboratory to laboratory. An asterisk (*) indicates that paediatric reference ranges differ from the adult range given.

Electrolytes/kidney	
Sodium	(135 - 145 mmol/L)
Potassium *	(3.5 - 5.0 mmol/L)
Chloride	(95 - 107 mmol/L)
Bicarbonate	(23 - 32 mmol/L)
Urea	(3 - 8.0 mmol/L)
Creatinine	(M 0.04 - 0.13, F 0.04 - 0.1 mmol/L)
eGFR	> 60 ml/min/1.72 m^2
Calcium * (total)	(2.10 - 2.60 mmol/L)
Phosphate	(0.90 - 1.35 mmol/L)
Magnesium *	(0.65 - 1.00 mmol/L)
Uric acid *	(M 0.17 - 0.45, F 0.12 - 0.40 mmol/L)
Liver function/pancreas	
Bilirubin (total) *	(< 20 µmol/L)
Bilirubin (direct) *	(< 3 µmol/L)
AST *	(<40 U/L)
GGT *	(F < 45; M < 65 U/L)
Alkaline phosphatase *	(< 120 U/L)
Total protein	(60 - 80 g/L)
Albumin	(38 - 50 g/L)
Amylase	(30 - 110 U/L)
Lipase	<80 U/L
Therapeutic drugs	
Digoxin *	(Ther. 1.3 - 2.6 nmol/L)
Phenytoin *	(Ther. 40 - 80 µmol/L)

Valproate*	(Ther. 300 - 700 µmol/L)
Carbamazepine*	(Ther. 10 - 50 µmol/L)
Gentamicin (pre)	(< 2.0 µg/mL)
Gentamicin (post)	(< 12.0 µg/mL)
Lithium	(Ther. 0.5 - 1.0 mmol/L)
Cardiac/lipids	
Troponin I or T	(< 0.1 µg/L)
AST*	(< 40 U/L)
CK (total)	(F < 200; M < 300 U/L)
CK - MB	(< 25 U/L)
Cholesterol*	(< 5.5 mmol/L)
Triglycerides*	(< 2.0 mmol/L)
HDL cholesterol	(> 1.00 mmol/L)
LDL cholesterol	(< 3.5 mmol/L)
Thyroid tests	
Free T4	(10.0 - 20.0 pmol/L)
Ultra sensitive TSH*	(0.3 - 5.0 mU/L)
Free T3	(3.3 - 8.2 pmol/L)
Other endocrine tests	
s Cortisol 8 am	(130 - 700 nmol/L)
s Cortisol 4 pm	(80 - 350 nmol/L)
FSH adult	(1.9 IU/L)
FSH ovulation	(10 - 30 IU/L)
FSH post menopausal	(4 - 200 IU/L)
Oestradiol menopausal	(< 200 pmol/L)
Testosterone	(M 10 - 35 F < 3.5 nmol/L)
Tumour markers	
PSA	(0 - 4.0 µg/L)
CEA	(< 7.5 µg/L)
AFP	(< 10 µg/mL)
CA-125	(< 35 U/mL)
Iron studies	
Ferritin	(20 - 250 µg/L)
Iron	(14 - 30 µmol/L)
Iron-binding capacity	(45 - 80 µmol/L)
Transferrin	2 - 3.5 g/L
Transferrin saturation	F20 - 55%
	M 20 - 60%
Blood gases/arterial	

PH*	(7.38 - 7.43)
PO_2*	(85 - 105 mmHg)
PCO_2*	(36 - 44 mmHg)
Bicarbonate*	(20 - 28 mmol/L)
Base excess*	(-3 to +3 mmol/L)
Glucose	
Glucose (fasting)	(3.5 - 6.0 mmol/L)
Glucose (random)	(3.5 - 9.0 mmol/L)
HbAîc	(4.7 - 6.1%)
Haematology	
Hb*	(F 115 - 165 M 130 - 180 g/L)
PCV*	(F 37 - 47 M 40 - 54%)
MCV*	(81 - 98 fL)
Reticulocytes	(0.5 - 2.0%)
Leucocytes*	(4.0 - 11.0 × 10^9/L)
Platelets	(150 - 400 × 10^9/L)
ESR	(< 20 mm)
Band neutrophils*	0.05 × 10^9/L
Mature neutrophils*	2.0 - 7.5 × 10^9/L
Lymphocytes*	1.0 - 4.0 × 10^9/L
Monocytes*	0.2 - 0.8 × 10^9/L
Eosinophils*	0.0 - 0.4 × 10^9/L
Coagulation	
Bleeding time	(2.0 - 8.5 min.)
Fibrinogen	(2.0 - 4.0 g/L)
Prothrombin time	seconds
Prothrombin ratio (INR)	1.0 - 1.2
APTT	25 - 35 seconds
D dimer	< 500 ng/ml
Others	
s Creatine (phospho) kinase	(< 90 U/L)
s Lead	(2 μmol/L)
s C-reactive protein	(< 10 mg/L)

REFERENCES

1. Levin M, Sikaris K. CRP and ESR. Aust Fam Physician, 2000; 29(10): 976 - 7.
2. Shaw AC. Serum C-reactive protein and neopterin concentration in patients with viral and bacteraemic infection. J Clin Pathol, 1991; 23: 596 - 9.
3. McPherson J, ed. *Manual of Use and Interpretation of Pathology Tests* (2nd edn). Sydney: RCPA, 1997.

4. Barratt M, Smith D. *Pathology Handbook*. Sydney: Barratt & Smith Pathologists, 1997.
5. Dunstan R, ed. *Abnormal Laboratory Tests*. Sydney: McGraw-Hill, 2001: 111 - 13.
6. Nicol D, McPhee SJ et al. *Pocket Guide to Diagnostic Tests*. Sydney: McGraw-Hill, 2003: 328 - 97.

CHAPTER 17. INSPECTION AS A CLINICAL SKILL

More mistakes, many more, are made by not looking than by not knowing.

SIR WILLIAM JENNER (1815 - 98)

INSPECTION AS A CLINICAL SKILL

GPs have an ideal opportunity to practise the art of careful observation and to notice all the signs and features characteristic of a patient from the time seen in the waiting room until the physical examination. We should be ‘like Sherlock Holmes’ in our analysis of the patient and accept the challenge of being astute diagnosticians and proud members of a noble profession.

It is important to stand back (so to speak) and look at the patient's general appearance and demeanour. We should be assessing their mood and affect as much as their physical appearance. The first assessment to make is ‘Does the patient look sick?’.

FIRST IMPRESSIONS

The first impression of the patient is always striking in some way and we should discipline ourselves to be as analytical as possible.

A rapid inspection from a trained observer may be all that is necessary to allow the observer to pinpoint specific disorders, such as anaemia, hyperthyroidism, jaundice, acromegaly and alcohol abuse. Such ‘spot’ diagnosis is not justifiable unless the original signs are supported by further examination, which must be comprehensive.

The following observations should therefore be made:

- facial characteristics
- abnormalities of the head and neck
- examination of the mouth
- character and distribution of hair
- examination of the skin (in general)
- height and weight
- posture and gait
- genitalia
- examination of extremities (hands, feet, nails, etc.)

PHYSIOGNOMY

Physiognomy, from the Greek physiognomonia, meaning the judging of one's nature, is the art of judging character from the features of the face. It flourished during the Middle Ages. According to Addison[1], ‘everyone is in some degree a master of that art which is physiognomy; and naturally forms to themselves the character of a stranger from the features of the face’. In reality, all doctors use a physiognomical approach to diagnose many medical conditions although we may not be as expert at the art as we should be.

The face is a person's most immediate means of communicating with others; it is a shield and banner, a mask and a mirror. It reveals mental faculties and emotional turmoil. It is the first perspective gained of patients as they walk into the consulting room.

THE FACE AS A MIRROR OF DISEASE

A fascinating aspect of the art of clinical medicine is the clinical interpretation of the patient's facies. Not only are specific skin lesions common on the face but the face may also mirror endocrine disorders and organ failure such as respiratory, cardiac, renal and liver failure.

Jaundice may be masked by the natural col-our of the cheeks but the yellow conjunctivae will be distinctive. A marked plethoric complexion may be seen in chronic alcoholics (alcohol may produce a pseudo-Cushing's syndrome), in Cushing's disease or in poly-cythaemia. Thickening of the sub-cutaneous tissues may be seen in chronic alcoholism, acromegaly and myxoedema, and the puffiness of the eyelids in the latter condition may simulate the true subcutaneous oedema of renal disease.

An individual's personality and mood rarely fail to leave an impression on the facial characteristics. This is partly due to the alteration in facial lines and wrinkles, which may become modified in anger, irritability, anxiety and stress. More profound changes occur with mental disease. Various CNS diseases, such as Parkinson's disease and myopathies, can affect facial expression (e.g. the immobile face of the patient with Parkinson's disease).

The appearance of the eyes can also be very significant and may reflect underlying systemic disease.

DIAGNOSTIC FACIES

Acromegalic

The enlarged characteristic face is due to a large supraorbital ridge that causes frontal bossing, a broad nose and a prominent broad and square lower jaw. Other features include an enlarged tongue and soft tissue swelling of the nose, lips and ears.

Adenoid facies

Due to mouth breathing in children: a narrow nose/nares, a high arched palate (the “Gothic” Palate), prominent incisor teeth, under, shot jaw with a perpetually open mouth and “stupid” expression.

Alcoholic (due to chronic use)

It is important to recognise the characteristic changes as early as possible - a plethoric face, thickened ‘greasy’ skin, telangiectasia, suffused conjunctivae and rosacea. Other features may include rhinophyma, parotid swelling and characteristic changes to the lips and corners of the mouth.

Bird-like (systemic sclerosis: CREST syndrome)

The bird-like features, beaking of the nose, limitation of mouth opening, puckering or furrowing of the lips and a fixed facial expression, are due to binding down of facial skin. Other features include telangiectasia on the face and hands.

Chipmunk (thalassaemia major)

There is bossing of the skull, hypertrophy of the maxillae (which tends to expose the upper teeth), prominent malar eminences and depression of the bridge of the nose. The major haemoglobinopathies cause hyperplasia of the skull and facial bones because of an increase in the bone marrow cavity.

Choleric facies

The patient with chdera has a pale face with cold dammes skin, sunken eyes, follow cheeks and a forelorn apathetic look (similar to the Hippocratic facies).

Cushingoid

The face has a typical ‘moon shape’, plethora, hirsutism (more obvious in women), acne.

Facial nerve palsy

Features include unilateral drooping of the corner of the mouth and flattening of the nasolabial fold.

- Upper motor neurone (UMN) type: the forehead movement is spared.
- Lower motor neurone (LMN) type (e.g. Bell's palsy, Ramsay-Hunt syndrome): lack of forehead muscle tone.

Hippocratic

This describes the deathly, mask-like features of advanced peritonitis - sunken eyes, ‘gaunt’ face, ‘collapsed’ temples, dry crusty lips and clammy forehead.

Obese

The distinguishing feature from the ‘moon face’ of Cushing's disease is the general roundness and uniform fatness of the face.

Thyrotoxic (hyperthyroidism)

The prominent eyes (sclera may not be covered by the lower eyelid) and conjunctivitis are features of the thyrotoxic patient. The thyroid stare (a frightened expression) may also be present.

Myxoedemic (hypothyroidism)

The face usually has an apathetic look and is ‘puffy’ with possible periorbital oedema. There is broadening of the lower part of the face. The skin (not the sclera) may appear yellow (due to hypercarotenaemia) and is generally dry and coarse. Other features may include thin, coarse, listless hair and loss or thinning of the outer third of the eyebrows. The tongue is usually enlarged and the patient speaks with a ‘thickened’, croaking, slow speech.

Marfanoid (Marfan syndrome)

The typical tall stature, arachnodactyly and chest deformities, combined with the facial features of a subluxation of the lens of the eye and high arched palate, help to pinpoint the diagnosis.

Mongoloid (Down syndrome)

The facial features include a flat profile, with crowded features, a round head, dysplastic lowset ears, protruding tongue, mongoloid slant of the eyes with epicanthic folds, mouth hanging open and peripheral silver iris spots (Brushfield's spots).

Mitral (mitral valve disease, especially mitral stenosis)

This is typically shown in flushed or rosy cheeks with a bluish tinge due to dilatation of the malar capillaries. It is associated with pulmonary hypertension.

Myotonic (dystrophia myotonia)

Typical features include frontal baldness, expressionless triangular facies, partial ptosis, cataracts and temporal muscle atrophy.

Myopathic (myopathy/myasthenia gravis)

Facial characteristics include an expressionless, ‘tired’-looking face with bilateral ptosis.

Pagetic (Paget's disease)

The main feature is skull enlargement, notably of the frontal and parietal areas (the head circumference is usually greater than 55 cm, which is abnormal) - the ‘hat doesn’t fit any more' hallmark. Other features include increased bony warmth and deafness.

Parkinsonian

Characteristic is the mask-like facies with lack of facial expression and fixed unblinking stare. There is immobility of the facial muscles.

Peutz-Jegher's

Pigmented macules (1 - 5 mm in diameter) occur on lips, buccal mucosa and fingers.

Smoker's facies

A face older than the years with premature gross wrinkling of the skin, stained teeth, deep raspy voice, 'loose' cough, smell of tobacco.

Turner's syndrome

The facial characteristics include ptosis - 'fishlike' mouth, small chin (micrognathia), low-set ears and deafness. Cardiac lesions include coarctation of the aorta and pulmonary stenosis. Webbing of the neck is the classic sign.

Uraemic facies

A sallow 'muddy' complexion will uraemic fetor - an ammoniacal halitosis.

SPECIFIC CHARACTERISTICS

Various facial signs may be present. The causes of these signs are listed below.

Butterfly 'rash'

- SLE
  Erythema, scaling with a discrete red advancing edge on the cheeks and bridge of the nose. The sharp border, lack of pustules and adherent scale make it differ from rosacea
- Rosacea
  Papules, pustules and telangiectasia on an erythematous background on cheeks, forehead and chin
- Erysipelas
  Painful, erythematous, indurated skin infection with a well-defined raised edge
- Seborrhoeic dermatitis
  Red and scaly rash involving eyebrows, eyelids, nasolabial folds
- Photosensitivity eruptions
  Erythematous on areas that are exposed to the sun

Chloasma/melasma

Increased browning pigmentation, usually confined to symmetrical areas of the cheeks. Caused by drugs:

- combined oral contraceptive pill
- hydroxychloroquine (Plaquenil)
- diphenylhydrazine

Malar flush

- Mitral stenosis
- Pulmonary stenosis
- Rosacea

- SLE
- Mesenteric adenitis

Spider naevi

- Pregnancy
- Liver disease
- Vitamin B deficiency in normal people

Enlarged tongue

- Acromegaly
- Hypothyroidism
- Amyloidosis

Cataracts

- Senility
- Corticosteroid therapy
- Diabetes
- Hypoparathyroidism
- Dystrophia myotonia
- Trauma
- Ocular disease (e.g. glaucoma)

Telangiectasia

- Systemic sclerosis
- CREST syndrome
- Liver disease (e.g. alcoholism)

CYANOSIS

Cyanosis is a bluish discolouration of the skin and mucous membranes due to deoxygenated haemoglobin concentrated in the superficial blood vessels. The arterial oxygen saturation is 80 - 85% before it is clinically apparent. It is classified as central or peripheral.

Central

Cyanosis is present in parts of the body with good circulation, such as the lips and tongue. The areas feel warm. The main causes are pulmonary disease, pulmonary oedema, cyanotic congenital heart disease (right to left shunt), respiratory depression, polycythaemia.

Peripheral

Cyanosis is in the extremities, such as the outer surface of the lips, nose and ears. The areas feel cold. The main causes are peripheral vascular disease, cardiac failure, ‘shock’, exposure to cold, left ventricular failure and all causes of central cyanosis.

CLUBBING OF FINGERS

Features

- Loss of usual angle between base of nail and nail fold.
- Curvature in two planes.
- Increased sponginess in base of nail.
- Increased convexity of nail.
- Mainly caused by respiratory disease.

Causes

1. Lung disease
    - carcinoma
    - bronchiectasis
    - cystic fibrosis
    - abscess/empyema
    - pulmonary fibrosis
2. Heart disease
    - bacterial endocarditis
    - cyanotic congenital heart disease
3. Liver disease
    - cirrhosis
4. Gastrointestinal disease
    - ulcerative colitis
    - Crohn's disease
    - coeliac disease
5. Congenital

INCREASED PIGMENTATION

Increased pigmentation is not common but if obvious in areas exposed to the sun look for ‘hidden’ areas, such as the inner aspect of the forearms. Causes include those listed below.

Increased melanocyte-stimulating hormone (MSH)

- Addison's syndrome
- Cushing's syndrome
- Ectopic ACTH syndrome

Metabolic

- Hyperthyroidism
- Haemochromatosis
- Cirrhosis of the liver
- Porphyria
- Chronic renal failure
- Malnutrition/malabsorption
- Pregnancy

Drugs

- Amiodarone
- Antibiotics (busulphan, bleomycin, minocycline)
- Antimalarials (chloroquine/hydroxychloroquine)
- Arsenic, gold, silver
- Chemotherapy
- Dapsone
- Oral contraceptive pill (OCP)
- Phenothiazines
- Photochemotherapy (PUVA)
- Psoralens
- Thiazides

Tumours

- Lymphomas
- Acanthosis nigricans
- Metastatic melanoma

REFERENCES

1. Addison T. A Collection of the Published Writings of Thomas Addison MD. New Sydenham Soc, 1818.

CHAPTER 18. A SAFE DIAGNOSTIC STRATEGY

For most diagnoses all that is needed is an ounce of knowledge, an ounce of intelligence, and a pound of thoroughness.

ANON, LANCET 1951

A SAFE DIAGNOSTIC STRATEGY

The discipline of general practice is probably the most difficult, complex and challenging of all the healing arts. Our field of endeavour is at the very front line of medicine and as practitioners we shoulder the responsibility of the early diagnosis of very serious, perhaps life-threatening, illness in addition to the recognition of anxiety traits in our patients.

The teaching of our craft is also an exciting challenge and presupposes that we have a profound comprehension of our discipline.

Our area is characterised by a wide kaleidoscope of presenting problems, often foreign to the classic textbook presentation and sometimes embellished by a ‘shopping list’ of seemingly unconnected problems or vague symptoms - the so-called undifferentiated illness syndrome.[1] Common undifferentiated symptoms include tiredness or fatigue, sleeping problems, anxiety and stress, dizziness, headache, indigestion, anorexia and nausea, sexual dysfunction, weight loss, loss of interest, flatulence, abdominal discomfort and chest discomfort.[2] It is important, especially in a busy practice, to adopt a fail-safe strategy to analyse such presenting problems. Such an approach is even more important in a world of increasing medical litigation and specialisation.

To help bring order to the jungle of general practice problems the author has developed a simple model to facilitate diagnosis and reduce the margin of error.

THE CONCEPT OF DIAGNOSTIC TRIADS

A most useful guide to learning or memorising diagnoses, especially of elusive and uncommon conditions, is to remember three key points to the condition. The cognitive process of learning these so-called ‘triads’ provides a useful template for the diagnostic methodology required in general practice. Some simple examples are shown below.

Examples of diagnostic triads

DxT: angina + dyspnoea + blackouts ° aortic stenosis

DxT: menstrual dysfunction + obesity + hirsutism ° polycystic ovary syndrome

DxT: malaise + night sweats + pruritus ° Hodgkin's lymphoma

DxT: abdominal pain + diarrhoea + fever ° Crohn's disease

DxT: vertigo + vomiting + tinnitus ° Meniere's syndrome

DxT: dizziness + hearing loss + tinnitus ° acoustic neuroma

DxT: fatigue + muscle weakness + cramps ° hypokalaemia

Examples such as these are interspersed throughout the text, especially in this chapter, and are prefixed by the symbol DxT.

THE BASIC MODEL

The use of the diagnostic model requires a disciplined approach to the problem with the medical practitioner quickly answering five self-posed questions. The questions are shown in Table 18.1.

Table 18.1 The diagnostic model for a presenting problem

1. What is the probability diagnosis?
2. What serious disorders must not be missed?
3. What conditions are often missed (the pitfalls)?
4. Could this patient have one of the 'masquerades' in medical practice?
5. Is this patient trying to tell me something else?

This approach, which is based on considerable experience, requires the learning of a predetermined plan which, naturally, would vary in different parts of the world but would have a certain universal application in the so-called developed world.

An excellent acronym on this theme 'PROMPT' was devised by a reader, Dr. Kelly Teagle:

- **P** probability
- **R** red flag
- **O** often missed
- **M** masquerades
- **P** patient wants to
- **T** tell me something

Another contribution by Flinders university medical student, fudals;

Things are not always cut and dried;

- **C** connective two disorders
- **U** UTI's particularly in v old and v young
- **T** thyroid disease *AND*
- **D** depression
- **R** remember to Rule out serious and Rate causes
- **I** iatrogenic causes
- **E** emotional needs
- **D** diabetics

Each of the above five questions will be expanded.

1 The probability diagnosis

The probability diagnosis is based on the doctor's perspective and experience with regard to prevalence, incidence and the natural history of disease. GPs acquire first-hand epidemiological knowledge about the patterns of illness apparent in individuals and in the community, which enables them to view illness from a perspective that is not available to doctors in any other disciplines. Thus, during the medical interview, the doctor not only is gathering information, allocating priorities and making hypotheses, but also is developing a probability diagnosis based on acquired epidemiological knowledge.

2 What serious disorders must not be missed?

While epidemiological knowledge is a great asset to the general practitioner, it can be a disadvantage in that he or she is so familiar with what is common that the all-important rare cause of a presenting symptom may be overlooked. On the other hand, the doctor in the specialist clinic, where a different spectrum of disease is encountered, is more likely to focus on the rare at the expense of the common cause. However, it is vital, especially working in the modern framework of a litigation-conscious society, not to miss serious, life-threatening disorders.

To achieve early recognition of serious illness the GP needs to develop a ‘high index of suspicion’. This is generally regarded as largely intuitive, but this is probably not so, and it would be more accurate to say that it comes with experience.

The serious disorders that should always be considered ‘until proven otherwise’ include malignant disease, acquired immunodeficiency syndrome (AIDS), coronary disease and life-threatening infections such as meningitis, Haemophilus influenza b infections, septicaemia and infective endocarditis (Table 18.2).

Table 18.2 Serious ‘not to be missed’ conditions

Neoplasia, especially malignancy
HIV infection/AIDS
Asthma
Severe infections, especially:
• meningoencephalitis
• septicaemia
• meningococcal infection(Figure 18.1)
• epiglottitis
• infective endocarditis
Coronary disease
• myocardial infarction
• unstable angina
• arrhythmias
Imminent or potential suicide
Intracerebral lesions (e.g. SAH)
Ectopic pregnancy



Figure 18.1 Meningococcal infection: complications of infarction (DIC) including gangrene from meningococcaemia

Myocardial infarction or ischaemia is extremely important to consider because it is so potentially lethal and at times can be overlooked by the busy practitioner. It does not always manifest as the classic presentation of crushing central pain but can present as pain of varying severity and quality in a wide variety of sites. These sites include the jaw, neck, arm, epigastrium and interscapular region. Coronary artery disease may manifest as life-threatening arrhythmias that may present as palpitations and/or dizziness. A high index of suspicion is necessary to diagnose arrhythmias.

Diagnostic triads for life-threatening conditions (examples)

DxT: fever + rigors + hypotension ° septicaemia

DxT: fever + vomiting + headache ° meningitis

DxT: fatigue + dizziness ± syncope ° cardiac arrhythmia

DxT: fever + drooling + stridor (child) ° epiglottitis

DxT: headache + vomiting + altered consciousness ° subarachnoid haemorrhage (SAH)

DxT: abdominal pain + amenorrhoea + abnormal vaginal bleeding ° ectopic pregnancy

Danger: think VIC

- Vascular
- Infection (severe)
- Cancer

3. What conditions are often missed?

This question refers to the common ‘pitfalls’ so often encountered in general practice. This area is definitely related to the experience factor and includes rather simple non-life-threatening problems that can be so easily overlooked unless doctors are prepared to include them in their diagnostic framework.

Classic examples include smoking or dental caries as a cause of abdominal pain; allergies to a whole variety of unsuspected everyday contacts; foreign bodies; occupational or environmental hazards as a cause of headache, respiratory discomfort or malaise; and faecal impaction as a cause of diarrhoea. We have all experienced the ‘red face syndrome’ from a urinary tract infection, whether it is the cause of fever in a child, lumbar pain in a pregnant woman or malaise in an older person. The dermatomal pain pattern caused by herpes zoster prior to the eruption of the rash (or if only a few sparse vesicles erupt) is a real trap.

Haemochromatosis can be a surprise diagnosis often by serendipity following a screening blood test for unexplained fatigue. Coeliac disease is a classic master of disguise in both children and adults. Research by derma-tologists[3] has highlighted that it can present in a number of ways that can affect the skin and hair. Apart from typical gastrointestinal symp-toms, such as chronic diarrhoea, steatorrhoea, weight loss, anorexia, and abdominal dis-tension, the following atypical symptoms have been described:

- nutritional presentations, including folate, zinc or iron (in particular) deficiency
- grouped blisters around the knees, elbows and buttocks (dermatitis herpetiformis)
- hair loss and mouth ulcers

Menopausal symptoms can also be over-looked as we focus on a particular symptom. Some important pitfalls are given in Table 18.3.

Table 18.3 Classic pitfalls

Abscess (hidden)
Allergies
Candida infection
Chronic fatigue syndrome
Coeliac disease
Domestic abuse, including child abuse
Drugs (see Table 18.4)
Herpes zoster
Faecal impaction
Foreign bodies
Giardiasis
Haemochromatosis
Lead poisoning
Malnutrition (unsuspected)
Menopause syndrome
Migraine (atypical variants)
Paget's disease
Pregnancy (early)
Sarcoidosis
Seizure disorders
Tourette's syndrome
Urinary infection

Table 18.4 The seven primary masquerades

1 Depression
2 Diabetes mellitus

3 Drugs
- Iatrogenic
- self-abuse
  - alcohol
  - narcotics
  - nicotine
  - others

4 Anaemia

5 Thyroid and other endocrine disorders
- hyperthyroidism
- hypothyroidism

6 Spinal dysfunction

7 Urinary tract infection (UTI)

Diagnostic triads for some ‘pitfalls’

DxT: fatigue + weight loss + diarrhoea ° coeliac disease

DxT: anorexia/nausea + faecal leaking + abdominal bloating ° faecal impaction

DxT: abdominal cramps + flatulence + profuse diarrhoea ° giardiasis

DxT: lethargy + tiredness + arthralgia ° haemochromatosis

DxT: lethargy + abdominal pains + irritability (in child) ° lead poisoning

DxT: aching bones + wadling gait + deafness ° Paget's disease

DxT: malaise + cough + fever (± erythema nodosum) ° sarcoidosis

DxT: (male child) snorting, blinking + oral noises (e.g. grunts) ± loud expletives ° Tourette's syndrome

4 The masquerades

It is important to utilise a type of fail-safe mechanism to avoid missing the diagnosis of these disorders. Some practitioners refer to consultations that make their ‘head spin’ in confusion and bewilderment, with patients presenting with a ‘shopping list’ of problems. It is in these patients that a checklist is useful. Consider the apparently neurotic patient who presents with headache, lethargy, tiredness, constipation, anorexia, indigestion, shortness of breath on exertion, pruritus, flatulence, sore tongue and backache. In such a patient we must consider a diagnosis that links all these symptoms, especially if the physical examination is inconclusive; this includes iron deficiency anaemia, depression, diabetes mellitus, hypothyroidism (Figure 18.2) and drug abuse.



Figure 18.2 Hypothyroidism in a 60 year old woman, a classic masquerade, with a slow subtle onset of facial changes

A century ago it was important to consider diseases such as syphilis and tuberculosis as the great common masquerades, but these infections have been replaced by iatrogenesis, malignant disease, alcoholism, endocrine disorders and the various manifestations of atherosclerosis, particularly coronary in-sufficiency and cerebrovascular insufficiency.

If the patient has pain anywhere it is possible that it could originate from the spine, so the possibility of spinal pain (radicular or referred) should be considered as the cause for various pain syndromes, such as headache, arm pain, leg pain, chest pain, pelvic pain and even abdominal pain. The author's experience is that spondylogenic pain is one of the most under-diagnosed problems in general practice.

A checklist that has been divided into two groups of seven disorders is presented in Tables 18.4 and 18.5. The first list, ‘the seven primary masquerades’, represents the more common disorders encountered in general practice; the second list includes less common masquerades although some, such as Epstein-Barr mono-nucleosis, can be very common masquerades in general practice.

Table 18.5 The seven other masquerades

1 Chronic renal failure

2 Malignant disease

- lymphomas
- lung
- caecum/colon
- kidney
- multiple myeloma
- ovary
- pancreas
- metastasis

3 HIV infection/AIDS

4 Baffling bacterial infections

- syphilis
- tuberculosis
- infective endocarditis
- the zoonoses
- *Chlamydia* infections
- atypical pneumonias (e.g. legionnaire's disease)
- others

5 Baffling viral (and protozoal) infections

- Epstein-Barr mononucleosis
- TORCH organisms (e.g. cytomegalovirus)
- hepatitis A, B, C, D, E
- mosquito-borne infections
    - malaria
    - Ross River fever
    - dengue fever
    - others

6 Neurological dilemmas

- Parkinson's disease
- Guillain-Barré syndrome
- seizure disorders
- multiple sclerosis
- myasthenia gravis
- space-occupying lesion of skull
- migraine and its variants
- others

7 Connective tissue disorders and the vasculitides

- Connective tissue disorders
    - SLE
    - systemic sclerosis
    - dermatomyositis
    - overlap syndrome
- Vasculitides
    - polyarteritis nodosa
    - giant cell arteritis/polymyalgia rheumatica
    - granulomatous disorders

Neoplasia, especially malignancy of the so-called ‘silent areas’, can be an elusive diagnostic problem. Typical examples are carcinoma of the nasopharynx and sinuses, ovary, caecum, kidney and lymphoietic tissue. Sarcoidosis is another disease that can be a real masquerade (p. 533).

SLE has been described as ‘the great pretender’.[4] The two most common symptoms are joint pain and fatigue but it is a multisystem disease which may present with involvement of any of these organ systems and may not initially be recognised as such.

As a practical diagnostic ploy, the author has both lists strategically placed on the surgery wall immediately behind the patient. The lists are rapidly perused for inspiration should the diagnosis for a particular patient prove elusive.

Diagnostic triads for some ‘masquerades’

DxT: malaise + fever + cough (± erythema nodosa) ° TB or sarcoidosis

DxT: fever + sore throat + cervical lymphadenopathy ° EB mononucleosis

DxT: fatigue + a/n/v + sallow skin ° chronic renal failure

DxT: polyuria + polydipsia + skin/orifice infections ° diabetes mellitus

DxT: FUO + cardiac murmur + embolic phenomena ° infective endocarditis

DxT: fatigue + polyarthritis + fever or skin lesions ° SLE

DxT: loin pain + haematuria + palpable loin mass ° carcinoma kidney

DxT: malaise + weight loss + cough ° lung carcinoma

DxT: fever + myalgia/headache + non-productive cough ° atypical pneumonia

DxT: malaise + night sweats + painless lymphadenopathy ° non-Hodgkin's lymphoma

DxT: arthralgia + Raynaud's phenomenon + GORD (± skin changes) ° systemic sclerosis

DxT: fatigue + headache + jaw claudication ° temporal arteritis

DxT: weakness + back pain + weight loss ° multiple myeloma

DxT: lethargy + physical/mental slowing + constipation ° hypothyroidism

Note: Diagnostic triads for neurological dilemmas are included in Chapter 33.

5. Is the patient trying to tell me something?

The doctor has to consider, especially in the case of undifferentiated illness, whether the patient has a ‘hidden agenda’ for the pres-entation.[5] Of course, the patient may be depressed (overt or masked) or may have a true anxiety state. However, a presenting symptom such as tiredness may represent a ‘ticket of entry’ to the consulting room. It may represent a plea for help in a stressed or anxious patient. We should be sensitive to patients' needs and feelings, and as listening, caring, empathetic practitioners provide the right opportunity for the patient to communicate freely.

Deep sexual anxieties and problems, poor self-esteem, and fear of malignancy or some other medical catastrophe are just some of the reasons patients present to doctors. The author has another checklist (Table 18.6) to help identify the psychosocial reasons for a patient's malaise.

Table 18.6 Underlying fears or image problems that cause stress and anxiety

1 Interpersonal conflict in the family
2 Identification with sick or deceased friends
3 Fear of malignancy
4 STIs, especially AIDS
5 Impending ‘coronary’ or ‘stroke’

6 Sexual problem
7 Drug-related problem
8 Crippling arthritis
9 Financial woes
10 Other abnormal stresses

In the author's experience of counselling patients and families the number of problems caused by interpersonal conflict is quite amazing and makes it worthwhile specifically exploring the quality of close relationships, such as those of husband-wife, mother-daughter and father-son.

Another common yet overlooked stressor is building[6] whether it is in the workplace, school, university, home or elsewhere.

It is a significant public health issue. The current fasliuon for tough, dynamic, ‘macho’ management styles has created a culture in which building can thrive. As GPs we should be more aware of the possibility that workplace building may be contributing to the stresses with which many patients present. A simple direct reaume greesseth ‘How are things at work?’ can create an opportunity to rasse the issue.

Identification and transference of illness, symptoms and death, in particular, are important areas of anxiety to consider. Patients often identify their problems with relatives, friends or public personalities who have malignant disease. Other somatoform disorders and the factitious disorders, including the fascinating Munchausen's syndrome, may be obvious or extremely complex and difficult to recognise.

A survey by researchers at Melbourne's Centre for Behavioural Research[7] has revealed that the three most feared diseases are cancer (81%), heart disease (32%) and HIV/AIDS (21%).

The bottom line is that patients are often desperately searching for security and we have an important role to play in helping them.

SOME EXAMPLES OF APPLICATION OF THE MODEL

Hiccough

Summary of diagnostic strategy model for abnormal hiccough

1. **Q. Probability diagnosis**
   A. Food and alcohol excess
   Psychogenic/functional
   Postoperative
     - gastric distension
     - phrenic nerve irritation
2. **Q. Serious disorders not to be missed**
   A. Neoplasia
     - CNS
     - neck
     - oesophagus
     - lung

   Subphrenic abscess
   Myocardial infarction/pericarditis

CNS disorders (e.g. CVA infection)
Chronic renal failure

3. **Q. Pitfalls**

   A. Alcohol excess
   Smoking
   Aerophagy
   GIT disorders

   - oesophagitis
   - peptic ulcer
   - hiatus hernia
   - cholecystitis
   - hepatomegaly

   Rarities
   Sudden temperature change
   Neck cysts and vascular abnormalities

4. **Q. Seven masquerades checklist**

A. Depression	-
Diabetes	-
Drugs	
Anaemia	-
Thyroid disorder	-
Spinal dysfunction	Possible
UTI	-

5. **Q. Is the patient trying to tell me something?**

   A. Emotional causes always to be considered.

Halitosis

Summary of diagnostic strategy model

1. **Q. Probability diagnosis**

   A. Dietary habits
   Orodental disease
   Dry mouth (e.g. on waking)
   Smoking/alcohol

2. **Q. Serious disorders not to be missed**

   A. Malignancy: lung, oropharynx, larynx, stomach, nose, leukaemia
   Pulmonary tuberculosis
   Quinsy
   Lung abscess
   Blood dyscrasias/leukaemia
   Uraemia
   Hepatic failure

3. **Q. Pitfalls**

   A. Nasal and sinus infection
   Systemic infection

Appendicitis
Bronchiectasis
Hiatus hernia
Rarities
Pharyngeal and oesophageal diverticula
Sjögren's syndrome
Scurvy

4. **Q. Seven masquerades checklist**

A. Depression	
Diabetes	acetone
Drugs	
Anaemia	-
Thyroid disorder	-
Spinal dysfunction	-
UTI	-

5. **Q. Is the patient trying to tell me something?**
   A. Possible manifestation of psychogenic disorder.

REFERENCES

1. Murtagh J. Common problems: a safe diagnostic strategy. Aust Fam Physician, 1990; 19: 733 - 42.
2. Frith J, Knowlden S. Undifferentiated illness. Med J Aust, 1992; 156: 472 - 6.
3. Nixon R. Cutaneous manifestations of coeliac disease. Australas J Dermatol 2001; 42: 136 - 8.
4. Hanrahan P. The great pretender: systemic lupus erythematosus. Aust Fam Physician, 2001; 30(7): 636 - 40.
5. Levenstein JH, McCracken EC, McWhinney IR et al. The patient-centred method. I. A model for the doctor-patient interaction in family medicine. Fam Practice, 1986; 3: 24 - 30.
6. McAvoy BR, Murtagh J. workplace building the silent epidemic' Editorial. BMJ 2003; 326; 776 - 777.
7. Borland R et al. Illnesses that Australians fear most in 1986 and 1993. Aust J Public Health, 1994; 18: 366 - 9.

CHAPTER 19. GENETIC CONDITIONS

> People love to oversimplify genetics, saying we have a ‘gene for cancer’ or a ‘gene for diabetes’. But the fact is, genes determine only so much. Identical twins are identical genomes, yet one may develop juvenile diabetes and the other typically doesn't. Understanding the role of genes should help pinpoint environmental factors… The genome is a history book showing the entire 6 billion-member human species traces back 7000 generations to a tiny founding population of some 60 000 people. Our species has only a modest amount of genetic variation - the DNA of any two humans is 99.9% identical.
>
> ERIC LANDER, THE HUMAN GENOME PROJECT, 3 JULY 2000[1]

GENETIC CONDITIONS

The family doctor has an important role to play in the exciting and rapidly expanding world of medical genetics. The role includes routine diagnosis, early detection, and community and ethical guidance. Virtually all of the three billion nucleotides of the human genome have been sequenced and the knowledge of their organisation into the known 30 000 - 35 000 functional units or genes continues to become more sophisticated.[2]

The genome project has commenced mapping out ‘single nucleotide poly-morphisms’ (SNPs) as signposts throughout the genome to assist in locating disease-associated genes and studying variations between individuals.[3] Any two unrelated individuals differ by one base per every thousand or so - these as SNPs - and it is believed that SNPs are the cause of the most common genetic disorders. If we carry the wrong set of SNPs, we can be predisposed to various diseases.

Genetic testing is now available for many common hereditary disorders, such as the HFE genes for haemochromatosis, presymptomatic DNA tests are available for the hereditary neurological disorders, such as Huntington disease, and predictive DNA testing is available for some forms of hereditary cancer, such as breast and colon cancer, and in the future for cardiovascular disease and diabetes.[4] Also in the future, pharmacogenetics, which predicts genetic-ally determined responses to pharmaceuticals, will greatly assist rational prescribing, and gene therapy is a futuristic treatment modality.

A summary of the prevalence of genetic disease is present in Table 19.1.[5]

Table 19.1 Prevalence of genetic disease (after Kingston)[5]

Type of genetic disease	Estimated prevalence per 1000 population
Single gene	
Autosomal dominant	2 - 10
Autosomal recessive	2
X-linked recessive	1 - 2
Chromosomal abnormalities	6 - 7
Common disorders with appreciable genetic component	7 - 10
Congenital malformations	20
Total	38 - 51

KEY FACTS AND CHECKPOINTS

- All people carry a small number of recessive genes, which are carried asymptomatically.
- The background risk that any couple will bear a child with a birth defect is about 4%. This risk is doubled for a first cousin (consanguineous) couple.[6]
- Although the majority of cancers are not inherited, some people carry inherited genetic mutations for certain cancers, notably breast and ovarian (linked), colorectal and others on a lesser scale, such as prostate and melanoma.
- GPs should look out for a family history of cancer, including the number of people with cancer on both sides of the family, the type of cancer and the age and onset of primary cancers.
- A GP caring for 1000 patients would expect to have 15 - 17 patients with a hereditary cancer predisposition.
- As genetic testing becomes more accessible it is prudent to be aware of the psychological consequences to patients of predictive or presymptomatic testing. Specialised genetic counselling is advisable.
- Carrier screening is now widely used for thalassaemia, Tay-Sachs disorder and cystic fibrosis.
- Prenatal screening and testing for genetic disorders is also a reality, especially for Down syndrome, foetal abnormalities and the haemoglobinopathies. Once again, careful selection, screening and counselling is important.

- Genetic services and familial cancer clinics provide an excellent service for referral, especially for genetic counselling expertise and advice about appropriate services and genetic testing.
- Pharmacogenomics and gene therapy are the future hope for targeted treatments based on a person's genetic profile.

Glossary of terms

Allele One of two different genes that occupy corresponding positions (loci) on paired (homologous) chromosomes (from the Greek allelon, of one another).

Haploid Normal state of genetics, containing one set of chromosomes (n).

Gametes Gametes (nature sex cells) are haploid cells

Diploid Normal state of somatic cells, containing two haploid sets of chromosomes (2n).

Zygote A diploid cell resulting from the fusion of a male and a female gamete.

Anuploid Chromosomal number that is not an exact multiple of the haploid set (e.g. 2n - 1 or 2n + 1).

Autosome Any of the 22 nuclear chromosomes other than the sex chromosomes.

Translocation An exchange of chromosomal material between two non-homologous chromosomes.

Phenotype The physical characteristics of a person (symptoms and signs) that reflect his or her genetic make-up and environmental influence.

Genotype The genetic make-up, that is, the information content inscribed on the DNA present in every cell of the body.

Gene The total DNA carried by the reproductive cell.

Genome All the DNA in an organism, including its genes.

Genomics Is the study of all genetic

Chromosomes The carriers on which the genes are carried from generation to generation.

Cytogenetics The study of chromosomes and their abnormalities in number, structure and function.

Dominant A trait or characteristic expressed in the offspring who are heterozygous for a particular gene.

Recessive A trait expressed in offspring who are homozygous for the gene (not heterozygous).

Heterozygous The presence of two different alleles at a particular focus in an individual.

Homozygous The presence of two identical alleles at a given focus in an individual.

Karyo type The chromosomal constitution of a cell or individual. It alto refers to a photographic or diagrammatic representation of the chromosomes from cell.

Monozygote Arising from one zygot Monozygote twins are kurun as identical twins.

Dizyotre Arising from two separate zygotic

Trisomy The presence of three copies of a chromosome instead of the normal two.

Mutation A spontaneous alteration in the nucleoside sequence of DNA.

Penetrance An expression of the frequency of appearance of a phenotype in the presence of one or more mutant alleles. If <100% pen trance is referred to as incomplete.

Multifactorial inheritance Characteristics resulting from a combination of more than one gene plus the influence of the environment.

Polymorphism Genetic characteristics with more than one common form in the population.

Oncogenes Genes with the potential to trigger cancer.

Consanguinity Where a couple shares one or more common ancestors.

First-degree relatives Mother, father, sisters, brothers, daughters and sons.

Second-degree relatives Grandparents, uncles, aunts, half-brothers, half-sisters, nieces, nephews and grandchildren.

Single nucleotide polymorphisms (SNPs) Base pairs of nucleotides that differ between individuals.

THE SPECTRUM OF GENETIC DISORDERS

The list of inherited disorders continues to grow as evidence supports intuition that certain conditions are hereditary. It is important for the family doctor to be aware of this potential. We are familiar with the more common classic disorders, such as cystic fibrosis, thalassaemia, Down syndrome and haemochromatosis, but it is incumbent on us to be conversant with the genetic basis of diseases such as cancer, particu-larly breast, ovarian and bowel cancer, in addition to the childhood chromosomal/microdeletion syndromes and various inborn errors of metabolism.

THE GENOGRAM

The genogram is a valuable pedigree chart that usually covers at least three generations of a family tree. It is a simple and disciplined means of gathering data about an individual couple or family to determine inheritance patterns. The data has to be gathered with tact and care. A useful strategy is to encourage patients to develop their own family genogram using charts as templates.

An example including the use of symbols is shown in Figure 19.1.


suicide
poor relationship
good relationship
=
patient
unaffected male
unaffected female
affected male
affected female
prospositus
(original case study)
or
dead
spontaneous abortion or stillbirth
induced abortion
carrier
=
consanguineous marriage
separated
divorced

Figure 19.1 Genogram: illustration of a family tree for an inherited disorder

SPECIFIC IMPORTANT GENETIC DISORDERS

Haemochromatosis

Hereditary haemochromatosis (HHC) is the most common serious single gene genetic disorder in our population.

It is a common condition in which the total body iron concentration is increased to 20 - 60 g (normal 4 g). The excess iron is deposited in and can damage several organs:

- liver - cirrhosis (10% develop cancer)

- pancreas - ‘bronze’ diabetes
- skin - bronze or leaden grey colour
- heart - restrictive cardiomyopathy
- pituitary - hypogonadism, impotence
- joints - arthralgia (especially hands), chrondrocalcinosis

It is usually hereditary (AR) or may be secondary to chronic haemolysis and multiple transfusions.

Note: Haemachromatosis is the genetic condition, haemosiderosis is the secondary condition.

Genetic profile[6]

Being an autosomal recessive disorder, the patient must inherit two altered (mutated) copies of the gene (see Figure 19.2 for an example). It is a problem mainly affecting Anglo-Saxons usually from middle age onwards. About 1 in 10 people are silent carriers of one mutated gene, while 1 in 200 are homozygous and are at risk of developing haemochromatosis. These people can have it to a variable extent (the penetrance factor), and some are asymptomatic while others have a serious problem. It is rare for symptoms to manifest before the third decade.


Both parents are carriers of β-thalassaemia
(i.e. thalassaemia minor)
no thalassaemia
25% chance
thalassaemia minor
50% chance
thalassaemia major
25% chance

Figure 19.2 Typical genogram of inheritance of an autosomal recessive disorder - thalassaemia

The two common identified specific mutations in the *HFE* geneare C282Y and H63D (another is S65C):

- homozygous C282Y - high risk for HHC
- homozygous H63D - unlikely to develop clinical HHC
- heterozygous C282Y and H63D - milder form of HHC

The key diagnostic sensitive markers are serum transferrin saturation and the serum ferritin level. The serum iron level is not a good indicator. An elevated ferritin level is not diagnostic of HHC but is the best serum marker of iron overload.

Symptoms

May have extreme lethargy, signs of chronic liver disease, polyuria and polydipsia, arthralgia, loss of libido and joint signs.

Signs

Look for hepatomegaly, very tanned skin, cardiac arrhythmias.

Diagnosis

- Increased serum transferrin saturation > 7090.
- Increased serum ferritin level > 250 µg/L.
- CT or MRI - increased iron deposition in liver.
- Liver biopsy (if liver function test enzymes are abnormal or ferritin > 1000 µg/L or hepatomegaly).MRI may take over from biopsy to evaluate liver status.
- Genetic studies: *HFE* gene - a C282Y and/or H63D mutation.
- Screen first-degree relatives (serum ferritin levels and serum transferrin saturation in older relatives and genetic testing in younger ones). No need to screen before adulthood.

Note: Full blood count (FBE) and erythrocyte sedimentation rate are normal.

Management

- Refer for specialist care.
- Weekly venesection 500 mL (250 mg iron) until serum iron levels are normal (may take at least 2 years), then every 3 - 4 months to keep serum ferritin level < 100 µg/L (usually 40 - 80 µg/L) and iron levels normal.
- Normal diet.
- Avoid alcohol.
- Life expectancy is normal if treated before cirrhosis or diabetes develops.

Thalassaemia

The thalassaemias, the most common human single-gene disorders in the world, are a group of hereditary disorders characterised by a defect in the synthesis of one or more of the globin chains (α or β) - there are two of each (α2, β2). This causes defective haemoglobin synthesis leading to hypochromic microcytic anaemia. α-Thalassaemia is usually seen in people of Asian origin while α-thalassaemia is seen in certain ethnic groups from the Mediterranean, the Middle East, South-East Asia and the Indian subcontinent. However, in our multicultural communities one cannot assume a person's origins. It is recommended that all women of child-bearing age be screened for thalassaemia.

DxT: pallor + jaundice + hepatosplenomegaly ° thalassaemia major

Genetics[6]

α-Thalassaemia is usually due to the deletion of one or more of the four genes for α-globin, the severity depending on the number of genes deleted: deletion of all four genes - α-thalassaemia (hydrops foetalis); of three genes - haemoglobin H disease, which results in lifelong anaemia of mild-to-moderate degree; of one or two genes - a symptomless carrier.

In β-thalassaemia, the β-chains are produced in decreased quantity rather than having large deletions. People who have two mutations (one in each β-globin gene) have β-thalassaemia major.

- β-thalassaemia minor - a single mutation (heterozygous) - the carrier or trait state
- β-thalassaemia major - two mutations (homozygous) - the person who has the disorder

Inheritance is illustrated in Figure 19.2. If both parents are carriers, there is a 1 in 4 chance that their child will have the disorder.

Clinical

Carriers are clinically asymptomatic and do not need treatment apart from counselling. Patients with thalassaemia major present with symptoms of severe anaemia (haemolytic anaemia). Without treatment, children with thalassaemia major are lethargic and inactive, show a failure to thrive or to grow normally and delayed puberty, hepatosplenomegaly and jaundice. Signs usually appear after 6 months and death from cardiac failure was common but with regular blood transfusions and iron-chelating treatment people can now live in good health.

Diagnosis

- FBE: in most carriers the mean corpuscular haemoglobin/mean corpuscular volume is low but can be normal. There is usually mild hypochromic microcytic anaemia but this is severe with the homozygous type.
- Haemoglobin electrophoresis: measures relative amounts of normal adult haemoglobin (HbA) and other variants (e.g. HbA_2, HbF). This will detect most carriers.
- Serum ferritin level: helps distinguish from iron deficiency, which has a similar blood film.
- DNA analysis: for mutation detection (mainly used to detect carriers).

Treatment for thalassaemia major

Treatment is based on a regular blood transfusion schedule for anaemia. Folate supplementation and a low iron diet are advisable. Excess iron is removed by iron chelation (e.g. deferoxamine). Allogenic bone marrow transplantation has been used with success.[7] Splenectomy may be appropriate.

Cystic fibrosis

Cystic fibrosis is the result of a defect in an ion channel protein, the cystic fibrosis transmembrane receptor, which is found in the membranes of cells lining the exocrine ducts. The defect affects the normal transport of chloride ions, leading to a decreased sodium and water transfer, thus causing viscid secretions that affect the lungs, pancreas and gut.

DxT: failure to thrive + chronic cough + loose bowel actions ° cystic fibrosis

Genetics[6]

- The most common AR paediatric illness.
- About 1 in 2500 Caucasians affected.
- About 1 in 20 - 25 are carriers.
- A mutation (δ-F508) of chromosome 7 is the most common of some 500 possible mutations of the gene. This deletes a single phenylalanine residue from a 1480-amino-acid chain.

Clinical features

- General: malaise, failure to thrive, exercise intolerance.
- Chronic respiratory problems: cough, recurrent pneumonia, bronchiectasis, sinus tenderness, nasal polyps.
- Gastrointestinal: malabsorption, pale bulky stools, jaundice (pancreatic effect), meconium ileus (10% of newborn babies).
- Infertility in males (atrophy of vas deferens).
- Pancreatic insufficiency.
- Early mortality but improving survival rates (mean age now 31 years).

Investigations

- Screening for immunoreactive trypsin in newborns detects 75%.
- Sweat test for elevated chloride and sodium levels.
- DNA testing for carriers identifies only the most common mutations (70 - 75%).

Treatment

- Early diagnosis and multidisciplinary team care are important.
- Physiotherapy for drainage of airway secretions.
- Hypertomic salive solution (by nebuliser) preceded by a bronchodilator.
- Treatment of infections: therapeutic and prophylactic antibiotics.
- Oral pancreatic enzyme replacement.
- Dietary manipulation.
- Lung and liver transplantation are considerations.

There is currently no cure for cystic fibrosis; treatment is based on correcting the nutritional deficiencies and minimising chest infections.

Neurofibromatosis

There are two types of this AD disorder:

- NF1 - peripheral neurofibromatosis (von Recklinghausen disorder)
- NF2 - central type, bilateral acoustic neuromas (rare)

The gene for NF1 is carried on chromosome 17 and NF2 on chromosome 22. Diagnostic genetic testing is not routinely available. Diagnosis is by clinical examination.

DxT: light-brown skin patches + skin tumours + axillary freckles ° NF1

Clinical features of NF1[6]

- Six or more café-au-lait spots (increasing with age).
- Freckling in the axillary or inguinal regions.
- Hypertension.
- Eye features (iris hamartomas).
- Learning difficulty.
- Musculoskeletal problems (e.g. scoliosis, fibrous dysplasia, pseudoarthrosis).
- Optic nerve gliomas.

General rule

- One-third asymptomatic, only have skin stigmata.
- One-third minor problems, mainly cosmetic.
- One-third significant problems (e.g. neurological tumours).

Management

- No special treatment available.
- Surgical excisions of neurofibromas as appropriate.
- Refer to a special clinic, including neurofibroma clinic.
- Careful surveillance - report new symptoms.
- Yearly examination for children and adults, including blood pressure, neurological, skeletal and ophthalmological examination.

Duchenne muscular dystrophy (DMD)

DMD is a progressive proximal muscle weakness disorder with replacement of muscle by connective tissue. Becker muscular dystrophy is a less severe variant.

Genetics

DMD is an X-linked recessive condition. It is caused by a mutation in the gene coding for dystrophin, a protein found inside the muscle cell membrane.

DxT: male child + gait disorder + bulky calves ° DMD

Clinical features[6]

- Usually diagnosed from2 - 5 years.
- Weakness in hip and shoulder gridles.
- Walking problems: delayed onset or starting in boys aged 3 - 7.
- Waddling gait, falls, difficulty standing and climbing stairs.
- Pseudohypertrophy of muscles, especially calves.
- Most in wheelchair by age 10 - 12.
- ± Intellectual retardation.
- Most die of respiratory problems by age 20.
- Gowers sign patient uses ‘trick’ method by using hands to climb up his or her legs when rising to an exact position from the flock.

Diagnosis

- Elevated serum creatinine kinase level
- Electromyography
- Direct dystrophin gene testing

- Muscle biopsy

Treatment

- Counselling, education, screening (especially mother).
- No specific treatment available. Corticosteroids delay progression.

INHERITED ADULT ONSET NEUROLOGICAL DISORDERS[9]

These disorders have the following common features:

- They are serious and usually eventually fatal.
- Onset is in adulthood.
- They are currently incurable.
- They affect successive generations.
- Most are multifactorial in causation.
- Most are inherited from a parent in an autosomal dominant fashion.
- Specific genetic testing is usually available.

Examples of inherited adult onset neurological conditions are:

- Huntington disease
- Creutzfeldt-Jakob disease and other prion diseases
- familial Alzheimer disease
- familial epilepsy
- familial motor neurone disease
- Friedreich ataxia
- hereditary peripheral neuropathies (Charcot Marie tooth disease)
- mitochondrial disorders
- hereditary spastic paraparesis
- muscular dystrophies
- myotonic dystrophy
- spinal muscular atrophy
- spinocerebellar ataxias

A minority of these conditions are due primarily to a dominantly inherited genetic alteration (mutation), e.g. Huntington disease, and are usually more accessible to genetic testing. Some gene alterations (polymorphism) may be associated with a higher risk of developing certain neurological conditions. Testing for polymorphisms is more complex. If concerned about a hereditary basis, GPs should refer their patients with these disorders to a neurologist or a neurogenetics clinic.

Huntington Disease

Genetic profile

- Inherited as an AD disorder.
- The responsible mutant gene has been located on the short arm of chromosome 4.
- One genetic mutation accounts for the vast majority of cases, which means that there is an accurate diagnostic test.
- Both sexes are equally affected.

DxT: chorea + abnormal behaviour + dementia ° Huntington disease

Clinical features

- Insidious onset and progression of chorea.
- Onset most often between 35 and 55 years.
- Mental changes - change in behaviour (can be as early as childhood or in very late life), intellectual deterioration leading to dementia.
- Family history present in the majority.
- Motor symptoms: flicking movements of arms, lilting gait, facial grimacing, ataxia, dystonia.
- Usually a fatal outcome 15 - 20 years from onset.

Treatment

- There is no cure or specific treatment.
- Supportive treatment with agents such as haloperidol.

Genetic testing and counselling

This is available, sensitive and important because offspring have a 1 in 2 risk and the onset may be late - after child-bearing years. It is appropriate to refer to expert centres for those seeking it. Of interest is that only 20% have undergone testing since it became available, indicating that those at risk generally prefer the uncertainty of not knowing to the reality.[9]

Familial Alzheimer disease (FAD)

Early onset familial Alzheimer's disease (EoFAD), which accounts for less than 5% of all Alzheimer's disease, is defined as the presence of two or more affected people with on set age < 65 years in more than one generation of a family with post-mortem pathologically proven Alzheimer's disease in at least one person. Mutations in any one of three different forms (alleles) of the susceptible ADOE gene are known to cause FAD.[10]

Parkinson's disease

Most cases are sporadic, and the majority of cases with a family history do not have a clear inheritance pattern and could be the result of several factors including a genetic predisposition or simply a chance aggregation. Consider referral to a neurogenetics clinic for families with unusual features, such as familial aggregation and/or early onset Parkinson's disease.

Motor neurone disease (amyotrophic lateral sclerosis)

Five to 10% of motor neurone disease is inherited, with an autosomal dominant inheritance pattern. Inherited MND shows familial aggregation and an earlier age of onset than average (40s or younger); otherwise clinical features are essentially the same as the sporadic form. If more than one family member presents with MND consider referral to a neurogenetic clinic.

The epilepsies

The epilepsies comprise a group of disorders with differing genetic components and the inheritance or genetic contribution relates to each specific disorder. Further studies on this subject are in progress. If there are two or more individuals in a family with epilepsy, referral for advice about the nature and inheritance about their form of epilepsy may be appropriate.

Psychiatric conditions

Serious psychotic and mood disorders can run in families, especially schizophrenia and bipolar disorder which have a clear genetic component that appears to be complex and poorly understood (Table 19.2). To date no genes causing schizophrenia have been identified, but large regions on some chromosomes have been associated with schizophrenia. Individuals with a first-degree relative with bipolar disorder or purely depressive (unipolar) have an increased risk of a mood disorder, but the genetics are not understood. Tourette's syndrome is another psychological disorder that has a hereditary basis through an autosomal dominant gene with variable expression (penetrance).

Table 19.2 Genetic risks (approximate) in schizophrenia and bipolar disorder

Affected relative	Schizophrenia (%risk)	Bipolar (%risk)
Nil (general population)	1	2 - 3
Parent	13	15
Both parents	45	50
Sibling	9	13
Monozygotic twin	40	70
Dizygotic twin	10	20

Source : Harper P. ***Practical Genetic Counselling*** . Oxford: Butterworth Heinemann, 1988.

HEREDITARY HAEMOGLOBINOPATHIES, HAEMOLYTIC DISORDERS, BLEEDING AND CLOTTING DISORDERS[9]

The commonest haemglobinopathies are the thalassaemias (see p. XXX), which are caused by a deficiency in the quality of globin chains whereas other haemoglobinopathies are caused by structural variations in the globin chain. These conditions include HbS (sickle cell), HbC, HbD, HbE, HbO and HbLepore.

Other inherited conditions that can cause haemolytic anaemia are those with a red cell membrane defect and include hereditary spherocytosis, hereditary elliptocytosis and hereditary stomatocytosis.

Sickle cell disorders

The most important abnormality in the haemoglobin (Hb) chain is sickle cell haemoglobin (HbS), which results from a single base mutation of adenine to thymine leading to a substitution of valine for glutamine at position 6 on the ß-globin chain. The defective Hb causes the red cells to become deformed in shape - ‘sickled’. The sickled cells tend to flow poorly and clog the microcirculation, resulting in hypoxia which compounds the sickling. Such attacks, which result in tissue infarction, are called ‘crises’. Sickling is precipitated by infection, hypoxia, dehydration, cold and acidosis, and may complicate operations. The disorder occurs mainly in Africans (25% carry the gene), but it is also found in India, South-East Asia, the Middle East and southern Europe.

- Heterozygous state for HbS ° sickle cell trait
- Homozygous state ° sickle cell anaemia/disease

Sickle cell anaemia

It varies from being mild or asymptomatic to a severe haemolytic anaemia and recurrent painful crises. It may present in children with anaemia and mild jaundice. Children may develop digits of varying lengths from the hand-and-foot syndrome due to infarcts of small bones.

Features of infarctive sickle crises include:

- bone pain (usually limb bones)
- abdominal pain
- chest-pleuritic pain
- kidney-haematuria
- spleen-painful infarcts

Hb electrophoresis is needed to confirm the diagnosis.

Long-term problems include chronic leg ulcers, susceptibility to infection, aseptic necrosis of bone (especially head of femur), blindness and chronic renal disease. The prognosis is variable. Children in Africa often die within the first year of life. Infection is the commonest cause of death.

Sickle cell trait

People with this usually have no symptoms unless they are exposed to prolonged anoxia, such as anaesthesia and flying in non-pressurised aircraft. The disorder is protective against malaria.

Hereditary spherocytosis

This is the commonest cause of inherited haemolytic anaemia in northern Europeans. It is an autosomal dominant disorder of variable severity, although in 25% of patients neither parent is affected, suggesting spontaneous mutation in some instances. Jaundice may present at birth or be delayed or occur not at all. Splenomegaly is a feature and splenectomy is considered to be the treatment of choice. Maintenance of folic acid levels is important.

Bleeding disorders

In inherited bleeding deficiency disorders, there are deficiencies of vital factors (see p. p. XXX). The common significant disorders are:

- haemophilia A (factor VIII deficiency) - X linked recessive
- haemophilia B (factor IX deficiency) - X linked recessive
- Von Willebrand's disease (deficiency of factor VIII:C + defective platelet factor) - autosomal dominant

Others to consider are:

- hereditary haemorrhagic telangiectasia (Osler-Weber-Rendu disease)
- inherited thrombocytopenia

Thrombophilia

This should be considered in patients with a past and/or family history of DVT or other thrombotic episodes (see p. XXX). There are several causes, including important inherited factors which are:

- factor V Leiden gene mutation (activated protein C resistance)
- prothrombin gene mutation

- protein C deficiency
- protein S deficiency
- antithrombin deficiency

It is important to be aware of these factors, especially in people with a past history of unexplained thrombotic episodes. Prescribing the OCP is an issue. In factor V Leiden, the most common factor in this group, there is a 35-fold increased risk of thrombosis for those taking the OCP.

CHROMOSOMAL/MICRODELETION SYNDROMES (CHILDHOOD EXPRESSION)[9]

The following disorders, whose clinical features manifest in children, present with developmental and intellectual disability.

Down syndrome[11]

Down syndrome (trisomy 21) is based on typical facial features (flat facies, slanting eyes, epicanthic folds, small ears), hypotonia, intellectual disability and a single palmar crease.

DxT: typical facies + hypotonia + single palmar crease ° Down syndrome

Facts

- 95% have extra chromosome of maternal origin (trisomy 21).
- Remainder due to either unbalances, translocations or mosaicism.
- Prenatal screening tests include early ultrasound (nuchal lucency) and maternal serum screening in first trimester. Karyotyping of chorionic villus sampling on amniocytes for pregnancies at risk is available.
- Prevalence 1 in 650 live births.

Associated disorders

- Seizures (usually later onset)
- Impaired hearing
- Leukaemia
- Hypothyroidism
- Congenital anomalies (e.g. heart, duodenal atresia, Hirschsprung's, TOF)
- Alzheimer-like dementia (fourth - fifth decade)
- Atlantoaxial instability

Management

- Assess child's capabilities.
- Refer to agencies for assessment (e.g. hearing, vision, developmental disability unit).
- Advise on sexuality, especially for females (i.e. menstrual management, contraception) as fertility must be presumed.
- Genetic counselling for parents.

Edward Syndrome

Trisomy 18

Clinical features: include

- Incidence in 2000 live births (approx.)
- Microcephaly
- Facial abnormalities of dept lip/palate
- Malformation of major organs
- Malformation hands feet - lenched hand posture.
- Neurce tabe defect

Prognosis is poor - About 1/3 die in first month, < 10% live beyond 12 months.

Prenatal diagnom available.

Paaatau syndrome[9]

Trisomy 13

Clinical features: include

- Incidence 1 in 7000(approx.)
- Microcephaly
- Brain and heart malfermation
- Cleft lip/palate
- polydactually
- Neural tube defeet

Prognosis is poor - 50% die within first month.

Fragile X syndrome (FXS)[11]

FXS presents as a classic physical phenotype with large prominent ears, long narrow face, macro-orchidism and intellectual disability. It is the most common inherited cause known of developmental disability and should always be considered. The cause is the result of an increase in the size of a trinucleotide repeat in the *FMR-I* gene on the X chromosome (the number of sequences determines carrier or full mutant status). Any individual with significant development delay should be tested for FXS.

DxT: facies + intellectual disability + large testes ° FXS

Facts

- M:F ratio 2:1.
- Prevalence 1 in 1000 - 4000.
- Variable spectrum of characteristic features, making detection difficult in some cases.
- Up to 1 in 300 females may be carriers.
- Family history of intellectual disability.
- Affects all ethnic groups.
- Females may appear normal but may be affected.

Diagnosis

- Cytogenetic testing (karyotyping)
- DNA test (specific for full mutation as well as carriers)

Associated disorders

- Intellectual disability (IQ < 70).
- Autism or autistic-like behaviour.
- Attention deficit in 10% (with or without hyperactivity).
- Seizures (20%).
- Connective tissue abnormalities.
- Learning disability and speech delay.
- Co-ordination difficulty.

Management

- Careful genetic appraisal and counselling.
- Assessment of child's capabilities.
- Multidisciplinary assessment, including developmental disability unit.
- Referral for integration of speech and language therapy, special education, behaviour management.
- Pharmacological treatment of any epilepsy, or attention or mood behaviour disorders.
- Medications may determine whether the child remains in the community or not.

Prader-Willi syndrome

This uncommon disorder (1 in 10 000 to 15 000) has classic features, especially a bizarre appetite and eating habits, of which the GP should be aware. It is probable that there are many undiagnosed cases in the community. The most common cause is a deletion of the short arm of chromosome 15.

DxT: neonatal hypotonia + failure to thrive + obesity ° Prader-Willi syndrome

Clinical features

- Usualies manifeets at 3 years.
- Hypotonic infants with weak suction and failure to thrive, then voracious appetite causing morbid obesity.
- Intellectual disability.
- Narrow forehead and turned-down mouth.
- Small hands and feet.
- Hypogonadism.

Management

- Early diagnosis and referral.
- Multidisciplinary approach.
- Expert dietetic control.

With proper care and support, longevity into the eighth decade is a reality.[11]

Williams syndrome

Williams syndrome (idiopathic hypercalcaemia or elfin face syndrome) is due to a microdeletion on chromosome 7, a deletion in the elastin gene.

DxT: 'elfin' face + intellectual disability + aortic stenosis ° Williams syndrome

The children have a distinctive elfin facial appearance, mild pre- and postnatal growth retardation, mild microcephaly and mild-to-moderate developmental delay. In the first 2 years of life feeding problems, vomiting, irritability, hyperacusis, constipation and failure to thrive may lead to presentation, but the children are rarely diagnosed at this stage.

Marfan syndrome[9]

This is a systemic connective tissue disorder characterised by abnormalities of the skeletal, cardiovascular and ocular systems. It has variable expressions and is a potentially lethal disorder. If untreated, death in the 30s and 40s is common.

DxT: tall stature + dislocated lens and myopia + aortic root dilatation ° Marfan syndrome

Genetic profile

- Mutations in the fibrillin gene on chromosome 15.
- Autosomal dominant.
- Prevalence about 5 per 100 000.
- No specific laboratory test to date.

Clinical features

- Disproportionally tall and thin.
- Long digits - arachnodactyly.
- Kyphoscoliosis.
- Joint laxity (e.g. genu recurvatum).
- Myopia and ectopic ocular lens.
- High arched palate.
- Aortic dilatation and dissection.
- Mitral valve prolapse.

Management

- Needs surveillance of eyes, heart and thoracic aorta.
- Echocardiography, possibly aortic root dilatation.
- Long-term β-blockade therapy reduces rate of dilatation.
- Consider prophylactic cardiovascular surgery.

- Genetic counselling for the family.

Tuberous sclerosis (epiloia)

This is an autosomal dominant disorder due to mutations in one of two genes located on chromosomes 9 and 16. A feature is tube-like growths that affect multiple systems including the brain.

DxT: facial rash + intellectual disability + seizures ° tuberous sclerosis

The above triad of features is classic but not applicable to all cases.

Noonan syndrome

This is an AD disorder with mutation of chromosome 11. It has been described as a male Turner's syndrome but affects both sexes.

DxT: facies + short stature + pulmonary stenosis ° Noonan's syndrome

Clinical features[11]

- Characteristic facies - down-slanting palpebral fissures, widespread eyes, low-set ears ± ptosis.
- Short stature.
- Pulmonary valve stenosis.
- Webbed neck.
- Failure to thrive, usually mild.
- Abnormalities of cardiac conduction and rhythm.
- ± Intellectual disability.

Treatment

- Refer to genetic service.
- Evaluation of cardiac status.
- Consider vision, hearing, clotting status, possible epilepsy.

SEX CHROMOSOME ABNORMALITIES

Klinefelter's syndrome

This is due to an extra X chromosome, resulting in a male phenotype and occurring in 1 in 800 live births.

DxT: lanky men + small testes + infertility ° Klinefelter's syndrome

Genetic profile

- XXY genotype.
- The extra X chromosome is usually of maternal origin.
- About 30 or more variants of the disorder.

Clinical features

Marked variation but usually:

- tall men with long limbs
- small firm testes ≤ 2 cm
- infertility (azoospermia)

There may be:

- learning difficulties
- intellectual disability
- gynaecomastia
- increased risk of breast cancer and diabetes

Treatment

- Transdermal testosterone.
- Excision of gynaecomastia.

Turner syndrome (gonadal dysgenesis)

This is due to only one X chromosome, occurring in 1 in 4000 live female newborns; 99% of conceptions are miscarried.[12]

DxT: short stature + webbed neck + facies ° Turner's syndrome

Genetic profile

- 45 chromosomes of XO karyotype (typical Turner's karyotype in 50% of cases).
- Many are mosaics (e.g. 45X/46XX chromosomes).
- Phenotypes vary.

Clinical features of typical XO karyotype

- Short stature.
- Primary amenorrhoea in XO patient (Turner variants may go through puberty and have early menopause).
- Webbing of neck.
- Typical facies.
- Lymphoedema of extremities.
- Multiple congenital abnormalities.
- Cardiac defects (e.g. coarctation of aorta)

Mental deficiency is rare.

Treatment

Treatment is hormone-based (e.g. growth hormone, hormone-replacement therapy).

FOETAL ALCOHOL SYNDROME[11]

This important syndrome is caused by the teratogenic effects of alcohol (not a chromosomal abnormality) and is estimated to involve 2 in 1000 live births. The phenotype varies with the dosage and gestational timing of the alcohol exposure.[13]

DxT: facies + growth retardation + microcephaly ° foetal alcohol syndrome

Clinical features[13]

- Markedly underweight until puberty.
- Central nervous system dysfunction and mental retardation.
- Microcephaly.
- Characteristic facies:
  - narrow frontal/forehead area
  - shortened palpebral fissures
  - low-set ears
  - mid-facial hypoplasia
  - long, smooth featureless philtrum
  - thin upper lip
  - upturned nose
  - micrognathia (in infant)
- Hyperactivity.
- Congenital heart disease often seen.
- Skeletal abnormalities.

Management

Management is by preventive strategies with community education about the harmful effects of drinking, especially in early pregnancy.

OTHER TYPES OF HEREDITARY DISORDERS

The many examples include familial hypercholesterolaemia (AD) and the AR disorders for which genetic testing is available, namely Gaucher's disease, Tay-Sachs disease, glycogen storage disease, phenylketonuria, galactosaemia, homocystinuria, the porphyrias and glucose-6-phosphate dehydrogenase (G-6-PD) deficiency.

Glucose-6-phosphate dehydrogenase deficiency

G-6-PD deficiency is a common disorder affecting over 200 million people worldwide. It is the most common red-cell enzyme defect that causes episodic haemolytic anaemia because of the decreased ability of red blood cells to cope with oxidative stresses. It is an X-linked recessive inherited disorder with a high prevalence among people of African, Medi-terranean or Asian ancestory. In some countries such as Malaysia there is a national screening program. Consider the disorder in male black people.

The important clinical features are:

- neonatal jaundice - infants at risk should be observed after delivery (at least 5 days)
- episodic acute haemolytic anaemia - triggered by antioxidants and infections, and drugs, especially antimalarials, sulph-onamides, nitrofurantoin, traditional medicines, vitamms candk, high dose aspirin, fava beans and naphthalene (e.g. moth balls).

There is no specific treatment. Known precipitants should be avoided. Avoid penicillin and probenecid.

Gaucher's disease

Gaucher's disease, which is due to a deficiency of the lysosomal enzyme glucocerebrosidase, leads to anaemia and thrombocytopenia as a result primarily of hypersplenism. There is chronic bone pain and ‘crises’ of bone pain. Consider it in children with fatigue, bone pain, delayed growth, epistaxis, easy bruising and hepatosplenomegaly. Replacement enzyme therapy is available.

Phenylketonuria (PKU)

This autosomal recessive disorder of the catabolism of the amino acid, phenylalanine, is caused by a deficiency of phenylalanine hydroxylase activity, leading to an elevation of plasma phenylalanine, which if untreated can cause intellectual disability (often very severe) and other neurological symptoms, such as seizures. Neonatal screening for high blood phenylalanine levels (the Guthrie test) is performed routinely.

Treatment aims to limit phenylalanine intake so that essential amino acid needs are met but not exceeded. Diet therapy must commence as soon as possible. Females who have been treated for PKU need pre-pregnancy counselling and dietary management during pregnancy to prevent damage to the foetus by high phenylalanine levels.

The porphyrias

The three most common porphyrias are acute intermittent porphyria, porphyria cutanea tarda (the commonest) and erythropoietic protoporphyria, which are caused by deficiencies of the third, fifth and eighth enzymes, respectively, of the heme biosynthesis pathway. Their clinical features are quite different.

Acute intermittent porphyria

This autosomal dominant disorder is the most serious of the porphyrias although it remains clinically silent in the majority of patients who carry the trait. It is due to a deficiency of porphobilinogen (PBG) deaminase.

DxT: severe abdominal pain + abnormal illness behaviour + ‘red’ urine ° acute intermittent porphyria

Clinical features

- Unexplained abdominal pain crises.
- Usually young women (teens or 20s).
- Recurrent psychiatric illnesses.
- Acute peripheral or nervous system dysfunction (e.g. peripheral neuropathy).
- PBG in urine during attack.
- Hyponatraemia.
- Attacks precipitated by various drugs (e.g. antiepileptics, alcohol, sulphonamides, barbiturates).

Diagnosis

- Urine PBGs (high) and serum sodium (very low) during ‘attack’.
- Erythrocyte PBG deaminase testing to screen relatives.

Treatment

- Avoid ‘unsafe’ drugs.
- High carbohydrate diet.

Glycogen storage disease (liver glycogenoses)

This is a group of inherited disorders caused by a deficiency of one or more enzymes involved in glycogen breakdown, leading to the deposition of abnormal amounts of glycogen in tissues, especially the liver. The best-known type is 1A (von Gierke's disorder), an autosomal recessive disorder due to deficiency of glucose-6-phosphatase (G-6-P). It is seen in several ethnic groups. It typically causes growth retardation, hepatomegaly, renomegaly, hypoglycaemia (can be severe), lactic acidosis and hyperlipidaemia. Children have characteristic morphological features - short, doll-like facies with fat cheeks, thin extremities and large abdomen (hepatomegaly).

Diagnosis is by abnormal plasma lactate and lipid levels, liver biopsy and recently by gene analysis for the G-6-P gene.

Treatment is aimed to prevent hypo-glycaemia and lactic acidosis via frequent carbohydrate feedings, such as uncooked cornstarch and overnight nasogastric glucose infusion. The prognosis is poor.

Tay-Sachs disease

About 1 in 25 Ashkenazi Jews is a carrier of Tay-Sachs disease (gangliosidosis), an AR disorder caused by a total deficiency of hexosaminidase A resulting in an accumulation of gangliosides in the brain.

The infantile form is fatal by age 3 or 4 with early progressive loss of motor skills, dementia, blindness, macrocephaly and cherry-red retinal spots. The juvenile onset form presents with dementia and ataxia, with death at age 10 - 15. The adult form has progression of neurological symptoms following clumsiness in childhood and motor weakness in adolescence. Carrier testing is available and prenatal diagnosis is available.

FAMILIAL CANCER

The majority of cancer is not inherited, rather it is acquired because of genetic mutations in several genes of a cell in a specific tissue during an individual's lifetime. 20 - 25 individuals in a population of 1000 a family history of bowl or breast cancer.[9]

However, some people carry inherited genetic mutations from conception that predispose them to developing certain cancers, particularly colorectal, breast and ovarian cancer, at a relatively young age. Up to 5% of some cancers are considered familial and the genetic basis is now understood. Most are AD with 50% of offspring being affected.[14]

The three most significant familial cancer syndromes are:

- hereditary breast - ovarian cancer syndrome (BRCA1 and BRCA2 genes)
- hereditary non-polyposis colorectal cancer (HNPCC)
- familial adenomatous polyposis (FAP)

Features of breast - ovarian cancer syndrome[6]

- Mutations in either of the two genes - BRCA1 and BRCA2 - result in a strong predisposition for both breast and ovarian cancer.
- Mutations present in about 1 in 800 of the general population (male and female), who are carriers.
- Dominant inheritance.
- The risk of developing breast cancer is 10-fold and 40 - 80% of cases occur before the age of 70 years.[14]
- The prognosis in these women is the same as for sporadic cases.
- Early age of onset of breast cancer.
- Male breast cancer (6% in males with BRCA2 gene mutation).
- Coexistence of ovarian and breast cancer in the same family.
- Carriers of mutations may be at an increased risk of prostate cancer, pancreatic cancer and colorectal cancer, although this is controversial for the latter two.

Risk indicators for familial breast - ovarian cancer

- Two first-degree or second-degree relatives on one side of the family with cancer.

- Individuals with age of onset of cancer < 50 years.
- Individuals with bilateral or multifocal breast cancer.
- Individuals with ovarian cancer.
- Breast cancer in a male relative.
- Jewish ancestry.

Colorectal cancer[14]

Both sexes have a risk of approximately 5% of developing bowel cancer in their lifetime. In some this risk is increased due to an inherited predisposition.

The two key disorders are HNPCC and FAP.

Hereditary non-polyposis colorectal cancer (Lynch syndrome)

- Caused by a defect in one of the genes responsible for DNA mismatch repair.
- Affects 1 in 1000 individuals.
- Autosomal dominant.
- Early age of onset.
- Increased risk of certain extracolonic cancers, including endometrial, stomach, ovary and renal tract cancers.

Familial adenomatous polyposis

- Less common than HNPCC; affects about 1 in 10 000.
- Caused by a mutation in the *APC* gene.
- Usually hundreds or thousands of polyps.
- Eventually almost 100% of cases develop colon cancer without prophylactic colectomy.
- Median age of diagnosis 40 years.
- Small increased risk of other cancers (e.g. thyroid, cerebral).

Individuals at risk

For HNPCC:

- Three or more close relatives with bowel cancer.
- Two or more close relatives with bowel cancer and:
    - more then one bowel cancer in same relative.
    - onset of bowel cancer before 50 years
    - a relative with endometrial cancer or ovarian cancer

For FAP:

- A relative with bowel cancer with polyposis.
- Individuals with multiple polyposis.

The rÍc nole of the GP in familial cancer[14]

The GP has an important role in identifying potential high-risk patients and families and in addressing their concerns.

- Take a family history and involve at least three generations.
- Map a family tree: include any diagnosed breast, ovarian or colorectal cancers in any relative and any type of cancer in first-degree or second-degree relatives on either side of the family.
- Record the age of onset and site of cancer in first-degree relatives.
- Confirm reports of cancer from medical records.
- Assess risk (high, low or intermediate) using guidelines from your Anti-Cancer Council.
- Reassure low-risk patients but provide general preventive and screening guidelines.
- Refer all patients at potentially high risk to a familial cancer clinic.

Services at these clinics involve:

- risk assessment
- genetic testing
- counselling, including pre- and post-testing
- surveillance advice

Management is based on early detection and potential prophylactic methods, for example:

- for breast cancer - regular imaging and clinical examination
- for ovarian cancer - transvaginal ultrasound and serum CA-125 detection
- for FAP and HNPCC - annual colonoscopy, faecal occult blood test

Patients at high risk will ask about prophylactic colectomy, mastectomy and oophorectomy, for which there are reasonable indications but it is necessary to refer to a cancer geneticist for expert evaluation before such decisions are made.

PREDICTIVE GENETIC TESTING

The Human Genetics Society of Australia strongly advises against predictive or presymptomatic genetic testing of children for disease where there is no pre-emptive treatment in childhood. It should only be carried out in children if an effective treatment or preventive strategy is available. It raises issues of confidentiality, informed consent and harmful effects on self-esteem.[9]

Routine genetic testing is only advisable for high-risk individuals, such as those with a family history.

The ethical issues for adults are also considerable and for the individual the decision is difficult and requires considerable counselling via a clinical genetics service. This applies particularly to those at risk of Huntington syndrome and other adult-onset neurodegenerative conditions for which no preventive treatment exists.

PRENATAL SCREENING AND DIAGNOSIS OF GENETIC DISORDERS

Approximately 2% of births are associated with congenital abnormalities, of which 1 in 7 are chromosomal, the most common of which is Down syndrome (trisomy 21). Antenatal screening tests that can now be performed for several conditions are mainly:

- screening tests for Down syndrome and other trisomies
- screening tests for thalassaemias/haemoglobinopathies
- second-trimester ultrasound scans for foetal abnormalities, such as neural tube defects (NTD) and abdominal wall defects (AWD)

Screening for Down syndrome

This has a live birth incidence of 1.4 per 1000 in Australia.[6] The risk of conceiving a child with Down syndrome increases proportionally with age (Figure 19.3). For a woman aged 21, it is 1 in 1000, while for a woman aged 35, it is 1 in 275 and for a woman aged 45, it is 1 in 20.



Figure 19.3 Risk of Down syndrome by maternal age

The tests available to test for Down syndrome can be divided into two types:

1. Screening tests (maternal serum screening, ultrasound) - safe but relatively low predictive values
2. diagnostic tests (chorionic villus sampling, amniocentesis)

The most reliable method is obtaining foetal tissue by these last means but there is a significant risk of miscarriage (1 in 100 for chor-ionic villus sampling and 1 in 200 for amniocentesis).

Usual antepartum timing of tests

First trimester serum screening tests:

- pregnancy-associated plasma protein (PAPP-A) - decreased in Down syndrome
- free β-human chorionic gonadotrophin (HCG) - increased in Down syndrome

If combined with first trimester ultrasound nuchal translucency it is termed the combined test.

Second trimester serum screening tests:

- α-fetoprotein - decreased
- unconjugated oestriol - decreased
- free β-HCG - increased
- inhibin A - increased

The tests can be calculated in conjunction with maternal age and ultrasound tests to give predictive values.

A quadruple test with ultrasound dating will have a 75% detection rate but 65 - 70% without it.[15]

CONSANGUINITY

Consanguinity is the situation where a couple shares one or more common ancestors. Consanguineous relationships occur in most societies and in some cultures are associated with particular advantages and religious traditions.[6]

The concern about the issue has been questioned by a study in the *Journal of Genetic Counselling*, which suggested ‘aside from a thorough history, there is no need to offer genetic testing on the basis of consanguinity alone’. In a comment on this study which indicated that ‘cousins run few risks in mating’, the author pointed out that people like Charles Darwin, Queen Victoria and Albert Einstein married first cousins and Darwin's children, in particular, were brilliant. In parts of Saudi Arabia 39% of all marriages are between first cousins.[16] On the other hand, there are substantiated records of problems with inherited disorders. Twenty-four US states ban the marriage of first cousins and the Catholic Church, with 2000 years of experience, has opposed cousin marriages for over 1000 years.

First cousins share a pair of grandparents and are statistically at an increased risk for autosomal recessive conditions. The baseline risk that any couple will have a baby with a birth defect is 3 - 4%.[6] In addition, empirical data shows that for first cousin marriage there is an additional 4% risk of having a child with a birth defect - this includes malformations such as intellectual impairment and many rare autosomal recessive disorders. So the combined risk is 8% irrespective of a positive family history, which is an important consideration in counselling these people.[6]

For siblings the risk of increase in birth defects is 30% and 19% for second cousins.

RARE BUT HELPFUL DIAGNOSTIC TIPS

- dark black urine on exposure ° alkaptonuria

DxT: arthritis + pigmentation of ear cartilage + black urine with alkalisation ° alkaptonuria

- red urine on exposure ° porphyria
- blue diapers ° tryptophan malabsorption syndrome
- maple syrup odour (urine and perspiration) ° maple syrup urine disease (AR) (an amino acid metabolic disorder)

DxT: odour + hypertonicity + seizures (infancy) ° maple syrup urine disease

- mousy body odour ° phenylketonuria

- ‘fish-like’ mouth with microganthia ° Turner syndrome
- ‘chipmunk’ facies ° thalassaemia major
- ‘doll-like’ facies ° glycogen storage disease (G-6-P deficiency)
- elfin face ° Williams syndrome
- butterfly-like facial rash in children ° tuberous sclerosis
- weak suction + delayed sitting and crawling ° Prader-Willi syndrome
- ‘happy puppet’ features ° Angelman syndrome
- characteristic eyes: wide-spaced, down-slanting openings, ptosis ° Noonan's syndrome
- long fingers and limbs ° Marfan syndrome
- high-pitched meowing cry low set ears, mental retardation ° ‘cat-cry’ syndrome (maladie du cri du chat)
- blue sclera ° osteogenesis imperfecta

REFERENCES

1. Golden F, Lemonick M. The race is over. Time (Asia Pacific edn), 2000; 3 July: 56 - 61.
2. Tierney LM et al. *Current Medical Diagnosis and Treatment* (41st edn). New York: McGraw-Hill & Lange, 2002: 1645.
3. Warner BJ, McArthur GA. Cancer and the genetic revolution. Aust Fam Physician, 2001; 30: 933 - 5.
4. Newstead J, Metcalfe S. Getting the gene into genetic practice. Aust Fam Physician, 2001; 30: 927.
5. Kingston H. Clinical genetic services. BMJ, 1989; 298: 306 - 7.
6. Craig J, Amor D, Macciocca I et al. Genetics. CHECK, 2001; issue 349.
7. Lucarelli G. Bone marrow transplantation in adult thalassaemia patients. N Engl J Med, 1999; 93: 1164.
8. Delatycki M, Tassicker R. Adult onset neurological disorders. Predictive genetic testing. Aust Fam Physician, 2001; 30(10): 948 - 52.
9. Gaff C, Newstead J, Metcalfe S. *The Genetics File*. Meltourne: Victorian Department of Human Services, 2003.
10. Levy-Lahad E, Bird T. Alzheimer's disease: genetic factors. In: Pulst SM, ed. Neurogenetics. New York: Oxford University Press, 2000.
11. Lennox N, Diggins J. *Management Guidelines: People with Development and Intellectual Disabilities* (1st edn). Melbourne: Therapeutic Guidelines Ltd, 2005.
12. Beers MH, Berkow R. *The Merck Manual of Diagnosis and Therapy* (17th edn). New Jersey: Merck Research Laboratories, 1999: 2238 - 9.
13. Bankier A. Syndrome Quiz. Fetal alcohol syndrome. Aust Fam Physician, 1990; 19: 1297.
14. Amor D. Familial cancers. Aust Fam Physician, 2001; 30: 937 - 45.
15. Sheffield L. Prenatal screening and diagnosis of genetic disorders. Curr Ther, 2002; April: 12 - 18.
16. Corliss R. Cousins: a new theory of relativity. Time, 2002; 15 April: 43.

CHAPTER 20. DEPRESSION

I am ignorant and impotent and yet, somehow or other, here I am, unhappy, no doubt, profoundly dissatisfied … In spite of everything I survive.

ALDOUS HUXLEY (1894 - 1963)

DEPRESSION

Depressive illness, which is probably the greatest masquerade of general practice, is one of the commonest illnesses in medicine and is often confused with other illnesses. It is a very real illness that affects the entire mind and body. Unfortunately, there is a social stigma associated with depression and many patients tend to deny that they are depressed.

It is a useful working rule to consider depression as an illness that seriously dampens the five basic activities of humans:

- energy for activity
- sex drive
- sleep
- appetite
- ability to cope with life

Many episodes of depression are transient and should be regarded as normal but 10% of the population have significant depressive illness. The lifetime risk of being treated for depression is approximately 12% for men and 25% for women.[1]

CLASSIFICATIONS

- Affective or mood disorders refer to those conditions in which there is a disturbance of affect or mood.
- The DSM-IV (TR) classification divides the disorder into the depressive disorders and bipolar disorders (both manic and depressive episodes).
- The depressive disorders include major depression, adjustment disorders with depressed mood, and dysthymia.
- Major depression includes those disorders with one or more major depressive episodes.[1]
- Dysthymia refers to long-standing (2 years or more) depression of mild severity (‘neurotic depression’).[1]
- Adjustment disorder with depressed mood is a less severe form of depression without sufficient criteria for major depression. It is very common and occurs in response to identifiable stressors (‘reactive depression’, e.g. loss of employment). Its duration is usually no longer than 6 months.

MAJOR DEPRESSION

The patient can experience many symptoms, both physical and mental. The DSM-IV (TR) diagnostic criteria for depression are outlined in the box on page 181.

These criteria can be extended to include:

- a feeling of not being able to cope with life
- continual tiredness (weariness with much sighing)
- loss of sense of humour
- tension and anxiety
- irritability, anger or fearfulness
- somatic symptoms, such as headache, constipation, indigestion, weight loss, dry mouth and unusual pains or sensations in the chest or abdomen

The symptoms may vary during the day, but are usually worse on waking in the morning. Some patients have psychotic features, usually only delusions but sometimes also hallucinations, and may be misdiagnosed as schizophrenic.

In practice the DSM-IV (TR) classification seems too rigid and the experienced doctor has to consider the global constellation of symptoms. Better management follows early diagnosis and intervention before the formal criteria for major depression develop.

DSM-IV (TR) diagnostic criteria for major depression

At least five of the following symptoms for 2 weeks (criterion 1 or 2 essential)

1. pervasive depressed mood
2. marked loss of interest or pleasure

3. significant appetite or weight loss or gain (usually poor appetite)
4. insomnia or hypersomnia (usually early morning waking)
5. psychomotor agitation or retardation
6. fatigue or loss of energy
7. feelings of worthlessness or excessive guilt
8. impaired thinking or concentration; indecisiveness
9. suicidal thoughts/thoughts of death or suicide

Source: *Diagnostic and Statistical Manual* (4th edn revised). Washington, DC: American Psychiatric Association, 2000.

Important points

- The essential feature of depression is mood change, which can vary in intensity from despondency to intense despair.[1]
- The other major feature is loss of interest or pleasure, including loss of interest in family, hobbies, sexual activity and personal appearance.
- There is an association between anxiety and depression and this possibility must be considered in management.

MINOR DEPRESSION

The diagnosis of minor depression is based on a total of two to four symptoms from the above list, including 1 and 2.

Minor depression is basically a condition where fluctuations of symptoms occur with some vague somatic symptoms and a transient lowering of mood that can respond to environmental influences. Suicidal feelings are fleeting, and delusions and hallucinations are absent. These patients usually respond in time to simple psychotherapy, reassurance and support. However, care should be taken lest they move into major depression.

MASKED DEPRESSION

This is a difficult yet common type of depression in practice and tends to be misdiagnosed. Patients do not complain of the classic symptoms and tend to deny depression, which is perceived as a social stigma and a sign of weakness. They usually have multiple minor complaints of the 'ticket of entry' type. Mood changes may be elicited only after careful questioning.

The classic affective features of depression are masked by a complex of somatic complaints. Such symptoms include fatigue, anorexia, weight loss, menstrual changes, unusual sensations in the abdomen, chest or head, bodily aches and pain, dry mouth, and difficulty in breathing.

If depression is not considered many fruitless, expensive and distressing in-vestigations may be performed. According to Davies,[2] nearly half of the patients with de-pressive illness report to the doctor with complaints that suggest physical illness. The family doctor has to suspect masked depression in a patient with a multitude of physical complaints or with complaints that do not fit any definite pattern of organic disease.

The differential diagnoses of depression are presented in Table 20.1.

Table 20.1 Differential diagnoses of depression

Psychiatric conditions
Anxiety disorder Schizophrenia Drug and alcohol abuse Dementia
Organic disorders
Malignancy (e.g. lung, pancreas, lymphoma) Hypothyroidism Hyperparathyroidism Other endocrine disorders (e.g. Cushing's, Addison's) Anaemia, especially pernicious anaemia Post-infective states (e.g. EBM) Cerebrovascular disease Parkinson's disease

- Congestive cardiac failure
- Systemic lupus erythematosus
- Drugs (which may cause depression)
    - antihypertensives
    - benzodiazepines (e.g. diazepam)
    - antiparkinson drugs
    - corticosteroids
    - cytotoxic agents
    - NSAIDs
    - oral contraceptives/progestogen

An Australian study on masked depression concluded:[3]

> It must be stressed that the masking of the depressive state occurs on the doctor's side as well as the patient's, and an awareness that this may be so leads us to recommend that, once organic lesions have been excluded, there is a place for the use of an adequate therapeutic trial of antidepressants.

The following additional points were made by a panel of psychiatrists at a symposium entitled 'Depression: masked or missed?' in Dallas, Texas:[4]

- Some patients dismissed as 'crocks' may go on to suicide if their depression is not treated.
- Masked depression would be missed much less frequently if the physician would look beneath symptoms that do not quite ring true.
- The patient with the 'tired blood syndrome' deserves something other than an iron tonic.
- Depression frequently accompanies organic diseases that are associated with nausea and other illness.
- A complete work-up may help to rule out organic disease but may result in iatrogenic disease if pursued overzealously.
- Alcoholism should be suspected as a cause of depression.

DEPRESSION IN THE ELDERLY

Severe depression affects 1 - 2% of the elderly population while 10% have significant depression affecting their life. Milder depression can affect a further 20%. Depression can have bizarre features in the elderly and may be misdiagnosed as dementia or psychosis. Agitated depression is the most frequent type of depression in the aged.[1] Features may include histrionic behaviour, delusions and disordered thinking.

Depression is often missed in the elderly because it is atypical and less expressive, and patients tend to be ashamed and reluctant to admit it. Four key guidelines help diagnosis:

- Are you basically satisfied with your life?
- Do you feel that your life is empty?
- Are you afraid that something bad is going to happen to you?
- Do you feel happy most of the time?

A useful clue is a change in sleep pattern, so a request for sleeping tablets may lead to the prescription for a more sedating antidepressant. Medical illness is an important precipitant of depression in the elderly. Tricyclic anti-depressants have to be used with caution in the elderly and most have some contraindications to their use. Electroconvulsive therapy (ECT) has a useful place in the treatment of severe cases.

DEPRESSION IN CHILDREN

Sadness is common in children but depression, although not as common, does occur and is characterised by feelings of helplessness, worthlessness and despair. Parents and doctors both tend to be unaware of depression in children.[5]

Major depression in children and adolescents (page) may be diagnosed using the same criteria as for adults, namely loss of interest in usual activities and the presence of a sad or irritable mood, persisting for 2 weeks or more.[6] The other con-stellation of depressive symptoms, including somatic complaints, may be present. Examples include difficulty in getting to sleep, not enjoying meals, poor concentration and low self-esteem. Poor motor skills and family instability are a proven association. Depression can present as antisocial behaviour or as a separation anxiety (e.g. school refusal). Although suicidal thoughts are common, suicide is rare before adolescence. Depressed adolescents are a serious suicide risk. Referral of these patients to an experienced child psychiatrist is advisable.

POST-PARTUM DEPRESSION[7]

Depressive symptoms in the postpartum phase range from normal ‘baby blues’ to the more serious postnatal depression (about 13% of work). Prompt psychiatric intervention is essential for the small percentage who exhibit psychotic symptoms, those who are suicidal, or where there is a risk of harm to the child. It is treated in the same way as major depression. Psychosocial treatments are important

THE DIAGNOSTIC APPROACH

Depression can be associated with many illnesses but it is important to realise that the somatic symptoms may be the presentation of depressive illness and thus ‘undifferentiated illness’ is a feature. The patient tends to complain of aches and pains, gastrointestinal symptoms and other similar symptoms rather than emotional problems.

There is a relationship between anxiety and depression so that many depressed patients are agitated and anxious - a feature that may mask the underlying depression.[8]

Questions to assess level of depression

- What do you think is the matter with you?
- Do you think that your feelings are possibly caused by nerves, anxiety or depression?
- Can you think of any reason why you feel this way?
- Do you feel down in the dumps?
- Do you feel that you are coping well?
- Do you have any good times?
- Has anything changed in your life?
- How do you sleep? Do you wake early?
- What time of the day do you feel at your worst?
- Where would you put yourself between 0 and 100%?
- Have you felt hopeless?
- Do you brood about the past?
- What is your energy like?
- What is your appetite like?
- Are you as interested in sex as before?
- Do you feel guilty about anything?
- Do you feel that life is worthwhile?
- Has the thought of ending your life occurred to you?
- Do you cry when no one is around? (especially for children)
- Two particularly good questions are:
- In the past month have you been bothered by feeling down, depressed or hopeless?
- In the past month have you often been bothered by little interest or pleasure in doing things?

Depression scales

Consider the use of depression scales, for example:

- Hamilton Depression Inventory
- Beck Depression Scale
- General Health Questionnaire
- Geriatric Depression Scale

These may be most useful as an aide-mémoire since studies show that no tests or questionnaires have been found to be practical in detecting early depression. Family and friends' awareness of the signs and symptoms and alerting the GP is helpful.[9]

MANAGEMENT

Important considerations from the outset are:

- Is the patient a suicide risk?
- Does the patient require inpatient assessment?
- Is referral to a specialist psychiatrist indicated?

If the symptoms are major and the patient appears in poor health or is a suicide risk, referral is appropriate.

The basic treatments are:

- Psychotherapy, including education, reassurance and support. All patients require minor psychotherapy. More sophisticated techniques, such as cognitive or behavioural therapy, may be used for selected patients. Cognitive behaviour therapy (CBT) basically involves teaching patients new ways of positive thinking, which have to be relevant and achievable for the patient. Patients need to be able to recognise their own negative cognitions, including their anxieties and worries.
- Pharmacological agents.
- ECT.

Note: Reassurance and support are needed for all depressed patients.

Useful guidelines

- Mild depression: psychotherapy alone, especially CBT, may suffice and may be more effective than drug therapy[10] but keep medication in mind.
- Moderate to severe depression: psychotherapy plus antidepressants is recommended.
- Reassurance and support are needed for all depressed patients.

Explanatory supportive notes for patients and relatives[11]

Most people feel unhappy or depressed every now and again, but there is a difference between this feeling and the illness of depression.

Depression is a very real illness that affects the entire mind and body. People cannot seem to lift themselves out of their misery or ‘fight it themselves’. Superficial advice like ‘snap out of it’ is unhelpful because the person has no control over it.

What is the cause?

The cause is somewhat mysterious but it has been found that an important chemical is present in smaller amounts than usual in the nervous system. It is rather like a person low in iron becoming anaemic.

Depression can follow a severe loss, such as the death of a loved one, a marital separation or financial loss. On the other hand, it can develop for no apparent reason, although it may follow an illness such as glandular fever or influenza, an operation or childbirth. Depression is seen more commonly in late adolescence, middle age (both men and women), retirement age and in the elderly.

What is the treatment?

The basis of treatment is to replace the missing chemicals with antidepressant medication. Antidepressants are not drugs of addiction and are very effective but take about 2 weeks before an improvement is noticed. Alcohol can interact with the tablets, so it is important not to drink and drive. If the person is very seriously depressed and there is a risk of suicide, admission to hospital will most likely be advised. Other more effective treatments can be used if needed. The depressed person needs a lot of understanding, support and therapy. Once treatment is started, the outlook is very good (an 80% cure rate).

Important points

- Depression is an illness.
- It is more common than realised.
- It just happens; no one is to blame.
- It affects the basic functions of energy, sex, appetite and sleep.
- It can be lethal if untreated.
- It can destroy relationships.
- The missing chemical needs to be replaced.
- It responds well to treatment.

Recommended reading

Paul Hauck. *Overcoming Depression*. London: The Westminster Press, 1987.

Gordon Parker. *Dealing with Depression*. Sydney: Allen & Unwin, 2002.

ANTIDEPRESSANT MEDICATION

The mode of action of available antidepressants is determined by receptor-mediated signal transduction in the serotonergic or nor adrenergic bathnap. The initial choice of an antidepressant depends on the age and sex of the patient, prior response to medication, safety in overdosage and the side-effect profile. All antidepressants are equally efficacious. The tricyclics and tetracyclics have been the first-line drugs but the newer drugs, the selective serotonin reuptake inhibitors (SSRIs) and moclobemide (a reversible monoamine oxidase inhibitor (MAOI) antidepressant) are equally effective, are better tolerated, have a wider safety margin[6] and are now considered first-line drugs.

Tricyclic antidepressants[6]

1. Amitriptyline and imipramine
    - the first generation tricyclics
    - the most sedating - valuable if marked anxiety and insomnia
    - strongest anticholinergic side-effects (e.g. constipation, blurred vision, prostatism)
2. Clomipramine, desipramine, dothiepin, doxepin, nortriptyline
    - less sedating and anticholinergic activity
    - nortriptyline is the least hypotensive of the tricyclics

Dosage

- 50 - 75 mg (o) nocte, increasing every 2 to 3 days to 150 mg (o) nocte by day 7.
- If no response after 2 to 3 weeks, increase by 25 - 50 mg daily at 2 to 3 week intervals (depending on adverse effects) to 200 - 250 mg (o) nocte. Trial for 6 weeks.

General adverse effects

- Dry mouth, weight gain, constipation, sedation
- Glaucoma, urinary retention, tremor
- Confusion and delirium in the elderly (caution in the elderly)
- Sexual dysfunction
- Postural hypotension
- Cardiac conduction impairment (caution in heart disease)
- Lowered seizure threshold

Tetracyclic antidepressants[6]

- Mianserin 30 - 60 mg (o) nocte increasing to 60 - 120 mg (o) nocte by day 7

Adverse effects

- Sedation, lethargy, dizziness, polyarthritis, dry mouth, headache
- Neutropenia (reversible) especially > 65 years (uncommon)
- Less anticholinergic effects than tricyclics
- Fewer cardiovascular side-effects

Selective serotonin reuptake inhibitors

- Fluoxetine and paroxetine 20 mg (o) mane. This dose is usually sufficient for most patients. If no response after 2 to 3 weeks, increase by 20 mg at 2 to 4 week intervals to 40 - 80 mg (o) daily in divided doses.
- Sertraline, 50 mg (o) daily, starting dose; can increase to 200 mg daily.
- Fluvoxamine, 50 mg (o) bd, starting dose; can increase to 200 mg daily.
- Citalopram 20 mg (o) daily, up to 60 mg (max.).
- Escitalopram 10 mg (o) daily, up to 20 mg.

These new drugs have a similar efficacy profile to the tricyclics. They do not appear to cause weight gain, interact with alcohol or cause serious cardiovascular effects. There are definitive differences between the SSRIs, and swapping when one is ineffective can be beneficial.

Adverse effects

Nausea, nervousness, fatigue, agitation, diarrhoea, headaches, insomnia. Possible effects include sexual dysfunction, mainly ejaculatory disturbances, allergic reactions and hypomania (in some manic depressives).

They should not be used with MAOIs or the tricyclics.

When withdrawing an SSRI, do it slowly and leave a recommended ‘wash-out’ period if changing to another class drug.

Monoamine oxidase inhibitors (MAOIs)

MAOIs are second-line antidepressants and are usually reserved for use by psychiatrists. The adverse effect is significant hypertension when combined with various foodstuffs, especially those containing large amounts of tigramine or other amines

- Phenelzine 45 - 90 mg daily in 2 or 3 divided doses
- Tranylcyromine 20 - 40 mg daily in 2 or 3 divided doses

Both drugs should be reduced to the lowest effective dose for maintainence once a response is obtained.

Moclobemide

- Moclobemide 150 mg (o) bd. If no response after 2 to 3 weeks, increase by 50 mg daily to maximum 300 mg (o) bd.

This is a reversible MAOI, which is less toxic than the irreversible MAOIs. It has minimal interaction with tyramine-containing foodstuffs, so that no dietary restrictions are necessary.

Adverse effects include nausea, headache, agitation, dizziness and insomnia.

The irreversible MAOIs, which should be reserved for second-line MAOI therapy, include phenelzine and tranylcypromine.

Serotonin noradrenaline reuptake inhibitors (SNRIs)

Venlafaxine

The first of the SNRIs to be released is venlafaxine. It is recommended for major depression where other therapy is in-appropriate. The starting dose is 37.5 mg (o) bd increasing to 75 mg bd after 2 weeks if necessary. A long-acting 75 mg once daily preparation is also available.

Side-effects appear to include nausea, dizziness, headache, sweating, insomnia and sexual dysfunction, giving the drug a similar side-effect profile to the SSRIs. It should not be used concomitantly with MAOIs and various ‘wash-out periods’ from other antidepressants are required.

Serotonin antagonist

Mirtazapine

Mirtazapine is considered to enhance central non-adrenergic and serotonin activity. Reported adverse effects include somnolence, increased appetite and weight gain. It is useful for patients with depression and insomnia. The dose is 15 mg (o) nocte increasing to 45 mg (o) nocte.

Noradrenaline reuptake inbitors

Reboxetine

Reboxetine is a selective noradrenaline reuptake inhibitor. It is recommended for the treatment of major depression.

The adult dose is 4 mg Bd increasing (if required) after 3 weeks to 10 mg daily.

Reported adverse effects include dry mouth, sweating, headache, insomnia, urinary retention, constipation and impotence.

NOTES ABOUT ANTIDEPRESSANTS[6]

- Tricyclics can be given once daily (usually in the evening).
- There is a delay in onset of action of 1 - 2 weeks after a therapeutic dose (equivalent to 150 mg imipramine at least) is reached.
- Each drug should have a clinical trial at an adequate dose for at least 4 - 6 weeks before treatment is changed.
- The SSRIs are probably now the first-line drugs of choice and the tricyclics second-line.
- Do not mix antidepressants.
- Combinations of antidepressants have not been shown to be more effective than monotherapy and there is the risk of severe adverse effects, such as the scerotonin syndrome.
- Consider referral if there is a failed (adequate) trial.
- Swapping from one antidepressant to another in those not responding is a proven beneficial strategy.
- Full recovery may take up to 6 weeks or longer (in those who respond).
- Continue treatment at maintenance levels for at least 6 to 9 months.[1] There is a high risk of relapse.
- For a second episode use antidepressants for 3 to 5 years.
- MAOIs are often the drugs of choice for neurotic depression or atypical depression.[1]

The serotonin syndrome[12]

This is a dangerous adverse reaction related to the use of the SSRIs and is most likely to occur with the combined use of MAOI drugs and other agents. The diagnosis is based on three criteria:

- Symptoms must coincide with the introduction or dose increase of a serotonergic agent.
- Other causes, such as infection, substance abuse or withdrawal, must be excluded.
- At least three of the symptoms or signs attributed to the syndrome must be present, i.e.
  - mental status/behaviour changes (e.g. agitation, confusion, hypomania, seizures)
  - altered muscle tone (e.g. tremor, shivering, myoclonus, hyper-reflexia)
  - autonomic instability (e.g. hypertension, tachycardia, fever, diarrhoea)

The offending agents should be withdrawn immediately and supportive therapy initiated.

COMPLEMENTARY THERAPY

St John's wort (Hypericum perforatium) has been found to be effective in mild-to-moderate depression[13] but a recent study showed that it was no better than placebo in treating moderately severe to major depression.[14] Considerable concern has been raised over the potential for St John's Wort to interact with prescription medication, including all antidepressants, warfarin, digoxin, anti-convulsants and the oral contraceptive pill.[15] Other herbal remedies, such as kava kava or valerian root, have not proved effective for the treatment of depression.

ELECTROCONVULSIVE THERAPY

ECT is safe, effective and rapidly acting.[1, 6, 16]

Indications

- Psychotic depression (e.g. delusions, hallucinations)
- Melancholic depression unresponsive to antidepressants
- Substantial suicide risk
- Ineffective antidepressant medication
- Severe psychomotor depression
  - refusal to eat or drink
  - depressive stupor
  - severe personal neglect

Immediate referral for hospital admission is necessary in most of these circumstances. The usual course is about 9 treatments over 3 to 5 weeks. Antidepressants are usually dis-continued during ECT but resumed (mood stabilisers can be an alternative) after ECT to prevent relapse.

Transcranial magnetic stimulation is an experimental procedure being explored as a less invasive alternative to ECT.

RECURRENT DEPRESSION

Lifelong antidepressant therapy may have to be considered. Lithium is an alternative medication for long-term use. New treatments are based on vagal nerve stimulation.

Recurrent brief depression

There is a high prevalence in general practice of patients presenting with recurrent episodes of depression of short duration, about 3 to 7 days, as often as monthly. PMT may be a factor. As a rule antidepressants are ineffective management is base on psychotherapy especially CBT.

Seasonal affective disorder

SAD or ‘winter blues’ is a recurrent disorder seen in people living in cold climates where the winters are bleak and dark. Features of depression include somnolence and increased appetite. Treatment is based on psychotherapy, phototherapy and medication (SSRIs). Refer http://www.sada.org.uk

SUICIDE

The risk of suicide is a concern in all depressed patients. Between 11% and 17% of people who have suffered a severe depressive disorder at any time will eventually commit suicide.[17] Referral for hospital admission should be arranged for patients who are at great risk for suicide. There is a distinction between patients who are determined to suicide and those who attempt suicide (parasuicide).

Risk factors for suicide include:

- male sex
- older age > 55 years
- adolescents
- young adults 15 - 25 years
- immigrant status
- isolation/living alone
- recent divorce, separation or bereavement
- recent loss of employment or retirement
- family history of psychiatric illness (including suicide)
- impulsive, hostile personality
- previous suicide attempt
- severe depression
- financial difficulties
- alcohol or other substance abuse
- psychosis
- early dementia
- physical illness, especially if chronic pain

A useful suicide risk assessment is the SAD PERSONS (mnemonic) index (Table 20.2). A score greater than 7 represents a very high risk that demands careful attention, including referral to an acute psychiatric service. The suicide rates in Australia, which demonstrate two peaks in males, are illustrated in Figure 20.1.

Table 20.2 SAD PERSONS Index: Suicide risk assessment[17]

Risk factor	Criteria	Score
Sex	Male	1
Age	< 20 years; > 45 years	1
Depression	Major (e.g. depressed mood)	2
Psychiatric history	Previous attempts	1
Excessive drug use	Ethanol or other drug abuse	1
Rationality loss	Psychosis, severe depression	2
Separated	Loss of spouse or other single	1
Organised plan	Determined suicide plan	2
No supports	No community back-up; generally isolated	1
Sickness	Chronic illness	1

Score > 7 ° high suicide risk



Figure 20.1 Suicide rates in Australia, 2000, 5-year age groups by sex Australian Bureau of Statistics Prepared by The Hunter Institute of Mental Health.

The rules of 7

- 1 in 7 people with recurrent depression suicide.
- 70% of suicides have depression.
- 70% of patients who suicide have seen a 6P within 7 weeks.
- Suicide is the 7th leading cause of death.

If there is concern about suicide risk and treatment is supervised outside hospital, provide closer supervision and considerable support, and prescribe drugs that are less toxic in overdosage (e.g. mianserin or fluoxetine). If tricyclics are prescribed, useful guidelines are that dangerous medical complications occur with an equivalent dosage of 1000 mg (40 tablets) of imipramine and a high risk of death with 2000 mg (80 tablets).[6]

WHEN TO REFER[1]

- Uncertainty about diagnosis
- Inpatient care obviously necessary
- Severe depression
- Inability to cope at home
- Psychotically depressed (with delusions or hallucinations)
- Substantial suicide risk
- Failure of response to routine antidepressant therapy
- Associated psychiatric or physical disorders
- Difficult problem in the elderly - where diagnosis including dementia' is doubtful
- Children with apparent major depression

REFERENCES

1. Burrows GD. Depressive disorders. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 139 - 41.
2. Davies B. An Introduction to Clinical Psychiatry. University of Melbourne, 1977: 76 - 7.
3. Serry DK, Serry M. Masked depression and the use of antidepressants in general practice. Med J Aust, 1969; 15 February: 35 - 7.
4. Depression: masked or missed? Patient Care, 1972; 1(3): 6 - 14.
5. Robinson MJ. Practical Paediatrics (2nd edn). Melbourne: Churchill Livingstone, 1990: 552 - 3.
6. Dowden J (Chairman). Therapeutic guidelines: Psychotropic (Version 5). Melbourne: Therapeutic Guidelines Ltd, 2003.
7. National prescribing service Ltd. Managing depression in primary care. NPS News, 2005; 42 1 - 6.

8. Davis A, Bolin T, Ham J. Symptom Analysis and Physical Diagnosis (2nd edn). London: Bailliere Tindall, 1990: 983.
9. Rosser WM, Shafir MS. Evidence-based Family Medicine. Hamilton: BC Decker Inc., 1998: 156 - 7.
10. Goodlee F et al. In Clinical Evidence (2nd edn). London: BMJ Publishing Group, 1999.
11. Murtagh J. Patient Education (3rd edn). Sydney: McGraw-Hill, 2000: 164.
12. Keltner N. Serotonin syndrome: a case of fatal SSRI/MAOI interaction. Perspect Psychiatr Care, 1994; 30(4): 26 - 31.
13. Linde K, Ramirez G, Mulrow CD et al. St John's Wort for depression - an overview and meta-analysis of randomised clinical trials. Br Med J, 1996; 313: 253 - 8.
14. Walsh TB. Effect of hypericum perforatum in management of depression. JAMA, 2002; 287: 1807 - 14, 1840 - 7.
15. National Prescribing Service Ltd. NPS NEWS, 2000; 11: 4 - 6.
16. Kumar PJ, Clark ML. Clinical Medicine (2nd edn). London: Saunders 2003; 1241 - 7.
17. Rogers I. Guidelines for the Management of Common Emergencies in the Emergency Department (2nd edn). Auckland Hospital, 1996: 43.

CHAPTER 21. DIABETES MELLITUS: DIAGNOSIS

Those labouring with this Disease, piss a great deal more than they drink. Authors who affirm the drink to be little or nothing changed are very far from the truth, because the urine very much differed both from the drink taken in and also in being wonderfullysweet as if it were imbued with honey or sugar.

THOMAS WILLIS (1621 - 75), THE PISSING EVIL

DIABETES MELLITUS: DIAGNOSIS

Diabetes comes from a Greek word meaning ‘to pass or flow through’ (i.e. excessive urination) and mellitus means ‘sweet’.

There are two main types of diabetes (see Table 21.1).

Table 21.1 Clinical differentiation between type 1 and type 2 diabetes

	Type 1	Type 2
Relative frequency (approx.)	15%	85%
Peak age incidence	10 - 30 years	>40 years
Age of onset	usually < 20	>40
Onset	rapid	insidious
Weight at onset	low (thin)	high (obese)
Ketoacidosis	yes	no
Familial	weak	strong
Insulin status	deficient	resistant
Complications	yes	yes

Note: These are generalisations and the clinical features may vary (e.g. type 2 diabetic patients may be thin and have a rapid onset; type 1 patients may exhibit a weak genetic link).

- Type 1 is also known as juvenile onset diabetes or insulin dependent diabetes mellitus (IDDM).
- Type 2 is also known as maturity onset diabetes or non-insulin dependent diabetes mellitus (NIDDM).

DIABETES: A REAL MASQUERADE

The onset of type 2 diabetes can be subtle and by stealth. Studies have demonstrated that it takes, on average, 7 - 9 years before a patient is diagnosed.[1]At any one given time 50% of patients with type 2 diabetesare undiagnosed. The Australian Diabetes, Obesity and Lifestyle Study (AUSDIAB) in 2000 showed that one in four adult Australians have abnormal glucose metabolism.[2] In general figures the prevalence of diagnosed and undiagnoseddiabetes is4% each and a further 16% have impaired fasting glucose or impaired glucose tolerance[3]. Very importantly, about 35% of newly diagnosed diabeticpatients are already harbouring complications of diabetes.[1] The challenge for general practitioners is to be on constant lookout for these patients especially those at risk. Type 2 diabetesis becoming more prevalent in industrial countries - partly due to the ageing population and partly because our lifestyle encourages us to ‘eatmore and walk less’.[3] Furthermore, 60% of our population are overweight or obese.

KEY FACTS AND CHECKPOINTS

- In Australians older than 25 years the prevalance of diabetes is 7.5%, with another 10.6% having impaired glucose tolerance.[2]
- About 30% of these people will develop clinical diabetes within 10 years.[3]
- Many type 2 diabetics are asymptomatic.
- Diabetes can exist for years before detection and complications may be evident.
- Type 2 diabetes is not a mild disease. About one-third of those surviving 15 years willrequire insulin injections to control symptoms or complications.[4] Complications occur in type 2 diabetes as well as in type 1.
- There are several causes of secondary diabetes that are very uncommon(see Table 21.2).

Table 21.2 Causes of secondary diabetes

Endocrine disorders •Cushing's syndrome •Acromegaly •Phaeochromocytoma •Polycystic ovary syndrome
Pancreatic disorders •Haemochromatosis •Chronic pancreatitis
Drug-induced diabetes (transient) •Thiazide diuretics •Oestrogen therapy (high dose - not with low-dose HRT) •Corticosteroids
Other transient causes •Gestational diabetes •Medical or surgical stress

- Asymptomatic people of high risk of undiagnosed diabetes should be screened by plasma glucose measurement.
- Blood glucose may be temporanly devatod during acute illness after trauma or surgery.

CLINICAL FEATURES

The classic symptoms of uncontrolled diabetes are:

- polyuria
- polydipsia
- loss of weight (type 1)
- tiredness and fatigue
- propensity for infections, especially of the skin and genitals (vaginal thrush)

The young type 1 diabetic person typically presents with a brief 2 - 10 week history of the classic triad of symptoms:

DxT: thirst + polyuria + weight loss ° type 1 diabetes

Other possible symptoms are:

- vulvovaginitis due to *Candida albicans*
- pruritus vulvae due to *Candida albicans*
- balanitis due to *Candida albicans*
- nocturnal enuresis (type 1)
- blurred vision/visual changes

Symptoms of complications (may be presenting feature) include:

- staphylococcal skin infections
- polyneuropathy
    - tingling or numbness in feet
    - pain (can be severe if present)
- impotence
- arterial disease
    - myocardial ischaemia
    - peripheral vascular disease

The clinical examination should follow the guidelines under the heading ‘Examinations’ in Table 128.5 (p. 200).

HISTORY

The history of a suspected or known diabetic patient should cover the following features including assessment of cardiovascular risks and end-organ damage:

- specific symptoms:
  - polyuria
  - polydipsia
  - loss of weight
  - polyphagia
  - tiredness/malaise/fatigue
  - nocturia
- related general symptom review:
  - cardiovascular, e.g. chest pain, dyspnoea
  - urinary function
  - sexual function
  - neurological, e.g. tingling in feet/hands
  - vision, e.g. blurred
  - infection tendency, e.g. skin, urine, genital
  - genital itching
- general:
  - family history
  - medication
  - smoking and alcohol
  - obstetric history (where applicable)
  - physical activity
  - nutrition/eating habits

EXAMINATION

The physical examination should ideally follow the protocol for annual review.

Initial screening for suspected diabetes should include:

- general inspection including skin
- BMI (weight/height)
- waist circumference
- visual acuity
- blood pressure - lying and standing
- test for peripheral neuropathy: tendon reflexes, sensation (e.g. cotton wool, 10 g monofilament, Neurotip)
- urinalysis: glucose, albumin, ketones, nitrites

INVESTIGATIONS

- Initial:
  - fasting or random blood sugar
  - follow up oral glucose tolerance test (OGTT) if indicated
- Other tests according to clinical assessment, e.g. HbA1c, lipids, renal function, ECG

PEOPLE PREDISPOSED TO DIABETES

- Age > 40 years
- Family history
- Overweight/obesity
- Sedentary lifestyle

- Positive obstetric history
- Women with polycystic ovarian syndrome (PCOS)
- Hypertension//ischaemic heart disease
- Medication causing hyperglycaemia
- Ethnic/cultural groups
    - Aboriginal and Torres Strait Islanders
    - Pacific Islanders
    - Indian Subcontinent; Chinese
    - Afro-Caribbeans

WHO AND WHEN TO SCREEN?

- People with impaired fasting glucose/impaired glucose tolerance
- Age ≥ 55 years
- Age > 45 years with:
    - family history (1 ° relative with type 2)
    - obesity (BMI > 30)
    - hypertension
- Age ≥ 35 years from high prevalence ethnic groups
- Previous gestational diabetes; history of large babies
- People on long-term steroids
- People on atypical antipsychotics
- Polycystic ovarian syndrome, especially if overweight
- Cardiovascular disease and other risk factors

The optimal frequency is every 3 years and annually in very high-riskgroups.

DIAGNOSIS OF DIABETES

Diabetes is diagnosed as follows:[5, 6]

1. If symptomatic (at least two of polydipsia, polyuria, frequent skin infections or frequent genital thrush):
    - fasting venous plasma glucose (VPG) ≥ 7.0 mmol/L
    - random VPG (at least 2 hours after last eating) ≥ 11.1 mmol/L
2. If asymptomatic:
    - At least two separate elevated values, either fasting, 2 or more hours post-prandial, or the two values from an oral glucose tolerance test (OGTT)

Note: If random or fasting VPG lies in uncertain range (5.5 - 11.0 mmol/L) in either a symptomatic patient or a patient with risk factors (over 50 years, overweight, blood relative withtype 2 diabetes or high blood pressure) perform an OGTT. The cut-off point for further testing has now been reduced to 5.5 mmol/L.[7, 8]

The 2 hour blood sugar on an OGTT is still the gold standard for the diagnosk of uncertain diabetes, ie. >11.1 mmol/L.

The OGTT should be reserved for true borderline cases and for gestational diabetes. A screeningtest at 26 - 30 weeks gestation is recommended.

The diagnostic criteria after a 75 g load of glucose are.[8]

Venous plasma glucose in mmol/L	Fasting	2 hours later
Normal	< 5.5	< 7.8
Impaired tolerance	<7.0	7.8 - 11.1
Diabetes	>7.0	>11.1
Gestational diabetes	≥ 5.5	≥ 7.8

Urinalysis is unreliable in diagnosis since glycosuria occurs at different plasma glucose values in patients with different renal thresholds.

For a summary of diagnosis of diabetic staesrefer to Fig 21.1


< 5.5
F: 5.5–6.9
R: 5.5–11.0
F: ≥ 7.0
R: ≥ 11.1
Diabetes unlikely
Diabetes uncertain
Diabetes likely
Re-test yearly if high risk
3 yearly if increased risk
Oral glucose tolerance test
2 hour glucose levels
< 7.8
7.8–11
≥ 11.1
Diabetes unlikely
Diabetes likely
Impaired glucose tolerance
F = fasting
R = random

Figure 21.1 Glucose levels - venous plasma (mmol/L)

Source: Diabetes Management in General Practice. Melbourne: Diabetes Australia & RACGP. Reproduced with permission

DIABETES IN CHILDREN

A study by Sinah et al. detected impaired glucose tolerance in 25% of55 obese children (4 to 10 years of age) and 21% of 112 obese adolescents (11 to 18 years of age).[9] Type 2 diabetes was identified in 4% of obese adolescents. However over 30% of newly diagnosed diabetics in childhood and adolescents is upon presentation with diabetic ketoacidosis. These children especially infants may present as the very ill child with vomiting, stupor, dehydration, rapid respiration (Kussmal breathing) andketotic breath. Children with type 1 diabetes usually exhibit the classic features of polyuria, polydipsia, weight loss and lethargy. Be aware of unusual presentations such as urinary disorders including enuresis or daytime wetting accidents when a midiagnosis of urinary infection or some other condition is sometimes forthcoming. The diagnosis can be made by an elevated random or fasting blood sugar. Oral glucose tolerance tests are inappropriate in the very young. Upon diagnosis it is appropriate to refer the child or adolescent to a multidisciplinary diabetes team.

GESTATIONAL DIABETES

Gestational diabetes is the onset or initial recognition of abnormal glucose tolerance during pregnancy. Pregnancy is diabetogenic for those with a genetic predisposition. All pregnant women should be screened at 28 weeks with an oral glucose challenge test. If the blood glucose level is > 7.8 mmol/L or if diabetes is otherwise suspected a diagnostic oral glucose tolerance test is indicated. The WHO definition of gestational diabetes is fasting blood sugar of ≥ 7 mmol/L or a 2 hour level of ≥ 7.8 mmol/L. Refer to page XXX for further information.

VIABETES IN THE ELDERLY

The incidence of diafetes use with age. The elderly have incteased montalely and morbidity form the desease and sequre caveful monidoving especiall with adveise drieg effects aggravated by polypharmary and coi morbidities special uses include diet, foot care and porural lypolensum.

COMPLICATIONS OF DIABETES

Complications may occur in patients with both type 1 and type 2 diabetes, even with early diagnosis and treatment (Figure 21.2). Type 1 diabetics still have a significantly reduced life expectancy. The main causes of death are diabetic nephropathy and vascular disease (myocardial infarction and stroke).


CVA (increased)
General
hypoglycaemia
hyperglycaemia/
ketoacidosis
dyslipidaemia
Eyes
cataract
retinopathy
ocular palsies (III, VI nerves)
Blood pressure effects
postural hypotension
hypertension
Heart
accelerated atherosclerosis
myocardial infarction (may be silent)
vagal neuropathy → arrhythmias
Kidney
diabetic nephropathy
urinary infection
Gastrointestinal
gastroparesis
diarrhoea
peripheral nerve palsies
Genitourinary
impotence
atonic bladder
small muscle wasting
amyotrophy
(painful wasting)
Skin
candidiasis
• balanitis
• vulvovaginitis
staphylococcal infections
necrobiosis lipoidica
acanthosis nigricans
vitiligo
Peripheral vascular disease
macrovascular
microvascular
Peripheral (poly) neuropathy
paraesthesia
muscle weakness
reduced reflexes
± pain
Feet problems
ischaemia (rest pain)
ulcers
gangrene

Figure 21.2 The complications of diabetes

Diabetes causes both macrovascular and microvascular complications but microvascular disease isspecific to diabetes. Special attention should be paid to the ‘deadly quartet’ associated with type 2 diabetes.[8,10]

Macrovasular complicationd incude:

- Ischaemic/coronary heart disease.
- Cerebrovas cular disease.
- Peripheral vascular disease.

An analysis of patients with type 2 diabetes in the HOPE study[11,12] showed a benefit of ramipril to reduce the risk of:

- death (24%)
- myocardial infarction (22%)

- stroke (33%)
- cardiovascular death (37%)
- overt nephropathy (24%)

Conider organs/problems affected by diaprtes under the mnenomic'KNIVES'.

- Kidney
- Nerves
- Infection
- Vessels
- Eyes
- Skin

Microvascular disease

The small vessels most affected from a clinical viewpoint are the retina, nerve sheath and renl glomerulus. In younger patients it takes about 10 to 20 years after diagnosis for the problems of diabetic retinopathy, neuropathy and nephropathy to manifest.

Nephropathy

Prevention of diabetic nephropathy is an essential goal of treatment. Early detection of the yardstick, which is microalbuminuria, is important as the process can bereversed with optimal control. The dipstick method is unreliable. Screening is done simply by an overnight collection (10 - 12 hours) of all urine, including the first morning sample. It is sent to the laboratory to determine the albumin excretion rate. Microalbuminuria is 20 - 200 μg/minute (2 out of 3 positive collections).

ACE inhibitors should be used for evidence of hypertension.

Retinopathy and maculopathy

Retinopathy develops as a consequence of microvascular disease of theretina. Its prevalence is related to the duration of illness but up to 20% of people with type 2 diabetes have diabetic retinopathy at the time of diagnosis. The European multi-centre study[13] showed that diabetes is the single most commoncause of blindness in European adults in the 16 - 64-years age groups. It is recommended that patients should undergo fundoscopy each year by anexpert. Assessment is by direct ophthalmoscopy(with dilated pupils), retinal photography and fluorescein angiography (depending on the state of the patient's fundi). Early diagnosis of serious retinopathy is vital since the early use of laser photocoagulation may delay and prevent visual loss.

Neuropathy

The following types of neuropathy may occur:

- rediculopa thy (diabetic lumbosacral radiculoplex oplexopathy) sensory polyneuropathy
- isolated mononeuropathy and multiple mononeuropathy
  - isolated peripheral nerve lesions (e.g. median nerve)
  - cranial nerve palsies (e.g. III, VI)
  - amyotrophy
- autonomic neuropathy, which may leed to
  - 0erectile dysfunctim
  - postural hypotension & synwlx
  - Inpaired gastic emptymg(gostroparesis)
  - Diarrhoea
  - Ddelgyed or inconplete bladder empty
  - Loss of cardiac pain® 'silent ischee
  - Hypoglycaemia ‘unawareness’
  - Suddeuarrest, oppeceall under Andesthetic

Infections

Poorly controlled diabetics are prone to infections, especially:

- skin
  - mucocutaneous candidiasis (e.g. balanitis, vulvovaginitis)
  - staphylococcal infections (e.g. folliculitis)
- urinary tract

  - cystitis (women)
  - pyelonephritis and perinephric abscess
- lungs
  - pneumonia: staphylococcal, streptococcal pneumonia, others
  - tuberculosis

Diabetic metabolic complications

- Hypoglycaemia
- Diabetic ketoacidosis
- Hypermolar hyperglycaemia
- Lactic acidosis

Other complications

- Cataracts
- Refractive errors of eye
- Sleep apnoea
- Depression
- Musculoskeletal
  - Neuropathic joint damagy (charcots type arthropathy)
  - Tendon rupture
- Foot wkers (related to neunopa thy)

PREVENTION OF DIABETES

Several large studies have demonstrated that it is possible to prevent or delay the onset of diabetes in those at risk.[13, 14] It involves intensive lifestyle intervention in individuals who are overweight with impaired glucose tolerance or raised fasting bloodglucose. One strategy is to follow the SNAP guidelines (Smoking, Nutrition, Alcohol, Physical activity). The essentials were healthy eating, weight loss and physical activity. This represents an important approach that GPs can recommend to their patients at risk.

REFERENCES

1. UKPDS Group. Complications in newly diagnosed type 2 diabetic patients and their association with different clinical and biochemical risk factors. Diabetes Res, 1990; 13: 1 - 11.
2. Dunstan D, Zimmet P, Welborn T et al. on behalf of the AusDiab Steering Committee. *Diabetes and Associated Disorders in Australia 2000: The Accelerating Epidemic. Australian Diabetes, Obesity and Lifestyle Report*. Melbourne: International Diabetes Institute, 2001.
3. Phillips P. *Diabetes*. CHECK Programme, unit 401. Melbourne: RACGP, 2005; 4 - 20.
4. Welborn TA. Diabetic mellitus. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 149 - 52.
5. Coleman PG, Thomas DW, Zimmet P et al. New classification and criteria for the diagnosis of diabetes mellitus. Med J Aust, 1999; 170: 375 - 8.
6. *Diabetes Management in General Practice*. Melbourne: Diabetes Australia & RACGP, 2005 - 6: 21 - 49.
7. Welborn T et al. National diabetic study. Metabolism, 1997; 1: 1 - 3.
8. Shenfield G.(Chair: Writing group). *Therapeutic Guidelines: Endocrinology* (Version 3). Melbourne: Therapeutic Guidelines Ltd, 2004; 2 - 5.
9. Sinah R et al. Diabetes in childhood obesity. NEJM, 2002; 346: 802 - 10.
10. UK Prospective Diabetes Study Group. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes (UKPDS 38). Br Med J, 1998; 317: 703 - 13.
11. Hypertension Expert Working Group. Evidence based guideline for the diagnosis and management of hypertension in type 2 diabetes. National evidence basedguidelines for the management of type 2 diabetes mellitus. Draft for public consultation. Sydney: Australian Centre for Diabetes Strategies, Prince of Wales Hospital, January 2001.
12. Heart Outcomes Prevention Evaluation (HOPE) Study Investigators. Effects of ramipril on cardiovascular andmicrovascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. Lancet, 2000; 355: 253 - 9.
13. Managing type 2 diabetes. NPS News, 2005; 39: 1 - VI.
14. Vaag A, et al. Metabolic impact of a family history of type 2 diabetes. Results from a European Multicentre study. Sisätautien Klinikka,

2001: 65.

CHAPTER 22. DRUG PROBLEMS

A custome loathsome to the eye, hateful to the nose, harmful to the braine, dangerous to the lungs and the blacke stinking fume thereof, neerest resembling the horrible Stigian smoke of the bottomless pit.

JAMES I (1566 - 1625), ON SMOKING

DRUG PROBLEMS

Drug-related problems are true masquerades in family practice. This includes prescribed drugs, over-the-counter drugs and social or illegal street drugs. It is important therefore that all prescribing doctors maintain a high index of suspicion that any clinical problem may be associated with their treatment of the patient.

ADVERSE DRUG REACTIONS

An adverse drug effect is defined as 'any unwanted effect of treatment from the medical use of drugs that occurs at a usual therapeutic dose'. Almost every drug can cause an adverse reaction, which must be elicited in the history. Any substance that produces beneficial therapeutic effects may also produce unwanted, adverse or toxic effects. The severity of the reaction may range from a mild skin rash or nausea to sudden death from anaphylaxis. A study has shown that the incidence of adverse reactions increases from about 3% in patients 10 - 20 years of age to about 20% in patients 80 - 89 years of age.[1]

Reactions can be classified in several ways - side-effects, overdosage, intolerance, hypersensitivity and idiosyncrasy. However, a useful classification of unwanted effects is divided into type A and type B.

Type A reactions are the most common and involve augmented pharmacology; that is, they are caused by unwanted, albeit predictable, effects of the drug. Examples include:

- constipation due to verapamil
- blurred vision and urinary outflow problems due to tricyclic antidepressants
- hyperuricaemia due to thiazide diuretics

Type A reactions are dose-dependent.

Type B reactions are by definition bizarre. The reactions are unpredictable from known properties of the drug. Examples include hepatotoxicity and blood dyscrasias.

Golden rules for prevention of adverse effects

Before prescribing any drug the prescriber should consider the following rules:

1. Is the drug really necessary?
2. What will happen if it is not used?
3. What good do I hope to achieve?
4. What harm may result from this treatment?

Common adverse effects

There is an extensive list of clinical problems caused by drugs as side-effects or interactions that are highlighted throughout this book. Common side-effects include:

- CNS - malaise, drowsiness, fatigue/tiredness, headache, dizziness
- CVS - palpitations, peripheral oedema, hypotension
- GIT - nausea, vomiting, dyspepsia, change in bowel habit (diarrhoea, constipation)
- Skin - rash, pruritus, flushing
- Psychiatric/emotional - insomnia, irritability, anxiety, depression, agitation

Drugs that commonly produce adverse effects

- Antimicrobials
    - penicillin/cephalosporins

  - sulphonamides
  - tetracyclines
  - streptomycin
  - ketoconazole
- Anticonvulsants
  - carbamazepine
  - phenobarbitone
  - phenytoin
  - sodium valproate
- Antidepressants
  - tricyclics
  - MAOIs
- Anti-inflammatories and analgesics
  - aspirin/salicylates
  - codeine/morphine, etc.
  - NSAIDs
  - gold salts
  - DMARDs
  - b DMARDs
- Antihypertensive agents
  - (several)
- Cardiac agents
  - digoxin
  - quinidine
  - amiodarone
  - other antiarrhythmics
- Diuretics
  - thiazides
  - frusemide
- Tranquillisers
  - phenothiazines
  - benzodiazepines
  - barbiturates
  - chlordiazepoxide
- Other drugs
  - cytotoxics
  - hormones
  - allopurinol
  - warfarin

NICOTINE

'Smoking is good for you', according to an old Arab proverb. 'The dogs will not bite you because you smell so bad; thieves will not rob you at night because you cough in your sleep and you will not suffer the indignities of old age because you will die when you are relatively young.'

Tobacco smoking is the largest single, preventable cause of death and disease in Australia. It has been estimated to have caused approximately 20 000 deaths in 1991, over six times the number of deaths from road accidents.[2] Diseases attributed to smoking are summarised in Figure 22.1. Signs of major dependence are smoking within 30 minutes of waking and ≥20 cigarettes a day.


Cancer of
mouth and throat
larynx and trachea
lungs
oesophagus and stomach
pancreas
bladder
Other diseases
brain → stroke
lungs → chronic bronchitis and emphysema
heart → coronary artery disease
stomach and duodenum } peptic ulcers
ovaries ♀
uterus
testes ♂
} reduced fertility
peripheral arteries → vascular disease

Figure 22.1 Possible serious adverse effects of nicotine smoking

GETTING PATIENTS TO QUIT

Several studies have highlighted the value of opportunistic intervention by the family doctor. It is important not only to encourage people to quit but also to organise a quitting program and follow-up. In Australia 80% of smokers (representing about 30% of the adult population) have indicated that they wish to stop smoking. Point out that it is not easy and requires strong will power. As Mark Twain said, ‘Quitting is easy - I've done it a thousand times'.

METHODS

- Educate patients about the risks to their health and the many advantages of giving up smoking, and emphasise the improvement in *health, longevity, money savings, looks* and *sexuality*.
  Point out the following advantages to quitting:
    - Food tastes better.
    - Sense of smell improves.
    - Exercise tolerance is better.

  - Sexual pleasure is improved.
  - Bad breath improves.
  - Risk of lung cancer drops: after 10 - 15 years of quitting it is as low as someone who has never smoked.
  - Early COPD can be reversed.
  - Decreases URTIs and bronchitis.
  - Chance of premature skin wrinkling and stained teeth is less.
  - Removes effects of passive smoking on family and friends.
  - Removes problem of effects on pregnancy.
- The extent of nicotine dependence can be assessed using a questionnaire (based on the Fagerstrom Test) and scoring system.[3] As a baseline ask about the number of cigarettes smoked per day, how soon after waking to smoking the first cigarette of the day and any difficulties with coping with antismoking venues (e.g. cinemas, plane travel).

Intervention: the 5A framework

- *Ask* about and document tobacco use at every opportunity.
- *Assess* motivation and confidence to quit: 'Are you interested in quitting?'
- *Advise* all smokers to stop.
- *Assist* the smoker to stop.
- *Arrange* followup to maintain non-smoking.
- Ask them to keep a smoker's diary.
- If they say no to quitting, give them motivational literature and ask them to reconsider.
- If they say yes, make a contract (example below).

A contract to quit

'I ______________ agree to stop smoking on ______________ I understand that stopping smoking is the single best thing I can do for my health and that my doctor has strongly encouraged me to quit.'

______________ (Patient's signature)

______________ (Doctor's signature)

- These motivated patients will require educational and behavioural strategies to help them cope with quitting. Ongoing support by their GP is very important.
- Organise joining a support group.
- Arrange follow-up (very important), at least monthly, especially during first 3 months.
- Going 'cold turkey' (stopping completely) is preferable but before making the final break it can be made easier by changing to a lighter brand, inhaling less, stubbing out earlier and reducing the number. Changing to cigars or pipes is best avoided.

QUITTING TIPS (ADVICE TO PATIENT)

- Make a definite date to stop (e.g. during a holiday).

After quitting

- Eat more fruit and vegetables (e.g. munch carrots, celery and dried fruit).
- Foods such as citrus fruit can reduce cravings.
- Chew low-calorie gum and suck lozenges.
- Increase your activity (e.g. take regular walks instead of watching television).
- Avoid smoking situations and seek the company of non-smokers.
- Drink more water and avoid substituting alcohol for cigarettes.
- Be single-minded about not smoking - be determined and strong.
- Take up hobbies that make you forget smoking (e.g. water sports).
- Put aside the money you save and have a special treat. You deserve it!!

WITHDRAWAL EFFECTS

The initial symptoms are restlessness, cravings, hunger, irritability, poor concentration, headache, tachycardia, insomnia, increased cough, tension, depression, dysphonia, tiredness and sweating. After about 10 days most of these effects subside but it takes about three months for a smoker to feel relatively comfortable with not smoking any more. Nicotine replacement therapy certainly helps patients cope.

PHARMACOLOGICAL TREATMENT

Nicotine replacement therapy (NRT), which should be used in conjunction with an educational support program, has been proved to be effective and is available as chewing gum, inhaler or transdermal patches (the preferred method). Ideally the nicotine should not be used longer than 3 months. Eight weeks of patch therapy is as effective as longer courses.

NRT should be directed at smokers who are motivated to quit. There is little evidence that drug treatment will benefit individuals with low levels of nicotine dependence who smoke fewer than 10 cigarettes a day.[4]

All forms of NRT are effective: a pooled analysis of all NRT trials showed an absolute increase in cessation at 1 year of 7% compared to placebo.[4]

Nicotine gum[3]

This is available as 2 mg and 4 mg.

- Low dependence (less than 10 cigarettes per day): use non-pharmacological methods rather than replacement.
- Moderate dependence (10 - 14 cigarettes per day): 2 mg, chew 8 - 12 pieces daily.
- High dependence (> 15 per day): 4 mg initially, 6 - 10 pieces chewed daily changing to 2 mg after 4 - 6 weeks.

USEFUL POINTS

- Chew each piece slowly for about 30 minutes.
- Ensure all the nicotine is utilised.
- Chew at least 6 pieces per day, replacing at regular intervals (not more than 1 piece per hour).
- Use for 3 months, weaning off before the end of this period.

Transdermal nicotine[3]

This is available as 16-hour or 24-hour nicotine patches in three different strengths. The patients should stop smoking immediately on use.

RECOMMENDATIONS

- Low dependence (10 - 20 cegarettes/day): 14 mg 24hour or long/16 hour patch, daily; aim to cease within 12 weeks.
- High dependence (> 20/day): 21 mg/24 hour or 15 mg/16 hour patch; change to 14 mg or 10 mg patch after 4 - 6 weeks; aim to cease within 12 weeks.

Apply to non-hairy, clean, dry section of skin on upper outer arm or upper chest and leave in place for 24 hours. Rotate sites with a 7-day gap for reuse of a specific site.

Nocotine inhaler

4mg capsules in a monthpece resembling smoking.

Nocotine lobenges

Logenges are available in 2 mg and 4 mg strengths, the strenth depending on the time of the first smoke after waking.

Combination therapy

Controlled trials have shown enhanced outcomes when nicotine patches are combined with gum or inhaler. Consider it for the highly addicted smokers.

CONTRAINDICATIONS

These are pregnancy and breastfeeding, children, severe myocardial ischaemia, arrhythmias or recent CVA.

Adverse reactions

- *Gum*: hiccoughs, orodental problems, jaw pain, gastrointestinal, including exacerbation of peptic ulcer
- *Patches*: local reaction, sleep disturbances (use 16-hour patch for this)
- *Both*: nervousness, sweating, dry mouth, dyspepsia, abdominal cramps, angina and arrhythmias

Bupropion for smoking cessation[5]

Bupropion (Zyban) was serendipitously discovered to relieve symptoms of cravings or urges to smoke when prescribed as an antidepressant in the United States. In the only comparative trial with NRT (patch) to date, bupropion 300 mg/day for 9 weeks achieved an 8.6% absolute increase in the continuous abstinence rate at 12 months. The findings have to be interpreted with caution but it may have a place in management; however more evidence is needed and NRT is preferred.

Adverse effects include insomnia and dry mouth (both common), while serious effects, such as allergic reactions and seizures, have been reported.[6] It is contralndicated in persons with a history of epilepsy.

Recommended dose: 150 mg (o) daily for 4 days then bd for 7 weeks.

There appears to be minimal gain by combined NRT and bupropion therapy.

Note: Regular follow-up for all methods is essantal for outcome.

ALCOHOL

Refer to page 1245 (Chapter 120) for alcohol dependence.

ILLICIT DRUGS

There are several psychotropic substances that are used for their effects on mood and other mental functions. Many of the severe problems are due to withdrawal of the drug. Symptomatic behaviour common to illicit drugs includes:

- rapid disappearance of clothing and personal belongings from home
- signs of unusual activity around hang-outs and other buildings
- loitering in hallways or in areas frequented by addicts
- spending unusual amounts of time in locked bathrooms
- inability to hold a job or stay in school
- rejection of old friends
- using the jargon of addicts

Newer drugs include ‘crack’, which is a cocaine base where the hydrochloride has mostly been removed, usually in a microwave oven. Crack can be inhaled or smoked. ‘ICE’ is the crude form of methamphetamine, a derivative of amphetamine. ‘Speed’ is dexamphetamine.

Ecstasy is another ‘designer’ drug which is an amphetamine derivative - methylenedioxy-methamphetamine (MDMA). It has high abuse potential, some hallucinogenic properties and a tendency to neurotoxicity, as proved on PET brain scans. It is popular in ‘rave’ parties. Deaths have occurred, reportedly in association with relative dehydration or excessive hydration. A popular party drug, it is implicated as a ‘date rape’ drug. There is no specific antidote.[8]

The drug list

Illicit drugs taken by injecting drug users in Victoria, Australia in 2005 are:

- heroin 89%
- cannabis 87%

- spreed 75%
- ice 29%
- cocaine 15%
- base 13%

(*Source*: Illicit drug reporting system)

A summary of the effects of illicit or 'hard' street drugs is presented in Table 22.1. (on p. 204).

Table 22.1 Illicit substance abuse: a summary of hallmarks

Drug	Physical symptoms	Look for	Dangers
LSD	Severe hallucinations. Feelings of detachment. Incoherent speech. Cold hands and feet. Vomiting. Laughing and crying.	Cube sugar with discolouration in centre. Strong body odour. Small tube of liquid.	Suicidal tendencies. Unpredictable behaviour. Chronic exposure causes brain damage. LSD causes chromosomal breakdown.
Amphetamines	Aggressive or overactive behaviour. Giggling. Silliness. Euphoria. Rapid speech. Confused thinking. No appetite. Extreme fatigue. Dry mouth. Shakiness.	Jars of pills of varying colours. Chain smoking.	Death from overdose. Hallucinations. Methamphetamines sometimes cause temporary psychosis.
Ecstasy	Anxiety, panic, sweating. 'Loving' feelings. Jaw clenching, teeth grinding. Bizarre behaviour, hallucinations, increased heart rate, BP and body temperature.	Small tablets of various colours, shapes, sizes and designs. Also comes in powder and capsules.	Convulsions, risk of death from heart attack, cerebral haemorrhage, hyperthermia, fluid imbalance with hyponatraimia, acute kidney failure, DIC. Liver toxicity.
Barbiturates	Drowsiness. Stupor. Dullness. Slurred speech. Drunk appearance. Vomiting.	Pills of various colours.	Death from overdose or as a result of withdrawal. Addictions. Convulsions.
Narcotics (a) opioids (e.g. heroin)	Stupor/drowsiness. Marks on body. Watery eyes. Loss of appetite. Bloodstain on shirt sleeve. Running nose.	Needle or hypodermic syringe. Cotton. Tourniquet - string. Rope, belt, burnt bottle, caps or spoons. Glassine envelopes.	Death from overdose. Mental deterioration. Destruction of brain and liver. Hepatitis. Embolisms.
(b) cocaine	Similar effects to amphetamines - muscle pains, irritability, paranoia, hyperactive, jerky movements.	Powder: in microwave ovens	Death from overdose - sudden death from arrhythmias. Seizures, mental disorders. Severe respiratory problems.
Phencyclidine (angel dust)	Lack of co-ordination. Feeling of increased physical strength. Hallucinations. Mood disorders.	White powder. Tablets - unbranded. Syringes. Smoked in conjunction with marijuana.	Suicidal tendencies. Death from overdose. Mental disorder. Self-injury.
Marijuana	Initial euphoria. Floating feeling. Sleepiness. Wandering mind. Enlarged eye pupils. Lack of co-ordination. Craving for sweets. Changes of appetite.	Strong odour of burnt leaves. Small seeds in pocket lining. Cigarette paper. Discoloured fingers.	Inducement to take stronger narcotics. Recent medical findings reveal that prolonged usage causes cerebral lesions.
Glue sniffing	Aggression and violence. Drunk appearance, slurred speech. Dreamy or blank expression.	Tubes of glue, glue smears. Large paper or plastic bags or handkerchiefs.	Lung/brain/liver damage. Death through suffocation or choking.

A list of street drugs and their slang names is presented in Table 22.2 (on p. 206).

Table 22.2 A street drug dictionary

Amphetamines or uppers	
• Benzedrine:	roses, beanies, peaches
• Dexedrine:	dexies, speed, hearts, pep pills, fast, go-ee
• Methedrine:	meth, crystals, white light, ice, whiz

• Drinamyl:	purple hearts, goof balls
Amphetamine derivatives	
• **Ecstasy:**	E, eggs, eckies, XTC, 'the love drug', Mitsubishis, MDMA, vitamin E, X
• **Crack:**	crack, split, base/space base(with phehcyclidene)
• **Crank:**	crystal M, crank
Hallucinogens	
• **LSD:**	acid, blue cheer, strawberry fields, barrels, sunshine, pentagons, purple haze, peace pills, blue light, trips
• **Cannabis** (Indian hemp)	
1. Hashish (the resin):	hash, resin
2. Marijuana (from leaves):	pot, tea, grass, hay, weed, locoweed, Mary Jane, rope, bong, jive, Acapulco gold.
• **Cigarettes:**	reefers, sticks, muggles, joints, spliffies, head, smoke, gunga
• **Smoking pot:**	blow a stick, blast a joint, blow, get high, get stoned
Narcotics	
Morphine:	Morph, Miss Emma
Heroin:	H, Big H, Big Harry, GOM (God's own medicine), crap, junk, horse dynamite (high-grade heroin), lemonade (low-grade heroin) *Injection of dissolved powder:* mainlining, blast, smack *Inhalation of powder:* sniffing
Cocaine:	coke, snow, lady of the streets, nose candy, toot, snort, C, flake, rock, blow, vitamin C
H & C:	speed balls
Miscellaneous	
Barbiturates:	devils, barbies, goof balls
Benzodiazepines	Rowies, moggies
Solvents	Chroming

NARCOTIC DEPENDENCE

This section will focus on heroin dependence.

Typical profile of a heroin-dependent person[7]

- Male or female: 16 - 30 years.
- Family history: often severely disrupted, such as parental problems, early death, separation, divorce, alcohol or drug abuse, sexual abuse, mental illness, lack of affection.
- Personal history: low threshold for toleration, unpleasant emotions, poor academic record, failure to fulfil aims, poor self-esteem.
- First experiments with drugs are out of curiosity, and then regular use follows with loss of job, alienation from family, finally moving into a 'drug scene' type of lifestyle.

Methods of intake

1. Oral ingestion
2. Inhalation
    - intranasal
    - smoking
3. Parenteral

- subcutaneous
- intramuscular
- intravenous (Figure 22.2)



Figure 22.2 Intravenous heroin injection signs: linear tracks and scarring from repeated venepunctures along the course of a vein. Less common sites are the lower leg, dorsum of foot, neck and dorsal vein of penis. *Photo courtesy John Jagoda*

Withdrawal effects[7, 8]

These develop within 12 hours of ceasing regular usage. Maximum withdrawal symptoms usually occur between 36 and 72 hours and tend to subside after 10 days.

- Anxiety and panic
- Irritability
- Chills and shivering
- Excessive sweating
- 'Gooseflesh' (cold turkey)
- Loss of appetite, nausea (possibly vomiting)
- Lacrimation/rhinorrhoea
- Tiredness/insomnia
- Muscle aches and cramps
- Abdominal colic
- Diarrhoea

A secondary abstinence syndrome is identified[7] at 2-3 months and includes irritability, depression and insomnia.

Complications

Medical

- Acute heroin reaction: respiratory depression - may include fatal cardiopulmonary collapse. There is an alarming increase in opioid deaths (including methadone).
- Injection site: scarring, pigmentation, thrombosis, abscesses, ulceration (especially with barbiturates).
- Distal septic complications: septicaemia, infective endocarditis, lung abscess, osteomyelitis, ophthalmitis.
- Viral infections: hepatitis B, hepatitis C (refer to Chapter 58), HIV infection (refer to Chapter 28).
- Neurological complications: transverse myelitis, nerve trauma.
- Physical disability: malnutrition.

Social

Alienation from family, loss of employment, loss of assets, criminal activity (theft, burglary, prostitution, drug trafficking).

Management

Management is complex because it includes the medical management not only of physical dependence and withdrawal but also of the individual complex social and emotional factors. The issues of impaired liver function, hepatitis B and C and HIV prevention also have to be addressed. Sociological tests for these illnesses should be considered.

Patients should be referred to a treatment clinic and then a shared care approach can be used. The treatments include cold turkey with pharmacological support, acupuncture, mega-doses of vitamin C, methadone substitution and drug-free community education programs.

Maintenance programs that include counselling techniques are widely used for heroin dependence. Acute toxicity requires injections of naloxone.

Opioid withdrawal[8]

Methadone withdrawal (short term) is used to prevent the emergence of a withdrawal syndrome in contradistinction from methadone maintenance, where there is an extended treatment period.

Initial dose

- buprenorphine 4 - 8 mg (o) as a single daily dose, increasing to 12 mg (max) on day 3, then reduce gradually over the next 3 - 5 days

Note: If autonomic signs, use clonidine 5 - 15 mg/kg/day (o) in 3 divided doses for 7 - 10 days then taper off. If anxiety and agitation, use diazepam 5 - 20 mg (o) qid (with care). Clonidine can be used as first-line treatment. Th ese drugs are preferred to methadone.

Maintenance programs for long-term opioid dependence[8]

There are currently four alternative programs - methadone, buprenorphine, naltrexone and slow-release morphine - which are substitutes for heroin and other opioids. Another agent is the methadone derivative leva-alpha-acetylmethadol.

Methadone

The dose needs to be determined individually according to past use and initial response to methadone.

- Methadone 20 mg (o) daily initially. Stabilise dose over 3 weeks. Beware of doses > 40 mg, especially in unwell patients. Maintenance 50 - 80 mg (o) daily.

Buprenorphine

- Buprenorphine 2 - 4 mg sublingual, once daily initially, increase to 8 - 24 mg daily or alternative days once stabilised. It is less dependent and prone to overdose than methadone but can precipitate withdrawal if used too soon.

Naltrexone

Care is required in giving naltrexone to a person physically dependent on opioids. A naloxone challenge test is used.[3] If no evidence of withdrawal give:

- Naltrexone 25 mg (o) initially increasing to 50 mg daily on day 2 if tolerated. Careful supervision with appropriate counselling is required.

The natural history of opioid dependence indicates that many patients do grow through their period of dependence and, irrespective of treatments provided, a high percentage become rehabilitated by their mid-thirties.

STIMULANT SUBSTANCE ABUSE

The stimulants include amphetamines and their analogues, ephedrine, cocaine and certain appetite suppressants.

Stimulant-induced syndrome[8]

- Aggressive behaviour
- Paranoid behaviour
- Irritability
- transient toxic psychosis
- Delirium

- Schizophrenic-like syndrome

Treatment

- withdrawal of drugs
- chlorpromazine 200 - 600 mg (o) daily for short term

Stimulant-withdrawal syndrome[8]

This syndrome should be suspected in people whose occupation involves shift work, interstate transport driving or multiple jobs presenting with the following symptoms:

- drowsiness
- hypersomnia, then insomnia
- irritability
- aggressive behaviour
- dysphoria
- urge to resume drugs

Treatment

- psychological support and encouragement
- desipramine (or similar tricyclic antidepressant) 75 mg (o) nocte (increasing as necessary)
- bromocriptine 1.25 mg (o) bd has also been used for cocaine withdrawal

HALLUCINOGEN ABUSE

Hallucinogens in use include lysergic acid (LSD), phencyclidine (angel dust), diethylamide and many synthetics. Symptoms include psychotic behaviour, including severe hallucinations. Withdrawal from these drugs is not usually a problem but ‘flashbacks’ can occur. Treatment, specially with fear or anxiety, is diazepam 10 - 20 mg (o) statim.

Medication to counter symptoms[8]

- Haloperidol 2.5 - 10 mg (o) daily *or*
- Diazepam 10 - 20 mg (o) repeated every 2 hours prn (to max 120 mg daily)

CANNABIS (MARIJUANA)

Cannabis is a drug that comes from the plant *Cannabis sativa* or the Indian hemp plant. It contains the chemical tetrahydrocannabinol, which makes people get ‘high’. It is commonly called marijuana, grass, pot, dope, hash or hashish. Other slang terms are Acapulco gold, ganga, herb, J, jay, hay, joint, reefer, weed, locoweed, smoke, tea, stick, Mary Jane, Panama red and spliffy (Table 22.2). Marijuana comes from the leaves, while hashish is the concentrated form of the resinous substances from the head of the female plant and can be very strong (it comes as a resin or oil). The drug is usually smoked as a leaf (marijuana) or a powder (hashish), or hashish oil is added to a cigarette and then smoked. The effects of taking cannabis depend on how much is taken, how it is taken, how often, whether it is used with other drugs and on the particular person.[8] The effects vary from person to person. The effects of a small-to-moderate amount include:

- feeling of well-being and relaxation
- decreased inhibitions
- woozy, floating feeling
- lethargy and sleepiness
- talkativeness and laughing a lot
- red nose, gritty eyes and dry mouth
- unusual perception of sounds and colour
- nausea and dizziness
- loss of concentration
- looking ‘spaced out’ or drunk
- lack of co-ordination

- delusions and hallucinations (more likely with larger doses)
- a new form called 'skunk' or 'mad weed' causes paranoia

The effects of smoking marijuana take up to 20 minutes to appear and usually last 2 to 3 hours and then drowsiness follows.[9] The main problem is habitual use with the development of dependence; dependence (addiction) is worse than originally believed.

Long-term use and addiction

The influence of 'pot' has a severe effect on the personality and drive of the users. They lose their energy, initiative and enterprise. They become bored, inert, apathetic and careless. A serious effect of smoking pot is the inability to concentrate and loss of memory. Some serious problems include:

- deterioration of academic or job performance
- anxiety and paranoia
- respiratory disease (more potent than nicotine for lung disease): causes COPD, laryngitis and rhinitis
- often prelude to taking hard drugs
- becoming psychotic (resembling schizophrenia): the drug appears to unmask an underlying psychosis
- impaired ability to drive a car and operate machinery

Withdrawal

Sudden withdrawal produces insomnia, night sweats, nausea, depression, myalgia, irritability and maybe anger and aggression. However, the effects are often mild with recovery within a few days in many but heavy users have a severe withdrawal.

Management

No specific treatment is available.

The best treatment is prevention. People should either not use it or limit it to experimentation. If it is used, people should be prepared to 'sleep it off' and not drive.

ANABOLIC STEROIDS

The apparent positive effects of anabolic steroids include gains in muscular strength (in conjunction with diet and exercise) and quicker healing of muscle injuries. However, the adverse effects, which are dependent on the dose and duration, are numerous.

Adverse effects in women are:

- masculinisation - male pattern beard growth
- suppression of ovarian function
- changes in mood and libido
- hair loss

In adult men, adverse effects are:

- feminisation: enlarged breasts, high-pitched voice
- acne
- testicular atrophy and azoospermia
- libido changes
- hair loss

Severe effects with prolonged use include:

- liver function abnormalities, including hepatoma
- tumours of kidneys, prostate
- heart disease

In prepubescent children there can be premature epiphyseal closure with short stature.

DRUGS IN SPORT

It is important for general practitioners to have a basic understanding of drugs that are banned and those that are permissible for elite sporting use. The guidelines formulated by the International Olympic Committee (IOC) Medical Commission and the World Anti-doping agency are generally adopted by most major sporting organisations.[10] Tables 22.3 and 22.4 provide useful guidelines. The IOC's list of prohibited drugs is regularly revised. Banned classes of drugs include stimulants, narcotics, anabolic agents, diuretics and various hormones. Banned methods include blood doping (the administration of blood, red blood cells and related blood products) enhancement of oxygen transfer (e.g. erythropoitin, efaproxinal), give droping and pharmaceutical, chemical and physical manipulation (substances or methods that alter the integrity and validity of the urine testing).

Table 22.3 Prohibited classes of substances with examples: World Anti-Doping agency 2004[10]

Classes	Examples
A Stimulants	Amiphenazole, amphetamines, terbutaline, *salmeterol, *salbutamol, *selegiline, pseudoephedrine, phenylpropanolamine
B Narcotics**	Diamorphine (heroin), methadone, morphine, pethidine, pentazocine
C Anabolic agents	Methandienone, nandrolone, stanozolol, testosterone, oxandrolone, DHEA, tetrahydrogestrinone, clenbuterol, zeranol
D Diuretics (Masking agents)	Acetazolamide, frusemide, hydrochlorothiazide, triamterene, indapamide, spironolactone (and related substances)
E Peptide and glycoprotein hormones and analogues	Growth hormone, corticotrophin, chorionic gonadotrophin (in males), erythropoietin, insulin *Note*: masking agents such as probenecid epitest osterone and plasma expandes are banned
Classes subject to certain restrictions	
A Alcohol	Restricted in certain sports (refer to regulations)
B Cannabinoids	Prohibited in competition (refer to regulations)
C Local anaesthetics	Most agents permissible except cocaine: route restricted to local or intra-articular injection
D Corticosteroids	Route of administration restricted to topical inhalation, local or intra-articular injection
E Beta-blockers	Restricted in certain sports

* Permitted by inhaler but only with permission. Caffeine is now permitted.

** Caffeine, codeine, dextromethorphan, dextropropoxyphene, dihydrocodeine, diphenoxylate and pholcodeine are permitted.

Source: < http//www.wada-ama.org/ > (world anti-doping agency)

Table 22.4 Guidelines for treatment of specific conditions: International Olympic Committee Medical Code 1996

Asthma	Allowed	Salbutamol inhaler, salmeterol inhaler, terbutaline inhaler, formoterol inhaler
	Banned	Sympathomimetic products (e.g. ephedrine, pseudoephedrine, isoprenaline, systemic beta-2 agonists)
Cough	Allowed	All antibiotics, steam and menthol inhalations, cough mixtures containing antihistamines, pholcodine, dextromethorphan, dihydrocodeine
	Banned	Sympathomimetic products (e.g. ephedrine, phenylpropanolamine)
Diarrhoea	Allowed	Diphenoxylate, loperamide, products containing electrolytes (e.g. Gastrolyte)
	Banned	Products containing opioids (e.g. morphine)
Hayfever	Allowed	Antihistamines, nasal sprays containing a corticosteroid or antihistamine, sodium cromoglycate preparations
	Banned	Products containing ephedrine, pseudoephedrine

Pain	Allowed	Aspirin, codeine, dihydrocodeine, ibuprofen, paracetamol, tramadol, all NSAIDs
	Banned	Products containing opioids (e.g. morphine) or caffeine
Vomiting	Allowed	Domperidone, metoclopramide

Restricted drugs include alcohol, marijuana, local anaesthetics, corticosteroids and betablockers. Practitioners can check the guidelines and provide written notification to the relevant authority.

REFERENCES

1. Kumar PJ, Clark ML. *Clinical Medicine* (2nd edn). London: Bailliere Tindall, 1990: 733 - 40.
2. Holman CDJ. *The Quantification of Drug-caused Morbidity and Mortality in Australia*. Canberra: Commonwealth Department of Community Services & Health, 1988.
3. Smith A (Chairman). *Therapeutic Guidelines: Cardiovascular* (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2003: 29 - 37.
4. Silagy C et al. Nicotine replacement therapy for smoking cessation (Cochrane review). In: The Cochrane Library, Issue 1, 2002. Oxford: Update software.
5. Managing drug and alcohol problems in primary care. NPS News, 2002; 22.
6. Murtagh JE. Alcohol abuse in an Australian community. Aust Fam Physician, 1987; 16: 20 - 5.
7. Jagoda J. *Drug Dependence and Narcotic Abuse: Clinical Consequences*. Course Handbook. Melbourne: Monash University of Community Medicine, 1987: 66 - 71.
8. Dowden J (Chair). *Therapeutic Guidelines: Psychotropic* (Version 5). Melbourne: Therapeutic Guidelines Ltd, 2003: 106 - 13.
9. Goldman L. *Handbook on Alcohol and Other Drug Problems for Medical Practitioners*. Canberra: Australian Government Publishing Service, 1991: 35.
10. Fricker PA. The anti-doping code in sport - update for 2004, Aust Prescriber, 2004: 27(4): 84 - 6.

CHAPTER 23. ANAEMIA

There's never none of these demure boys come to any proof; for their drink doth so over cool their blood, and making many fish-meals, that they fall into a kind of male green-sickness.

WILLIAM SHAKESPEARE (1564 - 1616), *KING HENRY IV*

ANAEMIA

Anaemia is a label, not a specific diagnosis. Anaemia is defined as a reduction in red blood cell numbers or a haemoglobin (Hb) level below the normal reference level for the age and sex of that individual. It is regarded as a masquerade because the problem can develop surreptitiously and the patient may present with many seemingly undifferentiated symptoms before the anaemia is detected. Once identified, a cause must be found.

KEY FACTS AND CHECKPOINTS

- In Australia, most people with anaemia will have iron deficiency ranging from up to 5% for children to 20% for menstruating females.[1]
- The remainder will mainly have anaemia of chronic disorders.
- The incidence of haemoglobinopathy traits, especially thalassaemia, is increasing in multicultural Western societies.
- If a patient presents with precipitation or aggravation of myocardial ischaemia, heart failure or intermittent claudication, consider the possibility of anaemia.
- The serum ferritin level, which is low in cases of iron-deficiency anaemia, is probably the best test to monitor iron-deficiency anaemia as its level reflects the amount of stored iron.
- Normal reference values for peripheral blood are presented in Table 23.1.

Table 23.1 Normal reference values for peripheral blood - adults

	Male	Female
Haemoglobin (g/L)	130 - 180	115 - 165
Red cells (×10^{12}/L)	4.5 - 6	4 - 5.5
PCV (haematocrit)	40 - 53	35 - 47
MCV (fL)		80 - 95 to 100
Platelets (×10^{9}/L)		150 - 400
White cell count (×10^{9}/L)		4 - 11
Neutrophils		2.5 - 7.5
Lymphocytes		< 4.5
Monocytes		< 0.8
Eosinophils		< 0.44
Reticulocytes (%)		0.5 - 2
ESR (mm/hour)		< 20

Reproduced with permission from Dr M. Gribble[2]

DxT : fatigue + palpitations + exertional dyspnoea ° anaemia

CLINICAL FEATURES OF ANAEMIA

Patients with anaemia may be asymptomatic. When symptoms develop they are usually nonspecific.

Symptoms

- Tiredness/fatigue
- Muscle weakness
- Headache
- Lack of concentration
- Faintness/dizziness
- Dyspnoea on exertion
- Palpitations
- Angina on effort
- Intermittent claudication
- Pica - usually brittle and crunchy food, e.g. ice (iron-deficiency anaemia)

Signs

Non-specific signs

- Pallor
- Tachycardia
- Systolic flow murmur

If severe

- Ankle oedema
- Cardiac failure

Specific signs

- Jaundice - haemolytic anaemia
- Koilonychia (spoon - shaped nails) - iron-deficiency anaemia

History

The history may indicate the nature of the problem:

- iron deficiency: inadequate diet, pregnancy, GIT loss, menorrhagia, NSAID and anticoagulant ingestion
- folate deficiency: inadequate diet especially with pregnancy and alcoholism, small bowel disease
- vitamin B_{12} deficiency: previous gastric surgery, ileal disease or surgery, pernicious anaemia, selective diets (e.g. vegetarians, fad)
- haemolysis: abrupt onset anaemia with mild jaundice
- possibly lead toxicity, especially in children

Classification of anaemia

The various types of anaemia are classified in terms of the red cell size - the mean corpuscular volume (MCV):

- microcytic - MCV ≤ 80 fL
- macrocytic - MCV > 98 fL
- normocytic - MCV 80 - 98 fL

Note: Upper limit of MCV varies from 95 - 100 fldepending on age and laboratory.

Table 23.2 outlines a classification of some of the more common causes of anaemia encountered in general practice.

There can be an interchange of disorders between the above groups, for example, the anaemia of chronic disorders (chronic infection, inflammation and malignancy) can occasionally be microcytic as well as normocytic; the anaemia of hypothyroidism can be macrocytic in addition to the more likely normocytic; the anaemia of bone marrow disorder or infiltration can also be occasionally macrocytic.

Table 23.2 Selected causes and investigations of anaemia

Causes/classification	Primary diagnostic feature	Secondary investigations
Microcytic (MCV < 80 fL)		
Iron deficiency	s.Fe ↓ s.ferr ↓ transferrin ↑	Therapeutic trial of iron; GIT evaluation for blood loss
Haemoglobinopathy (e.g. thalassaemia)	s.Fe N; s.ferr N or ↑	Haemoglobin investigation, e.g. electrophoresis
Sideroblastic anaemia (hereditary)	s.Fe N; s.ferr N or ↑	Bone marrow examination
Occasionally microcytic		
Anaemia of chronic disease (sometimes microcytic)	s.Fe ↓ s.ferr N or ↑ transferrin ↓	Specific for underlying disorder
Macrocytic (MCV > 98 fL)		
(a) With megaloblastic changes		
Vitamin B_{12} deficiency	s.B12 ↓; rc.Fol N or ↑	IF antibody assay; Schilling test
Folate deficiency	s.B_{12} N; rc.Fol ↓	Usually none
Cytotoxic drugs	Appropriate setting; s.B_{12} N; rc.Fol N	None
(b) Without megaloblastic changes		
Liver disease/alcoholism	Appropriate setting; uniform macrocytosis; s.B_{12} N; rc.Fol N	Liver function tests
Myelodysplastic disorders (including sideroblastic anaemia)	Specific peripheral blood findings; s.B_{12} N; rc.Fol N	Bone marrow examination
Normocytic (MCV 80 - 98 fL)		
Acute blood loss/occult	Isolated anaemia; Retic ↑	Dictated by clinical findings
Anaemia of chronic disease[1]	Appropriate setting; Retic ↓	s.Fe ↓ and s.ferr N or ↑
Haemolysis	Specific red cell changes; Retic ↑	s.Bil and s.LDH ↑ s.hapt ↓ specific tests for cause
Chronic kidney disease	Isolated anaemia; Retic ↓	Kidney function
Endocrine disorders (e.g. hypothyroidism)	Appropriate setting; isolated anaemia; Retic ↓	Specific endocrine investigation

Abbreviations: MCV ° mean corpuscular volume; s.Fe ° serum iron; s.ferr ° serum ferritin; s.B_{12} ° serum vitamin B_{12}; rc.Fol ° red cell folate; IF ° intrinsic factor; Retic ° reticulocyte count; s.Bil ° serum bilirubin; s.LDH ° serum lactate dehydrogenase; s.hapt ° serum haptoglobin; N ° normal; ↓ ° reduced; ↑ ° elevated; GIT ° gastrointestinal *Source*: Adapted from 'Anaemia', *MIMS Disease Index*,[1] with permission of MIMS Australia, a division of MediMedia Australia Pty Limited

MICROCYTIC ANAEMIA - MCV ≤ 80 FL

The main causes of microcytic anaemia are iron deficiency and haemoglobulinopathy, particularly thalassaemia. Consider lead poisoning.

IRON-DEFICIENCY ANAEMIA

Iron deficiency is the most common cause of anaemia worldwide. It is the biggest cause of microcytic anaemia, with the main differential diagnosis of microcytic anaemia being a haemoglobinopathy such as thalassaemia.

An understanding of the interpretation of iron studies is important in management. This is presented in Chapter 16.

Key features

- Microcytic anaemia
- Serum ferritin level low (NR: 20 - 250 μg/L)
- Serum iron level low
- Increased transferrin level
- Reduced transferrin saturation
- Response to iron therapy

Non-haematological effects of chronic iron deficiency

- Angular chelitis/stomatitis
- Glossitis
- Oesophageal webs
- Atrophic gastritis
- Brittle nails and koilonychia

Important causes[1]

Blood loss

- Menorrhagia
- Gastrointestinal bleeding (e.g. carcinoma, haemorrhoids, peptic ulcer, hiatus hernia, GORD, NSAID therapy)
- Frequent blood donations
- Malignancy

Increased physiological requirements

- Prematurity, infant growth
- Adolescent growth
- Pregnancy

Malabsorption

- Coeliac disease
- Postgastrectomy

Dietary

- Inadequate intake
- Special diets (e.g. fad, vegetarianism)

Investigations

Investigations are based on the history and physical examination, including the rectal examination. If GIT bleeding is suspected the faecal occult blood test is not considered very valuable but appropriate investigations include gastroscopy and colonoscopy, small bowel biopsy and small bowel enema.

Haemological investigations: typical findings

- Microcytic, hypochromic red cells
- Anisocytosis (variation in size), poikilocytosis (shape) - pencil-shaped rods
- Low serum iron level
- Raised iron-binding capacity
- Serum ferritin level low (the most useful index)
- Soluble transferrin receptor factor - this factor is increased in iron deficiency, but not in chronic disease. It is very helpful therefore in differentiating iron deficiency from other forms. It is an indirect marker of what is happening in the bone marrow.[2]

The state of the iron stores is assessed by considering the serum iron, the serum ferritin and the serum transferrin levels in combination. Typically, in iron deficiency, the serum iron and ferritin levels are low and the transferrin high, but the serum iron level is also low in all infections - severe, mild and even subclinical - as well as in inflammatory states, malignancy and other chronic conditions. Serum ferritin estimations are spuriously raised in liver disease of all types, chronic inflammatory conditions and malignancy; and transferrin is normally raised in pregnancy. Since each of these estimations can be altered in conditions other than iron deficiency, all three quantities have to be considered together to establish the iron status.[3]

Treatment[2]

- Correct the identified cause.
- Diet - iron-rich foods, vitamin C rich foods (see Table 23.3). Iron is present in meat and legumes as Fe^{+++} and therefore requires gastric acid for conversion to Fe^{++}.
- Iron preparations
  - oral iron (ferrous sulphate 1 - 2 tablets daily for 6 months) e.g. Ferro-Gradumet with orange juice
  - parenteral iron preferably by IV infusion is probably best reserved for special circumstances (there is a risk of an allergic reaction). Avoid blood transfusions if possible.

Table 23.3 Optimal adult diet for iron deficiency

Adults should limit milk intake to 500 mL a day while on iron tablets. Avoid excess caffeine, fad diets and excess processed bread. Eat ample iron-rich foods (especially protein).
Protein foods • Meats - beef (especially), veal, pork, liver, poultry • Fish and shellfish (e.g. oysters, sardines, tuna) • Seeds (e.g. sesame, pumpkin) • Eggs, especially egg yolk
Fruits • Dried fruit (e.g. prunes, figs, raisins, currants, peaches) • Juices (e.g. prune, blackberry) • Most fresh fruit
Vegetables • Greens (e.g. spinach, silver beet, lettuce) • Dried peas and beans (e.g. kidney beans) • Pumpkin, sweet potatoes
Grains

• Iron-fortified breads and dry cereals • Oatmeal cereal
For better iron absorption, add foods rich in vitamin C (e.g. citrus fruits, cantaloupe, brussel sprouts, broccoli, cauliflower).

Response

- Anaemia responds after about 2 weeks and is usually corrected after 2 months.[1]
- Oral iron is continued for 3 to 6 months to replenish stores.
- Monitor progress with regular serum ferritin levels.
- A serum ferritin level > 50 μg/L generally indicates adequate stores.

Failure of iron therapy

Consider:

- poor compliance
- continuing blood loss
- malabsorption (e.g. severe coeliac disease)
- incorrect diagnosis (e.g. thalassaemia minor, chronic disease)
- bone marrow infiltration

THALASSAEMIA

This inherited condition is seen mainly (although not exclusively) in people from the Mediterranean basin, the Middle East, north and central India and South-East Asia, including south China. The heterozygous form is usually asymptomatic; patients show little if any anaemia and require no treatment. The condition is relatively common in people from these areas. The homozygous form is a very severe congenital anaemia needing lifelong transfusional support but is comparatively rare, even among the populations prone to thalassaemia (refer to Chapter 19).[3]

The key to the diagnosis of heterozygous thalassaemia minor is significant microcytosis quite out of proportion to the normal Hb or slight anaemia, and confirmed by finding a raised HbA2 on Hb electrophoresis. DNA screening analysis is now available. The importance of recognising the condition lies in distinguishing it from iron-deficiency anaemia, for iron does not help thalassaemics and is theoretically contraindicated. Even more importantly it lies in recognising the risk that, if both parents have thalassaemia minor, they run a one in four chance of having a baby with thalassaemia major in every pregnancy, with devastating consequences for both the affected child and the whole family.

Treatment of thalassaemia major is transfusion to a high normal Hb with packed cells plus desferrioxamine.

HAEMOGLOBIN E

This Hb variant is common throughout South-East Asia.[2] It has virtually no clinical effects in either the homozygous or heterozygous forms, but these people have microcytosis, which must be distinguished from iron deficiency; moreover, if the HbE gene is combined with the thalassaemia gene, the child may have a lifelong anaemia almost as severe as thalassaemia major. Both genes are well established in the South-East Asian populations in Australia as well as in their own countries.

MACROCYTIC ANAEMIA - MCV > 98 FL

Alcohol and liver disease

Each individually, or in combination, leads to macrocytosis with or without anaemia. The importance of this finding lies in its often being the first indication of alcohol abuse, which can so frequently go unnoticed unless there is a firm index of suspicion. Chronic liver disease due to other causes may also be late in producing specific clinical symptoms.

Drug toxicity

Cytotoxic drugs, anticonvulsants in particular, and various others (see Table 23.4) may cause macrocytosis. This is of little clinical significance and does not need correction unless associated with anaemia or other cytopenia.

Table 23.4 Drugs causing macrocytosis[3]

Alcohol	
Cytotoxics/immunosuppressants	Azathioprine Methotrexate 5-fluorouracil
Antibiotics	Cotrimoxazole Pyrimethamine (incl. Fansidar and Malouprim) Zidovudine
Anticonvulsants	Phenytoin Primidone Phenobarbitone

Myelodysplastic syndromes

These conditions have been recognised under a variety of names, such as ‘refractory anaemia’ and ‘preleukaemia’, for a long time, but only relatively recently have they been grouped together. They are quite common in the elderly but may be seen in any age group (refer Table 23.2 on p. 212).

These conditions frequently present as a macrocytic anaemia with normal serum vitamin B_{12} and red cell folate, and are unresponsive to these or any other haematinics. They are usually associated with progressive intractable neutropenia or thrombocytopenia or both, and progress slowly but relentlessly to be eventually fatal, terminating with infection, haemorrhage or, less often, acute leukaemia.

Vitamin B_{12} deficiency (pernicious anaemia)

Although well recognised, this is a much less common cause of macrocytosis than the foregoing conditions. It is usually caused by lack of intrinsic factor due to autoimmune atrophic changes and by gastrectomy. Anaemia does not develop for about 3 years after total gastrectomy. Vitamin B_{12} deficiency may also be seen together with other deficiencies in some cases of malabsorption and Crohn's disease.

Vitamin B_{12} (cobalamin) is found in the normal diet but only in foods of animal origin and consequently very strict vegetarians may eventually develop deficiency. Causes of food Vitamin B_{12} deficiency are:[2]

- atrophic gastritis
- *H. pylori* infection
- H_2 receptor blockers
- PPI drugs
- other drugs, e.g. OCP, metformin
- chronic alcoholism
- HIV

The clinical features are anaemia (macrocytic), weight loss and neurological symptoms, especially a polyneuropathy. The serum vitamin B_{12} is below the normal level.

B_{12} > 220 pmol/L ° deficiency unlikely < 148 pmol/L ° deficiency

Replacement therapy[1]

- Vitamin B_{12} (1000 µg) IM injection; body stores (3 - 5 mg) are replenished after 10 - 15 injections given every 2 to 3 days.
- Maintenance with 1000 µg injections every third month.
- Can use crystalline oral B_{12}.

Folic acid deficiency

The main cause is poor intake associated with old age, poverty and malnutrition, usually associated with alcoholism. It may be seen in malabsorption and regular medication with antiepileptic drugs such as phenytoin.[4] It is rarely, but very importantly, associated with pregnancy, when the demands of the developing fetus together with the needs of the mother outstrip the dietary intake - the so-called ‘pernicious anaemia of pregnancy’ which, if not recognised and treated immediately, can still be a fatal condition. Unlike vitamin B_{12}, folic acid is not stored in the body to any significant degree and requirements have to be satisfied by the daily dietary intake, which invariably meets the requirement of 5 - 10 µg/day. Folic acid is present in most fruit and vegetables, especially citrus fruits and green leafy vegetables (see p. 82).

Replacement therapy

Oral folate 5 mg/day to replenish body stores (5 - 10 mg) in about 4 weeks.

NORMOCYTIC ANAEMIA[3] (ANAEMIAS WITHOUT CHANGE IN THE MCV)

Acute haemorrhage

This is the most common cause of normocytic anaemia and is usually due to haematemesis and/or melaena.

Chronic disease

Chronic inflammation

Intercellular iron transport within the marrow is suppressed in inflammation so that, despite normal iron stores, the developing red cells are deprived of iron and erythropoiesis is depressed. If the inflammation is short-lived, the fall in Hb is not noticeable but, if it continues, an anaemia may develop that responds only when the inflammation subsides.

Malignancy

Anaemia may develop for the same reasons that apply to chronic inflammation.

Kidney failure

This is often associated with anaemia due to failure of erythropoietin secretion and is unresponsive to treatment, other than by alleviating the insufficiency or until erythropoietin is administered.

Haemolysis

Haemolytic anaemias are relatively infrequent. The more common of the congenital ones are hereditary spherocytosis and deficiencies of the red cell enzymes, pyruvate kinase and G-6-PD, although most cases of G-6-PD deficiency haemolyse only when the patient takes oxidant drugs such as sulphonamides or eats broad beans - ‘favism’.

Acquired haemolytic anaemias include those of the newborn due to maternal haemolytic blood group antibodies passing back through the placenta to the fetus, and adult anaemias due to drug toxicity or to acquired autoantibodies. About half of the latter are ‘idiopathic’ and half associated with non-Hodgkin's lymphomas, and the anaemia may be the presenting sign of lymphoma. Some of these antibodies are active only at cool temperatures - cold agglutinin disease; others act at body temperature and are the more potent cause of autoimmune haemolytic anaemia.

Bone marrow replacement

This may be due to foreign tissue, such as carcinomatous metastases or fibrous tissue as in myelofibrosis; it may also be due to overgrowth by one or other normal elements of the bone marrow, as in chronic myeloid leukaemia, chronic lymphocytic leukaemia and lymphoma, as well as by acute leukaemic tissue. A leuco-erythroblastic picture, in which immature red and white cells appear in the peripheral blood, is often seen when the marrow is replaced by foreign tissue.

ANAEMIA IN CHILDREN

Haemoglobin reference range

- Infant term (cord blood): 135 - 195 g/L
  3 - 6 months: 95 - 135 g/L
- Child 1 year: 105 - 135 g/L
  3 - 6 years: 105 - 140 g/L
  10 - 12 years: 115 - 145 g/L

Important causes of anaemia in childhood include iron-deficiency anaemia (quite common), thalassaemia major, sickle-cell anaemia and drug-induced haemolysis. Consider one of the haemoglobinopathies in children of Mediterranean, South-East Asian, Arabic or African - American descent, especially with a family history, normal ferritin level or anaemia resistant to iron therapy. Investigate with Hb electrophoresis.

Drugs that can cause haemolysis (the film will have reticulocytosis, spherocytosis and fragmented red cells) include some antibiotics (e.g. sulphamethoxazole, antimalarials and some antiinflammatories).

Think of anaemia in adolescents, especially females with a rapid growth spurt at the menarche and a relatively poor diet.

Iron deficiency in children[5]

- Iron deficiency is present in up to 10 - 30% of children in high-risk groups.
- It is often subclinical and anaemia develops in relatively few.
- It can lead to reduced cognitive and psychomotor performance (even without anaemia).
- High-risk groups include those infants < 6 months who are premature and/or with low birth weight; toddlers 6 - 36 months with a diet high in cows milk and low in iron-containing foods; those exclusively breastfed after 6 months; those with delayed introduction of solids; those with general poor food intake; and those with lack of vitamin C in diet. Bottle-feeding encourages a high milk intake and reduces the appetite for solid food.
- Possible symptoms include irritability, lethargy, minor behavioural changes.

Prevention

- Introduce iron-containing solids early - at 4 to 5 months, e.g. cereals, vegetables, egg and meat.
- Avoid cow's milk in first 12 months.[5]
- Avoid excessive cow's milk up to 24 months.
- Use iron-fortified formulas and cereals.

Important sources of iron

Infant milk formulas, meat, especially red meat (also fish and chicken), green vegetables and legumes, dried fruit, juices, fortified cereals.

Treatment

Treatment is mainly with ferrous gluconate (1 mL/kg of 300 mg/5 mL mixture). Continue for 3 months after Hb has normalised.

Practice tips[2]

- Iron-deficiency anaemia is blood-loss anaemia until proved otherwise.
- It is possible to be tired from iron deficiency without anaemia.
- Blood-loss anaemia is usually due to menorrhagia or gastrointestinal loss until proved otherwise.
- Investigations for suspected anaemia should include a FBE, ESR and iron studies. Others to consider are Hb electrophoresis, vitamin B_{12} and folate levels and kidney function tests.
- Hypothyroidism can cause a normocytic or a macrocytic anaemia.
- A therapeutic trial of iron (without investigations) is indefensible.
- Intramuscular injections of iron can tattoo so use with care: an IM iron dose is not ‘stronger’ than an oral iron dose.

REFERENCES

1. Van Der Weyden M. Anaemia. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 26 - 9.
2. Coghlan D, Campbell P. Anaemia: how to treat. Australian Doctor, 8 November 2002: 1 - VIII.
3. Gribble M. Haematology. CHECK Unit 188. Melbourne: CHECK Programme, 1987: 3 - 12.
4. Kumar PJ, Clark ML. *Clinical Medicine* (2nd edn). London: Saunders, 2003: 410 - 22.
5. Smart J et al. *Paediatric Handbook* (6Thedn). Melbourne: Blackwell Science, 2000: 351 - 4.

CHAPTER 24. THYROID AND OTHER ENDOCRINE DISORDERS

> It would indeed be rash for a mere pathologist to venture forth on the uncharted sea of the endocrines, strewn as it is with the wrecks of shattered hypotheses, where even the most wary mariner may easily lose his way as he seeks to steer his bark amid the glandular temptations whose siren voices have proved the downfall of many who have gone before.
>
> William Boyd (1885 - 1969)

THYROID AND OTHER ENDOCRINE DISORDERS

Thyroid disorders can be a diagnostic trap in family practice and early diagnosis is a real challenge. A family practice of 2500 patients can expect one new case of thyroid disorder each year and ten ‘cases’ in the practice.[1] The diagnosis of an overactive or underactive thyroid can be difficult as the early clinical deviations from normality can be subtle.

The clinical diagnosis of classical Graves' is usually obvious with the features of exophthalmos, hyperkinesis and a large goitre but if the eye and neck signs are absent it can be misdiagnosed as an anxiety state. Elderly patients may present with only cardiovascular signs, such as atrial fibrillation and tachycardia, or with unexplained weight loss.

The hypothyroid patient can be very difficult to diagnose in the early stages, especially if the patient is being seen frequently. Hypo-thyroidism often has a gradual onset with general symptoms such as constipation and lethargy.

If suspected, serum thyroid-stimulating hormone (TSH) should be requested.

Other comman endocrine disorders include diabetes mellitus, hyperprolactinaemia, calcium metabolic disorder, PCOS, sexual dyfunction and subclininal hypogonadum. They may be difficult to diagnose in the early stages of development. Certain symptoms (Table 24.1) alert one to the possibility of an endocrine disorder.

Table 24.1 Symptoms related to the endocrine system

Psychogenic changes • depression • anxiety • psychosis
Constitutional symptoms • tiredness, lethargy • weakness
Sexual dysfunction • menstrual dysfunction (e.g. amenorrhoea) • loss of libido • loss of pubic hair • hirsutism
Diabetogenic symptoms • polyuria/polydipsia
Glycaemia fluctuations • hypoglycaemia • hyperglycaemia
Weight changes • weight loss • weight gain
Cardiac changes

• heart rate disturbances • myocardial ischaemia
Skeletal changes • osteoporosis • bone pain
Muscle changes
Hair changes
Skin changes • pigmentation • vitiligo • coarse dry skin • striae
Blood pressure fluctuations • postural hypotension • hypertension
Gastrointestinal • anorexia • nausea • constipation • diarrhoea
Tongue enlargement • hypothyroidism, myxoedema, acromegaly

TESTS FOR THYROID DISORDERS

Thyroid function tests

Advances in technology have allowed the biochemical assessment of thyroid function to change dramatically in recent years with the introduction of the serum free thyroxine (T_4) and the monoclonal TSH assays. With the highly sensitive TSH assays it is now possible to distinguish suppressed TSH levels (as in hyperthyroidism) from low normal levels of TSH. However, the new assays are not foolproof and require inter-pretation in the context of the clinical picture. The serum TSH level is the most sensitive index of thyroid function and is the preferred test for suspected thyroid dysfunction. If necessary recessary repeat TSH in 3-6 months.

Serum tri-iodothyronine (T_3) measurement and serum free thyroxine (T_4) can be useful in suspected T_3 toxicosis where serum T_4 level may be normal, and for monitoring patients with treated thyroid dysfunction.

The relative values are summarised in Table 24.2.

Table 24.2 Summary of thyroid function tests

	TSH	T_4	T_3
Hypothyroidism			
• primary	↑*	↓*	N or ↓ (not useful)
• secondary (pituitary dysfunction)	N or ↓	↓	N or↓(not useful)
Hyperthyroidism	↓*	↑*	↑*
Sick euthyroid	N or ↓	N or ↓	N or ↓

Note: Results similar to hyperthyroidism can occur with acute psychiatric illness.

[*] Main tests.

Thyroid autoantibodies

Raised autoantibodies (antimicrosomal or antithyroid peroxidase) are suggestive of Hashimoto's thyroiditis (autoimmune thyroiditis). Antithyroglobulin and anti-thyried peroxidase and TSH receptor antibodies are elevated in Graves' disease, the TSH receptor antibody being very specific for graves disease.

Fine needle aspiration

This is the single most cost-effective investigation in the diagnosis of thyroid nodules. It is the best way to assess a nodule for malignancy. Care needs to be taken in the interpretation of the cytology results in conjunction with an experienced cytologist/pathologist.

Thyroid isotope scan

The scan may help in the differential diagnosis of thyroid nodules and in hyperthyroidism. A functioning nodule is said to be less likely to be malignant than a non-functioning nodule (cyst, colloid nodule, haemorrhage carcinoma is a usually non-functioning) are not functioning.

Thyroid ultrasound

A thyroid ultrasound is usually more sensitive in the detection of thyroid nodules. A multinodular goitre may be diagnosed on ultrasound while the clinical impression may be that of a solitary nodule (the other nodules not being palpable clinically). A multinodular goitre is said to be less likely to be malignant than a solitary thyroid nodule. An ultrasound allows for follow-up of thyroid nodule(s) to note if there are any changes in size over a period of time and to then discuss appropriate intervention with the patient. It can also differentiate a solid from a cystic mass.

CT scan

CT scan of the thyroid may be used particularly to determine if there is significant compression in the neck from a large multinodular goitre with retrosternal extension. Again follow-up CT scans may allow one to determine the progression or otherwise of such a goitre.

HYPOTHYROIDISM (MYXOEDEMA)

Hypothyroidism, which is relatively common, is more prevalent in elderly women (up to 5%).[2] The term *myxoedema* refers to the accumulation of mucopolysaccharide in subcutaneous tissues. The early changes are subtle and can be misdiagnosed, especially if only a single symptom is dominant.

Patients at risk include those with:

- previous Graves' syndrome
- autoimmune disorders (e.g. rheumatoid arthritis)
- Down syndrome
- Turner syndrome
- drug treatment: lithium, amiodarone
- previous thyroid surgery
- previous radioactive iodine treatment of the thyroid

Clinical features

The main features are:

- constipation
- cold intolerance
- lethargy

- physical slowing
- mental slowing
- huskiness of voice
- puffiness of face and eyes
- pallor
- loss of hair
- weight gain

Physical examination

See Figure 24.1. The main signs are:


mental slowing
physical slowing
loss of hair
xanthelasmas
coarse dry hair
'peaches and cream' complexion
puffiness of eyes and face
large tongue
husky voice
goitre
dry, cool coarse skin
sinus bradycardia
cardiomegaly
cardiac failure
mild obesity
slow, low-volume pulse
constipation
cool peripheries
carpal tunnel syndrome
delayed relaxation of reflexes (ankle jerks)

Figure 24.1 Clinical features of hypothyroidism

- sinus bradycardia
- delayed reflexes (normal muscular contraction, slow relaxation)
- coarse, dry and brittle hair
- dry, cool skin

- obesity
- goitre

Other diverse presentations of thyroid disorders are given in Table 24.3(on p.221).

Table 24.3 Various diverse presentations of thyroid disorders[2]

	Hypothyroidism	Hyperthyroidism
General	Lethargy, tiredness Dry skin Husky voice	Weakness Sweaty skin, especially hands
Psychiatric	Depression Dementia Psychosis (myxoedema madness)	Anxiety/irritability Hyperkinesis Psychosis
Musculoskeletal	Myofibrositis Myalgia Joint effusions	Muscle weakness Proximal myopathy
Skin	Dry cool skin Vitiligo	Warm, thin, soft, moist skin Vitiligo Pretibial myxoedema
Cardiovascular	Ischaemia Cardiomegaly Pericardial effusion Bradycardia Hyperlipidaemia	Tachycardia Atrial fibrillation Heart failure Systolic hypertension
Endocrine	Galactorrhoea Goitre Infertility	Goitre Gynaecomastia
Gynaecological	Menstrual irregularity Menorrhagia (mainly) Oligomenorrhea	Oligomenorrhea
Neurological	Neuropathy Nerve entrapment (e.g. carpal tunnel) Ataxia	Periodic paralysis Tremor
Haematological	Anaemia	-
Emergency	Myxoedema coma Postanaesthetic hypoventilation	Thyroid crisis
Other	Reduced libido Weight gain Cold intolerance	Reduced libido Eye signs Fever (uncommon) Onycholysis Premature grey hair Weight loss

DxT: tiredness + husky voice + cold intolerance ° myxoedema

Hashimoto's thyroiditis (autoimmune thyroiditis)

Hashimoto's thyroiditis, which is an autoimmune thyroiditis, is the commonest cause of bilateral non-thyrotoxic goitre in Australia. Features are:

- bilateral goitre
- classically described as firm and rubbery
- patients may be hypothyroid or euthyroid

Diagnosis is confirmed by a strongly positive antithyroid microsomal antibody titre and/or fine needle aspiration cytology.[3]

Hashimoto's thyroiditis may present as postpartum hypothyroidism. The hypo-thyroidism may resolve in 6 - 12 months or may be permanent.[3]

Laboratory diagnosis of hypothyroidism

Thyroid function tests (Table 24.2):

- T_4 - subnormal
- TSH - elevated

If T_4 is low and TSH is low or normal, consider pituitary dysfunction (secondary hypothyroidism) or sick euthyroid syndrome.

Other abnormal tests

- Serum cholesterol level elevated
- Anaemia
    - usually normocytic
    - may be macrocytic
- ECG
    - sinus bradycardia, low voltage, flat T waves

Management principles

Confirm the diagnosis, provide appropriate patient education, and refer the patient where appropriate.

Exclude coexisting hypoadrenalism and ischaemic heart disease before T_4 replacement.

Note:Treatment as primary hypothyroidism when hypopituitarism is the cause may precipitate adrenal crisis.

Thyroid medication

- Thyroxine 100 - 150 µg daily (once daily)

Note: Start with low doses (25 - 50 µg daily) in elderly and those with ischaemic heart disease and 50 - 100 µg in others. Avoid overdosage.

- Aim to achieve TSH levels of 0.5 - 2 mU/L.
- Monitor TSH levels monthly at first. As euthyroidism is achieved, monitoring may be less frequent (e.g. 2 - 3 months). When stable on optimum dose of T_4, monitor every 2 - 3 years. Treatment is usually lifelong.

Special treatment considerations

- *Ischaemic heart disease*. Rapid thyroxine replacement can precipitate myocardial infarction, especially in the elderly.
- *Pregnancy and postpartum*. Continue thyroxine during pregnancy; watch for hypothyroidism (an increased dose of T_4 is after required).
- *Elective surgery*. If euthyroid, can stop thyroxine for one week. If subthyroid, defer surgery until euthyroid.
- *Myxoedema coma*. Urgent hospitalisation under specialist care is required. Intensive treatment is required, which may involve parenteral T_4 or T_3.

Neonatal hypothyroidism

Misdiagnosing this serious condition leads to failure to thrive, retarded growth and poor school performance. If untreated it leads to permanent intellectual damage (cretinism). The clinical features of the newborn include coarse features, dry skin, supra-orbital oedema, jaundice, harsh cry, slow feeding and umbilical hernia. It is detected by routine heel prick blood testing. Thyroxine replacement should be started as soon as possible.

When to refer[2]

- Doubt about diagnosis, diagnostic tests or optimum replacement dosage
- Apparent secondary hypothyroidism, severe illness and associated ischaemic heart disease
- Concurrent autoimmune disease
- Hypothyroidism with goitre, postpartum thyroid dysfunction and in the neonate
- Myxoedema coma

HYPERTHYROIDISM

Hyperthyroidism (thyrotoxicosis) is also relatively common and may affect up to 2% of women, who are affected four to five times more often than men. In elderly women the most common cause is nodular thyroid disease, while Graves' syndrome occurs mainly before middle age.

Causes of thyrotoxicosis[4, 5]

- Graves' syndrome (typical symptoms with a diffuse goitre and eye signs)
- Autonomous functioning nodules
- Subacute thyroiditis (de Quervain's thyroiditis) - viral origin
- Excessive intake of thyroid hormones - thyrotoxicosis factitia
- Iodine excess
- Amiodarone (beware of this drug)

DxT: anxiety + weight loss + weakness ° thyrotoxicosis

Important notes

- The classic symptoms may be lacking in elderly patients who may have only cardiovascular manifestations (e.g. unexplained heart failure or cardiac arrhythmias).
- Care has to be taken not to dismiss hyperthyroidism as severe anxiety.

Clinical features

- Heat intolerance
- Sweating of hands
- Muscle weakness
- Weight loss despite normal or increased appetite
- Emotional lability, especially anxiety
- Palpitations
- Loose bowel motions

Physical examination

See Figure 24.2. Signs are:


irritability
weight loss
hyperkinesis
heat intolerance
exophthalmos
lid lag
conjunctival oedema
proximal muscle wasting
goitre
tachycardia
or atrial fibrillation
systolic hypertension
collapsing pulse
warm, sweaty palms
vasodilated peripheries
fine tremor
brisk tendon reflexes
pretibial myxoedema

Figure 24.2 Clinical features of hyperthyroidism

- agitated restless patient
- warm and sweaty hands
- fine tremor (place paper on hands)
- goitre
- proximal myopathy
- hyperactive reflexes
- bounding peripheral pulse
- ± atrial fibrillation

Eye signs

- Lid retraction (small area of sclera seen above iris)
- Lid lag
- Exophthalmos
- Ophthalmoplegia in severe cases

Investigations

- T_4 (and T_3) elevated
- TSH level suppressed
- Radioisotope scan

The isotope scan enables a diagnosis of Graves' syndrome to be made when the scan shows uniform increased uptake. Increased irregular uptake would suggest a toxic multinodular goitre, while there is poor or no uptake with de Quervain's thyroiditis and thyrotoxicosis factitia.

Management principles

- Establish the precise cause before initiating treatment.
- Educate patients and emphasise the possibility of development of recurrent hyperthyroidism or hypothyroidism and the need for lifelong monitoring.
- Monitor for cardiovascular complications and osteoporosis.

Treatment modalities[5, 6]

- Radioactive iodine therapy (^{131}I)
- Thionamide antithyroid drugs (initial doses)
  - carbimazole 10 - 45 mg (o) daily
  - propylthiouracil 200 - 600 mg (o) daily
- Adjunctive drugs
  - beta-blockers (for symptoms in acute florid phase, e.g. propranolol 10 - 40 mg, 6 to 8 hourly)
  - lithium carbonate (rarely used when there is intolerance to thionamides)
  - Lugol's iodine: mainly used prior to surgery
- Surgery
  - subtotal thyroidectomy *or*
  - total thyroidectomy

Treatment of Graves' syndrome

There is no ideal treatment, and selection of antithyroid drugs, radioiodine or surgery depends on many factors, including age, size of goitre, social and economic factors and complications of treatment.

Guidelines[4,5]

- Younger patients with small goitres and mild case - 18-month course antithyroid drugs.
- Older patients with small goitres - as above or radioiodine (preferably when euthyroid).
- Large goitres or moderate-to-severe cases - antithyroid drugs until euthyroid, then surgery. or^{131} I.
- In Australia (as in the US) ^{131}I is being increasingly used.

Treatment of autonomous functioning nodules

Control hyperthyroidism with antithyroid drugs, then surgery or ^{131}I. Long-term remissions on antithyroid drugs in a toxic nodular goitre are rare.

Treatment of subacute thyroiditis[6]

Hyperthyroidism is usually transient and follows a surge of thyroxine after a viral-type illness. Symptoms include pain and/or tenderness over the goitre (especially on swallowing) and fever. In the acute phase treatment is based on rest, analgesics (aspirin 600 mg (o) 4 - 6 hourly) and soft foods. Rarely, when pain is severe, corticosteroids may be used. Antithyroid drugs are not indicated but beta-blockers can be used to control symptoms.

Thyroid crisis (thyroid storm)[6]

Clinical features are marked anxiety, weight loss, weakness, proximal muscle weakness, hyperpyrexia, tachycardia (> 150 per minute), heart failure and arrhythmias. It is usually precipitated by surgery or an infection in an undiagnosed patient.

It requires urgent intensive hospital management with antithyroid drugs; IV saline infusion, IV corticosteroids, anti-heart failure and antiarrhythmia therapy.

When to refer[4]

- Doubt about the diagnosis
- Severe hyperthyroidism, especially if there is coexisting thyrocardiac disease
- Pregnant patients with hyperthyroidism
- Progression of exophthalmos

THYROID CARCINOMA

The main presentations are a painless nodule, a hard nodule in an enlarged gland or lymphadenopathy. Papillary carcinoma is the most common malignancy. Although rare compared with non-malignant lesions (such as colloid nodules, cysts, haemorrhage and benign adenomas), it is important not to miss carcinoma because of the very high cure rate with treatment. This involves total thyroidectomy, ablative ^{131}I treatment, thyroxine replacement and follow-up with serum thyroglobulin measurements often, I131/thallium scanning and neck ultrasound. Fine needle aspiration is the investigation of choice.

ADRENAL CORTEX DISORDERS

It is worth keeping in mind the uncommon disorders of the adrenal cortex, which can also be difficult to diagnose in the early stages, namely:

- chronic adrenal insufficiency (Addison's syndrome) - deficiency of cortisol and aldosterone
- Cushing's syndrome - cortisol excess
- primary hyperaldosteronism (refer Chapter 126)

Addison's syndrome

Clinical features

- Lethargy/excessive fatigue
- Anorexia and nausea
- Diarrhoea/abdominal pain
- Weight loss
- Dizziness/funny turns
  - hypoglycaemia (rare)
  - postural hypotension (common)
- Hyperpigmentation, especially mucous membranes of mouth and hard palate.

If Addison's syndrome remains undiagnosed, wasting leading to death may occur. Severe dehydration can be a feature.

DxT: fatigue + a/n/v + abdominal pain (± skin discolouration) ° Addison's syndrome

Diagnosis

- Elevated serum potassium, low serum sodium
- Low plasma cortisol level (fails to respond to synthetic adrenocorticotrophic hormone [ACTH])
- The short synacthen stimulation test is the definitive test.

Addisonian crisis

An Addisonian crisis develops because of an inability to increase cortisol in response to stress, which may include intercurrent infection, surgery or trauma.

Clinical features

- Nausea and vomiting
- Acute abdominal pain
- Severe hypotension progressing to shock
- Weakness, drowsiness progressing to coma

Urgent management

- Establish IV line with IV fluids
- Hydrocortisone sodium succinate 200 mg IV and 100-200 mg 4-6 hourly
- Arrange urgent hospital admission

CUSHING'S SYNDROME

The four main causes are:

- iatrogenic - chronic corticosteroid administration
- pituitary ACTH excess (Cushing's disease)
- adrenal tumour (adenoma, adenocarcinoma)
- ectopic ACTH or (rarely) corticotrophin-releasing hormone (CRH) from non-endocrine tumours (e.g. oat cell carcinoma of lung)

The clinical features are caused by the effects of excess cortisol and/or adrenal androgens.



Figure 24.3 62-year-old woman with Cushing's syndrome, showing centripetal obesity, hirsutism and virilisation (wearing a wig). ‘moon’ face, thin extremities and plethoric appearance. *Photo courtesy David Dammery*

Clinical features

- Proximal muscle wasting and weakness
- Central obesity, buffalo hump on neck
- Cushing's facies: plethora, moon face, acne
- Weakness
- Hirsutism
- Abdominal striae

- Thin skin, easy bruising
- Hypertension
- Hyperglycaemia
- Menstrual changes (e.g. amenorrhoea)
- Osteoporosis
- Psychiatric changes, especially depression
- Backache

DxT: plethoric moon face + thin extremities + muscle weakness ° Cushing's syndrome

Diagnosis (apart from iatrogenic cause)

- Cortisol excess (plasma or 24-hour urinary cortisol)
- Dexamethasone suppression test
- Serum ACTH
- Radiological localisation
  - MRI for ACTH-producing pituitary tumours
  - CT scanning for adrenal tumours

OTHER ENDOCRINE DISORDERS

Acromegaly

Symptoms suggestive of acromegaly include:

- excessive growth of hands (increased glove size)
- excessive growth of tissues (e.g. nose, lips, face)
- excessive growth of feet (increased shoe size)
- increased size of jaw and tongue
- general: weakness, sweating, headaches
- sexual changes, including amenorrhoea and loss of libido
- disruptive snoring (sleep apnoea)

DxT: nasal problems + fitting problems (e.g. rings, shoes) + ↑sweating ° acromegaly

Diagnosis[6]

- Plasma growth hormone excess
- Elevated insulin-like growth factor 1 (IGF-1) (somatomedin) - the key test
- MRI scanning pituitary
- Consider associated impaired glucose tolerance/diabetes

Hypopituitarism

This rare disorder should be considered with:

- a history of postpartum haemorrhage
- symptoms of hypothyroidism
- symptoms of adrenal insufficiency
- symptoms suggestive of a pituitary tumour

DxT (♀): amenorrhoea + loss of axillary and pubic hair + breast atrophy ° hypopituitarism

DxT (♂): ↑libido + impotence + loss of body hair ° hypopituitarism

Primary hyperparathyroidism

Hyperparathyroidism is caused by an excessive secretion of parathyroid hormone and is usually due to a parathyroid adenoma. The classic clinical features of hyperparathyroidism are due to the effects of hypercalcaemia. Rarely, a parathyroid crisis in a misdiagnosed patient may result in death from severe hypercalcaemia.

Clinical features

- Kidney
    - polyuria with nocturia (and thirst)
    - Kidney colic
- Musculoskeletal
    - aching legs, especially shins
    - muscle aching
    - muscle weakness
    - back pain
- Gastrointestinal
    - anorexia, nausea
    - constipation
- Psychiatric
    - depression
    - personality changes

> *Classic mnemonic*: bones, moans, stones, abdominal groans

Diagnosis

- Exclusion of other causes of hypercalcaemia
- Serum parathyroid hormone (elevated)
- TC-99m sestamibi scan to detect tumour

Hypercalcaemia

Suspect hypercalcaemia if there is weakness, tiredness, malaise, anorexia, nausea or vomiting, constipation, thirst, polyuria, drowsiness, dizziness, muscle aches and pains, visual disturbances. Measure urea and electrolytes (especially calcium), creatinine, albumin.

Primary hyperparathyroidism, familial hypercalciuric hypercalcaemia and neoplasia, especially lung and breast (with metastases to bone), account for over 90% of cases. Other causes include Paget's disease, sarcoidosis, and milk-alkali syndrome. Investigations include ESR, serum parathyroid hormone, serum ACE levels, serum alkaline phosphatase, chest X-ray sestamibi scan and bone scan.

> **DxT**: weakness + constipation + polyuria ° hypercalcaemia
>
> **DxT**: cramps + confusion + tetany ° hypocalcaemia

Diabetes insipidus and SIADH

Impaired secretion of vasopressin (antidiuretic hormone) from the posterior pituitary leads to polyuria, nocturia and compensatory polydipsia resulting in the passage of 3 - 20 L of dilute urine per day. There are several causes of diabetes insipidus (DI), the commonest being postoperative (hypothalamic-pituitary), which is usually transient only. Other causes of cranial DI include tumours, infections and infiltrations. In nephrogenic DI the renal tubules are insensitive to vasopressin. Differential diagnosis includes compulsive (psychogenic) water drinking. The syndrome of secretion of inappropriate antidiuretic hormone (SIADH), is caused by cancer (e.g. lung, lymphomas, kidney, pancreas), pulmonary disorders, various intracranial lesions and drugs such as carbamezepine and many antipsychotic agents. Management of SIADH is essentially fluid restriction.

The treatment of DI is desmopressin, usually given twice daily intranasally.

DxT: weakness + polyuria + polydipsia ° diabetes insipidus

Hypoparathyroidism

Hypoparathyroidism, causes hypocalcaemia. Causes include postoperative thyroidectomy and parathyroidectomy. congenital deficiency (di George's syndrome), idiopathic (autoimmune) hypoparathyroidism and The main features are neuromuscular hyperexcitability, tetany and neuropsychiatric manifestations.

Two important signs of hypocalcaemia are Trousseau's sign and Chvostek's sign.

Treatment involves careful adjustments in dosage of calcitriol and calcium to correct hypocalcaemia and avoid hypercalcaemia and hypercalciuria (the latter may lead to renal impairment).

REFERENCES

1. Fry J. *Common Diseases* (4th edn). Lancaster: MTP Press Limited, 1985: 358 - 61.
2. Stockigt J, Topliss DJ. Hypothyroidism. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 267 - 9.
3. Yuen R. Common thyroid conditions. Curr Ther, 1992; 10: 23 - 9.
4. Yuen R. *Endocrinology 1*. RACGP: CHECK Programme, unit 243, 1992: 2 - 5.
5. Boyages SC. Thyrotoxicosis. In: *MIMS Disease Index*. Sydney: IMS Publishing, 1996, 507 - 10.
6. Shenfield G (Chair). *Therapeutic Guidelines: Endocrinology* (version 3). Melbourne: Therapeutic Guidelines Ltd, 2004: 57 - 70.

CHAPTER 25. SPINAL DYSFUNCTION

The spine is an ordered series of bones running down your back. You sit on one end of it,
sometimes too hard with ill effect, and your head sits on the other. Poor spine - what a load.

ANON, 19TH CENTURY

SPINAL DYSFUNCTION

Spinal or vertebral dysfunction can be regarded as a masquerade mainly because the importance of the spine as a source of various pain syndromes has not been emphasised in medical training. Practitioners whose training and treatment are focused almost totally on the spine may swing to the other extreme and some may attribute almost every clinical syndrome to dysfunction of spinal segments. The true picture lies somewhere in between.

The diagnosis is straightforward when the patient is able to give a history of a precipitating event such as lifting, twisting the neck or having a motor vehicle accident, and can then localise the pain to the midline of the neck or back. The diagnostic problem arises when the pain is located distally to its source, whether it is radicular (due to pressure on a nerve root) or referred pain. The problem applies particularly to pain in anterior structures of the body.

If a patient has pain anywhere it is possible that it could be spondylogenic and practitioners should always keep this in mind.

The various syndromes caused by spinal dysfunction will be presented in more detail under neck pain, thoracic back pain and lumbar back pain.

CERVICAL SPINAL DYSFUNCTION[1]

The cervical spine is the origin of many confusing clinical problems, such as headache, migraine-like headache, arm pain, facial pain, periauricular pain, anterior chest pain and even visual dysfunction and dizziness. If the cervical spine is overlooked as a source of pain (such as in the head, shoulder, arm, upper chest - anterior and posterior - and around the ear or face) the cause of the symptoms will remain masked and mismanagement will follow.

Dysfunction of the cervical spine can cause many unusual symptoms such as headache and vertigo, a fact that is oft en not recognised. Despite teaching to the contrary from some, the cervical spine is a common cause of headache, especially dysfunction of the facet joints at the C1 - 2 and C2 - 3 levels. The afferent pathways from these levels share a common pathway in the brain stem as the trigeminal nerve, hence the tendency for pain to be referred to the head and the face (see p. 657).

Manipulation of the cervical spine can be a dramatically effective technique, but it should be used with care and never used in the presence of organic disease and vertebrobasilar insufficiency. It should, therefore, be given only by skilled therapists. Two groups at special risk from quadriplegia are those with rheumatoid arthritis of the neck and Down syndrome, because of the instability of the odontoid process.

However, good results can be achieved by gentler techniques, such as mobilisation and muscle energy therapy (refer Chapter 62).

THORACIC SPINAL DYSFUNCTION

The most common and difficult masquerades related to spinal dysfunction occur with disorders of the thoracic spine (and also the low cervical spine), which can cause vague aches and pains in the chest, including the anterior chest.

Pain in the thoracic spine with referral to various parts of the chest wall and upper abdomen is common in all ages and can closely mimic the symptoms of visceral disease, such as angina pectoris and biliary colic. If a noncardiac cause of chest pain is excluded, then the possibility of referral from the thoracic spine should be considered in the differential diagnosis. People of all ages can experience thoracic problems and it is surprisingly common in young people, including children.

Pain of thoracic spinal origin may be referred anywhere to the chest wall, but the commonest sites are the scapular region, the paravertebral region 2 - 5 cm from midline and, anteriorly, over the costochondral region.

Thoracic pain of lower cervical origin [2]

The clinical association between injury to the lower cervical region and upper thoracic pain is well known, especially with ‘whiplash’; injuries. It should be noted that the C4 dermatome is in close proximity to the T2 dermatome.

The T2 dermatome appears to represent the cutaneous areas of the lower cervical segments, as the posterior primary rami of C5, C6, C7, C8 and T1 innervate musculature and have no significant cutaneous innervation.

The pain from the lower cervical spine can also refer pain to the anterior chest, and mimic coronary ischaemic pain. The associated autonomic nervous system disturbance can cause considerable confusion in making the diagnosis.

The medical profession tends to have a blind spot about various pain syndromes in the chest, especially the anterior chest and upper abdomen, caused by the common problem of dysfunction of the thoracic spine. Doctors who gain this insight are amazed at how oft en they diagnose the cause that previously did not enter their ‘programmed’ medical minds.

Physical therapy to the spine can be dramatically effective when used appropriately. Unfortunately, many of us associate it with quackery. It is devastating for patients to create doubts in their minds about having a ‘heart problem’ or an ‘anxiety neurosis’ when the problem is spinal and it can be remedied simply (Chapter 38).

LUMBAR SPINAL DYSFUNCTION

The association between lumbar dysfunction and pain syndromes is generally easier to correlate. The pain is usually located in the low back and referred to the buttocks or the backs of the lower limbs. Problems arise with referred pain to the pelvic area, groin and anterior aspects of the leg. Such patients may be diagnosed as suffering from inguinal or obturator hernial and nerve entrapment syndromes.

Typical examples of referral and radicular pain patterns from various segments of the spine are presented in Figure 25.1.


C
C
C
C
T
T
T
T
T
T
C
T
T
T
T
L
L
L
L
L
L
S
S
L
C
C
C
T
T
T
T
T
C
T
L
L
T
C
L
S
L
L
L
L

C = cervical; T = thoracic; L = lumbar; S = sacral

Figure 25.1 Examples of referred and radicular pain patterns from the spine (one side shown for each segment)

REFERENCES

1. Murtagh J. *Cautionary Tales*. Sydney: McGraw-Hill, 1992: 36 - 130.
2. Kenna C, Murtagh J. *Back Pain and Spinal Manipulation* (2nd edn). Oxford: Butterworth Heinemann, 1997: 213 - 18.

CHAPTER 26. URINARY TRACT INFECTION

Experience has taught them, as mine has me, that one must listen to reason and agree with Hippocrates, Galen, Avicenna and many others, ancient and modern, that there is no surer way to determine the temperaments and constitutions of people of either sex than to look at the urine.

DAVACH DE LA RIVIERE, 18TH CENTURY, THE MIRROR OF URINES

URINARY TRACT INFECTION

Urinary tract infection (UTI) is a common problem affecting all ages and accounts for approximately 1% of all attendances in general practice. It is very common in sexually active women but uncommon in men and children.

Organisms causing UTI in the community are usually sensitive to most of the commonly used antibiotics. The important decision to make is whether to proceed with further investigation of the urinary tract. The morbidity of urinary infections in both children and adults is well known but it is vital to recognise the potential for progressive renal damage, ending in chronic renal failure. The main task in the prevention of chronic pyelonephritis is the early identification of patients with additional factors, such as reflux or obstruction, which could lead to progressive renal damage.

KEY FACTS AND CHECKPOINTS

- Screening of asymptomatic women has shown that about 5% have bacterial UTI.[1]
- About 1% of neonates and 1 - 2% of schoolgirls have asymptomatic bacteriuria.[2]
- About one-third of women have been estimated to have symptoms suggestive of cystitis at some stage of their life.
- The vast majority of these women have anatomically normal renal tracts, are at no significant risk from the UTI and respond quickly to simple therapy. The prevalence of underlying abnormalities is estimated at around 4%.[3]
- UTIs are largely caused by organisms from the bowel that colonise the perineum and reach the bladder via the urethra. In many young women infections are precipitated by sexual intercourse. Ascending infection accounts for 93% of UTIs.
- Haematogenous infection can occur sometimes, especially with the immunocompromised patient.
- All males and females less than 5 years old presenting with a UTI require investigation for an underlying abnormality of the urinary tract.
- In the presence of a normal urinary tract there is no evidence that UTI leads to progressive renal damage.

UTI AS A MASQUERADE

UTI can be regarded as a masquerade when it presents with a constitutional problem or general symptoms, without symptoms suggestive of a urinary infection such as frequency, dysuria and loin pain. This applies particularly to infants and young children and the elderly but is not uncommon in adult women and in pregnancy. Acute UTI may occasionally present as acute abdominal pain.

In infants and children, presenting non-specific symptoms include:

- fever
- Poor feeding
- failure to thrive
- vomiting
- abdominal pain
- diarrhoea

In the elderly:

- confusion
- behaviour disturbance
- fever of undetermined origin

RISK FACTORS FOR URINARY INFECTION

- Female sex
- Sexual intercourse

- Diabetes mellitus
- Diaphragm contraception
- Pregnancy
- Immunosuppression
- Menopause
- Urinary tract obstruction/malformation
- Instrumentation

CLASSIFICATION AND CLINICAL SYNDROMES

Sterile pyuria

This is defined as the presence of pus cells but a sterile urine culture.[2] The common causes of sterile pyuria are:

- contamination of poorly collected urine specimens
- urinary infections being treated by antibiotics, i.e. inadequately treated infection
- analgesic nephropathy
- staghorn calculi
- other kidney disorders, e.g. polycystic kidney
- bladder tumours
- tuberculosis
- chemical cystitis, cytotoxic therapy
- app endicitis

Asymptomatic bacteriuria

This is defined as the presence of a significant growth of bacteria in the urine, which has not produced symptoms requiring consultation.[1] On close questioning many patients admit to mild urinary symptoms.

- This is common only in sexually active women, the elderly and those with urinary tract abnormalities. UTI can exist without any symptoms.[1]
- These patients are more likely to have a past history of symptomatic UTI or to develop symptoms in the future than subjects with sterile urine.[1]
- During pregnancy, asymptomatic bacteriuria leads to acute clinical UTI in up to 33% of women.

Symptomatic bacteriuria

This is defined as the presence of frequency, dysuria and loin pain alone or in combination, together with a significant growth of organisms on urine culture.[2]

The clinical differentiation between cystitis or lower UTI and renal or upper UTI cannot be made accurately on the basis of symptoms, except in those patients with well-defined loin pain and/or tenderness.

Acute cystitis (dysuria-frequency syndrome)[1]

- Inflammation of the bladder and/or urethra is associated with dysuria (pain or scalding with micturition) and/or urinary frequency.
- In severe cases, haematuria may be present, and the urine may have an offensive smell.
- Constitutional symptoms are minimal or absent.
- Other causes of dysuria and frequency include urethritis, prostatitis and vulvovaginitis, all of which can normally be distinguished clinically.

Acute pyelonephritis

- Acute bacterial infection of the kidney produces loin pain and constitutional upset, with fever, rigors, nausea and sometimes vomiting.
- The symptoms of acute cystitis are often also present.
- The differential diagnosis includes causes of the acute abdomen, such as appendicitis, cholecystitis and acute tubal or ovarian diseases. The presence of pyuria and absence of rebound tenderness are helpful in distinction.

The clinical manifestations of UTI are summarised in Figure 26.1.


Symptoms
Generalised
• fever
• chills
• sweating
• rigors
• headache
• nausea
• vomiting
• diarrhoea
indicate kidney infection
Upper tract
• loin pain
• ± abdominal pain
Lower tract
• dysuria
• frequency
• urgency
• feeling of incomplete bladder emptying
• suprapubic discomfort
± strangury
haematuria
offensive urine
Physical examination
(check the following)
General
• temperature
• pulse
• respiration
• blood pressure
loin tenderness
? mass
suprapubic tenderness
vaginal examination
rectal examination

Figure 26.1 Clinical manifestations of urinary tract infection

Uncomplicated urinary tract infection

This is cystitis occurring in the uninstrumented non-pregnant female without structural or neurological abnormalities.

Urethral syndrome

The urethral syndrome (sometimes termed abacterial cystitis) is that where the patient presents with dysuria and frequency but does not show a positive urine culture.[3]

- 30 - 40% of adult women with urinary symptoms have this syndrome.[3]
- Many actually have bacterial cystitis but a negative culture.
- The organisms may be anaerobic or fastidious in their culture requirements.
- The organisms may include Ureaplasma, Chlamydia and viruses.
- The urine may have antiseptic contamination or residual antibiotic.
- The infection may be undergoing spontaneous resolution at the time of the culture.

Interstitial cystitis[3]

This is an uncommon but important cause of the urethral syndrome.

- The classic symptoms are frequency day and night and a dull suprapubic ache relieved briefly by bladder emptying.
- The feature is small haemorrhages on distension of the bladder.

LABORATORY DIAGNOSIS

The laboratory diagnosis of UTI depends on careful collection, examination and culture of urine.

Collection of urine[1]

It is best to collect the first urine passed in the morning, when it is highly concentrated and any bacteria have been incubated in the bladder overnight. Preferably the urine should be taken to the laboratory immediately, but it can be stored for up to 24 hours at 4?C to prevent bacterial multiplication.

- Midstream specimen of urine (MSU). This is best collected from a full bladder, to allow at least 200 mL of urine to be passed before collection of the MSU. It is important that the urine flow is continuous, and the container is moved in and out of the stream collecting at least 20 mL.
    - In women, a tampon should first be inserted and the vulva washed with clean water (to avoid contamination with vaginal and vulval organisms). The labia are then held apart with the fingers to prevent contact with the urinary stream while the specimen is collected.
    - In males, the foreskin (if present) is retracted and the glans washed with clean water.
- Catheter specimen of urine (CSU). In women who have difficulty with collecting an uncontaminated MSU (as is commonly the case in the elderly, the infirm and the grossly obese), a short open-ended catheter can be inserted and a specimen collected after 200 mL has flushed the catheter.
- *Suprapubic aspirate of urine* (SPA). This is an extremely reliable way to detect bacteriuria in neonates and in patients where UTI is suspected but cannot be confirmed because of low colony counts or contamination in an MSU. Under local anaesthetic, a needle (lumbar puncture needle in adults) is inserted into the very full bladder about 1 - 2 cm above the pubic symphysis, and 20 mL is collected by a syringe. Any organisms in an SPA specimen indicate UTI (see Figure 26.2).

Urine specimen collection in children

All children with a first UTI require investigation. It is diagnosed by significant growth on MSU, CSU or SPA.

- Bag specimen: cannot diagnose UTI
- MSU - usually by 3 - 4 years when co-operative
- SPA - reliable
- CSU - for failed SPA or those unable to void on request



Figure 26.2 Suprapubic aspiration of urine in a child

Dipstick testing

Dipstick finding of urinary leucocytes or nitrite are suggestive of LITI.

The reagents in dipstick testing are generally sensitive but have to be interpreted with care. 'Leucocyte esterase dipsticks' are useful in detecting pyuria and give a good guide to infection with a specificity of 94 - 98% (2 - 6% false positive) and 74 - 96% sensitivity (4 - 26% false negatives).[4] Positive nitrite dipsticks give a useful guide to the presence of bacteria. Unexplained haematuria detected by 'dipstick' analysis needs investigation.

Microscopic examination

The urine is examined under a microscope to detect pyuria (more than ten pus cells - WBCs - per high-powered field) but should be examined in a counting chamber to calculate the number of WBCs/mL of urine. In the counting chamber pyuria is > 8000 WBC/mL in phase-contrast microscopy. Pyuria is a very sensitive sign of UTI.

Vaginal squames and debris indicate contamination.

Culture of the urine

The nature and number of organisms present in the urine are the most useful indicators of UTI.[1]

- Most common are enteric organisms. *Escherichia coli* (especially) and *Staphylococcus saprophyticus* are responsible for over 90% of UTIs with other Gram-negative organisms (*Klebsiella* sp. and *Proteus* sp.) and Gram-positive cocci (*Streptococcus faecalis* and other staphylococci) also occurring.
- Infections due to organisms other than *E. coli* (e.g. *Pseudomonas* sp.) are suggestive of an underlying renal tract abnormality.
- If > 10^5 colony forming units (cfu) per mL of bacteria are present in an MSU, it is highly likely that the patient has a UTI.
- On the other hand, it is most important to realise that up to 30% of women with acute bacterial cystitis have less than 10^5 cfu/mL in the MSU. For this reason, it is reasonable to treat women with dysuria and frequency even if they have < 10^5 cfu/mL of organisms in an MSU.

Summary

(Refer to Chapter 16). Significant levels for UTI:

- Microscopy: WBC > 10 per μL (10 × 10^6/L)
- Culture: counts > 10^5 organisms/mL (10^8/L)

MANAGEMENT

- The indications for investigation of UTIs are presented in Table 26.1.
- Appropriate antimicrobial therapy can be expected to cure 80 - 90% of uncomplicated UTIs.
- Optimal treatment includes:
    - high fluid intake
    - complete bladder emptying, including double voiding, especially at bedtime or after intercourse
    - urinary alkalinisation for severe dysuria

ACUTE UNCOMPLICATED CYSTITIS

Advice to women (especially if recurrent attacks):

- Keep yourself rested.
- Drink a lot of fluid: 2 - 3 cups of water at first and then 1 cup every 30 minutes.
- Try to empty your bladder completely each time.
- Gently wash or wipe your bottom from front to back with soft, moist tissues after opening your bowels (for prevention in recurrent attacks).
- Use analgesics such as paracetamol for pain.
- Make the urine alkaline by taking sodium citrotartrate (4 g orally 6 hourly) - not if taking nitrofurantoin.

Table 26.1 Investigation of urinary tract infections

Investigations are indicated in: - all children - all males - all women with: — acute pyelonephritis — recurrent infections — confirmed sterile pyuria — other features of renal disease, e.g. haematuria
Basic investigations include: - MSU - microscopy and culture (post-treatment) - Renal function tests: plasma urea and creatinine - Intravenous urogram (IVU) and/or ultrasound
Special considerations: - in children: micturating cystogram - in adult males: consider prostatic infection studies if IVU normal - in severe pyelonephritis: ultrasound or IVU (urgent) to exclude obstruction - in pregnant women: ultrasound to exclude obstruction

Antimicrobial regimen (for non-pregnant women)[5]

Multiple dose therapy is preferred to single dose therapy.

Use for 5 days in women (trimethoprim - 3 days).

Use for 10 days in women with known urinary tract abnormality:

- trimethoprim 300 mg (o) daily for 3 days or

- cephalexin 500 mg (o) 12 hourly (5 days) or
- amoxycillin/potassium clavulanate 500/125 mg (o) 12 hourly (preferred agent) or
- nitrofurantion 50 mg (0) 6 hourly or
- norfloxacin 400 mg (o) 12 hourly for 3 days (if resistance to above agents proven)

Follow-up: MSU 3 weeks later

Note:

- Avoid using important quinolones - norfloxacin or ciprofloxacin - as first-line agents.
- Cotrimoxazole is not first line because it has no advantage over trimethoprim and has more side-effects.
- Treatment failures are usually due to a resistant organism or an underlying abnormality of the urinary tract.

Treatment of men[5]

Use any one of the regimens for non-pregnant women but use for a minimum of 14 days.

Note: All males with a UTI should be investigated to exclude an underlying abnormality.

ACUTE PYELONEPHRITIS

Mild cases can be treated with oral therapy alone using double the dosage of drugs recommended for uncomplicated cystitis, except for trimethoprim when the same dosage is recommended. Treatment should be continued for 10 days. Ciprofloxacin (500 mg (o) 12 hourly) is used if resistance to these drugs is proven.

For severe infection with suspected septicaemia admit to hospital and treat initially with parenteral antibiotics for 2 to 5 days after taking urine for microscopy and culture and blood for culture. amoxycillin 2 g IV 6 hourly[5] plus gentamicin 4 - 6 mg/kg/day, single daily IV dose

Follow with oral therapy for a total of 14 days therapy. Drug levels of gentamicin require monitoring. Gentamicin can be replaced with IV cefotaxime or ceftriaxone.

All patients should be investigated for an underlying urinary tract abnormality.

RECURRENT OR CHRONIC URINARY TRACT INFECTIONS

Recurrent infections occur as a relapse of a previously treated infection or because of reinfection, often with differing organisms. Persistent (chronic) UTIs indicate that the organism is resistant to the antimicrobial agents employed or that there is an underlying abnormality such as a renal stone or a chronically infected prostate in the male patient. Such infections may be treated with prolonged courses of an appropriate antibiotic or removal of the focus of infection.

In men and children an anatomical abnormality is usual, while recurrent cystitis in women often occurs despite a normal tract. In men, instruction on perineal hygiene, more frequent bladder emptying and postintercourse voiding may assist in the prevention of reinfection.

Treatment of recurrent or chronic UTI[5]

A 10- to 14-day course of:

- amoxycillin/potassium clavulanate (500/125 mg) (o) 12 hourly or
- trimethoprim 300 mg (o) once daily or
- cephalexin 500 mg (o) 12 hourly or (if proven resistance to above agents)
- norfloxacin 400 mg (o) 12 hourly

Prophylaxis for recurrent UTI[5]

In some female patients a single dose of a suitable agent after intercourse is adequate but, in more severe cases, courses may be taken for 3 - 6 months or on occasions longer. Adult doses given:

- nitrofurantoin (macrocrystals) 50 mg (o) nocte or
- trimethoprim 150 mg (o) nocte or
- cephalexin 250 mg (o) nocte or

- norfloxacin 200 - 400 mg (o) nocte (if proven resistance to others)

Cranberries

A recent cochrane review on the use of cranberries (Vaccinium macrocarpon) for the prevention of LUTI concluded that there were evidence to recommended cranberry juice or tablets for the prevention of symptomatic LUTIs in women, but poor evidence for its use in the treatment of LUTI, in the management of a symptomatic bacteriuria or in the prevention of LUTIs in children.

An appropriate regimen is 150 ml cranberry juice or one tablet (1:30 parts concentrated juice) twice daily.

ASYMPTOMATIC BACTERIURIA

- In neonates and preschool children, treat and investigate for evidence of vesicoureteral reflux.
- In men less than 60 years old, treat and investigate, especially for chronic prostatitis.
- In women, give single dose therapy and investigate only those in whom UTI persists or recurs.
- In pregnant women, treat because of the risk of developing pyelonephritis (up to 40% risk).
- School-age children and elderly men and women (over 60 years) probably do not require treatment if their urinary tracts are normal.
- In patients with long-term indwelling catheters treatment is not usually required or useful.
- Prophylaxis should be given for recurrent asymptomatic bacteriuria in pregnant women, in patients with associated renal tract abnormality, and in those undergoing genitourinary instrumentation or surgery or intermittent catheterisation.

URINARY TRACT INFECTION IN CHILDREN

UTI in infants and very young children is often renal in nature and may be associated with generalised symptoms such as fever, vomiting, diarrhoea and failure to thrive. Offensive urine may be noted. Symptoms of dysuria and frequency appear only after the age of 2 years when the child is able to indicate the source of the discomfort. In a girl or boy (rare presentation) with symptoms of dysuria and frequency an underlying abnormality is likely to be present with a reported incidence of vesicoureteric reflux (VUR) as high as 40% and scarred kidneys (reflux nephropathy) in 27%.[3]

Thus the early detection of children with VUR and control of recurrent renal infection could prevent the development of scars, hypertension and chronic renal failure. Radiological investigation of children with UTIs shows normal kidneys in approximately 66% and reflux in approximately 33%.

It is best to perform the ultrasound first.

Guidelines for investigation[8]

- < 1 year - ultrasound; possibly micturating cystourethrogram (MCU)
    - if both negative, no further investigation
    - if abnormal, follow up referral/investigation
- > 1 year - ultrasound

The dimercaptosuccinic acid scintigraph scan is the gold standard for diagnosis of renal scarring and measurement of differential renal function. It is usually reserved for children with dilating VUR.

Antimicrobial treatment for acute cystitis in children[5]

Treatment should be continued for 5 days:

- trimethoprim 6 mg/kg (max. 300 mg) orally, daily (suspension is 50 mg/5 mL) or
- cephalexin 12.5 mg/kg (max. 500 mg) orally 6 hourly or
- cotrimoxazole 4/20 mg/kg, (max. 160/800 mg) orally bd or
- amoxycillin/potassium clavulanate 22.5 mg/kg (max. 500/125 mg) orally 12 hourly

Norfloxacin and ciprofloxacin should be avoided routinely in children.

Check MSU in 3 weeks.

Vulvovaginitis in children

Although vulvovaginitis can affect women of any age, it is more prevelant in girls between 2 and 8 years. It can be confused with a LITI where there is dysurea which is a common symptom from this type of dermatitis (see p.xxx).

URINARY INFECTIONS IN THE ELDERLY

The typical settings in which UTIs occur in the elderly are in the frail, those who are immobilised, and those with faecal incontinence and inadequate bladder emptying. The presenting symptoms may be atypical, especially with upper UTI where fever of undetermined origin and behaviour disturbances may be a feature. In men obstructive uropathy from prostatism should be excluded by ultrasound.

Uncomplicated infections should be treated the same way as for other age groups but no antimicrobial treatment is recommended for asymptomatic bacteriuria.

URINARY INFECTIONS IN PREGNANCY

UTI in pregnant women requires careful surveillance. Asymptomatic bacteriuria should always be excluded during early pregnancy because it tends to develop into a full-blown infection. Acute cystitis is treated for 10 to 14 days with any of the following antimicrobials: cephalexin, amoxycillin/potassium clavulanate or nitrofurantoin (if a beta-lactam antibiotic is contraindicated). The dosages are the same as for other groups. Asymptomatic bacteriuria should be treated with a week-long course. Refer to Chapter 101

GENITOURINARY TUBERCULOSIS

The genitourinary tract is involved in 3 - 5% of cases of tuberculosis.[9] The genital and urinary tracts are often involved together as a result of miliary spread.

The commonest presenting complaints are dysuria and frequency, which can be severe. Other symptoms include strangury when the bladder is severely affected, loin pain and haematuria. Routine urine culture shows sterile pyuria.

Diagnosis is made on specific culture for mycobacterium, or biopsy of bladder lesions or the typical X-ray appearance of distorted calyces and medullary calcification. Treatment is with antituberculosis drugs.

CANDIDURIA[5]

The presence of Candida albicans in the urine is common. Antifungal therapy is not recommended if associated with indwelling catheters but is recommended if associated with upper UTIs and/or systemic candidiasis.

PROSTATITIS

Consider bacterial prostatitis in men with few urinary symptom (frequency, urgency and dysuria), flu-like illness fever, low backache and perineal pain. The prostate is exquisitely tender on reetal examination. For ruild to modirate infection, give the same oral treatment as for cystitis, e.g. amoxycillin + clavalanate 500/125 mg (0) bd for 14days. If severe, use amoxy/ampiallin 29 iv 6hours plus gentaincim (see p.xxx).

COMMON TREATMENT ERRORS[1]

- Not treating women with dysuria and frequency merely because there are < 10^5 cfu/mL in an MSU
- Overtreating women with acute cystitis and normal urinary tracts; single-dose therapy is effective in 70 - 80% of cases, and overtreatment often leads to vaginal candidiasis or antibiotic-induced diarrhoea
- Using single-dose therapy in patients with known anatomical abnormalities
- Failing to consider chronic prostatic infection in men with recurrent UTI and a normal IVU

WHEN TO REFER

- It is wise to refer all patients with urinary tract abnormalities to a nephrologist or urologist for advice on specific management.
- Refer also if the simple methods outlined above do not control recurrent UTI.
- Refer males with urinary infections that are not clearly localised to the prostate.
- Refer patients with impaired renal function.

Practice tips

- Most symptomatic UTIs are acute cystitis occurring in sexually active women with anatomically normal urinary tracts.
- A clinical diagnosis based on experience, plus a positive nitrite dipstick test and the finding of pyuria by office microscopy will generally enable immediate curative treatment.
- A 3-day course of trimethoprim 300 mg daily for 3 days is a suitable first choice for acute uncomplicated cystitis in women.
- Avoid overinvestigation of patients in whom there is a low likelihood of demonstrating structural abnormalities.
- The ultrasound examination may not defeat calculi small tumours, clubbed calyces and paprllary neeroris.
- In males the prostate is the most common source of recurrent UTI.
- UTI is commonly associated with microscopic haematuria (occasionally macroscopic haematuria).
- Persisting haematuria should be investigated.
- Due to the rising level of E. coli resistance, amoxycillin is no longer recommended unless susceptibility of the organism is proven.[5]

REFERENCES

1. Becker GJ. Urinary tract infection. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 545 - 7.
2. Heale W. *Kidney Disease*. CHECK Programme. Melbourne: RACGP, 1987: 1 - 20.
3. Kincaid-Smith P, Larkins R, Whelan G. *Problems in Clinical Medicine*. Sydney: MacLennan & Petty, 1990: 280 - 3.
4. Devilte WLIM etal. The urine dipstick test useful to rule out infection. A meta-analysis of the accuracy BioMed Central Urology,2004;4: 4.
5. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006; 307 - 315.
6. Jepson R G, Mihaljevic L, Craig J. Cranberries for preventing urinary tract infections (Review). Cochrane Database. Sept. Rev. 2004, Issue 2. Art No. CD001321.
7. Wyndam R. Cranberry juice and urinary tract infections. Medicine Today, 2006, 7(5): 72 - 3.
8. Paxton G, Munro J (ed). *Paediatric Handbook* (7th edn). Melbourne: Blackwell Science, 2003: 547 - 50.
9. Bullock N, Sibley G, Whitaker R. *Essential Urology*. Edinburgh: Churchill Livingstone, 1989: 126 - 9.

CHAPTER 27. MALIGNANT DISEASE

Cancers of the tongue and mouth begin with a small hard lump, and sometimes with a little sore; both of which are attended with pricking pains, and they spread in the same manner with cancerous sores in other parts. It is so great an evil, that the slightest suspicion of it occasions very great uneasiness.

WILLIAM HEBERDEN (1710-1801)

MALIGNANT DISEASE

The terms *malignancy, cancer* and *neoplasia* are usually used interchangeably. The differences between a malignant tumour and a benign tumour are summarised in Table 27.1.

Malignant disease accounts for 1 in 8 deaths of people under 35 years in Australia and 1 in every 4 (25%) of deaths in those over 45 years.[1] Cancer is the only major cause of death in Australia that is increasing in both sexes. At current rates about 1 in 3 males and 1 in 4 females will develop a cancer, excluding non-melanoma skin cancers, by the age of 75.[1]

The six most common causes of death from cancer in Australia and the United States are cancer of the lung, bowel, breast, prostate, lymphoma and pancreas.

Neoplasia, especially malignancy of the silent areas, can present as undifferentiated illness and be a real masquerade. The so-called ‘silent’ malignancies that pose a special problem include carcinoma of the ovary, kidney, caecum and ascending colon, liver (hepatoma), melanoma and haematological tissue.

This chapter focuses on the general features of these malignancies in order to promote early diagnosis and urgent referral at the primary care level.

Table 27.1 Different characteristics of benign and malignant tumours

Benign	Malignant
Well differentiated	Undifferentiated
Non-invasive	Invasive
Slow growth	Rapid growth
Not anaplastic	Anaplastic
Not metastatic	Metastatic

Acute emergency problems that can develop with various malignancies include spinal cord compression, malignant effusions, dis-seminated intravascular coagulation and hypercalcaemia.

CANCER IN CHILDREN

Although uncommon in children under 15 years, cancer is the second most common cause of death in this age group. The most common cancers (in order) are leukaemias, especially acute lymphocytic leukaemia (34%), brain tumours, especially astrocytoma (20%), lymphomas, especially non-Hodgkin's (13%), neuroblastoma, Wilms' tumour, soft tissue tumours especially rhabdomyosarcoma, and bone tumours.

Survival has improved dramatically in recent decades, indicating the value of early diagnosis and referral for expert treatment.

Studies have highlighted the importance of GPs responding to concerns of parents even if they could find no abnormality after examination. Parents of children eventually diagnosed with cancer and who were in dispute with their GP were alerted by signs and symptoms which were often vague, non-specific and common or unusual or ‘scary'. They felt that their child ‘wasn't right’.[2]

CLINICAL MANIFESTATIONS

The clinical manifestations of malignancy are usually due to:

- Pressure effects of the growth
- Infiltration or metastases in various organs (e.g. liver, brain, lungs, bone, blood vessels)
- Systemic symptoms, including paraneoplastic effects

SYSTEMIC SYMPTOMS

These can be divided into general non-specific effects and paraneoplastic syndromes, which are the remote effects caused by the tumour.

Undifferentiated general symptoms

- Tiredness/fatigue/weakness
- Anorexia and nausea
- Weight loss
- Fever
- Thirst (hypercalcaemia)
- Drowsiness (hyponatraemia)

Paraneoplastic effects

The paraneoplastic effects or syndromes are very important clinically because they may provide an early clue to the presence of a specific type of cancer, in addition to the possible lethal effect of the metabolic or toxic effect (e.g. hyponatraemia). These effects include:

- ectopic hormone production
- skin abnormalities
- metabolic effects
  - fever/sweats
  - weight loss
- haematologic disorders
  - anaemia
  - erythrocytosis
  - coagulation disorder
  - others
- neuropathies and CNS abnormalities
- collagen vascular disorders

A summary of various paraneoplastic syndromes is presented in Table 27.2.

Table 27.2 Paraneoplastic syndromes and associated tumours: more common examples

Hormone excess syndrome	Tumour
Cushing's	Lung, kidney, adrenal, thymoma, pancreas
ACTH	Lung, kidney, thymoma, thyroid
Gonadotrophins	Lung, hepatoma, choriocarcinoma
Other syndromes	**Tumour**

Hypercalcaemia	Lung, breast, kidney, multiple myeloma, prostate, pancreas, adrenal, hepatoma
Fever	Kidney, hepatoma, lymphoma, pancreas, thymoma
Neurologic	Lung, breast, thymoma, Hodgkin's, prostate
Coagulopathy	Lung, breast, hepatoma, prostate, pancreas
Thrombophlebitis	Kidney, pancreas, prostate
Polycythaemia	Kidney, hepatoma
Dermatomyositis	Lung, breast, pancreas

CLINICAL APPROACH

A history of constitutional symptoms that are often quite undifferentiated (often bizarre) may provide the clue to the possibility of an underlying malignancy. An occupational history may be relevant to the clinical problem (see Table 27.3).

Table 27.3 Occupational causes of cancer

Agent	Occupation	Cancer
Arsenic	Chemical industry	Lung, skin, liver
Asbestos	Insulation worker	Mesothelioma
Benzene	Glue worker, varnisher	Leukaemia
Soot, coal tar	Chimney sweep	Skin
Radiation	Mining, watch dials	Various
Ultraviolet light	Farmer, sailor	Skin
Vinyl chloride	PVC manufacturing	Liver (angiosarcoma)

Familial cancer

Although the great majority of cancer is not inherited some individuals carry inherited genetic mutations from conception that predispose them to developing certain cancers, particularly colorectal, breast and ovarian cancers. Refer to Chapter 19 (pp. 175Ð6).

TUMOUR MARKERS

A tumour marker is an abnormal characteristic that is specific for a particular type of malignancy (e.g. the Philadelphia chromosome for chronic myeloid leukaemia). Other exam-ples include human chorionic gonadotrophin (HCG) (elevated in trophoblastic tumours and germ cell neoplasms of the testes and ovaries) and the oncofetal antigens - carcinoembryonic antigen (CEA) and alpha-fetoprotein (AFP).

CEA and AFP are not specific markers but are elevated in certain tumours and are very useful in monitoring tumour activity.

Tumour markers, some of which are as yet unidentified, would appear to have an important role in future diagnosis and management of malignant disease. The most valuable are those associated with testis cancer - AFP and beta-HCG. Markers may be an adjunct to diagnosis of certain malignancies, including CEA for bowel cancer and CA-125 for ovarian cancer. These markers are summarised in Table 27.4.

LUNG CANCER

Apart from non-melanoma skin cancer, lung cancer is the most common cancer in Australia in terms of both incidence and death, accounting for at least 20% of cancer deaths.[1] In the United States it accounts for 28% of cancer deaths in men and 24% of deaths in women. Only 10Ð25% are asymptomatic at the time of diagnosis but lung cancer can cause an extraordinary variety of clinical symptoms and signs with a reputation for several paraneoplastic syndromes. Refer to Chapter 42 (pp. 459Ð61) and Chapter 49 (p. 547).

The paraneoplastic syndromes include hypercalcaemia, Cushing's syndrome, carcinoid syndrome, dermatomyositis, visual loss progressing to blindness from retinal degeneration, cerebellar degeneration and encephalitis.

The presentation of cough and chest pain renders it less of an ‘occult’ malignancy than several other types.

Table 27.4 Common tumour markers

Tumour marker	Condition
AFP	Testicular cancer (non-seminomatous) Hepatocellular carcinoma GIT cancers with and without liver metastases
CA-125	Ovarian carcinoma (non-mucinous)
CA-15-3	Breast
CA-19-9	Pancreas colon ovary
CEA	Colorectal cancer Pancreatic, breast, lung, small intestine, stomach, ovaries
PSA[*]	Prostate cancer
hCG	Choriocarcinoma Hydatidiform mole Trophoblastic diseases

[*] PSA ° prostate specific antigen

DxT: malaise + weight loss + cough ° carcinoma lung

RENAL TUMOURS

The most important tumours of the kidney are adenocarcinoma (80% of all renal tumours)[3] and nephroblastoma (Wilms' tumour).

Adenocarcinoma of kidney

Adenocarcinoma of the kidney has a great diversity of presenting symptoms, including:

- general symptoms of neoplasia
- haematuria (60%)
- loin pain (40%)
- loin mass
- signs of anaemia
- left supraclavicular lymphadenopathy
- varicocele
- hypertension
- symptoms of metastases (to liver, lungs, brain, bones)
    - respiratory symptoms
    - neurological symptoms and signs
    - bone pain
    - pathological fracture (vertebral collapse)

- urinalysis - 67% positive for blood

Diagnosis confirmed by imaging e.g. CT scan.

Refer for radical nephrectomy.

The classic triad of symptoms (in 10Ð15% of patients) is.[3]

DxT: haematuria + loin pain + palpable kidney mass ° adenocarcinoma

Wilms' tumour

Wilms' tumour is responsible for 10% of all childhood malignancies. Clinical features include:[3]

- peak incidence 2Ð3 years
- general symptoms of neoplasia
- palpable mass 80%
- abdominal pain 30%
- haematuria 25%

Diagnosis confirmed by ultrasound or CT scan. Early diagnosis with nephrectomy and chemotherapy leads to a very favourable prognosis.

DxT: FUO + abdominal mass + malaise ° Wilms' tumour

Ovarian cancer

Ovarian cancer has the highest mortality rate of all the gynaecological cancers because the majority of patients present in the late stage of the disease. It is responsible for5% of deaths in females It is usually asymptomatic prior to the development of metastases. Epithelial tumours are the most common of malignant ovarian tumours. They are uncommon under 40 years of age and the average age of diagnosis is 50 years.[4]

The most common presentation is abdominal swelling (mass and/or ascites). abdominal bloating or discomfort Non-specific symptoms, which may be present for a long time before diagnosis, include abnormal uterine bleeding, urinary frequency weight loss, abdominal discomfort, reduced capacity for food, anorexia, nausea and vomiting.(refer to p.998).

Diagnosis confirmed by pelvic ultrasound.

DxT: abdominal discomfort + weight loss + abnormal uterine bleeding ° carcinoma ovary

CARCINOMA OF CAECUM AND ASCENDING COLON

Malignancy in this area is more likely to present with symptoms of anaemia without the patient noting obvious blood in the faeces or alteration of bowel habit. Refer to Chapter 41 (p. 439).

DxT: blood in stools + mucus discharge + change in bowel habit ° carcinoma colon

THE LEUKAEMIAS

The leukaemias are caused by an acquired malignant transformation in the stem cell in the haemopoietic system. Acute leukaemia has a rapidly fatal course if untreated, while chronic leukaemia has a variable chronic course with an inevitable fatal outcome. See Figure 27.1. The main features of each type are as follows.


raised intracranial pressure (2%)
miserable, irritable (80%)
increasing paleness (80%)
bruising (80%)
testicular enlargement (2%)
joint pain (10%)
pain on walking or being held (80%)

Figure 27.1 Clinical features of a child with leukaemia [5]

The usual age range for acute lymphatic leukaemia (ALL) is 2-10 year with a second peak at about 40 years. The median age of presentation of acute myeloid leukaemia (AML) is 55-60 years.

Acute leukaemia

Symptoms

- General constitutional (e.g. malaise)
- Symptoms of anaemia

- Susceptibility to infection (e.g. sore throat, mouth ulceration, chest infection)
- Easy bruising and bleeding (e.g. epistaxis, gingival bleeding)
- Bone pain (notably in children with ALL) and joint pain
- Symptoms due to infiltration of tissues with blast cells (e.g. gingival hypertrophy in AML)

DxT: malaise + pallor + bone pain ° ALL

DxT: malaise + pallor + oral problems ° AML

Signs

- Pallor of anaemia
- Petechiae, bruising
- Gum hypertrophy/gingivitis/stomatitis
- Signs of infection
- Variable enlargement of liver, spleen and lymph nodes
- Bone tenderness, especially sternum and liver

Diagnosis

- FBE and film
    - normochromic/normocytic anaemia
    - pancytopenia with circulatory blast cells
    - Platelets: usually reduced
- Bone marrow examination

NOTE: As a rule relapse of a cute leukaemia means imminent death innless bone marrow transplantation is successful. The mean 5-year survival rate for dsildhood all about70-80% for adult all 30%, for AML it varies with age with poorer survival, about 20%, oven 55 year of age.

Chronic myeloid leukaemia (CML)

Main clinical features

- A disorder of middle age
- Insidious onset
- Constitutional symptoms: malaise, weight loss, fever, night sweats
- Symptoms of anaemia
- Splenomegaly (very large)
- Priapism
- Gout
- Makedly elevated white cell count (granulocytes)
- Marked left shift in myeloid series
- Presence of Philadelphia chromosome

DxT: fatigue + night sweats + abdominal fullness (splenomegaly) ° CML

Chronic lymphocytic leukaemia (CLL)

Main clinical features

- A disorder of late middle age and elderly

- Insidious onset
- Constitutional symptoms: malaise, weight loss, fever
- Lymphadenopathy - neck, axilla, groin (80%)
- Moderately enlarged spleen and liver (about 50%)
- Mild anaemia
- Lymphocytosis > 15×10^9/L
- ‘Mature’ appearance of lymphocytes

DxT: fatigue + weight loss + lymphadenopathy ° CLL

THE LYMPHOMAS[6]

Lymphomas, which are malignant tumours of lymphoid tissue, are classified as Hodgkin's lymphoma and non-Hodgkin's lymphoma on the basis of histological appearance of the involved lymph tissue.

Hodgkin's lymphoma

Clinical features

- Painless (rubbery) lymphadenopathy, especially cervical nodes
- Constitutional symptoms (e.g. malaise, weakness, weight loss)
- Fever and drenching night sweats
- Pruritus
- Alcohol-induced pain in any enlarged lymph nodes
- Possible enlarged spleen and liver

Diagnosis is by lymph-node biopsy with histological confirmation.

DxT: malaise + night sweats + pruritus ° Hodgkin's lymphoma

Non-Hodgkin's lymphomas

Non-Hodgkin's lymphomas are a hetero-geneous group of cancers of lymphocytes derived from the malignant clones of B or T cells.

Main clinical features

- Painless lymphadenopathy - localised or wide spread
- Constitutional symptoms possible, especially sweating
- Pruritus is uncommon
- Extra nodal sites of disease (e.g. skin, GIT)
- Possible enlarged liver and spleen
- Possible nodular infiltration of skin (e.g. mycosis fungoides)

Diagnosis is by lymph-node biopsy.

DxT: malaise + night sweats + lymphadenopathy ° non-Hodgkin's lymphoma

ARMULTIPLE MYELOMA

Multiple myeloma is a clonal malignanas of the differentiated B lyinphocyde the plasma cell. It is regarded as a disease of the elderly, the mean age of presentation being 60 years.[6] The classic presenting triad in an older person is anaemia, back pain and elevated ESR.

Main clinical features

- Bone pain (e.g. backache) - in more than 80% of patients
- Bone tenderness
- Weight loss
- Recurrent infections
- Symptoms of anaemia
- Bleeding tendency
- Replacement of bone marrow by malignant plasma cells

> **DxT:** weakness + back pain + weight loss ° multiple myeloma

Diagnostic criteria

The presence of:[5]

- paraprotein in serum (on electrophoresis)
- Bence-Jones protein in urine
- bony lytic lesions

Treatment is with chemotherapy: 3 year median survival

CARCINOID SYNDROME

Hormone secretion by carcinoid cells causes the characteristic carcinoid syndrome long before local growth or metastatic spread of the tumour is apparent.

The syndrome

- Classic triad: skin flushing (especially face), diarrhoea (with abdominal cramps), valvular heart disease
- Other features: wheezing, telangiectasia, hypotension
- Sites of tumours: ileum, stomach, bronchi

POTENTIALLY CURABLE MALIGNANT TUMOURS

Several tumours are curable by chemotherapy even in the advanced stage. Such tumours are as follows.

Haematological tumours

- Some lymphomas
- Hodgkin's lymphoma
- Acute lymphatic leukaemia
- Acute myeloid leukaemia

Solid tumours

- Choriocarcinoma
- Testicular teratoma

- Neuroblastoma
- Wilms' tumour (nephroblastoma)
- Burkitt's tumour
- Embryonal rhabdomyosarcoma

Tumours curable by adjuvant chemotherapy

- Breast cancer (especially up to stage 2)
- Osteogenic cancer
- Soft tissue cancer
- Colorectal cancer

Survival rates

Common cancers and their seqear survival rates over a isyear penod (1987-1997) hare been puttished by the Victorian caner council (Table 27.5). These show improving trends for many cancers. The lowest survival rate was for meso thelioma at 4%.

Table 27.5 Common cancers and their 5-year survival rates[7]

Testicular	94%
Melanoma	91%
Thyroid	91%
Hodgkin's disease	84%
Breast	82%
Uterus	81%
Prostate	76%
Bladder	73%
Non-Hodgkin's Lymphoma	56%
Colon	55%
Ovary	42%
Stomach	21%
Lung	11%
Liver	8%
Panaceas	5%

METASTATIC TUMOURS

It is very helpful for the practitioner to have a working knowledge of possible primary sources of tumour when metastatic lesions are detected in various organs.

Common sites of metastatic presentation are the lymph nodes, liver, lung, mediastinum and bone. Other sites include the brain, bone marrow, peritoneum, retroperitoneum, skin and the spinal cord.

These important sites (listed below) are followed by likely primary sources with the most likely listed first.

- *Liver*. Colon, pancreas, liver, stomach, breast, lung, melanoma
- *Lung and mediastinum*. Breast, lung, colon, kidney, testes, cervix/uterus, Hodgkin's lymphoma, melanoma

- *Bone*. Breast, prostate, lung, Hodgkin's lymphoma, kidney, thyroid, melanoma
- *Brain*. Breast, lung, colon, lymphoma, kidney, melanoma, prostate
- *Skin*. Lung, colon, melanoma, Kaposi's sarcoma
- Lymph nodes:
  - High cervical. Hodgkin's lymphoma, lymphoma, squamous cell carcinoma, oropharynx, nasopharynx
  - Low cervical. Lung, stomach, lymphoma, Hodgkin's lymphoma, oropharynx, larynx, skin, tongue
  - Axillary. Breast, lung, lymphoma
  - Inguinal. Lymphoma, ovary, uterus, vulva, prostate, skin
- Retroperitoneum. Lymphoma, Hodgkin's lymphoma, ovary, uterus, testes, prostate

It is important to keep in mind those malignancies that are potentially curable and refer as soon as possible.

PREVENTION

Preventive measures for malignant disease are addressed in more detail in Chapter 9. The significant decrease in deaths from cancer of the stomach in this country in recent years is probably reflected in our improved diet with more fresh fruit and vegetables. Important preventive measures include an appropriate healthy diet, no smoking, sun protection and perhaps safe sex measures. Of concern is the rapid increase in incidence of prostate cancer, chronic myeloid leukaemia, myeloma and non-Hodgkin's lymphoma.

Triads to consider

- **DxT:** anorexia + weight loss + jaundice (? epigastric pain) ° carcinoma pancreas
- **DxT:** fatigue + dysphagia + weight loss ° carcinoma oesophagus
- **DxT:** anorexia + dyspepsia + weight loss ° carcinoma stomach
- **DxT:** headache + a/n/v + ataxia ° medulloblastoma (children)
- **DxT:** fever + malaise (extreme) + a/n/v (? anaemia) ° neuroblastoma
- **DxT:** mental dysfunction + vomiting + (waking) headache ° cerebral tumour (late)
- **DxT:** indrawn eye + small pupil + ptosis (? anhydrosis) ° Horner's syndrome (? carcinoma lung)

REFERENCES

1. Australia's Health 1990. Canberra: Australian Institute of Health, 1990: 57Ð64.
2. Halliday, J. Malignant disease in children: the view of a general practitioner and parent. In: Baum J, Dominica F, Woodward R. *Listen my Child has a Lot of Living to Do*. Oxford: Oxford University Press, 1990: 19Ð27.
3. Bullock N, Sibley G, Whitaker Q. *Essential Urology*. Edinburgh: Churchill Livingstone, 1989: 202Ð8.
4. Dalrymple C. Ovarian cancer: a sinister and silent disease. Mod Med Aust, 1992; July: 26Ð37.
5. Vowels MR. Common presentations and management of leukaemia in childhood. Aust Fam Physician, 1994; 23: 1519Ð21.
6. Kumar PJ, Clark ML. *Clinical Medicine* (5th edn). London: Sounder, 2003:501 - 3
7. English D. *Caner survival in Victorian*. Melbourne Victorian cancer counal, 2003.

CHAPTER 28. HIV/AIDS - COULD IT BE HIV?

> The verdict for him too was death, not the inevitable death that horrified and yet was tolerable because science was helpless before it, but the death which was inevitable because the man was a little wheel in the great machine of a complex civilisation.
>
> W. SOMERSET MAUGHAM (1874 - 1965), *OF HUMAN BONDAGE*

HIV/AIDS - COULD IT BE HIV?

HIV, the cause of the well-known AIDS, can rightly be included as one of the clinical masquerades of modern medicine. Public health measures in the Western world have limited the spread of the infection, particularly in Australia where the incidence of new cases has remained relatively constant in the past 10 years. By contrast the incidence in Africa and Asia continues to rise at an alarming rate. The introduction of combination treatment with the protease inhibitors in November 1995 has changed the previously understood natural history of the disease and has given rise to renewed hope that HIV will become a chronic manageable disease.

The benefit of early diagnosis has become even more impressive since the discovery that HIV is not a latent infection throughout most of its course. Soon after initial infection, an explosive replication of HIV occurs, which is brought under control by the immune system in 6 to 8 weeks as the host-versus-virus interaction reaches an active and dynamic equilibrium. This dynamic situation continues throughout a person's lifetime, with as many as 10 billion new virions produced and up to 2 billion CD_4 lymphocytes destroyed and replaced daily. Clinical immunodeficiency develops when the body's ability to replace CD_4 cells is finally exhausted, resulting in further uncontrolled viral replication. It has taken over a decade to get this understanding of the dynamics of HIV infection. Viral load assays based on molecular techniques have revolutionised our understanding of the natural history of HIV disease. These advances make it imperative to make the diagnosis early in the course of the disease in order to start combination treatment to lessen the viral load.

The management of HIV infection is a specialised field but the general practitioner is central in prevention, diagnosis, counselling, monitoring and shared management of HIV disease. The GP must be alert to the benefits of early diagnosis as summarised in Table 28.1.

Table 28.1 The benefits of early HIV diagnosis

To individual patients
• Prolongation of the asymptomatic period
• Delayed disease progression
• Prevention of opportunistic infections
• Optimal maintenance of health through patient education and counselling
• Cures are only likely with early intervention
To the cohort of HIV-positive individuals
• Monitoring of advances in treatment
• Increased participation in research and clinical trials
• Development of new services to meet changing patient needs
To the community
• Documentation of changes in epidemiology
• Reduced high - risk activities
• Contact tracin
• Control of HIV transmission
To the doctors
• Time to influence the course of disease
• Time to counsel the patient

Source: Penny R. Could it be HIV? 2. Benefits of early diagnosis of HIV infection. Med J Aust, 1993; 158: 35 - 6. © Copyright 1993, The Medical Journal of Australia - reproduced with permission

KEY FACTS AND CHECKPOINTS

- HIV is a retrovirus with two known strains that cause a similar spectrum of syndromes: HIV-1 and HIV-2 (mainly confired to West Africa). It infects T-helper cells bearing the CD_4 receptor.
- Always consider HIV in those at risk: enquire about history of STIs, injection of illicit drugs, past blood transfusions, sexual activities and partners.
- About 50% of patients develop an acute infective illness similar to glandular fever within weeks of acquiring the virus (the HIV seroconversion illness).[1] The main features are fever, lymphadenopathy, lethargy and possibly sore throat, and a generalised rash.
- If these patients have a negative infectious mononucleosis test, perform an HIV antibody test, which may have to be repeated in 4 weeks or so if negative.
- Patients invariably recover to enter a long period of good health for 5 years or more.[2]
- Pneumocystitis pneumonia (PCP) is the commonest presentation of AIDS.
- Approximately 15 - 40% of HIV-positive children are infected from HIV-infected mothers.[3]
- Infants born to these mothers may develop the disease within a few months, with 30% affected by the age of 18 months.
- The time for the onset of AIDS in HIV-affected adults varies from 2 months to 20 years or longer; the median time is around 10 years.
- In family practice the most common presentation of HIV-related illness is seen in the skin/oral mucosa.[4]
- HIV antibody testing is a two-stage process: enzyme linked immunosorbent assay (ELISA) for screening followed by another method (e.g. Western blot) if ELISA is positive.
- The seroconversion period from acquiring HIV infection to a positive antibody test varies between individuals: this period is known as the 'window period'.
- All HIV infected patients require regular monitoring for immune function and viral load. The viral load test monitors viral activity.
- The level of immune depletion is best measured by the CD_4 positive T-lymphocyte (helper T-cell) count - the CD_4 cell count. The cut-off points for good health and severe disease appear to be 500 cells/?L and 200 cells/?L respectively.[2]

OCCURRENCE AND TRANSMISSION

HIV can be isolated from blood, tissues, semen, saliva, breast milk, cervical and vaginal secretions and tears of infected persons. HIV is transmitted in semen, blood and vaginal fluids, transplanted organs and breast milk through:

- unprotected sexual intercourse (anal or vaginal) and in rare cases oral sex with an infected person
- infected blood entering the body (through blood transfusion or by IV drug users sharing needles/syringes)
- needle-stick injury
- artificial insemination, organ transplantation
- infected mothers (to babies during pregnancy, at birth or in breast milk)

Infection with HIV can occur via the vagina, rectum or open cuts and sores, including any on the lips or in the mouth. Social (non-sexual) contact and insect vectors have not been implicated in transmission.

CLINICAL MANIFESTATIONS[5]

There are a multiplicity of clinical findings in HIV infection (Figure 28.1). The clinical stages of HIV disease are summarised in Table 28.2.(on p. 247).[6]


General
fever
night sweats
lethargy/malaise
anorexia
pharyngitis
arthralgia
myalgia
lymphadenopathy
Neuropathic
cognitive/affective impairment
headache
photophobia
meningoencephalitis
radiculopathy
brachial neuritis
peripheral neuropathy
Guillain-Barré syndrome
Gastrointestinal
oral candidiasis
nausea/vomiting
diarrhoea
mucocutaneous ulceration
Dermatopathic
maculopapular rash
roseola-like rash
alopecia
urticaria (diffuse)
desquamation
herpes simplex reactivation
herpes zoster
Laboratory tests
WCC↓
Platelets↓
ESR ↑ (slight)
s. transaminases↑
HIV –ve
recovery phase:
lymphocytosis
atypical mononucleosis
inversion $CD_4 : CD_8$ ratio
3–12 weeks later
HIV +ve

Figure 28.1 Possible clinical features of primary HIV infection

Table 28.2 Clinical stages of HIV disease[6]

Clinical stage	Common clinical features	CD_4 count
Seroconversion illness (self-limited 1 - 3 weeks)	Fever, headache (may have aseptic meningitis), sore throat, maculopapular rash, lymphadenopathy, splenomegaly; Atypical lymphocytes on FBE cells	Transient decrease, commonly followed by a return to near normal levels
Asymptomatic	Headaches Persistent generalised lymphadenopathy	Usually > 500 cells/μL Gradual decrease of 50 - 80 cells/μL
Symptomatic - early	Oral and vaginal candidiasis, oral hairy leucoplakia, seborrhoeic dermatitis, psoriasis, recurrent varicella zoster infection, cervical dysplasia, unexplained fever, sweats, weight loss, diarrhoea, tuberculosis	Usually 150 - 500 cells/μL
Symptomatic - late	PCP, Kaposi's sarcoma, oesophageal candidiasis, cerebral toxoplasmosis, lymphoma, HIV-1 associated dementia complex, cryptococcal meningitis	Usually < 150 cells/μL
Advanced	CMV retinitis, cerebral lymphoma, Mycobacterium avium complex (MAC) infection	Usually < 50 cells/μL

Reproduced with permission[6]

Acute (seroconversion) illness

At least 50% of patients have an acute illness associated with seroconversion. The illness usually occurs within 6 weeks of infection and is characterised by fever, night sweats, malaise, severe lethargy, anorexia, nausea, myalgia, arthralgia, headache, photophobia, sore throat, diarrhoea, lymphadenopathy, generalised maculoerythematous rash and thrombo-cytopenia. The main symptoms are headache, photophobia and neurological manifestations, including meningoencephalitis and peripheral neuritis. Acute HIV infection should be considered in the differential diagnosis of illnesses resembling glandular fever. This illness is self-limiting and usually revolves within 1 to 3 weeks. However, chronic lethargy, depression and irritability may persist after the acute illness. Non-specific viraemic sequelae such as mucosal ulceration, desquamation, exacerbation of seborrhoea and recurrences of herpes simplex may occur (see Figure 28.1).

Acute illness may be accompanied by neutropenia, lymphopenia, thrombocytopenia, and mildly elevated ESR and serum trans-aminases. During recovery lymphocytosis may occur with appearance of atypical mononuclear cells and an inversion of the CD_4 +:CD_8 + ratio due to elevation of CD_8 + cells. It is seronefahre for EBV.

Differential diagnoses are given in Table 28.3.

Table 28.3 Differential diagnoses of primary HIV infection

Epstein-Barr mononucleosis
Syphilis: secondary
TORCH organisms
• toxoplasmosis
• rubella
• CMV (espeually)
• herpes simplex
Disseminated gonococcal infection
Hepatitis A, B, C, D or E
Influenza
Other virus infections

DxT: fever + severe malaise + lymphadenopathy ° acute HIV

AIDS - defining conditions

The original US Centers for Disease Control (CDC) classification has been modified with time to provide a more simplified scheme for defining AIDS. The HIV/AIDS case surveillance system simply specifies a list of clinical conditions associated with the late stages of HIV infection as being 'AIDS-defining'.[7]

The AIDS-defining conditions are:

- candidiasis of bronchi, trachea or lungs
- candidiasis, oesophageal
- cervical cancer, invasive
- coccidioidomycosis, disseminated or extrapulmonary
- cryptococcosis, extrapulmonary
- cryptosporidiosis, chronic intestinal (> 1 month's duration)
- cytomegalovirus (CMV) disease (other than liver, spleen, or nodes)
- CMV retinitis (with loss of vision)
- encephalopathy, HIV-related
- herpes simplex virus (HSV): chronic ulcer(s) (> 1 month's duration); or bronchitis, pneumonitis or oesophagitis
- histoplasmosis, disseminated or extrapulmonary
- isosporiasis, chronic intestinal (> 1 month's duration)
- Kaposi's sarcoma
- lymphoma, Burkitt's (or equivalent term)

- lymphoma, immunoblastic (or equivalent term)
- lymphoma, primary, of brain
- Mycobacterium avium complex of M. kansasii, disseminated or extrapulmonary
- Mycobacterium tuberculosis, any site (pulmonary or extrapulmonary)
- Mycobacterium, other species or unidentified species, disseminated or extrapulmonary
- Pneumocystis carinii pneumonia
- Salmonella septicaemia, recurrent
- toxoplasmosis of brain
- wasting syndrome due to HIV

The Australian AIDS surveillance case definition does not refer to the CD_4 cell count although in the United States AIDS is also defined by a CD_4 cell count of < 200/μL, regardless of clinical condition.

Common clinical presentations of HIV infection

Fever

Usually this is of unknown origin.

Weight loss

This is usually severe and muscle wasting.

Respiratory

- Sinusitis
- Non-productive cough, increasing dyspnoea and fever: due to opportunistic pneumonias

More than 50% of patients present with PCP which may have an abrupt or insidious onset.[6] With the insidious type of onset, examination and chest X-ray are often normal early. Many other agents (e.g. CMV, cryptococcosis and TB) can be responsible. Exclusion of PCP is important as this condition carries a high mortality if untreated.

Gastrointestinal

- Chronic diarrhoea (many causes) with weight loss or dehydration

Neurological

- Headache
- Progressive dementia (HIV encephalopathy)
- Ataxia due to myelopathy
- Seizures
- Mononeuritis
- Guillain-Barré type mononeuropathy
- Toxoplasma encephalitis
- Cryptococcal meningitis
- Peripheral neuropathy
- Progressive visual loss (CMV retinitis)
- CNS lymphoma

Oral cavity

- Aphthous ulcers
- Angular chelitis
- Periodontal/gingival disease
- Tonsillitis
- Oral candidiasis
- Oral hairy cell leukaemia (frequently mistaken for candidiasis but affects lateral border of tongue)

Genitourinary

- Cervical dysplasia
- Vaginal candidiasis
- Various STIs (e.g. HSV, HPV)

Skin

- Impetigo
- Warts
- HSV
- Shingles, especially multidermatomal
- Seborrhoeic dermatitis
- Cutaneous mycoses
- Kaposi's sarcoma (painless red-purple lesions on any part of the body including palms, soles, oral cavity and other parts of the GIT) (Figure 28.2).



Figure 28.2 Kaposi's sarcoma of the skin on the face of a man with AIDS. Photo courtesy Hugh Newton-John

Figure 28.3 presents the chronology of HIV-induced disease correlated with time since infection and CD_4 cell levels.



Figure 28.3 Chronology of HIV-induced disease correlated with time since infection G.J. Steward, Could It Be HIV? 1. The Challenge: Clinical Diagnosis of HIV. The Medical Journal of Australia, 1993: 158: 31 - 4 © Copyright 1993, The Medical Journal of Australia - reproduced with permission

Investigations and diagnosis[6]

The laboratory investigation of AIDS covers three broad areas:

1. Tests for HIV infection:
    - ELISA
    - Western blot technique (used for confirmation)
2. Tests of immune function:
    - CD_4 lymphocyte counts - the strongest predictor of possible clinical manifestations of HIV infection
    - low CD_4 cells (counts < 500 cells/μL) ° defective cell immunity[2, 4]
    - counts < 200 cells/μL ° severe immunodeficiency
3. Viral load: a measure of the serum level of RNA of the HIV virus - correlates with response to treatment and progression to AIDS and death
4. Tests for opportunistic infections and other problems: (e.g. other STIs, EBV, CMV, hepatitis, Mantoux test)

MANAGEMENT

Patients with HIV infection require con-siderable psychosocial support, counselling and regular assessment from a non-judgmental caring practitioner.

The holistic approach to life

Most people with HIV infection will take 'natural therapies'. This should be viewed as being complementary with the management suggested by the GP, and the patient should be encouraged to tell his or her doctors of the alternative medicines being taken. Anecdotal reports suggest that 75% of people with HIV regularly use 'natural therapies',[8] and in the setting of the long-term nature of the condition it is important for doctors to be supportive and create a climate of acceptance around these practices.

Positive lifestyle factors include:

- a very healthy balanced diet: high fruit and vegetable intake, pure fruit juices, high fibre, low fat, high complex carbohydrates

- toxic avoidance: processed foods, caffeine, illicit drugs, alcohol, cigarettes
- relaxation and meditation (reduction of and self-monitoring of stress levels)
- appropriate sleep and exercise
- consider supplementary antioxidants
- support groups and continuing counselling

Medication[8, 9, 10]

Optimal antiretroviral therapy now depends on the use of combinations of drugs. Monotherapy is no longer accepted practice. The rec-ommendations for the use of antiretroviral therapy are constantly changing. Updated guidelines can be found on the Internet at < http://www.hivatis.org >. Viral resistance is the limiting factor, no matter how potent an individual drug may be at reducing viral load initially. The trials of combined zidovudine and lamivudine demonstrated both a more sustained decrease in plasma viral load than either drug did alone, and a more delayed development of viral resistance. There are now many antiretroviral drugs available for use in Australia (Table 28.4) and clinicians have a much wider scope of treatments available. However, many questions remain about combination therapy and further trials using viral load as a clinical end point should provide pointers for treatment. Currently the use of three drugs is favoured (HAART). Sideeffects, after serve that effect the quality of life … a problem. There are many possible combinations. Subcutaneous injections of in-terleukin-2 have been shown to boost immunity.

Table 28.4 Currently available antiretroviral drugs[9]

Nucleoside RT* inhibitors (NRTIs)	Non-nucleoside RT inhibitors (NRTIs)	Protease inhibitors	HIV entry inhibitors
• abacavir	• nevirapine (NEV)	• saquinavir (SQV)	• enfuvirtide (T20)
• zidovudine (AZT)	• delavirdine (DLV)	• indinavir (IDV)	
• didanosine (DDI)	• efavirenz	• ritonavir (RTV)	
• emtricitabine		• nelfinavir (NFV)	
• zalcitabine (DDC)		• amprenavir (APV)	
• stavudine (D4T)		fosamprenavir (FAPV)	
• lamivudine (3TC)		• lopinavir/ritonavir (LPV)	
• AZT + 3TC (Combivir)		atazanavir (ATZ)	
• AZT + 3TC + abacavir (Trizivir)		• tipranavir (TPV)	
• Tenofir (TFV)**			

* RT ° reverse transcriptase ** a nuclotide analogue RTI

HAART (highly active antiretroviral therapy)

This is a combination of three (or more) agenst with one or more penetrating the blood-brain barrer.

Pneumocystis carinii[9]

This is an important cause of preumohia and not usually seen until the CD4 cell count is < 200/ml. It is usually treated with trimethoprim + sulfanitoxazole (co-trmoxazd) which is also given orally as prophyalxis when the cell count reaches <200.

Acute HIV Infection

Treatment of seroconversion illness is not of proven clinical benefit (to date) but is optional and some clinics offer it.

THE HIV TEST: THE ROLE OF THE FAMILY DOCTOR

The astute GP will use the opportunity of a request for an HIV test to explore preventive and sexual health issues. A full sexual history and drug history must incorporate the three C's of counselling, confidentiality and consent in the pretest interview.

Many HIV-positive patients have described how the results left them bewildered and devastated, especially with an unexpected positive result. Part of the reason given was the lack of any form of pretest counselling.

Initial consultation

- First establish why the patient is presenting ‘now’ for the test.

- Explore the ‘hidden component’ of the patient's consultation.
- Take a full sexual, medical and drug-taking history. It is recognised that this can be embarrassing for both the doctor and the patient, but those experienced in this process advise the following approach:
  - Establish a supportive, non-judgmental atmosphere. Encourage disclosure of history and patterns of partners and sexual practices in a gender-neutral situation. Make no assumptions about sexual preferences; they will be indicated by the patient as the history evolves, provided you allow this to happen; this will take time.
- Non-judgmental matter-of-fact questions such as ‘Have you injected yourself with drugs?’ and (to a male patient) ‘Do you have sex with men or women or both?’ may permit honest disclosure.
- Stress the importance of disclosure of prior, known infections with STIs. Assess the patient's risk for an STI.
- Assess the patient's coping strategies and social network.

Pretest counselling

- Give information on the test (what it tells and does not tell).
- Explain about the false negative and ‘window period’.
- Give appropriate information about HIV disease and other STIs.
- Dispel any myths about transmission of infection.
- Give preventive advice on safer practices (sex and IV drugs).
- Assess the possible coping mechanisms of the patient.
- Assess the patient's social support networks and interpersonal bonds.
- Reassure about confidentiality. This is a legal requirement.
- Discuss who to tell: informing sexual contacts.
- Offer tests other than STIs.

Finally

- Discuss how the patient ‘will cope with the test result’.
- Discuss legal requirements (check with state laws).
- Advise of need for informed consent (not only for HIV test but other STIs).
- Make arrangements to discuss the test results.

Consider the useful question: ‘How would your behaviour change, if at all, as a result of having this test today?’

Test result (about 2 weeks later)

This must be given in consultation (whether positive or negative): avoid the telephone.

The negative test result

- Provide reassurance.
- Emphasise the safe sex information.
- Counter any suggestion that current risk-taking behaviour is safe.
- Retest if in high-risk category or known HIV contact or in a ‘window period’ of 12 weeks.
- A test in 3 months helps rule out recent acquisition.
- Maintain confidentiality.

The intermediate test result[11]

Intermediate or equivocal renulls indicate a possible ‘window’ penod.

The renualls need ingoing monitoring including repeat testing. Ensure that a face-to-face meeting is arrounded to discum the defimetic result.

The positive test result

This is a very traumatic experience and good communication skills are essential. The result should be stated clearly. Good pretest history taking, risk assessment and counselling will make the task of informing the patient of a positive test result a lot easier. Being able to inform the patient of an improved prognosis due to combination drug therapy is also of assistance in breaking the news. Be warm and open. Many patients want to be touched for reassurance. The words ‘you are not dying’ can be firmly stated a few times. Educate the patient about the difference between HIV infection and the acquired immuno deficiency syndrome.

Discuss with the patient to whom he or she is going to tell the result and help the patient decide on the most supportive friend. Ask the patient what he or she is going to do aft er the consultation and make an appointment for the next day. Give the patient an HIV support line number for overnight telephone support if required. Discuss the issue of contact tracing briefly and in more depth at the next consultation, because sexual contacts should be notified. However, avoid an information overload.

Start immediate post-test counselling and education. Emphasise that an HIV-positive status does not equal AIDS and need not lead to it in the short term. Organise a consultant's appointment and provide sound patient education material.

Interview and full clinical assessment

- Assessment of general health
- Particular assessment of prior psychiatric history and confirm prior STIs and drugs
- Evidence of EBV (glandular fever), hepatitis B, HIV illness (acute seroconversion illness)
- Further counselling and discussion of specific problems
- Assessment of bonds and relationships with regard to support

Examination - to set a base level

- Full examination, skin, CNS - especially fundi, chest, abdomen and genitals; urine and lung function test
- Monitor temperature and weight

Blood tests - to set a base level and check immune status

- Repeat HIV antibody test (to eliminate possibility of error)
- CD_4 cells with FBE and a differential WCC
- Viral load test
- G-6-PD screen for enzyme deficiency
- Serology for syphilis (RPR), hepatitis A, B and C, toxoplasmosis, CMV
- Test for gonorrhoea and Chlamydia, herpes and thrush (if indicated)
- Mantoux test for tuberculosis

Encourage a holistic approach to health maintenance and enhancement. Explore their feelings, anxieties, fears and confidentiality concerns. Reinforce safer practices.

The second post-positive result consultation (1 - 2 weeks later)

- Give results of repeat HIV test and baseline tests.
- Explore patient's understanding, feelings, coping abilities and spiritual issues (if appropriate).
- Some of the common questions that patients with HIV infection ask are:
    - Am I going to get sick?
    - How long will I live for?
    - Are there any treatments available?
    - Is there a cure around the corner?
    - What is going to happen to me?
    - Should I tell my friends?
    - Should I tell my family?
    - What are the social and legal issues?

It is worth pre-empting these questions and having ready appropriate answers for that particular patient.

- Give appropriate reassurance: prognosis may be much better, in respect of long remission, than appreciated.
- Discuss support systems.
- Check personal prophylaxis (safer sex and needle-sharing habits).
- Reinforce lifestyle strategies and suggest how patients can help themselves: give case examples.
- Recommend meditation and appropriate literature.
- Provide appropriate referrals (if needed):
    - specialist counsellors
    - self-help and support groups
    - meditation classes

- Advise patients about their legal and ethical responsibilities not to pass on the infection to others.
- Organise contact tracing.
- Address the difficult issue of telling sexual partners.
- If the patient is unwilling to inform sexual partners or is uncertain of who they may be, then contact-tracing organisations run through state government offices are of assistance in tracing sexual partners who may be at risk.
- If a patient refuses to inform a sexual partner of the risk, then the doctor may disclose this information to the patient's partner in the following circumstances: if there is a clear risk of transmission; if the patient has been given education and counselling and this has been ignored; or if the doctor in the case has sought advice from colleagues and institutional ethics committee and before disclosure discussed it with the medical defence organisation. The doctor then should provide the patient with written advice that the patient must notify the partner, and if the patient still refuses to do so then the doctor has the right to do so.
- Discuss ‘safer sex’ guidelines. Condom usage is essential to prevent further viral loading and this needs to be explained, as well as the reason of protecting others.

Continuing maintenance consultations

- Provide appropriate support, encouragement and counselling.
- Frequency depends on CD_4 cell count (e.g. 3 to 6 monthly).
- Examination:
  - Check general condition, temperature and weight.
  - Look for unusual lung infection, diarrhoea, skin lesions, tongue and oropharynx, fevers, wasting and neurological signs.
  - Examine for signs of CMV retinitis (at risk if CD_4 count < 100 cells/?L).
  - Monitor for depressive illness.
  - Look for early signs of AIDS-related dementia.
- Tests
  - CD_4 cell count and syphilis serology
  - viral load test
  - chest X-ray and induced sputum (if cough, shortness of breath), faeces microculture if diarrhoea persists, Candida mouth swabs and herpes swabs appropriately
- Treat intercurrent illness (Table 28.5).

Table 28.5 General practice treatment of HIV-related dermathopathy

Dermatoses	Treatment
Common	
Oral candidiasis	miconazole oral gel, 6 hrly, 10 - 14 days or amphotericin lozenges or fluconazole 50 mg (o), daily, 10 - 14 days
Hairy leucoplakia on tongue	no treatment or aciclovir
Seborrhoeic dermatitis (e.g. nasolabial folds, hairy areas)	miconazole 2% or ketoconazole 2% hydrocortisone 1% (if very itchy)
Itchy folliculitis	oral antihistamines miconazole + hydrocortisone topical
Less common	
Perianal herpes simplex	Mild: topical aciclovir cream (if available) Betadine or silver nitrate Severe: aciclovir 200 mg 5 times daily
Molluscum contagiosum	needle incision then Betadine or liquid nitrogen
Candidal angular stomatitis	clotrimazole or miconazole cream
Herpes zoster	Mild: menthol in flexible collodion Severe: refer for IV aciclovir or other guanine analogue
Severe tinea pedis	topical clotrimazole or miconazole ketoconazole 200 mg (o) daily

Source: After Pohl[4]

- Prophylaxis - this is managed according to immune status: if CD_4 count < 200 cells/μL use cotrimoxazole to prevent opportunistic infections, particularly PCP.

Contact tracing

Contacts of HIV-positive patients should be traced and offered testing with counselling.[5] Patients with HIV infection must be advised of the risk they pose to seronegative sexual partners. A person who has HIV or is at risk of HIV infection must not make any blood, semen or tissue donation. Because of the probable association between genital ulcerative disease and HIV transmission, the effective management of STIs is part of the general strategy for HIV control.

PREVENTION OF HIV INFECTION

Counselling the person at risk re ‘safer practices‘

No effective vaccine has been developed. Modification of behaviour is the only valid strategy for prevention of HIV infection. Education programs to encourage sexual practices that reduce the exchange of genital secretions (safe sex) may achieve risk reduction for sexually active individuals. Condoms provide a barrier if used properly and consistently but may be too easily damaged to offer reliable protection during anal intercourse. A water-based lubricant such as KY gel or Lubafax should be used: oil-based lubricants such as Vaseline weaken condoms.

Discuss alternative sex practices, including touching, cuddling, body-to-body rubbing and mutual masturbation.

Emphasise the importance of being in control with drug taking, IV usage, safe sex practices and the needle-exchange program.

Of special importance is the finding that the most important biological risk factor for HIV transmission is the presence of other active STIs in either partner. This includes chlamydia, gonorrhoea, syphilis and genital herpes. Herpes is also likely to increase the risk of HIV transmission during both homosexual and heterosexual intercourse, even in the presence of condoms.[12, 13]

Health professionals

Care should be exercised whenever blood samples are taken or sharps have been used. Advise safe disposal of sharps and other disposables and appropriate sterilisation of material. Gloves should be worn for all invasive procedures. Management of needle-stick injuries and other at-risk exposures is described in Chapter 134. Blood donors need to be carefully screened.

COMMUNITY EDUCATION

Educating the community in a non-emotional, responsible way about AIDS should be a priority. While the personal, community and global benefits of effective AIDS education are generally acknowledged, the fear of addressing such a sensitive issue sometimes results in failure to act.[13] AIDS education in schools in particular can be an important strategy. People with HIV infection would be appropriate resource educators and the use of videos would be a most appropriate medium for education.

WHEN TO REFER[11]

Most patients with HIV disease need referral to a specialist or clinic that can manage the patient expertly and sympathetically.

Referral should take place at the time of:

- onset of a life-threatening opportunistic infection
- the need to initiate antiretroviral drug therapy
- administration of prophylactic pentamidine aerosol therapy
- serious psychological problems related to HIV-positive status.

ACKNOWLEDGMENT

Part of this text, including clinical mani-festations, prevention and contact tracing is reproduced from the Handbook on Sexually Transmitted Diseases.[5] Commonwealth of Australia copyright, reproduced by permission.

REFERENCES

1. Boyle MJ, McMurchie M, Tindall B, Cooper D. HIV seroconversion illness. Med J Aust, 1993; 158: 42 - 4.
2. Stewart GJ. The challenge: the clinical diagnosis of HIV. Med J Aust, 1993: 158: 31 - 4.
3. Kumar P, Clark M. Clinical Medicine (2nd edn). London: Saunders 2005; 132-133.
4. Pohl M. Managing HIV patients in general practice. Patient Management, 1989; June: 49 - 61.
5. NH&MRC. Handbook on Sexually Transmitted Diseases. Canberra: Department of Community Services Health, 1990: 1 - 55.
6. McCoy R. Alarm bells. When to worry about your patient with HIV. Aust Fam Physician, 1997; 26: 803 - 9.
7. http://www.code.gov (us centers for disease control and preventum update on HIV/AIDS developments)
8. Bradford D. Update on issues for HIV management. Aust Fam Physician, 1997; 26: 812 - 17.
9. Spicer J (Chair). Therapeutic Guidelines: Antibiotic (Version 12). Melbourne: Therapeutic Guidelines Ltd, 2005: 93 - 106.
10. Kidd M, McCoy R. Managing HIV/AIDS Part 2-Treatment. Medical observer, 1 March 2002: 36-7
11. HIV/AIDS ‘Let’s talk about it: Asitm, 2005. Available from < http://www.ashm.org.au >
12. Rogers G.L HIV care and prevention part 1. Australian Doctor, 2006; 3 February :25-32
13. World Health Organization report. AIDS Prevention through Health Promotion. Geneva: WHO, 203. < http://www.who.int/topics/hiv_intections >
14. Bradford DL. Acquired immune deficiency syndrome. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 1 - 5.

CHAPTER 29. BAFFLING VIRAL AND PROTOZOAL INFECTIONS

Our lot is a perilous age … but where shall we fly to escape from pestilences that come and pestilences that do not come, from ships that bring us yellow fever, from cattle diseases that can only be exterminated by exterminating the cattle, from infectious patients whose pulses must be felt with a pair of tongs and their chests explored with tarred stethoscopes.

JACOB BIGELOW 1860

BAFFLING VIRAL AND PROTOZOAL INFECTIONS

Almost any infection, especially if subacute or insidious in its onset, can be baffling and can belong to the 'fever of undetermined origin' group of infections. Syphilis and tuberculosis were the great mimics of the past. Now malaria and Epstein-Barr mononucleosis can be regarded as important mimics. EBM (infectious mononucleosis, glandular fever) can be a perennial baffler and can be confused with HIV infection in its primary clinical phase. Any of the febrile diseases can be confusing before declaring themselves with classic symptoms such as the jaundice of hepatitis or the rash of dengue fever, or before serological tests become positive.

Viral and protozoal infections that can present as masquerades include:

- HIV infection (especially primary)
- EBM
- TORCH organisms
    - toxoplasmosis
    - rubella
    - CMV
    - HSV
- hepatitis A, B, C, D, E
- mosquito-borne infections
    - malaria
    - dengue fever
    - yellow fever
    - Japanese encephalitis
    - Ross River fever

The TORCH organisms (TORCH being an acronym for toxoplasmosis, rubella, CMV and herpes) are well known for their adverse intrauterine effects on the fetus. Three are viral (toxoplasmosis is a protozoa) and the first three of these fetal pathogens are acquired by passage across the placenta. Most of these organisms are noted for being opportunistic infections in immunocompromised patients, especially in later stage HIV infection.

The mosquito-borne infections are mainly viral, apart from the protozoa causing malaria, and are of particular significance in travellers returning from endemic areas.

The major protozoal diseases of man are:

- blood:
    - malaria
    - trypanosomiasis
- GIT:
    - giardiasis
    - amoebiasis
    - cryptosporidium
- tissues:
    - toxoplasmosis
    - leishmaniasis

FOUR SIMILAR CLINICAL PRESENTATIONS

Four infections - EBM, primary HIV, CMV and toxoplasmosis - produce almost identical clinical presentations and tend to be diagnosed as glandular fever or pseudoglandular fever. It is important for the first contact practitioner to consider all four possibilities, especially keeping in mind the possibility of HIV infection.

A worthwhile approach is to make a provisional diagnosis based on the clinical variations as presented in Table 29.1.

Table 29.1 Clinical features differentiating HIV, EBV, CMV and toxoplasmosis infections (all can present with a similar illness)

Feature	EBV infection	HIV infection	CMV infection	Toxoplasmosis
Onset	Insidious	Acute	Insidious	Insidious or acute
Fever	A feature Intermittent	A feature	Quotidian (afternoon spikes)	Low grade
Fatigue/malaise	Common	Common, severe	Common	Common
Tonsillar hypertrophy	Common	Mild enlargement	Uncommon	Uncommon
Exudative pharyngitis	Common	Rare	Rare	Occurs
Mucocutaneous ulcers	Rare	Common	Unknown	Unknown
Skin rash	About 5%	Common	About 5%	About 10%
Jaundice	About 8%	Rare	Uncommon	Uncommon
Diarrhoea	Unknown	Occurs	Unknown	Unknown
Cervical lymphadenopathy	Common	Common	Uncommon	Common (a feature)
Hepatomegaly	About 8%	Rare	Common	Occasional
Splenomegaly	About 50%	Rare	About 50%	Up to 30%
Atypical lymphocytes	In 80 - 90%	In < 50%	Common	Uncommon

Screening tests are:

- FBE, especially WCC
- EBV-specific antibodies
- serological test for CMV (specific antibodies)
- serological test for toxoplasmosis (specificantibodies - acute and convalescent)
- HIV antibody test (ELISA)
- PCR testing

Epstein-Barr mononucleosis

EBM (infectious mononucleosis, glandular fever) is a febrile illness caused by the herpes (Epstein-Barr) virus. It can mimic diseases such as HIV primary infection, streptococcal tonsillitis, viral hepatitis and acute lymphatic leukaemia. There are three forms: the febrile, the anginose (with sore throat Figure 29.1) and the glandular (with lymphadenopathy).



Figure 29.1 Tonsillitis of Epstein-Barr mononucleosis. This is often confused with bacterial tonsillitis

It may occur at any age but usually between 10 and 35 years, commonest in 15 - 25 year old age group.

The typical clinical features are presented in Table 29.2 and Figure 29.2.

Table 29.2 Clinical features of EBM[1, 2]

Symptoms
• Slow onset malaise 1 - 6 weeks
• Fever
• Myalgia
• Headaches, anorexia
• Blocked nose - mouth breathing
• Nasal quality to voice
• Sore throat (85%)
• Nausea ± vomiting
• Rash - primary 5%
• Dyspepsia
Clinical findings
• Exudative pharyngitis (84%)
• Petechiae of palate (not pathognomonic) (11%)
• Lymphadenopathy, especially posterior cervical
• Rash - maculopapular
• Splenomegaly (50%)
• Jaundice ± hepatomegaly (5 - 10%)
• Clinical or biochemical evidence of hepatitis



Figure 29.2 Clinical features of Epstein-Barr mononucleosis

DxT: sore throat + fever + lymphadenopathy ° EBM

Epidemiology

EBM has an annual incidence of 4 - 5 new cases in a population of 2500.1 It usually affects people in their late teenage years or early twenties. It is endemic in most countries. Subclinical infection is common in young children. The incubation period is at least 1 month but data is insufficient to define it accurately.

EBV is excreted in oropharangeal secretions during the illness and for some months (sometimes years) after the clinical infection. EBM has a low infectivity and isolation is not necessary. It is apparently transmitted only by close contact, such as kissing and sharing drinking vessels.

Progress of the primary infection is checked partly by specific antibodies (which might prevent cellto-cell spread of the virus) and partly by a cellular immune response, involving cytotoxic T-cells, which eliminates the infected cells. This response accounts for the clinical picture. The virus is never eliminated from the body.

Second attacks and fatalities do occur and there is a possible association between EBM and lymphoma.[3]

The rash

The rash of EBM is almost always related to antibiotics given for tonsillitis. The primary rash, most often non-specific, pinkish and maculopapular (similar to that of rubella), occurs in about 5% of cases only.

The secondary rash is most often precipitated by one of the penicillins, especially ampicillin or amoxycillin. About 90 - 100% of patients prescribed ampicillin or amoxycillin will be affected; up to 50% of those given penicillin will develop the rash. It can be extensive and sometimes has a purplish tinge.

The complications of EBM are presented in Table 29.3 and the differential diagnoses in Table 29.4.

Table 29.3 Complications of EBM[1]

Common
• Antibiotic-induced skin rash
• Prolonged debility
• Hepatitis
• Depression
Rare
Cardiac:
• myocarditis
• pericarditis
Haematological:
• agranulocytosis
• haemolytic anaemia
• thrombocytopenia
Respiratory tract:
• upper airway obstruction (lymphoid hypertrophy)
Miscellaneous:
• ruptured spleen
Neurological:
• cranial nerve palsies, especially facial palsy
• Guillain-Barré syndrome
• meningoencephalitis
• transverse myelitis

Table 29.4 Differential diagnoses of EBM[1]

Other agents that cause typical EBM syndrome:
• HIV infection (acute initial illness)
• CMV
• toxoplasmosis
Exudative tonsillitis resembling EBM:
• acute streptococcal pharyngitis
• adenovirus infection
• diphtheria (unlikely in Australia)
Hepatitis A, B, C, D, E
Lymphadenopathy, fever and splenomegaly:
• lymphoma
• leukaemia
Complications of EBM without other manifestations:
• encephalitis
Others:
• drug reaction
• influenza

Laboratory diagnosis

The following confirm the diagnosis of EBM:

- WCC shows absolute lymphocytosis.
- Blood film shows atypical lymphocytes.
- Paul Bunnell test for heterophil antibody is positive (although positivity can be delayed or absent in 10% of cases).
- Diagnosis confirmed (if necessary) by EBVspecific antibodies, viral capsule antigen (VCA) antibodies - IgM, IgG and EB nuclear antigen (EBN-A).

Culture for EBV and tests for specific viral antibodies are not done routinely.

False positives for the Paul Bunnell test are:

- hepatitis
- Hodgkin's lymphoma
- acute leukaemia

Prognosis

EBM usually runs an uncomplicated course over 6 - 8 weeks. Major symptoms subside within 2 - 3 weeks. Patients should be advised to take about 4 weeks off work.

Treatment

- Supportive measures (no specific treatment)
- Rest (the best treatment) during the acute stage, preferably at home and indoors
- Aspirin or paracetamol to relieve discomfort
- Gargle soluble aspirin or 30% glucose to soothe the throat
- Advise against alcohol, fatty foods, continued activity, especially contact sports (risk of splenic rupture)
- Ensure adequate hydration
- Corticosteroids for:
    - - neurological involvement
    - - thrombocytopenia
    - - threatened airway obstruction

Post-EBM malaise

Some young adults remain debilitated and depressed for some months. Lassitude and malaise may extend up to a year or so.

Cytomegalovirus infection

CMV has a worldwide distribution and causes infections that are generally asymptomatic. The virus (human herpes virus 5) may be cultured from various sites of healthy individuals. It has its most severe effects in the immunocompromised, especially those with AIDS, and also in recipients of solid organ transplants and bone marrow grafts; 90% of AIDS patients are infected with CMV and 95% have disseminated CMV at autopsy. CMV infection can be an important development following massive blood transfusion, including those given to infants. The incubation period of CMV ranges from 20 to 60 days and the illness generally lasts about 2 to 6 weeks.[2]

Clinical features

Three important clinical manifestations are described.

1 Perinatal disease

Intrauterine infection may cause serious abnormalities in the fetus, including CNS involvement (microcephaly, hearing defects, motor disturbances), jaundice, hepatosplenomegaly, haemolytic anaemia and thrombocytopenia. Up to 30% of CMV-affected infants have mental retardation[3] (see p. 1047).

2 Acquired CMV infection

In healthy adults CMV produces an illness similar to EBM with fever, malaise, arthralgia and myalgia, generalised lymphadenopathy and hepatomegaly. However, cervical lymphadenopathy and exudative pharyngitis are rare.

The infection may be spread by blood transfusion, and CMV should be suspected on clinical grounds in a patient with a febrile illness resembling EBM following major surgery, such as open heart surgery or kidney transplantation, and where extensive transfusion has been necessary.

The fever often manifests as quotidian intermittent fever spiking to a maximum in the mid-afternoon and falling to normal each day (Figure 29.3). There is often a relative lymphocytosis with atypical lymphocytes but the heterophil antibody test is negative. Liver function tests are often abnormal.



Figure 29.3 Cytomegalovirus infection: typical quotidian intermittent fever pattern

Diagnosis: Specific diagnosis can be made by demonstrating rising antibody titres from acute and convalescent (2 weeks) sera. A four-fold increase indicates recent infection. PCR testing can be used. The virus can be isolated from the urine and blood.

3 CMV disease in the immunocompromised host

Disseminated CMV infection occurs in the immunedeficient person, notably HIV infection causing opportunistic severe pneumonia, retinitis (a feature of AIDS), encephalitis and diffuse involvement of the gastrointestinal tract.

Treatment

In the patient with normal immunity no treatment apart from supportive measures is required, as the infection is usually self-limiting. In immunosuppressed patients various antiviral drugs, such as ganciclovir and foscarnet, have been used with some benefit.

Toxoplasmosis

Toxoplasmosis, which is caused by *Toxoplasma gondii*, an obligate intracellular protozoan, is worldwide, albeit a rare infection. The definitive host in its life cycle is the cat (or pig or sheep) and the human is an intermediate host. However, clinical toxoplasmosis is very uncommon. Infection in humans usually occurs through eating foodstuffs contaminated by infected cat faeces. Its main clinical importance is an opportunistic infection.

The five major clinical forms of toxoplasmosis are:4

1. asymptomatic lymphadenopathy (the commonest)
2. lymphadenopathy with a febrile illness, similar to EBM
3. acute primary infection: a febrile illness similar to acute leukaemia or EBM; a rash, myocarditis, pneumonitis, chorioretinitis and hepatosplenomegaly can occur
4. neurological abnormalities - includes headache and neck stiffness, sore throat and myalgia
5. congenital toxoplasmosis - this is a rare problem but if it occurs it typically causes CNS involvement and has a poor prognosis

In the immunocompromised, clinical forms 3 and 4 are typical features with meningoencephalitis being a serious development.

Diagnosis

Diagnosis is by serological tests, which are sensitive and reliable.

Treatment

Patients with a mild illness or with asymptomatic infection require no treatment. Children under 5 years may be treated to avoid the possible occurrence of chorioretinitis. Symptomatic patients are treated with pyrimethamine plus sulphadiazine. Spiramycin is usually used in pregnant patients.

MALARIA

Malaria, a protozoal infection, is an important masquerade and mimic. The diagnosis requires a high index of suspicion in a traveller returning from an endemic area (see p. 122).

Five hundred million people are infected every year.

In humans malaria is caused by four species of plasmodium:

- *Plasmodium vivax* and *P. ovale* - tertian malaria
- *P. falciparum* - malignant tertian malaria
- *P. malariae* - quartan malaria

Some clinical features

- It has an incubation period of 7 to 40 days.
- Most present within 2 months of returning from malaria area.
- It can present after 2 or more years.
- It is an acute febrile illness with headache.
- It can have atypical presentations such as gastrointestinal symptoms (diarrhoea, nausea, vomiting, abdominal pains), dry cough and arthralgia.
- Typical relapsing patterns may be absent.

Special notes

- Beware of treatment-modified infection.
- Early diagnosis is essential as it must be treated within 4 days.

Diagnosis

Diagnosis is with thick and thin smears. At least three blood samples should be taken each day for 3 to 4 days and found to be negative before a patient is declared malaria-free. The presence of monocytosis is a helpful diagnostic pointer.

Various PCR methods are in evolution.

AMOEBIASIS

Amoebiasis can be diagnosed in a sick traveller returning from an endemic area with severe diarrhoea characterised by blood and mucus. Complications include fulminating colitis, amoebomas (a mass of fibrotic granulation tissue) in the bowel and liver abscess.

Amoebic liver abscess

Clinical features

- High swinging fever
- Profound malaise and anorexia
- Tender hepatomegaly
- Effusion or consolidation of base of right chest

There is often no history of dysentery, and jaundice is unusual. Diagnosis is by serological tests for amoeba and by imaging (CT scan). Treatment is with metronidazole and by percutaneous CT-guided aspiration.

DENGUE FEVER

Also known as 'breakbone fever', dengue is a mosquito-borne viral infection found mainly in Asia and Africa and also prevalent in tropical North Queensland (see p. 123). The features are the acute onset of fever, headache, retrobulbar pain, severe backache and aching of muscles and joints. Lymphadenopathy, petechiae on the soft palate and skin rashes may occur.

The diagnosis of dengue is based on clinical suspicion and also by dengue-specific IM antibody. Severe fatigue and depression (with suicidal risk) is a feature of convalescence. The patients are managed symptomatically and with support and reassurance.

JAPANESE ENCEPHALITIS

This viral illness is also acquired overseas, especially in South-East Asia and the Far East. The onset is heralded by severe rigors, fever, headache and malaise. The encephalitic stage is marked by fever, neck rigidity and neurological signs, such as altered consciousness and convulsions. Coma can follow. Diagnosis is by specific antibody detection. Treatment is symptomatic.

YELLOW FEVER

This viral illness, acquired in the tropical areas of Africa and South America, can have a variety of presentations. If yellow fever is mild it resembles influenza. However, the typical clinical features are a febrile illness - high fever, headache, flushed face, arthralgia, GIT symptoms, relative bradycardia (Faget's sign) - followed by apparent recovery and then the classic features of fever, deepening jaundice, proteinuria and bleeding from the gums.

ROSS RIVER FEVER

Epidemic polyarthritis or Ross River virus, which is an alpha virus, occurs in all states of Australia. It is most prevalent in mosquito-prone areas (especially during the summer) and in tropical and temperate coastal regions and inland riverine areas.[5] Subclinical infection is common with variable clinical manifestations.

Clinical features

- All age groups, especially 20 - 30 years
- Incubation period 3 - 21 days (usually 7 - 11)

Major symptoms

- Polyarthritis (75% of patients) - mainly fingers, wrists, feet, ankles and knees
- Maculopapular rash - widespread, often 'subtle', mainly trunk and limbs
- Myalgia

DxT: polyarthritis + fever + rash ° Ross River fever

Other symptoms

- Pyrexia (mild)
- Headache
- Nausea
- Fatigue with exercise

Signs (which may be present) include joint swelling (mainly hands and feet), tenosynovitis around the wrists and ankles (poor prognostic sign), the rash and mild lymphadenopathy.

Outcome

In many patients the illness resolves within 2 to 4 weeks and most feel normal within 3 months, but some with a more severe arthritis can enter a chronic phase lasting 18 months or more.

Diagnosis

Diagnosis is by antibody testing of serum. The differential diagnosis includes other viral infections that cause arthritis, such as hepatitis B, rubella, Barmah Forest virus (a mosquito-borne virus) and dengue, and early rheumatoid arthritis and rheumatic fever.

Treatment

Treatment is symptomatic with bed rest and simple analgesics such as aspirin. NSAIDs are used for more severe cases. Oral corticosteroids are effective but should be avoided if possible.

DEADLY INFECTIONS (WORLDWIDE)

The WHO in its 1996 report[6] estimated that the leading cause of approximately 50 million deaths in the world in 1995 was infectious diseases, which accounted for about one-third or 17 million deaths. Other figures included diseases of the circulatory system (15 million deaths, including ischaemic heart disease - 7.2 million - and stroke - 4.6 million) and cancer (6.2 million deaths, approximately 12%).

The report indicated that nearly 50 000 people are dying every day from infectious diseases such as cholera, malaria, tuberculosis and HIV (Table 29.5). At least 30 new infections have emerged in the past 20 years, and for many of them there is no treatment, cure or vaccine. These include rotavirus (which causes infant diarrhoea), *Legionella pneumophila*, *Lyme borreliosis* (Lyme disease), the Wantaan virus (which can cause a fatal haemorrhagic fever), HIV and hepatitis E and C.

Table 29.5 The world's deadliest infectious diseases

Infectious disease	Cause	Annual deaths
1 Acute lower respiratory infections (mostly pneumonia)	Bacterial or viral	3700000 (approx. 3.4 million children)
2 HIV/AIDS *	Viral	3000000
3 Tuberculosis	Bacterial	2900000
4 Diarrhoeal diseases	Bacterial or viral	2500000
5 Malaria	Protozoal	2100000
6 Hepatitis B	Viral	1100000
7 Measles	Viral	1000000
8 Neonatal tetanus	Bacterial	460000
9 Pertussis	Bacterial	355000
10 Intestinal worm disease		135000

Source: World Health Organization 1998 and 2001 * figures

The report also said that until recently the struggle for control over infectious diseases had seemed almost over, with smallpox eradicated and six other diseases, including polio, leprosy and guinea-worm disease, singled out for eradication within a few years. However, half the world's 6 billion were at risk of many infectious diseases since many that seemed under control such as tuberculosis and malaria were becoming more prevalent. Some such as yellow fever and cholera were striking in new regions. Other infections were resistant to drugs and virtually untreatable. The previously almost forgotten infections such as yersinia (the plague) had re-emerged.

The deadly haemorrhagic fevers that have broken out in isolated endemics include the zoonotic African diseases - Ebola haemorrhagic fever, Marburg haemorrhagic fever and Lassa fever. These are caused by filoviruses and for most no specific treatment is available.

Another serious infection that emerged sporadically was the so-called ‘flesh eating’ *Streptococcus A* infection, which was a particularly virulent strain causing localised destruction of soft tissue.

West Nile encephalitis caused by a mosquito transmitted virus and carried by birds has surfaced in the USA and beyond, causing thousands of cases and hundreds of deaths.

The lessons to be learned include careful surveillance, attention to prevention, especially with effective immunisation programs, rational antibiotic prescribing and care with travelling to developing tropical countries.

SEVERE ACUTE RESPIRATORY SYNDROME (SARS)

SARS is a respiratory syndrome of varying severity, usually presenting with symptoms of atypical pneumonia. It is caused by a unique corona virus. It should be suspected in a patient with a fever >38°C with one or more respiratory symptoms including cough and dyspnoea, and one or more of close contact or a history of travel within 10 days of onset of symptoms.

Other symptoms can be diarrhoea, myalgia, rhinorrhoea, sore throat and chills.

Severe cases (up to 10% mortality) can progress to acute respiratory distress syndrome (ARDS).

Plain X-rays may show pulmonary infiltrates while high resolution CT scans may show typical patterns.

Diagnosis is by PCR testing.

Treatment trials have been with ribavirin and corticosteroids but it is controversial. Intensive care is required for many cases.

AVIAN (BIRD) FLU[7]

This influenza illness has variable clinical features, depending on the viral strain. The deadly strain is H5N1 influenza A strain. It tends to occur in a pandemic.

The clinical spectrum includes fever, upper respiratory symptoms, rapidly progressive pneumonia (approximately 60%) and gastrointestinal tract symptoms.

Diagnosis is confirmed by PCR assay for avian influenza A (H5N1) RNA, viral culture or a four-fold rise in H5-specific antibody.

Treatment options include the M2 ion channel blockers (amantadine and rimantadine) and neuroaminidase inhibitors (oseltamivir and zanamivir).

PRION TRANSMITTED DISEASES[7]

Prions are proteinaceous infectious particles devoid of nucleic acid that can present with a wide spectrum of clinical neurological manifestations. The feature is transmissible spongiform encepalopathy with Creutzfeld-Jakob disease (CJD) being the classic example. The annual incidence is one per million people. Transmission is from contaminated human tissue (e.g. corneal graft), cadaver pituitary human gonadotrophin or eating contaminated beef. There is no specific treatment for prion disease.

DxT: fatigue + psychiatric symptoms + myoclonus ° CJD

Classic clinical features

- Progressive dementia (starts with memory loss)
- Myoclonus
- Fatigue and somnolence
- Variable neurological features (e.g. ataxia, chorea)

Others

- Persistent paraesthesia or dysaesthena
- Akinetic mutism

Diagnosis

- MRI: high-signal intensity in thalami

- CSF: positive 14-3-3 protein immunoassay

REFERENCES

1. Dwyer D. Mononucleosis syndrome. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 317 - 20.
2. Braunwald E et al. *Harrison's Principles of Internal Medicine* (15th edn). New York: McGraw-Hill, 2001: 112 - 14.
3. Tierney LM et al. *Current Medical Diagnosis and Treatment* (45th edn). New York: Lange, 2006: 1358 - 62.
4. Kumar PJ, Clark ML. *Clinical Medicine* (5th edn). London: Saunders, 2003: 140 - 1.
5. Whitby M. Ross River fever. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 452 - 3.
6. World Health Organization. *Annual Report*. Geneva: WHO, May 2003.
7. WHO diseases alert: 16 July 2004. Avian influenza: necessary precautions to prevent human infection of H5N1, need for virus sharing.
   < http://www.who.int/csr/don/2004_07_16/en/index.html >.

CHAPTER 30. BAFFLING BACTERIAL INFECTIONS

In its beginning the malady is easier to cure but difficult to detect, but later it becomes easy to detect but difficult to cure.

NICCOLÒ MACHIAVELLI (1469 - 1527), *On Tuberculosis*

BAFFLING BACTERIAL INFECTIONS

Bacterial infections can present diagnostic brainteasers, and a high index of suspicion is needed to pinpoint the diagnosis. Many are rarely encountered, thus making diagnosis more difficult yet demanding vigilance and clinical flexibility.

The list includes:

- syphilis
- tuberculosis
- infective endocarditis
- the zoonoses (e.g. brucellosis, Lyme disease)
- clostridial infections
  - tetanus
  - gas gangrene
  - puerperal infection
  - botulism
- hidden suppuration
  - abscess
  - osteomyelitis
- mycoplasma infections
  - atypical pneumonia
- *Chlamydia* infections
  - psittacosis
  - non-specific arthritis
  - pelvic inflammatory disease
  - trachoma
  - atypical pneumonia
- legionnaire's disease

Chlamydia and rickettsial organisms have been confirmed as being small bacterial organisms.

SYPHILIS

Although syphilis is uncommon but increasing in the general population it is extremely common in certain indigenous groups and is frequently acquired from homosexual activity or sexual contacts overseas.[1]

It presents either as a primary lesion or through the chance finding of positive syphilis serology. Family physicians should be alert to the various manifestations of secondary syphilis, which can cause difficulties in diagnosis. Congenital syphilis is rare where there is general serological screening of antenatal patients.

Clinical manifestations[2]

Primary syphilis

The primary lesion or chancre usually develops at the point of inoculation after an incubation period averaging 21 days. The chancre is typically firm, painless, punched out and clean. The adjacent lymph nodes are discretely enlarged, firm and non-suppurating. Anorectal changes may occur in homosexual men.

Untreated, early clinical syphilis usually resolves spontaneously within 4 weeks, leading to latent disease, which may proceed to late destructive lesions.

Secondary syphilis

The interval between the appearance of the primary chancre and the onset of secondary manifestations varies from 6 to 8 weeks. Constitutional symptoms, including fever, headache, malaise and general aches and pains, may precede or accompany the signs of secondary syphilis.

The most common feature of the secondary stage of infection is a rash, which is present in about 80% of cases. The rash is typically a symmetrical, generalised, coppery red maculopapular eruption on the face, palms and soles and is neither itchy nor tender. It can resemble any skin disease except those characterised by vesicles. Other features may be:

- condylomata lata, which are broad-based, moist, warty or papular growths occurring in skin folds or creases
- patchy alopecia (scalp, outer third of eyebrow)
- oral, pharyngeal or vulvovaginal ulcers or ‘mucous patches’, which are round lesions with a greyish-white base edged by a dull red areola that may coalesce to produce a serpiginous ulcer - the ‘snail-track ulcer’
- lymphadenopathy characterised by firm, enlarged painless nodes typically involving inguinal, suboccipital, posterior cervical, axillary and preauricular groups

Latent syphilis

Positive serology in a patient without symptoms or signs of disease is referred to as latent syphilis and is the commonest presentation of syphilis in Australia today. Possibly because of the widespread use of antibiotics the infection often proceeds to the latent stage without a recognised primary or secondary stage.

Late syphilis

Tertiary manifestation of syphilis, which is very rare, may be ‘benign’ with development of gummas (granulomatous lesions) in almost any organ, or more serious with cardiovascular or CNS involvement. Benign gummatous disease is rare but cardiovascular disease and neurosyphilis occasionally occur. Careful management and follow-up of patients with early or latent disease is essential to prevent late sequelae.

Late syphilis should be excluded in any patient with aortic incompetence or dilatation of the ascending arch of the aorta. Syphilis should be excluded as the cause of dementia, personality change, multifocal neurological disorders, nerve deafness, pupillary abnormalities, retinal disease or uveitis.

Think of syphilis

Syphilis should not be overlooked as a cause of oral or anorectal lesions. The diagnosis of syphilis depends on a detailed history, careful clinical examination and specific examinations.

Underlying these approaches is the need to think of the possibility of syphilis with concurrent STIs.

Syphilis and HIV infection[2]

HIV and syphilis are commonly associated. In patients with AIDS and syphilis, standard regimens for syphilis are not always curative. Seronegative syphilis has been reported in patients with HIV infection. Lymphadenopathy in a patient with HIV infection may be due to coexisting secondary syphilis.

Diagnostic tests

Dark field examination[2]

Spirochaetes can be demonstrated by microscopic examination of smears from early lesions using dark-field techniques and provide an immediate diagnosis in symptomatic syphilis. Antibiotics or antiseptics should not be used until satisfactory examination has been completed. Dark-field examination has relatively low sensitivity and is not suitable for oral lesions. The direct fluorescent antibody techniques (FTAABS) can be used on this smear.

Serology

Serological tests provide indirect evidence of infection, and the diagnosis of asymptomatic syphilis relies heavily on these tests. The two main types of tests are:

- reagin tests (VDRL and RPR) - not specific for syphilis but useful for screening
- treponemal tests (TPHA, FTA - ABS) - specific tests, with the latter being sensitive and widely used

Treatment

Refer to Chapter 110.

TUBERCULOSIS

Tuberculosis (TB), caused by *Mycobacterium tuberculosis*, still has a worldwide distribution with a very high prevalence in Asian countries where 60 - 80% of children below the age of 14 years are affected.[3] This has special implication in Australia where large numbers of Asian migrants are settling. The WHO estimates that one-third of the world's population is infected by the tubercle bacillus. It remains a deadly disease with 3 million people worldwide dying of TB every year.

TB can be a mimic of other diseases and a high level of suspicion is necessary to consider the diagnosis, especially if there are extrapulmonary manifestations. There may be no symptoms or signs, even in advanced disease.

DxT: malaise + cough + weight loss (± erythema nodosum) ° TB

Primary infection

The primary infection usually involves the lungs. Transmission is by droplet infection. Symptoms include cough, sputum production, haemoptysis, fever, Figure 30.1 Classic erythema nodosum involving the legs of a patient with pulmonary tuberculosis sweating, malaise and anorexia. The focus is usually subpleural in the upper to mid zones and is almost always accompanied by lymph node involvement.



Figure 30.1 Classic erythema nodosum involving the legs of a patient with pulmonary tuberculosis

Erythema nodosum may accompany the primary infection (Figure 30.1). Primary TB is symptomless in most cases although there may be a vague illness associated with a cough. In most people this pulmonary focus heals but leaves some surviving tubercle bacilli, even if it becomes calcified.

Progressive primary tuberculosis

If the immune response is inadequate, progressive primary TB develops, with constitutional and pulmonary symptoms. Rarely, haematogenous spread can occur to the lungs (‘miliary tuberculosis’), to the pleural space (tuberculosis pleural effusion) or to extrapulmonary sites such as the meninges and bone.

Post-primary or adult-type pulmonary TB

Most cases of TB in adults are due to reactivation of disease some years later and not to reinfection. The factors causing this include poor social living conditions with malnutrition, diabetes and other factors lowering natural immunity, such as immunosuppressant drugs, corticosteroids, lymphoma and HIV infection (later stage).

Reactivated pulmonary TB

This usually presents with constitutional symptoms of poor health and night sweats, and a cough that is initially dry but may become productive and be bloodstained (see p. 456). Sometimes the infection will be asymptomatic.

Extrapulmonary TB

The main sites of extrapulmonary disease (in order of frequency in Australia) are the lymph nodes (the commonest, especially in young adults and children), genitourinary tract (kidney, epididymis, Fallopian tubes), pleura and pericardium, the skeletal system (arthritis and osteomyelitis with cold abscess formation), CNS (meningitis and tuberculomas), the eye (choroiditis, iridocyclitis), the skin (lupus vulgaris), the adrenal glands (Addison's disease (p. 224)) and the GIT (ileocaecal area and peritoneum).

These sites are illustrated in Figure 30.2 (on p. 266).


meningitis
choroiditis
iridocyclitis
primary pulmonary TB with hilar lymph nodes
lymphadenopathy
miliary pulmonary TB
adult pulmonary TB (reactivation)
constrictive pericarditis
other pulmonary problems
e.g. • collapse
• effusion
• pneumonia
lupus vulgaris (skin)
adrenal destruction (Addison's disease)
kidney
peritoneum → ascites
gastrointestinal tract
tubal disease
arthritis and osteomyelitis
epididymitis
erythema nodosum

Figure 30.2 Pulmonary and extrapulmonary distribution of tuberculosis: the primary infection starts in the lung and then spread can occur throughout the body, especially to the lymph nodes

Miliary tuberculosis

This disorder follows diffuse dissemination of tubercle bacilli via the bloodstream and is fatal without treatment. It can occur within 3 years of the primary infection or much later because of reactivation.

Diagnosis of TB

A high index of suspicion is critical for the diagnosis of TB. Tests include:

- Mantoux test (a guide only)
- chest X-ray; CT scan if doubtful
- sputum for stain (acid-fast bacilli)
- sputum for culture (takes about 6 weeks but important)

- immunochromatographic fingerprick test (new and promising)
- biopsies on lesions/lymph nodes may be necessary
- fibre-optic bronchoscopy to obtain sputum may be necessary
- consider HIV studies

Tuberculin (Mantoux) testing and BCG vaccination

A tuberculin (Mantoux) test should be performed prior to BCG vaccination in all individuals over 6 months of age. (It is read at 48 - 72 hours.)

If area of induration:

- < 5 mm - negative (*Note*: may be negative in presence of very active pulmonary infection)
- 5 - 10 mm - typical of past BCG vaccination
- > 5 mm - significant in immunocompromised, close contacts and HIV infection
- > 10 mm - positive ° tuberculosis infection (active or inactive)
- > 15 mm - highly significant for ‘normal’ people

The BCG vaccination should be given if the reaction is < 5 mm induration. Do not give it for a reaction > 5 mm.

BCG vaccination is recommended for:

- Indigenous neonates in regions of high incidence
- neonates born to patients with leprosy
- children < 5 years travelling for long periods to countries of high TB prevalence

BCG vaccination should be considered for:

- neonates in household with immigrants or visitors recently arrived from countries of high prevalance (e.g. South-East Asia) (*Note*: tuberculin test not necessary for neonates < 14 days)
- children and adolescents < 16 years with continued exposure to active TB patient and where isonizid therapy is contraindicated
- others at increased risk (and where value of BCG vaccine uncertain), such as health care workers, travellers > 5 years with significant exposure

BCG vaccination is contraindicated for:

- tuberculin reactions > 5 mm
- immunocompromised or malignancies involving bone marrow lymphatics
- high-risk HIV infection
- significant fever or intercurrent illness
- generalised skin diseases, including keloid tendency
- pregnancy
- previous infection

Treatment

The current antimicrobial treatment is to use 4 antituberculous drugs initially (rifampicin + isoniazid + pyrazinamide + ethambutol) daily for 2 months, then rifampicin + isoniazid daily for 4 months. The usual precautions with adverse reactions are required. Pyridoxine 25 mg daily is recommended for adults taking isoniazid.

Referral to specialist care is appropriate.

INFECTIVE ENDOCARDITIS

Infective endocarditis can be a difficult problem to diagnose but must be considered in the differential diagnosis of fever, especially in patients with a history of cardiac valvular disorders. It is caused by microbial infection of the cardiac valves or endocardium. Previously referred to as bacterial endocarditis, the term *infective endocarditis* is preferred because not all the infecting organisms are bacteria.

It may present as a fulminating or acute infection but more commonly runs an insidious course and is referred to as subacute (bacterial) endocarditis. Its incidence is increasing, probably due to the increasing number of elderly people with degenerative valve disease, more invasive procedures, IV drug use and increased cardiac catheterisation.[4]

DxT: FUO + cardiac murmur + embolism ° endocarditis

Predisposing factors

- Past history of endocarditis
- Rheumatically abnormal valves
- Congenitally abnormal valves
- Mitral valve prolapse
- Calcified aortic valve
- Congenital cardiac defects (e.g. VSD, PDA)
- Prosthetic valves
- IV drug use
- Central venous catheters
- Temporary pacemaker electrode catheters

Note: Only about 50% of patients with infective endocarditis have previously known heart disease.[4]

Responsible organisms

- *Streptococcus viridans* (50% of cases)
- *Streptococcus bovis*
- *Enterococcus faecalis*
- *Staphylococcus aureus* (causes 50% of acute form)
- *Candida albicans/Aspergillus* (IV drug users)
- *Staphylococcus epidermidis*
- *Coxiella burnetti* (Q fever)

Presentations

- Acute endocarditis
- Subacute endocarditis
- Prosthetic endocarditis

Infective endocarditis without cardiac murmur is frequently seen in IV drug users who develop infection on the tricuspid valve.

Warning signs for development of endocarditis

- Change in character of heart murmur
- Development of a new murmur
- Unexplained fever and cardiac murmur ° infective endocarditis (until proved otherwise)
- A febrile illness developing after instrumentation (e.g. urethral dilatation) or minor and major surgical procedures (e.g. dental extraction, tonsillectomy, abortion)
- The 'classic tetrad' of clinical features[4]
  - signs of infection
  - signs of heart disease
  - signs of embolism
  - immunological phenomena

There is a significant high mortality and morbidity from infective endocarditis, which is often related to a delay in diagnosis.

A golden rule

Culture the blood of every patient who has a fever and a heart murmur.

Clinical features

The classic clinical features are summarised in Figure 30.3.


General symptoms
• intermittent fever*
• night sweats*
• malaise*
• myalgia*
• weight loss
? pallor of anaemia
neurological problems
cerebral emboli
retinal haemorrhages
(Roth's spots)
mucosal petechiae
cardiac murmurs
cardiac failure
thoracic
back pain
(pulmonary
infarction)
splenomegaly*
(slight)
petechiae*
abdominal
pain
Hand signs
Janeway's lesions
Osler's nodes
splinter haemorrhages
? clubbing
Urinalysis
blood (microscopic)*
proteinuria
arthritis
* common features

Figure 30.3 Infective endocarditis: possible clinical features

The patients are often elderly, appear pale and ill, with intermittent fever, and complain of vague aches and pains. The full clinical presentation takes time to develop. A febrile illness of 1 to 2 weeks duration is a common presentation.

Investigation

This includes:

- FBE and ESR: ESR ↑, anaemia and leucocytosis
- urine: proteinuria and microscopic haematuria
- blood culture: positive in about 75% [4]
    - at least 3 sets of samples (aerobic and anaerobic culture)
- echocardiography - to visualise vegetations
- chest X-ray
- ECG

Consider kidney function tests and C-reactive protein.

Management

The patient should be referred because optimal management requires close co-operation between physician, microbiologist and cardiac surgeon.

Any underlying infection should be treated (e.g. drainage of dental abscess). Bactericidal antibiotics are chosen on the basis of the results of the blood culture and antibiotic sensitivities. Four blood cultures should be sent to the laboratory within the first hour of admission and treatment should seldom be delayed longer than 24 hours.

Antimicrobial treatment[5, 6]

Consultation with an infectious disease physician or clinical microbiologist should be sought.

Empirical treatment

Once cultures have been taken, prompt empirical antimicrobial treatment should be commenced, especially in fulminating infection suspected to be endocarditis (usually due to *Staphylococcus aureus*). Benzylpenicillin, gentamicin and flucloxacillin/dicloxacillin are recommended, for example (adults):

benzylpenicillin 1.8 g IV, 4 hourly plus flucloxacillin/dicloxacillin 2 g IV, 4 hourly plus gentamicin 4 - 6 mg/kg IV, daily

For specific organisms isolated on culture and prosthetic valve endocarditis confer with a consultant.

Prevention of endocarditis

Antibiotic prophylaxis[6, 7]

The evidence for prophylaxis of endocarditis is not clear, and current international practice is not to treat low-risk cardiac abnormalities having procedures with a low incidence of bacteraemia.

It is advisable to discuss the relative risks with patients.

If in doubt, discuss the issue with the patient's cardiologist or an infectious disease specialist.[7]

Low risk patients

These include patients with murmurs not due to valve disorders, isolated secundum ASD, pacemakers, implanted defribrillators, previous rheumatic fever without valve dysfunction, previous CABGS, mitral valve prolapse without regurgitation, complete surgical or device closures of congenital heart defects.

No prophylaxis is recommended in these patients.

Moderate to high risk patients

These include past history of endocarditis, prosthetic heart valves, most congenital heart disease especially cyanotic types, hypertrophic cardiomyopathy, all acquired valvular heart disease, mitral valve prolapse with regurgitation and surgically fashioned systemicpulmonary shunts.

Procedures requiring prophylaxis

- Dental: invasive dental surgery - any procedure causing bleeding from gingiva, bone or mucosa, e.g. dental extractions, drainage of abscess, osteotomies, dental implants, periodontal procedures including probing, endodontic surgery, intraligamentary local anaesthetic infections
- Other procedures - examples are:
  - genitourinary procedures in the presence of infection, e.g. D&C, IUCD, urethral dilatation, circumcision, prostatic surgery, vaginal delivery in presence of infection or prolonged labour
  - gastrointestinal tract in presence of infection
  - respiratory tract procedures - tonsillectomy/adenoidectomy, rigid bronchoscopy
  - incision and drainage of local abscess, e.g. boils, perirectal, dacrocystitis

Recommended antibiotics

Dental procedures and URT interventions:

- amoxycillin 2 g (50 mg/kg up to adult dose) orally, 1 hour beforehand (if not on long-term penicillin), *or*
- (amoxy) ampicillin 2 g (50 mg/kg up to adult dose) IV just before procedure commences or IM 30 minutes before if having a general anaesthetic, *or*
- if hypersensitive to penicillin: clindamycin or vancomycin

Genitourinary and gastrointestinal procedures

- gentamicin (child: 2.5 mg/kg) 2 mg/kg IV (just before procedure) or IM (30 minutes beforehand) plus
- amoxy/ampicillin (child: 50 mg/kg up to) 2 g IV (just before procedure) followed by (child 5 mg/kg up to) 1 g IV, IM or orally, 6 hours later

If hypersensitive to penicillin: vancomycin or teicoplanin plus gentamicin.

ZOONOSES

Zoonoses are those diseases and infections that are naturally transmitted between vertebrate animals and humans (see Table 30.1). Zoonotic diseases (which are not restricted to farming communities) can present as a mild illness but are prolonged in duration and can have debilitating sequelae.8 There is a long list of diseases, which vary from country to country, and includes plague, rabies, scrub typhus, Lyme disease, tularaemia, hydatid disease, orf, anthrax, erysipeloid, listeriosis, campylobacteriosis and ornithosis (psittacosis).

Table 30.1 Major zoonoses in Australia

Zoonosis	Organism/s	Animal host	Mode of transmission	Main presenting features
Q fever	*Coxiella burnetti*	Various wild and domestic animals	• Inhaled dust • Animal contact • Unpasteurised milk	Fever, rigors, myalgia, headache, dry cough
Leptospirosis	*Leptospira pomona*	Various domestic animals	• Infected urine contaminating cuts or sores	Fever, myalgia, severe headache, macular rash
Brucellosis	*Brucella abortus*	Cattle	• Contamination of cuts or sores by nimal tissues • Unpasteurised milk	Fever (undulant), sweats, myalgia, headache, lymphadenopathy
Lyme disease	*Borrelia burgdorferi*	Marsupials (probable)	• Tick bites	Fever, myalgia, arthritis, backache, doughnutshaped rash
Psittacosis	*Chlamydia psittaci*	Birds: parrots, pigeons, ducks, etc.	• Inhaled dust	Fever, myalgia, headache, dry cough
Bovine tuberculosis	*Mycobacterium bovis*	Cattle	• Unpasteurised milk	Fever, sweats, weight loss, cough (as for human pulmonary TB)
Listeriosis	*Listeria monocytogenes*	Various wild and domestic animals	• Unpasteurised milk or cheese • Contaminated vegetables • Person to person	Mild febrile illness (in most) Meningoencephalitis in those susceptible (neonates, pregnancy, elderly etc.)

Diagnosis of zoonoses[9]

If a zoonosis is overlooked in the differential diag nosis, many will remain undiagnosed and untreated.

Fever and sweats (influenza-like illness)

Any patient with undiagnosed fever should be questioned about exposure to animals, recent travel both in and out of Australia, animal bites, cat scratches, consumption of raw milk, mosquito and tick bites, pets and occupation.

Rash

Consider rickettsial illness such as leptospirosis, Q fever, Lyme disease.

Cough or atypical pneumonia

Consider Q fever, psittacosis, bovine TB.

Arthralgia/arthritis

Consider Lyme disease, Ross River fever.

Meat workers

Consider Q fever, leptospirosis, orf, anthrax.

Papular/pustular lesions

Consider orf, anthrax (black).

BRUCELLOSIS

Brucellosis (undulant fever, Malta fever) has diminished in prevalence since the campaign to eradicate it from cattle. Entry is mainly by the mouth, or abraded or cut skin.

Features of acute brucellosis

- Incubation period 1 - 3 weeks
- Insidious onset: malaise, headache, weakness
- The classic fever pattern is undulant (refer to p. 578).

Possible:

- arthralgia
- lymphadenopathy
- hepatomegaly
- spinal tenderness
- splenomegaly (if severe)

Complications such as epididymo-orchitis, osteomyelitis and endocarditis can occur. Localised infections in sites such as bones, joints, lungs, CSF, testes and cardiac valves are possible but uncommon.

Symptoms of chronic brucellosis are virtually indistinguishable from the ‘chronic fatigue syndrome’ and can present with FUO.

DxT: malaise + headache + undulant fever ° brucellosis

Diagnosis

- Blood cultures if febrile (positive in 50% during acute phase)9
- *Brucella* agglutination test (rising titre) - acute and convalescent (3 - 4 weeks) samples

Treatment[10]

- Adults: doxycycline 100 mg (o) bd for 6 weeks plus either rifampicin 600 mg (o) daily for 6 weeks *or* gentamicin 4 - 6 mg/kg/day IV daily for 2 weeks
- Children: cotrimoxazole + rifampicin
- Relapses do occur.

Prevention and control

Involves eradication of brucellosis in cattle, care handling infected animals and pasteurisation of milk. No vaccine is currently available for use in humans.

Q FEVER

Q fever is a zoonosis due to *Coxiella burnetti*. It is the most common abattoir-associated infection in Australia and can also occur in farmers and hunters. Rash is not a major feature but can occur if the infection persists without treatment.

Clinical features

- Incubation period 1 - 3 weeks
- Sudden onset fever, rigors and myalgia
- Dry cough (may be pneumonia in 20%)10
- Petechial rash (if persisting infection)
- > ± Abdominal pain

Persistent infection may cause pneumonia or endocarditis so patients with valvular disease are at risk of endocarditis. It is a rare cause of hepatitis. The acute illness may resolve spontaneously. Untreated chronic infection is usually fatal.

DxT: fever + headache + prostration ° Q fever

Diagnosis

- Serodiagnosis is by antibody levels in acute phase and 2 - 3 weeks later.

Treatment[10]

- Doxycycline 100 mg (o) bd for 14 days
- For endocarditis: prolonged course of doxycycline plus clindamycin or rifampicin
- Children: > 8 same antibiotics according to weight

< 8 cotrimoxazole (instead of doxycycline)

Prevention

The disease can be prevented in abattoir workers by using Q fever vaccine.

LEPTOSPIROSIS

Leptospirosis follows contamination of abraded or cut skin or mucous membranes with leptospira-infected urine of many animals including pigs, cattle, horses, rats and dogs. In Australia it is almost exclusively an occupational infection9 of farmers and workers in the meat industry. There is a risk to dairy farmers splashed with urine during milking. Early diagnosis is important to prevent it passing into the immune phase.

Clinical features

- Incubation period 3 - 20 days (average 10)
- Fever, chills, myalgia
- Severe headache
- Macular rash
- Light-sensitive conjunctivitis (marked suffusion)

Some may develop the immune phase (after an asymptomatic period of 1 - 3 days) with aseptic meningitis or jaundice and nephritis (icterohaemorrhagic fever, Weil's syndrome) with a significant mortality.

DxT: abrupt fever + headache + conjunctivitis ° leptospirosis

Diagnosis

- High or rising titre of antibodies: can be cultured

Treatment[10]

- Doxycycline 100 mg (o) bd for 5 - 7 days *or*
- benzylpenicillin 1200 mg IV, 6 hourly for 5 - 7 days

LYME DISEASE

Lyme disease (known as *Lyme borreliosis*) was first described in 1975 and named after the town Lyme in Connecticut (US). It is widespread in the United States and is now appearing in Europe, Asia and Australia. Very infective, it is caused by a spirochaete, *Borrelia burgdorferi*, and transmitted by *Ixodes* ticks, so that people living and working in the bush are susceptible. It has been reported in deer farmers. Lyme disease presents in three phases.

Stages of Lyme disease

Stage 1

- Erythema migrans - a characteristic pathognomonic rash, usually a doughnutshaped, well-defined rash about 6 cm in diameter at the bite site.
- > ± Influenza-like illness: malaise, fever, myalgia, arthralgia, backache.

Stage 2

(weeks or months later)

- Neurological problems, such as limb weakness, muscle pain, aseptic meningitis, painful radiculopathies, cranial nerve palsies (e.g. Bell's palsy).
- Cardiac problems: arrhythmias, myocarditis.

Stage 3

- Arthritis (weeks or months later). In children it can be mistaken for juvenile chronic arthritis. In adults, it usually affects one or two joints, especially the knee joints. A LIMP syndrome (localised intermittent musculoskeletal pain) is described, which is severe pain without significant findings on examination.

Diagnosis

- Clinical pattern especially rash of erythema migrans. Serology to demonstrate Lyme antibodies, which take about 6 weeks to develop from the onset of the disease.

Treatment

- A typical regimen for adults is doxycycline 100 mg bd for 21 days or penicillin (the earlier the diagnosis the more effective the treatment).

PSITTACOSIS

Most patients are bird fanciers. Psittacosis accounts for 1 - 5% of hospital admissions for pneumonia. The disease may follow a low-grade course over several months but can have a dramatically acute presentation.

Clinical features

- Incubation period 1 - 2 weeks
- Fever, malaise, myalgia
- Headache
- Cough
- Minimal chest signs
- Splenomegaly (sometimes)

Mortality can be as high as 20% if untreated.

Diagnosis

- Serology - rising antibody

Treatment

- Tetracycline or erythromycin for 14 days

LISTERIOSIS[11]

Listeriosis is caused by *Listeria monocytogenes*, a bacterium widespread in nature that can contaminate food and has been found in many fresh and processed foods (e.g. dairy products, especially unpasteurised milk, soft cheese, processed meats and smoked seafood). Its significance lies in the mortality rate in high-risk groups such as pregnant women, the immunocompromised, frailaged, and very young but especially neonates and foetuses. Babies may be stillborn or aborted.

Clinical features

It may be subclinical but possible presentations include:

- influenza-like illness (usually mild)
- food poisoning (atypical)
- meningitis, especially infants, elderly
- septicaemia (in susceptible)
- pneumonia (in susceptible)

Diagnosis

- Microscopy or isolation of organism from infected site
- Serological tests available

Treatment

- Amoxycillin 1 g (o) 8 hourly for 10 - 14 days11

OTHER ZOONOSES

- Mosquito-transmitted infections - Murray Valley encephalitis, Ross River virus, Barmah Forest virus
- Infections from bites and scratches - cat-scratch disorder, rat bite fever
- Hydatid disease, orf, milker's nodules
- Toxoplasmosis, histoplasmosis, hookworm

CLOSTRIDIAL INFECTIONS

Tetanus

This sometimes misdiagnosed bacterial infection (*Clostridium tetani*) can appear from one day to several months after the injury, which can be forgotten. A total of 10 - 20% of patients with tetanus have no identifiable wound of entry.[12]

Clinical features

- Prodrome: fever, malaise, headache
- Trismus (patient cannot close mouth)
- Risus sardonicus (a grin-like effect from hypertonic facial muscles)
- Opisthotonos (arched trunk with hyperextended neck)
- Spasms, precipitated by minimal stimuli

Differential diagnosis: phenothiazine toxicity, strychnine poisoning, rabies

Management

- Give tetanus antitoxin
- Refer immediately to expert centre
- Intubate and ventilate if necessary

Gas gangrene

Gas gangrene (clostridial myonecrosis) is caused by entry of one of several clostridia organisms e.g. *C. perfringes* into devitalised tissue, such as exists following severe trauma to a leg.

Clinical features

- Sudden onset of pain and swelling in the contaminated wound
- Brownish serous exudate
- Gas in the tissue on palpation or X-ray
- Prostration and systemic toxicity
- Circulatory failure (‘shock’)

Management

- Refer immediately to surgical centre
- Start benzylpenicillin 2.4 g IV, 4 hourly

Botulism

Botulism is food poisoning caused by the neurotoxin of *Clostridium botulinum*. From 12 to 36 hours after ingesting the toxin from canned, smoked or vacuumpacked food (e.g. homecanned vegetables or meat) visual problems such as diplopia suddenly appear. General muscle paralysis and prostration quickly develop. Refer immediately for antitoxin.

PNEUMONIA

Surprisingly the initial presentation of pneumonia can be misleading, especially when the patient presents with constitutional symptoms such as fever, malaise and headache rather than respiratory symptoms. A cough, although usually present, can be relatively insignificant in the total clinical picture. Th is problem applies particularly to atypical pneumonia but can occur with bacterial pneumonia, especially lobar pneumonia (refer to pp. 451 - 2).

The atypical pneumonias

Clinical features

- Fever, malaise
- Headache
- Minimal respiratory symptoms, non-productive cough
- Signs of consolidation absent
- Chest X-ray (diffuse infiltration) incompatible with chest signs

> **DxT:** ‘flu’ + headache + dry cough ° atypical pneumonia

Serology tests

Blood tests are available for all the following causative organisms (refer to Chapter 16):

- *Mycoplasma pneumoniae* - the commonest:
  - adolescents and young adults: treat with roxithromycin (first line) 300 mg (o) daily *or* tetracycline (e.g. doxycycline 100 mg bd for 14 days)
- *Legionella pneumophila* (legionnaire's disease):
  - related to cooling systems in large buildings
  - incubation 2 - 10 days

  Diagnostic criteria include:
  - prodromal-like illness
  - a dry cough, influenza, confusion or diarrhoea
  - lymphopenia with marked leucocytosis
  - hyponatraemia

  Patients can become very prostrate with complications - treat with azithromycin (IV) or erythromycin (IV or o) plus (if very severe) ciprofloxacin or rifampicin for 21 days.
- *Chlamydia psittaci* (psittacosis)
  - treat with doxycycline 100 mg bd for 10 days

- *Coxiella burnetti* (Q fever)

HANSEN'S DISEASE (LEPROSY)

Hansen's disease is caused by the acid-fast bacillus *Mycobacterium leprae*. It is a disorder of tropical and warm temperate regions, especially South-East Asia. It is considered to be transmitted by nasal secretions with an incubation period of 2 - 6 years.

Clinical features

(WHO 1999) [13]

Diagnosis is one or more of:

- skin lesions - usually anaesthetic; hypopigmented or reddish maculopapules or annular lesions
- thickened peripheral nerves with loss of sensation, e.g. ulnar (elbow), median (wrist), common peroneal (knee) and greater auricular (neck)
- demonstration of acid-fast bacilli in a skin smear or on biopsy

It can be localised (tuberculoid) or generalised (lepromatous).

Diagnosis is by biopsy, the lepromin test, cultivation of the organisms or by PCR tests.

Referral to specialists or a specialist centre is advisable.

WHO treatment recommendations are multiple drug therapy, e.g. rifampicin, clofazimine and dapsone, but therapy is constantly being evaluated.

ACKNOWLEDGMENT

Part of this text, on the clinical manifestations of syphilis, is reproduced from the *Handbook on Sexually Transmitted Diseases.* [2] Commonwealth of Australia, copyright, reproduced by permission.

REFERENCES

1. Hart G. Syphilis. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 493 - 6.
2. NH&MRC. *Handbook on Sexually Transmitted Diseases*. Canberra: Department of Community Services & Health, 1990: 23 - 9.
3. Kumar PJ, Clark ML. *Clinical Medicine* (5th edn). London: Bailliere Tindall, 2003: 126 - 7.
4. Oakley C. Infective endocarditis. Med Int, 1986; 21: 872 - 8.
5. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 46 - 52.
6. Speed B. Endocarditis, infective. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing 1996: 167 - 9.
7. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 174 - 9.
8. Scott J. Zoonoses. Curr Ther, 1995: April: 42 - 5.
9. Benn R. Australian zoonoses. Curr Ther, 1990: July: 31 - 40.
10. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 299 - 305.
11. NH&MRC Statement. Listeria, advice to medical practitioners. Canberra: NH&MRC, 1992.
12. Braunwald E et al. *Harrison's principles of internal medicine* (15th edn). New York: 2001: 918 - 20.
13. World Health Organisation leprosy website < http://www.who.int/lep/ >

CHAPTER 31. CHRONIC KIDNEY DISEASE

I have never yet examined the body of a patient dying with dropsy attended by albuminous urine,
in whom some obvious derangement was not discovered in the kidneys.

RICHARD BRIGHT 1827

CHRONIC KIDNEY DISEASE

Chronic kidney disease (CKD) is defined as a glomerular filtration rate (GFR) less than 60 ml/min/1.73 m2 and/or evidence of kidney damage for a period of at least 3 months (Table 31.1).

Table 31.1 Definition of chronic kidney disease[2]

- GFR ˂60 mL/min/1.73 m^2 for ≥ 3 months with or without evidence of kidney damage *or*
- Evidence of kidney damage (with or without decreased GFR) for ≥ 3 months, as evidenced by any of the following:
  - microalbuminuria (urinary albumin excretion rate 30 - 300 mg/day)
  - macroalbuminuria (urinary albumin excretion rate < 300 mg/day)
  - persistent haematuria (where other causes such as urologic conditions have been excluded)
  - pathologic abnormalities (e.g. abnormal kidney biopsy)
  - radiologic abnormalities (e.g. scarring or polycystic kidneys on kidney ultrasound scan)

Chronic kidney or kidney failure (CKF) is defined as a severe reduction in nephron mass over an extended period of time, resulting in uraemia. It can present surreptitiously and be a real master of disguise in clinical practice. Asymptomatic CKF may be discovered on routine health screening, as a chance finding in hospitalised or hypertensive patients, or during follow-up of patients with known kidney disease.[1]

ACUTE KIDNEY FAILURE

Acute kidney failure (AKF, a forerunner of CKF) is defined as a sudden (days to weeks) decrease in kidney function (azotaemia) with or without oliguria. It results in dysfunctional fluid and electrolyte balance and nitrogenous waste excretion with a sudden increase in blood urea and creatinine levels.

AKF is usually classified into:

- prerenal (e.g. acute circulatory failure → kidney hypoperfusion)
- postrenal (e.g. obstruction)
- kidney (intrinsic) (e.g. acute glomerulonephritis)

Early diagnosis with hospital admission is important and this is achieved by being aware of the patient at risk and the early detection of hypovolaemia, hypertension or hypotension, oliguria or urine abnormalities.

DxT: malaise (extreme) + a/n/v + confusion (±oliguria) ° AKF

KEY FACTS AND CHECKPOINTS

- At least 95 people per million of the population are treated for end-stage kidney/kidney failure (ESKF) each year.
- Two-thirds of these are under 60 years of age.
- Important causes are glomerulonephritis, analgesic nephropathy, diabetes mellitus (30%) polycystic kidney disease, reflux nephropathy and hypertension (Table 31.2 on p. 276).[3]
- The commonest cause of ESKF in Australia is diabetes mellitus.
- The commonest cause of nephritis leading to kidney failure in Australia is IgA nephropathy.
- In children the incidence of chronic kidney failure is quite low (1 to 2 per million of the population).[3]
- Warmer climates, poorer living conditions and certain genetic predispositions are associated with a higher prevalence of kidney failure.

- Kidney failure should be considered in the diagnosis of patients with:
    - unexplained anaemia
    - unexplained poor health
    - unusually high analgesic intake[3]
- Uraemic symptoms are non-specific and usually are not recognised until the creatinine clearance is less than 20% of normal.
- CKF is characterised by the accumulation of uraemic toxins and a deficiency of kidney hormones that cause dysfunction of organs other than kidneys.
- This interaction can cause phosphate retention, secondary hyperparathyroidism and bone disorders such as osteomalacia.
- It is possible to identify stages of kidney failure (Table 31.3).

Table 31.2 Significant causes of chronic kidney failure (approximate order of prevalence)

Diabetes mellitus
Glomerulonephritis
• IgA nephropathy (commonest)
Hypertension
Vascular
• atherosclerosis, including kidney artery stenosis
Polycystic kidneys
Obstructive nephropathy/reflux
• bilateralureteric obstruction
• bladder outflow obstruction
- prostatic enlargement
- urethral stenosis
Drugs, including analgesic nephropathy
Lupus and other connective tissue disorders
Vasculidities, e.g. PAN
Gout
Amyloidosis
Hypercalcaemia

Table 31.3 Classification of CKD stages[2]

CKD stage	GFR (mL/min)	Description	Clinical action plan
1	>90	Evidence of kidney damage, e.g. scarring on ultrasound, proteinuria/haematuria	
2	60 - 89	Evidence of kidney damage Mild kidney failure	Further investigation for those at risk: • Assess proteinuria • Urinalysis • BP Cardiovascular risk reduction: • BP, cholesterol, blood glucose, smoking, obesity
3	30 - 59	Moderate kidney failure	As above. • Avoid nephrotoxic drugs. • Monitor eGFR 3 monthly. • Prescribe antiproteinuric drugs, ACE inhibitors or AIIRAs if appropriate. • Address anaemia, acidosis, hyperparathyroidism.

			• Ensure drug dosages are appropriate for level of kidney function. Consider referral to nephrologist.
4	15 - 29	Severe kidney failure	As above. • Referral to nephrologist. • Prepare for dialysis or transplantation (if appropriate).
5	<15	End-stage kidney failure	As above. • Institute dialysis or transplantation (if appropriate).

IMPORTANT CLINICAL ASSOCIATIONS

The possibility of CKF should be monitored in patients with:

- diabetes mellitus
- hypertension
- severe gout
- a history of urinary tract abnormality (e.g. vesico-ureteric reflux, bladder outflow obstruction)

The possibility of CKF should be considered and investigated in patients presenting with:

- unexplained poor health
- hypertension
- anaemia
- pruritus
- hyperparathyroidism
- pericarditis
- urinary tract symptoms or signs
  - proteinuria
  - haematuria
  - oedema
  - nocturia
  - loin pain
  - prostatic obstruction
- neurological disturbances
  - confusion
  - coma
  - peripheral neuropathy
  - seizures

Patients with CKF may present with features of acute kidney failure with the intervention of complicating factors such as:

- drug toxicity
- infection
- fluid imbalance

Urgent treatment of the following conditions, which can lead to rapid kidney failure, is essential:

- progressive nephritis
- systemic lupus erythematosus
- vasculitides (p. 289) e.g.
  - polyarteritis nodosa
  - Wegener's granulomatosis

THE CLINICAL APPROACH

History

A hallmark of early stage CKF is a non-specific history and examination, and the diagnosis is very difficult in the absence of a known past history of kidney disease. The diagnosis can be established only by kidney function tests. Symptoms from CKF are rare unless the creatinine clearance is less than 20% of normal and only become common when less than 10% of normal.

In patients with chronic kidney disease, symptomatic uraemia may be precipitated by prerenal factors, such as fluid loss from vomiting or diarrhoea, infection, antibiotic therapy especially tetracyclines, or increasing hypertension.

Symptoms and signs

The symptoms and signs of CKF are summarised in Figure 31.1.


Anaemia
• lethargy
• pallor
Haematological
• anaemia
• bruising
• epistaxis
Respiratory
• dyspnoea
• tachypnoea
Skin
• pruritus
• scratch marks
• pigmentation
• purpura
Nail changes
• proximal leuconychia
• distal brownish discoloration
Flapping tremor
Musculoskeletal
• muscle weakness
• bone pain
• restless legs
Periphery
• peripheral neuropathy
• oedema
CNS
• weakness
• drowsiness
confusion
twitching
fits
coma
severe uraemia
hyperparathyroidism (secondary)
Cardiovascular
• heart failure
• hypertension
• pericarditis
Gastrointestinal tract
• anorexia, nausea
• vomiting
• hiccoughs
• diarrhoea
Renal
• polyuria
• nocturia
Genital
• erectile impotence
• infertility
• amenorrhoea
haematuria
proteinuria

Figure 31.1 Clinical features of chronic kidney disease

The common early presenting symptoms are generally non-specific and referable to the GIT, presumably due to the formation of ammonia in the upper GIT.

Such symptoms include:

- anorexia
- nausea
- vomiting
- tiredness
- lethargy

If a patient presents with these symptoms and has a sallow 'lemon' tinge appearance due to a combination of anaemia and brownish pigmentation, then CKF should be highly suspected.

DxT: fatigue + anv + sallow skin ° CKF

Physical examination[4]

General inspection of the patient with CKF will usually reveal a sallow complexion with yellow-brown pigmentation in the skin, which is oft en dry and pruritic. The patient's mental state should be noted. The respiratory and pulse rates are usually rapid because of anaemia and metabolic acidosis. Other findings may include bruising, uraemic fetor, reduced mental status, pericarditis and peripheral neuropathy. The abdomen should be carefully palpated, especially in the kidney areas. A rectal examination is indicated to detect prostatomegaly or other rectal or pelvic pathology. Ophthalmoscopic examination may show hypertensive or diabetic retinopathy. Urinalysis should test glucose, blood and protein. Proteinuria should be confirmed with a 24-hour urine protein estimation or (preferably) a albumin creatinine ratio (ACR).

ACR guidelines:

Normoalbuminuria

- Men: < 2.5 mg/mmol
- Women: < 3.5 mg/mmol

Microalbuminuria

- Men: 2.5 - 25 mg/mmol
- Women: 3.5 - 35 mg/mmol

Macroalbuminuria

- Men: > 25 mg/mmol
- Women: > 35 mg/mmol

Investigations

- Kidney function tests (most appropriate for the GP):
  - plasma urea
  - plasma creatinine
  - creatinine clearance (more precise)
  - eGFR (the new standard)
- Plasma electrolytes:
  - sodium, potassium, chloride, bicarbonate
  - calcium and phosphate
- Consider:
  - magnesium, urate, glucose
  - lipids
  - prescribed drug level
  - cardiac studies
  - full blood count ? anaemia
  - protein electrophoresis (? myeloma)
  - ANA for lupus
  - ANCA for vasculitis
- Determination of underlying cause:
  - imaging of urinary tract - ultrasound
  - immunological tests
  - kidney biopsy

CHRONIC KIDNEY FAILURE IN CHILDREN

The incidence of CKF in children is about two per million of the total population per year. The commonest causes include chronic glomerulonephritis, obstructive nephropathy and reflux nephropathy. Identification of structural kidney abnormalities by obstetric ultrasound and early investigation of urinary tract infections may decrease the incidence of CKF. Dialysis and transplantation are normally considered for children over 2 years of age with end-stage CKF. For children under 2 years there are complex ethical, psychological and technical problems.[5] Nevertheless the prognosis for such treatment is poor.

MONITORING CKD

The traditional test in identifying and monitoring CKD is the serum creatinine level.[5] The normal range is about 40 - 120 μmol/L (0.04 - 0.12 mmol/L) but the laboratory will indicate their appropriate reference level. However, serum creatinine is an unreliable and insensitive marker of CKD. To improve detection and management guidelines laboratories now report on estimated glomerular filtration rate (eGFR) using the Modification of Diet in Kidney Disease (MDRD) formula with every request for serum creatinine concentration, which is required to calculate to GFR.[6] The eGFR measured by the Cockroft and Gault formula more reliable than the MDRH method.

DRUG PRESCRIBING IN CKD[1, 7]

Drugs that can damage the kidneys include:

- classic nephrotoxic drugs, e.g. gentamicin, vancomycin
- NSAIDs, COX-2 inhibitors
- ACE inhibitors and AIIRA antagonists
- aminoglycosides
- cephalosporins (various)
- tetracylcines
- lithium

Beware of the 'triple whammy':

- NSAIDs/COX-2 inhibitors
- ACE inhibitors
- diuretics

These three agents individually or in combination are implicated in over 50% of cases of iatrogenic acute kidney failure.[8]

Diuretics should be used with care.

Drugs causing hyperkalaemia

- NSAIDs/COX-2
- ACE inhibitors
- AIIR antagonists (AIIRA)
- Aldactone
- Moduretic
- Trimethoprim

Increased risk of adverse reaction

- Allopurinol:
    - vasculitis
    - liver dysfunction
- Statins:
    - liver dysfunction
    - myopathy
    - rhabdomyolysis
- Gemfibrozil:
    - rhabdomyolysis

Do not use statins and gemfibrozil together.

- Beta lactams:
    - interstitial nephritis
- LMW heparin:
    - ↑bleeding
- Aspirin/NSAID:
    - GIT bleeding
- Omeprazole and related agents:
    - interstitial nephritis

Dangerous drug accumulation is presented in Table 31.4.

Table 31.4 Dangerous drug accumulation in kidney impairment[7]

Drug	Problem
Acyclovir	Confusion, encephalopathy
Co-trimoxazole	Steven Johnson syndrome
Flagyl (long term)	Peripheral neuropathy
Penicillin (high dose IV)	Seizures
Quinalones	
• ciprofloxacin	Seizures
• norfloxacin	
Metformin	Lactic acidosis
Sulfonylureas	Prolonged hypoglycaemia
Insulin	Hypoglycaemia
Atenolol	Bradycardia/heart block
Digoxin	Nausea, bradycardia
Sotalol	Ventricular tachycardia (Mg required before conversion)
Codeine	Confusion, acute brain syndrome
Methotrexate	Liver dysfunction
	Bone marrow depression
Lithium	Tremor - confusion
	Thyroid dysfunction

Management

The patient should be referred to an appropriate specialist as early as possible. The underlying disease and any abnormalities causing progressive kidney damage must be corrected where possible. The management of CKF is based on the team approach involving specialists and paramedical personnel. The patient is usually faced with years of ongoing care so that an empathic support team based around the patient's general practitioner is very important to the patient, who will require considerable psychosocial support.

Optimum treatment includes:

- regular review
- good blood pressure control (the most effective way to slow progression)
- keeping plasma phosphate levels in normal range
- maintaining effective fluid and electrolyte balance
- prompt treatment of intercurrent illness
- judicious use of drugs
- avoiding treatment errors, especially with drugs
    - avoid potassium-sparing diuretics
    - other drugs that may cause problems include digoxin, tetracyclines, gentamicin, NSAIDs, nitrofurantoin and ACE inhibitors

(ACEIs)

- rapid treatment of complications, especially salt and water depletion and acute urinary tract infection
- diet - low protein, sodium and potassium
- treating anaemia with human recombinant erythropoietin and iron (iron infusions)

Targets: goals of management[1]

The following are optimal targets for patients with chronic kidney disease:

- blood pressure < 130/85 mm Hg if proteinuria < 1 g/day ≤ 125/75 mm Hg if proteinuria > 1 g/day
- cholesterol total < 4.0 mmol/L LDL < 2.5 mmol/L
- blood sugar pre-prandial 4.4 - 6.7 mmol/L Hb Alc ≤ 7%
- haemoglobin 110 - 120 g/L
- serum potassium ≤ 6 mmol/L
- BMI 25 kg/m^2
- proteinuria ≥ 50% reduction of baseline value
- acidosis $HCO^-_3 > 22$ mmol/L
- phosphate $PO^-_4 \leq 1.75$ mmol/L
- no smoking
- alcohol ≤ 2 standard drinks/day

Blood pressure control

- No added salt diet (with care)
- Drug control: none of the antihypertensive agents are specifically contraindicated but those eliminated mainly by the kidney (e.g. ACE inhibitors, atenolol, sotalol) should be given in lower dosage. ACE inhibitors should not be used in the presence of kidney artery stenosis; loop diuretics (e.g. frusemide) are effective in larger doses.[3] The first line agents are ACEIs or AIIRAs. Th ey should be ceased if the creatinine levels exceed 30% above baseline or if the serum K exceeds 6 mmol/L (despite dose reduction).[1] The non-dihydropyridine calcium channel blockers are next choice. Beta blockers can be used. Diuretics have a vital role in the patient with diastolic heart failure (p. 1337).[7]

Hyperphosphataemia control

- Balanced nutrition to reduce dietary phosphate
- Protein restriction
- Calcium carbonate (to bind phosphate)

Dialysis

Dialysis is indicated when all other methods fail. About two-thirds of patients receive haemodialysis and about 22% are on continuous ambulatory peritoneal dialysis and automated overnight peritoneal dialysis.

Transplantation

Transplantation is the treatment of choice for kidney failure except where contraindicated, such as with active malignancy or tuberculosis and perhaps the elderly. However, a critical shortage of donors remains a problem.

WHEN TO REFER[1]

- (Glomerular) haematuria
- eGFR < 30 (stage 4 or 5 CKD)
- Rapidly declining kidney function
- Significant proteinuria > 1 g/24 hours
- Kidney impairment + hypertension (poor control)
- Diabetes with kidney impairment or albuminuria/proteinuria

REFERENCES

1. Kumar PJ, Clark M. *Clinical Medicine* (5th edn). London: Saunders, 2003: 637 - 8.

2. Johnson DW, Usherwood T. Chronic kidney disease. Aust Fam Physician, 2005; 34(11): 915 - 21.
3. Dawborn JK. Chronic kidney failure. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 442 - 3.
4. Tally N, O'Connor S. *Clinical Examinations* (5th edn). Sydney: Maclennan & Petty, 2006: 186 - 7.
5. Robinson MJ, Roberton DM. *Practical Paediatrics* (5th edn). Melbourne: Churchill Livingstone, 2003: 610 - 1.
6. Johnson DW, Usherwood T. Automated reporting of GFR: coming to a laboratory near you. Aust Fam Physician, 2005; 34(11): 925 - 8.
7. Cunninghan M. Drug therapy in the renally challenged. Monash University Update for GPs. Seminar proceedings November, 2005: 1 - 8.
8. Th omas MC. Diuretics, ACE inhibitors and NSAIDs - 'the triple whammy'. MJA, 21 Feb 2000: 172.

CHAPTER 32. CONNECTIVE TISSUE DISEASE AND THE VASCULITIDES

In its more aggravated forms diffuse scleroderma is one of the most terrible of all human ills. Like Tithonus to 'wither slowly' and like him to be 'beaten down and marred and wasted', until one is literally a mummy, encased in an ever shrinking, slow contracting skin of steel, is a fate not pictured in any tragedy.

SIR WILLIAM OSLER 1898

CONNECTIVE TISSUE DISEASE AND THE VASCULITIDES

The connective tissue diseases and the related vasculitides are groups of disorders that are difficult to classify because their causation is generally unknown. They all cause joint and soft tissue inflammation and multiple other possible manifestations that create diagnostic difficulties.

AUTOIMMUNE DISEASES[1]

These are disorders in which the body's immune system damages specific organs or systems. The connective tissue diseases are a classic subgroup of autoimmune disease. Rheumatoid arthritis is the most common autoimmune disease. Organ-specific autoimmune diseases include diabetes type 1, Hashimoto's thyroiditis, pernicious anaemia, IgA glomerulonephritis, Graves' disease, autoimmune hepatitis and myasthenia gravis.

It is convenient to consider a working classification of joint pain (Table 32.1) that includes apparent joint pain (arthralgia), as some of the inflammatory disorders cause problems in the soft tissues around joints (e.g. giant cellarteritis and hydroxyapatite crystalo pathy of the tendons around the shoulder joint).

Table 32.1 A classification of rheumatological pain

Hyperacute (red hot) joints	Crystals	• Urate: gout • Calcium pyrophosphate • Hydroxyapatite
	Pus	e.g. staphylococcal septic arthritis
Inflammation of joints	Symmetrical	e.g. rheumatoid arthritis
	Asymmetrical	e.g. spondyloarthropathies
Non-inflammatory joint disorder	Typical	Primary (e.g. in hands)
• osteoarthritis	Atypical	e.g. post-trauma; haemochromatosis
Joint and soft tissue inflammation	Connective tissue disorders	• SLE • Systemic sclerosis • Polymyositis/dermatomyositis
	Vasculitides	• Polyarteritis nodosa • Giant cell arteritis • Polymyalgia rheumatica
Non-articular (soft tissue) inflammation	Generalised	e.g. fibrositis, fibromyalgia, polymyalgin
	Localised	e.g. plantar fasciitis; epicondylitis

Source: After Dr Stephen Hall

Vasculitis is, in fact, a condition common to the connective tissue disorders and to the so-called vasculitides (Table 32.2).

Table 32.2 List of connective tissue disorders and vasculitides

Connective tissue disorders
• SLE
• Scleroderma Limited Scleroderma
• Polymyositis/Dermatomyositis
• Sjögren's syndrome

• Raynaud's phenomenon
Vasculitides Large vessel predominantly • Giant cell arteritis/Temporal arteritis/polymyalgia rheumatica • Takayasu's arteritis • Behçet's syndrome Medium vessel • Polyarteritis nodosa • Kawasaki's disease Small vessel (mainly) • Henoch-Schönlein purpura • Hypersensitivity vasculitis • Essential cryoglobulinaemia ANCA associated • Wegener's granulomatosis • Churg-strauss vasculitis • Microscopic polyangiitis

A major concern to all is that the diagnosis of these conditions is elusive and often delayed.

CONNECTIVE TISSUE DISEASE

The term ‘connective tissue disease’ (CTD) is a generic label applied to a group of disorders characterised by inflammation, presumed initiated by an auto immune response to an autoantigen and perpetuated by unknown factors.[2] The vasculitides are a variety of CTD.

The CTDs comprise three distinct conditions, namely systemic lupus erythematosus (SLE), scleroderma and polymyositis/dermatomyositis (Figure 32.1).[1]


systemic lupus erythematosus
overlap syndrome (mixed connective tissue disorder)
polymyositis/ dermatomyositis
scleroderma

Figure 32.1 The connective tissue disorders

Mixed connective tissue disorder includes features of all three disorders and is sometimes referred to as 'overlap' syndrome. Other related disorders classified as CTDs include Sjögren's syndrome and Raynaud's syndrome.

Common features include:

- fatigue
- arthralgia or arthritis
- multisystem involvement
- vasculitis
- immunological abnormalities
- sicca (dry skin and mucous membranes)

INVESTIGATIONS IN CONNECTIVE TISSUE DISEASE

The rapidly growing list of investigations, particularly that of autoantibodies, can be confusing. Baseline investigations to consider are FBE, ESR, C-reactive protein and rheumatoid factor. The rheumatoid factor can be positive in a great range of disorders, including rheumatoid arthritis, SLE, systemic sclerosis, Sjögren's syndrome, chronic liver disease and various viral (e.g. hepatitis), bacterial (e.g. tuberculosis) and parasitic (e.g. malaria) infections. There is usually a low titre, except in rheumatoid arthritis. X-rays and HLA-B_{27} tests are not recommended.

Table 32.3 summarises the majority of autoantibodies that can be tested. The antinuclear antibody (ANA) test is very sensitive for SLE, but not absolutely specific (false positives occur with viral arthritis and others e.g. Sjögren's)

Table 32.3 Autoantibodies in connective tissue disease[2]

Antinuclear antibody (ANA)	High sensitivity > 95%, low specificity for SLE
Anti-dsDNA	High sensitivity and specificity for SLE: found in rheumatoid arthritis, scleroderma, Sjögren's syndrome, others

Antibodies to extractable nuclear antigens (ENA)	
Sm	Highly specific for SLE
U1 RNP	Found in mixed connective tissue disease, SLE
Ro (SSA)	Found in Sjögren's syndrome, SLE and some other connective tissue diseases
La (SSB)	Found in Sjögren's syndrome, SLE
Scl-70 (antitopoisomerase)	Found in 20 - 30% of patients with scleroderma
Jo1	Found in 30% of patients with polymyositis
Anti-centromere	High sensitivity and specificity for CREST syndrome
Anti-neutrophil cytoplasm	High sensitivity and specificity for Wegener's granulomatosis
Anti-phospholipids	Diagnostic in antiphospholipid syndrome
Anti-cardiolipin	Present in 5 - 10% of SLE
Anti-β_2-GPI antibodies	
Lupus anticoagulants	
Anti-prothrombin	

The more specific antibodies for SLE, namely to double-stranded DNA (dsDNA) and extractable nuclear antigens (ENA), should only be ordered if there is a significantly positive ANA.

Antiphospholipid antibody syndrome

This syndrome that may occur with SLE or in isolation and is responsible for recurrent arterial and/or venous thromboembolism, recurrent spontaneous abortions or thrombocytopenia in the presence of antiphos pholipid antibodies but without features of SLE. Livedo reticularis, skin ulcers and neuropsychiatric disturbances have also been noted. If suspected commence aspirin 150 - 300 mg (o) daily and refer to a consultant.

SYSTEMIC LUPUS ERYTHEMATOSUS

SLE (lupus) which is the commonest of the connective tissue disorders, is described as the ‘great pretender’.3 It is a multisystem autoimmune disorder with a wide variety of clinical features that are due to vasculitis (Figure 32.2). Arthritis is the commonest feature of SLE (90% of cases). Milder manifestations outnumber more severe forms.


General
fever
depression
malaise
alopecia
Nervous system
seizures
ataxia
psychoses
headache (intractable)
cranial nerve lesions
Eye disorders
inflammation
retinal vasculitis
mouth ulcers
Skin
butterfly rash
photosensitivity
urticaria
purpura
discoid rash
lymphadenopathy
Cardiac
pericarditis
endocarditis
Chest
pleuritis
effusion
splenomegaly
Renal disorder
glomerulonephritis
proteinuria > 0.5 g/day
abdominal pain
Raynaud's phenomenon
Arthritis
mainly PIP
joints of hand
nail-fold hyperaemia
Tendonitis
myalgia
(muscle pain)
Avascular necrosis
(e.g. femoral head)
Haematological
anaemia
leucopenia
lymphopenia
thrombocytopenia
peripheral neuropathy
Arthritis
small joints
of feet

Figure 32.2 Clinical features of SLE

Features

- Prevalence about 1 in 1000 of population
- Mainly affects women in 'high oestrogen' period (90% of cases)
- Peak onset between 15 and 40 years
- Fever, malaise, tiredness common
- Multiple drug allergies
- Problems with oral contraceptive pill and pregnancy

DxT: polyarthritis + fatigue + skin lesions ° SLE

Criteria for classification of SLE

(SLE ° four or more of these 11 criteria)

- Malar (butterfly) rash
- Discoid rash
- Photosensitivity
- Arthritis (non-erosive arthritis in ≥ 2 peripheral joints)

- Oral ulcers (usually painless)
- Serositis (pleurisy or pericarditis)
- Kidney features (proteinuria or cellular casts)
- Neurological features (intractable headache, seizures or psychosis)
- Haematological features (haemolytic anaemia, leucopenia, lymphopenia or thrombocytopenia)
- Immunological features (positive anti-DNA, antiphospholipid antibodies or anti-Sm tests and false positive syphilis serology)
- Positive antinuclear antibody (ANA) test

Note: Drugs that can cause a lupus-like syndrome are listed in Table 35.2 (on p. 336).

Diagnostic tests

- ESR/CRP - elevated in proportion to disease activity
- ANA test - positive in 95% (perform first) (key test)
- dsDNA antibodies - 90% specific for SLE but present in only 60% (key test)
- ENA antibodies, especially Sm - highly specific
- Rheumatoid factor - positive in 50%
- LE cell test - inefficient and not used

> For suspected SLE, the recommended approach is to perform an ANA test. If positive then order dsDNA and ENA antibodies.

The diagnosis cannot be made on blood tests alone. Supportive clinical evidence is necessary.

Management[4]

- Appropriate explanation, support and reassurance, use of sunscreens (refer to p. 1176).
- Refer to consultant for shared care in a multidisciplinary team.
- Drug treatments[2]
  - mild: NSAIDs (for arthralgia)
  - moderate (especially skin, joint serosa involved): low-dose antimalarials (e.g. hydroxychloroquine up to 6 mg/kg once daily), e.g. 400 mg (o) daily for 3 months, then 200 mg daily long term
  - consider: fish body oil 0.2 mg/kg (o) daily
  - severe: corticosteroids are the mainstay (e.g. prednisolone 7.5 - 15 mg (o) daily) immunosuppressive drugs (e.g. azathioprine, methotrexate)
- Avoid drugs in those in clinical remission and with normal complement levels.
- Other treatments, such as plasma exchange and immunosuppressive regimens, are available for severe disease.
- Keep in mind antiphospholipid antibody syndrome, especially with recurrent fetal loss and thrombotic episodes.

SCLERODERMA (SYSTEMIC SCLEROSIS)

This can present as a polyarthritis affecting the fingers of the hand in 25% of patients, especially in the early stages. Soft tissue swelling produces a ‘sausage finger’ pattern. Scleroderma mainly affects the skin, presenting with Raynaud's phenomenon in over 85% of patients (Figure 32.3).


tight skin
on chest
Characteristic facies
• taut & immobile
• puckering of mouth
• narrow nose
• thin lips
• radial wrinkling upper lip
dysphagia
heartburn
reflux
respiratory symptoms
(e.g. fibrosis)
Cardiac disorders
• arrhythmias
• pericarditis
renal failure
Fingers
'sausage' fingers
Raynaud's
phenomenon
telangiectasia

Figure 32.3 Clinical features of scleroderma

There are three clinical variants:

1. limited cutaneous disease
2. cutaneous with limited organ involvement (CREST)
3. diffuse systemic disease (systemic sclerosis)

Features

- Female to male ratio ° 3:1
- A progressive disease of multiple organs
- Raynaud's phenomenon
- Stiffness of fingers and other skin areas
- 'Bird-like' facies (mouth puckered)
- Dysphagia and diarrhoea (malabsorption)
- Respiratory symptoms
- Cardiac symptoms: pericarditis, etc.
- Look for tight skin on chest (Roman breastplate)

DxT: finger discomfort + arthralgia + GORD (± skin tightness) ° scleroderma

Diagnostic tests[2, 5]

- ESR may be raised
- Normocytic normochromic anaemia may be present
- ANA test - up to 90% positive (relatively specific)
- Rheumatoid factor - positive in 30%
- Anticentromere antibodies - specific (positive in 90% with limited disease and 5% with diffuse)
- Antitopoisomerase I (anti-Scl-70) antibody is specific but only positive in 20 - 40%
- Skin biopsy - increase in dermal collagen

Management

- Refer to consultant for shared care
- Empathic explanation, patient education
- Analgesics and NSAIDs for pain Avoid vasospasm (no smoking, betablockers, ergotamine): nifedipine may help Raynaud's
- Treat malabsorption if present; skin emollients
- D-penicillamine can help if significant systemic or cutaneous involvement[6]

Localised scleroderma

- Morphoea - plaques of erythema with violaceous periphery, feels hard; mainly on trunk
- linear - may be ‘en coup de sabre’ (a sabre stroke)

CREST syndrome (localised scleroderma)

Features

- Calcinosis
- Raynaud's phenomenon
- Oesophageal dysmotility
- Sclerodactyly
- Telangiectasia
- Anti-centromere antibody (invariably positive)

POLYMYOSITIS AND DERMATOMYOSITIS

Polymyositis is an uncommon systemic disorder whose main feature is symmetrical muscle weakness and wasting involving the proximal muscles of the shoulder and pelvic girdles.

Clinical features

- Any age group
- Peak incidence 40 - 60 years
- Female to male ratio ° 2:1
- Muscle weakness and wasting proximal limb muscles
- Main complaint is weakness
- Muscle pain and tenderness in about 50%
- Arthralgia or arthritis in about 50% (resembles distribution of rheumatoid arthritis)
- Dysphagia in about 50% due to oesophageal involvement
- Raynaud's phenomenon
- Consider associated malignancy: lung and ovary

DxT: weakness + joint and muscle pain + violaceous facial rash ° dermatomyositis

The rash

The distinctive rash shows features of photosensitivity. There is violet discolouration of the eyelids, forehead and cheeks, and possible erythema resembling sunburn and periorbital oedema. There is a characteristic rash on the hands, especially the fingers and nail folds. The knees and elbows are commonly involved.

The main features are summarised in Figure 32.4.


heliotrope-coloured photosensitivity rash
proximal muscle weakness and wasting
rash on dorsum of hands
rash on knees

Figure 32.4 Clinical features of polymyositis/dermatomyositis

Diagnosis

- Muscle enzyme studies (serum creatine kinase and aldolase)
- Biopsies - skin and muscle
- EMG studies - show characteristic pattern

Treatment includes corticosteroids and cytotoxic drugs. Early referral is appropriate.

SJÖGREN'S SYNDROME

The underdiagnosed syndrome of dry eyes (keratoconjunctivitis sicca) in the absence of rheumatoid arthritis or any other autoimmune disease is known as primary Sjögren's syndrome (SS):

- primary SS - limited or multisystem
- secondary SS - occurs in association with other CTDs including rheumatoid arthritis (accounts for 50%)

Features

- Fatigue
- Sicca (xerostomia, dry eyes, dry vagina)

- Difficulty swallowing food
- Increased dental caries; denture dysfunction
- Salivary gland enlargement
- Xerotrachea → chronic dry cough; hoarseness
- Dysparunia
- Arthralgia ± non-erosive arthritis

Although considered benign can transform into non-Hodgkin's lymphoma (44 times risk).

DxT: dry eyes + dry mouth + arthritis ° Sjögren's syndrome

Diagnosis

Autoantibody tests - positive ENA, Ro(SSA), La (SS-B)

Management

- Referral to rheumatologist
- Treatment is symptomatic for dry eyes, mouth and vagina; arthralgia
- Hydroxychloroquine or steroids for arthritis

RAYNAUD'S PHENOMENON[2]

(Refer to Chapter 64 on pain in the arm and hand) (pp. 690 - 1).

It is classified as either primary (without associated disease) or secondary (when associated with any CTD).

Patients with primary Raynaud's may progress to a CTD but the liklihood is low (5 - 15%) and the delay to diagnosis is long (average of 10 years).[2] The more severe the Raynaud's, the more likely it is to progress to systemic disease.

Raynaud's is a clinical syndrome of episodic arteriolar vasospasm usually involving the fingers and toes (one or two at a time). It may also involve the nose, ear or nipple.

THE VASCULITIDES

The vasculitides or vasculitis syndromes are a heterogeneous group of disorders involving inflammation and necrosis of blood vessels, the clinical effects and classification depending on the size of the vessels involved.

Small vessel vasculitis is the common type encountered in practice. Medium vessel vasculitis includes polyarteritis nodosa and large vessel vasculitis includes giant cell arteritis.

Small vessel vasculitis

This is associated with many important disorders, such as rheumatoid arthritis, SLE, bacterial endocarditis, Henoch-Schönlein purpura and hepatitis B. Skin lesions are usually associated with these disorders and the most common presentation is painless, palpable purpura, such as occurs with Henoch-Schönlein purpura.

Rarer but deadly causes

The major vasculitides called systemic vasculidities are polyarteritis nodosa (PN), polymyalgia rheumatica (PR), giant cell arteritis (GCA), Takayasu's arteritis, Bechet's syndrome, Churg-Strauss vasculitis and Wegener's granulomatosis (WG). Unfortunately, many of these patients die or become terminally ill before the diagnosis is suspected.

Practice tip

If a serious ANCA associated disease is suspected early diagnosis is life-saving because of sinister kidney damage. Perform a urine examination for haematuria and proteinuria. If positive order an antineutrophil cytoplasmic antibody (ANCA) test. If positive refer urgently.

Henoch-Schönlein purpura

More details are presented about this vasculitis disorder on pp. 405 - 6.

Takayasu's arteritis[2]

Known as ‘pulseless disease’ or ‘aortic arch syndrome’, this vasculitis involves the aortic arch and other major arteries. It typically affects young Japanese female adults. Features include absence of peripheral pulses and hypertension.

POLYARTERITIS NODOSA

The hallmark of PN is necrotising vasculitis of the small and medium arteries leading to skin nodules, infarctive ulcers and other serious manifestations (Figure 32.5). The cause is unknown but associations are found with drug abusers (especially adulterated drugs), B-cell lymphomas, other drugs and hepatitis B surface antigen. It should be suspected in any multisystemic disease of obscure aetiology.


General
malaise
weight loss
fever
Cardiac
myocardial infarction
arrhythmia
heart failure
hypertension
Lungs
pleurisy
asthma
Renal
glomerulonephritis
arthritis
nodules
ulcer
livedo reticularis

Figure 32.5 Clinical features of polyarteritis nodosa

Clinical features

- Young to middle-aged men
- Constitutional symptoms: fever, malaise, myalgia, weight loss
- Migratory arthralgia or polyarthritis
- Subcutaneous nodules along arterial lines

- Livedo reticularis and skin ulcers
- Kidney impairment and hypertension
- Cardiac disorders: arrhythmia, failure, infarction
- Diagnosis confirmed by biopsy or angiogram
- ESR raised
- Treatment with corticosteroids and immunosuppressants
- Death is usually from kidney disease

DxT: arthralgia + weight loss + fever (± skin lesions) ° polyarteritis nodosa

GIANT CELL ARTERITIS AND POLYMYALGIA RHEUMATICA

The basic pathology of this very important disease complex is GCA (synonyms: temporal arteritis, cranial arteritis). The clinical syndromes are polymyalgia rheumatica and temporal arteritis. The clinical manifestations of polymyalgia rheumatica invariably precede those of temporal arteritis, of which there is about a 20% association. The diagnosis is based on clinical grounds. No definite cause has been found.

Clinical features of polymyalgia rheumatica

- Pain and stiffness in proximal muscles of shoulder and pelvic girdle, cervical spine (Figure 32.6)



Figure 32.6 Polymyalgia rheumatica: typical sites of areas of pain and stiffness

- Symmetrical distribution
- Typical ages 60 - 70 years (rare < 50)
- Both sexes: more common in women
- Early morning stiffness
- May be systemic symptoms: weight loss, malaise, anorexia
- Painful restriction of movement of shoulders and hips
- Signs may be absent later in day

Differential diagnosis: Polymyalgic onset rheumatoid arthritis

DxT: malaise + painful shoulder girdle + morning stiffness (>50 years) ° polymyalgia rheumatica

Clinical features of temporal arteritis

- Headache - unilateral, throbbing
- Temporal tenderness
- Loss of pulsation of temporal artery
- Jaw claudication
- Biopsy of artery (5 cm) is diagnostic

DxT: fatigue + headache + jaw claudication ° temporal arteritis

Investigation

- No specific test for polymyalgia rheumatica
- ESR - extremely high, around 100
- C-reactive protein - elevated
- Mild anaemia (normochromic, normocytic)

Treatment

Prednisolone

- Starting dose
  - temporal (giant cell) arteritis: 40 - 60 mg (o) daily initially for 2 - 4 weeks (+ aspirin 100 mg/day) then gradual reduction according to ESR/CRP
  - polymyalgia rheumatica: 15 mg (o) daily for 2 - 4 weeks, then taper
- Taper down gradually to the minimum effective dose (often < 5 mg daily) according to the clinical response and the ESR and CRP. Aim for treatment for 2 years. Relapses are common.

Other drug

- Azathioprine or methotrexate can be used as steroid-sparing agents.

Practice tip

In giant cell arteritis, a delay in diagnosis after presenting with amaurosis fugax and non-specific symptoms can have tragic consequences, in the form of ischaemic events such as blindness and strokes.

Behçet's syndrome

Behçet's syndrome is systemic (multiorgan) vasculitis of unknown aetiology, affecting veins and arteries of all sizes. The main feature is painful oral ulceration and the hallmark is the ‘pathergy’ reaction whereby simple trauma such as a pin prick can cause a papule or pustule to form within a few hours at the site.

Features

- M:F ° 2:1
- Recurrent oral and/or genital ulceration

- Arthritis (usually knees)
- Ocular symptoms - pain, reduced vision, floaters (ocular inflammation)

There is no specific diagnostic test.

Associated problems/complications: repeated uveitis and retinitis → blindness, colitis, venous thrombosis, meningoencephalitis

Treatment: high dose steroids

Practice tip

Patients with Behcet's eye disease should be referred promptly for an opthalmological opinion, which may be sight-saving.[2]

Wegener's granulomatosis

In this rare granulomatous vasculitis of unknown cause there is a classic triad: upper respiratory tract (URT) granuloma, fleeting pulmonary shadows (nodules) and glomerulonephritis. Without treatment it is invariably fatal and sometimes the initial diagnosis is that made at autopsy. It is difficult to diagnose, especially as the patient (usually young to middleaged) presents with a febrile illness and respiratory symptoms, but early diagnosis is essential. It usually gets confused with benign nasal conditions.

Clinical features

- Adolescence to elderly, mean age 40 - 45 years
- Constitutional symptoms (as for PN)
- Lower respiratory tract (LRT) symptoms (e.g. cough, dyspnoea)
- Oral ulcers
- Upper respiratory symptoms: rhinorrhoea, epistaxis, sinus pain
- Eye involvement - orbital mass
- Polyarthritis
- Kidney involvement - usually not clinically apparent (about 75% get glomerulonephritis)
- Chest X-ray points to diagnosis - multiple nodes, cavitations
- Antineutrophil antibodies (c-ANCA) are a useful diagnostic marker (not specific)
- Diagnosis confirmed by biopsy, usually an open lung biopsy
- Better prognosis withearly diagnosis and treatment with cyclophosphamide.

DxT: malaise + URTs (e.g. rhinitis, sinusitis) + LRTs (e.g. wheeze, cough) ° Wegener's granulomatosis

REFERENCES

1. Kumar PJ, Clark ML. *Clinical Medicine* (5thedn). London: Saunders, 2005: 557 - 69.
2. Mashford L. *Therapeutic Guidelines: Rheumatology* (Version 1). Melbourne: Therapeutic Guidelines Ltd, 2006.
3. Hanrahan P. The great pretender: SLEs. Aust Fam Physician, 2001; 30(7): 636 - 40.
4. Ryan PFJ. Systemic lupus erythematosus. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 497 - 9.
5. Randell P. Scleroderma. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 458 - 60.
6. Steen VD et al. D-penicillamine therapy in progressive systemic sclerosis. Ann Intern Med, 1982; 97: 652 - 9.

CHAPTER 33. NEUROLOGICAL DILEMMAS

The disease is of long duration; to connect, therefore, the symptoms which occur in its later stages with those which mark its commencement, requires a continuance of observation of the same case, or at least a correct history of its symptoms, even for several years.

JAMES PARKINSON (1755 - 1824), AN ESSAY ON THE SHAKING PALSY

In general practice there are many neurological problems that present a diagnostic dilemma, with some being true masquerades for the non-neurologist. This applies particularly to various seizure disorders, space-occupying lesions in the cerebrum and the cerebellum, demyelinating disorders, motor neurone disorders and peripheral neuropathies.

The most common pitfall that occurs with neurological disorders is misdiagnosis, and the most common reason for misdiagnosis is an inadequate history. Failure to appreciate the neurological meaning of points elicited during the history is another reason for misdiagnosis.

Some very important neurological disorders are presented in this section: Parkinson's disease, which is common and can be easily misdiagnosed, especially when the classic ‘pill rolling’ tremor is absent or mild; multiple sclerosis (MS), because it is difficult to diagnose initially; and acute idiopathic demyelinating polyneuropathy (Guillain-Barré syndrome), because it can be rapidly fatal if misdiagnosed. MS can masquerade as almost anything - ‘If you don’t know what it is, think of MS’.

Another brain teaser for the family doctor is to diagnose accurately the various types of epilepsy. The most commonly misdiagnosed seizure disorders are complex partial seizures or atypical generalised tonic - clonic seizures (see Chapter 54).[1] Even more difficult is the differentiation of real seizures from pseudo- or non-epileptic seizures.

DIPLOPIA

The onset of diplopia (double vision) in adults is often acute, very distressing and invariably easy to diagnose. It can be divided into two distinct types: uniocular (confined to one eye) or binocular, which usually results from extraocular muscular imbalance or weakness. The type of binocular diplopia - vertical, horizontal or oblique - provides clues in identifying the affected muscle.

Causes of uniocular diplopia

- Early cataract (common in older patient - usually in dim light and at night)
- Dislocated lens
- Severe astigmatism
- Psychogenic/functional

Causes of binocular diplopia

- Ocular nerve palsies (3, 4 or 6); consider:
  - CVA or TIA
  - tumour (orbital or intracerebral)
  - aneurysm
  - diabetes mellitus
  - arteritis
  - head injury
  - ophthalmoplegic migraine (transient)
- muscle tethering (e.g. blow-out orbital fracture)
- concussion
- multiple sclerosis (recurrent diplopia)
- myasthenia gravis multiple muscle movement
- hyperthyroidism multiple muscle movement

Office tests

Test for double vision with each eye occluded. If diplopia persists it is uniocular. If, however, double vision disappears when either eye is covered, there is a defect of one of the muscles moving the eyeball. Determine whether diplopia occurs in any particular direction of gaze. It is most marked when moved in the direction of action of the weak muscle. Ask patient to follow your finger, red pin or penlight with both eyes and move it in an H pattern.

- 3rd nerve - eye turned out: divergent squint
- 6th nerve - failure to abduct: convergent squint

See Figure 33.1.



Figure 33.1 Direction of movement of the right eye indicating the responsible extra-ocular muscles and cranial nerves (3 ° oculomotor, 4 ° trochlear, 6 ° abducens)

Laboratory test

- ESR (consider arteritis)

Note:

- Exclude 3rd and 6th nerve palsies as they may be secondary to life-threatening conditions.
- Refer urgently if diplopia is binocular, of recent onset and persistent.

MOTOR WEAKNESS

Muscle weakness is a common feature of many disorders ranging from neurogenic and myogenic disorders to metabolic and psychiatric. It is very important clinically to be able to differentiate upper motor neurone (UMN) signs from lower motor neurone (LMN) signs (Table 33.1).

Table 33.1 Clinical differences between a lower motor neurone lesion and an upper motor neurone lesion

Manifestation	UMN	LMN
Weakness	Present	Present
Wasting	Absent or mild	Marked
Power	Reduced	Reduced
Tone	Usually increased (spastic paralysis) ±clonus	Absent or decreased (flaccid paralysis)
Fasciculations	Absent	May be present
Reflexes	Brisk tendon reflexes Abdominal absent Extensor plantar response	Absent or diminished

Upper motor neurone lesions

UMN signs occur when a lesion has interrupted a neural pathway at a level above the anterior horn cell.[2] Examples include lesions in motor pathways in the cerebral cortex, internal capsule, brain stem or spinal cord.

Clinical examples include stroke (thrombosis, embolism or haemorrhage in the brain), tumours of the various pathways, and demyelinating disease, such as multiple sclerosis and infection, for example, HIV.

Lower motor neurone lesions

LMN signs occur when a lesion interrupts peripheral neural pathways from the anterior horn cell, that is, the spinal reflex arc.

Clinical examples include peripheral neuropathy, Guillain-Barré syndrome, motor neurone disease, poliomyelitis and a thickened peripheral nerve (e.g. leprosy).

Note: A spinal cord lesion causes LMN signs at the level of the lesion and UMN signs below that level.

Neurogenic and myogenic muscle weakness

It is also important to distinguish between weakness caused by neurological conditions, especially those causing LMN lesions and muscular disorders. The features are compared in Table 33.2.

Table 33.2 Muscle weakness: main clinical differences between neurogenic and myogenic lesions

Myogenic weakness	Neurogenic weakness
Reflexes often present despite severe weakness	Reflexes often absent despite minimal weakness
Weakness out of proportion to wasting	Wasting out of proportion to weakness
Sensation normal	±Sensory changes
No fasciculation	Fasciculation a feature

MOTOR NEURONE DISEASE (MND)

MND is a progressive neuromuscular disorder resulting in muscular limb and bulbar weakness due to death of motor neurones in the brain, brain stem and spinal cord. The sensory system is not involved, nor the cranial nerves to the eye muscles. Five to 10% of MND is inherited with an autosomal dominant pattern; the rest is sporadic. The three main patterns are:

1. amyotrophic lateral sclerosis - combined LMN muscle atrophy plus UMN hyper-reflexia, leading to progressive spasticity
2. progressive muscle atrophy - wasting beginning in the proximal muscles; widespread fasciculation
3. progressive bulbar and pseudobulbar palsy - LMN lesions of the brainstem motor nuclei resulting in wasted fibrillating tongue, weakness of chewing and swallowing, and of facial muscles

Symptoms and signs

- Weakness or muscle wasting - first noticed in hands (weak grip) or feet
- Stumbling (spastic gait, foot drop)
- Difficulty with swallowing
- Difficulty with speech, e.g. slurring, hoarseness
- Fasciculation (twitching) of skeletal muscles and tongue
- Cramps
- Emotional instability, depression
- ±Muscle pain

The diagnosis is clinical. There are no diagnostic tests, but neurophysiological tests help differentiate from other conditions.

MND is incurable and progresses to death usually within 3 - 5 years from ventilatory failure/aspiration pneumonia.

No treatment is proven to influence outcome although Riluzole, a sodium channel blocker, appears to slow progression slightly. Baclofen 10 mg bd may help symptoms of cramp. Botulinum toxin may help spasticity and propantheline or amitriptyline for drooling.

TREMOR

Tremor is an important symptom to evaluate correctly. A list of causes is presented in Table 33.3. A common pitfall in patients presenting with tremor is for Parkinson's disease to be diagnosed as benign essential tremor and for benign essential tremor to be diagnosed as Parkinson's disease, but the clinical distinction is not always easy and it must be remembered that as many as 20% will experience both concurrently.

Tremors can be classified as follows:

Table 33.3 Causes of tremor

Physiological
Benign essential (familial) tremor
Senile
Anxiety, including hyperventilation
Hyperthyroidism
Toxic (e.g. alcohol, liver failure, uraemia)
Drugs (e.g. lithium, narcotic withdrawal)
Parkinson's disease
Drug-induced parkinsonism
Cerebellar disease
Cerebral tumour (frontal lobe)
Alzheimer's dementia
Wilson's syndrome
Miscellaneous (e.g. red-nucleus lesion, hypoglycaemia)

Resting tremor - Parkinsonian

The tremor of Parkinson's disease is present at rest. The hand tremor is most marked with the arms supported on the lap and during walking. The characteristic movement is ‘pill-rolling’ where movement of the fingers at the metacarpophalangeal joints is combined with movements of the thumb. The resting tremor decreases on finger - nose testing but a faster action tremor may supervene. The best way to evoke the tremor is to distract the patient, such as focusing attention on the left hand with a view to ‘examining’ the right hand or by turning the head.

Action or postural tremor

This fine tremor is noted by examining the patient with the arms outstretched and the fingers apart. The tremor may be rendered more obvious if a sheet of paper is placed over the dorsum of the hands. The tremor is present throughout movement, being accentuated by voluntary contraction.

Causes include:

- essential tremor (also called familial tremor or benign essential tremor)
- senile tremor
- physiological
- anxiety/emotional
- hyperthyroidism
- alcohol
- drugs, for example, drug withdrawal (e.g. heroin, cocaine, alcohol), amphetamines, lithium, sympathomimetics (bronchodilators), sodium valproate, heavy metals (e.g. mercury), caffeine, amiodarone
- phaeochromocytoma

Intention tremor (cerebellar disease)

This coarse oscillating tremor is absent at rest but exacerbated by action and increases as the target is approached. It is tested by ‘finger - nose - finger’ touching or running the heel down the opposite shin, and past pointing of the nose is a feature. It occurs in cerebellar lobe disease and with lesions of cerebellar connections.

Flapping (metabolic tremor)

A flapping or 'wing-beating' tremor is observed when the arms are extended with hyperextension of the wrists. It involves slow, coarse and jerky movements of flexion and extension at the wrists.

Note: Flapping (asterixis) is not strictly a tremor.

Causes

- Wilson's syndrome
- Hepatic encephalopathy
- Uraemia
- Respiratory failure
- Lesions of the red nucleus of the midbrain (the classic cause of a flap)

ESSENTIAL TREMOR

Essential tremor, which is probably the most common movement disorder, has been variously called benign, familial, senile or juvenile tremor. However, it is not always benign and there is no family history in about half the cases.

Features

- Autosomal dominant disorder (variable penetrance)
- Often begins in early adult life, even adolescence
- Begins with a slight tremor in one hand and spreads to the other with time
- May involve head (titubation), chin and tongue and rarely trunk and legs
- Interferes with writing (not micrographic), handling cups of tea and spoons, etc.
- Tremor most marked when arms held out (postural tremor)
- Tremor exacerbated by anxiety
- May affect speech if it involves bulbar musculature
- Relieved by alcohol
- Can swing arm and gait normal

Triad of features

- Positive family history
- Tremor with little disability
- Normal gait

Distinguishing essential tremor from Parkinson's disease

This is not always easy as a postural tremor can be present in Parkinson's disease although the hand tremor is most marked at rest with the arms supported on the lap. Parkinsonian tremor is slower at 4 - 6 Hz while essential tremor is much faster at around 8 - 13 Hz.

A most useful way to differentiate the two causes is to observe the gait. It is normal in essential tremor but in Parkinson's there may be loss of arm swing and the step is usually shortened.

Management of essential tremor

Most patients do not need treatment and all that is required is an appropriate explanation.1 If necessary, use propranolol (first choice) or primidone.[3] A typical starting dose of propranolol is 10 - 40 mg bd; many require 120 - 240 mg/day.3 If the tremor is only intrusive at times of increased emotional stress, intermittent use of benzodiazepines (e.g. lorazepam) 30 minutes before exposure to the stress may be all that is required. Modest alcohol intake (e.g. a glass of scotch) is very effective. A standard drink of alcohol often alleviates the tremor. Larger doses of alcohol have no additional effect.

PARKINSON'S DISEASE

Parkinson's is a disease of the automatic processor of the brain which relies on dopamine to maintain movements at a selected size and speed. Loss of dopamine causes movements to become smaller and slower. Lewy bodies in the neurones are the pathognomic sign of Parkinson's disease.4 Genetic factors occur in 5% of individuals.

One of the most important clinical aspects of Parkinson's disease, which has a slow and insidious onset, is the ability to make an early diagnosis. Sometimes this can be very difficult especially when the tremor is absent or mild, as occurs with the atherosclerotic degenerative type of parkinsonism. The lack of any specific abnormality on special investigation leaves the responsibility for a diagnosis based on the history and examination. As a general rule of thumb the diagnosis of Parkinson's disease is restricted to those who respond to levodopa - the rest are termed parkinsonism.

Key facts and checkpoints

- Parkinson's disease is a most common and disabling chronic neurological disorder.
- The prevalence in Australia is 100 - 120 per 100 000.5
- The mean age of onset is between 58 and 62 years.5
- The incidence rises sharply over 70 years of age.5
- The classic triad of Parkinson's disease (Figure 33.2) is:
    - tremor
    - rigidity
    - bradykinesia (poverty of movement)
- Non-motor automatic dysfunctions
    - cognition
    - behaviour
    - mood
- Hemi-parkinsonism can occur; all the signs are confined to one side and thus must be differentiated from hemiparesis. In fact, most cases of Parkinson's disease start unilaterally.
- Always consider drug-induced parkinsonism. The usual drugs are phenothiazines, butyrophenones and reserpine. Tremor is uncommon but rigidity and bradykinesia may be severe.

Refer to Table 33.4 (on p. 297).

Table 33.4 Parkinson's disease: symptoms and signs (a checklist)

General	• Tiredness Lethargy Restlessness
Tremor	• Present at rest Slow rate - 4 to 6 cycles per second Alternating, especially arms Pill-rolling (severe cases)
	Note: may be absent or unilateral
Rigidity	• ‘Cogwheel’ - ‘juddering’ on passive extension of the forearm Lead pipe - limbs resist passive extension through movement
Bradykinesia/hypokinesia	• Slowness of initiating a movement Difficulty with fine finger tasks Micrographia (Figure 33.3) Masked facies Relative lack of blinking Impaired convergence of eyes Excessive salivation (late) Difficulty turning over in bed and rising from a chair Slow monotonous speech/dysarthria
Gait disorder	• No arm swing on one or both sides Start hesitation

	Slow and shuffling Short steps (*petit pas*) Slow turning circle ‘Freezing’ when approaching an obstacle Festination
Disequilibrium	• Poor balance Impaired righting reflexes Falls (usually late)
Posture	• Progressive forward flexion of trunk Flexion of elbow at affected side
Autonomic symptoms	• Constipation (common) Postural hypotension Depression (early)
Psychiatric	• Progressive dementia in 30 - 40% usually after 10 years[6]

- Three major traps in missing early diagnosis:[5]
    - age: 10 - 15% are < 50 years at onset
    - belief that it is a disease of men: M ° F
    - absence of resting tremor (only 50% have it at onset)


masked facies
staring expression
slow, monotonous speech
failure to swing arm
pill-rolling tremor at rest
slow and shuffling gait short steps

Figure 33.2 Basic clinical features of Parkinson's disease

Physical signs

- Power, reflexes and sensation are usually normal.
- The earliest abnormal physical signs to appear are loss of dexterity of rapid alternating movements and absence of arm swing, in addition to increased tone with distraction.
- Positive frontal lobe signs, such as grasp and glabellar taps (only allow three blinks), are more common with parkinsonism.

Note: There is no laboratory test for Parkinson's - it is a clinical diagnosis. Hypothyroidism and depression, which also cause slowness of movement, may cause confusion with diagnosis.

Note: The Steele-Richardson syndrome (parkinsonism, mild dementia and vertical gaze dysfunction) is worth considering.

Principles of management

- Provide appropriate explanation and education.
- Explain that Parkinson's disease is slowly progressive, and is improved but not cured by treatment. It is associated with increased mortality (RR death compared with general population ranges 1.6 to 3). The question of whether treatment reduces mortality remains controversial.[7]
- Refer to a specialist for shared care.
- Support systems are necessary for advanced Parkinson's disease.

- Walking sticks with appropriate education into their use may be necessary to help prevent falls and constant care is required, so that admission to a nursing home for end-stage disease may be appropriate.



Figure 33.3 Micrographia, one of the signs of Parkinson's disease

Pharmacological management[8, 9]

Avoid postponing treatment. It should be commenced as soon as symptoms interfere with working capacity or the patient's enjoyment of life. This will be apparent only if the correct questions are asked as the patient may accept impaired enjoyment without appreciating that it is due to Parkinson's disease. Start low - Sinemet 100/25 (½ tab bd). There is usually no difference between the levodopa preparations. The dosage should be tailored so that the patient neither develops side-effects nor is on an inadequate dose of medication without significant therapeutic benefit (Table 33.5).

Table 33.5 Anti-Parkinson drugs[8, 9]

Agent	Usual dose	Main side effects
Dopaminergic (standard and slow release)		• Nausea and vomiting
		• Involuntary dyskinetic movements
• levodopa + benserazide	100/25 - 250/62.5 mg tds	• Psychiatric disturbances
• levodopa + carbidopa	100/25 - 250/50 mg tds	• On - off phenomena
		• End of dose failure
		• Constipation
Dopamine agonists		• Nausea and vomiting
• bromocriptine	5 - 15 mg bd	• Dizziness, fatigue
• carbergoline	0.5 - 6 mg daily	• Psychiatric disturbances
• pergolide	0.05 - 1.5 mg tds	• Pleuropulmonary changes
Anticholinergics		
• benzhexol	2 mg bd or tds	• Dryness of mouth
• benztropine	1 - 2 mg bd	• Confusion in elderly
• biperiden	1 - 2 mg bd	• Contraindicated in glaucoma and prostatism
• orphenadrine	100 mg bd	• Other anticholinergic effects (e.g. constipation, blurred vision)
COMT inhibitors		
• entacapone	200 mg with each dose levodopa	• Diarrhoea
		• Sleep problems
		• Potentiation of levodopa
Others		
• amantadine	100 mg bd	• Nausea and vomiting
		• Psychiatric disturbances

		• Ankle oedema • Livedo reticularis
• apomorphine	SC injection (refer to schedule)	• Nausea • Psychosis • Dyskinesia
• selegiline (a MAO-B inhibitor)	2.5 - 5 mg once daily or bd	• Dry mouth • Neuropsychiatric disturbances • Nausea • Dizziness, fatigue • Insomnia

The older drugs, such as anticholinergics and amantadine, have minimal usage in modern management as levodopa, which basically counters bradykinesia, is the best drug and the baseline of treatment. With the onset of disability (motor disturbances) levodopa in combination with a decarboxylase inhibitor (carbidopa or benserazide) in a 4:1 ratio should be introduced. Levodopa therapy does not significantly improve tremor but improves rigidity, dyskinesia and gait disorder. Consider benzhexol or benztropine if tremor is the feature, especially in young patients.

Pergolide can be used in treatment, especially with the levodopa 'on - off' phenomenon (fluctuations throughout the day). It appears to be most effective when used in combination. The major side-effects of pergolide are similar to levodopa. Dyskinesia and nausea are less problematic but severe psychiatric disturbances are more common with bromocriptine (a dopamine agonist). It should therefore be used with caution in patients with a history of confusion or dementia. Selegiline promises to be an effective first-line drug. If there is associated pain, depression or insomnia, the tricyclic agents (e.g. amitriptyline) can be effective.

Entacapone has the potential to increase 'on time' and reduce motor fluctuations in levodopa treated patients who are beginning to experience end of dose failure. The initial dose is 200 mg.

Treatment strategy[8, 9]

Mild

(minimal disability)

- Levodopa preparation (low dose) e.g. levodopa 100 mg + carbidopa 25 mg (½ tab bd - increase gradually as necessary to 1 tab (o) tds)
- Amantadine 100 mg (o) bd may help the young or the elderly for up to 12 months
- Selegiline (can be used as monotherapy and postpone need for levodopa)

Moderate

(independent but disabled, e.g. writing, movements, gait)

- Levodopa preparation
- Add if necessary - pergolide or cabergoline or bromocriptine
  - pergolide 50 μg (o) bd, gradually increasing to 1.5 mg (max.) bd
  - cabergoline 0.5 mg (o) daily, increasing to 6 mg (max.) once daily

Severe

(disabled, dependent on others)

- Levodopa (to maximum tolerated dose) + pergolide or cabergoline
- Add entacapone 200 mg (o) with each dose of levodopa, e.g. Statevo®
- Consider antidepressants

Long-term problems

After 3 - 5 years of levodopa treatment side-effects may appear in about one-half of patients:[5]

- involuntary movements - dyskinesia (use lower dose + pergolide or cabergoline)
- end of dose failure (reduced duration of effect to 2 - 3 hours only) - use entacapone

- ‘on - off ’ phenomenon (sudden inability to move with recovery in 30 - 90 minutes)
- early morning dystonia, such as clawing of toes (due to disease - not a side-effect); management of motor problems is summarised in Table 33.6

Table 33.6 Management of motor problems in treated Parkinson's disease[4]

Motor problem	Management
End of dose failure	• Dosages closer together • Slow-release preparations • MAO-B inhibitor (e.g. selegiline) • Dopaminergic agonist (e.g. pergolide)
‘On - off’ phenomenon	• Subcutaneous apomorphine for ‘off’ phase (1 h action) with domperidone (o) to prevent vomiting • Levodopa and ascorbic acid solution
Loss of efficacy	• Increase levodopa dose as high as possible • Dopaminergic agonist (e.g. pergolide)
Peak dose dyskinesia	• Decrease levodopa dose • MAO-B inhibitor, if efficacy lost • Dopaminergic agonist (e.g. pergolide)
Early morning dystonia	• Slow release levodopa • Dopaminergic agonists (e.g. pergolide)
Nocturnal akinesia	• Slow release levodopa • Dopaminergic agonist

Advanced disease[8]

- Apomorphine can be used for severe akinesia not responsive to levodopa:
  - apomorphine 600 μg to 6 mg (mean 3.4 mg) SC
- For nausea and vomiting side-effects:
  - domperidone 20 mg (o) tds 24 hours prior to apomorphine
- Better control may also be achieved with:
  - amantadine 100 mg (o) bd

Contraindicated drugs

- Phenothiazines
- Butyrophenones

Surgical treatment

The indication for surgery (pallidotomy or stereotactic thalamotomy)10 is the presence of tremor or rigidity not responding to chemical therapy. It is considered more appropriate for younger patients with a unilateral tremor.[8] A high success rate has been claimed. It alleviates tremor and rigidity but does not prevent progression of bradykinesia, dysarthria or dementia. However, two systematic reviews found limited evidence that pallidotomy reduces contralateral tremor and rigidity during ‘off’ time but carried significant risk of morbidity and mortality.[7]

When to refer

If the diagnosis is unclear at the time of initial presentation, it is appropriate either to review the patient at a later date or to refer the patient for more neurological assessment.

Once diagnosed or highly suspected it is best to refer to establish the diagnosis and to seek advice on initiation of treatment. Patients and families usually prefer this approach. In the initial years before motor fluctuations develop, management could be performed by the general practitioner according to an overall plan developed in liaison with a neurological colleague. When fluctuations develop and end-stage diseases manifest (e.g. gait disorders), specialist supervision is appropriate.[1]

Cognitive impairment with Parkinson's disease[4]

This may be due to multiple factors including Parkinson's associated dementia, Alzheimer's disease and medication, all of which can induce psychosis, but L-dopa is the least likely. Neuropsychiatric symptoms, which can be varied and bizarre and usually worse in the evening, can occur. Factors contributing to psychosis are illustrated in Figure 33.4. Management is based on monotherapy with gradual build up of L-dopa to maximum tolerated dose, e.g. 450 - 600 mg/day.



Figure 33.4 Factors contributing to psychosis

Management of psychotic problems

- Treat as an inpatient.
- Exclude and treat co-morbidities, e.g. UTI.
- Eliminate and wean off worst drugs.
- Increase L-dopa slowly to 150 mg tds or qid.
- Give quetiapine at night-time.

Practice tips

- Levodopa is the gold standard for therapy.
- Longer acting levodopa preparations may reduce the 'end of dose' failure effect but remember the possible need for a 'kick start' with short-acting preparations (e.g. first thing in the morning).
- Ensure that a distinction is made between drug-induced involuntary movements and the tremor of Parkinson's disease.
- Keep the dose of levodopa as low as possible to avoid these drug-induced involuntary movements.
- In the elderly with a fractured hip always consider Parkinson's disease (a manifestation of disequilibrium).
- Remember the balance of psychosis and Parkinson's disease in treatment.
- Keep in mind the 'sundown' effect - patients often go psychotic as the sun goes down.
- Don't fail to attend to the needs of the family, who often suffer in silence.
- Bromocriptine and pergolide should be used very cautiously in the elderly because of possible acute psychotic reactions.
- If drugs are to be withdrawn they should be withdrawn slowly.

MULTIPLE SCLEROSIS

Multiple sclerosis (MS) is the most common cause of progressive neurological disability in the 20 - 50 year age group.11 It is generally accepted that MS is an autoimmune disorder. Genetic and environmental factors are believed to play a role.12 Early diagnosis is difficult because MS is characterised by widespread neurologic lesions that cannot be explained by a single anatomical lesion, and the various symptoms and signs are subject to irregular exacerbations and remissions. The most important issue in diagnosis is the need for a high index of suspicion.

MS is a primary demyelinating disorder with demyelination occurring in plaques throughout the white matter of the brain, brain stem, spinal cord and optic nerves. The clinical features depend on their location.

Clinical features

(See Figure 33.5.)


impaired vision
diplopia
cerebellar signs
• intention tremor
• ataxia
dizziness
fatigue
band sensations around trunk or limbs
sexual difficulties
bladder urgency
incontinence
constipation
Features
UMN lesion
> one lesion
separate times
limb (any)
• weakness
• numbness
• tingling
• or unsteadiness
paraparesis

Figure 33.5 Basic clinical signs in multiple sclerosis

- More common in females
- Peak age of onset is in the fourth decade
- Transient motor and sensory disturbances
- UMN signs
- Symptoms develop over several days but can be sudden
- Monosymptomatic initially in about 80%
- Multiple symptoms initially in about 20%
- Common initial symptoms include:
  - visual disturbances of optic neuritis
    - blurred vision or loss of vision in one eye (sometimes both)
    - central scotoma with pain on eye movement (looks like unilateral papilloedema)
  - diplopia (brain-stem lesion)
  - weakness in one or both legs, paraparesis or hemiparesis
  - sensory impairment in the lower limbs and trunk
    - numbness, paraesthesia
    - band-like sensations
    - clumsiness of limb (loss of position sense)

        - feeling as though walking on cotton wool
    - vertigo (brain-stem lesion)
- Subsequent remissions and exacerbations that vary from one individual to another
- There is a progressive form, especially in women around 50 years
- Anxiety, depression and other mood disorders are relatively common

Neurological examination

The findings depend on the site of the lesion or lesions and include optic atrophy, weakness, hyperreflexia, extensor plantar responses, nystagmus (two types - cerebellar or ataxic), ataxia, incoordination and regional impairment of sensation.

Symptoms causing diagnostic confusion

- Bladder disturbances, including retention of urine and urgency
- ‘Useless hand’ due to loss of position sense
- Facial palsy
- Trigeminal neuralgia
- Psychiatric symptoms

In established disease common symptoms are fatigue, impotence and bladder disturbances.

Diagnosis

The diagnosis is clinical and depends on the following determinants:

- Lesions are invariably UMN.
- Lesions affect the CNS white matter.
- > 1 part of CNS is involved, although not necessarily at time of presentation.
- Episodes are separated in time (it is possible to make a diagnosis with the first episode).

Other neurological disorders such as infections (e.g. encephalitis), malignancies, spinal cord compression, spinocerebellar degeneration and others must be excluded.

Investigations

- Lumbar puncture: oligoclonal IgG detected in CSF in 90% of cases11 (only if necessary)
- Visual evoked potentials: abnormal in about 90% of cases
- CT scan: rarely demonstrates MS lesions but useful in excluding other pathology
- MRI scan: usually abnormal, demonstrating MS lesions in about 90% of cases11

Course and prognosis

- The course is variable and difficult to predict. An early onset (< 30 years) is usually ‘benign’ while a late onset (≥50 years) is often ‘malignant’.
- MS follows a classic history of relapses and remissions in 80 - 85% of patients.[13]
- The rate of relapse is about once in 2 years.
- About 20% have a progressive course from the onset with a progressive spastic paraparesis (applies mainly to late-age onset).
- The average duration of MS is about 30 years from diagnosis to death.[11]
- A benign course occurs in about 40% of patients with 10 - 20% never suffering major disability.
- The median time to needing a walking aid is 15 years.[8]
- The likelihood of developing MS after a single episode of optic neuritis is about 60%.

Management principles

- All patients should be referred to a neurologist for confirmation of the diagnosis, which must be accurate.
- Explanation about the disorder and its natural history should be given.
- Acute relapses require treatment if causing significant disability.
- Depression and anxiety, which are common, require early treatment.

Relapses

Mild relapses

Mild symptoms, such as numbness and tingling, require only confirmation, rest and reassurance.

Moderate relapses[8]

For symptoms that are unpleasant or disabling use a short course of high-dose prednisolone.

- prednisolone 75 mg (o) once daily for 4 days, then 50 mg (o) daily for 4 days, then 25 mg (o) daily for 4 days

Severe relapses or attacks[8, 14]

These attacks include optic neuritis, paraplegia or brain-stem signs. Admit to hospital for IV therapy:

- methylprednisolone 1 g in 200 mL saline by slow IV injection (3 hours) daily for 3 to 5 days

Observe carefully for cardiac arrhythmias.

Drugs to prevent relapses[14]

Currently available immunomodulators are the interferons and glatiramir.

Interferon beta-1b (SC injection) and beta-1a (IM injection) appear to be effective (but expensive) for those with frequent and severe attacks.

Evidence-based medicine

The best evidence to date indicates that the interferons and methylprednisolone are likely to be beneficial but glatiramir, IV immunolobulin, cytotoxic agents and plasma exchange have unkown effectiveness.[15] One RCF found a modest effect with glatiramir on a relapse rate over 2 years but found no evidence of an effect on disability.

Drugs to reduce progression[8, 16]

Current recommended treatment given under careful supervision:

- methotrexate 7.5 mg (o), on 1 day each week +
- folic acid 5 mg (o), 3 days after each methotrexate dose

Immunosuppression can also be given with cyclophosphamide or azathioprine.[17] The chemotherapeutic agent mitozantrane has been shown to arrest progression of MS but its use is limited by its cardiotoxicty.[12]

Treatment of symptoms[7]

Spasticity

- Physiotherapy
- Baclofen 10 - 25 mg (o) nocte
- For continuous drug therapy:
  - baclofen 5 mg (o) tds, increasing to 25 mg (o) tds +
  - diazepam 2 - 10 mg (o) tds

Paroxysmal (e.g. neuralgias)

- Carbamazepine or clonazepam

See references 8 and 16 for treatment of other symptoms.

ACUTE IDIOPATHIC DEMYELINATING POLYNEUROPATHY (GUILLAIN-BARRÉ SYNDROME)

Guillain-Barré syndrome is the best known of the peripheral neuropathies that have an acute onset, and it is potentially fatal. Early diagnosis of this serious disease by the family doctor is crucial as respiratory paralysis may lead to death. The underlying pathology is segmental demyelination of the peripheral nerves and nerve roots.

Clinical features

- Paraesthesia or pain in the limbs
- Weakness in the limbs (usually symmetrical)
- Both proximal and distal muscles affected

- Facial and bulbar paralysis
- Weakness of extraocular muscles (rarely)
- Reflexes depressed or absent
- Variable sensory loss but rare

Within 3 - 4 weeks the motor neuropathy, which is the main feature, progresses to a maximum disability, possibly with complete quadriparesis and respiratory paralysis.[18]

Investigations

- CSF protein is elevated; cells are usually normal.
- Motor nerve conduction studies are abnormal.

Management

- Admit to hospital.
- Respiratory function (vital capacity) should be measured regularly (2 - 4 hours at first).
- Tracheostomy and artificial ventilation may be necessary.
- Physiotherapy to prevent foot and wrist drop and other general care should be provided.
- Plasmapheresis is the mainstay of treatment.
- IV immunoglobulin (0.4 g/kg/day for 5 days) is also effective.[17]
- Corticosteroids are not generally recommended.

Outcome

About 80% of patients recover without significant disability. Approximately 5% relapse.[18]

MYASTHENIA GRAVIS

Myasthenia gravis (MG) is an acquired autoimmune disorder that usually affects muscle strength. Patients have fluctuating symptoms and variable distribution of muscle weakness. All degrees of severity, ranging from occasional mild ptosis to fulminant quadri-plegia and respiratory arrest, can occur.[19] See Table 33.7. It is associated with thymic tumour and other autoimmune diseases, e.g. RA, SLE.

Table 33.7 Clinical classification of acquired myasthenia gravis

Group I	Ocular MG
Group IIA	Mild generalised MG
Group IIB	Moderate to severe MG
Group III	Acute severe (fulminating) MG with respiratory muscle weakness
Group IV	Late (chronic) severe MG

Clinical features

- Painless fatigue with exercise
- Weakness also precipitated by emotional stress, pregnancy, infection, surgery
- Variable distribution of weakness:
  - ocular
    - ptosis (60%) and diplopia (Figure 33.6)
    - ocular myasthenia only remains in about 10%
  - bulbar: weakness of chewing, swallowing, speech (ask to count to 50), whistling and head lolling
  - limbs (proximal and distal)
  - generalised
  - respiratory: breathlessness, ventilatory failure



Figure 33.6 Myasthenia Gravis in a 40-year-old woman with a 12-month history of increasing muscular weakness including drooping of the eyelids. Ptosis, especially on the right side, is apparent

Note: The classic MG image is 'the thinker' - the hand used to hold the mouth closed and the head up.

Diagnostic tests

- ↑Serum anti-acetylcholine receptor antibodies
- Electrophysiological tests if antibody test negative
- CT scan and chest X-ray to detect thymoma
- Edrophonium test still useful

Management principles[8, 20]

- Detect possible presence of thymoma with CT or MR scan of thorax. If present, removal is recommended.
- Thymectomy is recommended early for generalised myasthenia, especially in all younger patients with hyperplasia of the thymus, even if not confirmed preoperatively.
- Plasmapheresis is useful for acute crisis or where temporary improvement is required or patients are resistant to treatment.
- Avoid drugs that are relatively contraindicated.
- Pharmacological agents:
    - anticholinesterase drugs (e.g. pyridostigmine, neostigmine or distigmine): should be used only for mild-to-moderate symptoms
    - corticosteroids: useful for all grades of MG; should be introduced slowly

Practice tips

- The combination of ocular and facial weakness should alert the family doctor to the possibility of a neuromuscular disorder, especially MG or mitochondrial myopathy.[20] Look for weakness and fatigue.
- Beware of facioscapulohumeral dystrophy.
- Ptosis may develop only after looking upwards for a minute or longer.
- Smiling may have a characteristic snarling quality.

PTOSIS

It is worth remembering that the four major causes of ptosis are:

1. 3rd cranial nerve palsy - ptosis, eye facing 'down and out', dilated pupil, sluggish light reflex
2. Horner's syndrome - ptosis, miosis (constricted pupil), ipsilateral loss of sweating
3. Mitochondrial myopathy - progressive external ophthalmoplegia or limb weakness, induced by activity
4. Myasthenia gravis - ptosis and diplopia

DYSTONIA

Definition

Dystonias are sustained or intermittent abnormal repetitive movements or postures resulting from alterations in muscle tone. The dystonic spasms may affect one (focal) or more (segmental) parts of the body or the whole body (generalised).

Key points

- Misdiagnosis is common as transient symptoms may be mistaken for an emotional or psychiatric disorder. Many cases take years to diagnose.
- Dystonias are often regarded as nervous tics.
- The cause is thought to be disorders of the basal ganglia of the brain, but mainly there is no known specific cause.
- Neuroleptic and dopamine receptor blocking agents (e.g. levodopa, metoclopramide) can induce a severe generalised dystonia (e.g. oculogyric crisis) which is treated with benztropine 1 - 2 mg IM or IV.[20]

Focal dystonias

- *Blepharospasm* is a focal dystonia of the muscles around the eye resulting in uncontrolled blinking, especially in bright light.
- *Oromandibular dystonia* affects the jaw, tongue and mouth, resulting in jaw grinding movements and grimacing. Proper speech and swallowing may be disrupted.
- *Meige's syndrome* is a combination of blepharospasm and oromandibular dystonia.

Note: It must be differentiated from the buccallingual-facial movements of tardive dyskinesia.

- *Hemifacial spasm* involves involuntary, irregular muscle contractions and spasms affecting one side of the face. It usually starts with twitching around the eye and then spreads to involve all the facial muscles on one side. It is usually due to irritation of the facial nerve in its intracranial course and surgical intervention may alleviate this problem.
- *Writer's cramp, typist's cramp, pianist's cramp, golfer's cramp* are all occupational focal dystonias of the hand and/or forearm initiated by performing these skilled acts.
- *Cervical dystonia or spasmodic torticollis* is a focal dystonia of the unilateral cervical muscles. It usually begins with a pulling sensation followed by twisting or jerking of the head, leading to deviation of the head and neck to one side. In early stages patients can voluntarily overcome the dystonia.
- *Laryngeal or spastic dystonia* is a focal dystonia of the laryngeal muscles resulting in a strained, hoarse or creaking voice. It may lead to inability to speak in more than a whisper.

Treatment of focal dystonias

The current treatment for focal or segmental (spread to an adjacent body region) dystonias is localised injection of purified botulinum A toxin into the affected muscle groups. The dosage is highly individualised and needs to be repeated at intervals of 3 and 6 months. The injections have to be given with great caution.

TICS

Motor and vocal tics are a feature of Tourette's disorder. If socially disabling treat with:

- haloperidol 0.25 mg (o) nocte, very gradually increasing to 2 g (max.) daily[7] or
- clonidine 25 μg (o) bd for 2 weeks, then 50 - 75 μg bd

FACIAL NERVE (BELL'S) PALSY[8, 15]

Facial (7th nerve palsy), which is an acute unilateral lower motor neurone paresis or paralysis, is the commonest cranial neuropathy. The classic type is Bell's palsy, which is usually idiopathic although attributed to an inflammatory swelling involving the facial nerve in the bony facial canal. In the Ramsay-Hunt syndrome, which is due to infection with herpes zoster causing facial nerve palsy, vesicles may be seen on the ipsilateralear.

Associations:

- herpes simplex virus (postulated)
- diabetes mellitus
- hypertension

- thyroid disorder, e.g. hyperthyroidism

Features

- abrupt onset (can worsen over 2 - 5 days)
- weakness in the face (complete or incomplete)
- preceding pain in or behind the ear
- impaired blinking

Less common:

- difficulty eating
- loss of taste - anterior two-thirds of tongue
- hyperacusis

Management

- prednisolone 50 mg (o) daily in divided doses for 4 days then taper to zero over next 7 days (start within 3 days of onset)

Note: This is controversial as evidence is not convincing[15] (better in severer cases).

- patient education and reassurance
- adhesive patch or tape over eye if corneal exposure (e.g. windy or dusty conditions, during sleep)
- artificial tears if eye is dry and at bedtime
- massage and facial exercises during recovery

Note:

- At least 70 - 80% achieve full spontaneous recovery; higher if mild.
- Electromyography and nerve excitability or conduction studies are a prognostic guide only.
- No evidence that nucleoside analogue, e.g. aciclovir, is useful but should be used for Ramsay-Hunt syndrome.
- No evidence that surgical procedures to decompress the nerve are beneficial.[15]

Diagnostic triads of neurological dilemmas

All triads show a chronic onset unless indicated by an asterisk (acute onset).

If you see this combination of signs:	Consider:
Charcot's triad:	
• dysarthria + intention tremor + nystagmus	° cerebellar disease (typical of MS)
• visual disturbance (blurred or transient loss) + weakness in limbs ±paraesthesia in limbs	° multiple sclerosis
Note: here are many combinations of MS (Charcot's has historical interest).	
• rigidity + bradykinesia + resting tremor	° Parkinson's disease
• tremor (postural or action) + head tremor + absence of Parkinsonian features	° essential tremor
• fatiguable and weakness of eyelids and eye movements + limbs + bulbar muscles (speech and swallowing)	° myasthenia gravis
• weakness of limbs + of face + areflexia *	° Guillain-Barré syndrome (GBS)
• (episodic) vertigo + tinnitus + hearing loss *	° Meniere's syndrome
• dementia + myoclonus + ataxia	° Creutzfeld-Jakob disease
• drowsiness + vomiting + headache (waking)	° ↑intracerebral pressure
• enophthalmos + meiosis + ptosis ±anhydrosis	° Horner's syndrome
• blank spell + lip-smacking (or similar automation) + olfactory/gustatory hallucination	° complex partial seizure
• gradual spread (Jacksonian march) of focal jerking (mouth, arm or leg) or sensory disturbance or (rarely) visual field disturbance	° simple partial seizure
• ↑intracranial pressure +/or focal signs +/or epilepsy	° cerebral tumour
• dysphagia + dysphonia/dysarthria + spastic tongue	° pseudobulbar palsy

• recurrent: headache (often unilateral) + nausea (±vomiting) + visualaura[*]	° migraine with aura (formerly 'classical migraine')
• recurrent: severe retro-orbital headache + rhinorrhoea + lacrimation[*]	° cluster headache
• instantaneous: headache ±vomiting ± neck stiffness	° subarachnoid haemorrhage until disproven
• headache + visual obscurations + papilloedema (often in obese young female)	° benign intracranial hypertension
• acute and transient: amaurosis fugax or dysphasia or hemiplegia[*]	° TIA (carotid)
• typical facies (temporalis atropy and frontal balding) + muscle weakness (±myotonia) + cataracts	° myotonic dystrophy
• ataxia + ophthalmoplegia + areflexia[*]	° Miller-Fisher variant of GBS
• vertigo + provoked by movement (especially rolling in bed) + Hall pike test + ve	° BPPV
• UMN signs + LMN signs + fasciculations	° motor neurone disease

REFERENCES

1. Iansek R. *Pitfalls in Neurology*. Melbourne: Proceedings of Monash University Medical School Update Course, 1992: 40 - 4.
2. Talley NJ, O'Connor S. *Clinical Examination* (5th edn). Sydney: Churchill Livingstone, 2005: 345 - 6.
3. Morris JGL. Essential tremor. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 179 - 80.
4. Iansek R. Parkinson's disease and dementia. Proceedings of Monash University Medical School Update Course, 2005: 1 - 23.
5. Selby G, Herkes G. Parkinson's disease. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 395 - 8.
6. Fung V, Hely M, Morris J. How to treat: Parkinson's disease and Parkinsonian syndromes. Australian Doctor, 2003; 14 November: 33 - 40.
7. Barton S et al. *Clinical Evidence* (Issue 5). London: BMJ Publishing Group, 2001: 906 - 13.
8. Tiller J (Chair). *Therapeutic Guidelines: Neurology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2002.
9. Fung VS, Hely MA et al. Drugs for Parkinson's disease. Aust Prescriber, 2001; 24: 92 - 5.
10. Selby G. Stereotactic surgery. In: Koller WC ed. *Handbook of Parkinson's Disease*. New York: Marcel Dekker Inc., 1987: 421 - 35.
11. McLeod JR. Multiple sclerosis. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 321 - 3.
12. Broadley S. Multiple sclerosis: update. Medical Observer, 2004; 16 October: 27 - 30.
13. Matthews B. Multiple sclerosis. Med Int, 1988; 48: 1961 - 7.
14. Burnham JA et al. The effect of high-dose steroids on MRI gadolinium enhancement in acute demyelinating lesions. Neurology, 1993; 43: 655 - 61.
15. Barton S et al. *Clinical Evidence* (Issue 5). London: BMJ Publishing Group, 2001: 894 - 904.
16. Jacobs L et al. Advances in specific therapy for multiple sclerosis. Curr Opin Neurol, 1994; 7: 250 - 4.
17. Milanese C et al. A double blind study on azathioprine in the treatment of multiple sclerosis. J Neurology, 1993; 240: 295 - 8.
18. Pollard J. Neuropathy, peripheral. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 346.
19. McLeod JG. Peripheral neuropathy. Med Int, 1988; 48: 1980 - 4.
20. Darveniza P. Myasthenia gravis. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 324 - 6.

CHAPTER 34. ABDOMINAL PAIN

> A great fit of the stone in my left kidney: all day I could do but three or four drops of water, but I drunk a draught of white wine and salet oyle, and after that, crabs' eyes in powder with the bone in the carp's head and then drunk two great draughts of ale with buttered cake; and I voyded with an hour much water and a stone as big as an Alexander seed. God by thanked!
>
> JOHN DEE 1594

ABDOMINAL PAIN

Abdominal pain represents one of the top 15 presenting symptoms in primary care[1] andvaries from a self-limiting problem to a life-threatening illness requiring immediate surgical intervention. Abdominal pain can be considered to be acute, subacute, chronic or recurrent. It can embrace all specialties, including surgery, medicine, gynaecology, geriatrics and psychiatry. For acute abdominal conditions it is important to make a rapid diagnosis in order to reduce morbidity and mortality. Most casesrequire surgical referral (Table 34.1). Lower abdominal pain in women adds another dimension tothe problem and will be presented in a separate chapter.

Table 34.1 Surgical causes of the acute abdomen

Process	Organ involved	Disorder
Inflammation	Bowel	Inflammatory bowel disease
	Appendix	Appendicitis
	Gall bladder	Cholecystitis
	Pancreas	Pancreatitis
	Fallopian tube	Salpingitis
	Colonic diverticulae	Diverticulitis
Perforation	Duodenum	Perforated duodenal ulcer
	Stomach	Perforated gastric ulcer
	Colon (diverticula or carcinoma)	Faecal peritonitis
	Gall bladder	Biliary peritonitis
	Appendix	Appendicitis
Obstruction	Gall bladder	Biliary colic
	Small intestine	Acute small bowel obstruction
	Large bowel	Acute large bowel obstruction
	Ureter	Ureteric colic
	Urethra	Acute urinary retention
	Mesenteric artery occlusion	Intestinal infarction
Haemorrhage	Fallopian tube	Ruptured ectopic pregnancy
	Spleen or liver	Ruptured spleen or liver (haemoperitoneum)
	Ovary	Ruptured ovarian cyst
	Abdominal aorta	Ruptured AAA
Torsion (ischaemia)	Sigmoid colon	Sigmoid volvulus
	Ovary	Torsion ovarian cyst
	Testes	Torsion of testes

KEY FACTS AND CHECKPOINTS

- The commonest causes of the acute abdomen in two general practice series were: Series 1 - acute appendicitis (31%) and the colics (29%);[2] Series 2 - acute appendicitis (21%), the colics (16%), mesenteric adenitis(16%).[3] The latter study included children.
- An international study involving referral to 26 seugecal departments 17 wuntnis weald nou-speufic obdoaunel paiw (34%), acule offeutecfes (28%)cholecystitis (10%) as the most common condefeous.[1]
- As a general rule upper abdominal pain is caused by lesions of the upper GIT and lower abdominal pain by lesions of the lower GIT.
- Colicky midline umbilical abdominal pain (severe) → vomiting → distension ° small bowel obstruction (SBO).

- Midline lower abdominal pain → distension → vomiting ° large bowel obstruction (LBO).
- If cases of acute abdomen have a surgical cause, the pain nearly always precedes the vomiting.
- Mesenteric artery occlusion must be considered in an elderly person with arteriosclerotic disease or in patients with atrial fibrillation presenting with severe abdominal pain or following myocardial infarction.
- Up to one-third of presentations of abdominal pain are considered to be non-specific, whereby no specific cause is found.

A DIAGNOSTIC APPROACH

A summary of the separate diagnostic models for acute abdominal pain and chronic abdominal pain are presented in Tables 34.2 and 34.3.

Table 34.2 Acute abdominal pain (adults): diagnostic strategy model (excluding trauma)

Q. Probability diagnosis
A. Acute gastroenteritis
Acute appendicitis
Mittelschmerz/dysmenorrhoea
Irritable bowel syndrome
Q. Serious disorders not to be missed
A. Cardiovascular
myocardial infarction
ruptured AAA
dissecting aneurysm aorta
mesenteric artery occlusion
Neoplasia
large or small bowel obstruction
Severe infections
acute salpingitis
peritonitis
ascending cholangitis
intra-abdominal abscess
pancreatitis
Ectopic pregnancy
Small bowel obstruction
Sigmoid volvulus
Perforated viscus
duodenal ulcer
colonic diverticulum
Meckel's diverticulum
colonic cancer
Q. Pitfalls (often missed)
A. Acute appendicitis
Myofascial tear
Pulmonary causes
pneumonia
pulmonary embolism
Faecal impaction (elderly)
Herpes zoster
Rarities
porphyria

lead poisoning haemochromatosis sickle cell anaemia tabes dorsalis Haemoglobinuria Addisons disease
Q. Seven masquerades checklist A. Depression Diabetes Ketoacidosis Drugs Anaemia sickle cell Thyroid disorder - Spinal dysfunction UTI including urosepsiis
Q. Is the patient trying to tell me something? A. May be very significant. Consider Munchausen's syndrome, sexual dysfunction and abnormal stress.

Table 34.3 Chronic or recurrent abdominal pain (adult): diagnostic strategy model

Q. Probability diagnosis
A. Irritable bowel syndrome Mittelschmerz/dysmenorrhoea Peptic ulcer/gastritis **Q. Serious disorders not to be missed**
A. Cardiovascular • mesenteric artery ischaemia • AAA Neoplasia • carcinoma bowel/stomach • carcinoma pancreas • ovarian tumours Severe infections • hepatitis • recurrent PID **Q. Pitfalls (often missed)** A. Food allergies Lactase deficiency Constipation Chronic pancreatitis Crohn's disease Endometriosis Diverticulitis Rarities • uraemia • lead poisoning • Crohn's disease

- porphyria
- sickle cell anaemia
- Addison's syndrome

Q. Seven masquerades checklist

A. Depression

Diabetes -

Drugs

Anaemia -

Thyroid disorder -

Spinal dysfunction

UTI

Q. Is the patient trying to tell me something?

A. Astrong possibility: consider hypochondriasis, anxiety, sexual dysfunction.

Probability diagnosis

The most common causes of acute abdomen are acute appendicitis, acutegastroenteritis, an irritable bowel syndrome, the various ‘colics’ and ovulation pain (mittelschmerz). Mesentericadenitis is common in children. The various causes of chronic or recurrent abdominal pain are presented in Table 34.3. A study on chronic abdominal pain[4] showed that the commonestreasons (approximate percentages) were: no discoverable causes (50%), minor causes including muscle strains (16%), irritable bowel syndrome (12%), gynaecological causes (8%), peptic ulcers and hiatus hernia (8%).

Serious disorders not to be missed

Most of the causes of the acute abdomen are serious and early diagnosis is mandatory to reduce mortality and morbidity.

It is vital not to misdiagnose a ruptured ectopic pregnancy, which causes lower abdominal or suprapubic pain of sudden onset, or the life-threatening vascular causes, such as a ruptured or dissecting aortic aneurysm, mesenteric artery occlusion and myocardial infarction (which can present as epigastric pain).

Perforated ulcers and strangulated bowel, such as volvulus of the sigmoid and entrapment of the small bowel in a hernial orifice or around adhesions, also demand an early diagnosis.

There are some important ‘red flag’ symp-toms and signs[1] of abdominal emergencies demanding urgent attention:

- collapsing at toilet (points to intra-abdominal bleeding)
- light-headedness
- progressive intractable vomiting
- progressive abdominal distension
- progressive intensity of pain
- prostration

Signs

- Pallor and sweating
- Hypotension
- Atrial fibrillation or tachycardia
- Fever
- Rebound tenderness and guarding
- Decreased urine output

Dangers of misdiagnosis

- Ectopic pregnancy → rapid hypovolaemic shock
- Ruptured abdominal aortic aneurysm (AAA) → rapid hypovolaemic shock
- Gangrenous appendix → peritonitis/pelvic abscess
- Perforated ulcer → peritonitis
- Obstructed bowel → gangrene

Pitfalls

A very common pitfall is misdiagnosing acute appendicitis, especially in the elderly, in children, in pregnancy and in those taking steroids, where the presentation may be atypical. Early appendicitis presents typically with central abdominal pain that shifts to the right iliac fossa (RIF) some 4 to 6 hours later. This causes confusion early on. It can cause diarrhoea with abdominal pain, especially if a pelvic appendix, and can be misdiagnosed as acute gastroenteritis.

Disaccharidase deficiencies, such as lactase deficiency, are associated with cramping abdominal pain, which may be severe. The pain follows some time, maybe hours, after the ingestion of milk and is accompanied by the passage of watery stool. The association with milk may be unrecognised by the patient.

Herpes zoster, especially in the elderly patient with unilateral abdominal pain in the dermatomal distribution, is a trap. Referred pain from conditions above the diaphragm, such as myocardial infarction, pulmonary embolism and pneumonia, can be misleading. The rare general medical causes, such as diabetes ketoacidosis, acute porphyria, Addison's syndrome, lead poisoning, tabes dorsalis, sickle cell anaemia, haemochromatosis and uraemia often create a diagnostic dilemma and should be kept in mind.

Specific pitfalls

- Misdiagnosing a ruptured ectopic pregnancy in the patient on contraception or with a history of normal menstruation or where the brownish vaginal discharge is mistaken for a normal period.
- Failing to examine hernial orifices in a patient with intestinal obstruction.
- Misleading temporary improvement (easing of pain) in perforation of gangrenous appendix or perforated peptic ulcer.
- Overlooking a perforation in the elderly or in patients taking corticosteroids, because of relative lack of pain.
- Overlooking acute mesenteric artery obstruction in an elderly patient with colicky central abdominal pain.
- Attributing abdominal pain, frequency and dysuria to a urinary infection when the cause could be diverticulitis, pelvic appendicitis, salpingitis or a ruptured ectopic pregnancy.

Seven masquerades checklist

Depression, diabetes, drugs, spinal dysfunction and UTI can all cause abdominal pain although the pain may be more subacute or chronic. Abdominal pain and even tenderness can accompany diabetic ketoacidosis. Drugs that can cause abdominal pain are listed in Table 34.4.

Table 34.4 Drugs to consider as a cause of abdominal pain

Alcohol
Antibiotics (e.g. erythromycin)
Aspirin
Corticosteroids
Cytotoxic agents
Tricyclic antidepressants (e.g. imipramine)
Iron preparations
Nicotine
NSAIDs/cox$_2$ inhibitors
Sodium valproate
Phenytoin

Spinal dysfunction of the lower thoracic spine and thoracolumbar junction can cause referred pain to the abdomen (Figure 34.1). The pain is invariably unilateral, radicular in distribution, and related to activity. It can be confused with intra-abdominal problems such as biliary disease (right-sided), appendicitis and Crohn's disease (right side), diverticular disorder (left-sided) and pyelonephritis.



Figure 34.1 Referred pain patterns to the anterior abdominal wall from dysfunction of the thoracic spine via anterior cutaneous branches of the nerve roots (right side). Note the possibility of confusion with biliary pain, appendicitis and inguinal disorders

Psychogenic considerations

Psychogenic factors can be most relevant, especially in recurrent or chronic abdominal pain where no specific cause can be identified in most cases.[5] Bain and Spaulding found that 40% of cases of adult abdominal pain had non-structural causes with 28% having psychiatric diagnoses and 6% spastic colon. They noted that ‘psychological disturbances are often fairly easy to identify if care is taken to obtain the personal history and to assess the patient's personality, but the diagnostic terms used to describe psychological disturbances lack precision'.

THE CLINICAL APPROACH

History

The urgency of the history will depend on the manner of presentation, whether acute or chronic. Pain has to be analysed according to its quality, quantity, site and radiation, onset, duration and offset, aggravating and relieving factors and associated symptoms and signs.

Special attention has to be paid to:

- anorexia, nausea or vomiting
- micturition
- bowel function
- menstruation

- drug intake

Key questions

Point to where the pain is and where it travels to. Questions to ask:

- What type of pain is it: is it constant or does it come and go?
- How severe would you rate it from 1 to 10?
- Have you ever had previous attacks of similar pain?
- What else do you notice when you have the pain?
- Do you know of anything that will bring on the pain? Or relieve it?
- What effect does milk, food or antacids have on the pain?
- Have you noticed any sweats or chills or burning of urine?
- Are your bowels behaving normally? Have you been constipated or had diarrhoea or blood in your motions?
- Have you noticed anything different about your urine?
- What medications do you take?
- How much aspirin do you take?
- Are you smoking heavily or taking heroin or cocaine?
- How much alcohol do you drink?
- How much milk do you drink?
- Have you travelled recently?
- What is happening with your periods? Is it mid-cycle or are your periods overdue?
- Does anyone in your family have bouts of abdominal pain?
- Do you have a hernia?
- What operations have you had for your abdomen?
- Have you had your appendix removed?

Physical examination

A useful checklist for conducting the examination is:

- general appearance
- oral cavity
- vital parameters: temperature, pulse, BP, respiratory rate (record these in the notes)
- chest: check heart and lungs for upper abdominal pain (especially if absent abdominal signs)
- abdomen: inspection, palpation, percussion and auscultation.

The abdominal examination should be performed with the patient lying flat with one pillow under the head and the abdomen uncovered from xiphisternum to groin. Ask the patient to breathe through the mouth during the examination. Consider the following:

- inguinal region (including hernial orifices) and femoral arteries
- rectal examination: mandatory
- vaginal examination (females): for suspected problems of the fallopian tubes, uterus or ovaries
- thoracolumbar spine (if referred spinal pain suspected)
- urine analysis: white cells, red cells, glucose and ketones, porphyrins
- special clinical tests
  - Murphy's sign (a sign of peritoneal tenderness with acute cholecystitis)
  - iliopsoas and obturator signs

Guidelines

- *Palpation*: palpate with gentleness - note any guarding or rebound tenderness
  - guarding indicates peritonitis
  - rebound tenderness indicates peritoneal irritation (bacterial peritonitis, blood). Feel for maximum site that corresponds to focus of the problem
- Patient pain indicator: the finger pointing sign indicates focal peritoneal irritation; the spread palm sign indicates visceral pain
- Atrial fibrillation: consider mesenteric artery obstruction
- Tachycardia: sepsis and volume depletion
- Tachypnoea: sepsis, pneumonia, acidosis
- Pallor and ‘shock’: acute blood loss
- *Auscultation*:note bowel activity or a sucussion splasn

causes of a ‘silent abdormen‘:

- Diffuse sepsis
- Ileus
- Mechancial dostrution (advancee)

Hypertympany inducates mehanical obstructure.

Physical signs may be reduced in the elderly, grossly obese, severely ill and patients taking corticosteroid therapy.

Investigations

The following investigations may be selected:

- haemoglobin - anaemia with chronic blood loss (e.g. peptic ulcer, carcinoma, oesophagitis)
- blood film - abnormal red cells with sickle cell disease
- WCC - leucocytosis with appendicitis (75%),[2] acute pancreatitis, mesenteric adenitis (first day only), cholecystitis (especially with empyema), pyelonephritis
- ESR - raised with carcinoma, Crohn's disease, abscess, but non-specific
- C-reactive protein (CRP) - use in diagnosing and momtori nginfection, inflammation (e.g. pancreatic) preferable to ESA
- liver function tests - hepatobiliary disorder
- serum amylase - and lipase & preterable if raised to greater than five times normal upper level acute pancreatitis is most likely; also raised partially with most intraabdominal disasters (e.g. ruptured ectopic, perforated peptic ulcers, ruptured empyema of gall bladder, ruptured aortic aneurysm)
- pregnancy tests - urine and serum β-HCG: for suspected ectopic
- urine:
    - blood: ureteric colic (stone or blood clot), urinary infection
    - white cells: urinary infection, appendicitis (bladder irritation)
    - bile pigments: gall bladder disease
    - porphobilinogen: porphyria (add Ehrlich's aldehyde reagent)
    - ketones: diabetic ketoacidosis
    - air (pneumaturia): fistula (e.g. diverticulitis, other pelvic abscess, pelvic carcinoma)
- faecal blood - mesenteric artery occlusion, intussusception ('redcurrant jelly'), carcinoma colon, diverticulitis, Crohn's disease and ulcerative colitis

Radiology

The following tests can be considered according to the clinical presentation:

- plain X-ray abdomen (erect and supine): look for (Figure 34.2):


free gas under diaphragm
gallstones or kidney stones
large oval gas shadow of sigmoid volvulus
ureteric calculi
small bowel obstruction with ladder pattern of fluid levels

Figure 34.2 The acute abdomen: signs to watch for on plain abdominal X-ray

  - renal/uteric stones - 70% opaque[2]
  - biliary stones - only 10 - 30% opaque
  - air in biliary tree
  - calcified aortic aneurysm
  - marked distension sigmoid → sigmoid volvulus
  - distended bowel with fluid level → bowel obstruction
  - enlarged caecum with large bowel obstruction
  - blurred right psoas shadow → appendicitis
  - a senital loop of gas in left upper quadrant (LUQ) → acute pancreatitis
- chest X-ray: air under diaphragm → perforated ulcer
- ultrasound: good for hepatobiliary system, kidneys and female pelvis: look for
  - gallstones
  - ectopic pregnancy
  - pancreatic pseudocyst
  - aneurysm aorta/dissecting aneurysm
  - hepatic metastases and abdominal tumours
  - thickened appendix
  - paracolic collection
- IVP
- contrast - enhanced X-rays (e.g. Gastrografin meal): diagnosis of bowel leakage
- barium enema
- HIDA or DIDA nuclear scan - diagnosis of acute cholecystitis
- CT scan: gives excellent survey of abdominal organs including masses and fluid collection
- ERCP: shows bile duct obstruction and pancreatic disease
- MRI scan

Other tests:

- ECG

- endoscopy upper GIT
- sigmoidoscopy and colonoscopy

DIAGNOSTIC GUIDELINES

General rules

- Upper abdominal pain is caused by lesions of the upper GIT.
- Lower abdominal pain is caused by lesions of the lower GIT or pelvic organs.
- Early severe vomiting indicates a high obstruction of the GIT.
- Acute appendicitis features a characteristic ‘march’ of symptoms: pain ? anorexia nausea → vomiting.

Pain patterns

The pain patterns are presented in Figure 34.3. Colicky pain is a rhythmic pain with regular spasms of recurring pain building to a climax and fading. It is virtually pathognomonic of intestinal obstruction. Ureteric colic is a true colicky abdominal pain, but so-called biliary colic and renal colic are not true colics at all.



Figure 34.3 Characteristic pain patterns for various causes of ‘colicky’ acute abdominal pain

Site of pain

Typical pain sites of abdominal pain (general guidelines only) are presented in Figures 34.4 and 34.5. Epigastric pain usually arises from disorders of the embryologic foregut, such as the oesophagus, stomach and duodenum, hepatobiliary structures, pancreas and spleen. However, as some disorders progress the pain tends to shift from the midline to the right (gall bladder and liver) or left (spleen). Periumbilical pain usually arises from disorders of structures of the embryologic midgut, while structures from the hindgut tend to refer pain to the lower abdomen or suprapubic region.


Organs
Conditions
Right hypochondrium
gall bladder
biliary tract
liver
biliary colic
cholecystitis
ascending cholangitis
Periumbilical (central)
jejenum/ileum
appendix
aorta
small bowel obstruction
appendicitis (early)
ruptured AAA
mesenteric artery occlusion
RIF
appendix
caecum/ileum
appendicitis
Crohn's disease
Meckel's diverticulitis
Groin
torsion of testis
check hernial orifices with SBO
Organs
Conditions
Epigastric
oeophagus
stomach
duodenum
gall bladder
liver
pancreas
perforated ulcer
cholecystitis
pancreatitis
myocardial infarction (referred)
lower lobe pneumonia (referred)
L hypochondrium and loin
kidney
spleen
kidney colic
urinary infection
splenic infarct
Lower abdomen
large bowel
large bowel obstruction
LIF
descending colon
sigmoid
diverticulitis
Suprapubic (hypogastric)
ovaries
fallopian tubes
bladder
ruptured ectopic
acute PID
ruptured ovarian cyst
torsion of ovary
NB lower abdominal and suprapubic pain overlap

Figure 34.4 Typical sites of various causes of acute abdominal pain


chronic cholecystitis
biliary colic
hepatitis
hepatic congestion
mesenteric artery ischaemia
recurrent appendicitis
Crohn's disease
peptic ulcer
oesophagitis/reflux
carcinoma stomach
chronic pancreatitis
irritable bowel syndrome
diverticulitis
constipation
gynaecological disorders
• PID
• mittelschmerz
• ovarian tumours

Figure 34.5 Typical sites of various causes of chronic or recurrent abdominal pain

The intra-abdominal sensory receptors can be considered as innervating visceral or parietal peritoneum. Visceral mechano-receptors are triggered by intestinal distension or tension on mesentry or blood vessels while nociceptors are triggered by mechanical, thermal and chemical stimuli. The pain from viscera is felt as diffuse and poorly localised while stimulation of parietal peritoneal nociceptors gives a pain that is experienced directly at the site of insult.

ABDOMINAL PAIN IN CHILDREN

Abdominal pain is a common complaint in children, especially recurrent abdominal pain, which is one of the most common complaints in childhood. The problem causes considerable anxiety in parents and it is important to differentiate the severe problems demanding surgery from non-surgical problems.

Acute abdominal pain

The causes of abdominal pain can be considered in the diagnostic model category.

1. Common causes/probability diagnosis
    - infantile colic
    - gastroenteritis (all ages)
    - mesenteric adenitis
2. Serious causes, not to be missed
    - intussusception (peaks at 6 - 9 months)
    - acute appendicitis (mainly 5 - 15 years)
    - bowel obstruction
3. Pitfalls
    - Child abuse
    - constipation
    - torsion of testes
    - lactose intolerance
    - peptic ulcer
    - infections
        - mumps
        - tonsillitis
        - pneumonia (especially right lower lobe)
        - EBM
        - UTI
4. Rarities
    - Meckel's diverticulitis
    - Henoch-Schönlein purpura
    - sickle crisis
    - lead poisoning
5. Seven masquerades checklist
    - diabetes mellitus
    - drugs
    - UTI
6. 5. Psychogenic consideration
    - important cause

Infantile colic

This is the occurrence in a well baby of regular, unexplained periods of inconsolable crying and fretfulness, usually in the late afternoon and evening, especially between 2 weeks and 16 weeks of age. No cause for the abdominal pain can be found and it lasts for a period of at least 3 weeks. It is very common and occurs in about one-third of infants.

Typical features

- Baby between 2 and 16 weeks old
- Prolonged crying - at least 3 hours
- Crying worst at around 10 weeks of age
- Crying during late afternoon and early evening
- Occurrence at least 3 days a week
- Child flexing legs and clenching fists because of the ‘stomach ache’
- Normal physical examination

Management

Reassurance and explanation to the parents. Advice for the parents:

- Use gentleness (such as subdued lighting where the baby is handled, soft music, speaking softly, quiet feeding times).
- Avoid quick movements that may startle the baby.
- Make sure the baby is not hungry - underfeeding can make the baby hungry.

- If the baby is breastfed, express the watery foremilk before putting the baby to the breast.
- Provide demand feeding (in time and amount).
- Make sure the baby is burped, and give posture feeding.
- Provide comfort from a dummy or pacifier.
- Provide plenty of gentle physical contact.
- Cuddle and carry the baby around (e.g. take a walk around the block).
- A carrying device such as ‘snuggly’ or ‘Meh Tai Sling’ allows the baby to be carried around at the time of crying.
- Make sure the mother gets plenty of rest during this difficult period.
- Do not worry about leaving a crying child for 10 minutes or so after 15 minutes of trying consolation.

Medication

Drugs are not generally recommended, but for very severe problems some preparations can be very helpful (e.g. simethicone and hyoscine (Donnalix) preparations).

Intussusception

Intussusception is the diagnosis that should be foremost in one's mind with a child aged between 3 months and 2 years presenting with sudden onset of severe colicky abdominal pain, coming at intervals of about 15 minutes and lasting 2 - 3 minutes. Early diagnosis, within 24 hours of the onset, is essential, for after this time there is a significant rise in morbidity and mortality. It is due to telescoping of the segment of bowel into the adjoining distal segment (e.g. ileocaecal segment), resulting in intestinal obstruction.

Typical clinical features[6]

See Figure 34.6.



Figure 34.6 Typical features with pain distribution of acute intussusception

- Male babies > female
- Age 6 - 12 months
- Range: birth to school age, usually 5 - 24 months
- Sudden-onset acute pain with shrill cry
- Vomiting
- Leth apgy
- Pallor with attacks

- Intestinal bleeding: redcurrant jelly (60%)[7]

DxT: pale child + severe 'colic' + vomiting ° acute intussusception

Physical signs

- Pale, anxious and unwell
- Sausage-shaped mass in right upper quadrant (RUQ), any where behaveen the line of wlon and umbelecus, especially during attacks (difficult)
- Signe de dance (i.e. emptiness in RIF to palpation)
- Alternating high-pitched active bowel sounds with absent sounds
- Rectal examination: ± blood

Diagnosis

- Ultrasound
- Air or barium enema also used for diagnosis and treatment

Treatment[7]

- Hydrostatic reduction by airor oxygen from the will supply (preferred) or barium enema
- Surgical intervention may be necessary

Differential diagnosis

- Acute gastroenteritis: can be difficult in those cases where there is some loose stool with intussusception and with blood and mucus without much watery stool in gastroenteritis. However, usually attacks of pain are of shorter duration, and there is loose watery stool, fever and no abdominal mass. If doubtful refer as possible intussusception.
- Impacted faeces can lead to spasms of colicky abdominal pain - usually an older child with a history of constipation.
- Other causes of intestinal obstruction (e.g. irreducible inguinal hernia, volvulus, intra-abdominal band).

Drugs

In any child complaining of acute abdominal pain, enquiry should be made into drug ingestion. A common cause of colicky abdominal pain in children is cigarette smoking (nicotine); consider other drugs such as marijuana, cocaine and heroin.

Acute appendicitis in children

This may occur at any age, being more common in children of school age and in adolescence, and uncommon under 3 years of age. Special problems of early diagnosis occur with the very young (less than 3 years) and in intellectually disabled children, many of whom present with peritonitis.

Vomiting occurs in at least 80% of children with appendicitis and diarrhoea in about 20%. The temperature is usually only slightly elevated but in about 5% of cases it exceeds 39°C.[2]

In children the physical examination, especially eliciting abdominal (including re-bound) tenderness, and the rectal examination demand considerable tact, patience and gentleness. Jumping or hopping induces pain.

A serious point of confusion can occur between pelvic appendicitis, causing diarrhoea and vomiting, and acute gastroenteritis. A particularly severe case of apparent gastroenteritis, especially if persistent, should be regarded as pelvic appendicitis until proved otherwise.

Mesenteric adenitis

This presents a difficult problem in differential diagnosis with acute appendicitis because the history can be very similar. At times the distinction may be almost impossible. In general, with mesenteric adenitis localisation of pain and tenderness is not as definite, rigidity is less of a feature, the temperature is higher, and anorexia, nausea and vomiting are also lesser features. The illness lasts about five days followed by a rapid recovery. Comparisons between the two are presented in Table 34.5 but if in any doubt it is advisable to consider the problem as acute appendicitis and perhaps proceed to laparotomy.[7]

Table 34.5 Comparison of the features of acute appendicitis and mesenteric adenitis in children (guidelines only)

	Acute appendicitis	Mesenteric adenitis
Typical child	Older	Younger
Site of onset of pain	Midline Shifting to right	RIF Can be midline

Preceding respiratory illness	Uncommon	Invariable: URTI or tonsillitis
Anorexia, nausea, vomiting	+ +	±
Colour	Usually pale	Flushed: malar flush
Temperature	N or ↑	↑↑ → ↑↑↑
Abdominal palpation	Tender in RIF Guarding ° Rigidity	Tender in RIF Minimal guarding Usually no rigidity
Rectal examination	Invariably tender	Often tender but lesser degree
Psoas and obturator tests	Usually positive	Usually negative

Mesenteric adenitis can sometimes present an anaesthetic risk and patients are usually quite ill in the immediate postoperative period.[7]

Recurrent abdominal pain

Recurrent abdominal pain (RAP) - three distinct episodes of abdominal pain over 3 or more months - occurs in 10% of school-aged children. In only 5 - 10% of children will an organic cause be found so that in most the cause remains obscure.[7, 8]

Organic causes

An organic cause, however, must be considered and excluded. Organic disease is more likely if:

- the pain is other than periumbilical
- the pain radiates rather than remains localised
- the pain wakens the child from sleep
- the pain is accompanied by vomiting
- the child is not completely well between attacks
- there is associated weight loss
- there is failure to thrive

Possible causes

- Constipation
- Childhood migraine equivalent (pain with extreme pallor)
- Lactose intolerance (symptoms related to milk ingestion)
- Intestinal parasites (may disturb child about 60 minutes after falling asleep)

Investigations

- Urine analysis and MSU
- FBE and ESR
- Plain X-ray (assesses faecal retention)

Non-organic RAP

Typical clinical features:

- acute and frequent colicky abdominal pain
- pain localised to or just above umbilicus
- no radiation of pain
- pain lasts less than 60 minutes
- nausea frequent and vomiting rare
- diurnal (never wakes the child at night)
- minimal umbilical tenderness
- anxious child
- obsessive or perfectionist personality
- one or both parents intense about child's health and progress

Psychogenic factors

Although psychogenic factors are very relevant in individual cases there is scant hard evidence to support the widely held hypothesis[8] that such factors account for the vast majority of RAP. Some children will have obvious psychological problems or even be school avoidant, a common factor being family disruption.

Management options[8]

- Give explanation, reassurance and support (ensurethat the patient is involved in the discussion).
- Reassurance can only be given following a careful examination and thoughtfully chosen investigations.
- Avoid investigations, epeceally radiological of possifly (FBE and MCU are ok)
- Acknowledge that the child has pain.
- Emphasise that the disorder is common, and usually traverses childhood without ill effects.
- Recommend simple measues as local warmth, brief rst for painful episode.
- Advise revew of epersiodes change in nature, painnew symptim.
- Identify any life stresses and provide insight therapy.
- Enquire about family structures and function, and school performance.
- Discourage identification with the sick role.
- Refer for psychological assessment and counselling if necessary.

ABDOMINAL PAIN IN THE ELDERLY

The elderly can suffer from a wide spectrum of disorders. Ischaemic events, emboli, cancer (in particular) and diverticulae of the colon are more common in old age; duodenal ulcer is less so. Those causes of abdominal pain that occur with more frequency include:

- vascular catastrophies
  - ruptured AAA
  - mesenteric artery occlusion
- perforated peptic ulcer
- biliary disorders: bilary pain and acute cholecystitis
- diverticulitis
- sigmoid volvulus
- strangulated hernia
- intestinal obstruction
- carcinoma, especially of the large bowel
- herpes zoster, causing unilateral root pain
- constipation and faecal impaction

Problems arise with management because the pain threshold is raised (colic in particular is less severe) and there is an attenuated response to infection so that fever and leucocytosis can be absent. Non-specific signs, such as confusion, anorexia and tachycardia, might be the only systemic evidence of infection.

ABDOMINAL AORTIC ANEURYSM

An AAA may be asymptomatic until it ruptures or may present with abdominal discomfort and a pulsatile mass noted by the patient. There tends to be a family history and thus screening is appropriate in such families. Ultrasound screening is advisable in first-degree relatives over 50 years.

The risk of rupture is related to the diameter of the AAA and the rate of increase in diameter. The normal diameter of the abdominal aorta, which is palpated just above the umbilicus, is 10 - 30 mm, being 20 mm on average in the adult; an aneurysm is greater than 30 mm in diameter.[9] Greater than 50 mm is significantly enlarged and is chosen as the arbitrary reference point to operate because of the exponential rise in risk of rupture with an increasing diameter. Refer all cases. The patency of a dacron graft after 5 years is approximately 95% (Figure 34.7).



Figure 34.7 Guidelines for normal and abnormal widths of the abdominal aorta in adults (to exact scale)

Investigations

- Ultrasound (good for screening) in relatives > 50 years
  - Obesity a problem
- CT scan (clearer imaging)
  - Helical/spiral scan is investigation choice
- MRI scan (best definition)

Rupture of aneurysm

This is a real surgical emergency in an elderly person who presents with acute abdominal and perhaps back pain with associated circulatory collapse (Figure 34.8). The patient often collapses at toilet because they feel the need to defecate and the resultant Valsalva manoeuvre causes circulatory embarrassment.



Figure 34.8 Typical pain distribution of a ruptured aortic abdominal aneurysm

DxT: intense pain + pale and 'shocked' ? back pain ° RAAA

The patient should be transferred immediately to a vascular surgical unit, which should be notified in advance. Two important emergency measures for the 'shocked' patient are intravenous access for plasma expanding fluid (a central venous line is best if possible) and he application of medical anti-shock trousers (MAST).

MESENTERIC ARTERY OCCLUSION

Acute intestinal ischaemia arises from superior mesenteric artery occlusion from either an embolus or a thrombosis in an atherosclerotic artery. Another cause is an embolus from atrial fibrillation. Necrosis of the intestine soon follows if intervention is delayed.

Typical clinical features

- Abdominal pain - gradually becomes intense (Figure 34.9)



Figure 34.9 Typical pain distribution of mesenteric arterial occlusion

DxT: anxiety and prostration + intense central pain + profuse vomiting ± bloody diarrhoea ° mesenteric arterial occlusion

- Profuse vomiting
- Watery diarrhoea - blood in one-third of patients (later)
- Patient becomes confused

Signs

- Localised tenderness, rigidity and rebound over infarcted bowel (later finding)
- Absent bowel sounds (later)
- Shock develops later
- Tachycardia (may be atrial fibrillation and other signs of atheroma)

Investigations

- CRP: May be elevated intestinal alkaline phosphatase.
- X-ray (plain): Shows 'thumb printing' due to mucosal oedema on gas-filled bowel. CT scanning gives the best definition while mesenteric arteriography is performed if embolus is suspected. However, it is commonly only diagnosed at laparotomy.

Management

Early surgery may prevent gut necrosis but massive resection of necrosed gut may be required as a life-saving procedure. Early diagnosis (within a few hours) is essential.

Note:

- Mesenteric venous thrombosis can occur but usually in patients with circulatory failure.
- Inferior mesenteric artery occlusion is less severe and survival more likely.

ACUTE RETENTION OF URINE

Acute retention of urine usually causes severe lower abdominal pain, which may not be apparent in a senile or demented person. Apart from the common cause of an enlarged prostate it can also result from bladder neck obstruction by faecal loading or other pelvic masses or anticholinergic drugs. It is often precipitated by extreme cold or an excess of alcohol.

Management

- Perform a rectal examination and empty rectum of any impacted faecal material.
- Catheterise with size 14 Foley catheter to relieve obstruction and drain.
- Have the catheter in situ and seek a urological opinion.
- If there is any chance of recovery (e.g. if the problem is drug-induced), withdraw drug, leave catheter in for 48 hours, remove and give trial of prazosin 0.5 mg bd or terazosin.

FAECAL IMPACTION

Faecal impaction is encountered typically in the aged, bedridden, debilitated patient. It may closely resemble malignant obstruction in its clinical presentation.[10] Spurious diarrhoea can occur, which is known as ‘faecal incontinence’.

ACUTE APPENDICITIS

Acute appendicitis is mainly a condition of young adults but it affects all ages (although uncommon under 3 years). It is the commonest surgical emergency and special care has to be taken with the very young and the very old. The symptoms can vary because of the different positions of the appendix.

Typical clinical features

See Figure 34.10. Typical clinical features are:


DxT: localised RIF pain + a/n/v + guarding = acute appendicitis
First
central abdominal pain
(visceral pain)
Later
RIF pain (somatic pain)
McBurney's point

Figure 34.10 Typical pain distribution of acute appendicitis

- maximum incidence 20 - 30 years
- initial pain is central abdominal (sometimes colicky)
- increasing severity and then continuous
- shifts and localises to RIF within 6 hours
- may be aggravated by walking (causing a limp) or coughing
- sudden anorexia
- nausea and vomiting a few hours after the pain starts

DxT: localised RIF pain + a/n/v + guarding ° acute appendicitis

Signs

- Patient looks unwell
- Flushed at first, then pale
- Furred tongue and halitosis
- May be febrile
- Tenderness in RIF, usually at McBurney's point
- Local rigidity and rebound tenderness
- Guarding
- ± Superficial hyperaesthesia
- ± Psoas sign: pain on resisted flexion of right leg, on hip extension or on elevating right leg (due to irritation of psoas especially with

retrocaecal appendix)

- ± Obturator sign: pain on flexing patient's right thigh at the hip with the knee bent and then internally rotating the hip (due to irritation of internal obturator muscle)
- Rovsing's sign: tenderness in RIF while palpating in LIF
- PR: anterior tenderness to right, especially if pelvic appendix or pelvic peritonitis

Variations and cautions

- Abscess formation ? localised mass and tenderness
- Retrocaecal appendix: pain and rigidity less and may be no rebound tenderness; loin tenderness; positive psoas test
- Pelvic appendix: no abdominal rigidity; urinary frequency; diarrhoea and tenesmus; very tender PR; obturator tests usually positive
- Elderly patients: pain often minimal and eventually manifests as peritonitis; can simulate intestinal obstruction
- Pregnancy (occurs mainly during second trimester): pain is higher and more lateral; harder to diagnose; peritonitis more common
- Perforation more likely in the very young, the aged and the diabetic

Investigations

Few investigations including imaging are of value:

- blood cell count shows a leucocytosis (75%) with a left shift
- urea electrolytes to assen hyaration prior to surging.
- CRP - elevated
- plain X-ray may show local distension, blurred psoas shadow and fluid level in caecum
- ultrasound shows a thickened appendix (80 - 90% accurate)[11]
- CT scan also accurate and allows other causes, especially in the female pelvis, to be evaluated[12]
- laparoscopy

Management

Immediate referral for surgical removal.

SMALL BOWEL OBSTRUCTION

The symptoms depend on the level of the obstruction (Table 34.6). The more proximal the obstruction the more severe the pain.

Table 34.6 Small bowel obstruction: difference between a high and a low obstruction

	High	Low
Frequency of spasms	3 - 5 minutes	6 - 10 minutes
Intensity of pain	+ + +	+
Vomiting	Early, frequent Violent	Later Less severe
Content:	Gastric juices then green	Feculent (later)
Dehydration and degree of illness	Marked	Less prominent
Distension	Minimal	Marked

Main causes

- Outside obstruction (e.g. adhesions - commonest cause, previous laparotomy), strangulation in hernia or pockets of abdominal cavity (Figure 34.11)



Figure 34.11 Operative findings (corrugated drainage material) in a 65 year old man with subacute bowel obstruction after a 21 year history of nagging abdominal pain following a cholecystectomy

- lumen obstructions (e.g. foreign body, trichobezoar, gallstones, intussusception, malignancy)

Typical clinical features

- Severe colicky epigastric and periumbilical (mainly) pain (Fig 34.12)



Figure 34.12 Typical pain distribution of small bowel obstruction

DxT: colicky central pain + vomiting + distension ° SBO

- Spasms last about 1 minute
- Spasms every 3 - 10 minutes (according to level)
- Vomiting
- Absolute constipation (nil after bowel emptied)
- No flatus
- Abdominal distension (especially if lower SBO)

Signs and tests

- Patient weak and sitting forward in distress
- Visible peristalsis, loud borborygmi
- Abdomen soft (except with strangulation)
- Tender when distended
- Increased sharp tinkling bowel sounds
- Dehydration rapidly follows, especially in children and elderly
- PR: empty rectum, may be tender

Note:check all hernial orifices, including umbilicus

- X-ray: plain erect film confirms diagnosis ‘stepladder’ fluid levels in 3 - 4 hours
    - Gastrografin follow-through for precise diagnosis(don't give it in a ‘virgin abdomen‘). It can cause serve diarrhoese and maybe therapentu in adheseve obstructure.

Management

- IV fluids and bowel decompression with nasogastric tube
- Laparotomy or hernia repair

LARGE BOWEL OBSTRUCTION

The cause is commonly colon carcinoma (75% of cases), especially on the left side, but it can occur in diverticulitis or in volvulus of the sigmoid colon (10% of cases) and caecum.[10] Sigmoid volvulus is more common in older men and has a sudden and severe onset. The pain is less severe than in SBO. Be wary of the non-surgical causes, simple constipation or acute pseudo-obstruction of the colon (Ogilvie's syndrome).

Typical clinical features

- Sudden-onset colicky pain (even with carcinoma)
- Each spasm lasts less than 1 minute
- Usually hypogastric midline pain (Figure 34.13)



Figure 34.13 Typical pain distribution of large bowel obstruction

- Vomiting may be absent (or late)
- Constipation, no flatus

DxT: colicky pain + distension ± vomiting ° LBO

Signs and tests

- Increased bowel sounds, especially during pain
- Distension early and marked
- Local tenderness and rigidity
- PR: empty rectum; may be rectosigmoid carcinoma or blood. Check for faecal impaction
- X-ray: distension of large bowel with separation of haustral markings, especially caecal distension
  - Sigmoid volvulus shows a distended loop.
  - Gastrografin enema confirms diagnosis.

Management

Surgical referral.

PERFORATED PEPTIC ULCER

Perforation of a peptic ulcer can cause acute abdominal pain both with and without a prior history of peptic ulcer. It is an acute surgical emergency requiring immediate diagnosis. Consider a history of drugs, especially NSAIDs and H_2-receptor antagonists. Perforated ulcers may follow a heavy meal. There is usually no back pain.

The maximal incidence is 45 - 55 years, most common in males, and a perforated duodenal ulcer is more common than a gastric ulcer.

Consider the clinical syndrome in three stages:

1. prostration
2. reaction (after 2 - 6 hours) - symptoms improve
3. peritonitis (after 6 - 12 hours)

Typical clinical features

See Figure 34.14. Typical clinical features are:



Figure 34.14 Typical features of perforated peptic ulcer

DxT: sudden severe pain + anxious, still, 'grey', sweaty + deceptive improvement ° perforated peptic ulcer

- sudden-onset severe epigastric pain
- continuous pain but lessens for a few hours
- epigastric pain at first, and then generalised to whole abdomen
- pain may radiate to one or both shoulders (uncommon) or right lower quadrant
- nausea and vomiting (delayed)
- hiccough is a common late symptom

Signs and tests

- Patient lies quietly (pain aggravated by movement and coughing)
- Pale, sweating or ashen at first
- Board-like rigidity
- Guarding
- Maximum signs at point of perforation
- No abdominal distension
- Contraction of abdomen (forms a 'shelf' over lower chest)
- Bowel sounds reduced (silent abdomen)
- Shifting dullness may be present
- Pulse, temperature and BP usually normal at first

- Tachycardia (later) and shock later (3 - 4 hours)
- Breathing is shallow and inhibited by pain
- PR: pelvic tenderness
- X-ray: chest X-ray may show free air under diaphragm (in 75%) - need to sit upright for prior 15 minutes
  - limited Gastografin meal can confirm diagnosis

Special problems

- Beware of easing of pain as peritoneal fluid accumulates.
- Elderly patients may have minimal pain.
- Painless perforation can occur with steroids.
- Avoid giving morphine or pethidine until diagnosis confirmed.

Management

- Pain relief
- Drip and suction (immediate nasogastric tube)
- Broad-spectrum antibiotics
- Immediate laparotomy after resuscitation
- Conservative treatment may be possible (e.g. later presentation and Gastrografin swallow indicates sealing of perforation)

URETERIC COLIC

Renal colic is not a true colic but a constant pain due to blood clots or a stone lodged at the pelvic-ureteric junction; ureteric colic, however, presents as severe true colicky pain due to stone movement and ureteric spasm. Fortunately, the majority of urinary calculi are small and will pass spontaneously.

Guideline :

- loin pain - stone in kidney.
- kidney/ureteric colic - ureteric stone.
- strangury - stone in bladden.

Typical clinical features

- Maximum incidence 30 - 50 years (M > F)
- Intense colicky pain: in waves, each lasting 30 seconds with 1 - 2 minutes respite
- Begins in loin and radiates around the flank to the groin, thigh, testicle or labia (Figure 34.15)



Figure 34.15 Ureteric colic: typical radiation of pain in left ureteric colic

- Usually lasts < 8 hours
- ± Vomiting

DxT: intense pain (loin) → groin + microscopic haematuria ° ureteric colic

Signs

- Patient restless: may be writhing in pain
- Pale, cold and clammy
- Tenderness at costovertebral angle
- ± Abdominal and back muscle spasm
- Smoky urine due to haematuria

Diagnosis

- Urine: microscopy; blood testing strip (negative does not exclude calculus)
- Plain X-ray: most stones (75%) are radio - opaque (calcium oxalate and phosphate)
- IVP: confirms opacity, level of obstruction, kidney function and any anatomical abnormalities
- Ultrasound: may locate calculus but will exclude obstruction
- Non-contrast spiral CT is the ‘gold standard’ (sensitivity 97%, specificity 96%)
  - will show easily missed radiolucent uric acid stones

Management

If the diagnosis is in doubt (especially if narcotic addiction is suspected) get the patient to pass urine in the presence of an examiner and test for haematuria. While awaiting passage of urine, an indomethacin suppository may be tried for pain relief.

Routine treatment (average size adult)

- Morphine 15 mg IM or 10 mg IV[13] *and* metoclopramide 10 mg IM *or* hyoscine 20mg IV
- Avoid high fluid intake - provokes distension of urethra and aggravates pain.
- Most cases settle and the patient can go home when pain relief is obtained and an IVP arranged for the next day.
- Further pain can be alleviated by indomethacin suppositories but should be limited to two a day.

- An effective alternative treatment is diclofenac 75 mg IM injection then 50 mg (o) tds for 1 week. Several clinical trials have shown that NSAIDs by IM injection, including ketoralac, are effective and at least as efficaceous as opioids. [14, 15]

Outcome and follow-up

- The calculus is likely to pass spontaneously if < 5 mm (90% < 4 mm pass spontaneously).[16]
- If > 5 mm intervention will usually be required by lithotripsy or surgery.
- If the patient passes the calculus, he or she should retrieve it and present it for analysis.
- A repeat IVP may be necessary if there is evidence of obstruction for more than 3 weeks.
- The cause of the ‘stone’ should be considered. Search for causes such as hyperparathyroidism, hypercalcaemia, hyperoxaluria and UTI.
- Fever with ureteric colic indicates an obstructed infected kidney.

When to refer[14]

Any of the following:

- stone > 5 mm in diameter
- high-grade obstruction
- gross hydronephrosis
- fever/UTI
- unremitting pain
- stone fails to progress
- type 2 diabetes
- staghorn calculus
- presence of solitary kidney

Facts about urinary tract calculi

- The prevalence is 1 to 3 per 1000 population per year[14]
- The lifelong incidence is 10%
- The recurrence is up to 75% (most within 2 years)
- The typical age range is 20 to 50 years (peaks at 28 years)
- Pregnancy is a risk factor
- Male to female ratio ° 3:1
- The incidence is inversely proportional to fibre intake and proportional to ingestion of animal protein and persistently low urinary volume
- Formed from urinary supersaturation with calcium (calcium oxalate, 75 - 80%), uric acid (7%) and cysteine (rare): also infected calculi (struvite) - Mg^{+}, NH_4^{+}, PO_4^{-} (5%)

‘Phony’ colic

Some patients who present with typical colic may be feigning their pain mainly because they are opioid dependent and seeking drugs by deception. As ureteric colic can affect young people (peak age 28 years) this can be a very difficult management issue, even for the experienced.

The use of CT scanning would help in locating calculi in such patients. If in doubt an appropriate agent would be ketoralac (Toradol) 10 - 30 mg IM.

It is advisable to obtain a specimen of urine passed in the presence of an examiner and then tested for microscopic blood. While awaiting passage of urine an indomethacin suppository may be tried for pain relief.

Recurrent urinary calculi

Investigations:

- serum electrolytes, urea, creatinine
- serum calcium, phosphate, uric acid, magnesium
- serum alkaline phosphatase
- urine sample - microbiology and culture
- at least two consecutive 24 hour urine samples
- stone analysis
- IVP

Dietary advice is given on page 84.

BILIARY PAIN

Abdominal pain can be produced by contraction of the biliary tree upon an obstructing stone or inspissated bile. Although the stereotyped patient is female, forty, fat, fair and fertile it can occur from adolescence to old age and in both sexes.

Typical clinical features

See Figure 34.16.


pain may radiate to R shoulder
pain radiation to scapula
localised tenderness (Murphy's sign)

Figure 34.16 Typical site of pain of biliary colic and acute cholecystitis

DxT: severe pain + vomiting + pain radiation ° biliary colic

Typical clinical features are:

- acute onset severe pain
- post-prandial or at night (often wakes 2 - 3 am)
- constant pain (not colicky)
- lasts 20 minutes to 2 - 6 hours
- maximal RUQ or epigastrium
- may radiate to tip of right shoulder or scapula
- painful episode builds to a crescendo for about 20 minutes; may recede or last for hours
- some relief by assuming flexed posture
- ± nausea and vomiting with considerable retching
- often a history of biliary pain (may be mild) or jaundice
- often precipitated by a fatty meal

Signs

- Patient anxious and restless, usually in a flexed position or rolling in agony

- Localised tenderness (Murphy's sign) over fundus of gall bladder (on transpyloric plane)
- Slight rigidity

Diagnosis

- Abdominal ultrasound/DIDA
- CT-helical
- Intravenous cholangiography if previous cholecystectomy
- LFTs may show elevated bilirubin and alkaline phosphatase

Management options

- Pain relief: morphire 10 - 15 mg IM, even 3 to 4 houre hyoscine 20 mg Imor ovel[13].
- Gallstone dissolution or lithotripsy (in those unable to have surgery)
- Cholecystectomy (main procedure)

Gallstone facts[16]

- Gallstones form from bile in the gallbladder and sometimes in the bile duct (especially post-cholecystectomy)
- Two main types - cholesterol and pigment (bilirubin)
- Lifetime risk in first world countries is 12 - 20%
- 70% of people with gallbladder stones are asymptomatic, but risk of developing symptoms is about 15% over 20 years
- Cholecystectomy almost never indicated for such stones
- Complications: acute cholecystitis (may lead to empyema, perforation, cholecystoenteric fistula), obstructive jaundice, cholangitis and acute pancreatitis (pancreatic duct obstruction)

Acute cholecystitis

Cholecystitis is associated with gallstones in over 90% of cases[17] and there is usually a past history of biliary pain. It occurs when a calculus becomes impacted in the cystic duct and inflammation develops. It is very common in the elderly. The acute attack is often precipitated by a large or fatty meal. The cousative organisins are usually aerolic bowd flor eg Eldi Klesiella speues and enterococcus faecalis.

Typical clinical features

- Steady severe pain and tenderness
- Localised to right hypochondrium or epigastrium
- Nausea and vomiting (bile) in about 75%
- Aggravated by deep inspiration

Signs

- Patient tends to lie still
- Localised tenderness over gall bladder (positive Murphy's sign)
- Muscle guarding
- Rebound tenderness
- Palpable gall bladder (approximately 15%)
- Jaundice (approximately 15%)
- ± Fever

Diagnosis

- Ultrasound: gallstones but not specific for cholecystitis
- HIDA scan: demonstrates obstructed cystic duct - the usual cause
- WCC and CRP: can be elevated

Treatment

- Bed rest
- IV fluids
- Nil orally
- Analgesics
- Antibiotics
- Cholecystectomy

It every of sepsis use aoxy lamicollin 1/9 IV, ohouly gentamicin 4 - 6 mg/Kg IV daly[18].

ACUTE PANCREATITIS

With acute pancreatitis there may be a past history of previous attacks or a past history of alcoholism (35%) or gallstone disease (40 - 50%).

Typical clinical features

See Figure 34.17.

DxT: severe pain + nausea & vomiting +
relative lack abdominal signs
= acute pancreatitis



Figure 34.17 Typical pain distribution of acute pancreatitis

DxT: sever pain + nausea & vomiting + relative lack of abdominal signs ° acute pancreatitis

Typical clinical features are:

- sudden onset of severe constant epigastric pain
- lasts hours or a day or so
- pain may radiate to back
- nausea and vomiting
- sweating and weakness

Signs

- Patient is weak, pale, sweating and anxious
- Tender in epigastrium

- Lack of guarding, rigidity or rebound
- Reduced bowel sounds (may be absent if ileus)
- ± adominal distension
- Fever, tachycardia

Diagnosis

- WCC - leucocytosis
- Serum amylase (usually five-fold increase) or Lipase.
- CRP - elevated
- Serum glucose ↑, calcium ↓
- Blood gases: $P_a o_2$(pulmonary comphcation)
- LFTs: ? obstructive pattern
- Plain X-ray, may be senital loop
- CT scan (especially for complications)
- Ultrasound better for detecting cysts and unsuspected gallstones

Management

- Arrange admission to hospital.
- Basic treatment is bed rest, nil orally, nasogastric suction (if vomiting), IV fluids and analgesics (morphine).
- Use morphine10 - 15mgimevery 3 - 4 hours
- or 10mg morphine IV follpaved by continwou infeij.
- May require ERCP if obstructive LFTs.

Chronic pancreatitis

In contrast to acute pancreatitis the pain of chronic pancreatitis is milder but more persistent. The patient with this problem is often labelled as ‘gastritis’, ‘ulcer’ or ‘neurotic’ because of the indeterminate nature of the pain. Malabsorption and diabetes may result from pancreatitis and weight loss and steatorrhoea become prominent features.

Pain associated with carcinoma of the pancreas is indistinguishable from that of chronic pancreatitis but generally tends to be more severe and more prominent in the back. Use paracetamol or codeine for pain.

ACUTE DIVERTICULITIS

The patient with acute diverticulitis is usually over 40 years of age, with long-standing grumbling left-sided abdominal pain and constipation, but can have irregular bowel habit. It occurs in less than 10% of patients with diverticular disorder.[2]

Typical clinical features

See Figure 34.18.



Figure 34.18 Typical pain distribution of acute diverticulitis

DxT: acute pain + left-sided radiation + fever ° acute diverticulitis

Typical clinical features are:

- acute onset of pain in the left iliac fossa
- pain increased with walking and change of position
- usually associated with constipation

Signs

- Tenderness, guarding and rigidity in LIF
- Fever
- May be inflammatory mass in LIF

Investigations

- FBE Leucocytosis
- Elevated ESR
- Pus and blood in stools
- Abdominal ultrasound/CT scan (especially can delte fistula or perforation)
- Erect chest X-ray
- Erect and supine abdominal X-ray.

Complications

- Bleeding (can be profuse, especially in elderly)
- Perforation (high mortality)
- Abscess
- Peritonitis
- Fistula (bladder, vagina, small bowel)
- Intestinal obstruction

Treatment[18]

- Antibiotics
    - mild cases: amoxycillin + clavulanate 500 mg (o) tds for 5 - 7 days or metronidazole + cephalexin
    - severe: (amoxy) ampicillin 2 g IV 6 hourly + gentamycin 5 - 7 mg/kg IV/day + metronidazole 500 mg IV 12 hourly or metronidazole + ceftriaxone 1g IV/day
- surgery for complications

PENTONITIS

Peronitis can be acute due to intra-abdominal sepsi notably following peroration of a viscuo sugual intervenuton, as approprale is useally regaled artiboore tretunent (intravewiely) is by amoxy/amipallin + genatamian + metronidagule[18].

ABDOMINAL ‘STITCH’

The common ‘stitch in the side’ is the experience of a sharp, stabbing pain in the epigastric or hypochondrium regions of the abdomen, usually during running. The sufferer should:

- stop and rest, then walk - don't run
- apply deep massage to the area with the palps of the middle three fingers
- perform slow or deep breathing

CHRONIC OR RECURRENT ABDOMINAL PAIN

Advances in technology have increased the opportunities for diagnosing chronic or recurrent abdominal pain[19] in adults:

- Ultrasound. This is the single most useful screening test. It can detect gallstones, pelvic pathology, such as carcinoma of the ovary, and retroperitoneal problems, such as abdominal aneurysm and carcinoma of the pancreas.
- Endoscopy. This is the next most useful investigation. The history dictates whether it is gastroscopy or colonoscopy.

If the patient has ‘red flag’ symptoms (Table 34.7) and the above investigations are absent, consider the possibility of conditions such as carcinoma of the pancreas, carcinoma of the ovary, small bowel tumours, mesenteric ischaemia, Crohn's disease, metabolic disorders such as lactase deficiency, and rarer conditions as outlined in Table 34.3.

Table 34.7 Red flags for organic disease[12]

Older patient
Nocturnal pain or diarrhoea
Progressive symptoms
Rectal bleeding
Fever
Anaemia
Weight loss
Abdominal mass
Faecal incontinence or urgency (recent onset)

Other investigations that may help:

- CT scanning
- laparoscopy - this may allow the identification of chronic adhesive obstruction, small bowel tumours or inflammation, or intra-abdominal malignancy

Chronic appendicitis

It is possible to have recurrent episodes of subacute inflammation of the appendix. If suspected, laparoscopy performed during or soon after an attack is diagnostic.

Adhesions

There is no firm evidence that intra-abdominal adhesions are painful apart from complications such as bowel obstruction. Sometimes patients are ‘cured’ by laparoscopic divisions of adhesions.

Irritable bowel syndrome

See pages 488 - 9.

At least 3 months of continuous or recurrent:

- cramping pain (relieved by defecation)
- central or left lower quadrant pain (more common) but can be at any site
- mucus in stool
- altered stool form or passage

Peptic ulcer (gastric or duodenal)

See pages 524 - 5.

- Usually central epigastric pain
- Burning pain
- Relieved by antacids or food or milk
- DU: usually 2 - 3 hours after meals or wakes from sleep
- GU: may occur after meals but inconsistent relationship to eating

WHEN TO REFER

- All cases of acute abdominal pain where urgent surgical intervention is required; special urgency and early diagnosis is important with:
  - ruptured ectopic pregnancy
  - ruptured AAA
  - mesenteric artery occlusion
  - ruptured viscus
  - perforated peptic ulcer
  - strangulated obstructed bowel
  - intussusception
- All cases where the diagnosis is not apparent
- All cases where surgery is necessary
- Medical causes, such as diabetic ketoacidosis and porphyria

Practice tips

- Special caution is required at the extremes of age when the symptoms and signs do not often reflect the seriousness of the underlying pathology.
- If an elderly patient presents with intense acute abdominal pain, inadequately relieved by strong parenteral injections, likely causes include mesenteric artery occlusion, acute pancreatitis and ruptured or dissecting aortic aneurysm.
- When an inflamed appendix ruptures, the abdominal pain improves for a significant period of time.
- Consider gallstones and duodenal ulcer if the patient is woken (e.g. 2 - 3 am) with abdominal pain.
- A false sense of security can also occur with a perforated ulcer.
- Pus cells and red cells may be present in the urine with appendicitis when a pelvic appendix involves the bladder and a retrocaecal appendix involves the ureter.
- Consider diabetic ketoacidosis in a patient with abdominal pain, tenderness and rigidity and deep sighing respiration.

REFERENCES

1. de Wit NJ. Acute abdominal pain. In: Jones R (et al.) eds. *Oxford Textbook of Primary Care*. Vol. 2. Oxford: Oxford University Press, 2004: 738 - 40.
2. Sandler G, Fry J. Acute abdominal pain. In: *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986: 137 - 76.
3. Murtagh J. *The Anatomy of a Rural Practice*. Melbourne: Monash University Monograph, 1980: 34.
4. Sandler G, Fry J. Chronic abdominal pain. In: *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986: 177 - 86.
5. Bain ST, Spaulding WB. The importance of coding presenting symptoms. Can Med Assoc J, 1967; 97(16): 953 - 9.
6. Hutson JM Woodward AA, Beasleyls SW, *Jones clinical paediotric suigery*. Oxford Blackwell publishing, 2003; 139 - 145.
7. Hodgkin K. *Towards Earlier Diagnosis* (5th edn). Edinburgh: Churchill Livingstone, 1985: 483 - 6.
8. Dilley A. Abdominel surgical problems in children medical observe, 23 April 2005: 31 - 4.

9. Appleberg M. Abdominal aortic aneurysms. In: *How to Treat*. Aust Doctor, 2001; 22 June: iii - iv.
10. Hunt P, Marshall V. *Clinical Problems in General Surgery*. Sydney: Butterworths, 1991: 193 - 243.
11. Lau L (ed). *Imaging Guidelines* (4th edn). Melbourne: RA&NZC Radiologists, 2001.
12. Rao P, Boland G. Imaging of acute right lower abdominal quadrant pain. Clin Radiol, 1998; 53: 639 - 49.
13. Mashford ML (chair), *Therapeule guidelines Analgense* (version 4)Melborne: Therapeto ccvideline Ltd, 2002:257 - 8
14. Laerum E, Murtagh J. Kidney colt and recurrent urinary calculi. Aust Fam Physician, 2001; 30.(1): 36 - 41.
15. Laerum E et al. Oral diclofenac in the treatment of recurrent kidney colic. A double-blind comparison with placebo. Eur Urol, 1995; 28: 108 - 11.
16. Tooceli J, Wright T. Gallstones. Med J Aust, 1998; 169: 166 - 71.
17. Tierney LM et al. Current Medical Diagnosis and Treatment (41st edn). New York: McGraw-Hill and Lange, 2002: 707.
18. Spicer J (Chair). *Therapeeutie Guidelines :Antibiotic* (version 13) Melbousine Therapeutic Guidelines Ltd, 2006:89 - 142
19. Hugh TB. The investigation and management of recurrent or chronic abdominal pain. Mod Med Aust, 1995; 38(5): 126 - 32.

CHAPTER 35. ARTHRITIS

Rheumatic disorders are common in old age: much of rheumatology is geriatric and much of geriatrics is rheumatology.

DR FRANK DUDLEY HART [1] 1983

ARTHRITIS

The clinical evaluation of the patient presenting with the complaint of arthralgia (painful joints) or arthritis (inflammation of the joints) can be a difficult and challenging exercise because it can be a presentation of many systemic disorders, some of which are rare. Important considerations are sex, age, the pattern of joint involvement (monoarticular or polyarticular), immediate and more remote history, family history and drug use - all of which may provide important diagnostic clues. Polyarthritis, which implies the active inflammation of five or more joints, presents a more challenging diagnostic problem.

KEY FACTS AND CHECKPOINTS

- In a UK National Morbidity Survey rheumatic disease composed just over 7% of all morbidity presenting to the family doctor.[2]
- The commonest cause was osteoarthritis (OA), which affects 5 - 10% of the population.
- The same study indicated an episode rate for arthritis/arthralgia of 38.6 per 1000 population.
- The population incidence of rheumatoid arthritis (RA) is 1 - 2%.
- There should be no systemic manifestations with OA.
- One-quarter of disability in elderly people is due to severe joint disease.
- Systemic diseases that may predispose to, or present with, an arthropathy include the connective tissue disorders, diabetes mellitus, a bleeding disorder, previous tuberculosis, the spondyloarthropathies such as psoriasis, SBE, hepatitis B, rheumatic fever, the various vasculitic or arteritic syndromes (the vasculitides) such as Wegener's granulomatosis, HIV infection, carcinoma of the lung, haemochromatosis, sarcoidosis, hyperparathyroidism and Paget's disease.
- The pain of inflammatory disease is worse at rest (e.g. on waking in the morning) and improved by activity.
- Early diagnosis and management of rheumatoid arthritis (RA) results in considerably better outcomes.
- Gout and septic arthritis have a recognised cause and cure.
- Acute gout is almost exclusive to males: in women, it is usually seen only in those who are postmenopausal or taking thiazide diuretics.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic strategy model is presented in Table 35.1.

Table 35.1 Arthralgia: diagnostic strategy model

Q. Probability diagnosis A. Osteoarthritis Viral polyarthritis
Q. Serious disorders not to be missed A. Rheumatoid arthritis Connective tissue disorders • SLE • systemic sclerosis • dermatomyositis Neoplasia • carcinoma bronchus • leukaemia/lymphoma HIV arthropathy Severe infections

- rheumatic fever
- endocarditis
- tuberculosis
- brucellosis
- pyogenic arthritis
  - gonococcus
  - Staphylococcus

Q. Pitfalls (often missed)

A. Fibromyalgia syndrome

Polymyalgia rheumatica

Crystal deposition

- gout
- pyrophosphate (pseudogout)

Haemarthrosis

Dengue fever

Lyme disease

Ross River virus

Avascular necrosis

Rarities

Other vasculitides

Haemochromatosis

Sarcoidosis

Hyperparathyroidism

Familial Mediterranean fever

Amyloidosis

Pigmented villonodular synovitis

Q. Seven masquerades checklist

A. Depression

Diabetes

Drugs

Anaemia -

Thyroid disorder

Spinal dysfunction spondyloarthropathies

UTI -

Q. Is the patient trying to tell me something?

A. Always a consideration with pain

Psychogenic factors aggravate chronic arthritic conditions.

Probability diagnosis

The probability diagnoses for the patient presenting with arthritis are:

- osteoarthritis (mono- or polyarthritis)
- viral arthritis (if acute and polyarthritis)

OA is very common in general practice. It may be primary, which is usually symmetrical, and can affect many joints. This clinical pattern is different from secondary OA, which follows injury and other wear-and-tear causes.

Viral polyarthritis is more common than realised. It presents usually within 10 days of the infection, and is usually mild.

Features of viral arthritis

- Acute onset
- A polyarthritis
- Symmetric inflammation
- Mainly of hands and feet
- Rash-persists for 24hrs minimum
- Terminates rapidly-over days
- FBE: lymphopaenia, lymphocytosis, ± atypical lymphocytes

It tends to terminate quickly and spontaneously without permanent damage to joints. It is caused by many viruses, including those causing influenza, mumps, rubella, varicella, hepatitis B and C, infectious mononucleosis (more muscle aching), cyto-megalovirus, parvovirus, Australian epidemic polyarthritis due to the alphaviruses, Ross River virus and Barmah Forest virus. Adenovirus is common in children.

Serious disorders not to be missed

These include RA, which can start as a monoarthritis; pyogenic arthritis, including gonococcus, *Staphylococcus* and *Streptococcus* infections; tuberculosis; rheumatic fever; and bacterial endocarditis. Early diagnosis of septic arthritis is very important as it can destroy a hip in 24 hours.

It is important to be forever watchful for rheumatic fever (RF). It presents typically as a migratory polyarthritis involving large joints sequentially, one becoming hot, red, swollen and very painful as the other subsides. It rarely lasts more than 5 days in any one joint.

A flitting polyarthritis can also occur with endocarditis in addition to a systemic upset and a cardiac murmur. Gonococcal infection may present in a single joint or as flitting polyarthritis, often accompanied by a rash. Brucellosis can cause arthritis and sacroiliitis and can be confused with the spondyloarthropathies.

HIV infection is becoming a great mimicker. It can present as a chronic oligoarticular asymmetrical arthritis.[3] It can also present as a rash very similar to psoriasis.

With the large influx of migrants from South-East Asia the possibility of tuberculosis presenting as arthritis should be kept in mind.

Connective tissue disorders may be involved. They include SLE, progressive systemic sclerosis (scleroderma), and dermatomyositis. It is most inappropriate to settle with a general diagnosis such as 'rheumatism' or 'arthritis' and where doubtful it is important to find the specific entity causing the problem.

In respect to malignant disease, arthralgia is associated with acute leukaemia, lymphoma and neuroblastoma in children and with carcinoma of the bronchus, which may cause hypertrophic osteoarthropathy, especially of the wrist and ankle (not a true arthritis but simulates it). Occasionally polyarthritis may be the first feature of an occult neoplasm. Monoarticular metastatic disease may involve the knee (usually from lung or breast).

Red flags for polyarthritis

- Fever
- Weight loss
- Profuse rash
- Lymphadenopathy
- Cardiac murmur
- Severe pain and disability
- Malaise and fatigue
- Vasculitic signs
- Two or more systems involved

Pitfalls

There are several pitfalls, most of which are rare. A common pitfall is gout. This applies particularly to older women taking diuretics, whose osteoarthritic joints, especially of the hand, can be affected. The condition is often referred to as nodular gout and does not usually present as acute arthritis.

Fibromyalgia syndrome is a real puzzle (see p. 398) as it can mimic the connective tissue disorders in its early presentation - typically a woman in the third or fourth decade.

Another ‘trap’ is haemarthrosis in a patient with a bleeding disorder.

Infective causes that may be overlooked are dengue fever, especially in travellers returning from a tropical or subtropical area, and Lyme disease, which is now surfacing in many countries, especially where ticks are found.

There are many rare causes of arthritis. Sarcoidosis causes two forms: an acute benign form, usually in the ankles and knees, and a chronic form with long-standing sarcoidosis that involves joints (large or small) adjacent to underlying bone disease.

Then there are the uncommon vasculitides, which can cause confusion in diagnosis. This group includes polyarteritis nodosa, hypersensitive vasculitis, polymyalgia rheumatica/giant cell arteritis, Wegener's granulomatosis, Henoch-Schönlein purpura and Behçet's syndrome.

Haemochromatosis can present with a degenerative arthropathy that characteristically affects the second or third metacarpophalangeal joints.[3]

Other rare causes of arthritis are erythema nodosum, serum sickness and Sjögren's syndrome.

General pitfalls

- Not searching beyond RA when an RA pattern polyarthritis may be part of another systemic disease
- Failing to search for some cause of arthritis other than OA in a patient, especially an elderly patient (i.e. underdiagnosing); an important example of this is polymyalgia rheumatica
- Failing to consider the various drug interactions between NSAIDs, over-the-counter medications and other drugs used by the elderly
- Underdiagnosing and misdiagnosing through lack of appreciation of the many causes of arthritis, especially those presenting as part of a systemic disease

Seven masquerades checklist

- Depression - unlikely but complaints of arthralgia are possible
- Diabetes - occasionally causes an arthropathy
- Drugs - yes, a major consideration
- Anaemia - no
- Thyroid disease - possible
- UTI - no
- Spinal dysfunction - only with the spondyloarthropathies

Drug-induced arthritis is the main feature of this important group of disorders. It usually affects the hands and is generally symmetrical. The drugs may induce autoantibodies (e.g. ANA, ANCA). Drugs that cause arthritis are listed in Table 35.2. The problem usually resolves promptly after withdrawal of the agent.[4]

Table 35.2 Drug-induced arthralgia

Commonest drugs inducing arthralgia
Note: Usually affects the hands and is symmetrical **Drugs inducing lupus syndrome** • hydralazine • procainamide, quinidine • antiepileptics (e.g. phenytoin) • chlorpromazine • isoniazid • methyldopa • ACE inhibitors
Others • HMG CoA reductase inhibitors (statins) • cotrimoxazole • amoxycillin

- minocycline
- mianserin
- carbimazole
- progesterone only OCP
- nitrofurantoin
- antihypertensives
- cimetidine, famotidine

Note: Diuretics, especially frusemide and thiazides, can precipitate gout.

Intravenous drug abuse may be associated with septic arthritis, hepatitis B and C, HIV-associated arthropathy, SBE with arthritis and serum sickness reactions.

Hyperthyroidism can uncommonly cause acropathy (clubbing and swelling of the fingers) and may present as pseudogout, while hypothyroidism can present with an arthropathy or cause proximal muscle pain, stiffness and weakness. Diabetes mellitus may cause an arthropathy that can be painless or mild to moderately painful.

The spondyloarthropathies may be a causative factor. They often present with an acute monoarthritis, particularly in teenagers some time before causing sacroiliitis and spondylitis.

Psychogenic considerations

Although ‘arthralgia’ is an uncommon complaint in psychoneurotic disorders, any pain syndrome can be a significant manifestation. The usual cause of arthralgia is inflammation in the joint, that is arthritis, but a functional cause is encountered from time to time.

Furthermore some patients who are unfortunate enough to acquire arthritis, especially the more serious disorders, certainly develop ongoing emotional and psychological problems that appear to aggravate their total problem.

So-called ‘growing pains’ of the lower limb are common in children, and the physical examination and investigations are normal. Parents need to be reassured that it is a benign condition while recognising that emotional factors may be quite significant. As Apley pointed out, ‘physical growth is not painful, but emotional growth can hurt like hell’.[5]

THE CLINICAL APPROACH

A priority is to determine whether or not the arthritis is caused by a primary rheumatic disorder or whether it is part of an underlying systemic disorder.

History

Very careful enquiry about the exact onset of the arthritis is important. This includes whether it was acute or insidious, and confined to specific joints or flitting as in rheumatic fever and sometimes in infective endocarditis. Is it a true polyarthritis or monoarthritis? Symmetrical or asymmetrical? It is also important to differentiate between arthralgia (pains in or around the joints) and arthritis (inflammation of the joints). Not all arthralgia is arthritis.

A family history is important because a positive family history is associated with conditions such as RA (rarely), ankylosing spondylitis, connective tissue disorders (rarely), psoriasis, gout, pseudogout and haemophilia.[6]

A very hot, red, swollen joint suggests either infection or crystal arthritis.

Key questions

- Can you carefully point out exactly where you feel the pain?
- Does the pain move from joint to joint or stay in the same joint?
- Are you aware of anything that brought on the pain?
- Does the pain disturb you at night?
- Do your joints feel very sore or stiff when you wake up in the morning?
- What effect does exercise or activity have on the pain or stiffness?
- Have you had an injury in the past to your painful joint(s)?
- Do you get pain over both your shoulders and upper arms?
- Have you got a skin rash? Is it new?

- Have you had a fever, sweats or chills?
- Do you get very tired, weak or out of sorts?
- Have you noticed any change in the colour of your urine?
- Have you had a sore throat?
- Have you had sinus trouble?
- Have you had acute pain in your big toe or in other joints before?
- Do you have a history of psoriasis?
- Do you have a history of rheumatic fever?
- Do you have pain in your neck or lower back or in other joints?
- Have you had any diarrhoea?
- Have you had a discharge from your penis?
- Have you had any problems with your eyes?
- What drugs are you taking? Are you taking fluid tablets (diuretics)?
- How much alcohol would you drink a day?
- Have you been visiting the country or exposed to ticks or have you been to a deer farm?
- Have you travelled overseas recently?
- Have you been at risk of getting an STI?
- Have you been drinking untreated milk recently?
- Have you had cats as pets, especially as a child (associated with RA)?

Physical examination

A systematic examination of the affected joint or joints should be performed, looking for signs of inflammation, deformity, swelling and limitation of movement. Tenderness and warmth indicates inflammatory activity. Erythema indicates gouty arthritis or other crystalopathy, rheumatic fever or septic arthritis.

Joint swelling:

- acute (1 - 4 hours) with intense pain ° blood infection or crystals (e.g. gout)
- subacute (1 - 2 days) and soft ° fluid (synovial effusion)
- chronic and bony ° osteoarthritis
- chronic and soft/boggy ° synovial proliferation

A coarse crepitus suggests OA. Each joint should be examined specifically. Inspection should note the presence of lumps or bumps such as Heberden's nodes on the osteoarthritic DIP joints of the hands, Bouchard's nodes on the osteoarthritic PIP joints of the hands, and rheumatoid nodules, which are the only pathognomonic finding of RA and gouty tophi. Signs that may be of diagnostic help are presented in Figure 35.1.


temporal arteritis
tophi in ear: gout
eye signs: spondyloarthropathies
rheumatoid arthritis
butterfly rash: SLE
aortic incompetence: ankylosing spondylitis
pleural effusion/rub
• SLE
• RA
cardiac valvular lesions: RF
pericardial rub: SLE
RA
rheumatoid nodules
inflammation colon: ulcerative colitis
Crohn's disease
sacroiliitis: spondyloarthropathies
carpal tunnel syndrome: RA
Raynaud's phenomenon
• systemic sclerosis
• RA
• SLE
pitted nails: psoriasis
doughnut-shaped rash: Lyme disease
erythema marginatum: RF
psoriasis
gout

Figure 35.1 Physical examination: possible findings to consider in diagnosis

The specific inflamed joint or joints may give an indication of the disease process. Typical joints affected by various arthropathies are illustrated in Figure 35.2.


TMJ
• RA
Cervical spine
• RA
• OA
• spondyloarthropathies
Acromioclavicular joint
• OA
Glenohumeral joint
• RA
Thoracic spine
• OA
• spondyloarthropathies
Wrist
• RA
• OA
Elbow
• RA
Lumbar spine
• OA
• spondyloarthropathies
PIP
• RA
• viral
• SLE
Sacroiliac
• spondyloarthropathies
Hip
• OA
• RA
DIP
• OA
• psoriasis
• gout
• RA
Knee
• OA
• pseudogout > 60 years
• RA
• haemophilia
• spondyloarthropathies
• sarcoidosis
• gout
MTP (first)
• OA
• gout
• RA
Other MTPs
• RA
• gout
Ankle
• RA
• spondyloarthropathies (Reiter's)
• gout
PIP
• psoriasis
• Reiter's)

Figure 35.2 Joints typically affected by various arthropathies

Investigations

Table 35.3 (on p. 340) lists the many investigations that are used to reach a diagnosis.[7] Clinical acumen permits a judicious selection of specific tests rather than ordering an expensive battery.

Table 35.3 Investigations for arthritis

Appropriate tests can be selected from the following:
• urine analysis: blood, protein, sugar
• synovial fluid: analysis, culture
• radiology - plain X-ray
• blood and other cultures
• haemoglobin and differential WCC
• ESR
• C-reactive protein
• serum uric acid, creatinine
• 24 hour urinary uric acid
• rheumatoid factor

- anti-CCP (cyclic citrinullated peptide)antibody
- antinuclear antibody (screening test for SLE)
- dsDNA antibodies
- extractable nuclear antigen (ENA) antibodies
- HLA-B_{27} (poor predictive value)
- various specific serological tests (e.g. Australian epidemic polyarthritis, rubella, Lyme disease, hepatitis B, Barmah Forest virus)
- HIV serology
- antistreptolysin O titre
- streptococcal anti-DNAse B
- streptococcal AHT
- arthroscopy and biopsy
- bone scan

It is important to keep in mind the many specific serological tests to detect infective causes of arthralgia. These include Australian epidemic polyarthritis, Lyme disease, rubella, brucella, hepatitis B, gonococcus, mycoplasma, HIV tests, parvovirus and Barmah Forest virus.

In reference to viral serology, a positive immunoglobulin M (IgM) antibody test is presumptive evidence of recent infection and is likely to be of diagnostic significance in this clinical context. However, sometimes IgM antibodies can persist for months or years.[8] A positive IgG antibody result indicates previous exposure to the virus but a single positive titre is of no diagnostic significance. Seroconversion or at least a fourfold rise on paired sera confirms recent infection (see Figure 35.3 on p. 340).



Figure 35.3 Time course of 1gG amd 1gM antibodies in viral arthritis

Plain X-ray is invaluable, although in some conditions radiological changes may be apparent only when the disease is well established. Typical X-ray changes for common conditions are presented in Figure 35.4(on p. 340). Arthrography has limited value in the diagnosis of polyarthritis but is very useful for specific joints such as the shoulder and the knee. Ultrasound examination for joints such as the shoulder and the hip can be very useful.


Normal joint
Rheumatoid arthritis
cysts
joint destruction
subluxation
erosion joint margins
osteoporosis
Osteoarthritis
periarticular bone sclerosis
marginal osteophytes
loss of joint space
cysts
Gout
punched out erosions
cyst

Figure 35.4 X-rays for common arthritic conditions: typical changes

HLA-B_{27} should not be used for arthritis screening. It has a high sensitivity for ankylosing spondylitis, but low specificity, and should rarely be ordered.[8]

The various immunological tests for diagnosis of the connective tissue disorders are outlined with the description of each condition. Such screening tests include:

- rheumatoid factor
- antinuclear antibodies
- dsDNA antibodies

The LE cell test has been superseded by the antinuclear, dsDNA and ENA (especially Sm) antibody tests but the latter should only be performed if there is an elevated ANA test.[8]

ARTHRITIS IN CHILDREN

Arthralgia (joint pain) is a common problem in childhood and, although arthritis is rare, the complaint demands considerable respect because of the many serious problems causing it. Arthritis may be part of an infectious disease such as rheumatic fever, rubella, mumps, varicella cytomegalovirus infection, erythema in-fectiosum (human parvovirus), influenza or other viral infection, and is occasionally encountered with Henoch-Schönlein purpura. Actually, viral arthritis is very common in children. A FBE is helpful as it may show lymphopaenia, lymphocytosis or a typical lymphocyctes. It is worth noting that underlying bone tumours can be present as joint pain if the tumour is adjacent to the joint. A checklist of causes is presented in Table 35.4.

Table 35.4 Arthritis in children: causes to consider

Infections
Rheumatic fever
Septic arthritis
Meningococcaemia
Osteomyelitis
Reactive arthritis (post-infectious)
Tuberculosis
Viral infections (e.g. rubella, HIV)
Inflammation - chronic arthritis
Juvenile chronic arthritis
• oligoarthritis
• polyarticular (juvenile rheumatoid arthritis)
• systemic disease (Still's syndrome)
• juvenile ankylosing spondylitis
Psoriatic arthritis
Inflammatory bowel arthritis
Haematological disorders
Thalassaemia
Sickle-cell anaemia
Haemophilia
Neoplasms
Leukaemia
Lymphoma
Neuroblastoma
Orthopaedic conditions
Perthes' disorder
Slipped upper femoral epiphyses
Chondromalacia
Others
Henoch-Schönlein syndrome
Kawasaki's syndrome
Scurvy
Traumatic arthritis
Osteochondritis
Psychogenic rheumatism
Malignant tumour
• bone
• cartilage
• synovium

Note: Acute-onset monoarticular arthritis associated with fever is septic until proven otherwise.

Juvenile idiopathic arthritis (JIA)

JIA, also known as juvenile chronic arthritis and juvenile rheumatoid arthritis (USA) is defined as a chronic arthritis persisting for a minimum of 6 weeks (some criteria suggest 3 months) in one or more joints in a child less than 16 years of age.[5] It is rare, affecting only about 1 in 1000 children, but produces profound medical and psychosocial problems.

The commonest types of JIA arthritis are oligoarticular arthritis (pauciarticular), affecting four or fewer joints (about 50%), and polyarticular arthritis, affecting five or more joints (about 40%). Systemic onset arthritis, previously known as Still's syndrome, accounts for about 10% of cases. It is usually seen in children under the age of 5 but can occur throughout childhood. The children present with a high remittent fever and coppery red rash, plus other features, including lymphadenopathy, splenomegaly and pericarditis. Arthritis is not an initial feature but develops ultimately, usually involving the small joints of the hands, wrists, knees, ankles and metatarsophalangeal joints.

These children should be referred once the problem is suspected or recognised. JIA is not a benign disease - 50% have persistent active disease as adults.

Rheumatic fever

Rheumatic fever typically occurs in children and young adults, the first attack usually occurring between 5 and 15 years of age.

Arthritis in perspective

Five per cent of all children complain of recurrent lower limb pain, which often awakens them from their sleep. There may be emotional factors involved and parents need appropriate reassurance. A careful history and physical examination are essential, and perhaps simple basic investigations may be appropriate. As Rudge[5] points out, we have to be vigilant against underdiagnosis, misdiagnosis and overdiagnosis. Refer to growing pains (p. 878).

ARTHRITIS IN THE ELDERLY

OA is very common with advancing age and for this reason care has to be taken not to simply attribute other causes of arthritis to OA. Other musculoskeletal conditions that become more prevalent with increasing age are:

- polymyalgia rheumatica
- Paget's disease of bone
- avascular necrosis
- gout
- pseudogout (pyrophosphate arthropathy)
- malignancy (e.g. bronchogenic carcinoma)

Pseudogout

This crystal deposition arthropathy (chondrocalcinosis) is noted by its occurrence in people over 60 years. It usually affects the knee joint but can involve other joints.

Rheumatoid arthritis

Although it usually begins between the ages of 30 and 40 it can occur in elderly patients, when it occasionally begins suddenly and dramatically. This is called ‘explosive’ RA and fortunately tends to respond to small doses of prednisolone and has a good prognosis.[9] RA in the elderly can present as a polymyalgia rheumatica syndrome.

RHEUMATIC FEVER

RF is an inflammatory disorder that typically occurs in children and young adults following a group A *Streptococcus pyogenes* infection. It is common in developing countries and Indigenous Australians (p. 1429) but uncommon in first world countries.[10]

Clinical features

- Young person 5 - 15 years (can be older)
- Acute-onset fever, joint pains, malaise
- Flitting arthralgia mainly in leg (knees, ankles) and elbows and wrists of the arm
- One joint settles as the other is affected

- May follow a sore thoat

However, the symptoms depend on the organs affected and arthritis may be absent.

Diagnosis

Based on clinical criteria

2 or more major criteria or 1 major + 2 or more minor criteria

in the presence of supporting evidence of preceding streptococcal infection.

Major criteria

- Carditis
- Polyarthritis
- Chorea (involuntary abnormal movements)
- Subcutaneous nodules
- Erythema marginatum

Minor criteria

- Fever (≥ 38°C)
- Previous RF or rheumatic heart disease
- Arthralgia
- Raised ESR > 30mm/hr or CRP > 30 mg/L
- ECG - prolonged PR intervals

Investigations

A selective combination of:

- FBC
- throat swab
- ESR
- streptococcal ASOT
- streptococcal anti-DNAse B (repeat in 10 - 14 days)
- streptococcal AHT
- C-reactive protein
  Plus ECG and CXR.

Treatment

- Rest in bed
- Benzathine pencillin 900 mg IM (450 mg in child < 20 kg) statim or phenoxymethyl penicillin 500 mg (o) bd, 10 days
- Paracetamol
- Diuretics for carditis (may be ACE inhibitor and corticosteroids)
- Prophylactic long-term penicillin

OSTEOARTHRITIS

OA is the most common type of arthritis, occurring in about 10% of the adult population and in 50% of those aged over 60.[10] It is a degenerative disease of cartilage and may be primary or secondary to causes such as trauma and mechanical problems, septic arthritis, crystalopathy or previous inflammatory disorders, or structural disorders such as SCFE and Perthes' disorder.

The arthritis

Primary OA is usually symmetrical and can affect many joints. Unlike other inflammatory disease the pain is worse on initiating movement and loading the joint, and eased by rest. OA is usually associated with stiffness, especially after activity, in contrast to RA.

Joints involved

In primary OA all the synovial joints may be involved, but the main ones are:

- first carpometacarpal (CMC) joint of thumb
- first metatarsophalangeal (MTP) joint of great toe
- distal interphalangeal (DIP) joints of hands

Other joints that are affected significantly are the proximal interphalangeal joints, the knees, hips, acromioclavicular joints and joints of the spine, especially the facet joints of the cervical (C5 - 6, C6 - 7) and lumbar regions (L3 - 4, L4 - 5, L5 - S1) (Figure 35.5).


Most common sites
Other common sites
cervical spine
acromioclavicular joints
lumbar spine
hip joints
first CMC joint
DIP joints
PIP joints
knees
first MTP joint
great toe

Figure 35.5 Osteoarthritis: typical joint distribution

Clinical features

- Pain:
  - worse by the end of the day
  - aggravated by use
  - relieved by rest
  - worse in cold and damp
- Variable morning stiffness
- Variable disability

Signs

(See Figure 35.6)


radial deviation of distal phalanx
Heberden's node
Bouchard's node
metacarpophalangeal joints uninvolved

Figure 35.6 Typical clinical features of osteoarthritis of the hand

- Hard and bony swelling
- Crepitus
- Signs of inflammation (mild)
- Restricted movements
- Joint deformity

Note: There should be no systemic mani-festations.

Crystal arthropathy can complicate OA, especially in the fingers of people taking diuretics (e.g. nodular gout).

Differentiation from an inflammatory arthropathy[11]

OA does not exhibit the typical inflammatory pattern. The clinical diagnosis is based on:

- gradual onset of pain after activity (worse towards the end of the day)
- the pattern of joint involvement
- the lack of soft tissue swelling
- the transient nature of the joint stiffness or gelling
- takes < 30 minutes to settle after vest while inflammatory arthritis takes at least 30 minutes

Diagnosis

The diagnosis is clinical and radiological but the degree of changes on X-ray do not always parallel levels of symptoms.[10]

X-ray findings

- Joint space narrowing with sclerosis of subchondral bone
- Formation of osteophytes on the joint margins or in ligamentous attachments

- Cystic areas in the subchondral bone
- Altered shape of bone ends

Principles of management

- Provide explanation and reassurance including patient education handouts.
- Control pain and maintain function with appropriate drugs.
- Suggest judicious activity, exercise and physical therapy.
- Consider factors lowering the coping threshold (e.g. stress, depression, anxiety, overactivity).
- Refer for surgical intervention for debilitating and intractable pain or disability. Examples include OA of hip, knee, shoulder, first CMC joint of thumb, and first MTP joint where surgery is now very successful. Osteotomies still have a limited place for a varus or valgus deformity of the knee.

Optimal treatment

- *Explanation*. Provide patient education and reassurance that arthritis is not the crippling disease perceived by most patients.
- *Exercise*. A graduated exercise program is essential to maintain joint function. Aim for a good balance of relative rest with sensible exercise. It is necessary to stop or modify any exercise or activity that increases the pain.
  Systematic reviews have found that both exercise and education may help reduce the pain and disability in people with OA of the hip or knee.[12]
- *Rest*. Rest during an active bout of inflammatory activity only; prolonged bed rest contraindicated.
- *Heat*. Recommended is a hot-water bottle, warm bath or electric blanket to soothe pain and stiffness. Advise against getting too cold.
- *Diet*. If overweight it is important to reduce weight to ideal level. Obesity increases the risk of OA of the knee approximately fourfold and weight loss may slow progression;[13] otherwise, no specific diet has been proven to cause or improve OA. Some people claim that their arthritis is improved by having a nutritious balanced diet consisting of fish, rice and vegetables and avoiding meat, dairy produce, alcohol, pepper and spices.
- *Correction of predisposing factors and aids*. Apart from weight reduction the following may help:
  - walking stick
  - heel raise for leg length disparity
  - back brace
  - elastic or hinged joint support (e.g. knee)
- *Physiotherapy*. Referral should be made for specific purposes such as:
  - correct posture and/or leg length disparity
  - supervision of a hydrotherapy program
  - heat therapy and advice on simple home heat measures
  - teaching and supervision of isometric strengthening
  - exercises (e.g. for the neck, back, quadriceps muscle)
- *Occupational therapy*. Refer for advice on aids in the home, more efficient performance of daily living activities, protection of joints, and on the wide range of inexpensive equipment and tools.
- *Simple analgesics* (regularly for pain). Use paracetamol/acetaminophen (avoid codeine or dextropropoxyphene preparations, and aspirin if recent history of dyspepsia or peptic ulceration). Take before activity.
  Systematic reviews found that NSAIDs and simple analgesics reduce the pain of OA but there is no good evidence that NSAIDs are superior to simple analgesics (e.g. paracetamol) or that any one of the many NSAIDs is more effective than others.[12]
- *NSAIDs and aspirin*. These are the first-line drugs for more persistent pain or where there is evidence of inflammation, such as pain worse with resting and nocturnal pain. The risk versus benefit equation always has to be weighed carefully. As a rule, NSAIDs should be avoided if possible. Significant risks of NSAIDs are:
  - gastric erosion with bleeding
  - gastric ulceration
  - depression of renal function (check renal function beforehand)
  - hepatotoxicity

  Note: Change to a suppository form will not necessarily render the upper GIT safe from irritation.
- *COX-2 specific inhibitors (CSIs)*. Refer to Chapter 12, p. 104. These new generation agents, represented mainly by celecoxib and rofecoxib, should be considered where there is an indication for an NSAID but when the risk of NSAID-induced ulceration and bleeding is high.[14] Evidence shows that they have a similar efficacy to other NSAIDs and a modest absolute reduction in GIT complications.[15] The usual precautions need monitoring
- *Intra-articular (IA) corticosteroids*. As a rule IA corticosteroids are not recommended but occasionally can be very effective for an

inflammatory episode of distressing pain and disability on a background of tolerant pain (e.g. a flare-up in an osteoarthritic knee).

- *Viscosupplementation*. Intra-articular hylans, especially for OA of knee. Supported by level 1 evidence.
- *Complementary therapy*. Glucosamine, a natural amine sugar, derived from chitin in shellfish shell, has had anecdotal claims of efficacy for the treatment of OA. However, level 2 evidence has shown a significant benefit in its management.[16] The studies are promising so far but further studies are desirable. It appears to be more effective in the postmenopausal female. The dose is 1500 - 2000 mg (0) daily in divided doses, with food. It is well tolerated. (It is contraindicated in those with a significant seafood allergy).
- Epidemiological studies (level 3 evidence) suggest that OA patients whose diet is rich in antioxidants, such as vitamin C, vitamin D and green tea, have slower progression of joint space narrowing on X-ray over long-term follow-up.[17]
- *Contraindicated drugs*. For OA these include the immunosuppressive and disease-modifying drugs such as oral corticosteroids, gold, anti-malarials and cytotoxic agents.

RHEUMATOID ARTHRITIS

RA is the commonest chronic inflammatory polyarthritis and affects about 1 - 2% of the population. The disorder can vary from a mild to a most severe debilitating expression. About 10 - 20% of patients have a relentless pro-gression and require aggressive drug therapy.[18]

Genetic factors may represent a risk of 15 - 70% of developing RA.

The arthritis

RA generally presents with the insidious onset of pain and stiffness of the small joints of the hands and feet. The pain is persistent rather than fleeting and mainly affects the fingers where symmetrical involvement of the PIP joints produces spindling while the meta-carpophalangeal joints develop diffuse thickening as does the wrist (Figure 35.7). In 25% of cases RA presents as arthritis of a single joint such as the knee,[9] a situation leading to confusion with Lyme disease or a spondyloarthropathy.



Figure 35.7 Chronic rheumatoid arthritis showing classic features of deformities including subluxation of joints and rheumatoid nodules

Joints involved

- Hands: MCP and PIP joints, DIP joints (30%)
- Wrist and elbows
- Feet: MTP joints, tarsal joints (not IP joints), ankle
- Knees (common) and hip (delayed - up to 50%)
- Shoulder (glenohumeral) joints
- Temporomandibular joints
- Cervical spine (not lumbar spine)

Refer to Figure 35.8.


Most common sites
Other sites
temporomandibular joints
cervical spine
shoulder joints
elbow joints
hip (delayed)
wrist
DIP joints
MCP joints
PIP joints
knees
ankles
tarsal joints
MTP joints

Figure 35.8 Rheumatoid arthritis: typical joint distribution

Clinical features

- Insidious onset but can begin acutely (explosive RA)
- Any age 10 - 75 years - peak 30 - 50 years but bimodal 25 - 50 (peak age) and 65 - 75
- Female to male ratio ° 3:1
- Joint pain
  - worse on waking, nocturnal pain, disturbed sleep
  - relieved with activity
- Morning stiffness - can last hours
- Rest stiffness (e.g. after sitting)
- General: malaise, weakness, weight loss, fatigue
- Disability according to involvement

Signs

- Soft swelling (effusion and synovial swelling), especially of wrist, MCP and PIP joints
- Warmth
- Tenderness on pressure or movement
- Limitation of movement

- Muscle wasting
- Later stages: deformity, subluxation, instability or ankylosing
- Look for swan necking, boutonniere and two Z deformities, ulnar deviation (Figure 35.9)


ulnar deviation
swollen, thickened
rheumatoid nodule
z deformity
muscular atrophy
boutonniere deformity
swan neck deformity

Figure 35.9 Chronic rheumatoid arthritis: typical signs

- Check for a number of everyday functions, e.g.:
  - power grip (lifting a jug of water)
  - precision grip (using a key or pen) undoing
  - hook grip (carrying a bag)

The various possible extra-articular mani-festations are summarised in Figure 35.10.


anaemia
inflammation of eye
Sjögren's syndrome
• dry eyes
• dry mouth
lymphadenopathy
Pulmonary
• pleural effusion
• fibrosing alveolitis
• nodules (Caplan's syndrome)
amyloidosis (kidneys)
Vasculitis
→ purpura
nail-fold infarcts
skin ulcers
Cervical spine
• atlantoaxial subluxation
Cardiac
• pericarditis
• myocarditis
splenomegaly (Felty's syndrome)
subcutaneous nodules
bursitis
tenosynovitis
carpal tunnel syndrome
Raynaud's phenomenon
Baker's cyst (popliteal fossa)
peripheral sensory neuropathy
mononeuritis multiplex

Figure 35.10 Rheumatoid arthritis: significant non-articular clinical manifestations

Investigations

- ESR usually raised according to active disease
- Anaemia (normochronic and normocytic) may be present
- Rheumatoid factor
  - positive in about 70 - 80%(less frequent in early disease)
  - 15 - 25% of RA patients will remain negative[18]
- Anti-cyclic citrullinated peptide (anti-CCP) antibodies: more specific for RA (96% specificity)
- X-ray changes:
  - erosion of joint margin: ‘mouse-bitten’ appearance
  - loss of joint space (may be destruction)
  - juxta-articular osteoporosis
  - cysts
  - advanced: subluxation or ankylosing
- MRI - helpful for early diagnosis

Key point

If the RA factor is positive, it is non-specific - order the anti-CCP antibody to confirm the diagnosis.

Criteria for the diagnosis of RA are presented in Table 35.5.

Table 35.5 Reviced C criteria for the diagnosis of rheumatoid arthritis

Symptom duration of > 6 weeks
Early morning stiffness of > 1 hour
Arthritis in three or more regions
Bilateral compression tenderness of the metatarsopahalangeal joints
Symmetry of the areas affected
Rheumatoid factor positivity
Anti-cyclic citrinullated peptide antibody positivity
Bony erosions evident on radiographs of the hands or feet, although these are uncommon in early disease

Management principles[10,19]

- Give patient education support and appropriate reassurance. The diagnosis generally has distressful implications, and so the patient and family require careful explanation and support. Some patients have little or no long-term problems but even in mild cases, continuing care and medical supervision is important.
- There has been a radical shift from palliation to early induction of disease remission, to prevent joint damage and reduce morbidly from malignancy (especially lymphouse) and cardiovascular disease.
- Since many studies show disease progression in the first 2 years, relative aggressive treatment with disease - modifying antirheumatic drugs (DMARDs) from the outset is advisable, rather than to start stepwise with analgesics and NSAIDs only.[20]
- Use a team approach where appropriate, including a consultant referral for diagnosis and collaborative support.
- Fully assess the patient's functional impairment and impact on home life, work and social activity. Involve the family in decision making.
- Make judicious use of pharmaceutical agents. For serious cases consultant collaboration is essential.
- Review the patient regularly, continually assessing progress and drug tolerance.

Specific advice

- *Rest and splinting*. This is necessary where practical for any acute flare-up of arthritis.
- *Exercise*. It is important to have regular exercise, especially walking and swimming. Have hydrotherapy in heated pools.
- *Referral*. Referring to physiotherapists and occupational therapists for expertise in exercise supervision, physical therapy and advice regarding coping in the home and work is important.
- *Joint movement*. Each affected joint should be put daily through a full range of motion to keep it mobile and reduce stiffness.
- *Diet*. Although there is no special diet that seems to cause or cure RA there is evidence that avoiding animal fats (dairy products and some meats) and using fish oils is beneficial.[21] A nourishing well-balanced diet is common sense and obesity must be avoided.

Therapies used in the management of rheumatoid arthritis are presented in Table 35.6.

Table 35.6 Therapies used in the management of rheumatoid arthritis[21]

• Education (rest, literature, weight loss, joint protection advice)
• NSAIDs
• Simple analgesics
• DMARDs:
Immunosuppressants
- azathioprine
- cyclosporin
- leftunamide
- methotrexate
cyto-kinase inhibitors (biological DMARDs)
- adalimumab anti TNF α agents
- entanercept anti TNF α agents

  - infliximab anti TNF α agents
  - anakinra (anti-Interlukin-1 agent)

  Gold salts

  Quinolones
  - hydroxychloroquine
  - chloroquine

  others
  - D-penicillamine
  - sulfasalazine
- Physical therapy (hydrotherapy, isometric exercises)
- Occupational therapy (splints, aids and appliances)
- Corticosteroids:

  oral prednisolone

  intra-articular

  intravenous (steroid 'pulses')
- Fiish body oil
- Orthopaedic surgery:

  (synovectomy, joint replacement, arthrodesis, plastic hand surgery)
- Chiropody, footwear, insoles

Source: After Reilly and Littlejohn [21]

Pharmacological management[10]

NSAIDs

These are effective but the adverse effects are a problem.

The use of paracetamol, fish oils and low-dose glucocorticoids may allow NSAID use to be kept to a minimum or avoided.

Glucocorticoids

Oval use should be considered in patients with severe disease and where other treatments have failed or are contraindicated.

The dose is prednisolone 5 - 10 mg (o) daily. Avoid doses higher than 15mg daily if possible.

Intra-articular infections of depot preparations are effective in larger joints.

Disease modifying antirheumatic drugs (DMARDs)

These agents target synovial inflammation and prevent joint damage. The choice depends on several factors, but is best left to the specialist co-ordinating care. In most patients with recently diagnosed RA, methotrexate is the cornerstone of management and should be commenced as early as possible.

Initial dose: methotrexate 5 - 10 mg (0) weekly, increasing to maximum of 25mg weekly depending on clinical response and toxicity. Add folic acid 5 mg tds.

Biological DMARDs (bDMARDs) are the never agents which should be considered if remission is not achieved with appropriate methotrexate monotherapy, 'triple therapy' or other combinations. All bDMARDs are more effective when combined with methotrexate.

Warning: All practitioners should be aware of the increased risk of infectioners disease such as the atypical pneumomas and listerosis while taking bDMARDs. All patients should report unusual fever or symptoms. Injection site reactions are common.

Standard Initial drug therapy

Monotherapy with methotrexate (or occasionally another DMARD) is standard. Less than 20% will reach disease remission and, if not achieved, increase the dose or consider combination therapy.

Combination therapy

Counder standard triple therapy:

- Methotrexate + sulfasalazise + hydroxychloroquine

Triple therapy can be used if has methotrexate monotherapy or initially on diagnosis, dependent on the severly of the disease. Monitoring for FBE, LFTs and annual eye checks is necessary.

Several other double combination may be used, eg. Methotrexate with cyclosporin, leftunomide or a dDMARD.

CONNECTIVE TISSUE DISEASES

The three connective tissue disorders have the common feature of arthritis or arthralgia. Refer to Chapter 32.

Systemic lupus erythematosus

Arthritis is the commonest clinical feature of SLE (over 90%).[9] It is a symmetrical poly-arthritis involving mainly small and medium joints, especially the proximal interphalangeal and carpal joints of the hand. It is usually non-erosive and non-deforming, although deformities of fingers and thumbs can occur due to laxity of ligaments, tendons and capsules causing joint instability.

The initial presentation is similar to RA. The fibromyalgia syndrome can cause confusion although it has a distribution confined mainly to the trunk, especially the back.

Investigations

The two main screening tests are:

1. antinuclear antibodies - positive in 95%
2. dsDNA antibodies - specific for SLE but present in only 60%

Other screening tests are:

- ENA antibodies, especially Sm
- ESR - elevated in proportion to disease activity
- rheumatoid factor - positive in 50%
- antiphospholipid antibodies

Drug treatment

- Mild - NSAIDs/aspirin ± hydroxychloroquine
- Moderate - low-dose antimalarials (e.g. hydroxychloroquine or chloroquine)
- Moderate to severe - corticosteroids; immunosuppressive drugs (e.g. azathioprine, methotrexate)

Note: Referral for shared care is recommended.

Scleroderma

Scleroderma can present as a polyarthritis affecting the fingers of the hand in 25% of patients, especially in the early stages. Soft tissue swelling produces a ‘sausage finger’ pattern. Systemic sclerosis mainly affects the skin, presenting with Raynaud's phenomenon in over 85% of cases.

Investigations

- ESR may be raised
- Normocytic normochronic anaemia may be present
- Antinuclear antibodies - up to 90% positive
- Rheumatoid factor - positive in 30%
- Hypergammaglobulinaemia - 50% positive
- Antinucleolar and anticentromere antibodies - specific (see Table 32.3, pp. 284 and 287).

Polymyositis and dermatomyositis

Arthralgia and arthritis occur in about 50% of patients and may be the presenting feature before the major feature of muscle weakness and wasting of the proximal muscles of the shoulder and pelvic girdles appear. The small joints of the hand are usually affected and it may resemble RA.

CRYSTAL ARTHRITIS

Arthritis, which can be acute, chronic or asymptomatic, is caused by a variety of crystal deposits in joints. The three main types of crystal arthritis are monosodium urate (gout), calcium pyrophosphate dihydrate (CPPD) and calcium phosphate (usually hydroxyapatite).[22] Refer to Table 35.7.

Table 35.7 Crystal-induced disorders

Crystals	Associated disease/syndrome>	Typical joints or region affected
Monosodium urate	• Acute gout	Metatarsophalangeal joint of big toe
	• Tophaceous gout	Also: other foot joints ankle knee and patellar bursa wrist fingers
	• Asymptomatic	
	• Chronic gouty arthritis	
Calcium pyrophosphate dihydrate (CPPD)	• Acute pseudogout	Knee, wrist
	• Destructive arthropathy (like RA)	In older people > 60 years (average age 72)
	• Asymptomatic (most common)	F > M (2.7:1)
Basic calcium phosphate	• Acute calcific periarthritis	Shoulder (supraspinatus)
	• Destructive arthropathy	
	• Acute arthritis	

Gout (monosodium urate crystal disorder)

Gout is an abnormality of uric acid metabolism resulting in hyperuricaemia and urate crystal deposition. Urate crystals deposit in:

- joints - acute gouty arthritis
- soft tissue - tophi and tenosynovitis
- urinary tract - urate stones

Four typical stages of gout are recognised:

- *Stage 1* - asymptomatic hyperuricaemia
- *Stage 2* - acute gouty arthritis
- *Stage 3* - intercritical gout (intervals between attacks)
- *Stage 4* - chronic tophaceous gout and chronic gouty arthritis

Asymptomatic hyperuricaemia

- 10 times more common than gout[9]
- Elevated serum uric acid (> 0.42 mmol/L in men, > 0.36 mmol/L in women)
- Absence of clinical manifestations
- Usually does not warrant treatment

Typical clinical features of gout[10]

- mainly a disorder of men (5 - 8% prevalence)
- onset earlier in men (40 - 50) than women (60+)[6]
- Acute attack
  - excruciating pain in great toe (Figure 35.11)



Figure 35.11 Gout showing typical red, shiny, swollen arthritis of the first MTP joint with desquamation of the skin

  - early hours of morning
- Skin over joint - red, shiny, swollen and hot
- Exquisitely tender to touch
- May be precipitated by
  - alcohol excess (e.g. binge drinking)
  - surgical operation
  - starvation
  - drugs (e.g. frusemide, thiazide diuretics)
- Relief with colchicine, NSAIDs, corticosteroids
- Can subside spontaneously (3 to 10 days) without treatment

The arthritis

Monoarthritis in 90% of attacks:

- MTP joint great toe - 75%
- other joints - usually lower limbs
  - other toes
  - ankles
  - knees

Polyarticular onset is more common in old men and may occur in DIP and PIP joints of fingers. No synovial joint is immune.

Refer to Figure 35.12.


Common sites
Less common
DIP joints
knees
metatarsophalangeal joint–great toe
ankles
toes

Figure 35.12 Gout: possible joint distribution

Other features

- Prone to recurrence
- Tophi in ears, elbows (olecranon bursa), big toes, fingers, Achilles tendon (take many years)
- Can cause patellar bursitis
- Can get cellulites (does not respond to antibiotics)

Nodular gout

Develops in postmenopausal women with renal impairment taking diuretic therapy who develop pain and tophaceous deposits around osteoarthritic interphalangeal (especially DIP) joints of fingers.[23]

Diagnosis

- Synovial fluid aspirate → typical uric acid crystals using compensated polarised microscopy; this should be tried first as it is the only real diagnostic feature
- Elevated serum uric acid (up to 30% can be within normal limits with a true acute attack)[22]
- X-ray: punched out erosions at joint margins

Management

Management of gout includes these principles:

- Good advice and patient education information
- Provision of rapid pain relief
- Preventing further attacks
- Prevention of destructive arthritis and tophi
- Dealing with co-morbid conditions, eg. Alcohol dependence, obesity

The acute attack[10]

NSAIDs, If full dosage, are first-line and effective.

- Indomethacin 50 mg (o) tds (if tolerated) until symptoms abate (up to 3 - 5 days), then tape to 25mg tds until creation of the attack

If extreme: indomethacin 100mg (o) statim, 75mg 2 hours later (if tolerated), then 50 mg (o) tds for 24 - 48 hours, then 50 - 75 mg/day.

Note: Any other NSAID can be used. Add on antiemetic (e.g. metoclopramide 10 mg (o) tds) Or

- Colchicine 0.5 mg (o) statim, then 0.5 mg every 6 or 8 hours until pain relief (usually 24 - 28 hours) or diarrhoea develops (max. 6mg/24 hours)

Note:

- Must be given early.
- Avoid if renal impairment.

Consider:

- corticosteroids: intra-articular following aspiration and culture (gout and sepsis can occur together); a digital anaesthetic block is advisable. An oral course can be used: start with prednisolone 40 mg/day for 4 days then decrease gradually over 10 days
- corticotrophin (ACTH) IM in difficult cases (e.g. synthetic ACTH: tetracosactrin 1 mg IM)

Note:

- Avoid aspirin.
- Monitor renal function and electrolytes.

Long-term therapy

When acute attack subsides preventive measures include:

- weight reduction
- a normal, well-balanced diet
- avoidance of purine-rich food, such as organ meats (liver, brain, kidneys, sweetbread), tinned fish (sardines, anchovies, herrings), shellfish and game
- reduced intake of alcohol
- good fluid intake (e.g. water - 2 litres a day)
- avoidance of drugs such as diuretics (thiazides, frusemide) and salicylates
- wearing comfortable shoes

Drug prophylaxis

Allopurinol (a xanthine oxidase inhibitor) is the drug of choice.

Dose: 100 - 300 mg daily

Indications:

- frequent acute attacks
- tophi or chronic gouty arthritis
- renal stones or uric acid nephropathy
- hyperuricaemia

Adverse effects:

- rash (2%)
- severe allergic reaction (rare)

Beware of renal insufficiency and elderly patients - use lower doses.

Beware of drug interactions:

- azathioprine - potentially lethal
- amoxycillin - prone to rashes

Method: treatment of intercritical gout

- Commence 6 weeks after last acute attack.
- Start with 50 mg daily for the first week and increase by 50 mg weekly to maximum 300 mg.
- Check uric acid level after 4 weeks: aim for level < 0.38 mmol/L
- Add colchicine 0.5 mg bd for 6 months (to avoid precipitation of gout) or indomethacin 25 mg bd or other NSAIDs.

Probenecid (uricosuric agent)

Good for hypoexcretion of uric acid.

Dose: 500 mg/day (up to 2 g)

Note: Aspirin antagonises effect.

Calcium pyrophosphate crystal disorder (pseudogout)[10,24]

The finding of calcification of articular cartilage on x-ray examination is usually termed chondrocalcinosis. This is mainly a disorder of the elderly superimposed on an osteoarthritic joint. The acute attack is similar to an acute attack of gout but it affects the following joints (in order):

- Knee
- 2nd and 3rd MCP joints
- wrist
- shoulder
- ankle
- elbow

It can affect tendons, especially Auxiliary tendon.

The crystals in synovial fluid are readily identified by phase-contrast microscopy. X-rays are helpful in showing calcification of the articular cartilage.

Management is based on aspiration and installation of a depot glucocorticosteroid by injection into the joint (if joint infection excluded) plus analgesia. Be cautious of using NSAIDs in the elderly - paracetamol is preferred. Colchicine can be used.

Other treatment includes:

- Indomethacin 50 mg (o) tds (if tolerated) until symptoms abate *and/or*
- colchicine 0.5 mg (0) tds until attack subsides *and*
- paracetamol 500 - 1000 mg (o) four times daily, if necessary

THE SPONDYLOARTHROPATHIES

The spondyloarthropathies are a group of related inflammatory arthopathies with common characteristics affecting the spondyles (vertebrae) of the spine. It is appropriate to regard them as synonymous with the seronegative spondyloarthropathies in contradistinction to RA, which is seropositive and affects the cervical spine only. Apart from back pain this group tends to present with oligoarthropathy in younger patients. The arthritis is characteristically peripheral, asymmetrical, affects the lower limbs and can exhibit dactylitis (e.g. ‘sausage’ digits).

Features[25]

- Sacroiliitis with or without spondylitis
- Enthesopathy, especially plantar fasciitis, Achilles tendinitis, costochondritis
- Arthritis, especially larger lower limb joints
- Extra-articular features (e.g. iritis, anterior uveitis, mucocutaneous lesions, psoriasiform skin and nail lesions, chronic 61TKGU inflammation)

- Absent rheumatoid factor
- Association with HLA-B_{27} antigen
- Familial predisposition

The group of disorders

1. Ankylosing spondylitis
2. Reiter's syndrome/reactive arthritis
3. Inflammatory bowel disease (enteropathic arthritis)
4. Psoriatic arthritis
5. Juvenile onset ankylosing spondylitis
6. Unclassified spondyloarthritis - partial features only

Ankylosing spondylitis

This usually presents with inflammatory back pain (sacroiliac joints and spine) and stiffness in young adults, and 20% present with peripheral joint involvement before the onset of back pain. It usually affects the girdle joints (hips and shoulders), knees or ankles. At some stage over 35% have joints other than the spine affected.

Key clinical features

- Insidious onset of discomfort
- Age less than 40 years
- Persistence for > 3 months
- Associated morning stiffness
- Improvement with exercise or physical activity
- Symptoms respond to NSAIDs

Reactive arthritis (Reiter's syndrome)

This is a form of reactive arthropathy in which non-septic arthritis and often sacroiliitis develop after an acute infection with specific venereal or dysenteric organisms.

DxT: NSU + conjunctivitis ± iritis + arthritis ° Reiter's syndrome

DxT: similar syndrome without ocular or mucocutaneous lesions ° reactive arthritis

The arthritis tends to affect the larger peripheral joints, especially the ankle (talocrural) and knees but the fingers and toes can be affected in a patchy polyarthritic fashion. Mucocutaneous lesions, including keratoderma blennorrhagia and circinate balanitis, may occur, although the majority develop peripheral arthritis only (see Figure 35.13).


conjunctivitis
uveitis
mouth ulcers
aortitis
pericarditis
rare
spondylitis
sarcroiliitis
urethritis
circinate balanitis
arthritis
Achilles tendonitis
plantar fasciitis
keratoderma blennorrhagia

Figure 35.13 Possible clinical features of Reiter's syndrome

Enteropathic arthritis

Inflammatory bowel disease (ulcerative colitis, Crohn's disease and Whipple's disorder) may rarely be associated with peripheral arthritis and sacroiliitis.

Psoriatic arthritis

Like Reiter's syndrome, this can develop a condition indistinguishable from ankylosing spondylitis. It is therefore important to look beyond the skin condition of psoriasis, for about 5% will develop psoriatic arthropathy. It can have several manifestations:

1. mainly DIP joints
2. identical RA pattern but RA factor negative
3. identical ankylosing spondylosis pattern with sacroiliitis and spondylitis
4. monoarthritis, especially knees
5. severe deformity or ‘mutilans’ arthritis

Unclassified spondyloarthritis

Patients in this category seem to be the most frequently encountered in family practice. They clearly have a spondyloarthropathy but fail to meet the criteria for any one of the individual entities within the group. A typical patient is a young male in his third decade with a painful knee or other joint, unilateral (or bilateral) back pain with one of the entheseal problems (e.g. plantar fasciitis).

Investigations

- X-rays
  - Radiological sacroiliitis is central to the diagnosis.
  - Changes include narrowing of SIJs, margin irregularity, sclerosis of periarticular bone and eventually bony fusion. Spondylitis usually follows.
- ESR and CRP
  - Most patients have an elevated ESR and CRP at some stage of their disease.
- HLA-B_{27}
  - This test has low specificity and has limited value except that it predicts risk to offspring if positive.
- Microbiology
  - In patients with a history of reactive arthritis cultures should be obtained from the urethra, faeces, urine and blood.[26]

MANAGEMENT PRINCIPLES

- Identify the most active elements of the disease and treat accordingly.
- Provide patient and family education with appropriate reassurance: this is vital. Stress that, although the disease is non-curable, treatment is effective and long-term prognosis generally good.
- Provide regular assessment and support.
- Give genetic counselling - in cases of ankylosing spondylitis with positive HLA-B_{27} the risk to offspring is significant.
- Give advice regarding work, especially with posture.
- Acute anterior uvertis requires prompt treatment and monitoring by an ophthalmolog.
- Refer for physiotherapy for exercises, postural exercises and hydrotherapy. Appropriate physiotherapy slows deterioration in spinal function.[27]
- Consider referral for occupational therapy.
- Pharmacological agents:
  - NSAIDs (e.g. indomethacin 75 - 200 mg or 100 mg rectallynocle daily to control pain, stiffness and synovitis)
  - sulphasalazine (if NSAIDs ineffective)
  - intra-articular corticosteroids for severe monoarthritis and intralesional corticosteroids for enthesopathy

Cautions

- Careful monitoring is required with NSAIDs and sulphasalazine.
- Systemic corticosteroids are not indicated.
- Immunosuppressants (low dose weekly methotrexate) and bDMARDs may be needed for severe intractable problems with psoriasis and reactive arthritis.
- These conditions should be managed in collaboration with a consultant.
- Although phenylbutazone is the most effective NSAID, its side-effects (especially aplastic anaemia) are a major problem.

THE VASCULITIDES

The vasculitides or vasculitis syndromes are a heterogeneous group of disorders involving inflammation and necrosis of blood vessels, the clinical effects and classification depending on the size of the vessels involved.

More common causes

These are the small vessel vasculitis effects associated with many important disorders such as RA, SLE, infective endocarditis, Henoch-Schönlein purpura and hepatitis B. Skin lesions and arthritis are usually associated with these disorders.

Rarer causes

The major vasculitides are polyarteritis nodosa, giant cell arteritis, polymyalgia rheumatica and Wegener's granulomatosis. Arthritis or limb girdle pain can be a component of the clinical presentation. The vasculitides are presented in more detail in Chapter 32.

WHEN TO REFER

- Consider referring most severe true inflammatory disorders for diagnosis and initiation of treatment (e.g. RA, spondyloarthropathy, connective tissue disorders and suspicion of a vasculitide)
- Osteoarthritis:
  - generalised joint pain
  - associated systemic symptoms
  - deteriorating joint function
  - intractable pain (especially at rest)
  - if surgical procedure is contemplated[10]
- Rheumatoid arthritis:
  - all patients initially
  - persistent inflammation of a joint or joints
  - patient ill and corticosteroids contemplated
  - if a surgical procedure is contemplated
- Spondyloarthropathies:
  - initial referral for confirmation of diagnosis and initiation of treatment
  - disease unresponsive to conventional treatment
  - sudden deterioration in symptoms, especially pain
  - onset of uveitis or other ocular complications
  - adverse drug reactions
- Undiagnosed arthritis in presence of constitutional symptoms
- Suspicion of a suppurative or serious infective condition (e.g. septic arthritis, endocarditis, brucellosis)
- Children with evidence of juvenile idiopathic arthritis (e.g. Still's syndrome)

Practice tips

- Morning stiffness and pain, improving with exercise ° RA.
- Flitting polyarthritis and fever ° rheumatic fever;? endocarditis;? SLE.
- Polyarthritis (usually PIPs) and rash ° viral arthritis or drug reaction.
- If rheumatoid arthritis involves the neck, beware of atlantoaxial subluxation and spinal cord compression.
- If the patient is young - think of SLE.
- If a patient returns from overseas with arthralgia, think of drug reactions, hepatitis, Lyme disease, but if the pain is intense consider dengue fever.
- Consider the possibility of Lyme disease in people with a fever, rash and arthritis who have been exposed to tick bites in rural areas.
- If a patient presents with Raynaud's phenomenon and arthritis, especially of the hands, consider foremost RA, SLE and systemic sclerosis.
- Avoid the temptation to apply on doubtful grounds a broad label such as arthritis or rheumatoid, or a precise diagnosis such as RA, and introduce drugs.[28] Table 35.8 presents the diagnostic guidelines.[29]

Table 35.8 Diagnostic guidelines for arthritis

Disorder	Sex ratio	Typical age	Typical common joints involved	Associated features
OA (generalised - primary)	FM 6:1	> 50	• DIP > PIP fingers • Base thumb (1st CMC) • 1st MTP joint • Cervical and lumbar spines • Hips and knees	• Pain worse in evenings, relieved by rest
RA	FM 3:1	30 - 50	• PIP, MCP hands	• Any joint: worse at rest, better with activity; morning

				stiffness
			• Wrist	
			• Base of toes (MTP joints)	• Constitutional symptoms
			• Symmetrical	• Carpal tunnel syndrome
				• Many other general effects
SLE	FM 9:1	15 - 35	• Symmetrical and variable	• Constitutional symptoms
			• Small joints fingers	• Fever
			• Often slight	• Adverse drug reactions
				• Any other system affected
				• Rash (80%)
				• Pleuritic symptoms (67%)
				• Raynaud's
Systemic sclerosis	FM 3:1	20 - 50	• Symmetrical	• Raynaud's (90%)
			• Polyarthritis fingers	• Other skin changes
				• Dysphagia
Viral arthropathies (excluding HIV)	M ° F	Children	• Transient	• Rash, fever
			• Usually PIP joints fingers	
Ankylosing spondylitis	MF 3:1	18 - 30	• Sacroiliacs	• Iridocyclitis
			• Vertebral column esp. lumbar, costovertebral	• Chest dysfunction
				• Enthesopathy (e.g. plantar fasciitis)
			• Also knees, hips or ankles	
Psoriatic arthritis	M ° F	Any age	• DIP joints - fingers and toes, sacroiliacs	• Psoriasis rash (pre-existing)
				• Pitted nails, ‘sausage digits’
Enteropathic arthritis	M ° F	Any age	• Lower extremity: knees, feet, ankles	• Ulcerative colitis
				• Crohn's disease
			• hips: sacroiliacs	
Reactive arthritis: Genitourinary (e.g. Reiter's) Post-dysentery (e.g. *Salmonella*)	MF 20:1	15 - 30	As above	• Preceding dysentery or urethritis
				• Entheseal problems
	M ° F			
Gout	MF 20:1	M 40 - 50 F > 60	• Big toe (MTP joint): any other possible, esp. lower limb	• Tophi
				• Raised serum uric acid
				• Urate crystals in joints
			• DIP - osteoarthritis	• Diuretics in elderly
Pseudogout	M ° F	> 60	• Knee	• Chondrocalcinosis

		especially > 70		• Pyrophosphate crystals in joint
Polymyalgia rheumatica	FM 3:1	> 60	• Morning stiffness and pain in girdles, esp. shoulder • Joints normal or osteoarthritic	ESR ↑↑↑

REFERENCES

1. Hart FD. *Practical Problems in Rheumatology*. London: Dunitz, 1985: 77.
2. Cormack J, Marinker M, Morrel D. *Practice: A Handbook of Primary Health Care*. London: Kluwer-Harrop Handbooks, 1980; 3(61): 1 - 12.
3. Lassere M, McGuigan L. Systemic disease presenting as arthritis - a diagnostic approach. Aust Fam Physician, 1991; 20: 1683 - 714.
4. Carroll GJ, Taylor AL. Drug-induced musculoskeletal syndromes. Curr Ther, 2000; Feb: 47 - 50.
5. Rudge S. Joint pain in children: assessing the serious causes. Mod Med Aust, 1990; May: 113 - 21.
6. Sandler G, Fry J. *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986.
7. Nash PT, Webb J. Polyarticular and monoarticular arthritis. A diagnostic approach. Aust Fam Physician, 1986; 15: 1265 - 75.
8. Barraclough D. Rheumatology symptoms: will investigation make a difference? Aust Fam Physician, 2001; 30(4); 322 - 6.
9. Kumar PJ, Clarke ML. *Clinical Medicine* (5th edn). London: Saunders, 2003: 538-540.
10. Mashford L (chair). *Therapeutic guidelines: Rheumatology* (Version 1). Melbourne: Therapeutic guidelines Ltd, 2006.
11. March LM. Osteoarthritis. Med J Aust, 1997; 166: 98 - 103.
12. Barton S ed. *Clinical Evidence*. London. BMJ Publishing Group, 2001: 808 - 18.
13. Felson DT. Weight and osteoarthritis. J Rheumatol Suppl, 1995; 43: 7 - 9.
14. Day R. COX-2 Specific inhibitors: should I prescribe them? Curr Ther, 2000; Feb: 9 - 11.
15. Osteoarthritis - have COX-2s changed its management. NPS News, 2001; 18: 1 - 6.
16. Reginster JY et al. A controlled trial of glucosamine for osteoarthritis of the knee. Lancet, 2001; 357: 251 - 6.
17. McAlindon J, Felson DT. Nutrition: risk factors for osteoarthritis. Ann Rheum Dis, 1997, 56: 397 - 442.
18. Shmerling RH, Delbanco TL. How useful is the rheumatoid factor? An analysis of sensitivity, specificity and predictive value. Arch Intern Med, 1992; 152: 2417 - 20.
19. Brooks P. Rheumatoid arthritis. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 446 - 9.
20. Ostor A, McColl G. What's new in rheumatoid arthritis. Aust Fam Physician, 2001; 30(4): 314 - 20.
21. Reilly P, Littlejohn G. Current treatment concepts in arthritis. Aust Fam Physician, 1989; 18: 1499 - 1509.
22. Hall S. Crystal arthritis: a clinician's view. Aust Fam Physician, 1991, 20: 1717 - 24.
23. MacFarlane DG, Dieppe PA. Diuretic induced gout in elderly women. Br J Rheumatol, 1985; 24: 155 - 9.
24. Tierney LM et al. *Current Medical Diagnosis and Treatment* (45th edn). New York: McGraw-Hill and Lange, 2006: 1458 - 9.
25. Creamer P, Edmonds J. Spondyloarthropathies. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 479 - 82.
26. Edmonds JP. Spondyloarthropathies. Med J Aust, 1997; 166: 214 - 18.
27. Vilitanen JV et al. Fifteen months follow up of intensive inpatient physiotherapy and exercise in ankylosing spondylitis. Clin Rheum, 1995; 14: 413 - 19.
28. Kincaid-Smith P, Larkins R, Whelan G eds. *Problems in Clinical Medicine*. Sydney: MacLennan & Petty, 1989: 391.
29. Hart FD. Early clinical diagnosis of 12 forms of arthritis. Mod Med Aust, 1989: March: 34 - 40.

CHAPTER 36. ANORECTAL DISORDERS

Duncan ill with very bad piles - operated on last night, or, since that sounds alarming, lanced.
Can't really sympathise with that particular disease, though the pain is terrible. Must laugh.

VIRGINIA WOOLF 1934, DIARY ENTRY

ANORECTAL DISORDERS

Anorectal problems are common in family practice and tend to cause anxiety in the patient that is often related to the fear of cancer. This fear may be well foundedfor many instances of rectal bleeding and lumps. It is important to keep in mind the association between haem-orrhoids and carcinoma of the large bowel.

Anorectal problems include:

- pain
- lumps
- discharge
- bleeding
- pruritus

Common anorectal conditions are illustrated in Figure 36.1.


rectal polyp
internal sphincter
external sphincter
ischiorectal abscess
dentate line
perianal abscess
ischiorectal fistula
anal fissure
perianal haematoma
prolapsed (4th-degree) haemorrhoid

Figure 36.1 Common anorectal conditions

ANORECTAL PAIN

The patient may complain that defecation is painful or almost impossible because of anorectal pain.

Summary of causes

Pain without swelling:

- anal fissure

- anal herpes
- ulcerative proctitis
- proctalgia fugax
- solitary rectal ulcer
- tenesmus

Painful swelling:

- perianal haematoma
- strangulated internal haemorrhoids
- abscess
  - perianal
  - ischiorectal
- pilonidal sinus

ANAL FISSURE

Anal fissures cause pain on defecation and usually develop after a period of constipation (may be brief period) and tenesmus. Sometimes the pain can be excruciating, persisting for hours andradiating down the back of both legs. Anal fissures, especially if chronic, can cause minor anorectal bleeding (bright blood) noted as spotting on the toilet paper.

Examination

On inspection the anal fissure is usually seen in the anal margin, situated in the midline posteriorly (6 o'clock). The fissure appearsas an elliptical ulcer involving the lower third of the anus from the dentate line to the anal verge.(Figure 36.2).[1]



Figure 36.2 Anal fissure situated in the mid posterior position of the anal verge - the 6 o'clock position

Digital examination and sigmoidoscopy are difficult because of painful anal sphincter spasm.

If there are multiple fissures Crohn's disease should be suspected. These fissures look different, being indurated, oedematous and bluish in colour.

In chronic anal fissures a sentinel pile is common and in long-standing cases a subcutaneous fistula is seen at the anal margin, with fibrosis and anal stenosis.[1]

Treatment

A high residue diet and avoidance of constipation (aim for soft bulkystools) may lead to resolution and long-term prevention. A combined local anaesthetic and corticosteroid ointment applied to the fissure can provide relief and promote healing. Hot baths relax the internal anal sphincter. Inacute anal fissure will usually heel spontaneouslyor welhen afew week of a freatment of lughfere deft size baths lecture[2]. The most promising conservative treatment is the application of diluted glyceryl trinitrate ointment (e.g. Rectogesicthree times daily to the lower anal canal).[3]Transient headache is the man adverse offciel.

Lateral internal sphincterotomy is indicated in patients with a recurrent fissure and a chronic fissure with a degree of fibrosis and anal stenosis. This surgical procedure is very effective. An alternative is injection of botulinum toxin into the sphincter.

PROCTALGIA FUGAX (LEVATOR ANI SPASM)

Main features:

- fleeting rectal pain
- varies from mild discomfort to severe spasm
- lasts 3 - 30 minutes
- often wakes patient at night
- a functional bowel disorder
- affects adults, usually professional males

Management

- Explanation and reassurance
- Salbutamol inhaler (2 puffs statim) worth a trial

Alternatives include glyceryl trinitrate spray for the symptom or prophylactic quinine sulphate before retiring.

SOLITARY RECTAL ULCER SYNDROME

These ulcers occur in young adults; they canpresent with pain but usually present as the sensation of a rectal lump causing obstructed defecation and bleeding withmucus. The ulcer, which is usually seen on sigmoidoscopy about 10 cm from the anal margin on the anterior rectal wall, can resemble carcinoma. Management is difficultand a chronic course is common. Treatment includes a high residue diet and the avoidance of constipation.

PERIANAL HAEMATOMA

A perianal haematoma is a purple tender swelling at the anal margin caused by rupture of an external haemorrhoidal vein following straining at toilet or some other effort involving a Valsalva manoeuvre. The degree of pain varies from a minor discomfort to severe pain. It has been described as the ‘five day, painful, self-curing pile’.

Management

Surgical intervention is recommended, especially in the presence of severe discomfort. The treatment depends on the time of presentation after the appearance of thehaematoma.

1. *Within 24 hours of onset*. Perform simple aspiration without local anaesthetic using a 19gauge needle while the haematoma is still fluid.
2. *From 24 hours to 5 days of onset*. The blood has clotted and a simple incision under local anaesthetic over the haematoma with deroofing with scissors (like taking the top off a bole egg) to remove the thrombosis by squezing is recommended. Removal of the haematoma reduces the chances of the development of a skin tag, which can be a source of anal irritation.
3. *Day 6 onwards*. The haematoma is bestleft alone unless it is very painful or (rarely) infected. Resolution is evidenced by the appearance of wrinkles in the previously stretched skin.

Follow-up

The patient should be reviewed in 4 weeks for rectal examination and proctoscopy, to examine for any underlying internal haemorrhoid that may predispose to further recurrence. Prevention includes an increased intake of dietary fibre and avoidance of strainingat stool.

STRANGULATED HAEMORRHOIDS

A marked oedematous circumferential swelling will appear if all the haemorrhoids are involved. If only one haemorrhoid is strangulated, proctoscopy will help to distinguish it from a perianal haematoma. Initial treatment is with rest and ice packs and then haem-orrhoidectomy at the earliest possible time. It is best to refer for urgent surgery.

PERIANAL ABSCESS

Clinical features

- Severe, constant, throbbing pain
- Fever and toxicity
- Hot, red, tender swelling adjacent to anal margin
- Non-fluctuant swelling

Careful examination is essential to make the diagnosis. Look for evidence of a fistula.

Treatment

Drain via a cruciate incision which key heed to be deep (with trimming of the corners) over thepoint of maximal induration. A drain tube can be inserted for 7 to 10days. Packing is not necessary.

Antibiotics

It a perianal or per rectal abscen is recalcitrant will cellulites use :

- Metronidazole 400mg (0) 12 hrly for 5 - 7 days plus
- cephalexin 500mg (0) 6 hrly for 5-7 days.[4]

ISCHIORECTAL ABSCESS

An ischiorectal abscess presents as a larger, more diffuse, tender dusky red swelling in the buttock. The presence of an abscess is usually very obvious but the precise focus is not always obvious on inspection. Antibiotics are of little help and surgical incision and drainage as soon as possible is necessary. A deep general anaesthetic is necessary.

PILONIDAL SINUS AND ABSCESS

Recurrent abscesses and discharge in the sacral region (at the upper end of the natal cleft) caused by a midline pilonidal sinus, which often presents as a painful abscess. Once the infection has settled it is important to excise the pits, allow free drainageof the midline cavity and lateral tracks and remove all ingrown hair. Antibiotics are given to complement surgical drainage only if there is severe surrounding cellulitis. Pilonidal means ‘anest of hairs’ and the problem is particularly common in hirsute young men.

TENESMUS

Tenesmus is an unpleasant sensation of incomplete evacuation of the rectum. It causes the patient to attempt defecation at frequent intervals. The most common causeis irritable bowel syndrome. Another common cause is an abnormal mass in the rectum or anal canal, such as carcinoma (e.g. prostate, anorectal), haemorrhoids or a hard faecal mass. In some cases, despite intensive investigation, no cause is found and it appears to be a functional problem.

ANORECTAL LUMPS

Anorectal lumps are relatively common and patients are often concerned because of the fear of cancer. A lump arising from the anal canal orrectum, such as internalhaemorrhoids, tends to appear intermittently upon defecation, and reduce afterwards.[1] Common prolapsing lesions include second- and third-degree haem-orrhoids, hypertrophied anal papilla, polyps and rectal prolapse. Common presenting lumps include skin tags, fourth-degree piles and perianal warts (Table 36.1).

Table 36.1 Common anal lumps

Prolapsing lumps
•second- and third-degree haemorrhoids
•rectal prolapse

•rectal polyp •hypertrophied anal papilla
Persistent lumps •skin tag •perianal warts (condylomata accuminata) •anal carcinoma •fourth-degree haemorrhoids •perianal haematoma •perianal abscess

SKIN TAGS

The skin tag is usually the legacy of an untreated perianal haematoma. It may require excision for aesthetic reasons, for hygiene or because it is a source of pruritus ani or irritation. A tag may be associated with a chronic fissure.

Method of excision

A simple elliptical excision at the base of the skin is made under local anaesthetic. Suturing of the defect is usually not necessary.

PERIANAL WARTS

It is important to distinguish the common viral warts from the condylomata lata of secondary syphilis. Local therapy includes the application of podophyllin every 2or 3 days by the practitioner or imiquimod.

INTERNAL HAEMORRHOIDS

Haemorrhoids or piles are common and tend to develop between the agesof 20 and 50 years. About one out of two Westerners suffers from them by the time 50 is reached.[3] Internal haemorrhoids are a complex of dilated arteries, branches of the superiorhaemorrhoidal artery and veins of the internal haemorrhoidal venous plexus (Figure 36.3). The commonest cause is chronic constipation related to a lack of dietary fibre.


1st-degree internal haemorrhoid
2nd-degree internal haemorrhoid
3rd-degree (prolapsed) internal haemorrhoid
perianal haematoma

Figure 36.3 Classification of haemorrhoids

Anatomically there are three classical sites, namely 3, 7 and 11 o'clock (Figure 36.4).



Figure 36.4 Three sites of primary haemorrhoids, looking into the anus from below

Clinical stages and pathology[3]

- Stage 1: First-degree internal haemorrhoids: three bulges form above the dentate line. Bright bleeding is common.
- Stage 2: Second-degree internal haemorrhoids: the bulges increase in size and slide downwards so that the patient is aware of lumps when straining at stool, but they disappear upon relaxing. Bleeding is a feature.
- Stage 3: Third-degree internal haemorrhoids: the pile continues to enlarge and slide downwards, requiring manual replacement to alleviate discomfort. Bleeding is also a feature.
- Stage 4: Fourth-degree internal haemorrhoids: prolapse has occurred and replacement of the prolapsed pile into the anal canal is impossible.

Symptoms

Bleeding is the main and, in many people, the only symptom. The word 'haemorrhoid' means flow of blood. Other symptoms include prolapse, mucoid discharge, irritation/itching, incomplete bowel evacuation and pain (Figure 36.5).



Figure 36.5 Severely prolapsed haemorrhoids - requiring surgery

Treatment

The treatment of haemorrhoids is based on four procedures: three main, rubber band ligation, cryotherapy and sphincterotomy. Surgery is generally reserved for large strangulated piles. The best treatment, however, is prevention, and softish bulky faeces that pass easily prevent haemorrhoids. People should be advised to have a diet with adequate fibre by eating plenty of fresh fruit, vegetables, whole grain cereals or bran. They should complete their bowel action within afew minutes and avoid using laxatives.

ANAL DISCHARGE

Anal discharge refers to the involuntary escape of fluid from or nearthe anus. The causes may be considered as follows.[5]

1. *Continent*
    - Anal fistula
    - Pilonidal sinus
    - STIs
        - anal warts
        - gonococcal ulcers
        - genital herpes
    - Solitary rectal ulcer syndrome
    - Carcinoma of anal margin
2. Incontinent
    - Minor incontinence - weakness of internal sphincter

    - Severe incontinence - weakness of levator ani and puborectalis

3. Partially continent
    - faecal impaction
    - rectal prolapse

RECTAL BLEEDING

Patients present with any degree of bleedingfrom a smear on the toilet tissue to severe haemorrhage. Various causes are presented in Figure 36.6. Common causes arepolyps, carcinoma of the colon, ischaemic colitis, diverticular disease and haemorrhoids.


ischaemic colitis
carcinoma
angiodysplasia
carcinoma of caecum
diverticulitis
polyp
colitis
carcinoma
solitary ulcer of rectum
anal fissure
haemorrhoids

Figure 36.6 Various causes of rectal bleeding

Local causes of bleeding include excoriated skin, anal fissure, a burst perianal haematoma and anal carcinoma. A characteristic pattern of bright bleeding is found with haemorrhoids. It is usually small non-prolapsing haemorrhoids that bleed.

The nature of the blood (e.g. bright red, dark red or black) and the nature of the bleeding (e.g. smear, streaked on stool, mixed with stool, massive) gives an indication of the source of the bleeding (Table 36.2). Black tarry (melaena) stool indicatesbleeding from the upper gastrointestinal tract and israre distal to the lower ileum. Patients with melaena should be admitted to hospital.

Table 36.2 Presentation and causes of rectal bleeding[6]

Bright red blood in toilet separate from faeces	•Internal haemorrhoids
Bright red blood on toilet paper	•Internal haemorrhoids •Fissure •Anal carcinoma •Pruritus •Anal warts and condylomata
Blood and mucus on underwear	•Third-degree haemorrhoids •Fourth-degree haemorrhoids •Prolapsed rectum •Mucosal prolapse •Prolapsed mucosal polyp
Blood on underwear (no mucus)	•Ulcerated perianal haematoma •Anal carcinoma
Blood and mucus mixed with faeces	•Colorectal carcinoma •Proctitis •Colitis, ulcerative colitis •Large mucosal polyp •Ischaemic colitis
Blood mixed with faeces (no mucus)	•Small colorectal polyps •Small colorectal carcinoma
Melaena (black tarry stools)	•Gastrointestinal bleeding (usually upper) with long transit time to the anus
Torrential haemorrhage (rare)	•Diverticular disorder Angiodysplasia
Large volumes of mucus in faeces (little blood)	•Villous papilloma of rectum •Villous papilloma of colon
Blood in faeces with menstruation (rare)	•Rectal endometriosis

Source: Orlay, G. Office Proctology, p. 11.[6] © Copyright 1987 George Orlay - reproduced withpermission

Frequent passage of blood and mucus indicates a rectal tumour or proctitis, whereas more proximal tumours or extensive colitis present different patterns.

Substantial haemorrhage, which is rare, can be caused by diverticulardisorder, angio-dysplasia or more proximal lesions such as Meckel's diverticulum and even duodenal ulcers. Angiodysplasias are 5 mm collections of dilated mucosal capillaries and thick-walled submucosal veins, found usually in the ascending colon of elderly patients who have noother bowel symptoms. The bleeding is persistent and recurrent. The site is identified by technetium-labelled red cell scan or colonoscopy.

The history should also include an analysis of any associated symptoms such as pain, diarrhoea or constipation, presence of lumps and a sensation of urgency or unsatisfied defecation. The latter symptoms point to a rectal cause. Associated change of bowel habit suggests a diagnosis of carcinoma of the rectum or left colon. Bleeding from right colon cancer is often occult, presenting as anaemia.

The examination includes a general assessment, anal inspection, digital rectal examination and proctosigmoidoscopy. Even if there is an anal lesion, proximal bleeding must be excluded in all cases by sigmoidoscopy[3] and by colonoscopy if there are any bowel symptoms or no obvious anal cause or a doubt abouta lesion causing the symptoms.

PRURITUS ANI

Pruritus ani, which is itching of the anus, can be a distressing symptom that is worse at night, during hot weather and during exercise. It is seen typically in adult males with considerable inner drive, often at times of stress and in hot weather when sweating is excessive. In children, threadworm infestation should be suspected. It may be part of general itching, such as with a skin disorder, or localised whereby various anorectal disorders have to be excluded.sebarrhoen dermatitin us a partianlary caman underlying factor.

Physical signs

The skin changes can vary from minimal signs to marked pathology thatcan show linear ulceration, maceration or lichenification (Figure 36.7). Superficial skin changes can be moist and macerated or dry and scaly. Full anorectal examination is necessary.



Figure 36.7 Lichen chronicus simplex. Lichenification from scratching with longstanding pruritus

Causes and aggravating factors

- Psychological factors:
  - stress and anxiety
  - fear of cancer
- Generalised systemic or skin disorders:
  - eczema
  - diabetes mellitus
  - candidiasis
  - psoriasis (look for fissures in natal cleft)
  - antibiotic treatment
  - worms: pinworm, threadworm
  - diarrhoea causing excoriation
- Local anorectal conditions:
  - piles
  - fissures
  - warts
- Zealous hygiene
- Contact dermatitis:
  - dyed or perfumed toilet tissue, soap, powder
  - clothing
- Excessive sweating:
  - e.g. tight pantyhose in summer

Diagnostic approach

- Urinalysis (? diabetes)
- Anorectal examination
- Scrapings and microscopy to detect organisms
- Stool examination for intestinal parasites

Rules of treatment: patient education

1. *Scratching*. Stop - it's taboo! If you scratch at night, wear light cotton gloves tobed.
2. *Bathing*. Avoid hot water. Excessive showering and scrubbing is also bad for this condition. Usea cream such as bland aqueous cream or Cetaphil lotion or a mild soap substitute for cleaning rather than soap.
3. *Drying*. Keep the area as dry and cool as possible. After washing, dry gently and thoroughly with a soft towel or soft tissue: do not rub. Warm air from a hair drier is very useful.
4. *Bowel movements*. Keep bowels regular and smooth by eating plenty of high-fibre foods such as bran, fresh carrots and apples. Some doctors claim that your bowel actions should be so smooth and complete that toilet paper should hardly be necessary.
5. *Toilet*. Clean gently after bowel movements. Soft paper tissue (avoid pastel tints) may be used, and then clean with tufts of cottonwool with aqueous cream or bland soap and water. The best way is to use cotton wool in warm water.
6. *Soaps and powder*. Do not use perfumed soaps and talcum powder, including baby powder. Zinc powder (e.g. Curash) is suitable to keep the area dry and relieve itching.
7. *Clothing*. Wear loose clothing and underwear. In men, boxer shorts should be used in preference to jockey shorts. Cottons should be used. Let the air circulate in the area. At times a skirt but no underpants (in women) is desirable. Avoid pantyhose if possible.
8. *Topical creams*. Do not use ointments unless your doctor has prescribed them. If a cream has to be used, simple creams may be the most soothing (e.g. toilet lanoline).

Topical treatment

- Treat the cause (if known).
- Avoid local anaesthetics, antiseptics.
- Advise aqueous cream to wash anus (instead of soap).
- Most effective preparations (for short[7] courses):
  methyprednisolone aceponate 0.1% in a fatty ointment base
  or
  hydrocortisone 1% cream/ointment
  or
  hydrocortisone 1% cream with clioquinol 3% or clotrimazole 1% (especially if dermatosis and canaida suspected)

If an isolated area and resistant, infiltrate 0.5 mL of triamcinolone intradermally. Fractionated X-ray therapy can be used if very severe.

ANAL (FAECAL) INCONTINENCE

Studies have revealed that up to 1 in 10 people suffer some degree offaecal incontinence, which is a common reason for in-stitutionalisation of the elderly.[8] Patients may be reluctant to seek medical advice and doctors often do not ask specifically about the con-dition. The problem is as common in men as in women.

Practice tips for pruritus ani

- Most cases of uncomplicated pruritus ani resolve with simple measures, including explanation and reassurance.
- Otherwise prescribe a cortico steroid, especially methylprecinisolone aceponate 0.1% once symptoins controlled use hydrocortisone 1%[7]
- Lifestyle stress and anxiety underlies mostcases.
- In obese patients with intertrigo and excessive sweating strap the buttocks apart with adhesive tape.

Apart from ageing, risk factors include perianal injury, such as childbirth injury, anal surgery, irritable bowel syndrome and neurological disorders.

Among the various possible treatments there are surgical possibilities, such as the Neosphincter artificial bowel sphincter.[9] It is worth keeping in mindasking patients about the possibility of this problem and knowing ‘to whom to refer’.

REFERENCES

1. Hunt P, Marshall V. *Clinical Problems in General Surgery*. Sydney: Butterworths, 1991: 311.
2. Utzig MJ, kroesen AS, Buhr HS. Conservative Amd Gastroenterology, 2003:98:968 - 74.
3. Lund JN, Scholefield JH. A randomised, prospective, double-blind, placebo-controlled trial of glyceryl trinitrate ointment in the treatment of anal fissure. Lancet, 1997; Jan 4: 11 - 13.
4. Spices s (chair), *Theapeuic Iudelences: Antibiotics* (version 12). Therapentio ∧ndelines Ltd,2003: 67 - 68.
5. Ryan P. *A Very Short Textbook of Surgery* (2nd edn). Canberra: Dennis & Ryan, 1990: 37 - 42.
6. Orlay G. *Office Proctology*. Sydney: Australasian Medical Publishing Company, 1987: 11 - 52.
7. Marley; (chare). *Therapeutic Indelines: Dermatology* (version2). Therapentic Indelines Limited, 2004:184.
8. Kalantar JS, Howell S, Talley NJ. Prevalence of faecal incontinence and associated risk factors. Med J Aust, 2002; 176: 54 - 7.
9. O'Brien P. Faecal incontinence: artificial sphincter a new treatment option. Aust Doctor, 1999; 22 October: 57.

CHAPTER 37. LOW BACK PAIN

Last Wednesday night while carrying a bucket of water from the well, Hannah Williams slipped upon the icy path and fell heavily upon her back. We fear her spine was injured for though she suffers acute pain in her legs she cannot move them. The poor wild beautiful girl is stopped in her wildness at last.

FRANCIS KILVERT 1874

LOW BACK PAIN

Low back pain accounts for at least 5% of general practice presentations. The most common cause is minor soft tissue injury, but patients with this do not usually seek medical help because the problem settles within a few days.

Most back pain in patients presenting to GPs is postulated to be due to dysfunction of elements of the mobile segment, namely the facet joint, the intervertebral joint (with its disc) and the ligamentous and muscular attachments. This problem, often referred to as mechanical back pain, will be described as vertebral dysfunction - a general term that, while covering radicular and non-radicular pain, includes dysfunction of the joints of the spine, although the specific origin in most instances cannot be determined. It is therefore appropriate to refer to this as 'non-specific back pain' [1].

KEY FACTS AND CHECKPOINTS

- Back pain accounts for at least 5% of all presenting problems in general practice in Australia and 6.5% in Britain.[2]
- In the United States it is the commonest cause of limitation of activity in those under the age of 45.[3]
- Approximately 85% of the population will experience back pain at some stage of their lives, while 70% of the world's population will have at least one disabling episode of low back pain in their lives.[3]
- At least 50% of these people will recover within 2 weeks and 90% within 6 weeks, but recurrences are frequent and have been reported in 40 - 70% of patients; 2 - 7% develop chronic pain.[4]
- The most common age groups are the 30s, 40s and 50s, the average age being 45 years.[4]
- The most common cause of back pain is a minor strain to muscles and/or ligaments, but people suffering from this type of back pain usually do not seek medical treatment as most of these soft tissue problems resolve rapidly.
- The main cause of back pain presenting to the doctor is dysfunction of the intervertebral joints of the spine due to injury, also referred to as mechanical back pain (at least 70%).
- The second most common cause of back pain is spondylosis (synonymous with osteoarthritis and degenerative back disease). It accounts for about 10% of cases of low back pain.
- L5 and S1 nerve root lesions represent most of the cases of sciatica presenting in general practice. They tend to present separately but can occur together with a massive disc protrusion.
- An intervertebral disc prolapse has been proven in only 6 - 8% of cases of back pain.[3]

CAUSES OF LOW BACK PAIN

To develop a comprehensive diagnostic approach, the practitioner should have a clear understanding of the possible causes of low back and leg pain and of the relative frequency of their clinical presentations. The major causes of low back pain in several hundred patients presenting to the author's general practice are summarised in Table 37.1.

Table 37.1 Major causes of low back ± leg pain presenting in the author's general practice

Patients	%
Vertebral dysfunction	71.8
Lumbar spondylosis	10.1
Depression	3.0
Urinary tract infection	2.2
Spondylolisthesis	2.0
Spondyloarthropathies	1.9
Musculoligamentous strains/tears	1.2
Malignant disease	0.8

Arterial occlusive disease	0.6
Other	6.4
	100.0

Relevant causes are illustrated in Figure 37.1.


Visceral and vascular
B—biliary disorders
U—penetrating duodenal ulcer
P—pancreatitis
R—renal disorders
aortic aneurysm (ruptured/dissecting)
retroperitoneal haemorrhage
female pelvic disorders (e.g. endometriosis)
prostatitis
abscess (e.g. pilonidal)
arterial embolism
claudication of arterial occlusive disease
R
P
U
B
Musculoskeletal
Scheuermann's disorder
pathological fracture
metastatic disease
spondylolisthesis
disc disruption
facet joint dysfunction spondylosis
sacroiliac dysfunction sacroiliitis
trochanteric bursitis gluteus medius bursitis
coccydynia
ischial bursitis
meralgia paraesthetica
neurogenic claudication
radicular pain
referred pain
from spinal dysfunction
radicular pain

Figure 37.1 Relevant causes of back pain with associated buttock and leg pain

ANATOMICAL AND PATHOPHYSIOLOGICAL CONCEPTS

Recent studies have focused on the importance of disruption of the intervertebral disc in the cause of back pain. A very plausible theory has been advanced by Maigne[5] who proposes the existence, in the involved mobile segment, of a minor intervertebral derangement (MID). He defines it as 'isolated pain in one intervertebral segment, of a mild character, and due to minor mechanical cause'.

The MID always involves one of the two apophyseal joints in the mobile segment, thus initiating nociceptive activity in the posterior primary dermatome and myotome.

Maigne points out that the functional ability of the mobile segment depends intimately upon the condition of the intervertebral disc. Thus, if the disc is injured, other elements of the segment will be affected. Even a minimal disc lesion can produce apophyseal joint dysfunction, which is a reflex cause of protective muscle spasm and pain in the corresponding segment, with loss of function (Figure 37.2).



Figure 37.2 Reflex activity from a MID in the intervertebral motion segment. Apart from the local effect caused by the disruption of the disc (A), interference can occur in the facet joint (B) and interspinous ligament (C) leading possibly to muscle spasm (D) and skin changes (E) via the posterior rami Reproduced with permission from C. Kenna and J. Murtagh. *Back Pain And Spinal Manipulation*. Sydney: Butterworths, 1989

In theory, any structure with a nociceptive nerve supply may be a source of pain. Such structures include the ligaments, fascia and muscles of the lumbosacral spine, intervertebral joints, facet joints, dura mater and sacroiliac joints.[6]

Actually, pain can theoretically arise from any innervated structure in the region of the spine. It can be neurogenic, spondylogenic, viscerogenic, vasculogenic or rarely psychogenic.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic strategy model is presented in Table 37.2.

Table 37.2 Low back pain: diagnostic strategy model

Q. Probability diagnosis
A. Vertebral dysfunction especially facet joint and disc
Musculoligamentous strain/sprain
Spondylosis (degenerative OA)
Q. Serious disorders not to be missed
A. Cardiovascular
• ruptured aortic aneurysm
• retroperitoneal haemorrhage (anticoagulants)
Neoplasia
• myeloma

• metastases Severe infections • osteomyelitis • epidural abscess • discitis • tuberculosis • pelvic abscess/PID Cauda equina compression
Q. Pitfalls (often missed) A. Spondyloarthropathies • ankylosing spondylitis • Reiter's syndrome • psoriasis • bowel inflammation Sacroiliac dysfunction Spondylolisthesis Claudication • vascular • neurogenic Prostatitis Endometriosis
Q. Seven masquerades checklist A. Depression Diabetes - Drugs - Anaemia - Thyroid disorder - Spinal dysfunction UTI
Q. Is this patient trying to tell me something? A. Quite likely. Consider lifestyle, stress, work problems, malingering, conversion reaction

Note: Associated buttock and leg pain included.

Probability diagnosis

The commonest cause of low back pain is vertebral dysfunction or mechanical pain, which then has to be further analysed. The term can embrace musculoskeletal strain, discogemc and postenor ligament pain, or facetogenic ciys function/pain.

Degenerative changes in the lumbar spine (lumbar spondylosis) are commonly found in the older age group. This problem, and one of its complications, spinal canal stenosis, is steadily increasing along with the ageing population.

Serious disorders not to be missed

It is important to consider malignant disease, especially in an older person. It is also essential to consider infection such as acute osteomyelitis and tuberculosis, which is often encountered in recent immigrants, especially those from Asia central and Africa. These conditions are considered in more detail under thoracic back pain. For pain or anaesthesia of sudden onset, especially when accompanied by neurological changes in the legs, consider cauda equina compression due to a massive disc prolapse and also retroperitoneal haemorrhage. It is important to ask patients if they are taking anticoagulants. See Table 37.3.

Table 37.3 'Red flag' pointers to serious low back pain conditions[7]

Age > 50 years

History of cancer

Temperature > 37.8°C

Constant pain - day and night

Weight loss

Symptoms in other systems,. e.g. cough, breast mass

Significant trauma

Features of spondyloarthropathy, e.q. peripheral arthritis

Neurological deficit

Drug or alcohol abuse

Use of anticoagulants

Use of corticosteroids

No improvement over 1 month

Possible cauda equina syndrome

- saddle anaesthesia
- recent onset bladder dysfunction
- severe or progressive neurological deficit

Pitfalls

The inflammatory disorders must be kept in mind, especially the spondyloarthropathies, which include psoriatic arthropathy, ankylosing spondylitis, Reiter's syndrome, inflammatory bowel disorders such as ulcerative colitis and Crohn's disease, and reactive arthritis. The spondyloarthropathies are more common than appreciated and must be considered in the younger person presenting with features of inflammatory back pain (i.e. pain at rest, relieved by activity). The old trap of confusing claudication in the buttocks and legs, due to a high arterial obstruction, with sciatica must be avoided.

General pitfalls

- Being unaware of the characteristic symptoms of inflammation and thus misdiagnosing one of the spondyloarthropathies.
- Overlooking the early development of malignant disease or osteomyelitis; if suspected, and an X-ray is normal, a radionuclide scan should detect the problem.
- Failing to realise that mechanical dysfunction and osteoarthritis can develop simultaneously, producing a combined pattern.
- Overlooking anticoagulants as a cause of a severe bleed around the nerve roots and corticosteroids leading to osteoporosis.
- Not recognising back pain as a presenting feature of the drug addict.

‘Red flag’ pointers

There are several so-called ‘red flag’ or precautionary pointers to a serious underlying cause of back pain (Table 37.3). Such symptoms and signs should alert the practitioner to a serious health problem and thus guide selection of investigations, particularly plain films of the lumbar spine.

Seven masquerades checklist

Of these conditions, depression and urinary tract infection have to be seriously considered. For the young woman with upper lumbar pain, especially if she is pregnant, the possibility of a urinary tract infection must be considered. These patients may not have urinary symptoms, such as dysuria and frequency.

Depressive illness has to be considered in any patient with a chronic pain complaint. This common psychiatric disorder can continue to aggravate or maintain the pain even though the provoking problem has disappeared. This is more likely to occur in people who have become anxious about their problem or who are under excessive stress. Many doctors treat such patients with a therapeutic trial of antidepressant medication, for example, amitriptyline or doxepin.

Psychogenic considerations

The patient may be unduly stressed, not coping with life or malingering. It may be necessary to probe beneath the surface of the presenting problem.

A patient with low back pain (LBP) following lifting at work poses a problem that causes considerable anguish to doctors, especially when the pain becomes chronic and complex. Chronic pain may be the last straw for patients who have been struggling to cope with personal problems; their fragile equilibrium is upset by the back pain. Many patients who have been dismissed as malingerers turn out to have a genuine problem. The importance of a caring, competent practitioner with an insight into all facets of his or her patient's suffering, organic and functional, becomes obvious. The tests for non-organic back pain are very useful in this context.

'Yellow flag' pointers[1]

This term has been introduced to identify psychosocial and occupational factors that may increase the risk of chronicity in people presenting with acute back pain. Counter psychological issues of:

- abnormal illness behaviour
- compensation issues
- unsatisfactory restoration of activities
- failure to return to work
- unsatisfactory response to treatment
- treatment refused
- a typical physical signs

NATURE OF THE PAIN

The nature of the pain may reveal its likely origin. Establish where the pain is worst - whether it is central (proximal) or peripheral. The following are general characteristics and guides to diagnosis:

- aching throbbing pain ° inflammation (e.g. sacroiliitis)
- deep aching diffuse pain ° referred pain (e.g. dysmenorrhoea)
- superficial steady diffuse pain ° local pain (e.g. muscular strain)
- boring deep pain ° bone disease (e.g. neoplasia, Paget's disease)
- intense sharp or stabbing (superimposed on a dull ache) ° radicular pain (e.g. sciatica)

A comparison of the significant features of the two most common types of pain - mechanical and inflammatory - is presented in Table 37.4.

Table 37.4 Comparison of the patterns of pain for inflammatory and mechanical causes of low back pain[7]

Feature	Inflammation	Mechanical
History	Insidious onset	Precipitating injury/previous episodes
Nature	Aching, throbbing	Deep dull ache, sharp if root compression
Stiffness	Severe, prolonged Morning stiffness	Moderate, transient
Effect of rest	Exacerbates	Relieves
Effect of activity	Relieves	Exacerbates
Radiation	More localised, bilateral or alternating	Tends to be diffuse, unilateral
Intensity	Night, early morning	End of day, following activity

THE CLINICAL APPROACH

History

Analysing the history invariably guides the clinician to the diagnosis. The pain patterns have to be carefully evaluated and it is helpful to map the diurnal variations of pain to facilitate the diagnosis (Figure 37.3).



Figure 37.3 Typical daily patterns of pain for conditions causing back pain. Note conditions that can wake patients from sleep and also the combined mechanical and inflammatory patterns

It is especially important to note the intensity of the pain and its relation to rest and activity. In particular, ask whether the pain is present during the night, whether it wakes the patient, is present on rising or whether it is associated with stiffness.

Continuous pain present day and night is suggestive of neoplasia or infection. Pain on waking also suggests inflammation or depressive illness. Pain provoked by activity and relieved by rest suggests mechanical dysfunction while pain worse at rest and relieved by moderate activity is typical of inflammation. In some patients the coexistence of mechanical and inflammatory causes complicates the pattern.

Pain aggravated by standing or walking that is relieved by sitting is suggestive of spondylolisthesis. Pain aggravated by sitting (usually) and improved with standing indicates a discogenic problem.

Pain of the calf that travels proximally with walking indicates vascular claudication; pain in the buttock that descends with walking indicates neurogenic claudication. This latter problem is encountered more frequently in older people who have a tendency to spinal canal stenosis associated with spondylosis.

Key questions

- What is your general health like?
- Can you describe the nature of your back pain?
- Was your pain brought on by an injury?
- Is it worse when you wake in the morning or later in the day?

- How do you sleep during the night?
- What effect does rest have on the pain?
- What effect does activity have on the pain?
- Is the pain worse when sitting or standing?
- What effect does coughing or sneezing or straining at the toilet have?
- What happens to the pain in your back or leg if you go for a long walk?
- Do you have a history of psoriasis, diarrhoea, penile discharge, eye trouble or severe pain in your joints?
- Do you have any urinary symptoms?
- What medication are you taking? Are you on anticoagulants?
- Are you under any extra stress at work or home?
- Do you feel tense or depressed or irritable?

Physical examination

The basic objectives of the physical examination are to reproduce the patient's symptoms, detect the level of the lesion and determine the cause (if possible) by provocation of the affected joints or tissues. This is done using the time-honoured method of joint examination - look, feel, move and test function. The patient should be stripped to a minimum of clothing so that careful examination of the back can be made. A neurological examination of the lower limb should be performed if symptoms extend below the buttocks. It is important to perform a rectal examination to check for flaccidity in a patient with suspected cauda equina syndrome.

A useful screening test for a disc lesion and dural tethering is the slump test.[8]

The main components of the physical examination are:

1. inspection
2. active movements
    - forward flexion (to reproduce the patient's symptoms)
    - extension (to reproduce the patient's symptoms)
    - lateral flexion (R & L) (to reproduce the patient's symptoms)
3. provocative tests (to reproduce the patient's symptoms)
4. palpation (to detect level of pain)
5. neurological testing of lower limbs (if appropriate)
6. testing of related joints (hip, sacroiliac)
7. assessment of pelvis and lower limbs for any deformity (e.g. leg shortening)
8. general medical examination, including rectal examination

Important landmarks

The surface anatomy of the lumbar region is the basis for determining the vertebral level. Key anatomical landmarks include the iliac crest, spinous processes, the sacrum and the posterior superior iliac spines (PSISs).

- The tops of the iliac crest lie at the level of the L4 - 5 interspace (or the L4 spinous process).
- The PSISs lie opposite S2 (Figure 37.4).


L1
L2
L3
L4
L5
line projected from tops of iliac crests usually bisects L4–5 interspace
posterior–superior iliac spine
sacral hiatus
coccyx

Figure 37.4 Surface anatomy and important landmarks of the lumbosacral spine

Inspection

Inspection begins from the moment the patient is sighted in the waiting room. A patient who is noted to be standing is likely to have a significant disc lesion. Considerable in-formation can be obtained from the manner in which the patient arises from a chair, moves to the consulting room, removes the shoes and clothes, gets onto the examination couch and moves when unaware of being watched.

The spine must be adequately exposed and inspected in good light. Patients should undress to their underpants; women may retain their brassiere and it is proper to provide them with a gown that opens down the back. Note the general contour and symmetry of the back and legs, including the buttock folds, and look for muscle wasting. Note the lumbar lordosis and any abnormalities, such as lateral deviation. If lateral deviation (scoliosis) is present it is usually away from the painful side.

Note the presence of midline moles, tufts of hair or haemangioma that might indicate an underlying congenital anomaly, such as spina bifida occulta.

Movements of the lumbar spine

There are three main movements of the lumbar spine. As there is minimal rotation, which mainly occurs at the thoracic spine, rotation is not so important. The movements that should be tested, and their normal ranges, are as follows:

- extension (20° - 30°) (Figure 37.5a)
- lateral flexion, left and right (30°) (Figure 37.5b)
- flexion (75° - 90°: average 80°) (Figure 37.5a)


(a)
Extension
spinous process C7
20°–30°
80°
Flexion
sacrum



Figure 37.5 (a) Degrees of movement of the lumbar spine: flexion and extension **(b)** degree of lateral flexion of the lumbar spine
Reproducedwith permission from C. Kenna and J. Murtagh. *Back Pain And Spinal Manipulation*. Sydney: Butterworths, 1989

Measurement of the angle of movement can be made by using a line drawn between the sacrum and large prominence of the C7 spinous process.

Palpation

Have the patient relaxed, lying prone, with the head to one side and the arms by the sides. The levels of the spinous processes are identified by standing behind the patient and using your hands to identify the L4 and L5 spinous processes in relation to the top of the iliac crests. Mark the important reference points.

Palpation, which is performed with the tips of the thumbs opposed, can commence at the spinous process of L1 and then systematically proceed distally to L5 and then over the sacrum and coccyx. Include the interspinous spaces as well as the spinous processes. When the thumbs (or other part of the hand such as the pisiforms) are applied to the spinous processes, a firm pressure is transmitted to the vertebrae by a rocking movement for three or four ‘springs’. Significant reproduction of pain is noted.

Palpation occurs at three main sites:

- centrally (spinous processes to coccyx)
- unilateral - right and left sides (1.5 cm from midline)
- transverse pressure to the sides of the spinous processes (R and L)

Slump test

The slump test is an excellent provocation test for lumbosacral pain and is more sensitive than the straight leg raising (SLR) test. It is a screening test for a disc lesion and dural tethering. It should be performed on patients who have low back pain with pain extending into the leg, and especially for posterior thigh pain.

A positive result is reproduction of the patient's pain, and may appear at an early stage of the test (when it is ceased).

Method

1. The patient sits on the couch in a relaxed manner.
2. The patient then slumps forward (without excessive trunk flexion), and then places the chin on the chest.
3. The unaffected leg is straightened.
4. The affected leg only is then straightened (Figure 37.6).



Figure 37.6 The slump test: one of the stages

5. Both legs are straightened together.
6. The foot of the affected straightened leg is dorsiflexed.

Note: Take care to distinguish from hamstring pain. Deflexing the neck relieves the pain of spinal origin, not hamstring pain.

Significance of the slump test

- It is positive if the back or leg pain is reproduced.
- If positive, it suggests disc disruption.
- If negative, it may indicate lack of serious disc pathology.
- If positive, one should approach manual therapy with caution.

Neurological examination[8]

A neurological examination is performed only when the patient's symptoms, such as pain, paraesthesia, anaesthesia and weak-ness, extend into the leg.

The importance of the neurological examination is to ensure that there is no compression of the spinal nerves from a prolapsed disc or from a tumour. This is normally tested by examining those functions that the respective spinal nerves serve, namely skin sensation, muscle power and reflex activity.

The examination is not daunting but can be performed quickly and efficiently in 2 to 3 minutes by a methodical technique that improves with continued use. The neurological examination consists of:

1. quick tests

  - walking on heels (L5)
  - walking on toes (S1)

2. dural stretch tests
  - slump test
  - straight leg raising

3. specific nerve root tests (L4, L5, S1)
  - sensation
  - power
  - reflexes

Main nerve roots

Refer to Figure 37.7.


L5
Neurologic
level
motor
extensor hallucis
longus
reflex
none
sensation
L5
S1
Neurologic
level
motor
peroneus longus
+ brevis
reflex
sensation
S1

Figure 37.7 The main motor, sensory and reflex features of the nerve roots L5 and S1 Reproduced With Permission From S. Hoppenfeld, *Physical Examination Of The Spine And Extremities*, Norwalk, Ct: Appleton & Lange 1976

L3:

- femoral stretch test (prone, flex knee, extend hip)
- motor - extension of knee
- sensation - anterior thigh
- reflex - knee jerk (L3, L4)

L4:

- motor - resisted inversion foot
- sensation - inner border of foot to great toe
- reflex - knee jerk

L5:

- motor
  - walking on heels
  - resisted extension great toe
- sensation - middle three toes (dorsum)

- reflex - nil

S1:

- motor
  - walking on toes
  - resisted eversion foot
- sensation - little toe, most of sole
- reflex - ankle jerk (S1, S2)

Other examination

The method of examining the sacroiliac and hip joints is outlined in Chapter 65.

INVESTIGATIONS

Investigations for back pain can be classified into three broad groups: front-line screening tests; specific disease investigations; and procedural and preprocedural tests.

Plain X-ray of the lumbar spine are not routinely recommended in acute non-specific LBP in the absence of ‘sea flags’ as they are of limiter diagnostic vaine and no benefits in physical function are observer.[1]

Screening tests

These are most important for the patient presenting with chronic back pain when serious disease such as malignancy, osteoporosis, infection or spondyloarthropathy must be excluded. The screening tests for chronic pain are:

- plain X-ray
- urine examination (office dipstick)
- ESR
- serum alkaline phosphatase
- prostatic specific antigen (in males > 50 years)

Specific disease investigation

Such tests include:

- peripheral arterial studies
- HLA-B_{27} antigen test for ankylosing spondylitis and Reiter's syndrome
- serum electrophoresis for multiple myeloma
- PSA for possible carcinoma of prostate
- Brucella agglutination test
- blood culture for pyogenic infection and bacterial endocarditis
- bone scanning to demonstrate inflammatory or neoplastic disease and infections (e.g. osteomyelitis) before changes are apparent on plain X-ray
- tuberculosis studies
- X-rays of shoulder and hip joint
- electromyographic (EMG) studies to screen leg pain and differentiate neurological diseases from nerve compression syndromes
- radioisotope scanning
- technetium pyrophosphate scan of SIJ for ankylosing spondylitis
- selective anaesthetic block of facet joint under image intensification
- selective anaesthetic block of medial branches of posterior primary rami and other nerve roots

Procedural and preprocedural diagnostic tests

These tests should be kept in reserve for chronic disorders, especially mechanical disorders, that remain undiagnosed and unabated, and where surgical intervention is planned for a disc prolapse requiring removal.

Depending on availability and merit, such tests include:

- CT scan
- myelography or radiculography
- discography
- MRI

SUMMARY OF DIAGNOSTIC GUIDELINES FOR SPINAL PAIN

- Continuous pain (day and night) ° neoplasia, especially malignancy or infection.
- The big primary malignancy is multiple myeloma.
- The big three metastases are from lung, breast and prostate.
- The other three metastases are from thyroid, kidney/adrenal and melanoma.
- Pain with standing/walking (relief with sitting) ° spondylolisthesis.
- Pain (and stiffness) at rest, relief with activity ° inflammation.
- In a young person with inflammation think of ankylosing spondylitis or Reiter's syndrome.
- Stiffness at rest, pain with or after activity, relief with rest ° osteoarthritis.
- Pain provoked by activity, relief with rest ° mechanical dysfunction.
- Pain in bed at early morning ° inflammation, depression or malignancy/infection.
- Pain in periphery of limb ° discogenic → radicular or vascular → claudication or spinal canal stenosis → claudication.
- Pain in calf (ascending) with walking ° vascular claudication.
- Pain in buttock (descending) with walking ° neurogenic claudication.
- One disc lesion ° one nerve root (exception is L5 - S1 disc).
- One nerve root ° one disc (usually).
- Two or more nerve roots - consider neoplasm.
- The rule of thumb for the lumbar nerve root lesions is L3 from L2 - 3 disc, L4 from L3 - 4, L5 from L4 - 5 and S1 from L5 - S1.
- A large disc protrusion can cause bladder symptoms, either incontinence or retention.
- A retroperitoneal bleed from anticoagulation therapy can give intense nerve root symptoms and signs.

BACK PAIN IN CHILDREN

The common mechanical disorders of the intervertebral joints can cause non-specific back pain in children, which must always be taken seriously. Like abdominal pain and leg pain, it can be related to psychogenic factors, so this possibility should be considered by dip-lomatically evaluating problems at home, at school or with sport. It is helpful to remember that tight hamstring is associated with non-specific back discomfort witgh poor forward flexion.

Especially in children under the age of 10, it is very important to exclude organic disease. Infections such as osteomyelitis and tuberculosis are rare possibilities, and ‘discitis’ has to be considered. This painful condition can be idiopathic, but can also be caused by the spread of infection from a vertebral body. It has characteristic radiological changes.

Tumours causing back pain include the benign osteoid osteoma and the malignant osteogenic sarcoma. Osteoid osteoma is a very small tumour with a radiolucent nucleus that is sharply demarcated from the surrounding area of sclerotic bone. Although more common in the long bones of the leg, it can occur in the spine.

In older children and adolescents (in whom back pain is common) the organic causes of back pain are more likely to be inflammatory, congenital or from de-velopmental anomalies and trauma.

A prolapsed intervertebral disc, which can occur in adolescents, can be very unusual in its presentation. There is often marked spasm, with a stiff spine and lateral deviation, which may be out of proportion to the relatively lower degree of pain.

Other important conditions to consider are Scheurmanni disease (which largely affects thoracic spine) and early onset ankylosing Spondylitis.

Spondylolisthesis can occur in older children, usually due to a slip of L5 or S1, because the articular facets are congenitally absent or because of a stress fracture in the pars interarticularis. It is necessary to request standing lateral and oblique X-rays.

BACK PAIN IN THE ELDERLY

Traumatic spinal dysfunction is still the most common cause of back pain in the elderly and may represent a recurrence of earlier dysfunction. It is amazing how commonly disc prolapse and facet joint injury can present in the aged. However, degenerative joint disease is very common and, if advanced, can present as spinal stenosis with claudication and nerve root irritation due to narrowed intervertebral foraminae.

Special problems to consider are malignant disease, degenerative spondylolisthesis, vertebral pathological fractures and occlusive vascular disease.

ACUTE BACK AND LEG PAIN DUE TO VERTEBRAL DYSFUNCTION

Mechanical disruption of the vertebral segment or segments is the outstanding cause to consider, while the main serious clinical syndromes are secondary to disruption with or without prolapse of the intervertebral disc, usually L4 - 5 or L5 - S1.

Table 37.5 presents the general clinical features and diagnosis in acute back pain (fractures excluded) following vertebral dysfunction: the symptoms and signs can occur singly or in combination.

Table 37.5 Clinical features and diagnosis of vertebral dysfunction leading to low back and leg pain[8]

Clinical features	Frequency	Diagnosis
Syndrome A(surgical emergency)	Very rare	Spinal cord (UMN) or cauda equina (LMN) compression
Saddle anaesthesia (around anus, scrotum or vagina) Distal anaesthesia Evidence of UMN or LMN lesion Loss of sphincter control or urinary retention Weakness of legs peripherally		
Syndrome B (probable surgical emergency) Anaesthesia or paraesthesia of the leg Foot drop Motor weakness Absence of reflexes	Uncommon	Large disc protrusion, paralysing nerve root
Syndrome C Distal pain with or without paraesthesia Radicular pain (sciatica) Positive dural stretch tests	Common	Posterolateral disc protrusion on nerve root or disc disruption
Syndrome D Lumbar pain (unilateral, central or bilateral) ± buttock and posterior thigh pain	Very common	Disc disruption or facet dysfunction or unknown (non-specific) causation

Fortunately, syndromes A and B are extremely rare but, if encountered, urgent referral to a surgeon is mandatory. Clinical features of the cauda equina syndrome are presented in Figure 37.8. Syndrome B can follow a bleed in patients taking anticoagulant therapy or be caused by a disc sequestration after inappropriate spinal manipulation.


Pain
backs of thighs and legs
Numbness
buttocks, back of legs,
soles of feet,
saddle anaesthesia
Weakness
paralysis of legs and feet
Atrophy
calves
Paralysis
bladder and bowel
urinary retention

Figure 37.8 Cauda equina syndrome due to massive prolapsed intervertebral disc

Vertebral dysfunction with non-radicular pain

This outstanding common cause of low back pain is considered to be due mainly to dysfunction of the pain-sensitive facet joint. The precise pathophysiology is difficult to pinpoint.

Typical profile[8]

Age:	• Any age - late teens to old age, usually 22 - 55 years
History of injury:	• Yes, lifting or twisting
Site and radiation:	• Unilateral lumbar (may be central)
	• Refers over sacrum, SIJ areas, buttocks
Type of pain:	• Deep aching pain, episodic
Aggravation:	• Activity, lifting, gardening, housework (vacuuming, making beds, etc.)
Relief:	• Rest, warmth
Associations:	• May be stiffness, usually good health
Physical examination (significant):	• Localised tenderness - unilateral or central L4, L5 or S1 levels, may be restricted flexion, extension, lateral flexion
Diagnosis confirmation:	• Investigation, which is usually inappropriate, invariably normal

Note: diagnosis made clinically

Management[1,7]

- Activity directed by degree of pain but normal activity encouraged from outset
- Back education program
- Analgesics - paracetamol
- Consider NSAIDs (if inflammatory pattern)
- Exercise program and swimming (as tolerated) - conflicting evidence re efficacy
- Physical therapy - mobilisation, manipulation (for persistent problems, conflicting evidence) (refer to pages 386 - 7)

Current evidence for acute low back pain can be summarised as follows:[4]

- beneficial - advice to stay active, NSAIDs
- likely to be beneficial - analgesics, spinal manipulation/stretching (reduces period of morbidity)
- unknown - back exercises, trigger point injections, acupuncture

For chronic low back pain (pain > 12 weeks):

- beneficial - back exercises, multidisciplinary treatment program
- likely benefit - analgesics, NSAIDs, trigger point injections, spinal manipulation

Radiculopathy

Radicular pain, caused by nerve root compression from a disc protrusion (most common cause) or tumour or a narrowed intervertebral foramina, typically produces pain in the leg related to the dermatome and myotome innervated by that nerve root. Leg pain may occur alone without back pain and vary considerably in intensity.

The two nerve roots that account for most of these problems are L5 and S1 and the commonest disc lesson is L4 - 5, closely followed by L5 - 51. A disc can be confined, extruded or sequestrated. Most settle with time (6 - 12 weeks). The management is outlined at the end of this chapter and under ‘Sciatica’ (Chapter 66).

SPONDYLOLISTHESIS

About 5% of the population have spon-dylolisthesis but not all are symptomatic. The pain is caused by extreme stretching of the interspinous ligaments or of the nerve roots. The onset of back pain in many of these patients is due to concurrent disc degeneration rather than a mechanical problem. The pain is typically aggravated by prolonged standing, walking and exercise. The physical examination is quite diagnostic.

- Physical examination (significant):
    - stiff waddling gait
    - increased lumbar lordosis
    - flexed knee stance
    - tender prominent spinous process of ‘slipped’ vertebrae
    - limited flexion
    - hamstring tightness or spasm
- Diagnosis confirmation:
    - Lateral X-ray (standing) (Figure 37.9).



Figure 37.9 Spondylolisthesis: illustrating a forward shift of one vertebra on another

Management

This instability problem can be alleviated with relief of symptoms by getting patients to follow a strict flexion exercise program for at least 3 months. The objective is for patients to 'splint' their own spine by strengthening abdominal and spinal muscles.

Extension of the spine should be avoided, especially hyperextension. Gravity traction might help. Recourse to lumbar corsets or surgery (for spinal fusion) should be resisted although it is appropriate in a few severe intractable cases.

LUMBAR SPONDYLOSIS

Lumbar spondylosis, also known as degenerative osteoarthritis or osteoarthrosis, is a common problem of wear and tear that may follow vertebral dysfunction, especially after severe disc disruption and degeneration.

Stiffness of the low back is the main feature of lumbar spondylosis. Although most people live with and cope with the problem, progressive deterioration can occur, leading to subluxation of the facet joints. Subsequent narrowing of the spinal and intervertebral foramen leads to spinal canal stenosis (Figure 37.10).


osteophytic
formation
at facet joint
narrowed
spinal canal
annular
bulge of disc
subluxation
of facet joint
narrowed
invertebral foramen

Figure 37.10 Lumbar spondylosis with degeneration of the disc and facet joint, leading to narrowing of the spinal canal and intervertebral foramen

Management

- Basic analgesics (depending on patient response and tolerance)
- NSAIDs (judicious use)
- Appropriate balance between light activity and rest
- Exercise program and hydrotherapy (if available) - physiotherapy supervision
- Regular mobilisation therapy may help
- Consider trials of electrotherapy, such as TENS and acupuncture

THE SPONDYLOARTHROPATHIES

The seronegative spondyloarthropathies are a group of disorders characterised by in-volvement of the sacroiliac joints with an ascending spondylitis and extraspinal manifestations, such as oligoarthritis and enthesopathies (Figure 37.11) (refer to Chapter 35). The pain and stiffness that are the characteristic findings of spinal involvement are typical of inflammatory disease: namely, worse in the morning, may occur at night and improves rather than worsens with exercise.


(a)
(b)



Figure 37.11 (a) Ankylosing spondylitis and psoriasis: main target areas on vertebral column and girdle joints (**b)** Crohn's disease and ulcerative colitis: main target areas of enteropathies **(c)** Reiter's syndrome: main target areas

The main disorders in this group are ankylosing spondylitis, psoriatic arthritis, Reiter's syndrome, reactive spondylo-arthropathies and the inflammatory bowel diseases. Hence the importance of searching for a history of psoriasis, diarrhoea, urethral discharge, eye disorders and episodes of arthritis in other joints. The following profile for ankylosing spondylitis serves as a typical clinical presentation of back pain for this group.

Typical profile of ankylosing spondylitis[7]

Age and sex:	• Young men 15 - 30 years (rare onset after 40)
History of injury:	• None, unless coincidental
	• Has a slow insidious onset
Site and radiation:	• Low back, may radiate to both buttocks or posterior thigh (rare below knees)
	• Can alternate sides
Type of pain:	• Aching, throbbing pain of inflammation
	• Commonly episodic
Aggravation:	• Often worse at night (can wake patient), turning over in bed and rising in the morning
Relief:	• Activity including exercise
	• Patient may walk around during night for relief

Associations:	• Back stiffness, especially in morning
	• Pain and stiffness in thoracic or cervical spine
	• Pain and stiffness in thoracic cage
	• Peripheral joint pain (up to 50% of cases)
	• Iritis (up to 25% of cases)
Physical examination (significant):	• Absent lumbar lordosis
	• Lateral flexion limited first, then flexion and extension
	• Positive sacroiliac joint stress tests
	• Positive Schober's test
Diagnosis confirmation:	• X-ray of pelvis (sacroiliitis)
	• Bone scans and CT scans
	• ESR usually elevated
	• HLA-B_{27} antigen positive in over 90% of cases (10% of population are positive)

Treatment

The earlier the treatment the better the outlook for the patient; the prognosis is usually good. The basic objectives of treatment are:

- prevention of spinal fusion in a poor position
- relief of pain and stiffness
- maintenance of optimum spinal mobility

The basic methods of management are:

- advice on good back care and posture
- general education and counselling
- exercise programs to improve the range of movement and maintain mobility
- referral to physiotherapist
- drug therapy, especially tolerated NSAIDs, preferably indomethacin in optimal dosage
- sulphasalazine - a useful second-line agent if the disease progresses despite NSAIDs
- cytotoxic therapy (methotreseate) may be necessary in severe progressive disease

MALIGNANT DISEASE

It is important to identify malignant disease and other space-occupying lesions as early as possible because of the prognosis and the effect of a delayed diagnosis on treatment.

With respect to the neurological features, more than one nerve root may be involved and major neurological signs may be present without severe root pain. The neurological signs will be progressive.

If malignant disease is proved and myeloma is excluded, a search should be made for the six primary malignancies that metastasise to the spine (Figure 37.12). If the bone is sclerotic consider prostatic secondaries, some breast secondaries or Paget's disease.



Figure 37.12 Important primary malignancies to the spine. Note the difference between sclerotic and osteolytic metastases; multiple myeloma also causes osteoporotic lesions

NON-ORGANIC BACK PAIN

Like headache, back pain is a symptom of an underlying functional, organic or psychological disorder. Preoccupation with organic causation of symptoms may lead to serious errors in the assessment of patients with back pain. Any vulnerable aching area of the body is subject to aggravation by emotional factors.

Depressed patients are generally less demonstrative than patients with extreme anxiety and conversion disorders and malingerers, and it is easier to overlook the non-organic basis for their problem.

A trial of antidepressants for a minimum of 3 weeks is recommended and quite often a positive response with relief of backache eventuates.

Failure to consider psychological factors in the assessment of low back pain may lead to serious errors in diagnosis and management. Each instance of back pain poses a stimulating exercise in differential diagnosis. A comparison of organic and non-organic features is presented in Table 37.6.

Table 37.6 Comparison of general clinical features of organic and non-organic based low back pain[6,7]

Symptoms	Organic disorders	Non-organic disorders

Presentation	Appropriate	Often dramatic
Pain	Localised	Bilateral/diffuse Sacrococcygeal
Pain radiation	Appropriate Buttock, specific sites	Inappropriate Front of leg/whole leg
Time pattern	Pain-free times	Constant, acute or chronic
Paraesthesia/anaesthesia	Dermatomal Points with finger	May be whole leg Shows with hands
Response to treatment	Variable Delayed benefit	Patient often refuses treatment Initial improvement (often dramatic) then deterioration (usually within 24 hours)
Signs		
Observation	Appropriate Guarded	Overreactive under scrutiny Inconsistent
Tenderness	Localised to appropriate level	Often inappropriate level Withdraws from probing finger
Spatial tenderness (Magnuson)	Consistent	Inconsistent
Active movements	Specific movements affected	Often all movements affected
Axial loading test	No back pain (usually)	Back pain
SLR ‘distraction’ test	Consistent	Inconsistent
Sensation	Dermatomal	Non-anatomical ‘sock’ or ‘stocking’
Motor	Appropriate myotome	Muscle groups (e.g. leg ‘collapses’)
Reflexes	Appropriate May be depressed	Brisk hyperactive

Assessment of the pain demands a full understanding of the patient. One must be aware of his or her type of work, recreation, successes and failures; and one must relate this information to the degree of incapacity attributed to the back pain.

Patients with psychogenic back pain, especially the very anxious, tend to overemphasise their problem. They are usually demonstrative, the hands being used to point out various painful areas almost without prompting. There is diffuse tenderness even to the slightest touch and the physical disability is out of proportion to the alleged symptoms. The pain distribution is often atypical of any dermatome and the reflexes are almost always hyperactive. It must be remembered that patients with psychogenic back pain - for example, depression and conversion disorders - do certainly experience back pain and do not fall for the traps set for the malingerer.

Tests for non-organic back pain[9]

Several tests are useful in differentiating between organic and non-organic back pain (e.g. that caused by depression or complained of by a known malingerer).

Magnuson's method (the ‘migratory pointing’ test)

1. Request the patient to point to the painful sites.
2. Palpate these areas of tenderness on two occasions separated by an interval of several minutes, and compare the sites.

Between the two tests divert the patient's attention from his or her back by another examination.

Paradoxical straight leg raising test

Perform the usual SLR test. The patient might manage a limited elevation, for example, 30°. Keep the degree in mind. Ask the patient to sit up and swing the leg over the end of the couch. Distract attention with another test or some question, and then attempt to lift the straight leg to the same level achieved on the first occasion. If it is possible, then the patient's response is inconsistent.

Burn's ‘kneeling on a stool’ test

1. Ask the patient to kneel on a low stool, lean over and try to touch the floor.
2. The person with non-organic back pain will usually refuse on the grounds that it would cause great pain or that he or she might overbalance in the attempt.

Patients with even a severely herniated disc usually manage the task to some degree.

The axial loading test

1. Place your hands over the patient's head and press firmly downward (Figure 37.13).



Figure 37.13 The axial loading test

2. This will cause no discomfort to (most) patients with organic back pain.

TREATMENT OPTIONS FOR BACK PAIN

General aspects of management[1,7]

The aim of treatment is to reduce pain, maintain function and minimise the risk of chronicity.

Advice to stay active

Evidence from randomised controlled trials confirms that, in people with acute low back pain, advice to stay active speeds symptomatic recovery, reduces chronic disability and results in less time off work compared with bed rest or usual care.[3]Encourage the patient to stay at work or return early if possible.

The caring knowledgeable therepest

Evidence supports the positive value of education and reassurance from a confident, supporture and knowing therapest.

Relative rest

For acutely painful debilitating back problems 2 days strict rest lying on a firm surface is optimal treatment.[8] Resting for longer than 3 days does not produce any significant healing, and patients should be encouraged to return to activities of daily living as soon as possible.

Patient education

Appropriate educational material leads to a clear insight into the causes and aggravation of the back disorder plus coping strategies.

Heat

The application of heat in some form, including heat bags, hot flannels and similar methods, can be of benefit especially in the first 2 - 4 weeks of acute LBP.

Exercises

An early graduated exercise program as soon as the acute phase settles has been shown to promote healing and prevent relapses.[11] All forms of exercise (extension, flexion and isometric) appear to be equally effective (see Figure 37.14). Swimming is an excellent activity for back disorders.



Figure 37.14 Examples of exercises for low back pain: **(a)** rotation exercise; **(b)** flexion exercise

Studies support the use of exercises for chronic back pain rather than acute pain.[4]

Pharmacological agents

Basic analgesics

Analgesics such as paracetamol and codeine plus paracetamol (acetaminophen) should be used for pain relief. Paracetamol is re-commended as the first-line analgesic.

NSAIDs

These are useful where there is clinical evidence of inflammation, especially with the spondyloarthropathies, severe spondylosis and in acute radicular pain, to counter the irritation on the nerve root. NSAIDs have been shown to be more effective than placebo in acute back pain for pain relief and overall improvement. There is no evidence to distinguish different NSAIDs.[4]

Injection techniques

Trigger point injection

This may be effective for relatively isolated points using 5 - 8 mL of local anaesthetic. Studies indicate that it is likely to be more beneficial for chronic back pain.[4]

Chymopapain

This enzyme has been advocated for the treatment of acute nuclear herniation that is still intact. The indications are similar for surgical discectomy. However, studies show that although it is more effective than placebo, it is less effective than surgical discectomy.[12]

Facet joint injection

Corticosteroid injection under radio image intensification is widely used in some clinics. The procedure is delicate and expertise is required. Best evidence to date does not support the use of these injections.

Epidural injections

Injections of local anaesthetic with or without corticosteroids are used for chronic pain, especially for nerve root pain. The author favours the caudal (trans-sacral) epidural injection for persistent sciatica using 15 mL of half-strength local anaesthetic only (e.g. 0.25% bupivacaine) (Figure 37.15).



Figure 37.15 Caudal epidural injection: the needle should lie free in the space and be well clear of the dural sac

Others favour the use of steroids and the lumbar epidural approach. The evidence regarding effectiveness is conflicting.[4]

Physical therapy

Active exercises are the best form of physical therapy (Figures 37.14a, 37.14b).

Passive spinal stretching at the end range is a safe, effective method (Figure 37.16). Spinal mobilisation is a gentle, repetitive, rhythmic movement within the range of movement of the joint. It is safe and quite effective and a variation of stretching.



Figure 37.16 Lumbosacral spinal stretching technique (for right sided pain): a traditional technique. Illustration of direction of line of force Reproduced With Permission From C. Kenna And J. Murtagh. *Back Pain And Spinal Manipulation*. Sydney: Butterworths, 1989

Spinal manipulation is a high velocity thrust at the end range of the joint. It is generally more effective and produces a faster response but requires accurate diagnosis and greater skill. It is more effective for uncomplicated dysfunctional low back pain (without radicular pain), especially acute pain (Figure 37.17). [4, 7, 13, 14] However, the evidence from controlled trials is conflicting. Adverse effects are uncommon, but can be severe.



Figure 37.17 Lumbar spinal stretching manipulation: illustration of the specific technique for the L4 - 5 level with arrows indicating the direction of applied force Reproduced With Permission From C. Kenna And J. Murtagh. Back Pain And Spinal Manipulation. Sydney: Butterworths, 1989

Other treatments

The following treatments have advocates for the management of back pain, although clear-cut evidence for the efficacy of these modalities is still lacking:

- hydrotherapy
- traction
- TENS
- facet joint injection
- posterior nerve root (medial branch) blocks with or without denervation (by cryotherapy or radiofrequency)
- deep friction massage (in conjunction with mobilisation and manipulation)
- acupuncture
- pain clinic (if unresponsive to initial treatments)
- biofeedback
- gravitational methods (home therapy)
- lumbar supports

MANAGEMENT GUIDELINES FOR LUMBOSACRAL DISORDERS (SUMMARY)

The management of ‘mechanical’ back pain depends on the cause. Since most of the problems are mechanical and there is a tendency to natural resolution, conservative management is quite appropriate. The rule is: ‘if patients with uncomplicated back pain receive no treatment, one-third will get better within 1 week and by 3 weeks almost all the rest of the other two-thirds are better’.[15] Practitioners should have a clear-cut management plan with a firm, precise, reassuring and conservative clinical approach.

The problems can be categorised into general conditions for which the summarised treatment protocols are outlined.

- Acute pain ° pain less than 4 weeks It is usually defined as pain that has been present for at least 3 months.
- Subacute pain ° pain 4 - 12 weeks
- Chronic pain ° pain greater than 3 months

Acute low back pain

The common problem of low back pain caused by facet joint dysfunction and/or limited disc disruption usually responds well to the following treatment. The typical patient is aged 20 - 55 years, is well and has no radiation of pain below the knee.[16]

- Back education program
- Encouragement of normal daily activities according to degree of comfort
- Regular non-opioid analgesics (e.g. paracetamol)
- Physical therapy: stretching of affected segment, muscle energy therapy, spinal mobilisation or manipulation (if no contraindication on first visit)[12, 14, 16]
- Review in about 5 days (probably best time for physical therapy)
- No investigation needed initially

Most of these patients can expect to be relatively pain free in 14 days and can return to work early (some may not miss work and this should be encouraged). The evidence concerning spinal manipulation is that it reduces the period of disability.

Sciatica with or without low back pain

Sciatica is a more complex and protracted problem to treat, but most cases will gradually settle within 12 weeks.

Acute

- Back education program
- Resume normal activities as soon as possible
- Regular non-opioid analgesics with review as the patient mobilises
- NSAIDs for 10 - 14 days, then cease and review
- Walking and swimming
- Weekly or 2-weekly follow-up

consider iv a course of corticosteroids fo severe pain, e.g. prednisolone 50 mg for 5 days then 25 mg for 5 days, gradually tapering to 3 weeks in total.*or* 30 mg daily m for 3 weeks. Tapering to 0 over next 2 weeks (efficiency not clearly established)

Chronic

- Reassurance that problem will subside (assuming no severe neurological defects)
- Consider epidural anaesthesia (if slow response)

General guidelines for surgical intervention for radiculopathy

Absolute

- Bladder/bowel control disturbance; perineal sensory change
- Progressive motor disturbance (e.g. significant foot drop, weakness in quadriceps)

Relative

- Severe prolonged pain or disabling pain
- Failure of conservative treatment with persistent pain (problem of permanent nerve damage)

Note: An important controlled prospective study comparing surgical and conservative treatment in patients with sciatica over 10 years showed that there was significant relief of sciatica in the surgical group for 1 to 2 years but not beyond that time. At 10 years both groups had the same outcome, including neurological deficits.[17]

Chronic back pain

The basic management of the patient with uncomplicated chronic back pain should follow the following guidelines:

- back education program and ongoing support
- encouragement of normal activity
- exercise program
- analgesics (e.g. paracetamol)
- NSAIDs for 14 days (if inflammation, i.e. pain at rest - relieved by activity) and review
- trial of mobilisation or manipulation (at least three treatments) - if no contraindications[13, 14]
- consider trigger point injection
- a multidisciplinary team approach is recommended/back schools

Prevention of further back pain

Patients should be informed that an ongoing back care program should give them an excellent outlook. Prevention includes:

- education about back care, including a good layperson's reference

- golden rules to live by: how to lift, sit, bend, play sport and so on
- an exercise program: a tailor-made program for the patient
- posture and movement training, such as
  - the Alexander technique[18]
  - the Feldenkrais technique[19]

WHEN TO REFER

Urgent referral

- Myelopathy, especially acute cauda equina compression syndrome
- Severe radiculopathy with progressive neurologic deficit
- Spinal fractures

Other referrals

- Neoplasia or infection
- Undiagnosed back pain
- Paget's disease
- Continuing pain of 3 months duration without a clearly definable cause

Practice tips

- Back pain that is related to posture, aggravated by movement and sitting, and relieved by lying down is due to vertebral dysfunction, especially a disc disruption.
- The pain from most disc lesions is generally relieved by rest.
- Plain X-rays are of limited use, especially in younger patients, and may appear normal in disc prolapse.
- Remember the possibility of depression as a cause of back pain; if suspected, consider a trial of antidepressants.
- If back pain persists, possibly worse during bed rest at night, consider malignant disease, depressive illness or other systemic diseases.
- Pain that is worse on standing and walking, but relieved by sitting, is probably caused by spondylolisthesis.
- If pain and stiffness is present on waking and lasts longer than 30 minutes upon activity, consider inflammation.
- Avoid using strong analgesics (especially opioids) in any chronic non-malignant pain state.
- Bilateral back pain is more typical of systemic diseases, while unilateral pain typifies mechanical causes.
- Back pain at rest and morning stiffness in a young person demand careful investigation: consider inflammation such as ankylosing spondylitis and Reiter's syndrome.
- A disc lesion of L5 - S1 can involve both L5 and S1 roots. However, combined L5 and S1 root lesions should still be regarded with suspicion (e.g. consider malignancy).
- A large central disc protrusion can cause bladder symptoms, either incontinence or retention.
- Low back pain of very sudden onset with localised spasm and protective lateral deviation may indicate a facet joint syndrome.
- The T12 - L1 and L1 - 2 discs are the groin pain discs.
- The L4 - 5 disc is the back pain disc.
- The L5 - S1 disc is the leg pain disc.
- Severe limitation of SLR (especially to less than 30°) indicates lumbar disc prolapse.
- A preventive program for dysfunctional back pain based on back care awareness and exercises is mandatory advice.
- Remember that most back problems resolve within a few weeks, so avoid overtreatment.

REFERENCES

1. NIT & MRC. *Evidence-based Management of Acute Musculoskeletal Pain. A Guide for Clinicians* . Australian Acute Musculoskeletal Pain Guidelines Group. Canberra: Australian Government, 2004.
2. Cormack J, Marinker M, Morrell D. *Practice: a Handbook of Primary Health Care*. London: Kluwer-Harrap Handbooks, 1980; 3(68): 1 - 10.
3. Sloane P, Slatt M, Baker R. *Essentials of Family Medicine*. Baltimore: Williams & Wilkins, 1988: 228 - 35.
4. Barton S ed. *Clinical Evidence*. London. BMJ Publishing Group, 2001: 772 - 87.
5. Maigne R. Manipulation of the spine. In: Basmajian JV ed. *Manipulation, Traction and Massage*. Paris: RML, 1986: 71 - 96.

6. Bogduk N. The sources of low back pain. In: Jaysom M ed. *The Lumbar Spine and Back Pain* (4th edn). Edinburgh: Churchill Livingstone, 1992: 61 - 88.
7. Mashford L (Chair). *Therapeutic guidelines: Rheumatolugy* (Version 4). Melbourne: Therapeutic guidelines Ltd, 2006: 85 - 100.
8. Kenna C, Murtagh J. *Back Pain and Spinal Manipulation* (2nd edn). Oxford: Butterworth Heinemann, 1997: 70 - 164.
9. Waddell G et al. Non-organic physical signs in low back pain. Spine, 1980; 5: 117 - 25.
10. Deyo RA, Diehl AK, Rosenthal M. How many days of bed rest for acute low back pain? A randomised clinical trial. N Engl J Med, 1986; 315: 1064 - 70.
11. Kendall PH, Jenkins SM. Exercises for backache: a double blind controlled study. Physiotherapy, 1968; 54: 154 - 7.
12. Gibson JNA et al. Surgery for lumbar disc prolapse. Cochrane Database of Systematic Reviews Issue 2, 2002.
13. Blomberg S, Svardsudd K, Mildenberger F. A controlled, multicentre trial of manual therapy in low back pain. Scand J Prim Health Care, 1992; 10: 170 - 8.
14. Royal College of General Practitioners et al. *Clinical Guidelines for the Management of Acute Low Back Pain*. London: RCGP, 1996.
15. Kuritzky L. Low back pain. Family Practice, Audio-Digest California Medical Association: 1996; 44:14.
16. Deyo RA. Acute low back pain: a new paradigm for management. Br Med J, 1996; 313: 1343 - 4.
17. Weber et al. A controlled prospective study with 10 years observation of patients with sciatica. Spine, 1983; 8: 131 - 40.
18. Hodgkinson L. *The Alexander Technique*. London: Piatkus, 1988: 1 - 97.
19. Feldenkrais M. *Awareness Through Movement*. New York: Harper & Row, 1972.

CHAPTER 38. THORACIC BACK PAIN

The maladies that afflict the clerks aforesaid arise from three causes; first constant sitting, secondly the incessant movement of the hand and always in the same direction, and thirdly the strain on the midline from the effort not to disfigure the books by error or cause loss to their employers.

THE PHYSICIAN RAMAZZINI IN 1713

THORACIC BACK PAIN

Thoracic (dorsal) back pain is common in people of all ages, including children and adolescents. It account for 10-15% of all spinal pain. Dysfunction of the joints of the thoracic spine, with its unique costovertebral joints (which are an important source of back pain), is very commonly encountered in medical practice, especially in people whose lifestyle creates stresses and strains through poor posture and heavy lifting. Muscular and ligamentous strains may be common, but they rarely come to light in practice because they are self-limiting and not severe.

This dysfunction can cause referred pain to various parts of the chest wall and can mimic the symptoms of various visceral diseases, such as angina, biliary colic and oesophageal spasm. In similar fashion, heart and gall bladder pain can mimic spinal pain.

KEY FACTS AND CHECKPOINTS

- The commonest site of pain in the spine is the costovertebral articulations especially the costotransverse articulation (Figure 38.1).



Figure 38.1 The functional unit of the thoracic spine

- Pain of thoracic spinal origin may be referred anywhere to the chest wall, but the commonest sites are the scapular region, the paravertebral region 2 - 5 cm from midline and, anteriorly, over the costochondral region.
- Thoracic (also known as dorsal) pain is more common in patients with abnormalities such as kyphosis and Scheuermann's disorder.
- Trauma to the chest wall (including falls on the chest such as those experienced in body contact sport) commonly lead to disorders of the thoracic spine.
- Unlike the lumbar spine the joints are quite superficial and it is relatively easy to find the affected (painful) segment.
- The intervertebral disc prolapse is very uncommon in the thoracic spine.
- The older patient presenting with chest pain should be regarded as having a cardiac cause until proved otherwise.
- If the chest pain is non-cardiac, then the possibility of referral from the thoracic spine should be considered.
- The thoracic spine is the commonest site in the vertebral column for metastatic disease.
- Scheuermann's disorder, which affects the lower thoracic spine in adolescents, is often associated with kyphosis and recurrent thoracic back pain. Always inspect the thoracic spine of the younger patient for kyphosis and scoliosis, ideally at 9 years of age.
- Palpation is the most important component of the physical examination.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 38.1.

Table 38.1 Thoracic back pain: diagnostic strategy model

Q. Probability diagnosis
A. Musculoligamentous strains (mainly postural) Vertebral dysfunction
Q. Serious disorders not to be missed
A. Cardiovascular
• myocardial infarction
• dissecting aneurysm
• pulmonary infarction
• epidural haematoma (blood-thinning agents)
Neoplasia
• myeloma
• lung (with infiltration)
• metastatic disease
Severe infections
• pleurisy
• infectious endocarditis
• osteomyelitis
Pneumothorax
Osteoporosis
Q. Pitfalls (often missed)
A. Angina
Gastrointestinal disorders
• oesophageal dysfunction
• peptic ulcer (penetrating)
• hepatobiliary
• pancreatic
Herpes zoster
Spondyloarthropathies

Costochondritis • Tietze syndrome Fibromyalgia syndrome Polymyalgia rheumatica Chronic infection • tuberculosis • brucellosis **Q. Seven masquerades checklist**
A. Depression ✔ Diabetes - Drugs - Anaemia - Thyroid disorder - Spinal dysfunction ✔ UTI ✔ **Q. Is this patient trying to tell me something?**
A. Yes, quite possible with many cases of back pain.

Probability diagnosis

The commonest cause of thoracic back pain is musculoskeletal, due usually to mus-culoligamentous strains caused by poor posture. However, these pains are usually transitory and present rarely to the practitioner. The problems that commonly present are those caused by dysfunction of the lower cervical and thoracic spinal joints, especially those of the mid-thoracic (interscapular) area.

Arthritic conditions of the thoracic spine are not relatively common although degenerative osteoarthritis is encountered at times; the inflammatory spondyloarthropathies are uncommon.

The various systemic infectious diseases such as influenza and Epstein-Barr mononucleosis can certainly cause diffuse backache but should be assessed in context.

Not to be missed

A special problem with the thoracic spine is its relationship with the many thoracic and upper abdominal structures that can refer pain to the back. These structures are listed in Table 38.2 but, in particular, myocardial infarction and dissecting aneurysm must be considered. A complex problem described by neurosurgeons is the patient presenting with severe sudden thoracic back pain caused by an epidural haematoma related to aspirin or warfarin therapy.

Table 38.2 Non-musculoskeletal causes of thoracic back pain

Heart	• myocardial infarction
	• angina
	• pericarditis
Great vessels	• dissecting aneurysm
	• pulmonary embolism (rare)
	• pulmonary infarction
	• pneumothorax
	• pneumonia/pleurisy
Oesophagus	• oesophageal rupture
	• oesophageal spasm
	• oesophagitis

Subdiaphragmatic disorders of	• oesophageal cancer • gall bladder
Miscellaneous infections	• stomach (including ulcers) • duodenum (including ulcers) • pancreas • subphrenic collection • herpes zoster
Psychogenic	• Bornholm disorder • infective endocarditis

Red flag pointers[1]

The red flag pointers are similar to that for low back pain (Table 38.3)

Table 38.3 Thoracic back pain: red flag pointers

Fracture Pointer
• Major trauma
• Minor trauma
- osteoporosis
- female>50year
- mail>60year
Malignaney Pointer
• Age >50
• Past history malignancy
• Qnexplaned weigh loss
• Pain at rest
• constent pain
• Night pain
• urns at multiple sites
Infeetion pointer
• Feren
• Night soviets
• Risk factors for infixion
Other serious conditions
• Chest pain/heaviness
• Shortness of breath, cough

Cardiopulmonary problems

The acute onset of pain can have sinister implications in the thoracic spine where various life-threatening cardiopulmonary and vascular events have to be kept in mind. The pulmonary causes of acute pain include spontaneous pneumothorax, pleurisy and pulmonary infarction. Thoracic back pain may be associated with infective endocarditis due to embolic phenomena. The ubiquitous myocardial infarction or acute coronary occlusion may, uncommonly, cause interscapular back pain, while the very painful dissecting or ruptured aortic aneurysm may cause back pain with hypotension.

Osteoporosis

Osteoporosis, especially in people over 60 years, including both men and women, must always be considered in such people presenting with acute pain, which can be caused by a pathological fracture. The association with pain following inappropriate physical therapy such as spinal manipulation should also be considered.

Acute infections

Infective conditions that can involve the spine include osteomyelitis, tuberculosis, brucellosis, syphilis and *Salmonella* infections. Such conditions should be suspected in young patients (osteomyelitis), farm workers (brucellosis) and migrants from South-East Asia and third world countries (tuberculosis). The presence of poor general health and fever necessitates investigations for these infections.

Neoplasia

Fortunately, tumours of the spine are uncommon. Nevertheless, they occur frequently enough for the full-time practitioner in back disorders to encounter some each year, especially metastatic disease.

The three common primary malignancies that metastasise to the spine are those originating in the lung, breast and the prostate (all paired structures). The less common primaries to consider are the thyroid, the kidney and adrenals and malignant melanoma.

Reticuloses such as Hodgkin's lymphoma can involve the spine. Primary malignancies that arise in the vertebrae include multiple myeloma and sarcoma.

Benign tumours to consider are often neurological in origin. An interesting tumour is the osteoid osteoma, which is aggravated by consuming alcohol and relieved by aspirin.

The tumours of the spine are summarised in Table 38.4.

Table 38.4 Significant tumours affecting the thoracic and lumbar spine

	Benign	Malignant
Of bone	• osteoid osteoma	Primary
	• haemangioma	• multiple myeloma
	• osteoblastoma	• lymphomas (e.g. Hodgkin's)
	• aneurysmal bone cyst	• sarcoma
	• eosinophilic granuloma	
Spinal	Extradural	Secondary
	• lipoma	• breast
	• neuroma	• lung
	• fibroma	• prostate
	Intradural	• adrenals/kidney
	• neuroma	• thyroid
	• ependymoma	• melanoma
	• chordoma	Direct spread
	• meningioma	• stomach
		• large bowel
		• pancreas
		• uterus/cervix/ovary

Source: After Kenna and Murtagh[2]

The symptoms and signs that should alert the clinician to malignant disease are:

- back pain occurring in an older person
- unrelenting back pain, unrelieved by rest (this includes night pain)
- rapidly increasing back pain

- constitutional symptoms (e.g. unexplained weight loss, fever, malaise)
- a history of treatment for cancer (e.g. excision of skin melanoma)

An ESR and a plain X-ray of the thoracic spine should be the initial screening test in the presence of these pointers.

A common trap for the thoracic spine is carcinoma of the lung, such as mesothelioma, which can invade parietal pleura or structures adjacent to the vertebral column.

Pitfalls

Pitfalls include ischaemic heart disease presenting with interscapular pain, herpes zoster at the pre-eruption stage and the various gastrointestinal disorders. Two commonly misdiagnosed problems are a penetrating duodenal ulcer presenting with lower thoracic pain and oesophageal spasm, which can cause thoracic back pain.

Inflammatory rheumatological problems are not common in the thoracic spine but occasionally a spondyloarthropathy such as ankylosing spondylitis manifests here, although it follows some time after the onset of sacroiliitis.

Seven masquerades checklist

Spinal dysfunction is the outstanding cause in this checklist, but urinary tract infection may occasionally cause lower thoracic pain. Depression always warrants consideration in any pain syndrome, especially back pain. It can certainly cause exaggeration of pre-existing pain from vertebral dysfunction or some other chronic problem.

Psychogenic considerations

Psychogenic or non-organic causes of back pain can present a complex dilemma in diagnosis and management. The causes may be apparent from the incongruous behaviour and personality of the patient, but often the diagnosis is reached by a process of exclusion. There is obviously some functional overlay to everyone with acute or chronic pain, hence the importance of appropriate reassurance to these patients that their problem invariably subsides with time and that they do not have cancer

ANATOMICAL AND CLINICAL FEATURES

The functional unit of the thoracic spine is illustrated in Figure 38.1. Although there is scant literature and evidence about the origins of pain in the thoracic spine,[2] the strongest evidence indicates that pain from the thoracic spine originates mainly from the apophyseal joints and rib articulations. Any one thoracic vertebra has ten separate articulations, so the potential for dysfunction and the difficulty in clinically pinpointing the precise joint at a particular level are apparent.

The costovertebral joints are synovial joints unique to the thoracic spine and have two articulations - costotransverse and costo-central. Together with the apophyseal joints, they are capable of presenting with well-localised pain close to the midline or as referred pain, often quite distal to the spine, with the major symptoms not appearing to have any relationship to the thoracic spine.

Generalised referral patterns are presented in Figure 38.2, while the dermatome pattern is outlined in Figure 38.3.


referral to upper thorax from lower cervical spine
same segmental referral
three segments distal referral
anterior abdominal referral
T12 referral to buttock

Figure 38.2 Examples of referral patterns for the thoracic spine



(a) Anterior **(b)** Posterior

Figure 38.3 Dermatomes for the thoracic nerve roots, indicating possible referral areas. Reproduced With Permission From C. Kenna And J. Murtagh. *Back Pain And Spinal Manipulation* (2 ed), Oxford: Butterworth-heinemann, 1997.

The pain pattern acts as a guide only because there is considerable dermatomal overlap within the individual and variation from one person to another. It has been demonstrated that up to five nerve roots may contribute to the innervation of any one point in the anterior segments of the trunk dermatomes, a fact emphasised by the clinical distribution of herpes zoster.

Upper thoracic pain[1]

Dysfunction of the joints of the upper thoracic spine usually gives rise to localised pain and stiffness posteriorly but also can cause distal symptoms, probably via the autonomic nervous system.

A specific syndrome called the T4 syndrome[4] has been shown to cause vague pain in the upper limbs and diffuse, vague head and posterior neck pain. It has been proven to respond to spinal manipulation.

However, most of the pain, stiffness and discomfort arises from dysfunction of the upper and middle thoracic segments with patients presenting with the complaint of pain between 'my shoulder blades'.

Costovertebral joint dysfunction[1]

The unique feature of the thoracic spine is the costovertebral joint. Dysfunction of this joint commonly causes localised pain approximately 3 - 4 cm from the midline where the rib articulates with the transverse process and the vertebral body. In addition, it is frequently responsible for referred pain ranging from the midline, posterior to the lateral chest wall, and even anterior chest pain.

When the symptoms radiate laterally, the diagnosis is confirmed only when movement of the rib provokes pain at the costovertebral joint. This examination will simultaneously reproduce the referred pain.

Figure 38.4 presents the pattern of referred pain from these joints and highlights the capacity of the thoracic spine to refer pain centrally to the anterior chest and upper abdomen. Confusion arises for the clinician when the patient's history focuses on the anterior chest pain and fails to mention the presence of posterior pain, should it be present. The shaded areas on Figure 38.4 represent those areas where the patient experiences pain following the injection of hypertonic saline into the posterior elements of the spine.



Figure 38.4 Kellegren's (1939) pain referral patterns after stimulation of deep joints of the thoracic spine

THE CLINICAL APPROACH

History

The history of a patient presenting with thoracic back pain should include a routine pain analysis, which usually provides important clues for the diagnosis. The age, sex and occupation of the patient are relevant. Pain in the thoracic area is very common in people who sit bent over for long periods, especially working at desks. Students, secretaries and stenographers are therefore at risk, as are nursing mothers, who have to lift their babies.

People who are kyphotic or scoliotic or who have 'hunchbacks' secondary to disease such as tuberculosis and poliomyelitis also suffer from recurrent pain in this area.

Older people are more likely to present with a neoplastic problem in the thoracic spine and with osteoporosis. Senile osteoporosis is usually a trap because it is symptomless until the intervention of a compression fracture. Symptoms following such a fracture can persist for 3 months.

Pain that is present day and night indicates a sinister cause.

Features of the history that give an indication that the pain is arising from dysfunction of the thoracic spine include:

- *Aggravation and relief of pain on trunk rotation*. The patient's pain may be increased by rotating (twisting) towards the side of the pain but eased by rotating in the opposite direction.
- *Aggravation of pain by coughing, sneezing or deep inspiration*. This can produce a sharp catching pain which, if severe, tends to implicate the costovertebral joint.
- *Relief of pain by firm pressure:* Patients may state that their back pain is eased by firm pressure such as leaning against the corner of a wall.

It is very important to be able to differentiate between chest pain due to vertebral dysfunction and that caused by myocardial ischaemia.

Key questions

- Can you recall injuring your back, such as by lifting something heavy?
- Did you have a fall onto your chest or back?
- Is the pain present during the night?
- Do you have low back pain or neck pain?
- Does the pain come on after walking or any strenuous effort?

- Does the pain come on after eating or soon after going to bed at night?
- Have you noticed a fever or sweating at any time, especially at night?
- Have you noticed a rash near where you have the pain?
- What drugs are you taking? Do you take drugs for arthritis or pain? Cortisone?
- What happens when you take a deep breath, cough or sneeze?

Physical examination

The examination of the thoracic spine is straightforward with the emphasis on palpation of the spine - central and laterally. This achieves the basic objective of reproducing the patient's symptoms and finding the level of pain. The ‘LOOK, FEEL, MOVE, X-RAY’ clinical approach is most appropriate for the thoracic spine.

Inspection

Careful inspection is important since it may be possible to observe at a glance why the patient has thoracic pain. Note the symmetry, any scars, skin creases and deformities, ‘flat spots’ in the spine, the nature of the scapulae or evidence of muscle spasm. Look for kyphosis and scoliosis.

Kyphosis may be generalised, with the back having a smooth uniform contour, or localised where it is due to a collapsed vertebra, such as occurs in an older person with osteoporosis. Generalised kyphosis is common in the elderly, especially those with degenerative spinal disease. In the young it may reflect the important Scheuermann's disorder.

The younger person in particular should be screened for scoliosis, which becomes more prominent on forward flexion (Figure 38.5). Look for any asymmetry of the chest wall, inequality of the scapulae and differences in the levels of the shoulders. A useful sign of scoliosis is unequal shoulder levels and apparent ‘winging’ of scapula. When viewed anteriorly a difference in the levels of the nipples indicates the presence of scoliosis, or other problems causing one shoulder to drop. Inspection should therefore take place with posterior, lateral (side) and anterior views.



Figure 38.5 Screening for adolescent idiopathic scoliosis: configuration of the thoracic spine on forward flexion

Palpation

The best position is to have the patient prone on the examination table with the thoracic spine preferably in slight flexion. This is achieved by lowering the top of the table.

Test passive extension of each joint with firm pressure from the pad of the thumbs or the bony hand (either the pisiform prominence or the lateral border of the fifth metacarpal). Spring up and down with a few firm oscillations, keeping the elbows straight, but being well above the patient. Ask the patient if the pressure reproduces the pain.

Apart from asking the patient 'Is that the pain?' note:

- the distribution of pain and its change with movement
- the range of movement
- the type of resistance in the joint
- any muscle spasm

Palpation must follow a set plan in order to reproduce the patient's pain. The sequence is as follows:

1. central - over spinous processes
2. unilateral - over apophyseal joints (2 - 3 cm from midline)

3. transverse - on side of spinous processes
4. unilateral - costotransverse junctions (4 - 5 cm from midline)
5. unilateral - over ribs (spring over posterior rib curve with ulnar border of hand, along axis of rib)

Movements

There are four main movements of the thoracic spine to assess, the most important of which is rotation, as this is the movement that so frequently reproduces the patient's pain where it is facetal joint or costovertebral in origin.

The movements of the thoracic spine and their normal ranges are:

1. Extension 30°
2. Lateral flexion L and R 30°
3. Flexion 90°
4. Rotation L and R 60°

Ask the patient to sit on the table with hands placed behind the neck and then perform the movements. Check these four active movements, noting any hypomobility, the range of movement, reproduction of symptoms and function and muscle spasm.

Investigations

The main investigation is an X-ray, which may exclude the basic abnormalities and diseases, such as osteoporosis and malignancy. If serious diseases such as malignancy or infection are suspected, and the plain X-ray is normal, a radionuclide bone scan may detect these disorders. CT scanning has a minimal role in the evaluation of thoracic spinal pain.

Other investigations to consider are:

- FBE and ESR
- serum alkaline phosphatase
- serum electrophoresis for multiple myeloma
- Bence-Jones protein analysis
- *Brucella* agglutination test
- blood culture for pyogenic infection and bacterial endocarditis
- tuberculosis studies
- HLA-B_{27} antigen for spondyloarthropathies
- ECG or ECG stress tests (suspected angina)
- gastroscopy or barium studies (peptic ulcer)
- MRI

THORACIC BACK PAIN IN CHILDREN

The most common cause of thoracic back pain in children is ‘postural backache’, also known as ‘TV backache’, which is usually found in adolescent schoolgirls and is a diagnosis of exclusion.

Important, although rare, problems in children include infections (tuberculosis, discitis and osteomyelitis) and tumours such as osteoid osteoma and malignant osteogenic sarcoma.

Dysfunction of the joints of the thoracic spine in children and particularly in adolescents is very common and often related to trauma such as a heavy fall in sporting activities or falling from a height (e.g. off a horse). Fractures, of course, have to be excluded.

Inflammatory disorders to consider are juvenile ankylosing spondylitis and spinal osteochondritis (Scheuermann's disorder), which may affect adolescent males in the lower thoracic spine (around T9) and thoracolumbar spine. The latter condition may be asymptomatic, but can be associated with back pain, especially as the patient grows older. It is the commonest cause of kyphosis.

It is important to screen adolescent children for idiopathic scoliosis, which may be without associated backache.

Scheuermann's disorder

This is a structural saggital plane deformity of unknown cause affecting the T7, 8, 9 or T11, 12 areas.

Typical features

- Age 11 - 17 years
- Males > females
- Lower thoracic spine
- Thoracic pain or asymptomatic
- Increasing thoracic kyphosis over 1 - 2 months
- Wedging of the vertebrae
- Pain in the wedge, especially on bending (only 20% present with pain)
- Short hamstrings
- Cannot touch toes
- Diagnosis confirmed by X-ray (lateral standing) - shows Schmorl's node and anterior vertebral body wedging

Treatment

- Explanation and support
- Extension exercises, avoid forward flexion
- Postural correction
- Avoidance of sports involving lifting and bending
- Consider bracing or surgery if serious deformity
- If detected early hyperextension body casts followed by a Milwaukee brace can prevent deformity

Adolescent idiopathic scoliosis

A degree of scoliosis is detectable in 5% of the adolescent population.[5] The vast majority of curves, occurring equally in boys and girls, are mild and of no consequence. Eighty-five per cent of significant curves in adolescent scoliosis occur in girls.[6] Inheritance is a factor. The highest incidence is in first-degree female relatives (12%). The scoliotic deformity de-velops at 10 years of age. Such curves appear during the peripubertal period, usually coinciding with the growth spurt. The screening test (usually in 12 - 14 year olds but ideally as early as 9) is to note the contour of the back on forward flexion (Figure 38.6). The routine physical screening check of an adolescent should include this area.



Figure 38.6 Screening for adolescent idiopathic scoliosis: testing asymmetry by forward flexion

The test

The subject stands with the feet parallel and together, and bends forward as far as possible with outstretched hands, palms facing each other, pointed between the great toes.

Investigation

A single erect PA spinal X-ray is sufficient;[7] the Cobb angle (Figure 38.7) is the usual measurement yardstick.


top of vertebra at top end of curve
50°
Cobb angle
bottom of vertebra at lower end of curve
V

Figure 38.7 Scoliosis: the Cobb method of curve measurement

Management

Aims

- To preserve good appearance - level shoulders and no trunk shift
- Prevent increasing curve in adult life: less than 45°
- Not to produce a straight spine on X-ray

Methods

- Braces:
  - Milwaukee brace (rarely used)
  - High density polyethylene underarm orthosis
  - To be worn for 20 - 22 hours each day until skeletal maturity is reached.
- Surgical correction: depends on curve and skeletal maturity

Guidelines for treatment

- Still growing:

< 20°	observe (repeat examination + X-ray)
20 - 30°	observe, brace if progressive
30 - 45°	brace
≥ 45 - 50°	operate

- Growth complete:

< 45°	leave alone
> 45°	operate

Referral to consultant: > 20°

THORACIC BACK PAIN IN THE ELDERLY

Thoracic back pain due to mechanical causes is not such a feature in the elderly although vertebral dysfunction still occurs quite regularly. However, when the elderly person presents with thoracic pain, a very careful search for organic disease is necessary. Special problems to consider are:

- malignant disease (e.g. multiple myeloma, lung, prostate)
- osteoporosis
- vertebral pathological fractures
- polymyalgia rheumatica
- Paget's disease (may be asymptomatic)
- herpes zoster
- visceral disorders:
  - ischaemic heart disease
  - penetrating peptic ulcer
  - oesophageal disorders
  - biliary disorders

DYSFUNCTION OF THE THORACIC SPINE

This is the outstanding cause of pain presenting to the practitioner and is relatively easy to diagnose. It is ofter referrned to as the thorauc hypomobility syndrom with the disorder aresene in the facet joints, costo venrteb the facet joints, costovertebrd joits and thoracie musculoligamentous structure singularly or costovertcbral. The most efficaceous treatment for painful dysfunctional problems varies according to practitioners with a special interest in this area. There are a paucity of studies and evidence supporting the multiplicity of therapies, especially focal injections and physical therapy. Many claim that special mobilisation and manipulation provides effective short-term, sometimes immediate, relief.

Typical profile[2]

Age:	• Any age, especially between 20 and 40 years.
History of injury:	• Sometimes slow or sudden onset
Site and radiation:	• Spinal and paraspinal - interscapular, arms, lateral chest, anterior chest, substernal, iliac crest
Type of pain:	• Dull, aching, occasionally sharp; severity related to activity, site and posture
Aggravation:	• Deep inspiration, postural movement of thorax, slumping or bending, walking upstairs, activities (e.g. lifting children, making beds), beds too hard or soft, sleeping or sitting for long periods
Association:	• Chronic poor posture
Diagnosis confirmation:	• Examination of spine, therapeutic response to manipulation

Management

First-line management

- Explanation with printed information
- Reassurance, including spontaneous recovery likely
- Continued activity according to pain level
- Back education program
- Analgesics (e.g. paracetamol)
- Posture education
- Spinal mobilisation and manipulation (if appropriate)

Spinal mobilisation and manipulation

Spinal mobilisation is helpful but the more forceful manipulative therapy produces better and quicker results. There are many techniques that can be employed, the choice depending on which part of the back is affected.[7] The sternal thrust (Nelson hold) technique is widely used for upper thoracic segments and the crossed pisiform technique (patient prone) or posteroanterior indirect thrust (patient supine - Figure 38.8) is claimed to be the most effective for the mid-thoracic spine. These is lover weii eardence that spinal manpulre complete with placebo[8]



Figure 38.8 Manipulation of the mid-thoracic spine by the posteroanterior indirect thrust technique Reproduced With Permission From C. Kenna And J. Murtagh. *Back Pain And Spinal Manipulation* (2ed). Oxford; Butterworth-Heinmann, 1997.

Thoracic disc protrusion

Fortunately, a disc protrusion in the thoracic spine is uncommon. This reduced incidence is related to the firm splintage action of the rib cage. Most disc protrusions occur below T9, with the commonest site, as expected, being T11 - 12.

The common presentation is back pain and radicular pain that follows the appropriate dermatome so disc protrusion should be considered in patients with neurological signs at thoracic levels. This may include a flaccid area of the lower abdominal musculature.

However, disc lesions in the thoracic spine are prone to produce spinal cord compression, manifesting as sensory loss, bladder in-continence and signs of upper motor neurone lesion. The disc is relatively inaccessible to surgical intervention, but over the past decade there has been a significant improvement in the surgical treatment of thoracic disc protrusions, due to the transthoracic lateral approach.

Syrinx

A syrinx usually coines to noter as aradulogeal fuding in the preseuce of thoracic back when it may in fact be asgnptomatic. It is a rase flued filled neurologual mcapter wittin the spinal card. It is ineally a wrgental anomaly but a neoplasw needs to be excluded. A syrinx usually luginsat the corrcal level and exdead doun. Treat conservatively but refen to an expert who may conseder interventim if it is arymptomatic.

Muscle injury

Muscular injuries such as tearing are uncommon in the chest wall. The strong paravertebral muscles do not appear to be a cause of chest pain, but strains of intercostal muscles, the serratus anterior and the musculotendinous origins of the abdominal muscles can cause pain. Injuries to these muscles can be provoked by attacks of violent sneezing or coughing or overstrain, for example, lifting a heavy suitcase down from an overhead luggage rack.

SCAPNLOTHORACIC JOINT DISORDERS[9]

The gliding plane between the scperla and thoracici wall permits a considerable range of scapnlar movement which contributes significantly to movement of the shoulder. Sevnal museles inoluding the rhomboids, serratus anterion and tevator scator help stabilse scapular movement and may be a souree of pain in the scapular region.

Snapping scapula

The patient complans of a loud cracking or snapping seckend upon abduction of the scapula. There is open associated repintus. Pain is felt along the medral scapular border. The palent may derdop a habit ctit to neurotieally click the shoulder back and forth.

On examination than is usually gennalised hypermobelety of the scapula, abnormal movement and tendernen to palpation along the medial edge on full abduction.

The cause cuncouineon may be an underlying bony abaormalely such as a bony spur or the supeuion border of the scapula to seauh for this poretilaly.

Treatment

- “Explanation and reassnance (of X-ray normal) other wree resect any bone abnormelely.
- Around repeated scapular morement and ‘trick’ movements.
- Appropriate exesuses under physiotherapy supervision.
- Infiltrate any very tenden area in the useley (with Care) with local anaerthetee and stevord.
- Deep unassage to the tender focus.

Scapulo costal syndrome

This wndrtim cause localised pain and tendernen, ofter terve along the upper part of the medeal scapnlar border with radration around the chest wall and shoulder gudle to the neck, pain is useally worse with pulonged whouldes use toevards the end of the day. It is coueuwnly use in typisets, gymnasts and other sportspesple. It is related to poor posture. The cause may include fricion between the scapula and the thoracie wall, scoliosis, trauma myofascial strain due to posture.

Treatmend

- Avoid the movements producinp the pin.
- Postine reducation ehesuses and scapula ststeling.
- Deep fricton massge.
- Local infections of local anaesthetee and corticosdered into the tender area.

Winging of the scapula

The asymmetry may not be apparent until the patien tries to contract the serratus anteurr agamst resistanu by pushory the outstsetched arun against the wall. Three way be parascapular discomforct. The comuers couse is neorgenic paralysis of the serratus anleum musle. Paralysis may tesult from ingway to the long thoracie here (5,6,7 wervs roots) such as a neek infnry or a direed blow to the suprascpular area from injury to the braclual pleus such as excessive carrying of heavy packs sevre traction on the arem, forcefulcervical manipulating, most cases settle spontanessuly althongh it may take 1-2 year.

FIBROMYALGIA, FIBROSITIS AND MYOFASCIAL TRIGGER POINTS

Fibromyalgia is relatively uncommon but when encountered presents an enormous man-agement problem. It is not to be confused with so-called fibrositis or tender trigger points. Referral to a specialist with expertise in this condition or to a multidisciplinary pain clinic for the definitive diagnosis is recommended.

Fibrositis is not a diagnosis but a symptom, indicating a localised area of tenderness or pain in the soft tissues, especially of the upper thoracic spine. It is probably almost always secondary to upper thoracic or lower cervical spinal dysfunction.

Myofascial trigger points

As described by Travell and Rinzler[10] a trigger point is characterised by:

- circumscribed local tenderness
- localised twitching with stimulation of juxtaposed muscle
- pain referred elsewhere when subjected to pressure

Trigger spots also tend to correspond to the acupuncture points for pain relief.

Treatment

Local injectio.n is relatively easy and may give excellent results. Identify the maximal point of pain and inject 5 - 8 mL of local anaesthetic (LA; e.g. lignocaine/lidocaine 1%) into the painful point (Figure 38.9). Post-injection massage or exercises should be performed.



Figure 38.9 Injection for myofascial trigger points

Don't:

- use large volumes of LA
- use corticosteroids
- cause bleeding

Do:

- use a moderate amount of LA (only)

Fibromyalgia syndrome[11]

The main diagnostic features are:

1. a history of widespread pain (neck to low back)
2. pain in 11 of 18 tender points on digital palpation

These points must be painful, not tender. Smythe and Moldofsky have recommended 14 of these points on a map as a guide for management[12] (see Figure 38.10).



Figure 38.10 Fibromyalgia syndrome: typical tender points (the tender point map represents the 14 points recommended for use as a standard for diagnostic or therapeutic studies)

Other features

- Female to male ratio ° 4:1
- Usual age 29 - 37 years: diagnosis 44 - 53 years
- Poor sleep pattern
- Fatigue (similar to chronic fatigue syndrome)
- Psychological disorders (e.g. anxiety, depression, tension headache, irritable digestive system)

This disorder is very difficult to treat and is usually unresponsive in the long term to passive physical therapy or injections.[13] Patients require considerable explanation, support and reassurance. Best evidence to date supports the value of educational programs and regular aerobic exercise.[14]

Treatment

- Explanation, reassurance and counselling
- Attention to sleep disorders, stress factors and physical factors
- Relaxation program
- Rehabilitation exercise program (e.g. walking, swimming or cycling)
- Use paracetamol for first-line analgesia

Medication (often disappointing):

- antidepressants (of proven short-term value) e.g. amitriptyline 10 - 75 mg (o) nocte or doxepin 25 - 100 mg (o) nocte[15, 16] *or*
- clonazepam (Rivotril) 0.5 mg bd

Note:NSAIDs are of no proven benefit. New agents the antidepressants duloxetine and the minacipran pregabalin are being exaludlid.

SERIOUS PITFALLS

The following points regarding serious vertebral organic disease are worth repeating in more detail.

Metastatic disease[17]

Secondary deposits in the thoracolumbar spine may be the first presenting symptoms of malignant disease. Any patient of any age presenting with progressive severe night pain of the back should be regarded as having a tumour and investigated with a technetium bone scan as part of the primary investigations.

Secondary deposits in the spine can lead to rapid onset paralysis due to spinal cord infarction. Many such metastases can be controlled in the early stages with radiotherapy.

Multiple myeloma

Osteoporotic vertebral body collapse should be diagnosed only when multiple myeloma has been excluded. Investigations should include an ESR, Bence-Jones protein analysis, and immunoglobulin electrophoresis.

Early treatment of multiple myeloma can hold this disease in remission for many years and prevent crippling vertebral fractures.

Infective discitis and vertebral osteomyelitis

Severe back pain in an unwell patient with fluctuating temperature (fever) should be considered as infective until proved otherwise. Investigations should include blood cultures, serial X-rays and nuclear bone scanning. Biphasic bone scans using technetium with either indium or gallium scanning for white cell collections usually clinch this diagnosis.

Strict bed rest with high-dose antibiotic therapy is usually curative. If left untreated, vertebral end plate and disc space collapse is common and extremely disabling.

WHEN TO REFER

- Persistent pain or dysfunction - refer to a physical therapist.
- Evidence or suspicion of a sinister cause (e.g. neoplasia, infective discitis/osteomyelitis in a child).
- Suspicion of cardiac or gastrointestinal referred (persistent) pain.
- Significant idiopathic adolescent scoliosis or kyphosis (e.g. Scheuermann's disorder).

Practice tips

- Feelings of anaesthesia or paraesthesia associated with thoracic spinal dysfunction are rare.
- Thoracic back pain is frequently associated with cervical lesions.
- Upper thoracic pain and stiffness is common after ‘whiplash’.
- The T4 syndrome of upper to mid-thoracic pain with radiation (and associated paraesthesia) to the arms is well documented.
- Symptoms due to a fractured vertebra usually last 3 months and to a fractured rib 6 weeks.
- The pain of myocardial ischaemia, from either angina or myocardial infarction, can cause referred pain to the interscapular region of the thoracic spine.
- Beware of the old trap of herpes zoster in the thoracic spine, especially in the older person.
- Consider multiple myeloma as a cause of an osteoporotic collapsed vertebra.
- Examine movements with the patient sitting on the couch and hands clasped behind the neck.
- Spinal disease of special significance in the thoracic spine includes osteoporosis and neoplasia, while disc lesions, inflammatory diseases and degenerative diseases (spondylosis) are encountered less frequently than with the cervical and lumbar spines.

- It is imperative to differentiate between spinal and cardiac causes of chest pain: either cause is likely to mimic the other. A working rule is to consider the cause as cardiac until the examination and investigations establish the true cause.
- Always X-ray the thoracic spine following trauma, especially after motor vehicle accidents, as wedge compression fractures (typically between T4 and T8) are often overlooked.

REFERENCES

1. NH&MRC. *Evidence based management of Sacute Musculoskeletal pause a guide clincians Australian Acute Musculoskeletal Pain - A Guide for Clinicians*. Australian Acute Musculoskeletal Pain Guidelines Group, Canberra: Australian Government, 2004: 30 - 4.
2. Kenna C, Murtagh J. *Back Pain and Spinal Manipulation* (2nd edn). Oxford: Butterworth-Heinemann, 1997: 165 - 74.
3. Chua WL. Thoracic spinal pain - a review. Australas Musculoskeletal Med, 1996; 1: 13 - 22.
4. McGuckin N. The T4 syndrome. In: Grieve GD (ed.) *Modern Manual Therapy of the Vertebral Column*. London: Churchill Livingstone, 1986: 370 - 6.
5. Stephens J. Idiopathic adolescent scoliosis. Aust Fam Physician, 1984; 13: 180 - 4.
6. Kane WJ, Moe JH. A scoliosis prevalence survey in Minnesota. Clin Orthop, 1970; 69: 216 - 18.
7. Anonymous. *The Easter Seal Guide to Children's Orthopaedics*. Toronto: The Easter Seal Society, 1982: 64 - 7.
8. Scholer L. Effectweness of spinal manipulation thapy in the treatment of mechanical thoracic back pain. J of Manipulative and Physiological Therapeutics, 2001; 24: 394 - 401.
9. Corrigan B, mattand G. *Practical Orthopedie Medicine*. Sydney Butterworths 1986:384 - 5.
10. Travell J, Rinzler SH. The myofascial genesis of pain. Postgrad Med, 1952; 11: 425 - 34.
11. Reilly P, Littlejohn G. Current thinking on fibromyalgia syndrome. Aust Fam Physician, 1990; 19: 1505 - 76.
12. Smythe HA, Moldofsky H. Two contributions to understanding of the ‘fibrositis’ syndrome. Bull Rheum Dis, 1977; 28: 928 - 31.
13. McIndoe R, Littlejohn G. Management of fibromyalgia and regional pain syndromes. Mod Med Aust, 1995; 36(2): 56 - 69.
14. McCain GA, Bell DA et al. A controlled study of the effects of a supervised cardiovascular fitness training program on the manifestation of primary fibromyalgia. Arthritis Rheum, 1988; 31: 1135 - 41.
15. Jaeschke R, Adachi J, Guyatt G et al. Clinical usefulness of amitriptyline in fibromyalgia: the results of 23 N-of-I randomised controlled trials. J Rheumatol, 1991; 18: 447 - 51.
16. Mashford ML (Chair). *Therapeutic Guidelines: Analgesic* (Version 3). Melbourne: Therapeutic Guidelines Ltd, 1998: 91 - 2.
17. Young D, Murtagh J. Pitfalls in orthopaedics. Aust Fam Physician, 1989; 18: 653 - 4.

CHAPTER 39. BRUISING AND BLEEDING

My pa is one mask of brooses both blue and green.

CHARLES DICKENS (1812 - 70), NICHOLAS NICKLEBY

BRUISING AND BLEEDING

Many patients present with the complaint that they bruise easily but only a minority turn out to have an underlying blood disorder. Purpura is bleeding into the skin or mucous membranes, appearing as multiple small haemorrhages that do not blanch on pressure. Smaller purpuric lesions that are 2 mm or less in diameter (pinhead size) are termed petechiae, while larger purpuric lesions are called ecchymoses (Figure 39.1).



Figure 39.1 Purpuric rash (petechiae and ecchymoses)

Bruises are large areas of bleeding that are a result of subcutaneous bleeding. If bruising is abnormal and out of proportion to the offending trauma, then a disturbance of coagulation is suggested (Figure 39.2).



Figure 39.2 Severe bleeding in a patient with systemic fibrinolysis in a diabetic patient. Note the bleeding following insulin injections into the abdominal wall and an injection into the shoulder joint. *Photo courtesy Hatem Salem*

DIFFERENTIAL DIAGNOSIS

'Palpable purpura' due to an underlying systemic vasculitis is an important differential problem. The petechiae are raised so finger palpation is important. The cause is an underlying vasculitis affecting small vessels (e.g. polyarteritis nodosa).

The decision to investigate is difficult because decisions have to be made about which patients warrant investigation and whether the haemostatic defect is due to local or systemic pathology.[1] The ability to identify a bleeding disorder is important because of implications for surgery, pregnancy, medication and genetic counselling.

KEY FACTS AND CHECKPOINTS

- Purpura ° petechiae + ecchymoses.
- Abnormal bleeding is basically the result of disorders of (1) the platelet, (2) the coagulation mechanism, or (3) the blood vessel.
- There is no substitute for a good history in the assessment of patients with bleeding disorders.
- An assessment of the personal and family histories is the first step in the identification of a bleeding disorder.
- When a patient complains of 'bruising easily' it is important to exclude thrombocytopenia due to bone marrow disease and clotting factor deficiencies such as haemophilia.
- The commonest cause of an acquired bleeding disorder is drug therapy (e.g. aspirin, NSAIDs, cytotoxics and oral anticoagulants).
- In general, bleeding secondary to platelet defects is spontaneous, associated with a petechial rash and occurs immediately after trauma or a cut wound.[1] The bleeding is usually mucosal (e.g. bleeding from gingiva, menorrhagia, epistaxis and petechiae).
- Laboratory assessment should be guided by the clinical impression.
- Bleeding caused by coagulation factor deficiency is usually traumatic and delayed (e.g. haemorrhage occurring 24 hours after a dental extraction in a haemophiliac).
- The routine screening tests for the investigation of patients with bleeding disorders can occasionally be normal despite the presence of a severe haemorrhagic state.

CAUSES OF CLINICAL DISORDERS

The three major mechanisms of systemic bleeding disorders are:

1. coagulation deficiencies (reduction or inhibition of circulatory coagulation factors)
2. platelet abnormalities: of platelet number or function

3. vascular defects: of vascular endothelium

A list of differential diagnoses of systemic bleeding disorders is presented in Table 39.1.[1]

Table 39.1 Differential diagnoses of systemic bleeding disorders

Vascular disorders (a) Inherited • hereditary haemorrhagic telangiectasia • Marfan syndrome (b) Acquired • purpura simplex • senile purpura • Henoch-Schönlein purpura • steroid purpura • scurvy
Coagulation factor deficiency or inhibitor (a) Inherited • haemophilia A • haemophilia B • von Willebrand disease (b) Acquired • disseminated intravascular coagulation • liver disease • vitamin K deficiency • oral anticoagulant therapy or overdosage
Thrombocytopenia (a) Inherited • Fanconi syndrome • amegakaryocytic thrombocytopenia (b) Acquired • immune thrombocytopenic purpura • drug-induced thrombocytopenia • disseminated intravascular coagulation • bone marrow replacement or failure • thrombotic thrombocytopenic purpura • post-transfusion purpura • splenic pooling
Functional platelet disorders (a) Inherited • Glanzmann's thrombasthenia • Bernard-Soulier syndrome • storage pool deficiency (b) Acquired • drug-induced • uraemia

• myeloproliferative disorders
• dysproteinaemias

Source: After Mitchell et al.[1] Adapted from Bleeding disorders, MIMS Disease Index (2nd edn) 1996 with permission of *MIMS* Australia, a division of MediMedia Australia Pty Limited.

THE CLINICAL APPROACH

Differentiation of coagulation factor deficiencies and platelet disorders as the cause of a bleeding problem can usually be determined by a careful evaluation of the history and physical examination.

History

Factors that suggest the presence of a systemic bleeding defect include:

- spontaneous haemorrhage
- severe or recurrent haemorrhagic episodes
- bleeding from multiple sites
- bleeding out of proportion to the degree of trauma

If a bleeding diathesis is suspected it is essential to determine whether local pathology is contributing to the blood loss (e.g. postoperative bleeding, postpartum bleeding, gastrointestinal haemorrhage).

Diagnostic tips

- Platelet abnormalities present as early bleeding following trauma.
- Coagulation factor deficiencies present with delayed bleeding after initial haemostasis is achieved by normal platelets.
- A normal response to previous coagulation stresses (e.g. dental extraction, circumcision or pregnancy) indicates an acquired problem.
- If acquired, look for evidence of drugs, malignancy and liver disease.
- A diagnostic strategy is outlined in Table 39.2.

Table 39.2 Purpura: diagnostic strategy model

Q. Probability diagnosis
A. Simple purpura (easy bruising syndrome)
Senile purpura
Corticosteroid-induced purpura
Immune thrombocytopenic purpura
Henoch-Schönlein purpura
Q. Serious disorders not to be missed
A. Malignant disease
• leukaemia
• myeloma
Aplastic anaemia
Myelofibrosis
Severe infections
• septicaemia
• meningococcal infection
• measles
• typhoid
Disseminated intravascular coagulation

Thrombotic thrombocytopenic purpura
Q. Pitfalls (often missed) A. Haemophilia A, B von Willebrand disease Post-transfusion purpura Trauma (e.g. domestic violence, child abuse) *Rarities* • hereditary telangiectasia (Osler-Weber-Rendu syndrome) • Ehlers-Danlos syndrome • scurvy • Fanconi syndrome
Q. The masquerades A. Drugs • chloramphenicol • corticosteroids • sulphonamides • quinine/quinidine • thiazide diuretics • NSAIDs • cytotoxics • oral anticoagulants Anaemia • aplastic anaemia
Q. Psychogenic factors A. Factitial purpura

Family history

A positive family history can be a positive pointer to the diagnosis:

- sex-linked recessive pattern
  - haemophilia A or B
- autosomal dominant pattern
  - von Willebrand disease
  - dysfibrinogenaemias
- autosomal recessive pattern
  - deficiency of coagulation factors V, VII and X

Enquire whether the patient has noticed blood in the urine or stools and whether menorrhagia is present in women. A checklist for a bleeding history is presented in Table 39.3. The actual size and frequency of the bruises should be recorded where possible and if none are present at the time of the consultation the patient should return if any bruises reappear.

Table 39.3 Checklist for a bleeding history

- Skin bruising
- Epistaxis
- Injury
- Domestic violence
- Tonsillectomy
- Other operations
- Childbirth
- Haematuria

• Menorrhagia	• Rectal bleeding
• Haemarthrosis	• Drugs
• Tooth extraction	• Family history
	• Co-morbidities, e.g.
	- liver disease
	- kidney disease

Key questions

- How long has the problem been apparent to you?
- Do you remember any bumps or falls that might have caused the bruising?
- What sort of injuries cause you to bruise easily?
- Have you noticed bleeding from other areas such as your nose or gums?
- Has anyone in your family had a history of bruising or bleeding?
- What is your general health like?
- Do you have any tiredness, weight loss, fever or night sweats?
- Did you notice a viral illness or sore throat beforehand?
- How much alcohol do you drink?
- What happened in the past when you had a tooth extracted?
- Have you ever had painful swelling in your joints?

Medication record

It is mandatory to obtain a complete drug history. Examples of drugs and their responses are:

- vascular purpura
  - prednisolone/other steroids
- thrombocytopenia
  - cytotoxic drugs
  - gold
  - heparin
  - phenylbutazone
  - sulphonamides
  - quinine, quinidine
  - thiazide diuretics
  - chloramphenicol
- functional platelet abnormalities
  - aspirin
  - NSAIDs
- coagulation factor deficiency
  - warfarin

Physical examination

Careful examination of the skin is important. Note the nature of the bleeding and the distribution of any rash, which is characteristic in Henoch-Schönlein purpura. Senile purpura in the elderly is usually seen over the dorsum of the hands, extensor surface of the forearms and the shins.

Note the lips and oral mucosa for evidence of hereditary telangiectasia. Gum hypertrophy occurs in monocytic leukaemia. Search for evidence of malignancy, such as sternal tenderness, lymphadenopathy and splenomegaly. Examine the ocular fundi for evidence of retinal haemorrhages. Urinalysis, searching for blood (microscopic or macroscopic), is important.

Investigations

The initial choice of investigations depends upon the bleeding pattern.

If coagulation defect suspected:

- prothrombin time (PT), i.e. INR

- activated partial thromboplastin time (APTT)
- fibrinogen level

If platelet pathology suspected:

- platelet count
- skin bleeding time (of doubtful value)
- platelet function analyser (PFA-100)

The full blood examination and blood film is useful in pinpointing the aetiology. Platelet morphology gives a diagnostic guide to inherited platelet disorders. The skin-bleeding time as a screening test of haemostasis has been shown recently to be severely limited by its lack of specificity and sensitivity and its routine use cannot be recommended. It is not a useful predictor of surgical risk of haemorrhage.[1, 2] Other sophisticated tests, such as von Willebrand's screen and platelet aggregation, e.g. PFA-100, can be advised by the consulting haematologist. One of considerable value is the bone marrow examination, which is useful to exclude the secondary causes of thrombocytopenia, such as leukaemia, other marrow infiltrations and aplastic anaemia.

A summary of appropriate tests is presented in Table 39.4 and of blood changes for some coagulation factor deficiencies in Table 39.5.

Table 39.4 Laboratory investigation checklist for the easy bruiser

- Full blood count
- Platelet count
- Prothrombin time (INR)
- Activated partial thromboplastin time

Table 39.5 Blood changes for specific coagulation factor disorders

	Haemophilia A	von Willebrand disease	Vitamin K deficiency
PT	Normal	Normal	↑
APTT	↑	↑	↑

ABNORMAL BLEEDING IN CHILDREN

Abnormal bleeding in children is not uncommon and once again the clinical history, particularly the past and family history, provides the most valuable information. It is important to keep non-accidental injury, such as child abuse, in mind in the child presenting with ‘easy bruising’. However, it is appropriate to exclude a bleeding disorder, especially a platelet disorder.

Coagulation disorders, including haemophilia and von Willebrand disease, are usually suspected on clinical grounds because of widespread bruising or because of prolonged bleeding following procedures such as circumcision and tonsillectomy.

A common condition is haemorrhagic disease of the newborn, which is a self-limiting disease usually presenting on the second or third day of life because of a deficiency of coagulation factors dependent on vitamin K. The routine use of prophylactic vitamin K in the newborn infant has virtually eliminated this problem.

Idiopathic (immune) thrombocytopenic purpura (ITP) is the commonest of the primary platelet disorders in children. Both acute and chronic forms have an immunological basis. The diagnosis is based on the peripheral blood film and platelet count. The platelet count is commonly below 50 000/mm^3 × (50 10^9/L). Spontaneous remission within 4 to 6 weeks occurs with acute ITP in childhood.[2]

The commonest vascular defects in childhood are:

- anaphylactoid (Henoch-Schönlein) purpura
- infective states
- nutritional deficiency (usually inadequate dietary vitamin C)

Henoch-Schönlein purpura (HSP)

HSP is the commonest vasculitis of children. It affects small vessels, producing a leucocytoclastic vasculitis with a classic triad of non-thrombocytopenic purpura, large joint arthritis and abdominal pain. It is diagnosed clinically by the characteristic distribution of the rash (which is a palpable purpura) over the lower limbs, extending onto the buttocks (Figure 39.3), but it can also involve the upper limbs, trunk and even the face.

The onset of HSP typically follows an upper respiratory tract infection including a group A streptococcal tonsillo-pharyngitis.


abdominal pain
haematuria
typical sites of purpura (palpable)
arthritis

Figure 39.3 Henoch-Schönlein purpura: typical distribution

The bleeding time, coagulation time and platelet counts are normal. The prognosis is good; most recover fully in a few months.

Features of HSP

- All ages, mainly in children
- Rash, mainly on buttocks and legs (Figure 39.4)[3]



Figure 39.4 Henoch-Schönlein purpura in a 5 year old boy showing the typical distribution of the rash on the lower limbs

- Rash can occur on hands, arms and trunk

- Arthritis: mainly ankles and knees
- Abdominal pain - colicky (vasculitis of GIT)
- Haematuria (reflects nephritis)

Associations

- Kidney involvement - deposition of IgA immune complex
- Melaena
- Intussusception
- Scrotal involvement

Investigations

- FBE
- urine - protein and blood

Management

- Largely symptomatic - analgesics
- No specific therapy
- Short course of steroids for abdominal pain (if intussusception excluded)
- If haematuria: follow up urine microscopy and kidney function

Tip: beware of CKF

DxT: arthralgia + purpuric rash ± abdominal pain ° Henoch Schonlein purpura

Infective states

The purpura associated with severe infections, such as meningococcaemia and other septicaemias, is due primarily to a severe angiitis. Disseminated intravascular coagulation usually follows.[2]

ABNORMAL BLEEDING IN THE ELDERLY

The outstanding causes are senile purpura and purpura due to steroids.[4] The cause in both instances is atrophy of the vascular supporting tissue.

VASCULAR DISORDERS

The features are:

- easy bruising and bleeding into skin
- ± mucous membrane bleeding
- investigations normal

Simple purpura (easy bruising syndrome)

This is a benign disorder occurring in otherwise healthy women usually in their twenties or thirties. The feature is bruising on the arms, leg and trunk with minor trauma. The patient may complain of heavy periods. Major challenges to the haemostatic mechanism, such as dental extraction, childbirth and surgery, have not been complicated by excessive blood loss.

Factitial purpura

Unexplained bruising or bleeding may represent selfinflicted abuse or abuse by others. In self-inflicted abuse the bruising is commonly on the legs or areas within easy reach of the patient.

PLATELET DISORDERS

The features are:

- petechiae ± ecchymoses
- bleeding from mucous membranes
- platelet counts > 50 000/mm^3 (50 × 10^9/L)

Immune thrombocytopenic purpura

Essential features:

- easy bruising
- epistaxis and menorrhagia common
- no systemic illness
- splenomegaly rare
- isolated thrombocytopenia
- other blood cells normal
- otherwise normal physical examination
- normal bone marrow with normal or increased
- megakaryocytes (acute leukemia should be excluded)

DxT: bruising + oral bleeding + epistaxis ° ITP

The two distinct types caused by immune destruction of the platelets are:

- acute thrombocytopenia of childhood - usually in children, usually post-viral
- chronic ITP - autoimmune disorder, usually in adult women; all cases should be referred to a specialist unit

Acute thrombocytopenia of childhood

This is caused by a reaction to a virus infection resulting in the production of cross-reacting antibodies against platelets.

There is an early risk of spontaneous haemorrhage, so refer/admit to hospital.

The prognosis is good, invariably self - limiting - 90% resolve in 6 months. The rest pass into chronic ITP.

Bleeding is treated with ã globulin or steroids.

Chronic ITP

Chronic ITP rarely undergoes spontaneous remission and may require treatment with prednisolone. Some require splenectomy but this operation is avoided where possible, especially in young children, because of the subsequent risk of severe infection, particularly with *Streptococcus pneumoniae*.[4] (Refer p. 408.)

Thrombotic thrombocytopenic purpura

TTP is an uncommon life-threatening syndrome of haemolytic anaemia, thrombocytopenia and extremely high LDH. Clinical features include fever (non-infectious), neurologic and kidney abnormalities.

COAGULATION DISORDERS

The features are:

- ecchymoses
- haemarthrosis and muscle haematomas
- usually traumatic and delayed

The inherited disorders such as haemophilia A and B are uncommon and involve deficiency of one factor only. The acquired disorders, such as disseminated intravascular coagulation (DIC), occur more commonly and invariably affect several anticoagulation factors (Table 39.6).

Table 39.6 International nomenclature of clotting factors

Factor	Common synonyms
I	Fibrinogen[*]

II	Prothrombin[*]
III	No longer used
IV	No longer used
V[*]	No longer used
VI	No longer used
VII[*]	Tissue factor
VIII	Antihaemophilic factor Antihaemophilic globulin
IX[*]	Christmas factor, plasma thromboplastin component
X[*]	Stuart-Prower factor
XI[*]	No longer used
XII	Hageman factor, contact factor
XIII	Fibrin stabilising factor

[*] Common terminology in use.

INHERITED COAGULATION DISORDERS

A list of the inherited bleeding disorders is included in Table 39.1. The better known disorders are von Willebrand disease, haemophilia A and haemophilia B (Christmas disease).

von Willebrand disease

This is the most common disorder of haemostasis (incidence 1% of population) and is usually a mild problem with an excellent prognosis.[5] There are about five types.

Essential features:

- autosomal dominant inheritance (common types)
- equal sex incidence
- prolonged bleeding time
- bleeding tendency exacerbated by aspirin
- defective platelet adhesion at site of trauma combined with factor VIII deficiency[5]
- APTT prolonged
- menorrhagia and epistaxis common
- haemarthroses rare

DxT: increased bleeding - 1. incisions 2. dental 3. mucosal ° von Willebrand disease

Treatment

- No specific treatment
- Avoid aspirin
- Be cautious of surgical and dental procedures
- Preparations that help include desmopressin acetate (DDAVP), factor VIII concentrates and tranexamic acid

Haemophilia A

Essential features:

- spontaneous haemarthroses, especially knees, ankles and elbows, are almost pathognomic
- X-linked recessive pattern of inheritance
- invariably only males affected (1 in 5000)
- females theoretically affected if haemophiliac father and carrier mother

- the human factor gene has long been identified severity levels:
  - severe - bleed spontaneously
  - moderate - bleed with mild trauma or surgery
  - mild - bleed after major trauma or surgery
- deficiency of factor VIII
- APTT prolonged
- normal prothrombin time and fibrinogen
- many seropositive for HIV, hepatitis B or (factor VIII concentrate transmission)
- low platelet count should suspect
- HIV-associated ITP5

DxT: spontaneous haemarthrosis + muscle bleeds + delayed bleeding ° haemophilia A

Treatment:

- Infusion of recombinant factor VIII concentrates5
- Avoid aspirin

Haemophilia B (Christmas disease)

- Identical clinical features to haemophilia A
- Also an X-linked recessive hereditary disorder
- Incidence of 1 in 30 000
- Deficiency of coagulation factor IX
- Same laboratory findings as haemophilia A apart from specific factor assays
- Treatment is with recombinant factor IX concentrates

SPLENECTOMY

Main indications:

- immune thrombocytopenic purpura
- haemolytic anaemias especially hereditary spherocytosis
- hypersplenism
- trauma
- Hodgkins/non-Hodgkins lymphoma

Post splenectomy management[6]

Immediate problem is thrombocytosis (↑ platelets to 600-1000 × 10^9/L) for 2-3 weeks with risk of thromboembolism.

Long-term risk is overwhelming infection (mainly pneumococcus), especially in young children in the first 2 years post-splenectomy.

Prophylaxis

- Education about risks and early recognition of infection (special care with malaria)
- Pneumococcal immunisation - give 2 - 3 weeks preoperative, repeat 5-yearly
- Haemophilus type B vaccine - once only if not immunised
- Meningococcus vaccine - every 5 years
- Long-term penicillin: amoxycillin early or
- phenoxymethypenicillin bd

MANAGEMENT PRINCIPLES[1]

- Make the correct diagnosis.
- Stop or avoid drugs affecting the haemostatic system.

- Control bleeding episodes with appropriate drugs, blood products and local measures, such as simple compression or topical haemostatic agents.
- Infuse appropriate blood components for the treatment of coagulation factor deficiencies and some platelet disorders (e.g. factor VIII for haemophilia A, fresh frozen plasma for multiple factor deficiency).

Practice tips

- A careful history and physical examination will usually pinpoint the cause of the bleeding disorder.
- Drug therapy can lead to unmasking of pre-existing haemostatic disorders (e.g. platelet dysfunction induced by aspirin may cause spontaneous bleeding in patients with underlying von Willebrand disease).
- Think of DIC in any acutely ill patient with abnormal bleeding from sites such as the mouth or nose, venepuncture or with widespread ecchymoses. The clinical situations are numerous, such as septicaemia, obstetric emergencies, disseminated malignant disease, falciparum malaria and snake bites.

- Refer patients with identified defects to a consultant haematologist or haemophilia centre.
- Supervise advanced planning in patients intending pregnancy, surgery or dental extraction.

When to refer[1]

- Management of haemorrhage is not amenable to simple measures such as local therapy with simple compression and other measures.
- Elective surgery or pregnancy is being planned.

REFERENCES

1. Mitchell CA, Dear A, Salem H. Bleeding disorders. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 69 - 71.
2. McPherson J, Street A. Tests of haemostasis: detection of the patient at risk of bleeding. Aust Prescriber, 1995; 18(2): 38 - 40.
3. Robinson MJ. *Practical Paediatrics* (2nd edn). Melbourne: Churchill Livingstone, 1990: 349 - 56.
4. Kumar PJ, Clark ML. *Clinical Medicine* (5th edn). London: Saunders, 2003: 458 - 60.
5. Tierney LM, McPhee SJ, Papadakis M. Current Medical Diagnosis and Treatment (45th edn). New York: The McGraw-Hill Companies, 2006: 526 - 8.
6. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 181.

CHAPTER 40. CHEST PAIN

> There is a disorder of the breast marked with strong and peculiar symptoms, considerable for the kind of danger belonging to it. The seat of it, and sense of strangling, and anxiety with which it is attended, may make it not improperly be called angina pectoris.
>
> WILLIAM HEBERDEN (1710 - 1801)

CHEST PAIN

The presenting problem of chest pain is common yet very threatening to both patient and doctor because the underlying cause in many instances is potentially lethal, especially with chest pain of sudden onset. The causes of acute chest pain are summarised and presented in Figure 40.1.

KEY FACTS AND CHECKPOINTS

- Chest pain represents myocardial infarction until proved otherwise.
- Immediate life-threatening causes of spontaneous chest pain are:
  - myocardial infarction (MI) and unstable angina
  - pulmonary embolism
  - aortic dissection
  - tension pneumothorax
- The main differential diagnoses of myocardial infarction include angina, aortic dissection, pericarditis, oesophageal reflux and spasm, and hyperventilation with anxiety.
- The history remains the most important clinical factor in the diagnosis of ischaemic heart disease. With angina a vital clue is the reproducibility of the symptom.


Life-threatening
Mediastinum
• mediastinitis
• oesophageal rupture
Great vessels
• dissecting aneurysm
• pulmonary embolus
Heart
• myocardial infarction
• angina
• pericarditis
Respiratory
• pulmonary—embolus
—infarct
• pneumothorax
• pneumonia/pleurisy
Non-life-threatening
Functional pain
• anxiety
• hyperventilation
Spinal dysfunction
• cervical dysfunction
• thoracic dysfunction
Chest wall
• trauma
• pathological fractures
• costochondritis
Infection
• herpes zoster
• Bornholm disorder
GIT
• oesophagitis/acid reflux
• oesophageal spasm
• peptic ulcer
• aerophagy
• gall bladder disease

Figure 40.1 Causes of acute chest pain

A DIAGNOSTIC APPROACH

The safety diagnostic model (Table 40.1) can be used to analyse chest pain according to the five self-posed questions.

Table 40.1 Chest pain: diagnostic strategy model

Q. Probability diagnosis

A. Musculoskeletal (chest wall)
Psychogenic
Angina

Q. Serious disorders not to be missed

A. Cardiovascular

- myocardial infarction/unstable angina
- aortic dissection
- pulmonary embolism

Neoplasia

- carcinoma lung
- tumours of spinal cord and meninges

Severe infections

- pneumonia/pleurisy
- mediastinitis
- pericarditis

Pneumothorax

Q. Pitfalls (often missed)

A. Mitral valve prolapse
Oesophageal spasm
Gastro-oesophageal reflux
Herpes zoster
Fractured rib (e.g. cough fracture)
Spinal dysfunction

Rarities

- Bornholm disorder (pleurodynia)
- cocaine inhalation
- hypertrophic cardiomyopathy

Q. Seven masquerades checklist

A. Depression	possible
Diabetes	-
Drugs	-
Anaemia	indirect
Thyroid disorder	-
Spinal dysfunction	
UTI	-

Q. Is the patient trying to tell me something?

A. Consider functional causes, especially anxiety with hyperventilation

Note: Chest pain is myocardial ischaemia until proved otherwise.

Probability diagnosis

The commonest causes encountered in general practice are musculoskeletal or chest wall pain and psychogenic disorders. The former is a very important yet often overlooked cause and sometimes inappropriately referred to as fibrositis or neuralgia. Causes include costochondritis, muscular strains, dysfunction of the sternocostal joints and dysfunction of the lower cervical spine or upper thoracic spine, which can cause referred pain to various areas of the chest wall. Angina is common and must always be considered. If angina-like pain lasts longer than 15 minutes myocardial infarction must be excluded.

Serious disorders not to be missed

The usual triad of malignancy, myocardial ischaemia and severe infections (Table 40.1) must be considered. In addition other cardiovascular catastrophes, such as a dissecting aortic aneurysm and pulmonary embolus, must be excluded, albeit uncommon, especially in those at risk. Spontaneous pneumothorax should also be considered, especially in a young male of slight build. Malignancies of the lung are relatively common and may present as pain when the previously asymptomatic tumour invades nerves or the spine.

The severe infections that cause chest pain include pneumonia/pleurisy, pericarditis and mediastinitis.

Pitfalls

Unfortunately, myocardial infarction and angina are often missed. Referred pain from spinal dysfunction, especially if referred anteriorly, is commonly overlooked. Other pitfalls include a cough fracture of a rib, herpes zoster (prior to the eruption) and gastrointestinal disorders, including oesophageal spasm, reflux and cholecystitis. Oesophageal problems may be clinically indistinguishable from angina. Mitral valve prolapse can cause chest pain although the mechanism is unclear: think of it in an unwell female prone to palpitations and chest pain. The pain tends to be sharp, fleeting, non-exertional and located near the cardiac apex.

General pitfalls include:

- not being ‘coronary aware’ in patients presenting with chest pain
- referred pain from spinal disorders, especially of the lower cervical spine - one of the great pitfalls in medical practice
- labelling chest pain as psychological in an anxious patient presenting with acute chest pain
- assuming that pain radiating down the inside of the leftarm is always cardiac in origin
- being unaware that up to 20% of myocardial infarctions are silent, especially in elderly patients, and that pulmonary embolism is often painless

Seven masquerades checklist

Of this group spinal dysfunction is possible. Disc lesions from the lower cervical spine are unlikely to cause chest wall pain, but dysfunction of the facet joints of this area of the spine and the upper thoracic spine is a common cause of referred pain to the chest wall. Nerve root pain from spinal problems is rarely found in the chest wall. Pathological fractures secondary to osteoporosis or malignancy in the vertebrae cause posterior wall pain.

Psychogenic considerations

With psychogenic causes the pain can occur anywhere in the chest, and tends to be continuous and sharp or stabbing rather than constricting. Associated symptoms include palpitations, deep breathing, fatigue, tremor, agitation and anxiety. Abnormal stress, tension, anxiety or depression may precipitate the pain, which often lasts hours or days.

THE CLINICAL APPROACH

History

A meticulous history of the behaviour of the pain is the key to diagnosis. The pain should be analysed into its usual characteristics: site and radiation, quality, intensity, duration, onset and offset, precipitating and relieving factors, and associated symptoms. Association with serious medical problems such as diabetes, Marfan syndrome, anaemia and SLE should be kept in mind. The ability to take a detailed history will obviously be limited with severe acute pain.

Associated symptoms

- *Syncope*. Consider myocardial infarction, pulmonary embolus and dissecting aneurysm.
- *Pain on inspiration*. Consider pleurisy, pericarditis, pneumothorax and musculoskeletal (chest wall pain).
- *Thoracic back pain*. Consider spinal dysfunction, myocardial infarction, angina, aortic dissection, pericarditis and gastrointestinal disorders such as a peptic ulcer, cholecystitis and oesophageal spasm.

Key questions

- Where exactly do you get the pain?
- Does the pain travel anywhere?
- Can you give me a careful description of the pain?
- How long did the pain last and could you do anything to relieve it?
- Is the pain brought on by exertion and relieved by rest?
- Do cold conditions bring it on?
- Do you have any other symptoms, such as breathlessness, faintness, sweating or back pain?
- Is the pain made worse by breathing or coughing, or by movement or pressing on that area?
- Is there any blood in any sputum you bring up?
- Is your pain associated with what you eat and drink? Or with a bitter taste in your mouth?
- Do you get the pain on stooping over and after lying in bed at night?
- Do antacids relieve your pain?
- Have you noticed a rash where you get the pain?
- Have you had a blow to your chest or an injury to your back?

Physical examination

The examination should focus on the following areas:

- general appearance - evidence of atherosclerosis (senile arcus, thickened vessels), pale and sweating (myocardial infarction, dissecting aneurysm or pulmonary embolus), hemiparesis (? aortic dissection)
- pulses - both radial and femoral - check for nature of pulse and absence of femoral pulses
- blood pressure
- temperature
- palpation of chest wall, lower cervical spine and thoracic spine - look for evidence of localised tenderness, pathological fracture, spinal dysfunction, herpes zoster
- palpation of legs - check for evidence of deep venous thrombosis
- examination of chest - check for evidence of pneumothorax
- auscultation of chest
    - reduced breath sounds, hyper-resonant percussion note and vocal fremitus → pneumothorax
    - friction rub →pericarditis or pleurisy
    - basal crackles →cardiac failure
    - apical systole murmur →mitral valve prolapse
    - aortic diastolic murmur →proximal dissection (aortic regurgitation)
- upper abdominal palpation - check for tenderness suggestive of gall bladder disease or peptic ulceration

Note: In the presence of a myocardial infarction, the examination may be normal but the patient, apart from being cold, clammy or shocked, may have muffled heart sounds, a gallop rhythm, a systolic murmur. With a aortic dissection the patient may also appear cold, clammy and shocked, but may show absent femoral pulses, hemiparesis and a diastolic murmur of aortic regurgitation.

Possible findings on examination of a patient with chest pain are presented in Figure 40.2.[1]


signs of pneumothorax
hypotension
• AMI
• DA
hypertension
• pain (early AMI)
pleural friction rub
• pleurisy
• pulmonary infarction
pale, clammy, 'shocked' → AMI
AD
pulmonary embolus
tenderness and limited movement → referred pain
• cervical spine
• thoracic spine
aortic diastolic murmur (AI) → proximal AD
3rd or 4th heart sound
muffled sounds
AMI
friction rub → pericarditis
tenderness of costochondritis
? herpes zoster
tenderness of peptic ulcer
bradyarrhythmia → AMI
absent femoral pulses → AD
hemiparesis → can occur with AD
deep venous thrombosis
→ pulmonary embolus
AMI = acute myocardial infarction
AD = aortic dissection
AI = aortic incompetence

Figure 40.2 Possible examination findings in a patient with chest pain

Investigations[1]

The following investigations to aid diagnosis are available, although the majority are sophisticated and confined to hospitals with high-technology imaging departments. The fundamental tests that are readily available to the GP - ECG, chest X-ray and cardiac enzymes - should help confirm the diagnosis in most instances.

Electrocardiogram (ECG)

This may be diagnostic for ischaemia and myocardial infarction, although it is important to bear in mind that it may be normal with both, including the early minutes to hours of an acute infarction.

It can be helpful to differentiate between myocardial infarction, pulmonary embolism and pericarditis (Figure 40.3). The ECG in pulmonary embolism may show right axis deviation. Pericarditis is characterised by low voltages and saddle-shaped ST segment elevation.


(a)
I
II
III
AVR
AVL
AVF
V1
V2
V3
V4
V5
V6
(b)
I
II
III
AVR
AVL
AVF
V1
V2
V3
V4
V5
(c)
I
II
III
AVR
AVL
AVF
V1
V2
V3
V4
V5
V6

Figure 40.3 Typical ECG pattern for specific causes of acute chest pain (**a**) acute pulmonary embolism; (**b**) acute pericarditis; (**c**) acute inferolateral myocardial infarction

Exercise stress test

This is the key test for defining chest pain as cardiac in origin. Physical stress, such as the motor-driven treadmill or a bicycle ergometer, is used to elicit changes in the ECG to diagnose myocardial ischaemia.

Exercise thallium scan

This radionuclide myocardial perfusion scan using thallium can complement the exercise ECG.

Ambulatory Holter monitor

This monitor is especially useful for silent ischaemia, variant angina and arrhythmias.

Chest X-ray

The routine CXR is taken in full inspiration. Ask for an expiration film if pneumothorax is suspected.

Blood glucose

Tests association with diabetes.

Haemoglobin and blood film

Anaemia is a possible associated factor.

Serum enzymes

Damaged necrosed myocardial tissue releases cellular enzymes, which are markers of this damage:

- troponin I and troponin T (the key marker)
- creatinine kinase (CK) and creatinine kinase-myocardial bound fraction (CK - MB)
- myoglobin

Echocardiography

This can be used in the early stages of myocardial infarction to detect abnormalities in heart wall motion, when ECGs and enzymes are not diagnostic. Stress echocardiography is a new technique which is useful where standard exercise testing has been unhelpful.

Isotope scanning

1. Technetium-99m pyrophosphate studies:
    - myocardium - to diagnose posterolateral myocardial infarction in the presence of bundle branch block
    - pulmonary - to diagnose pulmonary embolism
2. Gated blood pool nuclear scan (radionuclide ventriculography) - this scan tests leftventricular function at rest and exercise in patients with myocardial ischaemia.

Angiography (arteriography)

Angiography should be selective:

1. coronary - to evaluate coronary arteries
2. pulmonary - to diagnose pulmonary thromboembolism

Transoesophageal echocardiography (TOE)

This investigation is for dissecting aneurysm (immediate diagnosis).

Oesophageal studies

- Endoscopy
- Barium swallow
- Oesophageal manometry
- Radionuclide transit studies

Spine - X-ray

- Cervical spine
- Thoracic spine

SITE, RADIATION AND FEATURES OF CHEST PAIN SYNDROMES

Myocardial ischaemia[1]

Coronary artery disease includes the acute coronary syndromes (unstable angina and myocardial infarction) and other variants of angina.

The typical retrosternal distribution of myocardial ischaemia is shown in Figure 40.4. Retrosternal pain or pain situated across the chest anteriorly should be regarded as cardiac until proved otherwise.



Figure 40.4 Pain of myocardial ischaemia: typical site

The wide variation of sites of pain, umbilicus to jaw, including neck, inside of arms, epigastrium and interscapular, should always be kept in mind (Figure 40.5). Pain is referred into the leftarm 20 times more commonly than into the right arm.



Figure 40.5 Pain of myocardial ischaemia: other sites

The quality of the pain is usually typical. The patient often uses the clenched fist sign to illustrate a sense of constriction.

The radiation of pain will assist in differentiating ischaemic pain from that caused by pericarditis. Enquiry about precipitating and relieving factors will enable a differentiation to be made between ischaemic pain and the almost identical pain caused by reference from the spine. Associated symptoms include dyspnoea, nausea and vomiting and sweating.

If a retrosternal pain almost identical with that of myocardial ischaemia is precipitated not by exertion but by bending, lifting, straining or lying down, oesophageal reflux and spasm is a possible diagnosis. This is frequently confused with ischaemic heart disease and can cause radiation into the leftarm.

Stable angina. The pain of angina tends to last a few minutes only (average 3 - 5 minutes) and is relieved by rest and glyceryl trinitrate (nitroglycerine). The pain may be precipitated by an arrhythmia.

The types of acute coronary syndromes are summarised in Table 40.2.[2]

Table 40.2 Types of acute coronary syndromes

	Serum markers		
	Creatine kinase	**Troponin**	**ECG at evaluation**
Unstable angina:			
• low risk	normal	non-detectable	normal
• high risk	normal	detectable	ST depression
Myocardial infarction:			
• non-ST elevation (NSTEMI)	elevated	detectable	ST depression, no Q wave
• ST elevation (STEMI)	elevated	detectable	± Q wave

1. *Myocardial infarction*. Ischaemic pain lasting longer than 15 to 20 minutes is usually infarction. The pain is typically heavy and crushing, and can vary from mild to intense. Occasionally the attack is painless, typically in diabetics. Pallor, sweating and vomiting may accompany the attack.
2. *Unstable angina*. This term includes rest angina, new onset effort angina, post infarct angina and post coronary procedure angina. Severe ischaemic chest pain can last 15 - 20 minutes or more. It is classified as low risk or high risk 'minor myocardial damage'.

> For management purposes it is best to classify the clinical presentation of acute chest pain as STEMI or non-ST elevation acute coronary syndrome (NSTEAC).

Aortic dissection

The pain, which is usually sudden, severe and midline, has a tearing sensation and is usually situated retrosternally and between the scapulae (Figure 40.6). It radiates to the abdomen, flank and legs. An important diagnostic feature is the inequality in the pulses (e.g. carotid, radial and femoral). There may also be occlusion of the coronary or kidney arteries with appropriate symptoms and signs. Hemiplegia, aortic incompetence or cardiac tamponade can occur.



Figure 40.6 Pain of aortic dissection

Pulmonary embolism[3]

This has a dramatic onset following occlusion of the pulmonary artery or a major branch, especially if more than 50% of the cross-sectional area of the pulmonary trunk is occluded.

The diagnosis can present clinical difficulties, especially when dyspnoea is present without pain. Embolism usually presents with retrosternal chest pain (Figure 40.7) and may be associated with syncope and breathlessness. In addition, hypotension, acute right heart failure or cardiac arrest occurs with a massive embolus. The physical examination can be deceptively normal. Pulmonary infarction is generally less dramatic than embolism and it is usually accompanied by pleuritic chest pain and haemoptysis. It complicates embolism in about 10% of patients. The diagnosis is usually confirmed by V/Q scan and/or CT pulmonary angiogram.



Figure 40.7 Pain of pulmonary embolism

Pleurisy[3]

Inflammation of the pleura is due to underlying pneumonia (viral or bacterial), pulmonary infarction, tumour infiltration or connective tissue disease (e.g. SLE).

Features

- Often sudden onset
- Pain usually localised without radiation
- Sharp knife-like pain
- Continuous pain with sharp exacerbations
- Aggravated by inspiration and coughing
- May be associated dyspnoea, cough, haemoptysis

Epidemic pleurodynia (Bornholm's disease)

Unilateral knife-like chest pain (and upper abdominal pain) following an URTI. Caused by a Coxsackie B virus. CXR is normal; diagnosis is by exclusion. Settles within a week with simple analgesics.

Acute pericarditis

Pericarditis causes three distinct types of pain:

1. pleuritic (the commonest), aggravated by cough and deep inspiration, sometimes brought on by swallowing
2. steady, crushing, retrosternal pain that mimics myocardial infarction
3. pain synchronous with the heartbeat and felt over the praecordium and leftshoulder

Occasionally, two and rarely all three types of pain may be present simultaneously (Figure 40.8).



Figure 40.8 Pain of pericarditis

Spontaneous pneumothorax

The acute onset of pleuritic pain and dyspnoea in a patient with a history of asthma or emphysema is the hallmark of a pneumothorax. It is due to a rupture of a subpleural 'bleb' or a small air-containing cyst. It often occurs in young slender males without a history of lung disorders. The pain varies from mild to severe and can be felt anywhere in the chest, sometimes being retrosternal. Typical pain distribution is shown in Figure 40.9. The diagnosis is made on expiration film.



Figure 40.9 Pain of pneumothorax (right side)

If a tension pneumothorax becomes painful and dyspnoea becomes rapidly more intense, urgent decompression of air is essential. A comparison of the serious causes of acute chest pain is summarised in Table 40.3.

Table 40.3 A comparison of the serious causes of acute chest pain

	Myocardial infarction	Angina	Pulmonary embolus	Aortic Dissection	Pericarditis	Pneumothorax
Pain intensity	+ → +++++	+	+ → +++	+++++	+ → +++	+ → +++
Pain quality	Heavy	Heavy	Dull	Tearing	Heavy	Tightness
	Crushing	Aching	Heavy	Searing	Aching	Sharp
	Vice-like	Tightness			± sharp	Stabbing
	Burning	Burning				
Pain sitc	Deep retrosternal	Deep retrosternal	Retrosternal	Anterior chest	Sternal surface	Lateral chest
Pain radiation	Throat/lower jaw	As for infarction	Lateral chest (pleuritic)	Front to back of chest	Left arm (uncommon)	Lateral chest
	Left arm (often)			Down back to abdomen	Right arm (rare)	
	Right arm (uncommon)			Arms	Throat (rare)	
	Back (uncommon)					
History	Family, risk factors	Family, risk factors	Phlebitis	Atherosclerosis	Viral infection	Asthma
			Calf pain	Hypertension		COAD
			Immobility	? Marfan	MI	Old TB
			Surgery			

Associated symptoms	Pallor, nausea, sweating, vomiting, dyspnoea, syncope	Strangling in throat	Dyspnoea, syncope, sweating, vomiting, cyanosis, agitation, ? haemoptysis	Syncope, pallor, cyanosis,	Fever, malaise ± pleuritic pain	Dyspnoea, cough, ? Cyanosis
			Neurological			
			• hemiparesis			
			• paraplegia			
Pulse	variable ? arrhythmias	variable ? arrhythmias	tachycardia	unequal some ?absent	weak if effusion	tachycardia
Cardiac auscultation	± gallop rhythm murmur of MI	S3 during attack	↓pulmonary S2 S3 or S4	± murmur of AI	± pericardial friction rub	
Chest auscultation	basal crackles		± adventitious sounds			hyper-resonant ↓breath sounds ↑percussion
Chest X-ray			#x00B1; localised oligaemia or infarction	widening of mediastinum	↑cardiac silhouette if effusion	diagnostic - expiration film
ECG	Q waves	normal or ST depression	normal or R heart strain	may show myocardial infarction	elevated ST segments	
	ST elevation					
	T inversion (variable)		S1, Q3, T3			
Special definitive diagnostic tests	• serum enzymes: troponin I or T	• stress ECG	• lung scanning	• TOE	• echocardiography (if effusion)	
	• cardiac scanning	• coronary angiography	• pulmonary angiography	• ultrasound		
		• technetium scanning		• aortic angiography		
		• enzymes		• CT scan		

Oesophageal pain

Gastro-oesophageal reflux can cause oesophagitis characterised by a burning epigastric or retrosternal pain that may radiate to the jaw. The pain is aggravated or precipitated by lying flat or bending over, especially after meals, and is more frequent at night. The pain is worse if oesophageal spasm is present. Oesophageal motor disorders, including spasm, may occur in isolation. The pain may radiate uncommonly to the back (Figure 40.10). It may be precipitated by eating, especially hot or cold food and drink, and may be relieved by eating or by glyceryl trinitrate (nitroglycerine) and other nitrates. Features differentiating angina-like oesophageal pain and cardiac pain are presented in Table 40.4 on p. 419. Gastrointestinal causes of chest pain are summarised in Table 40.5 on p. 419.[1]

Table 40.4 Features differentiating angina-like, oesophageal pain and cardiac pain

	Favour oesophageal	Favour cardiac	Non-discriminating
Precipitating factors	Meals, posture	Consistently with exercise	Emotion
Relieving factors	Antacids		Rest, nitrates
Radiation	Epigastrium	Arm	Back
Associated symptoms	Heartburn, regurgitation, dysphagia	Dyspnoea	Sweating

Table 40.5 A comparison of gastrointestinal causes of chest pain

	Acid reflux	Oesophageal spasm	Peptic ulcer	Gall bladder disease
Site	• Epigastric	• Deep retrosternal	• Deep retrosternal	• Right hypochondrium
Radiation	• Retrosternal • Throat	• Back	• To back (DU)	• Below right scapula • Tip right shoulder
Quality	• Burning	• Constricting	• Gnawing	• Deep ache
Precipitation	• Heavy meals GU:30 min DU:2 - 3 hours	• Eating hot/cold food and drinks • Wine/coffee • Lying • Bending	• Eating	• vFatty food
Relief	• Standing • Antacids	• Antispasmodics • Nitroglycerine	• Antacids	
Associated symptoms	• Water brash	• Dysphagia • Dyspepsia	• Dyspepsia	• Flatulence

GU - gastric ulcer; DU - duodenal ulcer



Figure 40.10 Oesophageal pain

Spinal pain

The commonest cause of pain of spinal origin is vertebral dysfunction of the lower cervical or upper dorsal region. The spinal problem may be a disc prolapse (relatively common in the lower cervical spine, but rare in the upper thoracic spine) or dysfunction of the facet joints or costovertebral joints causing referred pain. This referred pain can be present anywhere in the chest wall, including anterior chest, which causes confusion with cardiac pain (Figure 40.11). The pain is dull and aching. It may be aggravated by exertion, certain body movements or deep inspiration. The old trap for unilateral nerve root pain is herpes zoster.



Figure 40.11 Possible pain sites for thoracic spinal dysfunction (left side)

Costochondritis[3]

This causes mild to moderate anterior chest wall pain that may radiate to the chest, back or abdomen. It is usually unilateral, sharp in nature and exaggerated by breathing, physical activity or a specific position. It may be preceded by exercise or a URTI and can persist for several months. It is diagnosed by eliciting tenderness at the chondrosternal junction of the affected ribs and needs to be differentiated from Tietze's syndrome, where there is a tender, fusiform swelling at the chondrosternal junction (refer to p. 970).

Psychogenic pain

Psychogenic chest pain can occur anywhere in the chest, but often it is located in the left submammary region, usually without radiation (Figure 40.12 at p. 419). It tends to be continuous and sharp or stabbing. It may mimic angina but tends to last for hours or days. It is usually aggravated by tiredness or emotional tension and may be associated with shortness of breath, fatigue and palpitations.



Figure 40.12 Typical sites of psychogenic pain

Da Costa's syndrome (effort syndrome) is recurrent attacks of stabbing left-sided submammary pain, usually associated with anxiety ± depression.

CHEST PAIN IN CHILDREN

Chest pain in children is rarely the result of serious pathology but is an important complaint, especially in adolescents. A US study has shown that the mean age for childhood chest pain is 11.9 years.[4] Most cases are of unknown aetiology (probably psychogenic), while common causes include musculoskeletal disorders, cough-induced pain, costochondritis, psychogenic disturbance (includes hyperventilation) and asthma. See Table 40.6.

Table 40.6 Common causes of chest pain in children

Cause	%
Idiopathic	21
Musculoskeletal	16
Cough	10
Costochondritis	9
Psychogenic	9
Asthma	7
Trauma	5
Pneumonia	4
GIT problems	4
Cardiovascular	4

Source: Adapted from Selbst[4]

Chest pain in children less than 12 years old is more likely to have a cardiorespiratory cause, such as cough, asthma, pneumonia or heart disease, while chest pain in adolescents is more likely to be associated with a psychogenic disturbance.

Causes of musculoskeletal pain include strains to pectoral, shoulder or back muscles after excessive exercise, and minor trauma from sports such as football or wrestling.

Breast problems can present as chest pain.

Cardiac causes

Myocardial ischaemia is very rare in children but should be considered in any child with exerciseinduced chest pain, adolescents with long-standing diabetes and children with sickle-cell anaemia.

Precordial catch (Texidor's twinge or stitch in the side)

This common complaint presents as a unilateral low chest pain that lasts a few seconds or minutes, typically with exercise, such as long-distance running. The pain is relieved by straightening up and taking very slow deep breaths followed by shallow breaths.

CHEST PAIN IN THE ELDERLY

Chest pain is a very important symptom in the elderly as the life-threatening cardiovascular conditions - myocardial infarction and angina, dissecting aneurysm and ruptured aorta - are an increasing manifestation with age. In a community survey in Glasgow 20% of men and 12% of women over 65 were found to have ischaemic heart disease.[5] The elderly patient presenting with chest pain is most likely to have angina or myocardial infarction. Other important disorders to consider are herpes zoster, cough fracture of the rib, malignancy, pleurisy, pulmonary embolus and gastro-oesophageal reflux.

ANGINA PECTORIS

Main features

- There is a 2 - 3% incidence between 25 and 64 years.[6]
- The history is the basis of diagnosis.
- Angina is an oppressive discomfort rather than a pain.
- It is mainly retrosternal: radiates to arms, jaw, throat, back.
- It may be associated with shortness of breath.
- It occurs during exercise, emotion, after meals or in cold.
- It is relieved within a few minutes with rest.
- Physical examination is usually not helpful, except during an attack.
- Mitral valve prolapse, oesophageal spasm and dissecting aneurysm are important differential diagnoses.
- The causes of angina are summarised in Table 40.7.

Table 40.7 Causes of angina

Coronary artery atheroma
Valvular lesions (e.g. aortic stenosis)
Rapid arrhythmias
Anaemia
Rarities ▪ trauma ▪ collagen disease

Note: Ensure that the patient is not anaemic or thyrotoxic. Fever and tachycardia also have to be excluded.

Variants of angina[1, 6]

- *Stable angina*. Pain occurs with exertion and is usually predictable.
- *Unstable angina* (also referred to as crescendo angina, pre-infarct angina and acute coronary insufficiency). It is increasing angina (severity and duration) over a short period of time, precipitated by less effort and may come on at rest, especially at night. It may eventually lead to complete infarction, often with relief of symptoms. It is due to unstable plaque.
  - *Nocturnal angina*. Pain occurs during the night. It is related to unstable angina.
  - *Decubitus angina*. The pain occurs when lying flat and is relieved by sitting up.

- *Variant angina or Prinzmetal angina or spasm angina.*[7] The pain occurs at rest and without apparent cause. It is associated with typical transient ECG changes of ST elevation (as compared with the classic changes of ST depression during effort angina). It can lead to infarction and cause arrhythmias. It is caused by coronary artery spasm.

Aids to diagnosis

ECG

This may be normal or show ischaemia or evidence of earlier infarction. During an attack it may be normal or show well-marked depression of the ST segment, symmetrical T-wave inversion (Figure 40.13) or tall upright T waves.



Figure 40.13 Typical ECG pattern for angina pectoris: this tracing is usually observed during an attack. *Note:* there is no specific ECG of angina; the most that can be said is that an ECG is consistent with angina

Exercise ECG

This is positive in about 75% of patients with severe coronary artery disease and should be performed if the diagnosis is in doubt, for prognostic reasons or to aid in the timing of additional investigations (e.g. coronary angiography). A normal stress test does not rule out coronary artery disease.

Exercise thallium-201 scan

This test is helpful in some difficult circumstances such as in the presence of leftbranch bundle block (LBBB), old infarction and Wolf-Parkinson-While (WPW) syndrome (when exercise test is of little use) and with mitral valve prolapse, which gives high false positive tests. It helps determine the presence and extent of reversible myocardial ischaemia since thallium is only taken up by perfused tissue.

Ambulatory Holter monitoring

This may be useful in some patients.

Gated blood pool nuclear scan

This test assesses the ejection fraction, which is a reliable index of ventricular function and thus aids assessment of patients for coronary artery bypass surgery.

Echocardiography

This assesses global and regional wall motion abnormalities and assesses valvular dysfunction and pericardium status.

Coronary angiography

This test accurately outlines the extent and severity of coronary artery disease. It is usually used to determine the precise coronary artery anatomy prior to surgery.

The relationship between the degree of angina and coronary artery disease (CAD) is not clear-cut. Some people with severe angina have normal coronary arteries.

Indications for coronary angiography are presented in Table 40.8.

'Ultrafast CT' may replace angiography.

Table 40.8 Indications for coronary angiography

• Strong positive exercise stress test
• Suspected left main CAD
• Angina resistant to medical treatment
• Suspected but not otherwise proven angina
• Acute coronary syndromes
• Angina after myocardial infarction
• Patients over 30 with aortic and mitral valve disease being considered for valve surgery

Management of stable angina

Preventions

This is especially important for those with a positive family history and an unsatisfactory lifestyle. Modification of risk factors:

- no smoking
- weight reduction
- optimal low-fat diet
- control of hypertension
- control of diabetes
- control of blood lipids

General advice for the stable angina patient

- Reassure patient that angina has a reasonably good prognosis:
  - 30% survive more than 10 years[8]
  - spontaneous remission can occur
- Attend to any risk factors.
- If inactive, take on an activity such as walking for 20 minutes a day.
- Take regular exercise to the threshold of angina.
- If tense and stressed, cultivate a more relaxed attitude to life - consider a stress management/relaxation course.
- Avoid precipitating factors.
- Don't excessively restrict lifestyle.

Medical treatment[9]

The acute attack

- glyceryl trinitrate (nitroglycerin) 600 μg tab or 300 μg (½ tab) sublingually (SL) *or*
- glyceryl trinitrate SL 400 μg metered dose spray: 1 - 2 sprays; repeat after 5 minutes if pain persists (maximum 2 doses) *or*
- isosorbide dinitrate 5 mg sublingually; repeat every 5 minutes if pain persists (maximum 3 tablets) *or*
- nifedipine 5 mg capsule (suck or chew) if intolerant of nitrates

Tips about glyceryl trinitrate tablets

- Warn patient about headache and other side effects.
- Sit down to take the tablet.
- Take ½ (initially) or 1 tablet every 5 minutes.
- Take a maximum of 3 tablets in 15 minutes.
- Tablets must be fresh.
- Discard the bottle opened for 3 months or after 2 days if carried on the person.
- Keep tablets out of light and heat.
- If pain relieved quickly, spit out residual tablet.

- Advise patient to get medical advice if no relief after 3 tablets.

Note: avoid nitrates if patient has taken sildenafil or vardenafil in the previous 24 hours or tadalafil in the previous 5 days.

Mild stable angina

Angina that is predictable, precipitated by more stressful activities and relieved rapidly:

- aspirin 150 mg (o) daily *or* (if intolerant of aspirin) clopidrogel 75 mg (o) daily
- gyceryl trinitrate (SL or spray) prn
- consider a beta-blocker or long-acting nitrate or nicorandil

Moderate stable angina

Regular predictable attacks precipitated by moderate exertion:

- *add* beta-blocker e.g. atenolol 25 - 100 mg (o) once daily *or* metoprolol 25 - 100 mg (o) twice daily
- glyceryl trinitrate (transdermal: ointment or patches) daily (use for 12 - 16 hours only) *or* isosorbide mononitrate 60 mg (o) SR tablet mane

Note: Aim for a daily nitrate-free interval.

Persistent angina

Not prevented by beta-blocker: *add* a dihydropyridine calcium-channel blocker (CCB) (must have beta-blocker cover)

- nifedipine 10 - 20 mg (o) bd or nifedipine CR 30 - 60 mg (o) once daily *or* amlodipine or felodipine 2.5 - 10 mg (o) once daily

If beta-blocker contraindicated (use a nondihydropyridine calcium-channel blocker)

- diltiazem 30 - 90 mg (o) tds or CR 180 - 360 mg (o) daily *or* verapamil (according to directions) *and/or*
- nicorandil 5 mg (o) bd, increasing after a week to 10 - 20 mg bd

Refractory stable angina

Replace CCB with perhexiline.

Unstable angina

Includes onset of angina at rest, abrupt worsening of angina and angina following acute myocardial infarction:

- Should be hospitalised for stabilisation and further evaluation. May need IV nitrate therapy.
- The objectives are to optimise therapy and consider coronary angiography with a view to a corrective procedure.

Rules of practice

- For variant angina (spasm) use nitrates and calcium antagonist (avoid beta-blockers).
- As a rule avoid the combination of verapamil and a beta-blocker.
- Tolerance to nitrate use is a problem, so 24 hour coverage with long-acting preparations is not recommended.
- Consider using the potassium channel opening vasodilator nicorandil 5 mg (o) bd to 10 - 20 mg (o) bd (after 1 week).
- Nitrates can be used prophylactically prior to any exertion that is likely to provoke angina, e.g. glyceryl trinitrate spray or tablet *or* isosorbide dinitrate 5 mg tablet
- Avoid nitrates if the patient has used a 5 phosphodiesterase inhibitor in the past 1 - 5 days.

Non-medical treatment

Percutaneous intervention (PCI) and coronary angioplasty

One current technique is dilating coronary atheromatous obstructions by inflating a balloon against the obstruction - percutaneous transluminal coronary angioplasty (PTCA) (Figure 40.14) and maintaining patency with intracoronary stent devices.

Two complications of the balloon inflation angioplasty are acute coronary occlusion (2 - 4%) and restenosis, which occurs in 30% in the first 6 months after angioplasty.[8]



Figure 40.14 Percutaneous transluminal angioplasty (PTCA) with an inflatable balloon

Intra-coronary stents

PTCA followed by stenting is now the most favoured procedure to maintain patency of the obstructed coronary vessel (Figure 40.15). Modern drug eluting stents, which include drugs such as primolimus, sirolimus or paclitaxel, aim to achieve longer term patency.


After plaque is removed or compressed,
stent positioned and expanded to keep artery open
Stent in place

Coronary artery surgery

The main surgical techniques in current use are coronary artery bypass grafting (CABG) using either a vein (usually the saphenous) (Figure 40.16) or internal mammary arterial implantation (Figure 40.17) or both and endarterectomy.

Symptomatic patients with significant left main coronary obstruction should undergo bypass surgery, and those with two or three vessel obstruction and good ventricular function are often considered for angioplasty or surgery. A significant improvement in the quantity and quality of life can be expected.


coronary artery bypass grafting
obstruction
left coronary artery

Figure 40.16 Coronary artery bypass grafting to relieve coronary obstruction



Figure 40.17 Internal mammary arterial transplantation to relieve obstruction

MYOCARDIAL INFARCTION

Clinical guidelines

- Variable pain; may be mistaken for indigestion
- Similar to angina but more oppressive
- So severe, patient may fear imminent death - *angor animi*
- About 20% have no pain
- 'Silent infarcts' in diabetics, hypertensives and elderly; one report (*European Heart Journal* 14/2/2006, online) reported that 40% of infarcts were silent and undetected in older patients
- 60% of those who die do so before reaching hospital, within 2 hours of the onset of symptoms
- Hospital mortality is 8 - 10% [10]
- Like CVA, seems to peak at 6 - 10 am
- Diagnosis is based on 2 out of 3 criteria: history of prolonged ischaemic pain, typical ECG appearance, and rise and fall of cardiac enzymes

Aetiology

- Thrombosis with occlusion
- Haemorrhage under a plaque
- Rupture of a plaque
- Coronary artery spasm

Physical signs

These may be:

- no abnormal signs
- pale/grey, clammy, dyspnoeic
- restless and apprehensive
- variable BP
  - ↑ with pain
  - ↓ heart pump failure
- variable pulse: watch for bradyarrhythmias
- mild cardiac failure: third or fourth heart sound, basal crackles

Investigations

1. *ECG*. The ECG is valuable with characteristic changes in a full thickness infarction. The typical features (Figure 40.18) are:



Figure 40.18 Typical ECG features of myocardial infarction, illustrating a Q wave, ST elevation and T-wave inversion

   - the Q wave: broad (> 1 mm) and deep > 25% length R wave
     - occurs normally in leads AVR and V1; III (sometimes)
     - abnormal if in other leads
     - occurs also with LBBB, WPW and ventricular tachycardia (VT)
     - usually permanent feature after full thickness AMI
   - T wave and ST segment
     - transient changes (inversion and elevation respectively)

   The typical progression is shown in Figure 40.19.


before
minutes afterwards
hours afterwards
days afterwards
weeks afterwards

Figure 40.19 Typical evolution of ECG changes with myocardial infarction

Note:

- Q waves do not develop in subendocardial infarction.
- The strategies for management of AMI are based on the distinction between Q wave (transmural) or non-Q wave (subendocardial) infarction.
- Q wave infarction has been proved to benefit from thrombolytic therapy but non-Q wave infarction has not.
- A normal ECG, especially early, does not exclude AMI. Q waves may take days to develop.

2. *Cardiac enzymes*. The typical enzyme patterns are presented in Figure 40.20. As a rule large infarcts tend to produce high serum enzyme levels. The elevated enzymes can help time the infarct:



Figure 40.20 Typical cardiac enzyme patterns following myocardial infarction

- Troponin I or T:
  - starts rising at 3 - 6 hours, peaks at 10 hours and persists for several days
  - now the preferred test
  - positive in unstable angina
  - may have to wait until 10 hours before recording a negative result
  - both proteins, I and T, provide same information
  - reference interval < 0.1 μg/L
- creatinine kinase
  - after delay of 6 - 8 hours from the onset of pain it peaks at 20 - 24 hours and usually returns to normal by 48 hours
  - CK - MB: myocardial necrosis is present if > 15% of total CK; unlike CK, it is not affected by intramuscular injections

3. *Technetium pyrophosphate scanning*
   - It is performed from 24 hours to 14 days after onset.
   - It scans for 'hot spots', especially when a posterolateral AMI is suspected and ECG is unhelpful because of pre-existing LBBB.
4. *Echocardiography*. This is used to assist diagnosis when other tests are not diagnostic.

Note: The clinical diagnosis may be the most reliable, as the ECG and enzymes may be negative.

Management of acute coronary syndromes

General principles:[10, 11]

- Aim for immediate attendance if suspected.
- Call a mobile coronary care unit.
- Achieve coronary perfusion and minimise infarct size.
- Prevent and treat cardiac arrest; have a defibrillator available to treat ventricular fibrillation.
- Optimal treatment is in a modern coronary care unit (if possible) with continuous ECG monitoring (first 48 hours), a peripheral IV line and intranasal oxygen.
- Pay careful attention to relief of pain and apprehension.

- Establish a caring empathy with the patient.
- Give aspirin as early as possible (if no contraindications).
- Prescribe a beta-blocker drug and an ACE inhibitor early (if no contraindications).

Note: For a STEMI it is important to re-establish flow ASAP, usually by either thrombolytic therapy or primary angioplasty (preferably with stenting). Rescue angioplasty is usually used when large infarcts have not perfused at 60 - 90 minutes.[9]

First-line management (e.g. outside hospital)

- Perform an ECG and classify ACS into STEMI or NSTEACS, and notify the medical facility who will receive the patient (discuss over the telephone). The ECG is the sole test required to select patients for emergency perfusion
- Oxygen 4 - 6 L/min
- Secure an IV line (withdraw blood for tests especially troponin levels)
- Glyceryl trinitrate (nitroglycerin) 300 μg (½ tab) SL (every 5 min as necessary to max. of 3 doses) (beware of sildenafil (Viagra) use and bradycardia - correct with atropine)
- Aspirin 300 mg
- Morphine 2 - 5 mg IV statim bolus: 1 mg/min until pain relief (up to 15 mg) ±metoclopramide 10 mg IV (as antiemetic) (If feasible it is preferable to give IV morphine 1 mg/min until relief of pain; this titration is easier in hospital.)

Hospital management[9, 11]

- As for first-line management.
- Confirm ECG diagnosis: STEMI or NSTEACS.
- Take blood for cardiac enzymes particularly troponin levels, urea and electrolytes.
- Organise an urgent cardiology consultation for risk stratification and a decision whether to proceed to coronary angiography and coronary reperfusion with PCI (or CABGS) or with thrombolysis.

Management of STEMI [11]

The optimal first-line treatment for the patient with a STEMI is urgent referral to a coronary catheter laboratory within 60 minutes of the onset of pain for assessment after coronary angiography for percutaneous transluminal coronary angioplasty (PTCA). If available and performed by an interventional cardiologist it has the best outcomes (Level 1 evidence).

The principle is to achieve rapid reperfusion with primary angioplasty with a stent (optimal stent status currently under evaluation).

Adjunct therapy will include aspirin/clopidogrel and heparin and possibly a glycoprotein IIb/IIIa platelet inhibitor such as tirofiban or abciximab.

Acceptable time delay guidelines to PCI (from first medical contact to ballon inflation)[11]

Symptom time:	< 1 hour	1 - 3 hours	3 - 12 hours	> 12 hours
Acceptable delay:	60 minutes	90 minutes	120 minutes	Not recommended

Management of NSTEAC

All patients with non-ST-segment elevation ACSs should have their risk stratified to direct management decisions.

Fibrinolytic therapy

If angioplasty is unachievable either through timing or the unavailability of the service (such as in rural locations) thrombolysis is an indication for STEMI and the sooner the better, but preferably within 12 hours of the commencement of chest pain.[11] The decision should be made by an experienced consultant, especially as PCI is not usually possible once fibrinolytic therapy has been given.

Second-generation fibrin-specific agents (reteplase, alteplase or tenecteplase) are the agents of choice. Streptokinase can be used but it is inappropriate for use in Indigenous people and those who have received it on a previous occasion. There are several other contraindications for the use of fibrinolytic agents.

Further management strategies include:

- Full heparinisation for 24 - 36 hours (after rt-PA - not after streptokinase), especially for large anterior transmural infarction with risk of embolisation, supplemented by warfarin.
- Use LMW heparin e.g. enoxparin 1 mg/kg SC bd or fractionated heparin 5000 - 7500 units SC 12 hourly.
- Beta-blocker (if no thrombolytic therapy or contraindications) as soon as possible: atenolol 5 - 10 mg IV or 25 mg (o) statim →

25 - 100 mg (o) daily *or* metoprolol 5 - 10 mg IV or 25 mg (o) statim → 50 mg (o) bd

- Consider glyceryl trinitrate IV infusion if pain recurs.
- Start early introduction of ACE inhibitors (within 24 hours) in those with significant left ventricular (LV) dysfunction (and other indications).
- Statin therapy to lower cholesterol.
- Treat hypokalaemia.
- Consider magnesium sulphate (after thrombolysis).
- Consider frusemide.

Post AMI drug management

Proven:[1, 12, 13]

- beta-blockers - within 12 hours
- ACE inhibitors - within 24 hours
- aspirin 160 - 325 mg ± clopedrigol
- lipid-lowering drugs (e.g. statins)
- warfarin

Possible/promising:

- folic acid, vitamins B6 and B12

Ongoing management

- Education and counselling
- Bed rest 24 - 48 hours
- Check serum potassium and magnesium
- Early mobilisation to full activity over 7 - 12 days
- Light diet
- Sedation
- Beta-blocker (o): atenolol or metoprolol
- Warfarin where indicated (certainly if evidence of thrombus with echocardiography)
- ACE inhibitors for left ventricular failure and to prevent remodelling

On discharge

- Rehabilitation program
- Continued education and counselling
- No smoking
- Reduce weight
- Encourage consumption of omega-3 fatty acids
- Regular exercise, especially walking
- Exercise test (to be considered)
- Continue beta-blockers for 2 years
- Continue ACE inhibitors
- Aspirin 100 - 300 mg daily or clopidogrel 75 mg daily
- Warfarin where indicated (at least 3 months)

Special management issues

Indications for coronary angiography

- Development of angina
- Strongly positive exercise test
- Consider after use of streptokinase

Management of the extensive infarction

- ACE inhibitors (even if no CCF)

- Radionuclide studies (to assess left ventricular function)
- Beta-blockers (proven value in severe infarction) if no contraindications or LV dysfunction
- Anticoagulation

Treating and recognising complications of STEMI

Acute left ventricular failure

- Signs:
  - basal crackles
  - extra (third or fourth) heart sounds
  - X-ray changes
- Treatment (according to severity):
  - oxygen
  - diuretic (e.g. frusemide)
  - morphine IV
  - glyceryl trinitrate: IV, SL, (o) or topical
  - ACE inhibitors

Cardiogenic shock (a major hospital management procedure)

- Dobutamine ± dopamine
- Intra-aortic balloon pump
- Urgent angiography ± angioplasty/surgery

Pericarditis

This occurs in first few days after AMI (usually anterior AMI), with onset of sharp pain.

- Signs:
  - pericardial friction rub
- Treatment:[9]
  - paracetamol 1 g (o), 4 hourly *or* anti-inflammatory medication e.g. aspirin 600 mg (o) 4 hourly *or* indomethacin 25 mg (o) 4 hourly (or by suppository) *or* corticosteroids (keep in reserve)

Note: Avoid anticoagulants.

Post-AMI syndrome (Dressler's syndrome)

This occurs weeks or months later, usually around 6 weeks.

- Features:
  - pericarditis
  - fever
  - pericardial effusion (an autoimmune response)
- Treatment:
  - as for pericarditis

Left ventricular aneurysm

This is a late complication.

- Clinical:
  - cardiac failure
- Features:
  - arrhythmias
  - embolisation
- Signs:
  - double ventricular impulse
  - fourth heart sound
  - visible bulge on X-ray
- Diagnosis:

  - 2D electrocardiography
- Treatment:
  - antiarrhythmic drugs
  - anticoagulants
  - medication for cardiac failure
  - possible aneurysmectomy

Ventricular septal rupture and mitral valve papillary rupture

This presents with severe cardiac failure and a loud pansystolic murmur. Both have a poor prognosis and early surgical intervention may be appropriate.

MANAGEMENT OF OTHER SERIOUS CAUSES OF CHEST PAIN

Aortic dissection

- Early definitive diagnosis is necessary: best achieved by transoesophageal echocardiography.
- 50% of patients are hypertensive; so need pharmacological control of hypertension with IV nitroprusside and beta-blockers.
- Emergency surgery needed for many, especially for type A (ascending aorta involved).

Note: There is an increased risk during pregnancy.

Pulmonary embolus

Investigations to diagnose suspected pulmonary embolus (choose from):[3, 14, 15]

- chest X-ray and ECG
- radionucleide imaging - the ventilation/perfusion (V/Q) study (first-line study)
- CT pulmonary angiography
- digital subtraction angiography (gold standard)
- D-dimer assay
- Doppler sonography of lower limbs
- arterial blood gases

Management

Needs supportive medical care and anti-coagulation:

- Heparin IV: 5000 U as immediate bolus continuous infusion 30 000 U over 24 hours *or* heparin 12 500 U (sc) bd

Note: The dose of heparin should then be adjusted daily to maintain the APTT between 1.5 and 2 times control.

- Continue heparin 5 - 10 days.
- Warfarin (o) after 3 to 4 days; then continue heparin for 3 days after international normalised ratio (INR) at desired level.

Note: Thrombolytic therapy either IV or into the pulmonary artery can be used for major embolism. Surgical embolectomy is rarely necessary but needed if very extensive.

Ruptured oesophagus

Refer for early thoracotomy and repair of the defect, usually a longitudinal repair in the supradiaphragmatic lower oesophagus - with mediastinal and pleural drainage.[16]

Pneumothorax

- Most episodes resolve spontaneously without drainage (at least 20% lung collapse).
- Drainage of the pleural space indicated for a large pneumothorax > 30% pleural area, with persistent dyspnoea.
- For recurrent attacks, excision of cysts or pleurodesis may be necessary.
- Statistics indicate a 30 - 50% recurrence rate of spontaneous pneumothorax (most within 12 months), 35% on the same side, 10 - 15% on the opposite side. Recurrence should not recur after a pleurodesis where the lung surface has been rendered adherent to the chest wall.

Methods

1. *Simple aspiration without underwater drainage.* Under strict sepsis insert a 16-gauge polyethylene IV catheter into the second intercostal space in the mid-clavicular line on the affected side (under local anaesthetic). Then aspirate air into a 20 mL syringe to confirm entry into the pleural space, remove the stilette, connect the catheter via a flexible extension tube to a three-way tap and a 50 mL syringe. Aspirate and expel air via the three-way tap until resistance indicates lung re-expansion. Obtain a follow-up X-ray. Observe for 3 - 4 hours. Repeat aspiration may be necessary but most patients do not require inpatient admission.
2. *Standard intercostal catheter insertion with connection to an underwater seal drainage in hospital.*

Acute tension pneumothorax

For urgent cases insert a 12 - 16 gauge needle into the pleural space through the second intercostal space on the affected side. Replace with a formal intercostal catheter connected to underwater seal drainage.

TREATMENT OF OESOPHAGEAL DISORDERS

Gastro-oesophageal reflux

- Achieve normal weight if overweight.
- Avoid coffee, alcohol and spicy foods.
- Avoid large meals and overeating (keep to small meals).
- Use antacids or alginate compounds (e.g. Gaviscon, Mylanta Plus). If persistent: acid suppression - H2-receptor blockers (e.g. cimetidine, ranitidine) *or* proton-pump inhibitors (e.g. omeprazole)

Oesophageal spasm

long-acting nitrates (e.g. isosorbide dinitrate 10 mg tds) *or* calcium-channel blockers (e.g. nifedipine 10 mg tds)

Note: Attend to lifestyle and dietary factors, as for reflux.

MUSCULOSKELETAL CAUSES OF CHEST WALL PAIN

There are many musculoskeletal causes, most of which can be eliminated by the history and physical examination. Some of the causes listed in Table 40.9 are very uncommon and often part of a general disorder, such as ankylosing spondylitis. Muscular tears or strains of the chest wall are quite common. A differential diagnosis is a fractured rib including a cough fracture.

Musculoskeletal chest pain is typically aggravated or provoked by movements such as stretching, deep inspiration, sneezing and coughing. The pain tends to be sharp and stabbing in quality but can have a constant aching quality.

Costochondritis is a common cause of anterior pain, which is generally well localised to the costochondral junction and may also be a component of an inflammatory disorder, such as one of the spondyloarthropathies.

Management is generally conservative with analgesics, gentle massage with analgesic creams and NSAIDs if there is an inflammatory component. Other measures that can help for very painful chest wall problems are localised injections of local anaesthetic with or without corticosteroids (with care not to penetrate the parietal pleura) and a modified support (especially for rib injuries) in the form of a special elasticised rib belt (called a universal rib belt) that gives excellent support and symptom relief while permitting adequate lung expansion.

Table 40.9 Musculoskeletal causes/origins of chest wall pain (front and back)

Injury to thoracic spine →dysfunction

Vertebral fracture

• trauma

• pathological

- osteoporosis

- metastatic disease

- multiple myeloma

Intercostal muscle strains/tears

Rib disorders

- fractures
- slipping rib

Costochondritis

Tietze's syndrome

Fibromyalgia

Posterior chest (thoracic back) pain

Disorders of the musculoskeletal system represent the most common cause of thoracic (dorsal) back pain, especially dysfunction of the joints of the thoracic spine. Refer to Chapter 38 for more detail. Probably the commonest cause is costovertebral dysfunction caused by overstress of rib articulations with vertebrae (the costovertebral joints). This fact is clearly demonstrated with the midline thoracic back pain following cardiac surgery when these joints are compressed during sternotomy and splaying of the chest walls.

The back pain may be associated with simultaneous referred anterior chest pain or abdominal pain (Figure 38.4 at p. 392).

Acute thoracic back pain

Although posterior pain is invariably caused by vertebral dysfunction, there are several other important causes, including serious bone disease (leading to compression fractures) and life-threatening visceral and vascular causes. Refer to Tables 38.3 and 38.4 at p. 389.

Note:

- Intervertebral disc protrusions are rare in the thoracic spine.
- Rarely, a penetrating peptic ulcer can present with mid to lower thoracic back pain.

Management

Management has to be according to the cause. The most appropriate treatment for thoracic spine dysfunction (in the absence of organic disease such as osteoporosis and inflammation, and if the patient is not taking anticoagulants) is physical therapy - mobilisation or manipulation, as described in Chapter 38.

WHEN TO REFER

- Obvious or suspected myocardial infarction, especially with extensive infarction
- Transfer to major centre with complications of AMI
  - rupture of septum or papillary muscle
  - aneurysm
  - refractory arrhythmias
  - cardiogenic shock
- Patients with persistent post-infarction angina
- Angina
  - patient with angina not responding to drug treatment
  - patient with unstable angina
  - angina lasting for longer than 15 minutes (unresponsive to sublingual nitrate) needs urgent hospital admission
- Suspected or proven pulmonary embolus or dissecting aneurysm or other serious lifethreatening problem (after initial first-line measures, e.g. decompression of tension pneumothorax)
- Suspected oesophageal or other gastrointestinal disorder (e.g. duodenal ulcer), for endoscopy or appropriate gastroenterological evaluation

Practice tips

- All sudden acute chest pain is cardiac (and potentially fatal) until proven otherwise.
- A careful history is the basis of the diagnosis.

- Mitral valve prolapse is often an undiagnosed cause of chest pain: keep it in mind, especially if pain is recurrent and intermittent (proved by echocardiography).
- Calcium antagonists can cause peripheral oedema so be careful not to attribute this to heart failure.
- The pain of oesophageal spasm can be very severe and mimic myocardial infarction.
- Oesophageal spasm responds to glyceryl trinitrate: do not confuse with angina.
- Intervertebral disc protrusions are a very rare cause of severe sudden thoracic pain (T2 - 9).
- Infective endocarditis can cause pleuritic posterior chest pain.
- Family doctors need to monitor carefully patients who are taking anticoagulants. The INR ratio, which needs to be kept between 2 and 3, should be tested at least monthly.
- The sudden onset of dyspnoea without chest pain can occur frequently with (painless) myocardial infarction and pulmonary embolism.
- If a patient recovering from an AMI suddenly develops shortness of breath, consider ventricular septal rupture, mitral valve papillary rupture (with mitral regurgitation), pulmonary embolus and other serious complications.
- Treat (indefinitely) all post MI patients with ACE inhibitors, and all post MI and acute ischaemic syndrome patients with betablockers. [12, 13]
- Use antiplatelet agents indefinitely - 100 - 300 mg aspirin daily or, if contraindicated, clopidogrel 75 mg daily or warfarin. Some studies favour combining aspirin and clopidogrel.

REFERENCES

1. Juergens C. How to treat: Chest pain. Australian Doctor, 2005; 2 September: 27 - 34.
2. Management of unstable angina guidelines 2000. Med J Aust, 2000; 173, (Suppl.).
3. Worsnop C, Pierce R. Pleuritic chest pain. Medicine Today, March 2005; 6: 3; 53 - 60.
4. Selbst S, Ruddy R, Clark B et al. Paediatric chest pain: a prospective study. Paediatrics, 1988; 82: 319 - 23. **431**
5. Kennedy RD, Andrews GR, Mitchell JRA. Ischaemic heart disease in the elderly. Br Heart J, 1977; 39: 1121 - 7.
6. Caspari P. Angina pectoris. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 35 - 8.
7. Kincaid-Smith P, Larkins R, Whelan G. *Problems in Clinical Medicine*. Sydney: MacLennan & Petty, 1990: 10 - 11.
8. Kumar P, Clark M. *Clinical Medicine*. London: Saunders, 2003: 769 - 74.
9. Smith A (Chair). *Therapeutic Guidelines*: *Cardiovascular* (Version 4). Melbourne. Therapeutic Guidelines Ltd, 2003; 96 - 119.
10. Thompson P. Myocardial infarction. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 327 - 30.
11. Aroney C, Aylward P (Co-chairs). Guidelines for the management of acute coronary syndromes 2006. MJA, 2006; 184(8): Supplement.
12. Barton S ed. *Clinical Evidence*. London. BMJ Publishing Group, 2001: 8 - 23.
13. American Heart Association Guidelines for myocardial ischaemia. Circulation, 2001; 104: 1577 - 9.
14. Rashford S. Acute pleuritic chest pain. Aust Fam Physician, 2001; 30(9): 841 - 5.
15. Lau L. *Imaging Guidelines* (4th edn). Melbourne: RANZC Radiologists, 2001: 70.
16. Hunt P, Marshall V. *Clinical Problems in General Surgery*. Sydney: Butterworths, 1991: 87 - 8.

CHAPTER 41. CONSTIPATION

I have finally kum to the konklusion, that a good reliable set of bowels iz wurth more tu a man, than enny quantity of brains.

HENRY SHAW (1818 - 85), 'JOSH BILLINGS '

CONSTIPATION

Constipation is the difficult passage of small hard stools. It has also been defined as infrequent bowel actions or a feeling of unsatisfied emptying of the bowel. However, the emphasis is on the consistency of the stool rather than on the frequency of defecation; for example, a person passing a hard stool with difficulty once or twice a day is regarded as constipated, but the person who passes a soft stool comfortably every two or three days is not constipated. Various causes of chronic constipation are summarised in Figure 41.1.


Psychogenic
• depression
• anorexia nervosa
Neurological disease
• CVA
• autonomic neuropathy
Drugs
Endocrine/metabolic
• hypothyroidism
• hypercalcaemia
Idiopathic constipation
• simple constipation
• slow transit constipation
• irritable bowel syndrome
Organic disease of large bowel
• carcinoma colon
• diverticular disorder
• inflammatory bowel disease
• impacted faeces
• anorectal disorders
carcinoma
fissure
haemorrhoids
• Others
adynamic bowel
Hirschsprung's senility
spinal cord injury
connective tissue disorders
(e.g. scleroderma)

Figure 41.1 Causes of chronic constipation

KEY FACTS AND CHECKPOINTS

- A UK study1 showed that the diagnostic range is 12 per 1000 patients per year.
- The survey showed 10% of adults and 6% of children reported constipation in the preceding 2 weeks.

- Up to 20% of British adults take regular laxatives.[2]
- Constipation from infancy may be due to Hirschsprung's disorder.
- Diet is the single most important factor in preventing constipation.
- Beware of the recent onset of constipation in the middle aged and the elderly.
- Bleeding suggests carcinoma, haemorrhoids, diverticular disorder and inflammatory bowel disease.
- Unusually shaped stools (small pellets or ribbonlike) suggest the irritable bowel syndrome.
- Always examine the abdomen and rectum.
- Plain abdominal X-rays are generally not useful in the diagnosis of chronic constipation.
- The flexible sigmoidoscope is far superior to the rigid sigmoidoscope in investigation of the lower bowel.

A DIAGNOSTIC APPROACH

Using the safe diagnostic strategy model (Table 41.1 on p. 434), the five self-posed questions can be answered as follows.

Table 41.1 Chronic constipation: diagnostic strategy model

Q. Probability diagnosis
A. Simple constipation: low-fibre diet and bad habit
Q. Serious disorders not to be missed
A. Intrinsic neoplasia: colon, rectum or anus, especially carcinoma of colon Extrinsic malignancy (e.g. lymphoma, ovary) Hirschsprung's (children)
Q. Pitfalls (often missed)
A. Impacted faeces Local anal lesions Drug/purgative abuse Hypokalaemia Depressive illness Acquired megacolon Diverticular disease *Rarities* • Lead poisoning • Hypercalcaemia • Hyperparathyroidism • Dolichocolon (large colon) • Chagas' disease • Systemic sclerosis
Q. Seven masquerades checklist
A. Depression Diabetes rarely Drugs Anaemia - Thyroid disorder hypo Spinal dysfunction severe only UTI -
Q. Is the patient trying to tell me something?
A. May be functional (e.g. depression, anorexia nervosa).

Probability diagnosis

The commonest is ‘idiopathic’ constipation where there is no structural or systemic disease. This is also referred to as ‘functional’ constipation.

Probably the most frequent single factor causing constipation in Western society is deficiency in dietary fibre, including fruit, green leafy vegetables and wholemeal products. The amount of fibre in our diet is directly related to stool weight and to colonic transit time. The average colonic transit time in the large bowel for Westerners is 60 hours; for a rural African on a very high fibre diet it is 30 hours. Constipation is also a common problem in pregnancy.

Serious disorders not to be missed

Neoplasia

It is obvious that colonic or anorectal neoplasms must not be missed in a patient, especially middle-aged or elderly, presenting with constipation or other change in bowel habit. Most cases present with either complete or incomplete bowel obstruction.

Extrinsic malignancy, such as lymphoma or carcinoma of the ovary, compressing or invading the rectum, also has to be considered. Carcinoma of the large bowel is most prevalent in our society and appropriate screening examinations, including rectal examination, sigmoidoscopy and colonoscopy, must be considered where appropriate.

Megacolon

In children it is important to detect the presence of megacolon, for example megacolon secondary to Hirschsprung's disorder. Symptoms dating from birth suggest Hirschsprung's disorder, which occasionally may present for the first time in adult life.[3]

Neurological disorders

Constipation, often with faecal impaction, is a common accompaniment to paraplegia, multiple sclerosis, cerebral palsy and autonomic neuropathy.

Alarm symptoms

- Recent constipation in > 40 year old
- Rectal bleeding
- Family history of cancer

Pitfalls

The pitfalls can be summarised as follows:

- impacted faeces
- depressive illness
- purgative abuse
- local anal lesions
- drugs

Although patients with impacted faeces usually pre-sent with spurious diarrhoea, it is a form of idiopathic constipation and is very commonly encountered in general practice, especially in bedridden elderly people.

Anal pain or stenosis, such as fissure-in-ano, thrombosed haemorrhoids, perianal haematoma, or ischiorectal abscess, leads to constipation when the patient is hesitant to defecate.

General pitfalls and tips

- Ensure the patient is truly constipated, and not having unreal expectations of regularity.
- Ensure that the anthraquinone group of laxatives, including ‘Ford pills’, is never used longterm because they cause melanosis coli and associated megacolon.
- Be very wary of alternating constipation and diarrhoea (e.g. carcinoma of colon).
- In a busy practice be careful not to let ‘familiarity breed contempt’ (e.g. onset of hyperparathyroidism, carcinoma).
- Avoid relying solely on the rectal examination to exclude carcinoma.

Seven masquerades checklist

Three of the primary masquerades (Table 41.1) are important causes of constipation, namely drugs, depression and hypothyroidism.

Many drugs (Table 41.2) may be associated with constipation, especially codeine and its derivatives, antidepressants, aluminium and calcium antacids. A careful drug history is thus mandatory. Fortunately the constipation is usually relieved once the drug is withdrawn. Constipation can be a significant symptom in all types of depressive illness and may be aggravated by treatment with antidepressants.

Table 41.2 Drugs associated with constipation

Analgesics (inhibitors of prostaglandin synthesis)
Antacids (containing calcium carbonate or aluminium hydroxide)
Anticholinergic agents
Antidiarrhoeal agents
Antiepileptics
Antihistamines (H_1-receptor blockers)*
Antiparkinsonian drugs*
Barbiturates
Barium sulphate
Benzodiazepines
Calcium supplements
Cholestyramine
Clonidine
Cough mixtures
Cytotoxic drugs
Diuretics that cause hypokalaemia
Ganglionic blocking agents
Heavy metal (especially lead)
Iron supplements
Laxatives (chronic use)
Monoamine oxidase inhibitors
Muscle relaxants
Opioid analgesics (e.g. codeine)
Phenothiazines*
SSRIs
Tricyclic antidepressants*
Calcium-channel blockers (verapamil)

*Denotes anticholinergic effect.

The metabolic causes of constipation include hypothyroidism, hypercalcaemia and porphyria. We occasionally encounter the patient with hypercalcaemia, for example, hyperparathyroidism, but thyroid dysfunction is relatively common in general practice.

Diabetes rarely can be associated with constipation when an autonomic neuropathy can lead to alternating bouts of constipation and diarrhoea.

Psychogenic considerations

Constipation may be a manifestation of an underlying functional problem and psychiatric disorder, such as depression, anorexia nervosa, schizophrenia or drug abuse. Drug abuse must always be considered, keeping in mind that narcotics and laxatives present with rebound constipation. More commonly, it may reflect the inactive lifestyle of the patient and provide a good opportunity for appropriate counselling.

THE CLINICAL APPROACH

History

It is important to ask patients to define exactly what they mean by constipation. Some people believe that just as the earth rotates on its axis once a day, so should their bowels open daily to ensure good health. As always, a careful history is appropriate, including stool consistency, frequency, ease of evacuation, pain on defecation and the presence of blood or mucus. A dietary history is very relevant in the context of constipation.

Key questions

- How often do you go to the toilet?
- What are your bowel motions like?
- Are they bulky, hard, like rabbit pellets or soft?
- Is there pain on opening your bowels?
- Have you noticed any blood?
- Have you noticed any lumps?
- Do you have any soiling on your underwear?
- How do you feel in yourself?
- What medications are you taking?

Diary

Ask the patient to keep a 10-day diary recording frequency and nature of stools, and whether any difficulty was experienced when passing stool.

Physical examination

The important aspects are abdominal palpation and rectal examination. Palpation may reveal the craggy mass of a neoplasm, faecal retention (especially in the thin patient) or a tender spastic colon. The perianal region should be examined for localised disease. The patient should be asked to bear down to demonstrate perianal descent, haemorrhoids or mucosal prolapse.[3] Perianal sensation and the anal reflex should be tested. Digital rectal examination is mandatory, and may reveal a rectal tumour and faecal impaction, as well as testing for rectal size and tone. If there is a history from infancy, a normal or narrow rectum suggests congenital megacolon (Hirschsprung's disorder) but, if dilated, acquired megacolon.

General signs that may be significant in the diagnosis of constipation are summarised in Figure 41.2.


Irregular liver
• metastases
(carcinoma of colon)
Distension
Mass in RIF
• carcinoma
• Crohn's disease
Mass
• faecal impaction
• carcinoma

Figure 41.2 Possible significant abdominal signs in the patient with constipation

The rectal examination

The most important first step is to *do* the examination.

Method

- Explain to the patient what will happen.
- After inspection with the patient in the left lateral position and with knees drawn up, a lubricated gloved index finger is placed over the anus.
- Ask the patient to concentrate on slow deep breathing.
- With gentle backwards pressure the finger is then inserted slowly into the anal canal and then into the rectum (it helps patient comfort if they push down or squeeze to accommodate the finger).
- Rotate the finger anteriorly to feel the prostate in males and the cervix in females.
- The finger will reach to about 7 - 8 cm with gentle thrusting into the perineum.
- Gently withdraw the finger and examine the whole circumference of the rectum by sweeping the finger from posterior on both sides.

Points to note:

- any pain - fissure, proctitis, excoriation from diarrhoea (a rectal examination will not be possible in the presence of a fissure)
- induration from a chronic fissure or fistula in the anal canal

- the sphincter tone
- the nature of the faeces (? impaction)
- the rectal wall
  - carcinoma is usually indurated, elevated and ulcerated
  - a villous adenoma has a soft velvety feel
- posteriorly - the sacrum and coccyx
- laterally - the side walls of the pelvis
- anteriorly
  - cervix and pouch of Douglas in the female
  - prostate and rectovesical pouch in the male (Figure 41.3)



Figure 41.3 Rectal examination in the male: the normal prostate is bilobed with a central sulcus

The prostate

- It feels larger if the patient has a full bladder.
- The normal prostate is a firm smooth rubbery bilobed structure (with a central sulcus) about 3 cm in diameter.
- A craggy hard mass suggests carcinoma.
- An enlarged smooth mass suggests benign hypertrophy.
- A tender, nodular or boggy mass suggests prostatitis.

A common pitfall

In the female the cervix or a vaginal tampon can be mistaken for a mobile extrarectal tumor (Figure 41.4).



Figure 41.4 Rectal examination in the female: the cervix or vaginal tampon may be mistaken for a rectal mass

Endoscopy

Sigmoidoscopy - in particular, flexible sigmoidoscopy with examination of the rectosigmoid - is important in excluding local disease; search for abnormalities such as blood, mucus or neoplasia. The insufflation of air sometimes reproduces the pain of the irritable bowel syndrome.

It is worth noting that 60% of polyps and cancers will occur in the first 60 cm of the bowel4 and diverticular disorder should be evident with the flexible sigmoidoscope.

The presence of melanosis coli is an important sign - it may give a pointer to the duration of the constipation and the consequent chronic intake (perhaps denied) of anthraquinone laxatives.

Investigations

These can be summarised as follows.

Haematological

- Haemoglobin
- ESR

Stools for occult blood Biochemistry (where suspected)

- Thyroid function tests
- Serum calcium
- Serum potassium
- Carcinoembrogenic antigen (a tumour marker)

Radiological

- Double contrast barium enema (especially for primary colonic disease, e.g. megacolon)
- Bowel transit studies, using radio-opaque shapes taken orally and checking progress by abdominal X-ray or stool collection

Physiological tests

- Anal manometry - test anal tone
- Rectal sensation and compliance, using an inflatable rectal balloon
- Dynamic proctography, to determine disorders of defecation
- Rectal biopsy, to determine aganglionia

CONSTIPATION IN CHILDREN

Constipation is quite common in children but no cause has been discovered in 90 - 95%. The most common factor is diet. Constipation often begins after weaning or with the introduction of cow's milk. It is rare with breastfeeding. Low fibre intake and a family history of constipation may be associated factors.[5] Most children develop normal bowel control by 4 years of age (excluding any physical abnormality). It is normal to have a bowel movement every 2 - 3 days, providing it is of normal consistency and is not painful. It is important to differentiate between encopresis and constipation.

- *Encopresis* is the inappropriate passage of normal stool, which usually indicates a psychological disorder or stress.
- *Constipation* is difficulty or delay in passing the stool with incomplete emptying of the rectum. This can present as *soiling*, due to faecal retention with overflow of liquid faeces.

Other important conditions

Hirschsprung's disorder

- Consider if delay in passing first meconium stool and subsequent constipation.

Anal fissure in infants

- Consider if stool hard and associated with pain or bleeding.
- The mainstay of treatment is dietary manipulation.

Principles of treatment of functional constipation

- Encourage relaxed child - parent interaction with toilet training, such as appropriate encouragement, ‘after breakfast habit’ training.
- Introduce psychotherapy or behaviour modification program, especially where ‘fear of the toilet’ exists.
- Establish an empty bowel: remove any severely impacted faeces with microenemas (e.g. Microlax), and even disimpaction under anaesthesia if necessary. A good guide if faecal ‘rocks’ are visible on x-ray.
- Advice for parents of children over 18 months:
  - Drink ample non - milk fluids each day - several glasses of water, unsweetened fruit juice or milk.
  - Use prune juice, which contains sorbitol.
  - Get regular exercise - walking, running, outside games or sport.
  - Provide high-fibre foods - high-fibre cereals, wholegrain bread, brown rice, wholemeal pasta, fresh fruit with skins left on where possible, dried fruits such as sultanas, apricots or prunes, fresh vegetables.
- Use a pharmaceutical preparation as a last resort to achieve regularity:
  First line:
  - bulk-producing agent (e.g. psyllium granules, 1 teaspoonful, once or twice daily)

  Second line: osmotic laxative (e.g. lactulose):
  - 1 - 5 years: 5 mL bd
  - 6 - 12 years: 10 mL bd
  - > 12 years: 15 mL bd

  or Stool softeners:
  - paraffin oil e.g. Parachoc
  - lactulose
  - poloxamer drops e.g. Coloxyl

CONSTIPATION IN THE ELDERLY

Constipation is a common problem in the elderly with a tendency for idiopathic constipation to increase with age. In addition the chances of organic disease increase with age, especially colorectal carcinoma, so this problem requires attention in the older patient. Faecal impaction is a special problem in the aged confined largely to bed. Constipation is often associated with Parkinson's disease. In the elderly an osmotic laxative such as sorbitol or lactulose may be required for long-standing refractory constipation but stimulant and other non-osmotic laxatives should be avoided.

Practice tip on treatment

A suitable method of doing a rectal examination on a home visit (in the absence of gloves in the doctor's bag) is to apply moist soap around the finger and caked under the nail (in case of breakage), then plastic wrap and finally petroleum jelly (e.g. Vaseline).

Before resorting to a good old-fashioned '3H' enema (hot water, high and a hell of a lot), use a sorbitol compound (e.g. Microlax 5 mL enema). It can be carried in the doctor's bag, is very easy to insert and is most effective.

Manual disimpaction

If manual disimpaction should be necessary, the unpleasant procedure can be rendered virtually odourless if the products are 'milked' or scooped directly into a container of water. A large plastic cover helps restrict the permeation of the smell.

Discomfort and embarrassment are reduced by this method and by adequate premedication (e.g. IV midazolam and IV fentanyl) if large faecaliths are present.

IDIOPATHIC CONSTIPATION

It is best to classify idiopathic constipation into three subgroups:[3]

1. simple constipation
2. slow transit constipation
3. normal transit constipation (irritable bowel syndrome)

Of these, the commonest is simple constipation, which is essentially related to a faulty diet and bad habit. Avery-Jones,[6] who defined the disorder, describes it as being due to one or more of the following causes:

- faulty diet
- neglect of the call to stool
- unfavourable living and working conditions
- lack of exercise
- travel

Dyschezia, or lazy bowel, is the term used to describe a rectum that has become unresponsive to faecal content, and this usually follows repeated ignoring of calls to defecate.

Slow transit constipation occurs primarily in women with an apparently normal colon, despite a high-fibre intake and lack of the other causes described by Avery-Jones. Many are young, with a history dating from early childhood or, more commonly, adolescence. Constipation may follow childbirth, uncomplicated abdominal surgery or a period of severe dieting. However, in the majority no precipitating cause is evident.

Management

Most patients have simple constipation and require reassurance and education once an organic cause has been excluded.

Basic advice to the patient

- Adequate exercise, especially walking, is important.
- Develop good habit: answer the call to defecate as soon as possible. Develop the 'after breakfast habit'. Allow time for a good relaxed breakfast and then sit on the toilet.
- Avoid laxatives and codeine compounds (tablets or mixture).
- Take plenty of fluids, especially water and fruit juices e.g. prune juice.
- Eat an optimal bulk diet. Eat foods that provide bulk and roughage, such as vegetables and salads, cereals (especially wheat fibre), fresh and dried fruits, and wholemeal bread. Enough fibre should be taken to convert stools that sink to stools that float.[3]

Examples of food with good bulk properties are presented in Table 41.3.[7] Fruit has good fibre, especially in the skin, and some have natural laxatives (e.g. prunes, figs, rhubarb, apricots).

Table 41.3 Foods with bulk-forming properties (from least to most)

• Potato
• Banana
• Cauliflower
• Peas
• Cabbage
• Lettuce
• Apple
• Carrot
• Bran

Pharmaceutical preparations

Some patients may not tolerate unprocessed bran but tolerate pharmaceutical preparations better (Table 41.4). An appropriate choice would be one of the hydrophilic bulk-forming agents such as ispaghula or psyllium. Avoid stimulant laxatives except for short sharp treatments.

Table 41.4 Therapeutic agents to treat constipation (with examples)

Hydrophilic bulk-forming agents
psyllium mucilloid (Agiofibe, Metamucil)
sterculia (Granocol, Normacol)
ispaghula (Agiolax, Fybogel)
methylcellulose (Cellulone)
wheat bran
crude fibre (Fibyrax Extra)
Stimulant (irritant) laxatives
phenolphthalein (Laxettes, Figsen, Ford pills)
senna (Senokot)
cascara
castor oil
bisacodyl (Durolax)
Osmotic laxatives
magnesium sulphate (Epsom salts)
magnesium hydroxide (milk of magnesia)
lactulose (Duphalac)
mannitol
sorbitol (Sorbilax)
Stool-softening agents
liquid paraffin (Agarol)
dioctyl sodium sulphosuccinate (Coloxyl)
glycerin suppositories
sorbital/sodium compounds (Microlax)

- *First-line therapy* [8]
  Use a general bulking agent e.g. psyllium or ispaghula granules 1 - 2 teaspoonsful (o) once or twice daily
- *Second-line therapy*
  Use an osmotic laxative or a fibre-based stimulant preparation e.g. sorbitol 70% liquid, 10 mL (o) bd increasing as required *or* lactulose syrup, 15 - 30 mL (o) daily until response, then 10 - 20 mL daily *or* dried fruits with senna leaf (Nu-Lax) 10 g nocte
- *Third-line therapy*

(Recheck cause.) magnesium sulphate, 1 - 2 teaspoons in water once or twice daily (if normal kidney function) *or* a stimulating agent, e.g. bisacodyl 10 - 20 mg (o) nocte *or* glycerin suppository (retain for 15 - 20 minutes) *or* sodium citrate or phosphate enema

COLORECTAL CANCER

Clinical features

- Commonest GIT malignancy
- Second most common cause of death from cancer in Western society
- Generally men over 50 years (90% of all cases)
- Mortality rate about 60%
- Good prognosis if diagnosed early
- Two-thirds in descending colon and rectum

Refer to section on genetics of colorectal cancer (p. 179).

Predisposing factors

- Ulcerative colitis (long-standing)
- Familial
    - familial adenomatous polyposis (FAP)
    - hereditary non-polyposis colorectal cancer
- Colonic adenomata
- Decreased dietary fibre

Lifetime risk

- General population 1 in 25 females, 1 in 20 males
- Family history (first-degree relative) 1 in 8
- Familial polyposis coli 100% by age 50

Consider referral to a familial cancer clinic for assessment

Symptoms

- Blood in the stools
- Mucus discharge
- Recent change in bowel habits (constipation more common than diarrhoea)
- Alternating constipation with spurious diarrhoea
- Bowel leakage when flatus passed
- Unsatisfactory defecation (the mass is interpreted as faeces)
- Abdominal pain (colicky) or discomfort (if obstructing)
- Rectal discomfort
- Symptoms of anaemia
- Rectal examination - this is appropriate because many carcinomas are found in the lowest 12 cm and most can be reached by the examining finger

Obstruction (distension with ↑ pain)

If obstructing, there is a risk of rupture of the caecum.

Surgery is needed to circumvent the closed loop obstruction.

Spread

- Lymphatics → epigastric and para-aortic nodes
- Direct → peritoneum
- Blood → portal circulation

Various forms of presentation of large bowel cancers are shown in Figure 41.5.


B. Left colon carcinoma
• pain (varying degrees)
• obstruction (partial)
• altered bowel habit
A. Right colon carcinoma
• anaemia
• dyspepsia
• palpable mass
pain
umbilicus
pain
pain
C. Sigmoid colon carcinoma
• obstruction
• ± acute abdomen
• altered bowel habit
• rectal bleeding
D. Distal bowel and rectal carcinoma
• mass
• rectal bleeding
• tenesmus

Figure 41.5 Various forms of presentation of large bowel carcinoma

Investigations

- FOBT: immunochemical tests, e.g. inform, do not require dietary or medication restriction
- Sigmoidoscopy, especially flexible sigmoidoscopy
- Barium enema (accurate as a double contrast study) but may miss tumours
- Colonoscopy: essential if suspicion on clinical grounds remains and barium enema normal (more useful if rectal bleeding)
- Ultrasonograph and CT scanning not useful in primary diagnosis; valuable in detecting spread especially hepatic metastases

If FOBT is positive - investigate by colonoscopsy or by flexible sigmoidoscopy.

Screening[9]

An FOBT every 2 years is now recommended for all people 50 - 80 years (see guidelines on p. 77). Colonoscopy is recommended for those at high risk yearly or 2 yearly commencing at 25 years, and every 12 months from 10 - 15 years of age if a strong family history of FAP. In addition, flexible sigmoidoscopy and rectal biopsy for those with ulcerative colitis. Refer to a bowel cancer specialist to plan appropriate surveillance.

Management

Early surgical excision is the treatment, with the method depending on the site and extent of the carcinoma. Duke's classification gives a guide to prognosis (Table 41.5).

Table 41.5 Modified Duke's classification of colorectal cancer

Stage	Pathologic description	Approx. 5 year survival %
A	Cancer limited to mucosa and submucosa	90+
B	Cancer extends into muscularis or serosa	75 - 85
C	Cancer involves regional lymph nodes	30 - 40
D	Distant metastases (e.g. liver)	< 5

MEGACOLON

Hirschsprung's disorder (aganglionosis)

Features

- Congenital
- Constipation from infancy
- Abdominal distension from infancy
- Possible anorexia and vomiting
- Male to female ratio ° 8:1
- Rectal examination - narrow or normal rectum
- Abdominal X-ray/barium enema - distended colon full of faeces to narrow rectum
- Diagnosis, confirmed by full thickness biopsy, shows absence of ganglion cells
- Absent rectoanal reflex on anal manometry

Treatment

Resect narrow segment after preliminary colostomy.

ACQUIRED MEGACOLON

Features

- In older children and adults
- Mainly due to bad habit
- Can be caused by:
    - chronic laxative abuse
    - milder form of Hirschsprung's disorder
    - Chagas' disease (Latin America)[2]
    - hypothyroidism (cretinism)
    - systemic sclerosis
- Marked abdominal distension
- Rectal examination - dilate loaded rectum, lax sphincter
- Abdominal X-ray/barium enema - distended colon full of faeces but no narrowed segment

Treatment

Re-education of bowel habit is required.

WHEN TO REFER[4]

Patients with constipation or change in bowel habit of recent onset without obvious cause need further investigation.

- Patients with chronic symptoms that do not respond to simple measures should be referred.

Practice tips

- The objectives of treatment should be to exclude organic disease and then reassure and re-educate the patient about normal bowel function.
- Discourage long-term use of laxatives, suppositories and microenemas.
- The laxatives to discourage should include anthraquinone derivatives, bisacodyl, phenolphthalein, magnesium salts, castor oil and mineral oils.
- First-line treatment of functional constipation (unresponsive to simple measures) is a bulking agent. An osmotic laxative is good second-line therapy.
- Bleeding with constipation indicates associated organic illness - exclude carcinoma of the bowel. Bright red blood usually means haemorrhoids.
- Beware of hypokalaemia causing constipation in the elderly patient on diuretic treatment.
- If carcinoma can be felt on rectal examination an abdominal perineal procedure with colostomy usually follows; if not, an

anterior resection is generally the rule.

An appropriate management plan for the patient presenting with constipation is that given by Barnes.[3]

REFERENCES

1. Cormack JJC. *Practice: A Handbook of Primary Medical Care*. London: Kluwe-Harrap, 1980; 3(47): 1 - 2.
2. Sandler G, Fry J. *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986: 209.
3. Barnes PRH. Assessment and management of constipation. What's new in gastroenterology? Gastroenterol Soc Aust, 1989; 34: 1 - 4.
4. Bolin T. Constipation. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 127 - 9.
5. Barton S ed. *Clinical Evidence*. London: BMJ Publishing Group, 2001: 231 - 5.
6. Avery-Jones F, Godding FW. *Management of Constipation*. Oxford: Blackwell Scientific Publications, 1972: 16.
7. Sali A. Preventive initiatives in medicine and surgery. Aust Fam Physician, 1985; 14: 1314.
8. Moulds RFW (Chair). *Therapeutic Guidelines: Gastrointestinal* (2nd edn.) Melbourne: Therapeutic Guidelines Ltd, 1999: 83 - 91.
9. *Guidelines for Preventative Activities in General Practice* (6th edn). Sydney: RACGP, 2005: 54.

CHAPTER 42. COUGH

I bounded into bed. The bound made me cough - I spat - it tasted strange - it was bright red blood
- I don't want to find this is real consumption - I shan't have my work written. That's what matters.
How unbearable it would be to die - nothing real finished.

KATHERINE MANSFIELD (1888 - 1923), DIARY ENTRY 1918

COUGH

Cough is one of the five most common symptoms presenting in family practice. There is a wide range of causes (Table 42.1) with the great majority being minor and self-limiting, although the possibility of serious causes such as bronchogenic carcinoma should always be kept in mind.

Table 42.1 Significant causes of cough

Non-productive (dry cough)
Upper respiratory tract infection
Lower respiratory tract infection
- viral
- mycoplasma
Inhaled irritants
- smoke
- dust
- fumes
Inhaled foreign body
Bronchial neoplasm
Pleurisy
Interstitial lung disorders
- fibrosing alveolitis
- extrinsic allergic alveolitis
- pneumoconiosis
- sarcoidosis
Tuberculosis
Left ventricular failure (esp. nocturnal cough)
Whooping cough (pertussis)
GORD and hiatus hernia
Postnasal drip
Productive cough
Chronic bronchitis
Bronchiectasis
Pneumonia
Asthma
Foreign body (later response)
Bronchial carcinoma (dry or loose)
Lung abcess
Tuberculosis (when cavitating)

It is a feature of smokers, who often have a morning cough with little sputum. Coughing can also be initiated by pleural irritation. It is a reflex that provides an essential protective service. It serves to remove substances that may have been accidentally inhaled and removes excess secretions or exudates that may accumulate in the airway.

KEY FACTS AND CHECKPOINTS

- Cough is the commonest manifestation of lower respiratory tract infection.
- Cough is the cardinal feature of chronic bronchitis.
- Cough is a feature of asthma with sputum production, especially at night.
- Cough can have a psychogenic basis.
- Cough may persist for many weeks following an acute URTI as a result of persisting bronchial inflammation and increased airway responsiveness.[1]
- Postnasal drip is the commonest cause of a persistent or chronic cough, especially causing nocturnal cough due to secretions (mainly from chronic sinusitis) tracking down the larynx and trachea during sleep.
- The commonest causes of haemoptysis are URTI (24%), acute or chronic bronchitis (17%), bronchiectasis (13%), TB (10%). Unknown causes totalled 22% and carcinoma 4% (figures from a UK study).[2]

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 42.2.

Table 42.2 Cough: diagnostic strategy model

Q. Probability diagnosis
A. Upper respiratory infection Postnasal drip Smoking Acute bronchitis Chronic bronchitis
Q. Serious disorders not to be missed
A. Cardiovascular •left ventricular failure Neoplasia •carcinoma of lung Severe infections •uberculosis •pneumonia •influenza •lung abscess •HIV infection Asthma Cystic fibrosis Foreign body Pneumothorax
Q. Pitfalls (often missed)
A. Atypical pneumonias Gastro-oesophageal reflux (nocturnal) Smoking (children/adolescents) Bronchiectasis Whooping cough (pertussis) Interstitial lung disorders

Sarcoidosis
Q. Seven masquerades checklist
A. Depression -
Diabetes -
Drugs
Anaemia
Thyroid disorder -
Spinal dysfunction -
UTI -
Q. Is the patient trying to tell me something?
A. Anxiety and habit

Probability diagnosis

The most common cause of cough is an acute respiratory infection, whether a URTI or acute bronchitis.[3] Persistent coughing with a URTI is usually due to the development of sinusitis with a postnasal drip.

Chronic bronchitis is also a common cause of cough.

Serious disorders not to be missed

Bronchial carcinoma must not be overlooked. A worsening cough is the commonest presenting problem. A bovine cough is suggestive of carcinoma: the explosive nature of a normal cough is lost when laryngeal paralysis is present, usually resulting from bronchial carcinoma infiltrating the left recurrent laryngeal nerve.

Chronic cough may be the first presentation of *Pneumocystis carinii* pneumonia in an HIV-infected patient. Careful but tactful questioning of the patient in relation to IV drug use, sexual practice and previous blood transfusions is important. Important causes of a chronic cough are summarised in Table 42.3.

Table 42.3 Some causes of chronic cough[2, 4, 5]

Normal chest X-ray (includes most causes)
• Chronic postnasal drip
• Asthma
• Asthma + postnasal drip
• Postinfective bronchial hyper-responsiveness
• Gastro-oesophageal reflux
- symptomatic
- asymptomatic
• Chronic bronchitis
• Chronic heart failure
• Drugs (e.g. ACE inhibitors, inhaled steroids)
• Snoring and obstructive sleep apnoea
• Irritants: occupational and household
• Somekind cough
• Whooping cough (pertussis)
• Habit
• Functional
• Idiopathic
Abnormal chest X-ray
• Bronchiectasis

- Carcinoma: bronchogenic, larynx
- Cardiac failure
- COPD
- Cystic fibrosis
- Inhaled foreign body
- Interstitial lung disorders (e.g. sarcoidosis)
- Tuberculosis

The possibility of a foreign body should always be kept in mind, especially in children, and severe infections such as TB and pulmonary abscess must not be misdiagnosed.

It is also important not to overlook asthma in which a nocturnal cough, without wheezing, is a feature in children.

Pitfalls

Causes that tend to be overlooked, especially in the presence of a normal X-ray, are gastro-oesophageal reflux, postnasal drip and asthma. Gastro-oesophageal reflux is more common as a cause of reflex coughing, especially at night, than appreciated. Whooping cough, especially immunisation-modified, can be difficult to diagnose, particularly if the characteristic whoop is absent.

General pitfalls

- Attributing cough due to bronchial carcinoma in a smoker to ‘smoker's cough’
- Overlooking TB, especially in the elderly, by equating symptoms to old age, bronchitis or even smoking
- Overlooking the fact that bronchial carcinoma can develop in a patient with other pulmonary conditions, such as chronic bronchitis
- Being slow to order a chest X-ray
- Failing to recognise that pertussis presents in adults

Seven masquerades checklist

The applicable masquerade is drugs, many of which can produce a wide variety of disorders of the respiratory tract that cause a cough. Pulmonary infiltration with fibrosis may result from some cytotoxic drugs, especially bleomycin. Over 20 different drugs are known to produce an SLE-like syndrome, sometimes complicated by pulmonary infiltrates and fibrosis. Cough can be a feature of some of the ACE inhibitors and betablockers, inhaled steroids and salazopyrin.

Psychogenic considerations

A cough can occur for psychosocial reasons. Coughing is under cerebral control and a slight cough before commencing a speech is normal and presumably assists in clearing mucus from around the vocal cords.[6] This can readily become a nervous habit or mannerism. A typical ‘psychogenic’ cough is barking in quality - the ‘Cape Barren goose’ cough. It does not occur during sleep.

THE CLINICAL APPROACH

History

The nature of the cough may provide important diagnostic clues but it is the associated symptoms, such as the nature of the sputum, breathlessness, wheezing and constitutional symptoms, that provide the most helpful diagnostic value. A history of smoking habits, past and present, is essential and an occupational and hobby history requires investigation. Significant occupations (past or present) include mining (pneumoconiosis), aircraft manufacturing (asbestosis and mesothelioma), farming (‘farmer's lung’ - allergic pneumonitis from mouldy hay) and bird handling (‘bird fancier's lung’ - allergic alveolitis or psittacosis from pigeons or budgerigars). A past history of recurrent lung infections from childhood is suggestive of cystic fibrosis and bronchiectasis; a history of hay fever and eczema suggests asthma, while a family history involves asthma, cystic fibrosis, emphysema (α_1-antitrypsin deficiency) and tuberculosis.

Key questions[7]

- How would you describe the cough?
- How long has the cough been present?
- Do you cough up sputum?
- Describe the sputum, especially its colour.

- Is there any blood in the sputum?
- How much sputum do you produce - a teaspoonful, an eggcupful or more?
- Is there a burning sensation in your throat or chest when you cough?
- Have you noticed any other symptoms?
- What about chest pain or fever, shivers or sweats?
- Do you have a wheeze?
- Have you had previous attacks of wheezing or hay fever?
- Is there a history of asthma in your family?
- Have you lost weight?
- Has anyone in the family had TB or a persistent cough?
- How much do you smoke?
- Are you exposed to any smoke or fumes?
- What kind of work do you do?
- Where have you worked in the past?
- Is there a chance you have been exposed to asbestos?
- Do you keep birds at home?
- Do you have any birds nesting outside your bedroom?
- Is there a possibility of a foreign body such as a peanut ‘having gone down the wrong way’?
- Have you had an operation recently or been confined to bed?
- Have you noticed any swelling of your legs?
- Have you been exposed to birds such as pigeons?

Physical examination

Physical examination includes a general examination with a search for features such as enlarged cervical or axillary glands, which may indicate bronchial carcinoma, as would Horner's syndrome (constricted pupil, ptosis). A careful examination of the lungs and cardiovascular system is also appropriate. Fine crackles on auscultation indicates functionary oedema of heart failure, interstitial pulmonary fibrosis and early lobar pneumonia, while coarse crackles indicate resolving pneumonia, bronchiesctasis and TB. Careful inspection of the sputum forms an important part of the physical examination of the lungs. This should include its colour and consistency, presence of particulate matter and a 24 hour sputum watch.

Investigations

This applies particularly to patients with haemoptysis. Investigations include:

- haemoglobin, blood film and white cell count
- sputum cytology and culture
- ESR (elevated with bacterial infection, bronchiectasis, TB, lung abscess and bronchial carcinoma)
- respiratory function tests
- radiology
    - plain chest X-ray (shows many problems)
    - tomography
        - helps more precise localisation of lesion
        - may show cavitation
    - bronchography
        - shows bronchiectasis (a very unpleasant procedure)
    - CT scanning (more sensitive than plain X-ray)
    - ventilation/perfusion isotope scan
        - for pulmonary infarction
- skin tests
- lung biopsy
- bronchoscopy (best at time of haemoptysis)

A schema for the investigation of chronic cough is presented in Figure 42.1.[8]


Cough
Cause not apparent
Cause apparent
Treat
Chest X-ray
Unresolved
Resolved
Localised opacity
Generalised opacity
Normal chest X-ray
Examination of sputum by:
• microscopy
• culture (including examination for acid-fast bacilli)
• cytology
• PCR tests
Fibre-optic bronchoscopy
CT scan or thorax
Exclude pulmonary oedema
Spirometry
• total lung capacity
• diffusion capacity of lung for carbon monoxide
Fibre-optic bronchoscopy with:
• bronchoalveolar lavage
• transbronchial biopsy
Sinus X-rays
Spirometry
± bronchial provocation test
Barium swallow or gastroscopy
Oesophageal pH monitoring and/or manometry
Pneumonia
Lung neoplasms
• benign
• malignant
• primary
• metastatic
Inhaled foreign body
Bronchiectasis
No clear diagnosis
Referral to thoracic surgeon for:
• mediastinoscopy
• thoracotomy
Undiagnosed asthma
Gastro-oesophageal reflux
Chronic bronchitis
Postnasal drip
No clear diagnosis
CT scan of thorax
Fibre-optic bronchoscopy
Left ventricular failure
Interstitial lung diseases
Extensive pneumonic processes
Opportunistic infection
No clear diagnosis
Open lung biopsy

Figure 42.1 A recommended schema for investigation of chronic cough. Adapted With Permission From T.J. Williams And G. Bowes, *Mod Med Aust*, June 1992

However, all that is needed initially is a plain chest X-ray.

Diagnostic characteristics

There are important characteristics of cough that may point to the causation. Table 42.1 compares typical causes of dry and productive cough.

Character of the cough

- Brassy →
  - tracheitis and bronchitis (major bronchi)
  - extrinsic pressure on trachea
- Barking → laryngeal disorders (e.g. laryngitis)
- Croupy (with stridor) → laryngeal disorders (e.g. laryngitis, croup)
- Bovine (no power) → vocal cord paralysis (left recurrent laryngeal nerve)
- Weak cough → indicates bronchial carcinoma
- Paroxysmal with whoops → whooping cough
- Painful →
  - tracheitis

  - left ventricular failure

Timing

- Nocturnal cough →
  - asthma
  - left ventricular failure
  - postnasal drip
  - chronic bronchitis
  - whooping cough
- Waking cough →
  - bronchiectasis
  - chronic bronchitis
  - GORD

Associations

- Changing posture →
  - bronchiectasis
  - lung abscess
- Meals →
  - hiatus hernia (possible)
  - oesophageal diverticulum
  - tracheo-oesophageal fistula
- Wheezing →
  - asthma
- Breathlessness →
  - asthma
  - left ventricular failure
  - COPD

Sputum

A healthy non-smoking individual produces approximately 100 - 150 mL of mucus a day. This normal bronchial secretion is swept up the airways towards the trachea by the mucociliary clearance mechanism and is usually swallowed. The removal from the trachea is assisted also by occasional coughing although this is carried out almost subconsciously.[6]

Excess mucus is expectorated as sputum. The commonest cause of excess mucus production is cigarette smoking. Mucoid sputum is clear and white.

Character of sputum

- Clear white (mucoid) → normal or uninfected bronchitis
- Yellow or green (purulent) → due to cellular material (neutrophils or eosinophil granulocytes)
  - ± infection (not necessarily bacterial infection)
  - asthma due to eosinophils
  - bronchiectasis (copious quantities)
- Rusty → lobar pneumonia (*S. pneumoniae*): due to blood
- Thick and sticky → asthma
- Profuse, watery → alveolar cell carcinoma
- Thin, clear mucoid → viral infection
- Redcurrant jelly → bronchial carcinoma
- Profuse and offensive →
  - bronchiectasis
  - lung abscess
- Thick plugs (cast-like) →
  - allergic bronchopulmonary Aspergillus
  - bronchial carcinoma
- Pink frothy sputum → pulmonary oedema

Haemoptysis

Bloodstained sputum (haemoptysis), which varies from small flecks of blood to massive bleeding, requires thorough investigation. Often the diagnosis can be made by chest X-ray. Causes are presented in Table 42.4. Haem-optysis must be distinguished from blood-stained saliva caused by nasopharyngeal bleeding or sinusitis and also from haema-temesis.[6] Acute bronchitis preduces strcalcy haemoptysis.

Table 42.4 Causes of haemoptysis (bloodstained sputum)

Acute infection
• URTI (commonest cause[1])
• acute bronchitis (commonest cause[1])
Chronic bronchitis
Bronchiectasis
Lobar pneumonia (rusty sputum)
Tuberculosis
Neoplastic
• bronchogenic carcinoma
• metastatic carcinoma
Pulmonary infarction/embolism
Foreign body
Cardiac
• left ventricular failure
• mitral stenosis
Anticoagulant therapy
Unknown
Rarer causes
Idiopathic pulmonary haemosiderosis
Goodpasture's syndrome
Blood disorders including anticoagulants
Trauma
Iatrogenic (e.g. endotracheal tubes)

Note: Haemoptysis must be distinguished from bloodstained saliva caused by nasopharyngeal bleeding or sinusitis.[6] Copious haemoptysis is due to bronchiectasis or TB.

Productive cough

- Chronic bronchitis
  - mucoid or purulent
  - rarely exceeds 250 mL per day[6]
- Bronchiectasis
  - purulent sputum
  - up to 500 mL/day
- Asthma
  - mucoid or purulent
  - tenacious sputum
- Lung abscess
  - purulent and foul-smelling
- Foreign body
  - can follow impaction

COUGH IN CHILDREN[3]

Cough in children is a very common symptom, but troublesome persistent cough is a great cause of anxiety among parents and probably the commonest symptom for which the family doctor is consulted. Age-related causes of chronic cough (present at least 4 weeks) are presented in Table 42.5. Most children with chronic cough do NOT have asthma. A chest X-ray is advisable for a persistent cough.

Table 42.5 Chronic cough in children: age-related causes to consider

Early months of life
• milk inhalation/reflux
• asthma
Toddler/preschool child
• asthma
• bronchitis
• whooping cough
• cystic fibrosis
• croup
• foreign body inhalation
• tuberculosis
• bronchiectasis
Early school years
• asthma
• bronchitis
• *Mycoplasma* pneumonia
Adolescence
• asthma
• psychogenic
• smoking

Source: After Selecki and Helman[9]

Common causes are:

- asthma
- recurrent viral bronchitis
- acute URTIs
- allergic rhinitis
- croup

Disorders not to be missed are:

- asthma
- cystic fibrosis
- inhaled foreign body
- tracheo-oesophageal fistula
- pneumonia

Several clinicians describe the catarrhal child syndrome as the commonest cause of cough.[3] This refers to children who develop a postnasal drip following acute respiratory infection and allergic rhinitis. Recurrent caugh in children can usually be explainet by recurrent vanal respiratory infections which commonly acute when first exposed to other children. This airways ftend to be overacture. Three is a slight predisposition to asthma.

If asthma is suspected a therapeutic trial of salbutamol 200 µg 4 hourly via a spacer may be worthwhile.

Psychogenic causes

Habit cough can occur in children, especially those with a history of school phobia. The cough does not occur during sleep and remains unchanged with exertion or infection.

Croup (laryngotracheobronchitis)

Clinical features:

- characteristic barking cough with stridor
- sounds like a dog barking or a seal
- children 9 months to 3 years
- usually 11 pm to 2 am
- asucultation contirms inspiratory striodr
- occurs in small local epidemics

Management - refer to Chapter 86.

COUGH IN THE ELDERLY

Important causes of cough to consider in the elderly include chronic bronchitis, carcinoma of the lung, bronchiectasis and left ventricular failure, in addition to the acute upper and lower respiratory infections to which they are prone. It is important to be surveillant for bronchial carcinoma in an older person presenting with cough, bearing in mind that the incidence rises with age. One study found the causes of chronic cough in the elderly to be postnasal drip syndrome 48%, GORD 20% and asthma 17%.[10]

COMMON RESPIRATORY INFECTIONS

Respiratory infections, especially those of the upper respiratory tract, are usually regarded as trivial, but they account for an estimated one-fifth of all time lost from work and three-fifths of time lost from school, and are thus of great importance to the community.[11] The majority of respiratory infections are viral in origin and antibiotics are therefore not indicated.

URTIs are those involving the nasal airways to the larynx, while lower respiratory tract infections (LRTIs) affect the trachea downwards.

Combined URTI and LRTI include influenza, measles, whooping cough and laryngotracheo-bronchitis.

THE COMMON COLD (ACUTE CORYZA)

This highly infectious URTI, which is often mistakenly referred to as ‘the flu’, produces a mild systemic upset and prominent nasal symptoms (Figure 42.2).


Main symptoms
headache
malaise
fever (possible)
tender eyes
runny nose
sneezing
sore throat
cough
myalgia
(possible)
Main complications
sinusitis
otitis media
bronchopneumonia

Figure 42.2 The main symptoms and complications of the common cold

Typical clinical features

- Malaise and tiredness
- Sore, runny nose
- Sneezing
- Sore throat
- Slight fever

Other possible symptoms:

- headache
- hoarseness
- cough

The watery nasal discharge becomes thick and purulent in about 24 hours and persists for up to a week. Secondary bacterial infection is uncommon.

Management

Advice to the patient includes:

- rest - adequate sleep and rest

- analgesics - paracetamol (acetaminophen) or aspirin (max. 8 tablets a day in adults)
- steam inhalations (as per Figure 52.4) for a blocked nose
- cough mixture for a dry cough
- gargling aspirin in water or lemon juice for a sore throat
- vitamin C powder or tablets (e.g. 2 g daily) may aid recovery, however clinical trials are inconclusive
- clinical trials of zinc lozenges and echinacea give contradictory results to date[11]

INFLUENZA

Influenza causes a relatively debilitating illness and should not be confused with the common cold. The differences are presented in Table 42.6. The incubation period is usually 1 - 3 days and the illness commences abruptly with a fever, headache, shivering and generalised muscle aching (Figure 42.3).

Table 42.6 Comparison of common cold and influenza

	Common cold	Influenza
Incubation period	12 hours to 5 days	1 - 3 days
Fever	±	+ +
Cough	(later)	+
Sore throat	+ +	±
Rhinitis sneezing rhinorrhoea	+	-
Muscle aches	-	+
Toxaemia	-	±
Causes	Rhinoviruses Parainfluenza Influenza B, C Coronavirus RSV	Influenza A Influenza B


Main symptoms
prostration
fever/chills
cough
muscle aches and pains
Possible symptoms
headache
rhinorrhoea (late)
sore throat

Figure 42.3 The main features of influenza

Clinical criteria[12]

During an influenza epidemic:

- Fever >38 ° C plus at least one respiratory symption and one systemic symptom
- cough (dry)
- sore throat
- coryza
- prostration or weakness
- myalgia
- headache
- rigors or chills

Complications

- Secondary bacterial infection
- Pneumonia due to *Staphylococcus aureus* (mortality up to 20%)[1]
- Encephalomyelitis (rare)
- Depression (a common sequela)

Management

Advice to the patient includes:

- rest in bed until the fever subsides
- analgesics: aspirin is effective or codeine and aspirin (or paracetamol), especially if a dry cough
- fluids: maintain high fluid intake

Antiviral agents[12]

- Neuraminidase inhibitors (cover influenza A and B):
  - zanamivir (Relenza) 75 mg by inhalation bd
  - oseltamivir (Tamiflu) 10 mg (o) bd

  Both should be commenced within 36 hours of onset and given for 5 days
- Amantadine 100 mg (o) bd, until 48 hours after recovery. Must be given within 48 hours of onset and covers influenza A only.

Note: All the above agents have been shown in RCTs to be beneficial, with a reduction in symptoms by at least 24 hours compared with placebo.[12, 13]

Prophylaxis

Influenza vaccination offers some protection for up to 70% of the population for about 12 months.[1] (See Chapter 9.)

BRONCHITIS

Acute bronchitis

This is acute inflammation of the tracheobronchial tree that usually follows an upper respiratory infection. Although generally mild and self-limiting, it may be serious in debilitated patients.

Features

Features of acute infectious bronchitis are:

- cough and sputum (main symptoms)
- wheeze and dyspnoea
- usually viral infection
- can complicate chronic bronchitis - often due to *Haemophilus influenzae* and *Streptococcus pneumoniae*
- scattered wheeze on auscultation
- fever or haemoptysis (uncommon)

Outcome

- It improves spontaneously in 4 - 8 days in healthy patients.

Treatment[12, 14, 15]

- Symptomatic treatment
- Inhaled bronchodilators for airflow limitation
- Antibiotics usually not needed in previously healthy adult or child
- Use antibiotics only if evidence of acute bacterial infection with fever, increased spcitum volume and spictum purcilence:
  - amoxycylline 500 mg (o) 8 hourly for 5 days *or*
  - doxycycline 200 mg (o) statim then 100 mg daily for 5 days

Chronic bronchitis

This is a chronic productive cough for at least 3 successive months in 2 successive years:

- wheeze, progressive dyspnoea
- recurrent exacerbations with acute bronchitis
- occurs mainly in smokers

Refer to COPD (pp. xxx).

PNEUMONIA

This is inflammation of lung tissue. It usually presents as an acute illness with cough, fever and purulent sputum plus physical signs and X-ray changes if consolidation.

However, the initial presentation of pneumonia can be misleading, especially when the patient presents with constitutional symptoms (fever, malaise and headache) rather than respiratory symptoms. A cough, although usually present, can be relatively insignificant in the total clinical picture. This diagnostic problem applies particularly to atypical pneumonia but can occur with bacterial pneumonia, especially lobar pneumonia.

Community-acquired pneumonia (CAP)[12, 14]

This pneumonia occurs in people who are not or have not been in hospital recently, and who are not institutionalised or immuno-compromised. The choice of antibiotic is initially empirical. CAP is usually caused by a single organism, especially *Streptococcus pneumoniae*, which is now becoming resistant to antibiotics.[15] Treatment is usually for 5 - 10 days for most bacterial causes, 2 weeks for *Mycoplasma* or *Chlamydia* infection and 2 - 3 weeks for *Legionella*.

Typical pneumonia

The commonest community-acquired infection is with *Streptococcus pneumoniae* (majority), now becoming resistant to antibiotics, or *Haemophilus influenzae*.[14]

Clinical features

- Often history of viral respiratory infection
- Rapidly ill with high temperature, dry cough, pleuritic pain
- 1 - 2 days later may be rusty-coloured sputum
- Rapid and shallow breathing follows
- X-ray and examination: consolidation

The atypical pneumonias

(Refer to Chapter 30.)

Clinical features

- Fever, malaise
- Headache
- Minimal respiratory symptoms, non-productive cough
- Signs of consolidation absent
- Chest X-ray (diffuse infiltration) incompatible with chest signs

Causes

Mycoplasma pneumoniae - the commonest:

- adolescents and young adults
- treat with roxithromycin 300 mg (o) daily or
- doxycycline 100 mg bd for 10 - 14 days

Legionella pneumophila (legionnaire's disease):

- related to cooling systems in large buildings
- incubation 2 - 10 days

Diagnostic criteria include:

- prodromal influenza-like illness
- a dry cough, confusion or diarrhoea
- very high fever (may be relative bradycardia)
- lymphopenia with moderate leucocytosis
- hyponatraemia

Patients can become very prostrate with complications - treat with azithromycin IV (first line) or erythromycin (IV or 0) plus (if very serve) ciprofloxacin or rifampicin.

Chlamydia pneumoniae:

- similar to Mycoplasma

Chlamydia psittaci (psittacosis):

- treat with roxithromycin or erythromycin or doxycycline

Coxiella burnetti (Q fever):

- treat with doxycycline

Antibiotic treatment according to severity[11, 13, 14]

Mild pneumonia

This does not require hospitalisation.

- amoxycillin/clavulanate 875/125 mg (o) 12 hourly for 7 days
    - especially if *S. pneumoniae* isolated or suspected

  plus (especially if atypical pneumonia suspected)
- doxycycline 200 mg (o) loading dose, then 100 mg bd or roxithromycin 300 mg (o) daily for 7 days

Moderately severe pneumonia

This requires hospitalisation - see box.

Pneumonia: guidelines for hospitalisation[15]

Age over 65 years

Coexisting illness

High temperature: > 387°C

Clinical features of severe pneumonia

Involvement of more than one lobe

Inability to tolerate oral therapy

- benzylpenicillin 1.2 g IV 4 - 6 hourly for 5 - 10 days *or*
- procaine penicillin 1.5 g IM daily (drugs of choice for S. *pneumoniae*) *or*
- cephalothin 1 g IV 6 hourly for 5 - 10 days (in penicillin-allergic patient)
- If not so severe and oral medication tolerated can use amoxycillin/clavulanate or cefaclor or doxycycline
- If atypical pneumonia use doxycycline, erythromycin or roxithromycin

Severe pneumonia

The criteria for severity are presented in the box.

Guidelines for severe pneumonia[15]

Altered mental state

Rapidly deteriorating course

Respiratory rate > 30 per minute

BP < 90/60 mmHg

Hypoxia $P_a O_2 < 60$ mm Hg

Leucocytes < 4 10^9/L or > 20° 10^9/L

- erythromycin 500 mg IV slowly 6 hourly (covers *Mycoplasma, Chlamydia and Legionella*) plus
- cefotaxime 1 g IV 8 hourly *or* ceftriaxone 1 g IV daily

Pneumonia in children

Features

- Tachypnoea, expiratory grunt
- Possible focal chest signs
- Diagnosis often only made by chest X-ray

Pathogens

- Viruses are the most common cause in infants.
- *Mycoplasma* are common in children over 5 years.
- *S. pneumoniae* is a cause in all age groups.
- Pathogens are difficult to isolate - may need blood culture.

Treatment

Almost all those under 48 months should be admitted to hospital.

- Minimal handling
- Careful observations including pulse oximetry
- Attend to hydration
- Antibiotics indicated in all cases. Refer to guidelines.[12]
  - Mild to moderate (general guidelines only):[16] < 24 months - penicillin IV or IM initially > 24 months - penicillin or roxithromycin
  - Severe:[15] flucloxacillin IV + cefotaxime IV ± roxithromycin

CHRONIC PERSISTENT COUGH

A cough associated with a viral respiratory infection should last no more than 2 weeks. If it does it is termed persistent. A cough lasting 2 months or more is defined as a chronic cough. A cough that lasts longer than 3 to 4 weeks requires scrutiny. Table 42.3 includes some causes of chronic cough.

A chronic cough can be divided into productive and non-productive. If productive, the presence of pus is significant, as purulent sputum usually means bacterial infection in the bronchi and/or sinuses.[4] The main organisms are *Haemophilus influenzae* (the most common), *S. pneumoniae* and *Moraxella*. Such infections are most susceptible to amoxycillin or amoxycillin/clavulanate or parenteral cephalosporins.

Non-productive cough

Some of the many causes of a non-productive cough are included in Table 42.1 and more than one may be operative in a patient simultaneously; for example, an allergic snorer with oesophageal reflux taking an ACE inhibitor for hypertension may have a viral respiratory infection.[4] It has been shown that a non-productive or irritating cough is usually caused by persistent stimulation of irritant receptors in the trachea and major bronchi, and may result in the production of small amounts of mucoid sputum.

Investigations to be considered in intractable chronic cough include a chest X-ray, spirometry, CT scan of the thorax (searching in particular for a tumour) and ambulatory oesophageal pH monitoring.

If bronchial hyper-responsiveness is proven a trial of inhaled or oral corticosteroids can be used in these patients.[5]

Gastro-oesophageal reflux

This common condition is the most likely cause of a persistent, non-productive cough in an apparently well patient with a history of reflux. Recent studies utilising 24 hour ambulatory oesophageal pH monitoring have demonstrated that in patients with persistent unexplained cough the predominant cause is asymptomatic gastro-oesophageal reflux.[4, 5] In the absence of evidence of aspiration the cough is considered to be due to stimulation of a distal oesophageal-tracheobronchial reflex. Other studies have established a relationship between bronchial asthma and reflux or swallowing disorders whereby microaspiration can initiate an inflammatory response in the airways.

Indications for 24 hour ambulatory oesophageal pH monitoring in chronic cough include:

- unexplained chronic cough after clinical assessment
- symptomatic gastro-oesophageal reflux

- chronic cough with known aetiology unresponsive to therapy

If reflux is proven or suspected, treat with routine conservative antireflux measures and at least a 4 week trial of histamine H_2-receptor antagonists. If cough persists, refer for further evaluation.[5]

If all investigations and treatment trials (including antireflux and corticosteroids) prove unrewarding, a trial of ipratropium bromide both nebulised (500 µg, qid) and to a lesser degree metered dose inhaler (4 × 20 µg puffs qid) has been found to be effective.[5, 17]

However, for patients with idiopathic chronic cough it is important to provide ongoing interest and support, and not eventually to dismiss them as 'just a cough'.

BRONCHIAL CARCINOMA

Lung cancer accounts for 25% of cancer deaths in men and 24% of cancer deaths in women (rapidly rising), with cigarette smoking being the most common cause of lung cancer in both sexes.[11] It is also the most common lethal cancer in both sexes in Australia. Bronchogenic carcinoma accounts for over 95% of primary lung malignancies. The prognosis is poor - the 5 year overall survival is 12 - 14%.[18] The mesothelioma nadince continues to rise.

Clinical features

- Most present between 50 and 70 years (mean 67)
- Only 10 - 25% asymptomatic at time of diagnosis[9]
- If symptomatic - usually advanced and not resectable

Local symptoms

- Cough (42%)
- Chest pain (22%)
- Wheezing (15%)
- Haemoptysis (7%)
- Dyspnoea (5%)

General

- Anorexia, malaise
- Weight loss unexplained

Others

- Unresolved chest infaction
- Hoarseness
- Symptoms from metastases

The possible physical findings are summarised in Figure 42.4.


Local complications
Horner's syndrome
• small pupil
• ptosis
• loss of sweating (ipsilateral)
• enophthalmos
rib pain
hoarseness
bovine cough
severe inner arm pain (brachial plexus)
dysphagia
pleural effusion
cough
haemoptysis
Metastatic complications
brain
• headache
• personality change
• seizures
• limb weakness/sensory loss
• vomiting
bone
• back pain (vertebrae)
• rib pain
liver
• jaundice (late)
Also
• anaemia
• paraneoplastic syndromes
• paralysis diaphragm

Figure 42.4 Possible physical findings of bronchial carcinoma

Investigations

- Chest X-ray
- CT scanning
- Fibre-optic bronchoscopy
- PET scanning
- Fluorescence bronchoscopy (helps early detection)
- Tissue diagnosis where pussible

Note: No proven effect of any screning for a symptomatic patients.

Causes of a solitary pulmonary nodule on X-ray are presented in Table 42.7.

Table 42.7 Causes of a solitary pulmonary nodule (on X-ray)[19]

Common
• Bronchogenic carcinoma
• Solitary metastasis
• Granuloma (e.g. TB)

• Hamartoma
Less common
• Bronchial adenoma
• AVM
• Hydatid
• Others (e.g. haemotoma, cyst)

Management

Refer to a respiratory physician to determine the type of carcinoma. They are usually classified as small cell lung (oat cell) poorly differentiated carcinoma (15 - 20% incidence) (SCLC) and non-small cell lung cancer (NSCLC), which includes squamous cell carcinoma, adenocarcinoma and large cell carcinoma (approximately 20 - 30% of each). The main aim of management is a curative resection of NSCLC in those who can benefit from it. Sargery is not an option for SCLC since it inctastasises so rapedly (80% have metastasised at the time of diagnosis).[14] Chemotherapy is suitable for the deadly SCLC but currently only extends life expectancy from 3 to 20 months (at best). It also has an important little place in treating NSCLC. The main role of radiotherapy is palliative.

BRONCHIECTASIS

Bronchiectasis is dilatation of the bronchi when their walls become inflamed, thickened and irreversibly damaged, usually following obstruction followed by infection. Predisposing causes include whooping cough, measles, TB, inhaled foreign body (e.g. peanuts in children), bronchial carcinoma, cystic fibrosis and congenital ciliary dysfunction (Kartagener's syndrome). The left lower lobe and lingula are the commonest sites for localised disease.

Clinical features

- Chronic cough - worse on waking
- Mild cases: yellow or green sputum only after infection
- Advanced
    - profuse purulent offensive sputum
    - persistent halitosis
    - recurrent febrile episodes
    - malaise, weight loss
- Episodes of pneumonia
- Sputum production related to posture
- Haemoptysis (bloodstained sputum or massive) possible

Physical examination

- Clubbing
- Coarse crackles over infected areas (usually lung base)
- Other respiratory signs are given in Table 49.6

Investigations

- Chest X-ray (normal or bronchial changes)
- Sputum examination
    - for resistant pathogens
    - to exclude TB
- Main pathogens: *Streptococcus pneumoniae, Pseudomonas aeruginosa, Haemophilus influenzae* (commonest)
- CT scan: can show bronchial wall thickening
- Bronchograms: very unpleasant and used only if diagnosis in doubt or possible localised disease amenable to surgery (rare)

Management[14]

- Explanation and preventive advice.
- Postural drainage (e.g. lie over side of bed with head and thorax down for 10 - 20 minutes 3 times a day).

- Antibiotics according to organism - it is important to eradicate infection to halt the progress of the disease. Amoxycillin 500 mg (o) tds for 2 - 3 weeks is recommended for first presentation.[14]
- Bronchodilators if evidence of bronchospasm.

TUBERCULOSIS

Pulmonary TB may be symptomless and detected by mass X-ray screening.[8]

(Refer to Chapter 30, p. 164)

Typical clinical features

Respiratory symptoms

- Cough
- Sputum: initially mucoid, later purulent
- Haemoptysis
- Dyspnoea (especially with complications)
- Pleuritic pain

General (usually insidious)

- Anorexia
- Fatigue
- Weight loss
- Fever (low grade)
- Night sweats

Physical examination

- May be no respiratory signs or signs of fibrosis, consolidation or cavitation (amphoric breathing)
- Finger clubbing

Investigations

- Chest X-ray
- Micro and culture sputum (for tubercle bacilli)
- ESR
- Tuberculin test (misleading if previous BCG vaccination)

Management

TB is a notifiable disease and must be reported to state (and local) health departments. Hospitalisation for the initial therapy of pulmonary TB is not necessary in most patients. Monthly follow-up is recommended, including sputum smear and culture. Multiple drug therapy is initiated primarily to guard against the existence and/or emergence of resistant organisms. Standard initial therapy consists of rifampicin + ethambutol + isoniazid + pyrazinamide daily for at least 2 months, followed by rifampicin + isoniazid for 4 months if the organism is susceptible to these drugs.[11] If isoniazid resistance is suspected, streptomycin (with care) is added.

SYMPTOMATIC TREATMENT OF COUGH

Symptomatic treatment of cough should be reserved for patients who have acute self-limiting causes of cough, especially an acute viral infection. There are many cough mixtures available and the major constituents of these mixtures are shown in Table 42.8 [8]. The recommended mixture should be tailored to the patient's individual requirements. These mixtures should be used only in the short term.[20] Cough medicines are generally not recommended in children but for a very dry irritating cough in children 5 - 12 years use pholcodine 5 - 10 mg (o) tds or qid.[14]

Table 42.8 Cough mixtures: major constituents

Cough suppressants

Opioid
Codeine
Dihydrocodeine
Hydrocodone
Pholcodine
Ethylmorphine
Normethadone
Other
Carbetapentane
Dextromethorphan
Oxolamine
Expectorants/mucolytics
Senega
Ammonia
Guaiphenesin
Bromhexine
Analgesics/antipyretics
Paracetamol (acetaminophen)
Salicylates (e.g. aspirin)
Decongestants
Sympathomimetic
Ephedrine
Pseudoephedrine
Phenylephrine
Phenylpropanolamine
Methoxyphenamine
Antihistamine
Promethazine
Pheniramine
Chlorpheniramine
Diphenhydramine
Dexchlorpheniramine
Brompheniramine
Triprolidine
Anticholinergic
Atropine
Isopropamide

Source: After Williams and Bowes[8]

WHEN TO REFER

- Patients in whom bronchoscopy is necessary to exclude bronchial carcinoma
- Persistent hoarseness in a patient who requires expert laryngeal examination
- Evidence of pulmonary TB

Practice tips

- Unexplained cough over the age of 50 is bronchial carcinoma until proved otherwise (especially if there is a history of smoking).
- Consider TB in the presence of an unusual cough ± wheezing
- Bronchoscopy is essential to exclude adequately a suspicion of bronchial carcinoma when the chest X-ray is normal.
- Bright red haemoptysis in a young person may be the initial symptom of pulmonary TB.
- Avoid settling for a diagnosis of bronchitis as an explanation of haemoptysis until bronchial carcinoma has been excluded.
- Coughing may be so severe that it terminates in vomiting or loss of consciousness (post-tussive syncope).
- Large haemoptyses are usually due to bronchiectasis or TB.
- The presence of white cells in the sputum renders it yellow or green (purulent) but does not necessarily imply infection.

REFERENCES

1. Kumar PJ, Clarke ML. *Clinical Medicine* (2nd edn). London: Bailliere Tindall, 1990: 637 - 8.
2. Walsh TD. *Symptom Control*. Oxford: Blackwell Scientific Publications, 1989; 81: 81 - 8, 235 - 9.
3. Fitzgerad D. Children with chronic or recurrent cough. Medical obsener. 22 nobember 2002;32 - 3.
4. Burns M. Chronic cough. Aust Fam Physician, 1996; 25: 161 - 7.
5. Ing A. Intractable cough. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 130 - 3.
6. Kincaid-Smith P, Larkins R, Whelan G. *Problems in Clinical Medicine*. Sydney: MacLennan&Petty, 1990: 105 - 8.
7. Davis A, Bolin T, Ham J. *Symptom Analysis and Physical Diagnosis* (2nd edn). Sydney: Pergamon Press, 1990: 56 - 60.
8. Williams TJ, Bowes G. Cough as a symptom in adult life. Mod Med Aust, 1992; June: 84 - 92.
9. Selecki Y, Helman A. Chronic cough in children. Aust Dr Weekly, 1989; 18 Sept.: i - iv.
10. Smyrinos NA et al. From a prospective study of chronic cough: diagnostic and therapeutic aspects in older adults. Arch Intern Med, 1988; 158: 1222 - 8.
11. Tierney LM et al. *Current Medical Diagnosis and Treatment* (45th edn) McGraw-Hill and Lange, 2006: 266 - 72.
12. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 199 - 221.
13. Barton S ed. *Clinical Evidence*. London: BMJ Publishing Group, 2001: 319 - 22.
14. Dowden J, (Chair). *Therapeutic Guidelines: Respiratory* (Version 3). Melbourne: Therapeutic Guidelines Ltd, 2005:164-202.
15. Christiansen K. Antibiotics for common respiratory infections. Aust Fam Physician, 1995; 24: 49 - 56.
16. Smart J. *Paediatric Handbook* (6th edn). Melbourne: Blackwell Science, 2000, 499 - 501.
17. Holmes PW, Barter CE, Pierce RJ. Chronic persistent cough: use of ipratropium bromide in undiagnosed cases following URTI. Respir Med, 1992; 86: 425 - 9.
18. NH&MRC. *Assessment and Management of Long Cancer: Evidence based guidelines. A guide for GPs*. NH&MRC,2005.
19. Lau L ed. *Imaging Guidelines* (4th edn). Melbourne: RANZC Radiologists, 2001: 64.
20. Schroeder K, Fahey T. Systematic review of randomised controlled trials of over the counter cough medicines for acute cough. Br Med J, 2002; 324: 329 - 31.

CHAPTER 43. DEAFNESS AND HEARING LOSS

There are two kinds of deafness. One is due to wax and is curable; the other is not due to wax and is not curable.

Sir William Wilde (1815 - 1876)

DEAFNESS AND HEARING LOSS

Deafness is defined as impairment of hearing, regardless of its severity.[1] It is a major community health problem requiring a high index of suspicion for diagnosis, especially in children. Deafness may be conductive, sensorineural or a combination of both (mixed).

KEY FACTS AND CHECKPOINTS

- Deafness occurs at all ages but is more common in the elderly (Figure 43.1). Fifty per cent of people over 80 years have deafness severe enough to be helped by a hearing aid.



Figure 43.1 Prevalence of hearing problems with increasing age

- The threshold of normal hearing is from 0 to 20 decibels (dB), about the loudness of a soft whisper.
- One in seven of the adult population suffers from some degree of significant hearing impairment (over 20 dB in the better hearing ear).[2]
- One child in every 1000 is born with a significant hearing loss.
- Degrees of hearing impairment:[2, 3]
  - mild ° loss of 20 - 40 dB (20 dB is soft-spoken voice)
  - moderate ° loss of 40 - 70 dB (40 dB is
  - normal-spoken voice)
  - severe ° loss of 70 - 90 dB (shout)
  - profound ° loss of over 90 dB
- More women than men have a hearing loss.
- People who have worked in high-noise levels (> 85 dB) are more than twice as likely to be deaf.
- There is a related incidence of tinnitus with deafness.

A DIAGNOSTIC APPROACH

It is useful to consider the causes of deafness in terms of pathophysiology (conductive or sensorineural hearing loss) and anatomical sites (Figure 43.2).


Conduction deafness
Sensorineural deafness
cochlear (sensory) type
retrocochlear (neural) type
cerumen (wax)
osteomata
otitis externa
Eustachian tube dysfunction
congenital meatal atresia
chronic otitis media
'glue' ears
barotrauma
perforated TM
otosclerosis
Paget's disease of bone
cholesteatoma
trauma
viral infection
syphilis and TB
presbyacusis
noise-induced deafness
ototoxic drugs
Meniere's syndrome
congenital cochlear deafness
cochlear otosclerosis
acoustic neuroma
other cerebellopontine angle tumours
herpes zoster
central causes

Figure 43.2 Causes of deafness according to anatomical site

Conductive hearing loss is caused by an abnormality in the pathway conducting sound waves from the outer ear to the inner ear,[1] as far as the footplate of the stapes.

Sensorineural hearing loss (SNHL) is a defect central to the oval window involving the cochlear (sensor), cochlear nerve (neural) or, more rarely, central neural pathways.[1]

Congenital deafness is an important consideration in children, while presbyacusis is very common in the aged. The commonest acquired causes of deafness are impacted cerumen (wax), serous otitis media and otitis externa. Noise-induced deafness is also a common problem.

It is important not to misdiagnose an acoustic neuroma, which can present as acute deafness, although slow progressive loss is more typical. A summary of the diagnostic strategy, which includes several important causes of deafness, is presented in Table 43.1 and a checklist of ototoxic drugs in Table 43.2.

Table 43.1 Deafness and hearing loss: diagnostic strategy model

Q. Probability diagnosis
A. Impacted cerumen
Serous otitis media

Otitis externa Congenital (children) Presbyacusis
Q. Serious disorders not to be missed A. Neoplasia • acoustic neuroma • temporal lobe tumours (bilateral) • otic tumours Severe infections • generalised infections (e.g. mumps, measles) • meningitis • syphilis Perforated tympanic membrane Cholesteatoma Perilymphatic fistula (post-stapedectomy) Meniere's syndrome
Q. Seven masquerades checklist A. Depression - Diabetes Drugs Anaemia - Thyroid disorder hypo Spinal dysfunction - UTI -
Q. Is this patient trying to tell me something? A. Unlikely.

Table 43.2 Known ototoxic drugs

Alcohol
Aminoglycosides
• amikacin
• gentamicin
• kanamycin
• neomycin
• streptomycin
• tobramycin
Diuretics
• ethacrynic acid
• frusemide
Chemotherapeutic agents
Quinine and related drugs
Salicylates

Symptoms

The symptoms vary so that some barely notice a problem while others are severely disabled.

Common symptoms include inability to:

- hear speech and other sounds loudly enough
- hear speech and music clearly, even when loud enough
- understand speech even when loud enough - a problem of language reception

People with mild hearing loss notice only subtle differences and may have trouble hearing certain high-frequency sounds, such as ‘s’, ‘f’ or ‘th’. They may also have trouble hearing in certain situations, such as at a party or in a crowd where there is a lot of background noise. Those with moderate hearing loss have trouble hearing in many situations.

THE CLINICAL APPROACH

History

The history should include an account of the onset and progression of any deafness, noise exposure, drug history, a history of swimming or diving, air travel and head injury and family history. A recent or past episode of a generalised infection would be relevant and the presence of associated aural symptoms, such as ear pain, discharge, tinnitus and vertigo. Vertigo may be a symptom of Meniere's syndrome, multiple sclerosis, acoustic neuroma or syphilis.

Several important clues can be obtained from the history. The often sudden onset of hearing loss in an ear following swimming or showering is suggestive of wax, which swells to block the ear canal completely.

Patients with conductive loss may hear better in noisy conditions (paracusis) because we raise our voices when there is background noise. Conversely, people with sensorineural deafness (SND) usually have more difficulty hearing in noise as voices become unintelligible.

Examination

Inspect the facial structures, skull and ears. The ears are inspected with an otoscope to visualise the external meatus and the tympanic membrane (TM) and the presence of obstructions such as wax, inflammation or osteomata.

The examination requires a clean external auditory canal. Gentle suction is useful for cleaning pus debris. Syringing is reserved for wax in people with an intact TM and a known healthy middle ear.

It is an advantage to have a pneumatic attachment to test drum mobility. Reduction of TM mobility is an important sign in secretory otitis media.

There are several simple hearing tests. The distance at which a ticking watch can be heard can be used but the advent of the digital watch has affected this traditional method.

The hair-rubbing method

In children and in adults with a reasonable amount of hair grab several hairs close to the external auditory canal between the thumb and index finger. Rub the hairs lightly together to produce a relatively highpitched ‘crackling’ sound (Figure 43.3). If this sound cannot be heard, a moderate hearing loss is likely (usually about 40 dB or greater). Like the whisper test, this test is a rough guide only.



Figure 43.3 The simple hair-rubbing method of testing possible deafness

Tuning fork tests

If deafness is present, its type (conduction or sensorineural) should be determined by tuning fork testing. The most suitable tuning fork for preliminary testing is the C_2 (512 cps) fork. The fork is best activated by striking it firmly on the bent elbow.

Weber test

The vibrating tuning fork is applied firmly to the midpoint of the skull or to the central forehead or to the teeth.

This test is of value only if the deafness is unilateral or bilateral and unequal (Figure 43.4). Normally the sound is heard equally in both ears in the centre of the forehead. With sensorineural deafness the sound is transmitted to the normal ear, while with conduction deafness it is heard better in the abnormal ear.


In which ear is the sound louder?

Figure 43.4 Weber test

Rinne's test

The tuning fork is held:

- outside the ear (tests air conduction) and
- firmly against the mastoid bone (tests bone conduction)

It therefore compares air and bone conduction in the same ear (Figure 43.5). A variation of the test includes placing the tuning fork on the mastoid process and the patient indicates when it can no longer be heard. The fork is then placed at the external auditory meatus and the patient indicates whether the sound is now audible. Normally air conduction is better than bone conduction and the sound will again be heard.



Figure 43.5 Rinne's test comparing air conduction (a) with bone conduction (b)

A comparison of the interpretation of these tests is summarised in Table 43.3.

Table 43.3 A comparison of Rinne's and the Weber tests

State of the hearing	Rinne's test	Weber test
Normal	Positive: AC > BC	Equal in both ears
Conduction deafness	Negative: BC > AC	Louder in the deaf ear
Very severe conduction deafness	Negative: BC > AC May hear BC only	Louder in the deaf ear
Sensorineural deafness	Positive: AC > BC	Louder in the better ear
Very severe sensorineural deafness	'False' negative (without masking)	Louder in the better ear

AC ° air conduction; BC ° bone conduction

Audiometric assessment

Audiometric assessment includes the following:

- pure tone audiometry
- impedance tympanometry
- electric response audiometry
- oto-acoustic emission testing

Pure tone audiometry[4]

Pure tone audiometry is a graph of frequency expressed in hertz versus loudness expressed in decibels. The tone is presented either through the ear canal (a test of the conduction and the cochlear function of the ear) or through the bone (a test of cochlear function).

Figures 43.6 and 43.7 are typical examples of pure tone audiograms.



Figure 43.6 Pure tone audiogram for severe conductive deafness in right ear. After B. Black [4]



Figure 43.7 Pure tone audiogram for unilateral (right) sensorineural deafness. Suspect a viral or congenital origin in children; check adults for acoustic neuroma

The difference between the two is a measure of conductance. If the two ears have different thresholds, a white noise masking sound is applied to the better ear to prevent it hearing sound presented to the test ear. The normal speech range occurs between 0 and 20 decibels in soundproof conditions across the frequency spectrum.

Tympanometry

Tympanometry measures the mobility of the tympanic membrane, the dynamics of the ossicular chain and the middle ear air cushion. The test consists of a sound applied at the external auditory meatus, otherwise sealed by the soft probe tip.

DEAFNESS IN CHILDREN

Deafness in childhood is relatively common and often goes unrecognised. One to two of every 1000 newborn infants suffer from SND.[1] Congenital deafness may be due to inherited defects, to prenatal factors such as maternal intrauterine infection or drug ingestion during pregnancy, or to perinatal factors such as birth trauma, and haemolytic disease of the newborn.

Deafness may be associated with Down syndrome and Waardenburg's syndrome. Waardenburg's syndrome, which is dominantly inherited, is diagnosed in a patient with a white forelock of hair and different coloured eyes.

Acquired deafness accounts for approximately half of all childhood cases. Purulent otitis media and secretory otitis media are common causes of temporary conductive deafness. However, one in 10 children will have persistent middle ear effusions and mild to moderate hearing loss in the 15 - 40 decibel range.[5]

Permanent deafness in the first few years of life may be due to virus infections, such as mumps or meningitis, ototoxic antibiotics and several other causes.

Screening[1]

The aim of screening should be to recognise every deaf child by the age of 8 months to 1 year - before the vital time for learning speech is wasted. High-risk groups should be identified and screened, for example, a family history of deafness, maternal problems of pregnancy, perinatal problems, survivors of intensive care, very low birthweight and gestation < 33 weeks, cerebral palsy and those with delayed or faulty speech. The guidelines for early signs of normal hearing are presented in Table 43.4.

Table 43.4 Early signs of normal hearing

Age	Typical response
1 month	Should notice sudden constant sounds (e.g. car motor, vacuum cleaner) by pausing and listening.
3 months	Should respond to loud noise (e.g. will stop crying when hands areclapped).
4 months	Should turn head to look for source of sound, such as mother speaking behind the child.
7 months	Should turn instantly to voices or even to quiet noises made across the room.
10 months	Should listen out for familiar everyday sounds.
12 months	Should show some response to familiar words and commands, including his or her name.

Optimal screening times:

- 8 to 9 months (or earlier)
- school entry

There has been a proposition from the National Universal Hearing Screening Committee to introduce testing of newborn hearing following results[6] supporting this strategy after the finding of a higher incidence than appreciated. The average age at which these deaf children were diagnosed was 26 months.

Early signs of hearing loss

A high index of suspicion is essential in detecting hearing loss in children and any parental concern should be taken seriously. The presentation of hearing loss will depend on whether it is bilateral or unilateral, its severity and age of onset.

Typical presentations include:

- malformation of skull, ears or face
- failure to respond in an expected way to sounds, especially one's voice
- preference for, or response only to, loud sounds
- no response to normal conversation or to television
- speech abnormality or delay
- absence of ‘babbling’ by 12 months
- no single words or comprehension of simple words by 18 months
- learning problems at school
- disobedience
- other behavioural problems
- inability to detect sound direction (unilateral loss)
- inability to follow simple commands or less than 20 spoken words by 2 years

Screening methods

Hearing can be tested at any age. No child is too young to be tested and this includes the newborn. Informal office assessments, such as whispering in the child's ear or rattling car keys, are totally inadequate for excluding deafness and may be potentially harmful if they lead to false reassurance.

Pneumatic otoscopy is essential to exclude middle ear effusions.

Pure tone audiometry is unreliable in children under 4 years of age, so special techniques such as tympanometry are required. Tympanometry assesses TM compliance, and is highly sensitive and specific for detecting middle-ear pathology in children beyond early infancy.

Auditory brain stem response testing is used to evaluate children (particularly young infants) for whom information on behavioural hearing tests is either unobtainable or unreliable.[5]

Management

Children with middle ear pathology and hearing loss should be referred to a specialist. All children with SNHL (even those with profound deafness), as well as children with conductive losses not correctable by surgery, benefit from amplification. All children need referral to a specialist centre skilled in educational and language remediation.

DEAFNESS IN THE ELDERLY

The prevalence of hearing loss increases exponentially with age. The commnoest reason for bilateral progressive SND is presbyacusis, which is the highfrequency hearing loss of advancing age (Figure 43.8). There appears to be a genetic predisposition to presbyacusis.



Figure 43.8 Presbyacusis: bilateral high-frequency sensorineural deafness

Some features include:

- loss of high-frequency sounds
- usually associated with tinnitus
- intolerance to very loud sounds
- difficulty picking up high-frequency consonants, e.g. 'f', 's' - these sounds are often distorted or unheard, and there is confusion with words such as 'fit' and 'sit', 'fun' and 'sun'

Deafness is associated with various types of mental illness in the aged, including anxiety, depression, paranoid delusions, agitation and confusion because of sensory deprivation. The possibility of deafness should be kept in mind when assessing these problems.

Signs indicating referral for hearing test

Possible indications for referring the older person:

- speaking too loudly
- difficulty understanding speech
- social withdrawal
- lack of interest in attending parties and other functions
- complaints about people mumbling
- requests to have speech repeated
- complaints of tinnitus

- setting television and radio on high volume

SUDDEN DEAFNESS

Sudden deafness refers to sudden SNHL of greater than 30 - 35 dB with an onset period of between 12 hours and 3 days.[6] It specially excludes gradual progressive causes of SND, such as cumulative noise trauma or presbyacusis and also excludes causes of sudden deafness that may be related to pathology in the external auditory canal, TM or middle ear.

The main causes are given in Table 43.5.

Table 43.5 Causes of sudden deafness

Trauma
• head injury
• diving
• flying
• acoustic blast
Postoperative
• previous stapedectomy
Viral infections (e.g. mumps, measles, herpes zoster)
Ototoxic drugs (e.g. aminoglycosides)
Cerebellopontine angle tumours (e.g. acoustic neuroma)
Vascular disease
• polycythaemia
• diabetes
Meniere's syndrome
Cochlear otosclerosis

In several instances, despite a careful clinical examination and investigation, an explanation for sudden SND cannot be found. The cause of deafness in these cases is thought to be either vascular obstruction of the end artery system or viral cochleitis.[5, 6] Fortunately, spontaneous recovery usually results.

Patients with sudden SND require immediate referral. It is a difficult problem both in diagnosis and management. Early diagnosis and a high index of suspicion are fundamental.[7] Two important conditions that deserve special reference are perilymphatic fistula, which occurs after stapedectomy, and an acoustic neuroma presumably causing compression of the internal auditory artery by the tumour in the internal auditory meatus.

OTOSCLEROSIS

Otosclerosis is a disease of the bone surrounding the inner ear and is the most common cause of conductive hearing loss in the adult with a normal tympanic membrane. The normal middle ear bone is replaced by vascular, spongy bone that becomes sclerotic.

Features[3]

Usually:

- a progressive disease
- develops in the 20s and 30s
- family history (autosomal dominant)
- bilateral or unilateral
- female preponderance
- affects the footplate of the stapes

- may progress rapidly during pregnancy
- conductive hearing loss
- begins in lower frequencies, then progresses
- SND may be present
- impedance audiometry shows characteristic features (Figure 43.9)



Figure 43.9 Otosclerosis: the conductive loss is commonly associated with a mild sensorineural loss at 2000 cycles per second (Carhardt's notch)

- may be associated with Meniere's syndrome

Management

- Referral to an ENT consultant
- Stapedectomy (approximately 90% effective)
- Hearing aid (less effective alternative)

CHOLESTEATOMA[7]

A cholesteatoma is a sac of keratinising squamous epithelium that arises from a perforation involving the periphery of the TM. In other words, it is a 'big sac of skin' (refer to p. 545). It is dangerous to the ear because it tends to expand and destroy adjacent structures, including the TM, ossicular chain and cochlear. Destruction of the first two may result in conductive hearing loss of up to 60 dB. Irreversible SND caused by otic capsule erosion may also occur. Surgical correction is mandatory.

NOISE-INDUCED HEARING LOSS

Features

- Onset of tinnitus after work in excessive noise
- Speech seems muffled soon after work

- Temporary loss initially but becomes permanent if noise exposure continues
- High-frequency loss on audiogram

Sounds exceeding 85 dB are potentially injurious to the cochlea, especially with prolonged exposures. Common sources of injurious noise are industrial machinery, weapons and loud music.

TINNITUS

Tinnitus is defined as a sound perceived by the ear that arises from an internal source. When pathology in the inner ear is the cause, the tinnitus is non - pulsating, continuous and may have variable frequencies and intensity.

A thorough history and examination should be conducted so that tinnitus can be classified as objective (e.g. heard with stethoscope) or non-objective and pulsatile, or non-pulsatile.

Precautions:

- exclude wax, drugs including marijuana, NSAIDs, salicylates, quinine and aminoglycosides,[8] vascular disease, depression, aneurysm vascular tumours (e.g. glomus tumour), venous hum (jugular vein), acoustic neuroma, Meniere's syndrome and infections (e.g. viral cochleitis)
- beware of lonely elderly people living alone (suicide risk)

Note: Otosclerosis in young adults causes deafness and tinnitus.

Investigations

- Audiological examination by audiologist
- Tympanometry and speech discrimination
- MRI (if serious cause suspected)

Management

- Treat any underlying cause and aggravating factors. Otherwise, minimise symptoms.
- Educate and reassure the patient (tinnitus is nearly always amenable to treatment).

Holistic approach (options)

Mainly based on acoustic de-sensitisation

- Relaxation techniques
- Tinnitus retraining therapy (clinical psychologist)
- Cognitive behaviour therapy
- Background ‘noise’ (e.g. music played during night for masking)
- Tinnitus maskers
- Hearing aids (based on audiologist assessment)
- Consider hypnotherapy

Medical (trials of options)

- Clonazepam 0.5 mg nocte
- Minerals (e.g. zinc and magnesium)
- Betahistine (Serc) 8 - 16 mg daily (max. 32 mg)
- Carbamazepine
- Antidepressants

Acute severe tinnitus

- Lignocaine 1% IV slowly (up to 5 mL)

HEARING AIDS

Hearing aids are most useful in conductive deafness. This is due to the relative lack of distortion, making amplification simple.

In SND the dual problem of recruitment and the hearing loss for higher frequencies may make hearing aids less satisfactory. Modern aids selectively amplify higher frequencies and ‘cut out’ excessive volume peaks that would cause discomfort. A trial of such aids should be made by a reliable hearingaid consultant following full medical assessment.

Cochlear implants[9]

The cochlear implant or ‘bionic ear’ is used in adults and children with severe hearing loss unresponsive to powerful hearing aids. The implant consists of an array of 22 electrodes, inserted into the cochlear following mastoidectomy, attached to a receiver implanted in the skull next to the ear. External sounds are detected by an external processor worn behind the ear and connected to the implanted receiver with an external induction coil. Near normal speech and hearing may be achieved in children with congenital or acquired deafness with early implantation. The device is most suitable for children over 2 years and adults with severe deafness.

ADVICE FOR FAMILIES

Relatives and close friends need considerable advice about coping with deaf members. They should be told that the deaf person may hear in a quiet room but not in a crowd, and advised of the range of aids and services available and the importance of proper maintenance of any hearing aids (especially with aged people).

Do

- Face the light when speaking to them.
- Speak directly to them.
- Speak clearly and naturally.
- Speak at a uniform pitch: avoid lowering your voice during or at the end of a sentence.
- Speak within 2 metres.
- Be tolerant and relaxed.
- Be patient with mistakes.
- Write key words on a paper pad when necessary.

Don't

- Speak with your back to them.
- Mumble your words.
- Use exaggerated lip movements.
- Shout.
- Put your hand or fingers over your mouth when talking.
- Repeat one word over and over.

WHEN TO REFER

- Sudden deafness.
- Any child with suspected deafness, including poor speech and learning problems, should be referred to an audiology centre.
- Any child with middle ear pathology and hearing loss should be referred to a specialist.
- Unexplained deafness.

Practice tips

- A mother who believes her child may be deaf is rarely wrong in this suspicion.
- Suspect deafness in an infant with delayed development and in children with speech defects or behavioural problems.
- Audiological assessment should be performed on children born to mothers with evidence of intrauterine infection by any of the TORCH organisms (toxoplasmosis, rubella, cytomegalovirus and herpes virus).
- No child is too young for audiological assessment. Informal office tests are inadequate for excluding hearing loss.
- Sounds tend to be softer in a conductive hearing loss and distorted with sensorineural loss.
- People with conductive deafness tend to speak softly, hear better in a noisy environment, hear well on the telephone and have good speech discrimination.

- People with SND tend to speak loudly, hear poorly in a noisy environment, have poor speech discrimination and hear poorly on the telephone.

REFERENCES

1. Ludman H. *ABC of Otolaryngology* (3rd edn). London: British Medical Association, 1993: 10 - 22.
2. Fagan P. Assessing hearing in clinical practice. Medical Observer, 10 March 2006: 23 - 5.
3. Tierney LM et al. *Current Medical Diagnosis and Treatment* (45th edn). New York: The McGraw-Hill Companies, 2006: 181 - 2.
4. Black B. Pure tone audiograms. Aust Fam Physician, 1988; 17: 906 - 7.
5. Jarman R. Hearing impairment. Aust Paediat Rev, 1991; 4: 2.
6. Fortnum HM et al. Prevalence of permanent childhoood hearing impairment in the United Kingdom and implications for universal neonatal hearing screening: questionnairebased ascertainment study. Br Med J, 2001; 323: 536 - 40.
7. Pohl DV. Sudden deafness. Mod Med Aust, 1990; June: 72 - 8.
8. Black B, Harvey L. Tinnitus: update. Medical Observer, 13 August 2004: 31 - 3.
9. Atlas M, Lowinger D. The GP's essential guide to hearing loss. Med Today, 2000; June: 59.L

CHAPTER 44. DIARRHOEA

A dirty cook gives diarrhoea quicker than rhubarb.

Tung-su Pai (time uncertain)

DIARRHOEA

Diarrhoea is defined as the frequent passage of loose or watery stools.

Essential features are:

- an increase in frequency of bowel action
- an increase in softness, fluidity or volume of stools

Acute self-limiting diarrhoea, which is very common and frequently not seen by the medical practitioner, is usually infective and mild, and resolves within days. In Australia most infective cases are viral. The causes of diarrhoea are numerous, thus making a detailed history and examination very important in leading to the diagnosis. Important causes are presented in Figure 44.1.


General
• diet
• antibiotics
• laxatives
• irritable bowel syndrome
• lactose intolerance
Organic disease of bowel
chronic infections (e.g. *Giardia lamblia*)
carcinoma of bowel
diverticular disorder
inflammatory bowel disease
• ulcerative colitis
• Crohn's disease
• pseudomembranous colitis
impacted faeces with spurious diarrhaea
'gay bowel'

Figure 44.1 Important causes of chronic diarrhoea

The terminology for acute infective diarrhoea can be confusing. A simple classification is:

- vomiting and diarrhoea ° gastroenteritis
- diarrhoea (only) ° enteritis

KEY FACTS AND CHECKPOINTS

- The characteristics of the stool provide a useful guide to the site of the bowel disorder.
- Disorders of the upper GIT tends to produce diarrhoea stools that are copious, watery or fatty, pale yellow or green.
- Colonic disorder tends to produce stools that are small, of variable consistency, brown and may contain blood or mucus.
- Acute gastroenteritis should be regarded as a diagnosis of exclusion.
- Chronic diarrhoea is more likely to be due to protozoal infection (amoebiasis, giardiasis or *Cryptosporidium*) than bacillary dysentery.
- A history of travel, especially to countries at risk of endemic bowel infections, is essential.
- Certain antibiotics can cause an overgrowth of *Clostridium difficile*, which produces pseudomembranous colitis.
- Coeliac disease, although a cause of failure to thrive in children, can present at any age.
- In disorders of the colon the patient experiences frequency and urgency but passes only small amounts of faeces.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 44.1.

Table 44.1 Diarrhoea: diagnostic strategy model

Q. Probability diagnosis A. Acute • Gastroenteritis/infective enteritis • Dietary indiscretion • Antibiotic reaction Chronic • Irritable bowel syndrome • Drug reactions (e.g. laxatives) • Chronic infections
Q. Serious disorders not to be missed A. Neoplasia • Carcinoma of bowel HIV infection (AIDS) Infections • Cholera • Typhoid/paratyphoid • Amoebiasis • Malaria • Enterohaemorrhagic *E. coli* enteritis Inflammatory bowel disease • Crohn's/ulcerative colitis • Pseudomembranous colitis Intussusception Pelvic appendicitis/pelvic abscess
Q. The pitfalls A. Coeliac disease Faecal impaction with spurious diarrhoea Lactase deficiency *Giardia lamblia* infection *Cryptosporidium* infection Malabsorption states (e.g. coeliac disease) Vitamin C and other oral drugs Radiotherapy Diverticulitis Post GIT surgery 'Gay bowel' Ischaemic colitis (elderly) *Rarities* • Addison's disease (p. 224) • Carcinoid tumours • Short bowel syndrome • Amyloidosis

- Toxic shock
- Zollinger-Ellison syndrome

Q. Seven masquerades checklist

A. Depression -
Diabetes
Drugs
Anaemia -
Thyroid disorder hyper
Spinal dysfunction -
UTI -

Q. Is the patient trying to tell me something?

A. Yes, diarrhoea may be a manifestation of anxiety state or irritable bowel syndrome.

Probability diagnosis

Acute diarrhoea

Common causes are:

- gastroenteritis/enteritis
    - bacterial
        - *Salmonella* sp.
        - *Campylobacter jejuni*
        - *Shigella* sp.
        - enteropathic *Escherichia coli*
        - *Staphylococcus aureus* (food poisoning)
    - viral
        - rotavirus (50% of children hospital admissions[1])
        - norovirus
- dietary indiscretions (e.g. binge eating)
- antibiotic reactions

Chronic diarrhoea

Irritable bowel syndrome was the commonest cause of chronic diarrhoea in a UK study.[2]

Drug reactions are also important. These include ingestion of laxatives, osmotic agents such as lactose and sorbitol in chewing gum, alcohol, antibiotics, thyroxine and others.

Acute gastroenteritis that persists into a chronic phase is relatively common, especially in travellers returning from overseas. Important considerations are *Giardia lamblia, C. difficile, Yersinia, Entamoeba histolytica, Cryptosporidium* and HIV infection.

Serious disorders not to be missed

Colorectal carcinoma must be considered with persistent diarrhoea, especially if of insidious onset.

AIDS due to symptomatic HIV infection needs consideration, especially in those at risk. The serious infectious disorders that can affect international travellers, such as cholera, typhoid, paratyphoid and amoebiasis, should also be kept in mind.

In children coeliac disease and fibrocystic disease can present as chronic diarrhoea while intussusception, although not causing true diarrhoea, can present as loose redcurrant jelly-like stools and should not be misdiagnosed (as gastroenteritis). Appendicitis must also be considered in the onset of acute diarrhoea and vomiting.

Infection with enterohaemorrhagic strains of *E. coli* (e.g. O157:H7, O111:H8), may lead to the haemolytic uraemic syndrome or thrombotic thrombocytopenic purpura, particularly in children. What appears to be simple enteritis can eventuate to be fatal.

Clue: Think of it with atypical gastroenteritis and bloody diarrhoea.

Pitfalls

There are many traps in evaluating the patient with diarrhoea, including drug ingestion, especially vitamin C (sodium ascorbate powder), which causes diarrhoea. Faecal impaction with spurious diarrhoea is an age-old pitfall, as is lactase deficiency, which may go undiagnosed for many years. In recent times infection with *G. lamblia* may smoulder on for months with watery offensive stools before diagnosis.

General pitfalls

- Not considering acute appendicitis in acute diarrhoea - can be retrocaecal or pelvic appendicitis
- Missing faecal impaction with spurious diarrhoea
- Failing to perform a rectal examination
- Failing to consider acute ischaemic colitis in an elderly patient with the acute onset of bloody diarrhoea stools (following sudden abdominal pain in preceding 24 hours)

Seven masquerades checklist

The significant masquerades include diabetes, when an autonomic neuropathy may cause alternating bouts of constipation and diarrhoea, thyrotoxicosis and drugs. Drugs that can cause diarrhoea are summarised in Table 44.2.

Table 44.2 Drugs that can cause diarrhoea

Alcohol, esp. chronic abuse (often overlooked!)
Antibiotics, esp. penicillin derivatives
Antihypertensives, selected (e.g. methyldopa)
Arcarbose
Cardiac agents (e.g. digoxin, quinidine)
Chenodeoxycholic acid
Cisapride
Colchicine
Cytotoxic agents (e.g. methotrexate)
Heavy metals
H_2-receptor antagonists
Iron-containing compounds
Laxatives
Magnesium-containing antacids
Metformin
Misoprostol
NSAIDs
Prostaglandins
Quinidine
Salicylates
Statins
Theophylline
Thyroxine

Pseudomembranous colitis (antibioticassociated diarrhoea)

This colitis can be caused by the use of any antibiotic, especially clindamycin, lincomycin, ampicillin and the cephalosporins (an exception is vancomycin). It is usually due to an overgrowth of *C. difficile*, which produces a toxin that causes specific inflammatory lesions, sometimes with a pseudomembrane. It may occur, uncommonly, without antibiotic usage.

Features

- Profuse watery diarrhoea
- Abdominal cramping and tenesmus, may be fever
- Within 2 days of taking antibiotic (can start up to 4 to 6 weeks after usage)
- Persists 2 weeks (up to 6) after ceasing antibiotic

Diagnosed by characteristic lesions on sigmoidoscopy and a tissue culture assay for *C. difficile* toxin.

Treatment[3]

- Cease antibiotic
- Choice 1: metronidazole 400 mg (o) tds for 7 - 10 days *or*
- Choice 2: vancomycin 125 mg (o) qid for 7 - 10 days (in consultation with specialist)

Psychogenic considerations

Anxiety and stress can cause looseness of the bowel. The irritable bowel syndrome, which is a very common condition, may reflect underlying psychological factors and most patients find that the symptoms are exacerbated by stress. Look for evidence of depression.

In children chronic diarrhoea can occur with the so-called ‘maternal deprivation syndrome’, characterised by growth and developmental retardation due to adverse psychosocial factors.

THE CLINICAL APPROACH

History

As always, the history is the key to the diagnosis. First establish what the patient means by the term ‘diarrhoea’, his or her normal pattern and how the presenting problem varies from normal.

It is important to analyse the nature of the stools, the frequency of diarrhoea, associated symptoms, including abdominal pain, and constitutional symptoms, such as fever. Food intake in the past 72 hours and recent travel abroad may give a clue to acute gastroenteritis or food poisoning (an acute self-limiting illness of diarrhoea and vomiting). The difference between food poisoning and infective gastroenteritis is presented in Table 44.3. However, there can be an overlap of features from a specific organism and the exercise may be semantic, but it may provide a clue to foodborne causation. A summary of non-microbial food poisoning is presented in Table 44.4.

Table 44.3 Comparison of acute diarrhoea due to bacterial food poisoning and infective gastroenteritis

	Food poisoning	Infective gastroenteritis
Responsible organisms	**Toxins from:**	
	Staphylococcus aureus	Viral
	Salmonella sp.	Bacterial
	Clostridium perfringens	e.g. *Campylobacter jejuni Escherichia coli Shigella* sp. *Salmonella* sp.
	Vibrio parahaemolyticus	
	Aeromonas hydrophilia	
	Bacillus cereus	
Incubation period (onset from contact)	Short - within 24 hours	3 - 5 days
	Average - 12 hours	
	S. aureus - 2 - 4 hours	
Diarrhoea	Watery	Diarrhoea ± blood
Other features	Abdominal cramps (milder)	Abdominal cramps
	Dehydration	

	Headache Vomiting	
Typical foods	Chicken Meat Seafood Rice Custards and cream (*S. aureus*)	Milk Water Chicken

Table 44.4 Non-microbial food poisoning[4]

Food (specific types)	Toxin	Onset	Features (symptoms)
Mushrooms Toadstools	Muscarine	Minutes to hours	• N, V, D, P • multiple CNS symptoms
Immature or sprouting potatoes	Solanine	Within hours	• N, V, D, P • throat constriction
Fish	Ichthysarcotoxin Various, e.g. Ciguatera, scombrotoxin	10 - 60 minutes (occasionally longer)	• N, V, D, P • circumoral tingling • CNS symptoms • collapse
Mussels	Mytilotoxism	5 - 30 minutes	• N, V, P • CNS: paralysis
Grain, esp. rye	Ergot fungus	Minutes to 24 hours	• N, V, P • circulatory and CNS
Fava beans (favism)	Enzyme deficiency	Rapid	• V, D • acute haemolysis

N ° nausea; V ° vomiting; D ° diarrhoea; P ° abdominal pain

A drug history is relevant and a family history of diarrhoea, which may be significant for coeliac disease, Crohn's disease and cystic fibrosis.

Patients at risk from HIV infection should be discreetly evaluated.

Key questions

Acute diarrhoea

- Where did you eat in the 24 hours before the diarrhoea started?
- What food did you eat during this time?
- Did you have chicken or seafood recently? (Chicken may be contaminated with *Salmonella* or *Campylobacter* and seafood with *Vibrio parahaemolyticus*.)
- Did any other people get the same problem?
- Have you travelled overseas recently? Where?
- Have you noticed any blood or mucus in your motions?
- Have you had any previous attacks?
- Have you noticed fever, weakness or other symptoms?

Chronic diarrhoea

- Have you noticed any blood or mucus in the motion?
- Have you travelled overseas recently? Where?
- Do you have pain and is it relieved by opening your bowels or passing wind?

- Does anyone else in your family have diarrhoea?
- Have you had any operations on your abdomen recently?
- What medications are you taking?
- Are you taking antibiotics?
- Do you take vitamin C for your health?
- Do you take laxatives?
- How much alcohol do you drink?
- How much milk do you drink?
- What about thick shakes, ice-cream and yoghurt?
- Do you get clammy or shaky, or have you lost weight?
- Have you had trouble with pain in your joints, back pain, eye trouble or mouth ulceration?
- Do you have trouble flushing your motions down the toilet?
- Do you get diarrhoea during the night?
- Are you under a lot of stress?

Significance of symptoms

Abdominal pain

Central colicky abdominal pain indicates involvement of the small bowel, while lower abdominal pain points to the large bowel.

Nature of stools

If small volume consider inflammation or carcinoma of colon, and if large volume consider laxative abuse and malabsorption.

If there is profuse bright red bleeding consider diverticulitis or carcinoma of colon, and if small amounts with mucus or mucopus consider inflammatory bowel disorder. The presence of blood in the stools excludes functional bowel disorder. Diarrhoea at night suggests organic disease. In steatorrhoea the stools are distinctively pale, greasy, offensive, floating and difficult to flush. It is exacerbated by fatty foods.

The consistency of the stool as an aid to diagnosis[2,4] is summarised in Table 44.5, and the characteristics that distinguish between small and large bowel diarrhoea[1] are presented in Table 44.6.

Table 44.5 Stool consistency as an aid to diagnosis

Consistency	Probable cause
Liquid and uniform	Small bowel disorder (e.g. gastroenteritis)
Loose with bits of faeces	Colonic disorder
Watery, offensive, bubbly	*Giardia lamblia* infection
Liquid or semiformed, mucous ± blood	*Entamoeba histolytica*
Bulky, pale, offensive	Malabsorption
Pellets or ribbons	Irritable bowel syndrome

Table 44.6 Distinction between small and large bowel diarrhoea

	Small bowel	Large bowel
Volume	Large	Small
Pain	Central	Lower/LIF
Borborygmi	+ +	-
Undigested food	+	-
Steatorrhoea	+/-	-
Blood	-	+
Mucus	-	+
Urgency	-	+
Tenesmus	-	+

Physical examination

The extent of the examination depends on the nature of the presenting problem. If it is acute, profuse and associated with vomiting, especially in a child, the examination needs to be general to assess the effects of fluid, electrolyte and nutritional loss. An infant's life is in danger from severe gastroenteritis and this assessment is a priority. The general nutritional and electrolyte assessment is also relevant in chronic diarrhoea with malabsorption, and this includes looking for evidence of muscle weakness (e.g. hypokalaemia, hypomagnesaemia, tetany [hypocalcaemia], bruising [vitamin K loss]).

The examination should also focus on the abdomen (systematic palpation), the rectum and the skin. Possible helpful signs are included in Figure 44.2.


pigmentation of Addison's disease
postural hypotension of Addison's disease
liver disorder
• amoebiasis
• metastases
mass in RIF
carcinoma
Crohn's disease
pelvic abscess
skin changes of inflammatory bowel disorders
iritis—inflammatory bowel disorders
hyperthyroidism
cardiac disorder of carcinoid syndrome
splenomegaly (amyloid)
spondyloarthropathy of inflammatory bowel disorders
mass of carcinoma colon
tenderness of inflammation
warm moist palms and tremor of hyperthyroidism
peripheral neuropathy
• diabetes
• amyloidosis

Figure 44.2 Possible significant signs in the patient with diarrhoea

The stool

In all cases the stool should be examined. Note the presence of blood or mucus or steatorrhoea.

Investigations[4]

The following list includes a range of tests that may be required. Appropriate tests should be judiciously selected and in some instances, such as acute selflimiting diarrhoea, no investigations are necessary.

- Stool tests

  - microscopy for parasites and red and white cells (warm specimen for amoebiasis)
  - cultures: may need special requests for *Campylobacter* sp., *C. difficile* and toxin, *Yersinia* sp., *Cryptosporidium* sp., *Aeromonas* sp. (stools must be collected fresh on three occasions)
- Blood tests
  - haemoglobin; MCV, WCC, ESR, iron, ferritin, folate, vitamin B_{12}, calcium, electrolytes, thyroid function, HIV tests
- Antibody tests (e.g. IgA antiendomysial, IgA transglutaminase for coeliac disease)
- Haemagglutination tests for amoebiasis
- *C. difficile* tissue culture assay
- Malabsorption studies
- Endoscopy
  - proctosigmoidoscopy
  - flexible sigmoidoscopy/colonoscopy (with biopsy)
  - small bowel biopsy (coeliac disease)
- Radiology
  - plain X-ray abdomen - of limited value
  - small bowel enema
  - barium enema, especially double contrast

Note: HIV patients should be investigated in specialist centres.

Principles of treatment[5]

When an underlying cause of diarrhoea can be identified, apart from some common infections, management should be directed at that cause. There are a few situations in which the causative bacterial or parasitic pathogen requires specific treatment, for example, giardiasis. The management is determined by the nature of the pathogen and the severity of the illness.

However, in Australia most infective cases are viral. The basic principle therefore is to achieve and maintain adequate hydration until the illness resolves. In adults and children oral rehydration is indicated unless there is evidence of impending circulatory 'shock' demanding intravenous therapy. Oral rehydration solution containing sodium, potassium and glucose should be considered for patients with mild to moderate dehydration. Adults should drink 2 to 3 L of the solution in 24 hours.

In general, treatment should not be directed specifically at altering the frequency and consistency of the stools. The antimotility drugs (loperamide, diphenoxylate and codeine) have a role restricted to short-term control of symptoms in adults during periods of significant social inconvenience, such as travel. It must be emphasised that antimotility drugs are never indicated for management of acute diarrhoea in infants and children.[5]

The traditional absorbent agents, such as kaolin/pectin mixtures, activated charcoal and other mineral clays, have not been shown to be of value and may interfere with absorption of other drugs. They should not be used.

Specific antibiotics are reserved for the treatment of giardiasis, amoebiasis, antibiotic-associated diarrhoea, cholera and typhoid. Although antibiotics are usually unnecessary in most cases they may be indicated for severe cases of *Campylobacter* enteritis, *Salmonella* enteritis, shigellosis and traveller's diarrhoea.

MALABSORPTION

It is important to distinguish the steatorrhoea of various malabsorption syndromes from diarrhoea. Important causes are presented in Table 44.7.

Table 44.7 Important causes of malabsorption

Primary mucosal disorders
• gluten-sensitive enteropathy (coeliac disease)
• tropical sprue
• lactose intolerance (lactase deficiency)
• Crohn's disease (regional enteritis)
• Whipple's disease
• parasite infections (e.g. *Giardia lamblia*)

• lymphoma
Maldigestion states Lumenal abnormalities • postsurgery (e.g. gastrectomy, ileal resection) • systemic sclerosis
Pancreatic disorders • chronic pancreatitis • cystic fibrosis • pancreatic tumours (e.g. Zollinger-Ellison)

The common causes are coeliac disease, chronic pancreatitis and postgastrectomy.

Clinical features

- Bulky, pale, offensive, frothy, greasy stools
- Stools difficult to flush down toilet
- Weight loss
- Prominent abdomen
- Failure to thrive (in infants)
- Increased faecal fat
- Signs of multiple vitamin deficiencies (e.g. A, D, E, K)
- Sore tongue (glossitis)
- Hypochromic or megaloblastic anaemia (possible)

Refer for specific investigations (e.g. FBE, barium studies, small bowel biopsy, faecal fat [>21 g/3 days]).

Coeliac disease[5]

Synonyms: coeliac sprue, gluten-sensitive enteropathy

Note: It can appear at *any* age; refer to coeliac disease in children (p. 477).

It is widely underdiagnosed because most patients present with non-GIT symptoms, such as tiredness.

There is a genetic factor in this autoimmune disorder with a one in 10 chance if a first degree relative is affected. Consider screening under 2 years if there is such an association.

Diagnosis

- Classic triad: diarrhoea, weight loss, iron/folate deficiency
- Elevated faecal fat
- Characteristic duodenal biopsy (key test)
- IgA antigliadin antibodies (screening - limited)
- IgA antiendomysial antibodies (> 90% sensitivity and specificity)
- IgA transglutaminase antibodies (> 90% sensitivity and specificity)

Associations

- Type 1 diabetes
- Dermatitis herpatiformis
- IgA deficiency
- Autoimmune thyroid disease
- Osteoporosis

Management

- Diet control: high complex carbohydrate and protein, low fat, gluten-free (no wheat, barley, rye and oats)

- Treat specific vitamin deficiencies
- Give pneumococcal vaccination

Gluten-free diet

Avoid foods containing gluten either as an obvious component (e.g. flour, bread, oatmeal), or as a hidden ingredient (e.g. dessert mix, stock cube).

Forbidden foods include:

- standard bread, pasta, crispbreads, flour
- standard biscuits and cakes
- breakfast cereals made with wheat or oats
- oatmeal, wheat bran, barley/barley water
- ‘battered’ or bread-crumbed fish, etc.
- meat and fruit pies
- most stock cubes and gravy mixes

Whipple's disease

This is a rare malabsorption disorder usually affecting white males. It is caused by the bacillus *Tropheryma whippelii*. It may involve the heart, lungs and CNS.

Features

- Males > 40 years
- Chronic diarrhoea (steatorrhoea)
- Arthralgia
- Weight loss
- Lymphadenopathy
- ± Fever

Diagnosis

- PCR for *T. whippelii*
- Jejunal biopsy

Treatment

Co-trimoxazole or tetracycline for up to 12 months. This produces a dramatic improvement.

DIARRHOEA IN THE ELDERLY

The older the patient the more likely a late onset of symptoms that reflect serious underlying organic disease, especially malignancy. Colorectal cancer needs special consideration. The older the patient, especially the bedridden elderly patient, the more likely the presentation of faecal impaction with spurious diarrhoea. The possibility of drug interactions, including digoxin, should also be considered. Ischaemic colitis must be considered in an elderly patient.

Ischaemic colitis

This is due to atheromatous occlusion of mesenteric vessels.

Clinical features

Clinical features include:

- sharp abdominal pain in an elderly patient with bloody diarrhoea *or*
- periumbilical pain and diarrhoea about 15 - 30 minutes after eating
- maybe loud bruits over central abdomen
- other evidence of generalised atherosclerosis
- barium enema shows ‘thumb printing’ sign due to submucosal oedema
- the definitive test is aortography and selective angiography of mesenteric vessels

- most episodes resolve - may be followed by a stricture

DIARRHOEA IN CHILDREN

The commonest cause of diarrhoea in children is acute infective gastroenteritis, but there are certain conditions that develop in childhood and infancy and require special attention. The presentation of small amounts of redcurrant jelly-like stool with intussusception should also be kept in mind. Of the many causes only a few are commonly seen. The two commonest causes are infective gastroenteritis and antibiotic-induced diarrhoea.

Important causes of diarrhoea in children are:

- infective gastroenteritis
- antibiotics
- overfeeding (loose stools in newborn)
- dietary indiscretions
- sugar (carbohydrate) intolerance
- food allergies (e.g. milk, soya bean, wheat, eggs)
- maternal deprivation
- malabsorption states
  - cystic fibrosis
  - coeliac disease

Note: Exclude surgical emergencies (e.g. acute appendicitis), infections (e.g. pneumonia), septicaemia, otitis media < 5 years.

Acute gastroenteritis

Note: Dehydration from gastroenteritis is an important cause of death, particularly in obese infants (especially if vomiting accompanies the diarrhoea).

Definition

It is an illness of acute onset, of less than 10 days duration associated with fever, diarrhoea and/or vomiting, where there is no other evident cause for the symptoms.[6]

Features

Causes:

- mainly rotavirus (developed countries) and adenovirus: viruses account for about 80%
- bacterial: *C. jejuni* and *Salmonella* sp. (two commonest), *E. coli* and *Shigella* sp.
- protozoal: *G. lamblia*, *E. histolytica*, *Cryptosporidium*
- food poisoning - staphylococcal toxin

Differential diagnoses. These include septicaemia, urinary tract infection, intussusception, appendicitis, pelvic abscess, partial bowel obstruction, diabetes mellitus and antibiotic reaction[4] (see Table 44.8).

Table 44.8 Differential diagnosis of acute diarrhoea and vomiting in children

Bowel infection
• viruses
• bacteria
• protozoal
• food poisoning - staphylococcal toxin
Systemic infection
Abdominal disorders
• appendicitis
• pelvic abscess
• intussusception

• malrotation
Urinary tract infection
Antibiotic reaction
Diabetes mellitus

Note: Exclude acute appendicitis and intussusception in the very young.

Symptoms

- Anorexia, nausea, poor feeding, vomiting, fever, diarrhoea (fever and vomiting may be absent)
- Fluid stools (often watery) 10 - 20 per day
- Crying - due to pain, hunger, thirst or nausea
- Bleeding - uncommon (usually bacterial)
- Anal soreness

Viral indication: large volume, watery, typically lasts 2 - 3 days, systemic symptoms uncommon

Bacterial indication: small motions, blood, mucus, abdominal pain and tenesmus

Dehydration must be assessed (see Table 44.9).

Table 44.9 Assessment of hydration

	Mild	Moderate	Severe
Body weight loss	• 4 - 5%	• 6 - 9%	• ° ≥ 10%
Symptoms/general observations	• Thirsty • Alert • Restless	• Thirsty • Restless • Lethargic • Irritable	• *Infants*: drowsy, limp, cold, sweaty, cyanotic limbs, comatose • *Older*: apprehensive, cold and sweaty, cyanotic limbs
Signs	• Normal	• Dry mucous membranes, absent tears	• Rapid feeble pulse • Hypotensive • Sunken eyes and fontanelles • Very dry mucous membranes
Pinched skin test	• Normal	• Retracts slowly (1 - 2 seconds)	• Retracts very slowly > 2 seconds
Urine output	• Normal	• Decreased	• Nil
Treatment	• *Oral rehydration* - small amounts of fluids often - continue breast-feeding - solids after 24 hours - provide maintenance fluid and loss	• *Oral rehydration* - consider nasogastric tube for steady fluid infusion or - IV infusion	• Urgent IV infusion: isotonic fluid

Complications:

- febrile convulsions
- sugar (lactose) intolerance (common)
- septicaemia, especially *Salmonella*

Management

Management is based on the assessment and correction of fluid and electrolyte loss.[6,7] Since dehydration is usually isotonic with equivalent loss of fluid and electrolytes, serum electrolytes will be normal.

Note: The most accurate way to monitor dehydration is to weigh the child, preferably without clothes, on the same scale each time. However, the easiest is clinical assessment (e.g. vomiting, no urine, lethargy and thirst).

Avoid

- Drugs: antidiarrhoeals, antiemetics and antibiotics
- Lemonade: osmotic load too high, can use if diluted 1 part to 4 parts water but sugar may be poorly tolerated

To treat or not to treat at home

- Treat at home - if family can cope, vomiting is not a problem and no dehydration.
- Admit to hospital - if dehydration or persisting vomiting or family cannot cope; also infants < 6 months and high-risk patients.

Advice to parents (for mild-to-moderate diarrhoea)

If applicable, remove child from daycare or school. Advise about hygiene, including handwashing and napkin disposal.

General rules

- Give small amounts of fluids often
- Start solids after 24 hours
- Continue breastfeeding (may be increased in frequency) *or*
- Continue formula feeding if tolerated or resume it after 24 hours.
- Consider stool culture and test for rotavirus for symptoms that persist and worsen

Day 1

Give fluids, a little at a time and often (e.g. 50 mL every 15 minutes if vomiting a lot). A good method is to give 200 mL (about 1 cup) of fluid every time a watery stool is passed or a big vomit occurs.

The ideal fluid is Gastrolyte or New Repalyte. Other suitable oral rehydration preparations are WHO recommended solutions Electrolade and Glucolyte. A new product is Hydralyte paediatric rehydration, which is a solution as an ‘ice-block’ formulation:

- children 1 - 2 years - one ‘ice-block’ per hour
- older children - as often as desired.

Alternatives are:

• lemonade (not low-calorie)	1 part to 6 parts water
• sucrose (table sugar)	1 teaspoon to 120 mL water
• glucose	1 teaspoon to 120 mL water
• cordials (not low-calorie)	1 part to 16 parts water
• fruit juice	1 part to 4 parts water

Warning: Do *not* use straight lemonade or mix up Gastrolyte with lemonade or fluids other than water.

Method of assessing fluid requirements[5]

- Fluid loss (mL) ° % dehydration × body weight (kg) × 10
- Maintenance (mL/kg/24 h): 1 - 3 mo: 120 mL; 4 - 12 mo: 100 mL; > 12 mo: 80 mL
- Allow for continuing loss.

Example: 8 month 10 kg child with 5% dehydration:

Fluid loss ° 5 × 10 × 10 ° 500 mL

Maintenance ° 100 × 10 ° 1000 mL

Total 24 hour requirement (min.) ° 1500 mL

Approximate average hourly requirement ° 60 mL

- Aim to give more (replace fluid loss) in the first 6 hours.
- Rule of thumb: 100 mL/kg (infants) and 50 mL/kg (older children) in first 6 hours.

Days 2 and 3

Reintroduce your baby's milk or formula diluted to half strength (i.e. mix equal quantities of milk or formula and water).

Do not worry that your child is not eating food. Solids can be commenced after 24 hours. Start with bread, plain biscuits, jelly, stewed apple, rice, porridge or non-fat potato chips. Avoid fatty foods, fried foods, raw vegetables and fruit, and wholegrain bread.

Day 4

Increase milk to normal strength and gradually reintroduce the usual diet.

Breastfeeding. If your baby is not vomiting, continue breastfeeding but offer extra fluids (preferably Gastrolyte) between feeds. If vomiting is a problem, express breast milk for the time being while you follow the oral fluid program.

Note: Watch for lactose intolerance as a sequela - explosive diarrhoea after introducing formula. Replace with a lactose-free formula.

It acute invasive or persistent *Salmonella* are present, give antibiotics (ciprofloxacin or azithromycin).

CHRONIC DIARRHOEA IN CHILDREN

Sugar intolerance

Synonyms: carbohydrate intolerance, lactose intolerance.

The commonest offending sugar is lactose.

Diarrhoea often follows acute gastroenteritis when milk is reintroduced into the diet (some recommend waiting for 2 weeks). Stools may be watery, frothy, smell like vinegar and tend to excoriate the buttocks. They contain sugar.

A simple test

- Line the napkin with thin plastic and collect fluid stool.
- Mix 5 drops of liquid stool with 10 drops of water and add a Clinitest tablet (detects lactose and glucose but not sucrose).
- A positive result indicates sugar intolerance.

Treatment

- Remove the offending sugar from the diet.
- Use milk preparations in which the lactose has been split to glucose and galactose by enzymes, or use soya protein.

Note: Most milk allergies improve with age.

Cows milk protein intolerance[8]

This is not as common as lactose intolerance. Diarrhoea is related to taking a cows milk formula and relieved when it is withdrawn.

Allergic responses to cows milk protein may result in a rapid or delayed onset of symptoms. Delayed onset may be more difficult to diagnose, presenting with diarrhoea, malabsorption or failure to thrive.

It is diagnosed by unequivocal reproducible reactions to elimination and challenge. If diagnosed, remove cows milk from the diet and replace with either soy milk, a hydrolysed or an elemental formula.

Inflammatory bowel disorder

These disorders, which include Crohn's disease and ulcerative colitis, can occur in childhood. A high index of suspicion is necessary to make an early diagnosis. Approximately 5% of cases of chronic ulcerative colitis have their onset in childhood.[6]

Chronic enteric infection

Responsible organisms include *Salmonella* sp., *Campylobacter*, *Yersinia*, *G. lamblia* and *E. histolytica*. With persistent diarrhoea it is important to obtain microscopy of faeces and aerobic and anaerobic stool cultures. *G. lamblia* infestation is not an uncommon finding and may be associated with malabsorption, especially of carbohydrate and fat. Giardiasis can mimic coeliac disease.

Coeliac disease

Clinical features in childhood:

- usually presents at 9 - 18 months, but any age
- previously thriving infant
- anorexia, lethargy, irritability
- failure to thrive
- malabsorption - abdominal distension
- offensive frequent stools

Diagnosis: duodenal biopsy

Treatment: remove gluten from diet

Cystic fibrosis

Cystic fibrosis which presents in infancy is the commonest of all inherited disorders (1 per 2500 live births). Refer to p. 170.

ACUTE GASTROENTERITIS IN ADULTS

Features

- Invariably a self-limiting problem (1 - 3 days)
- Abdominal cramps
- Possible constitutional symptoms (e.g. fever, malaise, nausea, vomiting)
- Other meal sharers affected → food poisoning
- Consider dehydration, especially in the elderly
- Consider possibility of enteric fever

Traveller's diarrhoea

The symptoms are usually as above but very severe diarrhoea, especially if associated with blood or mucus, may be a feature of a more serious bowel infection such as amoebiasis. Possible causes of diarrhoeal illness are presented in Table 14.1, pp. 114 - 5. Most traveller's diarrhoea is caused by *E. coli*, which produces a watery diarrhoea within 14 days of arrival in a foreign country. For specific treatment refer to pages 114.

Persistent traveller's diarrhoea

Any traveller with persistent diarrhoea after visiting less developed countries, especially India and China, may have a protozoal infection such as amoebiasis or giardiasis.

If there is a fever and blood or mucus in the stools, suspect amoebiasis. Giardiasis is characterised by abdominal cramps, flatulence and bubbly foulsmelling diarrhoea.

Principles of treatment

Acute diarrhoea

- Maintenance of hydration
- Antiemetic injection (for severe vomiting) prochlorperazine IM, statim *or* metoclopramide IV, statim
- Antidiarrhoeal preparations (avoid if possible: loperamide preferred) loperamide (Imodium) 2 caps statim then 1 after each unformed stool (max: 8 caps/day) *or* diphenoxylate with atropine (Lomotil) 2 tabs statim then 1 - 2 (o) 8 hourly

General advice to patient

Rest

Your bowel needs a rest and so do you. It is best to reduce your normal activities until the diarrhoea has stopped.

Diet

It is vital that you starve but drink small amounts of clear fluids such as water, tea, lemonade and yeast extract (e.g. Marmite) until the diarrhoea settles. Then eat low-fat foods such as stewed apples, rice (boiled in water), soups, poultry, boiled potatoes, mashed vegetables, dry toast or bread, biscuits, most canned fruits, jam, honey, jelly, dried skim milk or condensed milk (reconstituted with water).

Avoid alcohol, coffee, strong tea, fatty foods, fried foods, spicy foods, raw vegetables, raw fruit (especially with hard skins), Chinese food, wholegrain cereals and cigarette smoking.

On the third day introduce dairy produce, such as a small amount of milk in tea or coffee and a little butter or margarine on toast. Add also lean meat and fish (either grilled or steamed).

Antimicrobial drugs[3,5]

It is advisable not to use these except where the following specific organisms are identified. The drugs should be selected initially from the list below or modified according to the results of culture and sensitivity tests.[3] Only treat if symptoms have persisted for more than 48 hours. Adult doses are shown.

Shigella dysentery (moderate to severe)

- cotrimoxazole (double strength) 1 tab (o) 12 hourly for 5 days: use in children (children's doses) *or*
- norfloxacin 400 mg (o) 12 hourly for 5 days (preferred for adults) *or*
- ampicillin 1 g (o) 6 hourly for 5 days

Campylobacter jejuni (if prolonged)

- norfloxacin 400 mg (o) 12 hourly for 5 days (adults) *or*
- erythromycin 500 mg (o) qid for 7 days (preferable) in adults and children

Giardiasis

- tinidazole 2 g (o), single dose *or*
- metronidazole 400 mg (o) tds for 7 days

(in children: 30 mg/kg/day [to max. 1.2 g/day] as single daily dose for 3 days)

Salmonella enteritis

Antibiotics are not generally advisable but if severe or prolonged use:

- ciprofloxacin 500 mg (o) bd for 5 - 7 days or azithromycin for 7 days

Note: Salmonella is a notifiable disease; infants under 15 months are at risk of invasive *Salmonella* infection.

Amoebiasis (intestinal)

- metronidazole 600 - 800 mg (o) tds for 6 - 10 days plus
- diloxanide furoate 500 mg (o) tds for 10 days

Blastocystitis hominus

Pathogenicity is disputed: give therapy only if severe. Associated with poor hygiene (travel, pets, dam/tank water, oysters).

- metronidazole for 7 days

Specialist advice should be sought.

Special enteric infections

Typhoid/paratyphoid fever

- ciprofloxacin 500 mg (o) 12 hourly for 7 - 10 days (use IV if oral therapy not tolerated) If ciprofloxacin is contraindicated (e.g. in children) or not tolerated, then use:
- ceft riaxone 3 g IV daily until culture and sensitivities available, then choose oral regimens

Alternative oral regimens (based on sensitivity)

- chloramphenicol 500 - 750 mg (o) 6 hourly for 14 days *or*
- cotrimoxazole (DS) 1 tablet (o) 12 hourly for 14 days *or*
- amoxycillin 1 g (o) 6 hourly for 14 days

If severe: administer same drug and dosage IV for first 4 - 5 days.

Cholera

Antibiotic therapy reduces the volume and duration of diarrhoea. Rehydration is the key.

- doxycycline 100 mg (o) 12 hourly for 3 days

For pregnant women and children:

- amoxycillin (child: 10mg/kg up to) 250 mg (o) 6 hourly for 4 days

INFLAMMATORY BOWEL DISEASE

Inflammatory bowel disease should be considered when a young person presents with:

- bloody diarrhoea and mucus
- colonic pain and fever
- constitutional symptoms including weight loss and malaise
- extra-abdominal manifestations such as arthralgia, low back pain (spondyloarthropathy), eye problems (iridocyclitis), liver disease and skin lesions (pyoderma granulosum, erythema nodosum)

Two important disorders are ulcerative colitis (UC) and Crohn's disease which have equal sex incidence and can occur at any age, but onset peaks between 20 and 40 years.

Ulcerative colitis

Features

- Mainly a disease of Western societies
- Mainly in young adults (15 - 40 years)
- High-risk factors - family history, previous attacks, low-fibre diet
- Recurrent attacks of loose stools
- Blood, or blood and pus, or mucus in stools
- Abdominal pain slight or absent
- Fever, malaise and weight loss uncommon
- Begins in rectum (continues proximally) - affects only the colon
- An increased risk of carcinoma after 7 - 10 years

Main symptom

- Bloody diarrhoea

Diagnosis

- Proctosigmoidoscopy: a granular red proctitis with contact bleeding
- Barium enema: characteristic changes

Prognosis

- 5% mortality in an acute attack

- Recurrent attacks common

Crohn's disease

Synonyms: regional enteritis, granulomatous colitis

Features

- Recurrent diarrhoea in a young person (20 - 40 years)
- Blood and mucus in stools (less than ulcerative colitis)
- Colicky abdominal pain (small bowel colic)
- Right iliac fossa pain (confused with appendicitis)
- Constitutional symptoms (e.g. fever, weight loss, malaise)
- Signs include perianal disorders (e.g. anal fissure, fistula, ischiorectal abscess), mouth ulcers
- Skip areas in bowel: 1/2 ileocolic, 1/4 confined to small bowel, 1/4 confined to colon

Main symptom

- Colicky abdominal pain

Diagnosis

- Sigmoidoscopy: ‘cobblestone’ appearance (patchy mucosal oedema)
- Colonoscopy: useful to differentiate from UC

Prognosis

Less favourable than UC with both medical and surgical treatment.

Management principles

- Treat under consultant supervision.
- Treatment of acute attacks depends on severity of the attack and the extent of the disorder.
  - mild attacks: manage out of hospital
  - severe attacks: hospital, to attend to fluid and electrolyte balance
- Role of diet controversial: consider a high-fibre diet but maintain adequate nutrition.
- Pharmaceutical agents (the following can be considered):
  - 5-aminosalicylic acid derivatives (mainly UC)
    - sulfasalazine (mainstay)
    - olsalazine; mesalazine
  - corticosteroids (mainly for acute flares):
    - oral
    - parenteral
    - topical (rectal foam, suppositories or enemas)
  - immunomodifying drugs (e.g. azathioprine, cyclosporin, methotrexate)
- Surgical treatment: reserve for complications.

ALTERNATING DIARRHOEA AND CONSTIPATION

Alternating diarrhoea and constipation are wellknown symptoms of incomplete bowel obstruction (carcinoma of colon and diverticular disease) and irritable bowel syndrome.

Irritable bowel syndrome (IBS)[5,9,10]

Clinical features

- Typically in younger women (21 - 40 years)
- Any age or sex can be affected
- May follow attack of gastroenteritis/traveller's diarrhoea

- Cramping abdominal pain (central or iliac fossa) - Figure 44.3



Figure 44.3 Classic symptoms of irritable bowel syndrome

- Pain usually relieved by passing flatus or by defecation
- Variable bowel habit (constipation more common)
- Diarrhoea usually worse in morning - several loose, explosive bowel actions with urgency
- Often precipitated by eating
- Faeces sometimes like small, hard pellets or ribbon-like
- Anorexia and nausea (sometimes)
- Bloating, abdominal distension, ↑ borborygmi
- Tiredness common

The Rome II diagnostic criteria for IBS is presented in Table 44.10.

Table 44.10 Rome II diagnostic criteria for irritable bowel syndrome[*10]

In the preceding 12 months, the patient has had at least 12 weeks (not necessarily consecutively) of abdominal discomfort or pain with two of the following three features: • relieved by defaecation and/or • onset associated with a change in stool frequency and/or • onset associated with a change in form (appearance) of stool
Symptoms that cumulatively support the diagnosis of irritable bowel syndrome

- Abnormal stool frequency (for research purposes may be defined as more than three bowel movements per day and less than three bowel movements per week)
- Abnormal stool form (lumpy/hard or watery/mushy)
- Abnormal stool passage (straining, urgency or feeling of incomplete evacuation)
- Passage of mucus
- Bloating or feeling of abdominal distension

*in absence of structural or metabolic abnormalities to explain symptoms

IBS is a diagnosis of exclusion. A thorough physical examination, investigations (FBE, ESR and stool microscopy or culture) and sigmoidoscopy are necessary. Insufflation of air at sigmoidoscopy may reproduce the abdominal pain of IBS.

Possible related causes

Bowel infection, food irritation (e.g. spicy foods), lactose (milk) intolerance, low-fibre diet, high fatty foods, laxative overusage, use of antibiotics and codeine-containing analgesics, psychological factors.

Management

The patient must be reassured and educated with advice that the problem will not cause malignancy or inflammatory bowel disease and will not shorten life expectancy. The basis of initial treatment is simple dietary modification and non-drug therapy. Drugs that have been used with mixed success include mebeverine, dicyclomine and peppermint oil. The new serotonin 5-H5 agonist, tegaserol (Zelmac) is promoted for use in women with IBS. Exacerbations of diarrhoea can be treated with loperamide 2 mg (o) one to three times daily.

Self-help advice to the patient

Anyone with IBS should try to work on the things that make the symptoms worse. If you recognise stresses and strains in your life, try to develop a more relaxed lifestyle. You may have to be less of a perfectionist in your approach to life.

Try to avoid any foods that you can identify as causing the problem. You may have to cut out smoking and alcohol and avoid laxatives and codeine (in painkillers). A high-fibre diet and 1.5 - 2 L of water a day may be the answer to your problem.

Diverticular disorder

Diverticular disorder is a problem of the colon (90% in descending colon) and is related to lack of fibre in the diet. It is usually symptomless.

Clinical features

- Typical in middle aged or elderly - over 40 years
- Increases with age
- Present in one in three people over 60 years (Western world)
- Diverticulosis - symptomless
- Diverticulitis - infected diverticula and symptomatic (Refer p. 330)
- Constipation or alternating constipation/diarrhoea
- Intermittent cramping lower abdominal pain in LIF
- Tenderness in LIF
- Rectal bleeding - may be profuse (± faeces)
- May present as acute abdomen or subacute obstruction
- Usually settles in 2 - 3 days

Complications (of diverticulitis)

- Abscess
- Perforation
- Peritonitis
- Obstruction (refer p. 331)
- Fistula - bladder, vagina

Investigations

- WBC and ESR - to determine inflammation
- Sigmoidoscopy
- Barium enema

Management

- It usually responds to a high-fibre diet.
- Avoidance of constipation.

Advice to the patient

The gradual introduction of fibre with plenty of fluids (especially water) will improve any symptoms you may have and reduce the risk of complications. Your diet should include:

1. cereals, such as bran, shredded wheat, muesli or porridge
2. wholemeal and multigrain breads
3. fresh or stewed fruits and vegetables

Bran can be added to your cereal or stewed fruit, starting with 1 tablespoon and gradually increasing to 3 tablespoons a day. Fibre can make you feel uncomfortable for the first few weeks, but the bowel soon settles to your improved diet.

WHEN TO REFER

Children with diarrhoea

- Infant under 3 months
- Moderate to severe dehydration
- Diagnosis of diarrhoea and vomiting in doubt (e.g. blood in vomitus or stool, bilestained vomiting, high fever or toxaemia, abdominal signs suggestive of appendicitis or obstruction)
- Failure to improve or deterioration
- A pre-existing chronic illness

Adults

- Patient with chronic or bloody diarrhoea
- Any problem requiring colonoscopic investigation
- Patients with anaemia
- Patients with weight loss, abdominal mass or suspicion of neoplasia
- Patients with anal fistulae
- Patients not responding to treatment for giardiasis
- Infection with *E. histolytica*
- Long-term asymptomatic carrier of typhoid or paratyphoid fever
- Patient with persistent undiagnosed nocturnal diarrhoea
- Patients with IBS with a significant change in symptoms
- Patients with inflammatory bowel diseases with severe exacerbations, possibly requiring immunosuppressive therapy and with complications
- Patients with ulcerative colitis of more than 7 years duration (screening by colonoscopy for carcinoma)

Practice tips

- Oral antidiarrhoeal drugs are contraindicated in children; besides being ineffective they may prolong intestinal recovery.
- Antiemetics can readily provoke dystonic reactions in children, especially if young and dehydrated.
- Acute diarrhoea is invariably self-limiting (lasts 2 - 5 days). If it lasts longer than 7 days, investigate with culture and microscopy of the stools.
- If diarrhoea is associated with episodes of facial flushing or wheezing, consider carcinoid syndrome.
- Recurrent pain in the right hypochondrium is usually a feature of the IBS (not gall bladder disease).
- Recurrent pain in the right iliac fossa is more likely to be IBS than appendicitis.

- Beware of false correlations or premature conclusions (e.g. attributing the finding of diverticular disorder on barium meal to the cause of the symptoms).
- Undercooked chicken is a common source of enteropathic bacterial infection.
- Consider alcohol abuse if a patient's diarrhoea resolves spontaneously on hospital admission.

REFERENCES

1. Bolin T, Riordan SM. Acute and persistent diarrhoea. Current therapeutics, 2001, May: 47 - 57.
2. Sandler G, Fry J. *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986: 25 - 30.
3. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 79 - 89.
4. Dalton C. How to treat: Foodborne illness. Australian Doctor, 2005; 15 April: 39 - 46.
5. Birkett D (Chair). *Therapeutic Guidelines: Gastrointestinal* (version 3). Melbourne: Therapeutic Guidelines Ltd, 2003.
6. Robinson MJ, Roberton DM. Practical paediatrics (5th edn). Edinburgh: Churchill Livingstone; 2003: 675 - 90.
7. Oberklaid F. Management of gastroenteritis in children. In: *The Australian Paediatric Review*. Melbourne: Royal Children's Hospital, 1990: 1 - 2.
8. Smart J. *Paediatric Handbook* (6th edn). Melbourne: Blackwell Science, 2000: 334 - 5.
9. Schmidt G. Irritable bowel syndrome. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 279 - 80.
10. Thompson WG, Longstreth G, Drossman DA, Heaton K, Irvine EJ, Muller-Lissner S. Functional bowel disorders and functional abdominal pain. In: Drossman DA, Corazziari E, Talley J, Thompson WG, Whitehead WE eds, *Rome II. The Functional Gastrointestinal Disorders. Diagnosis, Pathophysiology and Treatment: a Multinational Consensus* (2nd edn). USA: Degnon Associates, 2000: 360.

CHAPTER 45. THE DISTURBED PATIENT

There is not a sight in nature so mortifying as that of a Distracted Person, when his imagination is troubled, and his whole soul is disordered and confused.

JOSEPH ADDISON (1672 - 1719)

THE DISTURBED PATIENT

The disturbed and confused patient is a complex management problem ingeneral practice. The cause may be a single one or a combination of several abnormal mental states (see Table 45.1).[1] The cause may be an organic mental disorder, which may be a long-term insidious problem such as dementia, or an acute disorder (delirium), often dramatic in onset. Onthe other hand, the causeof the disturbance may be a psychiatric disorder such as panic disorder, mania, major depression or schizophrenia.

Table 45.1 A general classification of psychiatric disorders[1]

Organic mental disorders [1] • Acute organic brain syndrome (delirium) • Chronic organic brain syndrome (dementia)
Psychoactive and substance use disorders • Toxic states • Drug dependency • Withdrawal states
Schizophrenic disorders
Mood disorders • Major depression • Bipolar (manic depressive) disorder • Adjustment disorders with depressed mood • Dysthymia
Anxiety disorders • Generalised anxiety disorder • Panic disorder • Obsessive - compulsive disorder • Phobic disorders • Post-traumatic stress disorder
Disorders specific to children
Other disorders • Postpartum psychiatric illness • Eating disorders • Personality disorders • Body dysmorphic disorder

The manifestations of the disturbance are many and include perceptualchanges and hallucinations, disorientation, changes in consciousness, changes in mood from abnormally elevated to gross depression, agitation and disturbed thinking, including delusions.

KEY FACTS AND CHECKPOINTS

- Depression affects 15% of people over 65 and can mimic or complicate any other illness, including delirium and dementia.[1]
- Elderly patients with depression are at a high risk of suicide.
- Always search vigorously for the cause or causes of delirium.
- Seeing patients in their home is the best way to evaluate their problem and support systems. It allows opportunities for a history from close contacts and for checking medication, alcohol intake and other factors.

- The diagnosis of dementia can be overlooked: a Scottish study showed that 80% of demented patients were not diagnosed by their GP.[2]
- Patients with a chronic brain syndrome (dementia) are at special risk of an acute brain syndrome (delirium) in the presence of infections and many prescribed drugs.[1]
- Consider prescribed and illicit substances, including the severe anticholinergic delirium syndrome.
- The key feature of dementia is impaired memory.
- The two key features of delirium are disorganised thought and attention.

Definitions

Delirium

(also termed 'toxic confusional state') A relatively acute disorder in which impaired consciousness is associated with abnormalities of perception or mood.

Confusion

Disorientation in time, place and person. It may be accompanied by a disturbed conscious state (Table 75.1, p. 817).

Dementia

An acquired, chronic and gradually progressive deterioration of memory, intellect and personality. Presenile dementia or early onset dementia is dementia under 65 years of age. Senile dementia refers to older patients (usually over 80 years).

Alzheimer's disease

A term used for both senile and presenile dementia which has characteristic pathological degenerative changes in the brain.

Cognition

The mental functions of perception, thinking and memory. It is the process of 'knowing'.

Hallucinations

Disorders of perception quite divorced from reality. Features:

- mostly auditory or visual
- a false perception - not a distortion
- perceived as normal perceptions
- independent of the person's will

Illusions

False interpretations of sensory stimuli such as mistaking people or familiar things.

Delusions

Abnormal, illogical or false beliefs that are held with absolute conviction despite evidence tothe contrary.

Obsessions

Recurrent or persistent thoughts, images or impulses that enter the mind despite efforts to exclude them.

Compulsions

Repeated, stereotyped and seemingly purposeful actions that the person feels compelled to carryout but resists, realising they are irrational (most are associated with obsessions).

Somatisation

The conversion of mental experiences or states into bodily symptoms, with no physical causation.

Dysmorphophobia

The belief that one has a significant deformity or the dread of such a deformity.

Conversion

The process by which thoughts or experiences unacceptable to the mindare repressed and converted into physical symptoms.

Dissociation

A psychological disorder in which unpleasant memories or emotions aresplit off from consciousness and the personality and buried into the unconsciousness.

A DIAGNOSTIC APPROACH

A summary of the diagnostic strategy model for the disturbed or confused patient is presented in Table 45.2.

Table 45.2 The disturbed mind: diagnostic strategy model

Q. Probability diagnosis A. The 4 D's • dementia • delirium (look for cause) • depression • drugs - toxicity - withdrawal
Q. Serious disorders not to be missed A. Cardiovascular • CVAs • cardiac failure • arrhythmia • acute coronary syndromes Neoplasia • cerebral • carcinoma (e.g. lung) Severe infections • septicaemia • HIV infection • infective endocarditis Hypoglycaemia Bipolar disorder/mania Schizophrenia states Anxiety/panic Subdural haematoma
Q. Pitfalls (often missed) A. Illicit drug withdrawal Fluid and electrolyte disturbances Faecal impaction (elderly) Urinary retention (elderly) Hypoxia Pain syndromes (elderly) *Rarities* • Hypocalcaemia • Kidney failure • Hepatic failure • Prion diseases (e.g. Creutzfeldt-Jakob disease)
Q. Seven masquerades checklist A. Depression ✔ ✔ Diabetes ✔ Drugs ✔ ✔ Anaemia ✔ Thyroid disorder ✔

Spinal dysfunction ✔ (severe pain in elderly) UTI ✔
Q. Is the patient trying to tell me something?
A. Consider anxiety, depression, emotional deprivation or upset, change in environment, serious personal loss.

Probability diagnosis

The diagnosis depends on the age and presentation of the patient. In a teenager the probable causes of acute confusion or irrational behaviour include drug toxicityor withdrawal, schizophrenia, severe depression or a behavioural disorder.

It is the elderly who commonly present with confusion. The questions that must be asked are:

- Is the problem one of the 4 D's - dementia, delirium, depression, drugs or something else?
- If delirium is the problem, what is the cause?

Depression affects 15% of people over 65 and can mimic other causes of confusion and behavioural disturbance.

Significant prescribed drugs include hypnotics, sedatives, oral hypoglycaemics, antihypertensives, digoxin, antihistamines, anticholinergic drugs and antipsychotics.

Serious disorders not to be missed

There are many serious underlying disorders that must be considered, especially with delirium (Table 45.3). Cerebral organic lesions, including space-occupying lesions (e.g. cerebral tumour, subdural haematoma), severe infection (systemic or intracerebral) and carcinoma at any site, especially lung, breast, lymphoma or bowel, must be ruled out.

Table 45.3 Important causes of delirium (typical examples of each group)

Drug intoxication and drug sensitivity • anticholinergics • antidepressants • sedatives • alcohol, opioids, etc.
Withdrawal from substances of abuse and prescribed drugs • alcohol • opioids • amphetamines • cannabis • sedatives and anxiolytics
Infections *Specific* **Infections** • lower respiratory (e.g. pneumonia) • otitis media • cellulitis *Intracranial* • meningitis • encephalitis *Systematic* • infective endocarditis • septicaemia • HIV virus • other viral infections • malaria
Metabolic disturbances • uraemia, hepatic failure • electrolyte disturbances

• dehydration
Endocrine disturbances • diabetes ketosis, hypoglycaemia • hypothyroidism/hyperthyroidism
Nutritional and vitamin deficits • Wernicke's encephalopathy
Hypoxia • respiratory failure, cardiac failure, anaemia
Vascular • CVA
Head injury and other intracranial problems *Seizures* • complex partial seizures
'Subtle' causes • pain (e.g. herpes zoster) • emotional upset • environmental change • perioperative • faecal impaction • urinary retention

The sudden onset of delirium may suggest angina, myocardial infarction or a cerebrovascular accident. Twenty per cent of patients with delirium also have underlyingheart failure.[3]

Pitfalls

There are many pitfalls, especially with drug toxicity or withdrawal from the so-called illicitdrugs. In the elderly in particular, fluid and electrolyte disturbances, such as dehydration, hypokalaemia, hyponatraemia and hypocalcaemia, can cause delirium. Bowel disturbances such as faecal impaction or constipation can cause delirium and incontinence ofboth faeces and urine.

Seven masquerades checklist

All the following disorders can cause disturbed or confused behaviour, particularly in the elderly:

- depression: a very important cause of 'pseudodementia'
- drugs: toxicity or withdrawal (see Table 45.4)

Table 45.4 Prescribed drugs that can cause delirium

Anticholinergic • antiparkinsonian (e.g. benztropine) • tricyclic antidepressants
Tranquillisers and hypnotics • major tranquillisers (e.g. chlorpromazine) • minor tranquillisers (e.g. diazepam) • hypnotics • lithium
Antiepileptics
Antihistamines 1 and 2
Antihypertensives
Corticosteroids
Cardiac drugs • digoxin

• diuretics • beta-blockers
Opioids **Sympathomimetics**

- diabetes: especially hypoglycaemia, which can occur with type 2
- anaemia: often from self-neglect or chronic blood loss
- thyroid disorders: both hyperthyroidism andhypothyroidism can present with disturbed behaviour; ‘myxoedemic madness’ may be precipitated by atropine compounds
- urinary tract infection: causes or contributes to 20% of cases of hallucinations or illusions[2]
- spinal dysfunction: with its many severe pain syndromes, such as sciatica, it can be a significant factor

Psychogenic factors

Apart from the primary psychiatric disorders of anxiety, depression, mania and schizophrenia, relatively simple and subtle social problems, such as loneliness, boredom, a domestic upset, financial problem or similar issue, can trigger a confusional state.

THE CLINICAL APPROACH

History

Developing rapport with the disturbed or confused patient is essential and can be helped by a warm handshake or a reassuring pat on the shoulder. The basis of the history is a careful account from relatives or witnesses about the patient's behaviour.

When communicating with the patient, speak slowly and simply (avoid shouting), face the patientand maintain eye contact. Important features are the past history, recent psychosocial history including recent bereavement, family upsets and changes in environment. Search for evidence of depression and note any organic symptoms such as cough, constipation and so on.

Mental status examination

The most practical bedside screening test of mental function is the Mental Status Questionnaireof Kahn,[4] which includes ten simple questions.

1. What is the name of this place?
2. What city are you in now?
3. What year is it?
4. What month is it?
5. What is the date today?
6. What year were you born?
7. When is your birthday?
8. How old are you?
9. Who is the prime minister/president?
10. Who was the prime minister/president before him?

(Interpretation: normal 9 - 10; mildly impaired 8 - 9; confused/demented 7 or less.)

Other MMSEs are presented in Chapter 8, pp. 60 - 2.

If the patient has appropriate mental function ask questions related to possible depressive illness, such as:

- Do you feel hopeful about your future?
- Do you have good things to look forward to?
- Do you think life is worth living?
- Have you ever thought of taking your life?

Physical examination

The patient's general demeanour, dress and physical characteristics should be noted at all times. Assess the patient's ability to hear, see, speak, reason, obey commands, stand and walk. Any problems related to the special senses can cause confusion.

Look for features of alcohol abuse, Parkinson's disease and hypothyroidism.

Examine the neurological system and keep in mind the possibility of asubdural haematoma, whichmay have followed a forgotten fall.

Don't omit the rectal examination to exclude faecal impaction, melaena, cancer and prostatomegaly (in males) and also check the bladder for evidence of chronic retention.

Investigations

Investigations to consider for delirious or demented patients (unknown cause):

- urinalysis and microscopy
- cultures of blood and urine
- total and differential blood count; ESR
- blood glucose
- urea and creatinine and electrolytes
- calcium and phosphate
- thyroid function tests
- liver function tests
- serum vitamin B_{12} and folate levels
- ECG/troponin (? acute coronary syndrome)
- chest X-ray
- cerebral CT scan
- syphilis serology
- HIV
- arterial blood gases

MANAGEMENT OF THE ACUTELY DISTURBED PATIENT

Delirious or psychotic patients can be paranoid and respond defensively to the world around them. This behaviour can include aggressive and violent behaviour resulting in danger to themselves, their friends and family and to their medical attendants.

Dangerousness should be assessed from features such as the patient's past history (especially previous dangerous behaviour), age andsex, recent stress, victim behaviour, muscle bulk, presence of weapons, degree of overactivity and the manner of handling of the present distress by others. The patient may be in a state of acute panic and trying to flee a situation or in an agitated psychotic state prepared to confront the situation. It should be emphasised that most violent individuals are not mentally ill.

Most cases require an injection (the ideal intravenous administrationcan be extremely difficult and hazardous) which is often interpreted as a physical attack. It may not be possible to diagnose the cause of the problem before giving the injection.

Approach to management

- Assess the environment and don't move into the patient's space until in a position of control.
- React calmly. Communicate calmly and simply.
- State your task firmly and simply.
- Try to control the disturbed patient gently.
- Ensure the safety of all staff and make certain that heroics are not attempted in dangerous circumstances.
- An adequate number of staff to accompany the doctor is essential - six is ideal(one for immobilisation of each limb, one for the head and one to assist with drugs).[1]
- Patients should be placed on the floor in the prone position.

Principles of sedative administration[1]

- Use the safest possible route of administration whenever possible (i.e. oral in preference to parenteral). Intravenous administration has the lowest margin of safety.
- Parenteral administration should be restricted to severely disturbed patients.
- Closely monitor vital signs during and after sedative administration.
- Avoid intramuscular diazepam because of poor absorption.
- Avoid intravenous midazolam (Hypnovel) in such patients because of the risk of respiratory depression.
- Avoid benzodiazepines in patients with respiratory insufficiency. Haloperidol is an alternative.
- Patients have died from cardiopulmonary arrest after repeated sedative administration (especially benzodiazepines), so intensive monitoring is essential.

Monitor the following adverse effects:

- respiratory depression
- hypotension
- dystonic reactions, including choking
- neuroleptic malignant syndrome

Treatment options[1]

- diazepam 5 - 20 mg (o) as single dose (if patient cooperates); repeat every 2 - 6 hours (up to 120 mg daily) depending on response
- if intramuscular benzodiazepine required - midazolam (Hypnovel) 2.5 - 5 mg IM as single dose
- if intravenous benzodiazepine required
  - diazepam 5 - 20 mg IV (slow injection over several minutes) as single dose

Alternatives

- droperidol (Droleptan) 5 - 10 mg IM *or*
- haloperidol 5 - 10 mg IM
  (These injections can be repeated in 15 - 30 minutes if required. Droperidol is similar to haloperidol but more sedating. Keep in mind the rare but potentially fatal laryngeal dystonia with high doses - cover with benztropine 2 mg IM.) *or*(orally)
- chlorpromazine 100 mg (o) by tablet or syrup; repeat in 60 minutes if required. Avoid in older patients. IV use is dangerous (Syrup may be preferred as it is less likely to be hidden in the mouth.) *or*
- haloperidol 5 - 10 mg (o), repeat every 2 - 6 hours *or*
- combination of diazepam and haloperidol

Postdisturbance evaluation

Determine the likely cause, such as:

- acute organic brain syndrome
  - toxic causes
  - infection
- alcohol or drugs (illicit or prescribed)
  - intoxication
  - withdrawal
- manic illness
- severe depression
- schizophrenic syndrome
- severe panic

ACUTE ORGANIC BRAIN SYNDROME (DELIRIUM)

The many labels of acute organic brain syndrome include:

- delirium
- acute confusional state
- toxic confusional state
- confusional episode
- acute brain syndrome

Main features

- Clouding of conscious state
- Disorientation
- Impaired attention
- Impaired memory
- Global cognitive defect - onset over days/hours Refer to the box below.

DSM-IV criteria for delirium

Diagnosis of delirium requires evidence of:

1. Disturbance of consciousness
2. A change in cognition

  - perceptual disturbance
  - incoherent speech
  - disorientation
  - memory impairment
3. Clinical features appearing over a short period
4. Evidence of a cause

Other clinical features[1]

- The patients are usually elderly.
- Anxiety and agitation can be severe but in hypoactive deliria (usually due to metabolic disturbance) the conscious state can vary from drowsiness to coma.
- Odd behaviour with mood swings can occur.
- Psychotic symptoms can occur.
- Delusions are usually fleeting.
- The disturbance is usually worse at night and may be aggravated by sedation.
- Visual hallucinations are a feature of alcohol withdrawal.
- Attacks on bystanders may result (uncommon).

Always seek a cause.[1] A list of causes is presented in Table 45.3. The most important causes are:

- infections (usually in urinary tract, lungsor ear, or systemic in young or elderly)
- prescribed drugs

Anticholinergic delirium

Consider this cause (from drugs with anticholinergic properties or illicit substances). Features include hyperactivity, marked thought disorder, vivid visual hallucinations and very disturbed behaviour.

Differential diagnosis of delirium

In the earlier stages it may mimic the various psychiatric disorders, including anxiety, depression, various hallucinatory states, particularly agitated schizophrenia (rarely), extreme manic states, complex partial seizures, dementia. Consider deafness. Delirium is common in the hospital setting.

Investigations

Investigations are those listed under clinical approach (pp. 487 - 8).

Treatment

Principles:

- Acute delirium is a medical emergency.
- Establish normal hydration, electrolyte balance and nutrition.
- Attend to helpful environmental factors (e.g. calm atmosphere, a night-light, orientation clues, presence of friends and relatives).

Medication

Medication[1] may not beneeded but will be in the presence of anxiety, aggression or psychotic symptoms.

For anxiety and agitation:

- diazepam 5 - 10 mg (o) as a single dose *or*
- midazolam 2.5 - 5 mg IM *or*
- diazepam 5 - 20 mg IV (over several minutes) if a strong indication for rapid sedation exists For psychotic behaviour, add:
- haloperidol 1.5 - 10 mg (o) according to response *or*
- olanzapine 2.5 - 10 mg (o) daily in 1 or 2 doses

For severe symptoms, when parenteral medication is required (cover with benztropine 2 mg (o) or IM):

- haloperidol 5 - 10 mg (IM) as single dose *or*
- droperidol 5 - 10 mg (IM) as single dose (more sedating)

For anticholinergic delirium:

- tacrine hydrochloride 15 - 30 mg with caution by slow IV injection (an antidote)

Note:

- Benzodiazepines should be avoided in children and in patients with respiratory insufficiency.
- Consider necessity for pain relief.
- Use lower doses of parenteral medications in the very old and frail.

DEMENTIA (CHRONIC ORGANIC BRAIN SYNDROME)

Dementia is an important diagnosis to consider in the elderly patient. The DSM-IV criteria for dementia are presented in Table 8.2, p. 68 and elaborated in more detail in Chapter 8.

The main feature of dementia is impairment of memory, especially recent memory, when the person cannot remember what has happened a few hours (or even moments) earlier but may clearly remember the events of the past.

The more serious behavioural changes encountered with dementia tend to occur in the advanced stages. However, these disturbances may be precipitated by illness suchas infections, emotional upset and drugs.

These serious disturbances include:

- uninhibited behaviour
- hallucinations (generally uncommon)
- paranoid delusions

If a stable patient becomes acutely disturbed, delirium should be suspected.

Presenile dementia - Alzheimer's type

The main features are:

- onset in late 50s and early 60s
- insidious onset
- early loss of short-term memory
- progressive decline in intellect
- death in 5 - 10 years
- more common in Down syndrome

Differential diagnosis of dementia

There are two approaches to the differential diagnosis, including consideration of the classic causes of disturbed behaviour as summarised in the mnemonic in Table 45.5 [5] and those more everyday subtle causes presented in Table 8.4, p. 68 in the chapter on theelderly patient.

Table 45.5 Differential diagnosis of dementias

Ddelirium drugs (see toxic)
Eemotional disorder ° depression endocrine ° thyroid
Mmemory ° benign forgetfulness
Eelective ° anxiety disorders/neuroses
Nneurological
- CVA
- head trauma
Ttoxic
- drugs/medication
- metabolic disease
Iintellect - low or retarded
Aamnesic disorders - Korsakov's
Sschizophrenia (chronic)

*Source:*After McLean[5]

However, the foremost differential diagnosis should be ‘pseudodementia’ caused by severe depression.

A simple comparison between schizophrenia and dementia is shown in Table 45.6.

Table 45.6 Comparison of schizophrenia and dementia

	Dementia	Schizophrenia
Onset	Middle-aged or elderly	Young
Memory	Always impaired	Usually unaffected
Delusions	Rare	Frequent
Hallucinations	Uncommon	Frequent
Thought broadcasting	Never	Frequent

A vigorous search for a possible cause of dementia is warranted sincethere are a significant number of reversible causes. In particular, it is important to exclude the psychiatric conditions that may mimic dementia.

Treatment

See pp. 492 - 3.

- To control psychotic symptoms or disturbed behaviour:
  - risperidone 0.5 - 2 mg (o) daily *or*
  - olanzapine 5 - 10 mg (o) daily in 1 or 2 doses
- To control symptoms of anxiety and agitation - oxazepam 15 mg (o) 1 to 4 timesdaily. Avoid benzodiazepines for more than 2 weeks.

THE ACUTE PSYCHOTIC PATIENT

> Acute psychosis is the presence of the mental state where appreciationof reality is impaired as evidenced by the presence of typical psychotic symptoms such as delusions, hallucinations, mood disturbance and bizarre behaviour.[6]

The differential diagnoses of patients presenting with psychoses is presented in Table 45.7.

Table 45.7 Causes of psychoses[6]

Functional psychoses
• schizophrenia • schizoaff ective disorder (core symptoms of schizophrenia + mood symptoms) • bipolar mood disease (depressed or manic phase)
Drug-induced psychoses
Organic based psychoses
Other
• delusional disorder (paranoid psychoses) • brief psychotic disorder • folie à deux (psychosis occurring simultaneously in two close associates)

Early diagnosis

Early recognition of a psychosis, particularly schizophrenia, is so important as early intervention leads to improved outcomes. Early or prodromal symptoms include the following:

- social withdrawal
- reduced attention and concentration
- reduced drive and motivation
- depressed mood
- anxiety
- irritability/agitation
- suspiciousness
- sleep disturbance
- deterioration in role functioning

It is appropriate to ask the correct questions in order to elicit psychotic symptoms. These arepresented in Table 45.8.

Table 45.8 Questions for eliciting psychotic symptoms[6]

Anxiety	Have you been feeling especially nervous or fearful? Have you felt tense and shaky, or experienced palpitations?
Depressed mood	Have you been feeling sad or 'down in the dumps' recently, not enjoying activities as much as before?
Elevated mood	Have you been feeling especially good in yourself, more cheerful than usual and full of life?
Auditory hallucinations	Do you hear voices of people talking to you even when there is no-one nearby?
Thought insertion	Have you felt that thoughts are being put into your mind? Do you experience telepathy?
Thought withdrawal	Have you experienced thoughts being taken out of your mind?
Thought broadcasting	Have you felt that other people are aware of your thoughts?
Thought echo	Have you experienced voices or people echoing your thoughts?
Delusion of control	Have you felt under the control or influence of an outside force?
Delusions of reference	Do programs on the television or radio hold special meaning for you?
Delusions of persecution	Do you feel that you are being singled out for special treatment? Is there a conspiracy against you?
Delusions of grandeur	Do you feel special, with unusual abilities or power?
Delusions of guilt	Do you believe that you have sinned or have done something deserving punishment?

Reproduced with permission[6]

SCHIZOPHRENIA AND ASSOCIATED DISORDERS

The term schizophrenia (Bleuler 1911) refers to a group of severe psychiatric illnesses characterised by severe disturbances of emotion, language, perception, thought processes, volition and motor activity. The causes of schizophrenia disorders are unknown, but genetic factors and drug abuse are implicated.

Signs and symptoms of schizophrenia

Positive	- delusions
	- hallucinations
	- thought disorder
	- disorganised speech and behaviour
Negative	- flat affect
	- poverty of thought
	- lack of motivation
	- social withdrawal
	- reduced speech output
Cognitive	- distractibility
	- impaired working memory
	- impaired executive function e.g. planning
	- impaired insight
Mood	- mania (elevation)
	- depression

Other features include:

- bizarre behaviour
- subject to tension, anxiety or depression
- deterioration in work and study performance

- peak incidence 15 - 25 years[7] - smaller peak at 40 years
- lifetime prevalence 1 in 100
- equal sex incidence
- high risk of suicide

Differential diagnosis

Organic factors need to be excluded, especially drugs:

- amphetamines
- hallucinogens (e.g. LSD)
- marijuana

A comparison of delirium, dementia and functional psychosis is presented in Table 45.9.

Table 45.9 Comparison of the clinical features of delirium, dementia and acute functional psychoses[7]

Feature	Delirium	Dementia	Acute psychosis
Onset	Rapid	Slow - insidious	Rapid
Duration	Hours to weeks	Months to years	Depends on response to treatment
Course over 24 hours	Fluctuates - worse at night	Minimal variation	Minimal variation
Consciousness	Reduced	Alert	Alert
Perception	Misperceptions common, especially visual	Misperceptions rare	May be misperceptions
Hallucinations	Common, visual (usually) or auditory	Uncommon	Common, mainly auditory
Attention	Distractable	Normal to impaired	Variable - may be impaired
Speech	Variable, may be incoherent	Difficulty finding correct words	Variable: normal, rapid or slow
Organic illness or drug toxicity	One or both present	Often absent	Usually absent

Management

Drug treatment is only a part of the total management. Explanation and appropriate reassurance to the family with patient and family supportive care is obviously essential. Supportive psychotherapy is important in all phases. A team approach is necessary to cope with the disorder, which usually has a devastating effect on the family. Referral for specialist care is appropriate.

Acute phase

- Hospitalisation usually necessary.
- Drug treatment for the psychosis.[1]
  Drug treatment may include the first-generation (typical) antipsychotics such as haloperidol and chlorpromazine, which are effective for managing the 'positive' symptoms, or the second generation (atypical) antipsychotics such as risperidone, olanzapine, quetiapine, clozapine, amisulpride and aripiprazole, which in addition are more effective at treating the negative andother symptoms of schizophrenia.[8]
- The rule is to start with a low dose and titrate upwards at a rate and to a level that is optimal for the patient.

1. When oral medication is possible, first-line treatment is:[1]
    - olanzapine 5 - 10 mg (o) nocte *or*
    - risperidone 1 mg (o) at night increasing to1 mg bd and then titrated in 1 mg steps over several days to 2 - 4 mg (o) bd (beware of hypotension) *or*
    - quetiapine 25 mg (o) bd (day 1) increasing gradually as tolerated to 300 - 450 mg day in divided doses
2. When parenteral medication required:
    - haloperidol 5 - 10 mg IV or IM, initially, up to 20 mg in 24 hours, depending on the response *add*
    - benzotropine 1 - 2 mg (o) bd (to avoid dystonic reaction) *or*
    - zuclopenthixol acetate 50 - 150 mg IM as a single dose

   If dystonic reaction:

  - benztropine 1 - 2 mg IV or IM

If very agitated use:

  - diazepam 10 - 20 mg (o) or 5 - 10 mg IV

Chronic phase of schizophrenia

Long-term antipsychotic medication is recommended to prevent relapse.[1]

- Examples of oral medication regimens:[1]
  - olanzapine 10 - 20 mg (o) nocte *or*
  - risperidone 1 - 2 mg (o) bd *or*
  - quetiapine 150 mg (o) bd
- Aim for lowest possible dose to maintain control.
- Chlorpromazine is not recommended for longterm use because of photosensitivity reactions.
- Use depot preparations if compliance is a problem:
  - fluphenazine decanoate 12.5 - 75 mg IM, every 2 - 4 weeks (12.5 mg statim test dose) *or*
  - haloperidol decanoate 50 - 300 mg IM, every 4 weeks (50 mg statim) *or*
  - flupenthixol decanoate 20 - 80 mg IM, every 2 - 4 weeks (10 mg statim) *or*
  - zuclopenthixol 200 - 400 mg IM (100 mgstatim)

Tips with depot preparations:

- Start with IM test doses and then titrate to recommended controlling levels (half or full starting dose).
- May take 2 - 4 months to produce a stable response, so oral supplements may be necessary.
- Not as effective as oral therapy.
- Give as deep IM injection with 21 gauge needle in buttock.
- Use lowest possible dose to avoid tardive dyskinesia.
- Reassess at least every 3 months.
- Closely monitor patient for movement disorders.

Drug-resistant schizophrenia

Consider other causes (e.g. substances abuse). ECT may help the agitated patient, especially if catatonic. Consider a trial of clozapine (300 - 600 mg daily) with strict monitoring for blood dyscrasias or olanzapine (5 - 20 mg daily).

Movement disorders from antipsychotic medication[1]

Acute dystonias

- Usually bizarre muscle spasms affect face, neck, tongue and trunk
- Oculogyric crises, opisthotonos and laryngeal spasm

Treatment: benztropine 1 - 2 mg IV or IM

Akathisia

- Subjective motor restlessness of feet and legs
- Generally later onset in course of treatment

Treatment:

- reduce dosage until akathisia less troublesome or substitute thioridazine
- can use oral propranolol, diazepam or benztropine as a short-term measure

Parkinsonian

- Seen relatively early in treatment
- The akinesia can be confused with druginduced depression

Treatment:

- use lower dose or substitute a phenothiazine in low dosage
- alternatively, use benztropine or benzhexol

Tardive dyskinesia

Tardive dyskinesia is a syndrome of abnormal involuntary movements ofthe face, mouth, tongue, trunk and limbs. This is a major problem with the use of long-term antipsychotic drugs and may occur months or years (usually) after starting treatment.

Differential diagnosis:

- spontaneous orofacial dyskinesia
- senile dyskinesia
- ill-fitting dentures
- neurological disorders causing tremor and chorea

There is no specific treatment for tardive dyskinesia. The risks and benefits of continuing therapy have to be weighed.

Note: Because of the inability to manage tardive dyskinesia, prevention in the form of using the lowest possible dosage of antipsychotic medication is essential. This involves regular review and adjustment if necessary.

Neuroleptic (antipsychotic) malignant syndrome

This is a potentially fatal adverse effect that can develop at any time. It develops in hours to days.

Syndrome: high temperature, muscle rigidity, altered consciousness. Milder variants can occur.

Treatment:

- discontinue medication
- ensure adequate hydration with IV fluids
- if life-threatening:
    - bromocriptine 2.5 mg (o) bd, gradually increasing to 5 mg (o) tds and
    - dantrolene 50 mg (IV) every 12 hours for up to 7 doses
- consultant referral

Cardiac dysfunction

Various psychotrophic agents, particularly the phenothiazines, are prone to cause the adverse effect of prolongation of the QT interval with potential severe outcomes.

BIPOLAR DISORDER

The mood disorders are divided into depressive disorders and bipolar disorders. The swing in moods in bipolar disorders (manic depressive disorders) is illustrated in Figure 45.1. It affects 1% - 2% of the population.



Figure 45.1 Bipolar disorder (manic depression): possible mood swings

The symptoms of mania may appear abruptly.

The main clinical featuresof mania are:

- elevated, expansive or irritable mood
- accelerated speech
- agitation
- racing thoughts or flights of ideas
- increased activity
- reduced sleep

Other features include:

- grandiose ideas, sometimes paranoid
- reckless behaviour, overspending
- hasty decisions (e.g. job resignation, hasty marriages)
- impaired judgment
- increased sexual drive and activity
- poor insight into the problem
- variable psychotic symptoms - paranoia, delusions, auditory hallucinations

*Note:*The peak onset is in early adult life. There is a strong hereditary basis. Episodes may be precipitated by stress.

Hypomania is the term used to describe the symptoms of mania that are less severe and of shorter duration.

Management of acute mania

Hospitalisation

- For protection of patient and family
- Usually involuntary admission necessary

Drugs of choice [1, 9]

The basis of treatment is the mood stabilisers lithium carbonate, sodium valproate and carbamazepine.

1 Co-operative patient

lithium carbonate 750 - 1000 mg (o) daily[1]

- this is the initial dose
- give in 2 to 3 divided doses
- can increase by increments of 250 - 500 mg per day
- monitor by plasma levels
- therapeutic plasma level 0.8 - 1.4 mmol/L
- required daily dosage usually 1000 - 2500 mg
- elderly patients may require reduced dosage *or*
  (if not tolerated or for rapid cycling disorder)sodium valproate 400 - 800 mg (o) daily
- give in 2 divided doses
- introduce stepwise every 2 - 3 days
- check plasma levels after 7 days
- therapeutic level 350 - 700 μmol/L *or*
  carbamazepine 200 - 400 mg (o) daily
- give as above
- therapeutic plasma level 20 - 50 μmol/L *or*
  olanzapine 5 - 20 mg (o) daily, in 1 single dose at night or 2 divided doses

2 Unco-operative patients and manic behaviour problematic

haloperidol 10 - 20 mg (o) as single dose

- can be repeated up to 40 mg daily, depending on response
- use minimum possible dose to achieve control
- there is a risk of tardive dyskinesia

If parenteral antipsychotic drug required:

haloperidol 5 - 20 mg IM or IV *or*

droperidol 5 - 10 mg IM (more sedating)

- repeat in 15 - 30 minutes if necessary
- change to oral medication as soon as possible

*Note:*Benzodiazepines (e.g. diazepam)can be used with lithium and also the antiepileptics.

If not responding to medication consider ECT.

Maintenance

- Lithium carbonate - continue for 6 months; if not tolerated or ineffective use the antiepileptics, carbamazepine or sodium valproate. Give at minimum dose and for minimum time to gain control. Measure lithium levels every 1 to 3 months once desired level is achieved.
- Antiepileptics are claimed to be more effective in patients with rapid cycling illness (4 or more episodes per year).
- Remember to provide supportive psychotherapy with appropriate psychosocial interventions.

Prophylaxis for recurrent bipolar disorder

(Over 90% will have a recurrence at some time: consider medication iftwo or more episodes of either mania or depression in the previous 4 years.)

- Use long-term lithium (e.g. 3 - 5 years). Target plasma level for maintenance is usually 0.6 - 0.8 mmol/L. A recent study recommended lithium as the prime mood stabiliser.[10]
- If poor response use carbamazepine or sodium valproate.
- Unwanted side-effects of lithium include
    - a fine tremor
    - muscle weakness
    - weight gain
    - gastrointestinal symptoms
- With antiepileptics adjust dosage accordingto clinical response and toxicity.

BODY DYSMORPHIC DISORDER[9]

Body dysmorphic disorder is characterised by a preoccupation with thebelief that some aspect of physical appearance is abnormal, unattractive or diseased. The person's concern and distress is out of proportion to any imagined or actual defect and usually not amenable to reassurance. This preoccupation causes significant functional impairment. The condition rarely presents directly and may be over represented in the area of dermatology orplastic surgery. It begins in late childhood or early adolescence. The person's focus is on the face, head or secondary sexual characteristics.

Patients may be helped by counselling and psychotherapy including CBT. There is clinical evidence that SSRIs help if the symptoms suggest an obsessive compulsive disorder. An antipsychotic agent may help where beliefs are delusional or in the context of a psychotic disorder.

DEPRESSION

Depression is very common and presents in a great range of severity. In the context of 'the disturbed patient' depression can be confused with dementia or a psychosis, particularly if the following are present:

- psychomotor agitation
- psychomotor retardation
- delusions
- hallucinations

Assessment[1]

The following questions need to be addressed:

- Is the depression primary (i.e. not secondary to another psychiatric condition such as schizophrenia or anxiety disorder)?
- Is it part of a bipolar disorder? Has there been a previous manic or hypomanic episode? If so, a different approach to treatment is required.
- Is the depression caused by another illness or physical factor (e.g. hypothyroidism, cerebrovascular disease or medication)?

- Is the patient psychotic?
- Is the patient a suicide risk?

The treatment of depression is presented in Chapter 20.

PSYCHOACTIVE SUBSTANCE USE DISORDERS

It is important for the GP to be aware of the effects of self-administration of psychoactive substances, especially their toxic or withdrawal effects. They form significant consideration for the differential diagnosis of disturbed patient behaviour. The following substances can cause these effects.

Alcohol

Toxic and withdrawal effects, including delirium tremens, are outlined in Chapter 117. Abrupt withdrawal can cause symptoms ranging from tremors, agitation and dysphoria (feeling thoroughly miserable) to fully developed delirium tremens. Epileptic seizures may also occur.

Barbiturate dependence

Tolerance and symptoms on withdrawal are the main features. Barbiturate withdrawal is a very serious life-threatening problem and may be encountered in elderly people undergoing longstanding hypnotic withdrawal. Symptoms include anxiety, tremor, extreme irritability, twitching, seizures and delirium.

Management[1]

Undertake withdrawal with medical supervision as an inpatient.

Transfer the patient to phenobarbitone or diazepam.

- phenobarbitone 120 mg (o) hourly until sedation *or*
- phenobarbitone 30 mg for each 100 mg of shorter-acting barbiturate reduce the dose gradually over 10 to 14 days *or*
- diazepam 20 - 40 mg orally, daily
- reduce the dose gradually over 10 to 14 days

Benzodiazepine dependence

Withdrawal symptoms in the dependent patient include anxiety, restlessness, irritability, palpitation and muscle aches and pains, but delirium and seizures are uncommon except with very high doses. The shorter the half-life the greater the dependence.

Withdrawal is best achieved by supervising a gradual reduction in dosage aided by relaxation techniques and behavioural strategies to help patients cope with insomnia and anxiety.

Refer to Chapter 22 for the effects of opioid dependence, stimulant substance abuse p. 206, hallucinogen abuse p. 207 and cannabis use and dependence p. 207.

PSYCHIATRIC DISORDERS OF CHILDHOOD AND ADOLESCENCE[1]

The following disturbance problems do occur and must be taken seriously, especially the potential for suicide in the second decade. Many of these disorders are presented in more detail in Chapter 84.

Attention deficit hyperactivity disorder

Clinical features:

- short attention span
- distractability
- overactivity
- impulsiveness
- antisocial behaviour

Depression

Major depression follows the same criteria as for adults. Suicidal ideation has to be considered and taken very seriously if present. Imipramine is probably the drug of choice.

Bipolar disorders

Mania is seldom diagnosed before puberty. Adolescents may present (uncommonly) with symptoms of mania or hypomania.

Schizophrenia and related disorders

Schizophrenia is rare before puberty. The criteria for diagnosis are similar to adults:

- delusion
- thought disorder
- hallucinations
- 6 months or more of deterioration in functioning

Autism

Aggression and irritability can be a feature, especially during adolescence.

Tourette's syndrome

Behavioural problems can be part of this syndrome, which requires the attention of an experienced consultant.

Obsessive-compulsive disorders

In about one-third of cases the onset is between 5 and 15 years of age.

VIOLENCE AND DANGEROUSNESS

Dangerousness has been defined as a ‘propensity to cause serious physical injury or lasting psychological harm to others’ and, in the context of the mentallyabnormal, ‘the relative probability of their committing a violent crime’.[11]

Dangerousness is not related only to mental illness and, interestingly, most offenders have no sychiatric diagnosis. It is not an inherited, immutable characteristic of an individual but tends to surface on impulse in a particular context given a whole range of situational factors. Prediction of the risk of violence is not straightforward.

Various groups have been identified as contributing risk factors forviolent conduct.[11]

- schizophrenic psychoses, including:
  - older male paranoid schizophrenics
  - younger males prone to act violently and impulsively, presumably due to hallucinatory commands
- morbid jealousy:
  - associated with delusions of infidelity
- antisocial personality disorder
- mood disorder:
  - violence, usually associated with depression (rarely mania)
  - married women with severe depression (violence against young children)
  - history of suicide attempts in depression
- episodic discontrol syndrome (similar to intermittent explosive disorder)
- intellectual disability combined with personality disorder and behavioural disturbances
- alcohol abuse or dependency
- amphetamine or benzodiazepine abuse or dependency

From a management viewpoint, homicidal threats must be taken very seriously.

SUICIDE AND PARASUICIDE

The haunting issue of suicide and parasuicide is presented in Chapter 20. The disturbed patientis always a suicide risk rather than a homicide risk. The importance of recognising depression with an associated suicide risk in the elderly patient has been emphasised heavily in this chapter.

Facts and figures[12]

- More than 90% of suicides occur without underlying chronic conditions but most people are significantly depressed at the time.
- In Australia suicide is the second most common cause of death between the ages of 11and 25 years. Children as young as 5

years of age have committed suicide.

- Those who talk about suicide may attempt it later.
- About half those committing suicide have seen a doctor within their last month of life.
- Around 80 - 90% of suicides have given clear or subtle warnings to family, friends or doctors.
- There is no evidence that asking patients about suicidal ideation provokes suicidal acts.
- Doctors in Australia and other Western countries have a high suicide rate.

Suicide risk

Blumenthal's[13] overlapping model lists five groups of risk factors (Figure 45.2):

1. Psychiatric disorders:
    - affective disorder and alcohol abuse in adults
    - schizophrenia
    - depression and conduct disorder in young people
2. Personality traits:
    - impulsiveness and aggression
3. Environmental and psychosocial factors:
    - poor social supports
    - chronic medical illness (e.g. AIDS)
    - significant loss
4. Family history and genetics (both nature and nurture):
    - emulation of relatives
    - specific ethnic groups in custody
5. Biological factors:
    - possible serotonin deficiency


Overlap model
(five domains)
5. Biology
3. Psychosocial life events, and chronic medical illness
1. Psychiatric disorder
4. Family history and genetics
2. Personality traits

Figure 45.2 Overlap model for understanding suicidal behaviour. After Blumenthal and Kupfer.

Parasuicide

Parasuicide is attempted suicide; in many cases patients are drawing attention to themselves as a 'plea for help'. It is important for the GP to take an active role in the support of the patient and family after discharge from hospital, but preferably in conjunction with a psychiatric or counselling service. Arrange frequent consultationsat first and ensure adequate follow-up, especially for missed appointments.

PERSONALITY DISORDERS

People with personality disorders may become very distressed and acutely disturbed under stress or provocation, and this may involve dramatic scenes, including public suicide threats. It is important to recognise personality disorders because they usually cause considerable distress to the patients, their family, society and GPs.

In practice the personality disorders of most concern are those that present with hostility, either verbal or physical, particularly if a suicide or homicide threatis involved. It is a mistake to assume that those patients who manifest violent or psychopathicbehaviour have a personality disorder or, conversely, the meek and mild are free from personality disorder.

The diagnosis of personality disorder can be difficult. As practitioners we tend to have a 'gut feeling' about the diagnosis but often find it difficult to classify the personality of the patient and then to manage it appropriately.

The main characteristics of a personality disorder are:[12]

- lack of confidence and low self-esteem
- long history from childhood
- difficulties with interpersonal relationships and society
- recurrent maladaptive behaviour
- relatively fixed, inflexible and stylised reaction to stress
- minimal insight
- perception of difficulties as external to themselves

The medical/psychiatric significance:

- maladaptive relationships with GPs and society
- problem of sexually dysfunctional lives
- risk of substance abuse and self-destructive behaviour
- prone to depression and anxiety (usually low grade)
- susceptible to 'breakdown' under stress

Personality is the result of a genetic template and the continuing interaction of the person with outside influences (peer pressures, family interactions, influential events) and personal drives in seeking an identity. The outcome is a unique behaviour pattern manifest as a personality trait or character reflective of the individual's self-image and fundamental to his or her sense of personal identity.[14]

Although personality is unique, it is possible to make a hypothesis that one is normal or abnormal. If abnormal, it is possible to stereotype it according to the predominant symptoms or behaviours.

Using the International Classification of Disease (ICD-10) and the DSM-IV classification[15] various subtypes are readily identifiable (Table 45.8),[15] which can be considered in four main groups. There is a considerable overlap between the subtypes within the group[16] and it is more important to understand the specific features of a person's personality rather than categorise them.[17]

The antisocial group tend to come to the attention of GPs more frequently with some individuals representing 'heart-sink' patients because of demanding, angry or aggressive behaviour. The withdrawn group are typically withdrawn, suspicious and socially isolated but fall short of a true psychotic syndrome. GPs have problems communicating with them because they are often suspicious, which can make proper physical examination and management difficult.

In the dependent and inhibited groups, which may overlap with an anxiety state, the main features are nervousness, timidity, emotional dependence and fear of criticism and rejection. They are frequent attenders (the 'fat file' syndrome) and are often accompanied by friends and relatives because of their insecurity.

Management

The best treatment is a supportive 'therapeutic' community and an understanding and supportive GP. It is vital to understand that people with personality disorders perceive the world from a fundamentally different perspective. Problematic patients, if agreeable, may respond well to psychological intervention and behavioural techniques, especially operant conditioning (reinforcing acceptable behaviour) and averse conditioning (correcting inappropriate behaviour).[14] CBT has the most to offer.

The borderline and narcissistic disorders in particular respond well to specific types of psychotherapeutic intervention. Patients' self-esteem needs carefulsupport while maladaptive modes of behaviour are confronted. Hospitalisation is rarely requiredexcept for those at risk of suicide (e.g. antisocial patients).

Medication has limitations but may be useful to treat those individuals who temporarily decompensate into a psychosis, an anxiety state or depression. One study has shown that antipsychotic medication in low dosage (e.g. haloperidol 5 mg daily) is effective in treating the problematic behaviours in paranoid and some antisocial personality disorders.[18]

There are dangers to the therapist and it is important not to ‘buy into’ a particular psychopathy, especially with seductive, manipulative or paranoid patients.[17]

WHEN TO REFER[12]

Indications for referral to a psychiatrist:

- severe depression
- high suicide risk
- actual suicide attempt: recent or in the past
- suspected psychiatric disorders in the elderly:
    - ? depression or schizophrenia
    - ? depression or dementia
- failure to improve with treatment
- poor family and social supports

Table 45.10 Summary of main personality disorders

Main cluster group	Subtypes	Main features of disorder
A *Withdrawn*	• Paranoid	• Suspicious, oversensitive, argumentative, defensive, hyperalert, cold and humourless
synonyms odd eccentric	• Schizoid	• Shy, emotionally cold, introverted, detached, avoids close relationships
	• Schizotypal	• Odd and eccentric, sensitive, suspicious and superstitious, socially isolated, odd speech, thinking and behaviour. Falls short of criteria for schizophrenia
B *Antisocial*	Antisocial (sociopathic, psychopathic)	Impulsive, insensitive, selfish, callous, superficial charm, lack of guilt, low frustration level, doesn't learn from experience, relationship problems (e.g. promiscuous)
synonyms dramatic emotional sociopathic flamboyant erratic	Histrionic (hysterical)	Self-dramatic, egocentric, immature, vain, dependent, manipulative, easily bored, emotionalscenes, inconsiderate, seductive, craves attention and excitement
	Narcissistic (‘prima donna’)	Morbid self-admiration, exhibitionist, insensitive, craves and demands attention, exploits others, preoccupied with power, lacks interest in others, bullying, insightless
	Impulsive (‘mad dog’)	Mood swings, impulsiveness, prone to violence and threats (especially if criticised) but doesn't lack guilt
	Borderline (‘hell-raiser’)	Confused self-image, impulsive, reckless, ‘all or nothing’ relationships - unstable and intense, damaging reckless behaviour, full of anger and guilt, lacks self-control ± uncontrolled gambling, spending etc. *Note:*High incidence suicide and parasuicide; drug abuse
C *Dependent*	Avoidant (anxious)	Anxious, self-conscious, fears rejection, timid and cautious, low self-esteem, overreacts to rejection and failure
synonyms anxious fearful inhibited	Dependent	Passive, weak willed, lacks vigour, lacks self-reliance and confidence, overaccepting, avoids responsibility, seeks support
	Obsessional (obsessive-compulsive)	Rigid, perfectionist, pedantic, indecisive, egocentric, preoccupied with orderliness and control
	Passive-aggressive	Procrastinates, childishly stubborn, dawdles, sulks, argumentative, clings, deliberately inefficient and hypercritical of authority figures
D *Other*	Hypochondrial	Health-conscious, disease fearing, symptom preoccupation
	Depressive (dysthymic,	Pessimistic, anergic, low self-esteem, gloomy, chronic mild depression

	cyclothymic)

REFERENCES

1. Dowden J (Chair). *Therapeutic Guidelines: Psychotropic* (Version 5). Melbourne: Therapeutic Guidelines Ltd, 2003: 65 - 94.
2. Biro G. Dementia. Aust Dr Weekly, 1990; 16 February: I - VIII.
3. Biro G. Delirium in the elderly. Aust Dr Weekly, 1989; 1 December: I - VIII.
4. Kahn Rlet al. Brief objective measures of the determination of mental status in the aged. Am J Psychiatry, 1960; 117: 326 - 9.
5. McLean S. Is it dementia? Aust Fam Physician, 1992; 21: 1762 - 6.
6. Keks N, Blashki G. The acutely psychotic patient: assessment and initial management. Aust Fam Physician, 2006; 35(3): 90 - 4.
7. Norman T, Judd F. Schizophrenia. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 455 - 7.
8. Lovric K. Schizophrenia: update. Medical Observer, 2004; 17 September: 31 - 2.
9. Dowden J (Chair). *Therapeutic Guidelines: Psychotrophic* (Version 5). Melbourne: Therapeutic Guidelines Ltd, 2003: 119 - 185.
10. Sachs GS. A 25-year-old woman with bipolar disorder. JAMA, 2001; 285: 454 - 62.
11. Beaumont PJV, Hampshire RB. *Textbook of Psychiatry*. Melbourne: Blackwell Scientific Publications, 1989: 283 - 4.
12. Biro G. Suicide. Aust Dr Weekly, 1991; 26 April: I - VIII.
13. Blumenthal S. Suicide - a guide to risk factors, assessment and treatment of suicidal patients. Med Clin North Am, 1988; 72: 937 - 63.
14. Tierney LM et al. *Current Medical Diagnosis and Treatment* (43rd edn). New York: The McGraw-Hill Companies, 2006; 1054 - 6.
15. American Psychiatric Association: *Diagnostic and Statistical Manual of Mental Disorders* (4th edn). Washington DC; American Psychiatric Association, 2000.
16. Pullen I, Wilkinson G et al. *Psychiatry and General Practice Today*. London: RC Psych & RCGP, 1994: 180 - 3.
17. Kaplan R. Personality disorders: diagnoses and treatment. Medical Observer, 2001; 24 August: 32 - 3.
18. Soloff PH et al. Progress in pharmacotherapy of borderline disorders: a double blind study of amitriptyline, haloperidol and placebo. Arch Gen Psychiatry, 1986; 43: 691 - 7.

CHAPTER 46. DIZZINESS

I got my giddiness in 1690 (at the age of 23) by eating 100 golden pippins at a time at Richmond. Four years later at a place 20 miles further on in Surrey I got my deafness; and these two ‘friends’ have visited me one or other year since, and being old acquaintances have often sought fit to come together.

Jonathan Swift (1667 - 1745), describing his Meniere's syndrome

DIZZINESS

When patients complain of ‘dizziness’, they can be using this term to describe many different phenomena, and hence a careful history is required to unravel the problem. Other patients may use different terms to explain the same sensation, for example, ‘giddiness’, ‘swimming in the head’, ‘my brain spinning’, ‘whirling’ and ‘swinging’.

‘Dizzy’ comes from an old English word, *dysig*, meaning foolish or stupid. Strictly speaking, it means unsteadiness or lightheadedness - without movement or motion or spatial disorientation.

‘Vertigo’, on the other hand, comes from the Latin word for turning. The modern medical definition of vertigo is ‘a sudden sense of movement’.[1] It should describe a hallucination of rotation of self or the surroundings in a horizontal or vertical direction.

The term ‘dizziness’, however, is generally used collectively to describe all types of equilibrium disorders and, for convenience, can be classified as shown in Figure 46.1.



Figure 46.1 Classification of dizziness

KEY FACTS AND CHECKPOINTS

- Approximately one-third of the population will have suffered from significant dizziness by age 65 and about a half by age 80.[2]
- The commonest causes in family practice are postural hypotension and hyperventilation.
- The ability to examine and interpret the sign of nystagmus accurately is important in the diagnostic process.
- A drug history is very important, including prescribed drugs and others such as alcohol, cocaine, marijuana and illicit drugs.
- Meniere's syndrome is overdiagnosed. It has the classic triad: vertigo - tinnitus - deafness (sensorineural).
- Vertebrobasilar insufficiency is also overdiagnosed as a cause of vertigo. It often causes dizziness and sometimes vertigo but rarely in isolation.

DEFINED TERMINOLOGY

Vertigo

Vertigo is an episodic sudden sensation of circular motion of the body or of its surroundings. Other terms used by the patient to describe this symptom include ‘everything spins’, ‘my head spins’, ‘the room spins’, ‘whirling’, ‘reeling’, ‘swaying’, ‘pitching’ and ‘rocking’.

Vertigo is characteristically precipitated by standing or turning the head or by movement. Patients have to walk carefully and may become nervous about descending stairs or crossing the road and usually seek support. Therefore the vertiginous patient is usually very frightened and tends to remain immobile during an attack.

Patients may feel as though they are being impelled by some outside force that tends to pull them to one side, especially while walking.

True vertigo is a symptom of disturbed function involving the vestibular system or its central connections. It invariably has an organic cause. Important causes are presented in Table 46.1, while Figure 46.2 illustrates central neurological centres that can cause vertigo.


temporal lobe
• tumour
• focal epilepsy
eye (disorders causing diplopia)
labyrinth
• labyrinthitis
• Meniere's syndrome
vestibular nerve
• acoustic neuroma
• vestibular neuronitis
cerebellum
• tumours
• inflammation
brain stem
• multiple sclerosis
• vascular insufficiency
posterior inferior cerebellar artery
• thromboembolism
vertebral artery
• thromboembolism

Figure 46.2 Diagrammatic illustration of central centres that can cause vertigo

Table 46.1 Causes of vertigo

1 Peripheral disorders
Labyrinth
• labyrinthitis: viral or suppurative

• Meniere's syndrome
• benign paroxysmal positional vertigo (BPPV)
• drugs
• trauma
• chronic suppurative otitis media
8th nerve
• vestibular neuronitis
• acoustic neuroma
• drugs
Cervical vertigo
2 Central disorders
Brain stem
• vertebrobasilar insufficiency
• infarction
Cerebellum
• degeneration
• tumours
Migraine
Multiple sclerosis

Nystagmus is often seen with vertigo and, since 80 - 85% of causes are due to an ear problem, tinnitus and hearing disorders are also associated. In acute cases there is usually a reflex autonomic discharge producing sweating, pallor, nausea and vomiting.

Giddiness

Giddiness is a sensation of uncertainty or ill-defined lightheadedness. Other terms used by patients include ‘a swimming sensation’, ‘walking on air’ and ‘ground going beneath me’. It usually contains no elements of rotation, impulsion, tinnitus, deafness, nausea or vomiting.

The patient with giddiness, although fearful of falling or swooning, can nonetheless walk without difficulty if forced to do so.

Giddiness is a typical psychoneurotic symptom.

Syncopal episodes

Syncope may present as a variety of dizziness or lightheadedness in which there is a sensation of impending fainting or loss of consciousness. Common causes are cardiogenic disorders and postural hypotension, which are usually drug-induced.

Disequilibrium

Disequilibrium implies a condition in which there is a loss of balance or instability while walking, without any associated sensations of spinning. Other terms used to describe this include ‘unsteadiness on feet’, ‘the staggers’, ‘swaying feeling’ and ‘dizzy in the feet’.

Disequilibrium is usually of neurogenic origin.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 46.2.

Table 46.2 Dizziness/vertigo: diagnostic strategy model

Q. Probability diagnosis
A. Anxiety-hyperventilation (G)

Postural hypotension (G/S) Simple faint - vasovagal (S) Ear infection - acute labyrinthitis (V) Vestibular neuronitis (V) Benign paroxysmal positional vertigo (V) Motion sickness (V) Post head injury (V/G) Cervical dysfunction/spondylosis
Q. Serious disorders not to be missed A. Neoplasia • acoustic neuroma • posterior fossa tumour • other brain tumours, primary or secondary Intracerebral infection (e.g. abscess) Cardiovascular • arrhythmias • myocardial infarction • aortic stenosis Cerebrovascular • vertebrobasilar insufficiency • brain stem infarct, e.g. PICA thrombosis Multiple sclerosis
Q. Pitfalls (often missed) A. Ear wax - otosclerosis Arrhythmias Hyperventi lation Alcohol and other drugs Cough or micturition syncope Vertiginous migraine Parkinson's disease Meniere's syndrome (overdiagnosed) *Rarities* • Addison's disease (p. 224) • Neurosyphilis • Autonomic neuropathy • Hypertension • Subclavian steal • Perilymphatic fistula • Shy-Dragar syndrome
Q. Seven masquerades checklist A. Depression Diabetes possible Drugs

Anaemia
Thyroid disorder possible
Spinal dysfunction
UTI possible

Q. Is the patient trying to tell me something?

A. Very likely. Consider anxiety and/or depression.

G ° giddiness; S ° syncope; V ° vertigo

Probability diagnosis

In medical school we gain the wrong impression that the common causes of dizziness or vertigo are the relatively uncommon causes, such as Meniere's syndrome, aortic stenosis, Stokes-Adams attacks, cerebellar disorders, vertebrobasilar disease and hypertension. In the real world of medicine one is impressed by how often dizziness is caused by relatively common benign conditions, such as hyperventilation associated with anxiety, simple syncope, postural hypotension due to drugs and old age, inner ear infections, wax in the ears, post head injury, motion sickness and alcohol intoxication. In most instances making the correct diagnosis (which, as ever, is based on a careful history) is straightforward, but finding the underlying cause of true vertigo can be very difficult.

The common causes of vertigo seen in general practice are benign paroxysmal positional vertigo (BPPV, so often related to cervical vertebral dysfunction), vestibular neuronitis and acute viral labyrinthitis.

Viral labyrinthitis is basically the same as vestibular neuronitis, except that the whole of the inner ear is involved so that deafness and tinnitus arise simultaneously with severe vertigo.

Serious disorders not to be missed

Neoplasia

The important serious disorders to keep in mind are space-occupying tumours, such as acoustic neuroma, medulloblastoma and other tumours (especially posterior fossa tumours) capable of causing vertigo, intracerebral infections and cardiovascular abnormalities.

It is important to bear in mind that the commonest brain tumour is a metastatic deposit from carcinoma of the lung.[3]

Acoustic neuroma

This uncommon tumour should be suspected in the patient presenting with the symptoms shown in the diagnostic triad below. Headache may occasionally be present.

DxT: (unilateral) tinnitus + hearing loss + unsteady gait ° acoustic neuroma

Diagnosis is best clinched by high-resolution MRI. Audiometry and auditory evoked responses are also relevant investigations.

Cardiac disorders

Cardiac disorders that must be excluded for giddiness or syncope are the various arrhythmias, such as Stokes-Adams attacks caused by complete heart block, aortic stenosis and myocardial infarction.

Cerebrovascular causes

The outstanding cerebrovascular causes of severe vertigo are vertebrobasilar insufficiency and brain stem infarction. Vertigo is the commonest symptom of transient cerebral ischaemic attacks in the vertebrobasilar distribution.[1]

Severe vertigo, often in association with hiccoughs and dysphagia, is a feature of the variety of brain stem infarctions known as the lateral medullary syndrome due to posterior inferior cerebellar artery (PICA) thrombosis. There is a dramatic onset of vertigo with cerebellar signs, including ataxia. There are ipsilateral cranial nerve (brain stem) signs with contralateral spinothalamic sensory loss. Diagnosis is by CT or MRI scanning.

Neurological causes

Important neurological causes of dizziness are multiple sclerosis and complex partial seizures.

The lesions of multiple sclerosis may occur in the brain stem or cerebellum. Young patients who present with a sudden onset of vertigo with 'jiggly' vision but without auditory symptoms should be considered as having multiple sclerosis. Five per cent of cases of multiple sclerosis present with vertigo.

Pitfalls

A list of conditions causing dizziness that may be misdiagnosed is presented in Table 46.2. Wax in the ear certainly causes dizziness, though its mechanism of action is controversial. Cough and micturition syncope do occur, although they are uncommon.

Meniere's syndrome is a pitfall in the sense that it tends to be overdiagnosed.

Seven masquerades checklist

Of these conditions, drugs and vertebral dysfunction (of the cervical spine) stand out as important causes. Depression demands attention because of the possible association of anxiety and hyperventilation.

Diabetes mellitus has an association through the possible mechanisms of hypoglycaemia from therapy or from an autonomic neuropathy.

Drugs

Drugs usually affect the vestibular nerve rather than the labyrinth. Drugs that cause dizziness are presented in Table 46.3.

Table 46.3 Drugs that can cause dizziness

Alcohol
Aspirin and salicylates
Antibiotics: streptomycin, gentamicin, kanamycin, tetracyclines
Antiepileptics: phenytoin
Antidepressants
Antihypertensives
Antihistamines
Cocaine
Quinine-quinidine
Tranquillisers: phenothiazines, phenobarbitone, benzodiazepines
Diuretics in large doses: intravenous frusemide, ethacrynic acid
Glyceryl trinitrate

Cervical spine dysfunction

It is not uncommon to observe vertigo in patients with cervical spondylosis or post cervical spinal injury. It has been postulated4 that this may be caused by the generation of abnormal impulses from proprioceptors in the upper cervical spine, or by osteophytes compressing the vertebral arteries in the vertebral canal. Some instances of benign positional vertigo are associated with disorders of the cervical spine.

Psychogenic considerations

This may be an important aspect to consider in the patient presenting with dizziness, especially if the complaint is giddiness or light-headedness. An underlying anxiety may be the commonest cause of this symptom in family practice and clinical investigation of hyperventilation may confirm the diagnosis. The possibility of depression must also be kept in mind.[5] Many of these patients harbour the fear that they may be suffering from a serious disorder, such as a brain tumour or multiple sclerosis, or face an impending sroke or insanity. Appropriate reassurance to the contrary is often positively therapeutic for that patient.

The clinical approach

The essentials of the diagnostic approach include careful attention to the history and physical examination, and judicious selection of specific office tests and special investigations.

History

It is important to get patients to explain the precise nature of the symptoms, even asking their opinion as to the cause of their dizziness.

The following questions should be addressed:

- Is it vertigo or pseudovertigo?
- Symptom pattern:
    - paroxysmal or continuous?
    - effect of position and change of posture?
- Any aural symptoms?
    - tinnitus?
    - deafness?
- Any visual symptoms?
- Any neurological symptoms?
- Any nausea or vomiting?
- Any symptoms of psychoneurosis?
- Any recent colds?
- Any recent head injury (even trivial)?
- Any drugs being taken?
    - alcohol?
    - marijuana?
    - hypotensives?
    - psychotropics?
    - other drugs?

Physical examination

A full general examination is appropriate with particular attention being paid to the cardiovascular and central nervous systems and the auditory and vestibular mechanisms.

Examination guidelines are:

1. ear disease
    - auroscopic examination: wax? drum?
    - hearing tests
    - Weber and Rinne's tests
2. the eyes
    - visual acuity
    - test movements for nystagmus
3. cardiovascular system
    - evidence of atherosclerosis
    - blood pressure: supine, standing, sitting
    - cardiac arrhythmias
4. cranial nerves
    - 2nd, 3rd, 4th, 6th and 7th
    - corneal response for 5th
    - 8th - auditory nerve
5. the cerebellum or its connections
    - gait
    - co-ordination

  - reflexes
  - Romberg test
  - finger nose test: ? past pointing

6. the neck, including cervical spine
7. general search for evidence of:
   - anaemia
   - polycythaemia
   - alcohol dependence

Special office tests for dizziness

- Ask the patient to perform any manoeuvre that may provoke the symptom.
- Carry out head positional testing to induce vertigo and nystagmus (Figure 46.3).



Figure 46.3 Positional testing for benign positional vertigo (head taken rapidly from sitting position to hanging position with head turned to right: one of the three positions) - Hallpike's manoeuvre

- Take blood pressure measurements in three positions.
- Perform forced hyperventilation (20 to 25 breaths per minute) for 2 minutes.
- Carry out palpation of carotid arteries and carotid sinus (with care).

Investigations

Appropriate laboratory tests should be selected from Table 46.4.

Table 46.4 Investigations

Haemoglobin
Blood glucose
ECG: ? Holter monitor
Audiometry
Brain-stem evoked audiometry
Caloric test
Visual evoked potentials (MS)
Electrocochleography
Electro-oculography (electronystagmography)
Rotational tests
Radiology
• chest X-ray (? bronchial carcinoma)
• cervical spine X-ray
• CT scan
• MRI (the choice to locate acoustic neuroma or other tumour - may detect MS)

Diagnostic guidelines

- A sudden attack of vertigo in a young person following a recent URTI is suggestive of vestibular neuronitis.
- Dizziness is a common symptom in menopausal women and is often associated with other features of vasomotor instability.
- Phenytoin therapy can cause cerebellar dysfunction.
- Postural and exercise hypotension are relatively common in the older atherosclerotic patient.
- Acute otitis media does not cause vertigo but chronic otitis media can, particularly if the patient develops a cholesteatoma, which then erodes into the internal ear causing a perilymph fistula.

Dizziness in children

Dizziness is not a common symptom in children. Vertigo can have sinister causes and requires referral because of the possibility of tumours, such as a medulloblastoma. A study by Eviatar and Eviatar6 of vertigo in children found that the commonest cause was a seizure focus particularly affecting the temporal lobe. Other causes included psychosomatic vertigo, migraine and vestibular neuronitis.

Apart from the above causes it is important to consider:

- infection (e.g. meningitis, meningoencephalitis, cerebral abscess)
- trauma, especially to the temporal area
- middle ear infection
- labyrinthitis (e.g. mumps, measles, influenza)
- benign paroxysmal vertigo (short-lived attacks of vertigo in young children between 1 and 4 years of age: tends to precede adulthood migraine)[7]
- hyperventilation
- drugs - prescribed
- illicit drugs (e.g. cocaine, marijuana)
- cardiac arrhythmias
- alcohol toxicity

A common trap is the acute effect of alcohol in curious children who can present with the sudden onset of dizziness.

Dizzy turns in girls in late teens

- These are commonly due to blood pressure fluctuations.

- Give advice related to reducing stress, lack of sleep and excessive exercise.
- Reassure that it settles with age (rare after 25 years).

Dizziness in the elderly

Dizziness is a relatively common complaint of the elderly. Common causes include postural hypotension related mainly to drugs prescribed for hypertension or other cardiovascular problems. Cerebrovascular disease, especially in the areas of the brain stem, is also relevant in this age group. True vertigo can be produced simply by an accumulation of wax in the external auditory meatus, being more frequent than generally appreciated.

Middle ear disorder is also sometimes the cause of vertigo in an older person but disorder of the auditory nerve, inner ear, cerebellum, brain stem and cervical spine are common underlying factors.

Malignancy, primary and secondary, is a possibility in the elderly. The possibility of cardiac arrhythmias as a cause of syncopal symptoms increases with age.

Dizzy turns in elderly women

If no cause such as hypertension is found, advise them to get up slowly from sitting or lying, and to wear firm elastic stockings.

Acute vestibulopathy (vestibular failure)

Acute vestibulopathy covers both vestibular neuronitis and labyrinthitis, which are considered to be a viral infection of the vestibular nerve and labyrinth respectively, causing a prolonged attack of vertigo that can last for several days and be severe enough to require admission to hospital.[8]

DxT: acute vertigo + nausea + vomiting ° vestibular neuronitis

DxT: same symptoms + hearing loss ± tinnitus ° acute labyrinthitis

It is analagous to a viral infection of the 7th nerve causing Bell's palsy. The attack is similar to Meniere's syndrome except that there is no hearing disturbance. Characteristic features are:

- single attack of vertigo without tinnitus or deafness
- usually preceding ‘flu-like’ illness
- mainly in young adults and middle age
- abrupt onset with vertigo, nausea and vomiting
- generally lasts days to weeks
- examination shows nystagmus - rapid component away from side of lesion (no hearing loss)
- caloric stimulation confirms impaired vestibular function

It is basically a diagnosis of exclusion.

Treatment

- Rest in bed, lying very still.
- Gaze in the direction that eases symptoms.

The following drugs can be used:

- dimenhydrinate (Dramamine) 50 mg (o) 4 - 6 hourly *or*
- prochlorperazine (Stemetil) 12.5 mg IM (if severe) but may slow recovery or (recommended as best)
- diazepam (which decreases brain-stem response to vestibular stimuli)[2] 5 - 10 mg IM for the acute attack, then 5 mg (o) tds and possibly droperidol

A short course of corticosteroids often promotes recovery.

Outcome

Both are self-limiting disorders and usually settle over 5 - 7 days or several weeks. Labyrinthitis usually lasts longer and during recovery rapid head movements may bring on transient vertigo.

Benign paroxysmal positional vertigo

BPPV is a common type of acute vertigo that is induced by changing head position - particularly tilting the head backwards, changing from a recumbent to a sitting position or turning to the affected side.

Characteristic features:

- It affects all ages, especially the elderly.
- The female to male ratio is 2:1.
- It recurs periodically for several days.
- Each attack is brief, usually lasts 10 - 60 seconds, and subsides rapidly.
- Attacks are not accompanied by vomiting, tinnitus or deafness (nausea may occur).
- In one large series 17% were associated with trauma, 15% with viral labyrinthitis while about 50% had no clear predisposing factor other than age. One accepted theory of causation is that fine pieces of debris that are loose in the labyrinth settle on the bottom of the ear and generate endolymphatic movement.8 It may also be a variation of cervical dysfunction.
- Diagnosis is confirmed by head position testing. From a sitting position the patient's head is rapidly taken to a head-hanging position 30° below the level of the couch - do three times, with the head (1) straight, (2) rotated to the right, (3) rotated to the left. Hold on for 30 seconds and observe the patient carefully for vertigo and nystagmus. There is a latent period of few seconds before the onset of the symptoms - see Figure 46.3.
- Tests of hearing and vestibular function are normal.
- There is usually spontaneous recovery in weeks (most return to regular activity after 1 week).
- Recurrences are common: attacks occur in clusters.

Management

- Give appropriate explanation and reassurance.
- Avoidance measures: encourage the patient to move in ways that avoid the attack.
- Drugs are not recommended.
- Special exercises.
- Cervical traction may help.

Positional vestibular exercises

Most patients appear to benefit from exercise, such as the Brandt and Daroff procedure[9] or the Cawthorne - Cooksie exercises[10] that consist essentially of repeatedly inducing the symptoms of vertigo. Rather than resorting to avoidance measures, the patient is instructed to perform positional exercises to induce vertigo, hold this position until it subsides, and repeat this many times until the manoeuvre does not precipitate vertigo. The attacks then usually subside in a few days.

Surgical treatment

Rarely surgical treatment is required; it involves occlusion of the posterior semicircular canal rather than selective neurectomy.

MENIERE'S SYNDROME

This is caused by a build-up of endolymph.

- It is commonest in 30 - 50 years age group.
- It is characterised by paroxysmal attacks of:
    - vertigo
    - tinnitus
    - nausea and vomiting
    - sweating and pallor
    - deafness (progressive)
- Onset is abrupt - patient may fall and then be bedridden for 1 - 2 hours. Patient doesn't like moving head.
- Attacks last 30 minutes to several hours.
- There is a variable interval between attacks (twice a month to twice a year).
- Nystagmus is observed only during an attack (often to side opposite affected inner ear).
- Examination:
    - sensorineural deafness (low tones)
    - caloric test: impaired vestibular function

  - audiometry
    - sensorineural deafness
    - loudness recruitment
  - special tests
- There are characteristic changes in electrocochleography.

DxT: vertigo + vomiting + tinnitus ° Meniere's syndrome

Treatment

Acute attack[11]

Anticipation of attack (fullness, tinnitus):

- prochlorperazine 25 mg suppository
- 30 g urea crystals in orange juice (preferably 30 minutes before in prodromal phase)

Treatment:

- diazepam 5 mg IV ±prochlorperazine 12.5 mg IM
- consider betahistine 8 mg (o) tds if persistent or episodic

Long term

- Reassurance with a careful explanation of this condition to the patient, who often associates it with malignant disease
- Excess intake of salt, tobacco and coffee to be avoided
- A low-salt diet is the mainstay of treatment
- Alleviate abnormal anxiety by using stress management, meditation or possibly long-term sedation (fluid builds up with stress)
- Referral for neurological assessment
- Diuretic (e.g. hydrochlorothiazide/amiloride daily) - check electrolytes regularly

Surgery may be an option for intractable cases.

VESTIBULAR MIGRAINE

Migraine is a relatively common cause of vertigo and often unrecognised because of its many guises. It should be strongly suspected if there is a past and/or family history of migraine and also where there is a history of vertigo or ataxia that persists for hours or days in the absence of aural symptoms.[12] Vertigo, which is usually not violent, may take the place of the aura that precedes the headache or may be a migraine equivalent whereby the vertigo replaces the symptoms of headache. Pizotifen or propranolol are recommended for prophylaxis.

WHEN TO REFER

- Vertigo of uncertain diagnosis, especially in children
- Possibility of tumour, or bacterial infection
- Vertigo in presence of suppurative otitis media despite antibiotic therapy
- Presumed viral labyrinthitis not abating after 3 months
- Vertigo following trauma
- Presumed Meniere's syndrome, not responding to conservative medical management
- Evidence of vertebrobasilar insufficiency
- BPPV persisting for more than 12 months despite treatment with particle repositioning exercises

Practice tips

A careful drug history often pinpoints the diagnosis.

- Always consider cardiac arrhythmias as a cause of acute dizziness.
- Consider phenytoin therapy as a cause of dizziness in an epileptic patient.

- If an intracerebral metastatic lesion is suspected, consider the possibility of carcinoma of the lung as the primary source.
- Three important office investigations to perform in the evaluation are blood pressure measurement (lying, sitting and standing), hyperventilation and head positional testing.
- Cervical vertigo is very common and appropriate cervical mobilisation methods should be considered.
- BPPV is also common and prescribing a set of exercises to desensitise the labyrinth is recommended. Use either the Brandt and Daroff procedure or the Cawthorne - Cooksie program.[10]

REFERENCES

1. Kincaid-Smith P, Larkins R, Whelan G. *Problems in Clinical Medicine*. Sydney: MacLennan & Petty, 1989: 165.
2. Sloane PD, Slatt LM, Baker RM. *Essentials of Family Medicine*. Baltimore: Williams & Wilkins, 1988.
3. Sandler G, Fry J. *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986: 361 - 82.
4. Lance JW. *A Physiological Approach to Clinical Neurology* . London: Butterworths, 1970: 162 - 79.
5. Paine M. Dealing with dizziness. Aust Prescr, 2005; 28: 94 - 7.
6. Eviatar L, Eviatar A. Vertigo in children. Differential diagnosis and treatment. Paediatrics, 1977; 59: 833 - 7.
7. Tunnessen WW Jr. *Signs and Symptoms in Paediatrics*. Philadelphia: Lippincott, 1988: 591 - 4.
8. Waterson J. How to treat: Dizziness. Australian Doctor, 2003; 7 March: 1 - 8.
9. Brandt T, Daroff DB. Physical therapy for BPPV. Arch Otolaryngol, 1980; 106: 484 - 5.
10. Tiller J. *Therapeutic Guidelines: Neurology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2002: 55 - 63.
11. Tonkin JP. Meniere's disease. Curr Ther, 1995; 36: 39 - 43.
12. Pohl D. Vertigo. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 568 - 71.

CHAPTER 47. DYSPEPSIA (INDIGESTION)

Half the patients who get you up in the middle of the night and think they are dying are suffering from wind!

Francis Young (1884 - 1954), *Advice to a Younger Doctor*

DYSPEPSIA (INDIGESTION)

Dyspepsia or indigestion is a difficult, sometimes vague, symptom to define or evaluate and requires very careful questioning to clarify the exact nature of the complaint.

Dyspepsia embraces the following:

- nausea
- heartburn/regurgitation upper abdominal discomfort
- lower chest discomfort
- acidity
- epigastric fullness or unease
- abdominal distension

The discomfort can sometimes amount to pain. Diagnoses to consider in dyspeptic patients are summarised in Table 47.1.[1]

Table 47.1 Diagnoses to consider in dyspeptic patients

Gastrointestinal disorders
• Gastro-oesophageal reflux, including hiatus hernia
• Functional (non-ulcer) dyspepsia
• Oesophageal motility disorders (dysmotility)
• Peptic ulcer
• Upper GIT malignancies (e.g. oesophagus, stomach, pancreas)
• Hepatobiliary disease (e.g. hepatitis, biliary dyskinesia, cholelithiasis)
• Pancreatitis
• Upper GIT inflammation
- gastritis
- giardiasis
- Crohn's disease
• Irritable bowel syndrome
Non-gastrointestinal disorders
• Myocardial ischaemia
• Drug reaction
• Alcohol effect
• Somatisation
• Anxiety/stress
• Depression

Definitions

Dyspepsia

Pain or discomfort centred at the upper abdomen that is chronic or recurrent in nature.

Heartburn

A central retrosternal or epigastric burning sensation that spreads upwards to the throat.

Flatulence

Excessive wind. It includes belching, abdominal bloating or passing excessive flatus.[2]

FLATULENCE

Excessive belching (aerophagia)

- Usually functional
- Organic disease uncommon
- Due to air swallowing (aerophagy)
- Common in anxious people who gulp food and drink
- Associated hypersalivation

Management tips

- Make patient aware of excessive swallowing
- Avoid fizzy (carbonated) soft drinks
- Avoid chewing gum
- Don't drink with meals
- Don't mix proteins and starches
- Eat slowly and chew food thoroughly before swallowing
- Eat and chew with the mouth closed

If persistent: Simethicone preparation (e.g. Mylanta II, Phazyme)

If desperate: Place one small cork between the back teeth after meals for 30 minutes.

Excessive flatus

Flatus arises from two main sources:

- swallowed air
- bacterial fermentation of undigested carbohydrate

- malabsorption
- irritable bowel syndrome
- anxiety → aerophagy
- drugs, especially lipid-lowering agents
- lactose intolerance

Management:

- assess diet (e.g. high fibre, beans and legumes, cabbage, onions, grapes and raisins)
- avoid drinking with eating, especially with leafy vegetables
- cook vegetables thoroughly
- trial a lactose-free diet
- consider simethicone preparations (e.g. No Gas)

KEY FACTS AND CHECKPOINTS

- Dyspepsia or indigestion is a common complaint; 80% of the population1 will have experienced it at some time.
- Consider heartburn as ischaemic heart disease until proved otherwise.
- The presence of oesophagitis is suggested by pain on swallowing hot or cold liquids (odynophagia).
- All reflux is not due to hiatus hernia.
- Many patients with hiatus hernia do not experience heartburn.
- All patients with dysphagia must be investigated to rule out malignancy.
- Ten per cent of people in the community develop peptic ulcer (PU) disease.
- The major feature of PU disease is epigastric pain.
- The pain of duodenal ulcer (DU) classically occurs at night.

- At any time 10 - 20% of chronic NSAIDs users have peptic ulceration (greater than non-users).[3]
- NSAIDs mainly cause gastric ulcers (GU, gastric antrum and prepyloric region) with the duodenum affected to a lesser extent.
- Dyspeptic symptoms correlate poorly with NSAID-associated ulcer.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 47.2.

Table 47.2 Dyspepsia: diagnostic strategy model

Q. Probability diagnosis A. Irritable upper GIT (functional dyspepsia) Gastro-oesophageal reflux Oesophageal motility disorder (dysmotility)
Q. Serious disorders not to be missed A. Neoplasia • carcinoma: stomach, pancreas, oesophagus Cardiovascular • ischaemic heart disease • congestive cardiac failure Pancreatitis Peptic ulcer
Q. Pitfalls (often missed) A. Myocardial ischaemia Food allergy (e.g. lactose intolerance) Pregnancy (early) Biliary motility disorder Other gall bladder disease Post vagotomy Duodenitis *Rarities* • Hyperparathyroidism • Mesenteric ischaemia • Zollinger-Ellison syndrome • Kidney failure
Q. Seven masquerades checklist A. Depression Diabetes rarely Drugs Anaemia - Thyroid disorder - Spinal dysfunction - UTI -
Q. Is this patient trying to tell me something? A. Anxiety and stress are common associations of which patients are often unaware. Consider irritable bowel syndrome.

It is best to consider dyspepsia as:

- ulcer-like - localised pain

- dysmotility-like - diffuse discomfort, feeling full after meals (early satiety), nausea, bloating
- acid-reflux-like - indigestion or heartburn with acid reflux or regurgitation

The ulcer-like category may be due to an ulcer and if not is termed functional (non-ulcer) dyspepsia.

Pitfalls

Perhaps the commonest mistake is to attribute the discomfort of myocardial ischaemia to a disorder of the GIT. A sense of fullness or pressure in the epigastrium can certainly accompany ischaemia.

General pitfalls

- Reflux oesophagitis and peptic ulcer can mimic ischaemic heart disease.
- Overlooking gastric carcinoma as a cause of dyspepsia.
- Failing to stress that weight reduction to ideal level will generally alleviate gastro-oesophageal reflux.
- Overlooking drugs as a cause (Table 47.3).

Table 47.3 Drugs that may cause dyspepsia

Alcohol
Anticholinergics
Aspirin
Biphosphonates, esp. alendronate
Calcium-channel blockers
Corticosteroids
Digitalis
Lipid-lowering agents
Narcotics
Nicotine
NSAIDs
Potassium supplements (slow release)
Theophylline
Tricyclic antidepressants
Tetracycline

THE CLINICAL APPROACH

History

It is worthwhile spending some time clarifying the exact nature of the presenting complaint: what the patient means by ‘indigestion’ or ‘heartburn’.[4] The relationship of the symptom to eating is very important, and whether it occurs after each meal or after specific meals.

In particular, care should be taken to consider and perhaps exclude ischaemic heart disease.

Key questions

- How would you describe the discomfort?
- Can you show me exactly where it is and where it radiates?
- What makes your discomfort worse?
- What relieves your discomfort?
- What effect do food, milk and antacids have?
- What effect do coffee, onions or garlic have?
- What effect does a big meal have?
- What about drinking alcohol? Wine?
- What effect does exercise have?

- Do fried or fatty foods make it worse?
- Do hot spicy foods affect it?
- Does the problem come on at night soon after you go to bed?
- Does it wake you up at night?
- Does bending over (e.g. gardening) make it worse?
- Do you have periods of freedom from the problem?
- Are you under a lot of stress or have a lot of worry?
- Do you go flat out all day?
- Do you rush your meals?
- Do you chew your food properly?
- What drugs or medicines do you take?
- How much alcohol do you have? Do you smoke?
- Have you noticed anything else when you have the problem?
- Do you get constipated or have diarrhoea?
- Have you lost weight recently?
- Do you feel the discomfort between your shoulder blades, or in your shoulders or throat?

Symptoms analysis

Site and radiation

The site and radiation of pain or discomfort can provide a lead to the diagnosis.[5] Refer to Figure 40.10 at p. 418. If it is felt in the interscapular area consider oesophageal spasm, gall bladder disease or a DU. Retrosternal discomfort indicates oesophageal disorders or angina, while epigastric discomfort suggests disorders of the biliary system, stomach and duodenum.

Character of the pain

There tends to be considerable overlap in the character of the pain from the various disorders but some general characteristics apply:

- burning pain → gastro-oesophageal reflux (GORD)
- constricting pain → ischaemic heart disease or oesophageal spasm
- deep gnawing pain → PU
- heavy ache or ‘killing’ pain → psychogenic pain

Aggravating and relieving factors

Examples of these factors include:

- eating food may aggravate a GU but relieve a DU
- eating fried or fatty foods will aggravate biliary disease, functional dyspepsia and oesophageal disorders
- bending will aggravate GORD
- alcohol may aggravate GORD, oesophagitis, gastritis, PU, pancreatitis

Associated symptoms

Relevant examples:

- difficulty in swallowing → oesophageal disorders
- lump or constriction in throat → psychogenic
- acid regurgitation → GORD, oesophagitis
- anorexia, weight loss → carcinoma of the stomach
- water brash → GORD, hiatus hernia, PU
- symptoms of anaemia → chronic oesophagitis or gastritis, PU, carcinoma (stomach, colon)
- flatulence, belching, abnormal bowel habits → irritable bowel syndrome
- diarrhoea 30 minutes after meal → mesenteric ischaemia

Physical examination

The physical examination does not often provide the key to the diagnosis but it is important to perform very careful palpation and inspection. Look for evidence of clinical anaemia and jaundice. Diffuse mild abdominal tenderness and a pulsatable abdominal aorta are common findings but do not necessarily discriminate between organic and functional problems. Specific epigastric tenderness suggests peptic ulceration while tenderness over the gall bladder area (Murphy's sign) indicates gall bladder disease. An epigastric mass indicates carcinoma of the stomach.

Investigations

Do not overinvestigate. Investigations tend to be unrewarding in most instances of dyspepsia and could be postponed if the history is suggestive of a functional cause and the symptoms are not severe.[1] A trial of treatment such as changing adverse lifestyle factors, dietary modification and antacids could be the initial approach. Age is important in determining the extent of investigations, which are more relevant in those over 40 years.

The investigation of choice is endoscopy, which is superior to barium studies in investigation of the upper GIT. Gastroscopy is indicated for the *alarm* symptoms:

- abnormal symptoms of reflux/dyspepsia
- change of symptoms
- dysphagia
- unexplained weight loss
- GIT bleeding
- pain radiating to back
- pain waking at night
- abnormal signs on examination

Helicobacter pylori tests[6]

Helicobacter pylori has been proved to cause ulcers.

Non-invasive tests:

- serological - IgC antibodies (sensitivity 85 - 90%, specificity 90 - 99%); excellent for diagnosis, not for follow-up
- urea breath test (high sensitivity 97% and specificity 96%), good for follow-up
- Stool test (sensitivity 96%, specificity 97%) Invasive tests:
- mucosal biopsy during endoscopy can detect *H. pylori* through histology or rapid urease testing or *H. pylori* culture

DYSPEPSIA IN THE ELDERLY

An organic disorder is more likely in the older patient, in whom it is important to consider carcinoma of the stomach. Symptoms such as anorexia, vomiting and weight loss point to such a problem.

Other conditions causing dyspepsia that are more prevalent in this age group are:

- constipation
- mesenteric artery ischaemia
- congestive cardiac failure

DYSPEPSIA IN CHILDREN

Dyspepsia is an uncommon problem in children but can be caused by drugs, oesophageal disorders and gastro-oesophageal reflux in particular.[7] Reflux can be considered to be physiological or pathological.

Gastro-oesophageal reflux

Regurgitation of feeds because of gastro-oesophageal reflux is a common physiological event in newborn infants. A mild degree of reflux is normal in babies, especially after they burp; this condition is called *posseting*.

Symptoms

Milk will flow freely from the mouth soon after feeding, even after the baby has been put down for a sleep. Sometimes the flow will be forceful and may even be out of the nose.

Despite this vomiting or regurgitation, the babies are usually comfortable and thrive. Some infants will cry, presumably because of heartburn.[7]

In a small number the reflux may be severe enough (pathological) to cause serious problems such as oesophagitis with haematemesis or anaemia, stricture formation, failure to thrive, apnoea and aspiration.

Prognosis

Reflux gradually improves with time and usually ceases soon after solids are introduced into the diet. Most cases clear up completely by the age of 9 or 10 months, when the baby is sitting. Severe cases tend to persist until 18 months of age.

Investigations

These are not necessary in most cases but in those with persistent problems or complications referral to a paediatrician is recommended. The specialist investigations include barium meal with cine scanning, oesophageal pH monitoring or endoscopy and biopsy.

Management

Appropriate reassurance with parental education is important. It should be pointed out that changes in feeding practice and positioning will control most reflux.

The infant should be placed on the left side for sleeping with the head of the cot elevated about 10 - 20 degrees. The old bucket method, in which the child is placed in a bucket, is not necessary. Suspending the child in one of the new suspended swings for periods of 30 - 60 minutes after feeds will help.

Smaller, more frequent feeds and thickening agents are appropriate.

Thickening of feeds

Giving the baby thicker feeds usually helps those with more severe reflux. The old-fashioned remedy of using cornflour blended in bottles is still useful.

Bottle-fed babies (powdered milk formula)

Carobel: Add slightly less than 1 full scoop per bottle.

Gaviscon: Mix slightly less than ½ teaspoon of Infant Gaviscon Powder with 120 mL of formula in the bottle.

Cornflour (maize based): Mix 1 teaspoon with each 120 mL of formula. Check with your doctor or nurse for the proper method.

Karicare (or S26 AR): Very simple to use but more expensive.

Breastfed babies

Carobel: Add slightly less than 1 full scoop to 20 mL cool boiled water or 20 mL expressed breast milk and give just before the feed.

Gaviscon: Mix slightly less than ½ teaspoon of Infant Gaviscon Powder with 20 mL cool boiled water or expressed breast milk and give just after the feed.

For persistent or complicated reflux, including oesophagitis, specialist-monitored treatment will include the use of antacids, proton-pump inhibitors e.g. omeprazde, H_2-receptor blocking agents e.g. ranitidine.

GASTRO-OESOPHAGEAL REFLUX DISEASE (GORD)[8, 9]

Features

- Heartburn
- Acid regurgitation, especially lying down at night
- Water brash
- Diagnosis usually made on history
- Investigation usually not needed (reserve for danger signs and non-responsive treatment)

Complications

- Oesophagitis
- Iron-deficiency anaemia
- Stricture
- Respiratory: chronic cough, asthma, hoarseness

- Barrett's oesophagus (from prolonged reflux)

Barrett's oesophagus

- Usually a metaplastic response to prolonged reflux
- A premalignant condition (adenocarcinoma)
- Lower oesophagus lined with gastric mucosa (at least 3 cm)
- Prone to ulceration
- Needs careful management
- Consider 2 yearly endoscopies with biopsies

Management[8, 9, 10]

Stage 1

- Patient education/appropriate reassurance
- Consider acid suppression or neutralisation
- Attend to lifestyle:
    - weight reduction if overweight (this alone may abolish symptoms)
    - reduction or cessation of smoking
    - reduction or cessation of alcohol (especially with dinner)
    - avoid fatty foods (e.g. pastries, french fries)
    - reduction or cessation of coffee, tea and chocolate
    - avoid coffee and alcohol late at night
    - avoid gaseous drinks
    - leave at least 3 hours between the evening meal and retiring
    - increase fibre intake (e.g. high fibre cereals, fruit and vegetables)
    - small regular meals and snacks
    - eat slowly and chew food well
    - sleep on the left side
    - have main meal at midday with light evening meal
    - avoid spicy foods and tomato products
- Drugs to avoid
    - anticholinergics, theophylline, calciumchannel blockers, doxycycline. Pill-induced oesophagitis occurs, especially with tetracyclines, slow-release potassium, iron sulphate, corticosteroids, NSAIDs - avoid taking dry; use ample fluids
- Antacids (see Tables 47.4 and 47.5)
    - best is liquid alginate/antacid mixture e.g. Gaviscon/Mylanta plus 20 mL on demand or 1½ - 2 hours before meals and bedtime
- Elevation of head of bed or wedge pillow
    - if GORD occurs in bed, sleep with head of bed elevated 10 - 20 cm on wooden blocks or use a wedge pillow (preferable)

Table 47.4 Antacids in common use

Antacids
Water soluble Calcium carbonate Sodium
- bicarbonate
- citrotartrate
Note: Excess is prone to cause alkalosis - apathy, mental changes, stupor, kidney dysfunction, tetany
Water insoluble Aluminium
- hydroxide
- glycinate
- phosphate
Magnesium
- alginate
- carbonate

- hydroxide - trisilicate
Combination antacids Antacid + alginic acid Antacid + oxethazaine Antacid + simethicone

Table 47.5 Side-effects of common antacids

Aluminium hydroxide	- constipation
Magnesium trisilicate	- diarrhoea
Sodium bicarbonate	- alkalosis
Calcium carbonate	- milk alkali syndrome - alkalosis
	- constipation - milk alkali syndrome - hypercalcaemia

Stage 2 [8, 11]

If no relief after several weeks, the following approaches are recommended by the Gastroenterological Society of Australia (GESA).

Reduce acid secretion. Select from:

- H_2-receptor antagonists (oral use for 8 weeks)
  - cimetidine 400 mg bd pc or 800 mg nocte *or*
  - famotidine 40 mg nocte *or*
  - nizatadine 150 mg bd or 300 mg nocte *or*
  - ranitidine 150 mg bd pc or 300 mg nocte
- Proton-pump inhibitor (PPI) for 4 weeks
  - lansoprazole 30 mg mane *or*
  - omeprazole 20 - 40 mg mane *or*
  - pantoprazole 40 mg mane *or*
  - esomeprazole 30 mg mane *or*
  - rabeprazole 30 mg mane

Although the more traditional step-up approach of 1. Antacids → 2. H_2-receptor antagonists → 3. PPI can be used, there has been a change to favour a high level (more potent) initial therapy with PPIs at standard dose (a step-down approach; see Figure 47.1). This is based on the grounds of outcomes, speed of response and total cost.


Step up
Proton-pump inhibitor (PPI)
H2-receptor antagonist
lifestyle modification ± antacids
Step down

Figure 47.1 The stepwise approach to the management of dyspeptic symptoms

[9]

Surgery is usually for young patients with severe reflux. The gold standard is a short loose 360-degree fundoplication.

FUNCTIONAL (NON-ULCER) DYSPEPSIA[11]

This term applies to the 60% of patients presenting with dyspepsia in which there is discomfort on eating in the absence of demonstrable organic disease. This can be considered in two categories (although there is overlap):

- ulcer-like dyspepsia *or*
- dysmotility-like dyspepsia

Ulcer-like dyspepsia

Treat as for GORD. A practical approach is to commence with a 4-week trial of an H_2-receptor antagonist and cease if symptoms resolve. If there is a failed response, there is little evidence to suggest that a PPI will improve response.[9, 12]

Dysmotility-like dyspepsia

Features of dysmotility

- Discomfort with early sense of fullness on eating
- Nausea
- Overweight
- Emotional stress
- Poor diet (e.g. fatty foods)
- Similar lifestyle guidelines to GORD

Management

- Treat as for GORD (stage 1).
- Include antacids.
- If not responsive:
  - *Step* 1: H_2-receptor antagonists
  - *Step* 2: prokinetic agents domperidone 10 mg tds *or* metoclopramide 10 mg tds

PEPTIC ULCER DISEASE[6, 13]

General features

- Common: 10 - 20% incidence over a lifetime
- Point prevalence of ulcer disease: 3 - 5%
- DU:GU ° 4:1
- DUs common in men 3:1
- Cumulative mortality of 10%
- Risk factors
  - male sex
  - family history
  - smoking (cause and delayed healing)
  - stress
  - common in blood group O
  - NSAIDs 2 - 4 times increase in GU and ulcer complications
  - *H. pylori*: if absent and no NSAIDS, no ulcer
- Unproven risk factors
  - corticosteroids
  - alcohol (except for gastric erosion)
  - diet (does reduce recurrence of PU)
- Types of ulcers

  - lower oesophageal
  - gastric
  - stomal (post gastric surgery)
  - duodenal

Clinical features

- Episodic burning epigastric pain related to meals (1 - 2 hours after)
- Relieved by food or antacids (generally)
- Dyspepsia common
- May be 'silent' in elderly on NSAIDs
- Physical examination often unhelpful

Investigations

- Endoscopy (investigation of choice)14 - 92% predictive value
- Barium studies - 54% predictive value
- Serum gastrin (consider if multiple ulcers)
- *H. pylori* test: serology or urea breath test; diagnosis usually based on urease test performed at endoscopy

Complications

- Perforation
- Bleeding → haematemesis and melaena
- Obstruction - pyloric stenosis
- Anaemia (blood loss)
- Carcinoma (in GU)
- Oesophageal stenosis

Bleeding peptic ulcer

This can be treated with endoscopic haemostasis with heater probe or injection of adrenaline or both. Also IV omeprazole 80 mg bolus, then 8 mg/hr IV infusion for 3 days. Surgery is an option.

Management

Aims of treatment:

- relieve symptoms
- accelerate ulcer healing
- prevent complications
- minimise risk of relapse

The treatment of a GU is similar to that for a DU except that GUs take about 2 weeks longer to heal and the increased risk of malignancy has to be considered.

Stage 1: Treatment

General measures:

- same principles as for GORD
- stop smoking
- avoid irritant drugs
  - NSAIDs
  - aspirin
- normal diet but avoid foods that upset
- antacids

Stage 2: H_2-receptor antagonists (refer p. 516)

- 8-week oral course

Use with caution in:

- the elderly
- those on drugs, especially warfarin, anticonvulsants, beta-blockers
- liver disease

Outcome[14, 15]

- DU: 80% healing rate
  - prone to spontaneous remissions and relapse
  - once healed:
    - 60% relapse within 1 year
    - 20% persist but asymptomatic
- GU: 50% do not recur other 50% recurrence within 2 years

Note: Use maintenance or continuous treatment for two or more exacerbations in 12 months.

Proton-pump inhibitors

See p. 516.

PPIs provide more potent acid suppression and heal GUs and DUs more rapidly than H_2-receptor antagonists

Other possible agents

Cytoprotective agents

- Sucralfate 1 g tab (o) qid, 1 hour ac and nocte

Prostaglandin analogue

- Misoprostol 800 μg daily (divided doses)

Colloidal bismuth subcitrate (CBS)

- bismuth subcitrate (De-Nol) 2 tabs (chewed) bd for 6 - 8 weeks
  - effective for relapsing ulcers
  - appears to be effective against *H. pylori*

Therapy to eradicate Helicobacter pylori

This organism has a proven link with PU disease (both DU and benign non-drug induced GU), gastric carcinoma and maltoma (a gastric lymphoma) because of mucosal infection. This hypothesis is supported by a very low relapse of DU in subjects eradicated of *H. pylori*. Treatment is based on combination triple or quadruple therapy.

Treatment regimens (examples)

1. First-line therapy PPI (e.g. omeprazole 20 mg) + clarithromycin 500 mg + amoxycillin 1 g (all orally twice daily for 7 days) *or*
2. PPI + amoxycillin + metronidazole (all twice daily for 7 days) *or*
3. PPI + clarithromycin + metronidazole (twice daily for 7 days) - if hypersensitive to penicillin *or* other combinations, e.g. bismuth + PPI + tetaracycline + metronidazole (for failed triple combination)

Note: Resistance to metronidazole is common and to clarithromycin is increasing but uncommon with tetracycline and amoxycillin.[6]

Surgical treatment

Indications (now uncommon) include:

- failed medical treatment after 1 year
- complications
  - uncontrollable bleeding
  - perforation
  - pyloric stenosis
- suspicion of malignancy in GU
- recurrent ulcer after previous surgery

NSAIDS AND PEPTIC ULCERS[6, 13]

1. *Ulcer identified in NSAID user*:
    - stop NSAID (if possible)
    - check smoking and alcohol use
    - try alternative anti-inflammatory analgesic:
        - paracetamol
        - enteric-coated, slow-release aspirin
        - corticosteroids intra-articular or oral
    - H_2-receptor antagonist (full dose) for 4 - 6 weeks *or* misoprostol 800 µg a day (used for GU) *or* PPI for 2 - 4 weeks (gives best results)

    Note: Healing time is doubled if NSAID continued.[3] About 90% heal within 12 weeks. Check healing by endoscopy at 12 weeks.
2. *Prevention of ulcers in NSAID user*: Try alternatives (as above). Prophylactic drugs are rarely justified but reasonable in those over 75 years and in those with a past history of PU.
    - misoprostol (prevents GU recurrence)
    - H_2-receptor antagonist (prevents DU, not GU)

Increased dietary fibre assists DU healing and prevention.

Note: If *H. pylori* is present, it should be eradicated with combination therapy.

CARCINOMA OF STOMACH

Clinical features

- Male to female ratio ° 3:1
- Usually asymptomatic early
- Consider if upper GIT symptoms in patients over 40 years, especially weight loss
- Recent-onset dyspepsia in middle age
- Dyspepsia unresponsive to treatment
- Vague fullness or epigastric distension
- Anorexia, nausea, ± vomiting
- Dysphagia - a late sign
- Onset of anaemia
- Changing dyspepsia in GU
- Changing symptoms in pernicious anaemia
- *H. pylori* now implemented as a cause

Risk factors: ↑ age, blood group A, smoking, atropic gastritis

Limited physical findings

- Palpable abdominal mass (20%)
- Signs (Figure 47.2) in advanced cases


anaemia
enlarged supraclavicular lymph node (Troisier's sign)
hepatomegaly —hard and irregular
epigastric mass

Figure 47.2 Late signs of carcinoma of the stomach

Investigations

- Endoscopy and biopsy is optimal test
- Barium meal - false negatives

Treatment

- Surgical excision: may be curative if diagnosed early but overall survival is poor.

WHEN TO REFER

- Infants with persistent gastro-oesophageal reflux not responding to simple measures
- Failure to respond to stage 1 therapy for heartburn, when endoscopy is required
- Patients with persistent or recurrent ulcers
- Any patient with a PU complication, such as haemorrhage, obstruction or perforation

Practice tips

- Scleroderma is a rare but important cause of oesophagitis.

- Advise patients never to ‘dry swallow’ medications.
- Dysphagia always warrants investigation, not observation.
- Beware of attributing anaemia to oesophagitis.
- Epigastric pain aggravated by any food, relieved by acids ° chronic GU.
- Epigastric pain before meals, relieved by food ° chronic DU.
- Keep in mind the malignant potential of a GU.
- A change in the nature of symptoms with a GU suggests the possibility of malignant change.
- Avoid the long-term use of water-soluble antacids.
- Investigate the alarm symptoms - dysphagia, bleeding, anaemia, weight loss, waking at night, pain radiating to the back.

REFERENCES

1. Smallwood R. Dyspepsia. Med Observer, 1991; 27 September: 33 - 4.
2. Holvey DN. *The Merck Manual* (12th edn). Rahway: Merck Sharp & Dohme, 1975: 679 - 82.
3. Pritchard P. The management of upper gastrointestinal problems in patients taking NSAIDs. Aust Fam Physician, 1991; 20: 1739 - 41.
4. Kincaid-Smith P, Larkins R, Whelan G. *Problems in Clinical Medicine*. Sydney: MacLennan & Petty, 1989: 207 - 12.
5. Sandler G, Fry J. *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986: 99 - 135.
6. McGarity B, Morgia M. Peptic ulcer disease: an update on diagnosis and treatment. Med Today, 2001; December: 33 - 7.
7. Sewell J. Gastro-oesophageal reflux. Aust Paediat Rev, 1991; 3: 2.
8. *Gastro-Oesophageal Reflux Disease in Adults: Guidelines* (3rd edn). Sydney: Gastroenterological Society of Australia, 2001.
9. *GORD and Non-Ulcer Dyspepsia*. NPS News, 2001; PPR review No. 11.
10. Katelaris P. Dyspepsia: update. Australian Doctor, 2005; 7 October: 23 - 5.
11. Birkitt D. (Chair). *Therapeutic Guidelines*: *Gastrointestinal* (version 3). Melbourne: Therapeutic Guidelines Ltd, 2002: 39 - 50.
12. Tolley NJ et al. Gastroenterology, 1998; 114: 582 - 95.
13. Madge S, Yeomans N. Stomach and duodenal ulcers. Curr Ther, 2001; September: 69 - 72.
14. Korman M, Sievert W. Peptic ulcers. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 400 - 2.
15. Sandler G. *Common Medical Problems*. London: ADIS, 1984: 139 - 76.

CHAPTER 48. DYSPHAGIA

We swallow approximately 1200 times daily, largely subconsciously. While we take the fundamental function for granted, disordered swallowing can be a devastating condition, with substantial morbidity for those affected.

IAN COOK 1996

DYSPHAGIA

Dysphagia is difficulty in swallowing. It is usually associated with a sensation of hold-up of the swallowed bolus and is sometimes accompanied by pain.

Its origin is considered as either oropharyngeal or oesophageal. Oropharyngeal dysphagia is usually related to neuromuscular dysfunction and is commonly caused by stroke. Oesophageal dysphagia is usually due to motor disorders, such as achalasia or diffuse oesophageal spasm, and to peptic oesophageal strictures often secondary to reflux. In this type of dysphagia there is a sensation of a hold-up, which may be experienced in either the cervical or retrosternal region.1 Causes are usually classified as functional, mechanical and neurological (Table 48.1).

Table 48.1 Causes of dysphagia

Functional	e.g. muscle tension, 'express swallowing'
Neurological	e.g. stroke, myasthenia, MND
Mechanical	
Luminal	e.g. foreign body
Mural	e.g. stricture, tumour
Extramural	e.g. extrinsic compression (i.e. goitre)

Dysphagia must not be confused with globus sensation, which is the sensation of the constant 'lump in the throat' although there is no actual difficulty swallowing food.

There are only a few common causes of dysphagia and these are usually readily diagnosed on the history and two or three investigations. A careful history is very important, including a drug history and psychosocial factors.

DIAGNOSTIC GUIDELINES

- Any disease or abnormality affecting the tongue, pharynx or oesophagus can cause dysphagia.
- Patients experience a sensation of obstruction at a definite level with swallowing food or water; hence, it is convenient to subdivide dysphagia into oropharyngeal and oesophageal.
- Pain from the oropharynx is localised to the neck.
- Pain from the oesophagus is usually felt over the T2 - 6 area of the chest.
- Oropharyngeal causes: difficulty initiating swallowing; food sticks at the supersternal notch level; regurgitation; aspiration.
- Oesophageal causes: food sticks to mid to lower sternal level; pain on swallowing solid foods, especially meat, potatoes and bread, and then eventually liquids.
- A pharyngeal pouch usually causes regurgitation of undigested food and gurgling may be audible over the side of the neck.
- Neurological disorders typically result in difficulty swallowing or coughing or choking due to food spillover, especially with liquids.
- Dysphagia for solids only indicates a structural lesion, such as a stricture or tumour.
- Dysphagia for liquids and solids is typical of an oesophageal motility disorder, namely achalasia.[2]

- GORD tends to exclude achalasia.
- Gastroenterologists claim that the big three common causes referred to them are benign peptic stricture, carcinoma and achalasia.[3]
- Intermittent dysphagia for both liquids and solids is characteristic of a motility disorder such as oesophageal achalasia.
- Malignant oesophageal obstruction is usually evident when there is a short history of rapidly progressive dysphagia and significant weight loss.[4]

A summary of the safety diagnostic model is presented in Table 48.2.

Table 48.2 Dysphagia: diagnostic strategy model (excluding oropharyngeal infections and strokes)

Q. Probability diagnosis
A. Functional (e.g. 'express' swallowing, psychogenic) Tablet-induced irritation Pharyngo-tonsillitis Reflux oesophagitis
Q. Serious disorders not to be missed
A. Neoplasia - carcinoma pharynx, oesophagus, stomach - extrinsic tumour AIDS (opportunistic oesophageal infection) Stricture, usually benign peptic stricture Scleroderma Neurological causes - pseudobulbar palsy - multiple sclerosis - motor neurone disease (amyotrophic sclerosis) - Parkinson's disease
Q. Pitfalls (often missed)
A. Foreign body Drugs (e.g. phenothiazines) Subacute thyroiditis Extrinsic lesions (e.g. lymph nodes, goitre) Upper oesophageal web (e.g. Plummer-Vinson syndrome) Eosinophilic oesophagitis Radiotherapy Achalasia Upper oesophageal spasm (mimics angina) *Rarities* (some) • Sjögren's syndrome • Aortic aneurysm • Aberrant right subclavian artery • Lead poisoning • Cervical osteoarthritis (large osteophytes) • Other neurological causes • Other mechanical causes
Q. Seven masquerades checklist
A. Depression Diabetes -

Drugs

Anaemia -

Thyroid disorder

Spinal dysfunction -

UTI -

Q. Is this patient trying to tell me something?

A. Yes. Could be functional? Globus sensation.

Examination

It is worthwhile focusing on the following features:

- general examination including hands and skin
- mouth, pharynx, larynx (look for paralysis)
- neck, especially for lymph nodes and thyroid
- neurological, especially cranial nerve function and muscle weakness disorders
- special oesophageal obstruction test:
  - hand the patient a glass of water and place a stethoscope over the left upper quadrant of abdomen
  - measure time between swallowing and murmur produced by bolus passing the cardia (normal: 7 - 10 seconds)

Investigations

- Full blood examination: ? anaemia
- Neurological cause: oesophageal motility study (manometry)
- Mechanical:
  - extrinsic compression (e.g. barium swallow, CT scan, chest X-ray)
  - intrinsic (e.g. endoscopy ± barium swallow)

The primary investigation in suspected pharyngeal dysphagia is a video barium swallow[5] while endoscopy is generally the first investigation in cases of suspected oesophageal dysphagia. Barium swallow should precede endoscopy in the latter when there is a suspected oesophageal ‘ring’ and suspected oesophageal dysmotility. If endoscopy and radiology are negative, consider oesophageal motility studies to look specifically for achalasia or other less common motility disorders.

SPECIFIC CONDITIONS

Benign peptic stricture

- Fibrous stricture of lower third oesophagus (can be higher)
- Follows years of reflux oesophagitis
- Usually older patients
- Dysphagia with solid food
- Diagnosis confirmed by endoscopy and barium swallow

Treatment

- Dilate the stricture.
- Treat reflux vigorously.

Carcinoma of oesophagus

- Dysphagia at beginning of meal

- Dysphagia for solid food steadily progressive over weeks
- Can remain silent and tend to be invasive when diagnosed
- Discomfort or pain - throat, retrosternal, interscapular
- Weight loss can be striking
- Diagnosis confirmed by barium swallow and endoscopy
- Both SCC (commonest) and adenocarcinoma
- Adenocarcinoma associated with Barrett's mucosa
- Treatment is usually palliative surgery

Achalasia

- A disorder of oesophageal motility
- Widely dilated oesophagus
- Empties poorly through a smoothly tapered lower end
- Gradual onset of dysphagia for both liquids and solids
- Fluctuating symptoms
- Diagnosis confirmed by barium swallow or manometry
- Manometry is the only way to diagnose with certainty[1]

Treatment

- Conservative in the elderly (e.g. nifedipine/or endoscopic botulinum toxin injection into the sphincter)[6]
- Pneumatic dilatation of lower oesophageal sphincter or surgical myotomy

Note: Prokinetic drugs have no place in treatment.

Drug-induced oesophageal ulceration[3]

- Tetracycline, especially doxycycline, can cause painful ulceration in all age groups.
- Delayed passage of some drugs (due to pre-existing disorders) can cause local ulceration, even perforation (especially in elderly) (e.g. iron tablets, slow-release potassium, aspirin, NSAIDs, biphosphonates, zidovudine, antibiotics).
- The elderly are prone to the problem if they ingest drugs upon retiring to bed with insufficient liquid washdown.

Globus sensation

Also referred to as ‘globus hystericus’ or ‘lump in the throat’, it is the subjective sensation of a lump in the throat. It appears to be associated with psychological stress (e.g. unresolved hurt, grief, non-achievement). Suppression of sadness is most often implicated.[6] No specific aetiology or physiological mechanism has been established. The symptom can be associated with GORD, from frequent swallowing or emotionally based dry throat.

Features

- Sensation of being ‘choked up’ or ‘something stuck’ or lump
- Not affected by swallowing
- Eating and drinking may provide relief
- Normal investigations

Approach to patient

- Careful history and examination
- Exclude organic cause (refer Table 48.2)
- May require investigations if doubtful diagnosis

Management

- Usually settles with education, reassuring support and time (up to several months)
- No drug of proven value
- Treat any underlying psychological disorder

Odynophagia

Pain on swallowing is basically caused by irritation of an inflammed or ulcerated (in particular) mucosa by the swallowed food bolus.

Important casues include:

- GORD (the commonest cause) with associated oesophagitis
- oesophageal candidiasis, especially in the immunosuppressed
- herpes simplex oesophagitis, also in the immunosuppressed
- cytomegalovirus oesophagitis, in the immunosuppressed
- pill-induced oesophagitis/ulceration
- carcinoma of the oesophagus

Eosinophilic oesophagitis[7]

Eosinophilic oesophagitis is increasingly being recognised as a cause of dysphagia and acute food bolus obstruction in both children (particularly) and adults. It should be considered in those who regularly experience food getting stuck in their throat. It is associated with allergic disorders such as hay fever and asthma. Refer to gastroscopy. Treatment of the acute attack includes IM buscopan and a swallowed topical corticosteroid aerosol e.g. fluticasone.

Practice tips

- Although dysphagia is a common psychogenic symptom it must always be taken seriously and investigated.
- Mechanical dysphagia represents carcinoma until proved otherwise.
- Progressive dysphagia and weight loss in an elderly patient is oesophageal carcinoma until proved otherwise.
- Carcinoma of the oesophagus usually causes pain, wasting and regurgitation.
- Globus sensation or hystericus, an anxiety disorder, should not be confused with dysphagia. It is the subjective sensation of a lump or mass in the throat. Usually seen in young women.
- Carcinoma-induced achalasia occurs with tumours at the gastro-oesophageal junction usually due to adenocarcinoma of the stomach.
- Severe oesophageal reflux predisposes to adenocarcinoma.
- Oesophageal strictures can be benign, usually secondary to chronic reflux oesophagitis, or due to malignancy.
- Be careful of a change of symptoms in the presence of long-standing reflux. Consider stricture or carcinoma.

REFERENCES

1. Tally NJ, Martin CJ. *Clinical Gastroenterology: A Practical Problem Based Approach.* Sydney: MacLennan & Petty, 1996: 31 - 2.
2. Trate DM, Parkman HP, Fisher RS. Dysphagia. Evaluation, diagnosis and treatment. Primary Care, 1996; 23: 417 - 32.
3. Breen K. A practical approach to patients with dysphagia or pain on swallowing. Mod Med Aust, 1992; 3: 50 - 6.
4. Abeygunasekera S. Difficult and painful swallowing: a guide for GPs. Medicine Today, 2003; 4(10): 33 - 40.
5. Cook I. Swallowing disorders. Curr Ther, 1996; 37: 81 - 5.
6. Beers M, Berkow R eds. *The Merck Manual* (18th edn). Merck Research Laboratories, 2006: 69 - 70.
7. Kakakios A, Heine R. Eosinophilic oesophagitis. MJA. 2006; 185: 7: 401.

CHAPTER 49. DYSPNOEA

When man grows old … there is much gas within his thorax, resulting in panting and troubled breathing.

HUANG TI (2697-2597 BC), THE YELLOW EMPEROR'S CLASSIC OF INTERNAL MEDICINE

DYSPNOEA

Dyspnoea is the subjective sensation of breathlessness that is excessive for any given level of physical activity. It is a cardinal symptom affecting the cardiopulmonary system and can be very difficult to evaluate. Appropriate breathlessness following activities such as running to catch a bus or climbing several flights of stairs is not abnormal but may be excessive due to obesity or lack of fitness.

KEY FACTS AND CHECKPOINTS

- Determination of the underlying cause of dyspnoea in a given patient is absolutely essential for effective management.
- The main causes of dyspnoea are lung disease, heart disease, obesity and functional hyperventilation.[1]
- The most common cause of dyspnoea encountered in family practice is airflow obstruction, which is the basic abnormality seen in chronic asthma and chronic obstructive pulmonary disease (COPD).[2]
- Wheezing, which is a continuous musical or whistling noise, is an indication of airflow obstruction.
- Some patients with asthma do not wheeze and some patients who wheeze do not have asthma.
- Other important pulmonary causes include restrictive disease, such as fibrosis, collapse and pleural effusion.
- Dyspnoea is not inevitable in lung cancer but occurs in about 60% of cases.[3]
- Normal respiratory rate is 12 - 16 breaths/minute.

TERMINOLOGY

It is important to emphasise that dyspnoea or breathlessness is a subjective sensation of the desire for increased respiratory effort and must be considered in relation to the patient's lifestyle and individual tolerance of discomfort. It also depends on the age, physical fitness and physical expectations of the person. Patients may complain of tightness in the chest and this must be differentiated from angina.

The New York Heart Association functional and therapeutic classification applied to dyspnoea is:

- Grade 1 No breathlessness
- Grade 2 Breathlessness on severe exertion
- Grade 3 Breathlessness on mild exertion
- Grade 4 Breathlessness at rest

Definitions

Orthopnoea Breathlessness lying down flat.

Paroxysmal nocturnal dyspnoea Inappropriate breathlessness causing waking from sleep.

Tachypnoea An increased rate of breathing.

Hyperpnoea An increased level of ventilation (e.g. during exertion).

Hyperventilation Overbreathing.

DIFFERENCE BETWEEN HEART AND LUNG CAUSES

The distinguishing features between dyspnoea due to heart disease and to lung disease are presented in Table 49.1.

Table 49.1 Comparison of distinguishing features between dyspnoea due to heart disease and lung disease

Lung disease	Heart disease
History of respiratory disease	History of hypertension, cardiac ischaemia or valvular heart disease
Slow development	Rapid development
Present at rest	Mainly on exertion
Productive cough common	Cough uncommon and then ‘dry’
Aggravated by respiratory infection	Usually unaffected by respiratory infection

Source: After Sandler[1]

The history is a good indication and a useful guideline is that dyspnoea at rest is typical of lung disease, especially asthma, while it tends to be present on effort with heart disease as well as with COPD.

WHEEZING

Wheezing is any continuous musical expiratory noise heard with the stethoscope or otherwise. Wheeze includes stridor, which is an inspiratory wheeze.

Common causes of wheezing

Localised:

- partial bronchial obstruction
  - impacted foreign body
  - impacted mucus plugs
  - extrinsic compression

Generalised:

- asthma
- obstructive bronchitis
- bronchiolitis

‘CARDIAC ASTHMA’ AND BRONCHIAL ASTHMA

The term ‘cardiac asthma’ is used to describe a wheezing sensation such as that experienced with paroxysmal nocturnal dyspnoea. Differentiating features are presented in Table 49.2.

Table 49.2 Comparison of distinguishing features between ‘cardiac asthma’ and bronchial asthma

	Cardiac	Bronchial
Dyspnoea	Mainly inspiratory	Mainly expiratory
Cough	Follows dyspnoea	Precedes dyspnoea
Sputum	Pink and frothy	Thick and gelatinous
Relief	Standing up (by an open window) Intravenous diuretic/CPAP morphine	Coughing up sputum Bronchodilator
Lung signs	Mainly crackles	Mainly wheezes

Source: After Sandler[1]

Is it asthma or COPD?

This question is often asked, especially in the middleaged or elderly person with dyspnoea. Differentiating features are presented in Table 49.3.

A DIAGNOSTIC APPROACH

A summary of the diagnostic strategy model is presented in Table 49.4.

Probability diagnosis

The common causes of dyspnoea are lung disease, heart disease, obesity, anaemia (tissue hypoxia) and functional hyperventilation. More specifically, bronchial asthma, COPD, acute pulmonary infections and left heart failure (often insidious) are common individual causes.

Serious disorders not to be missed

Severe cardiovascular events such as acute heart failure, which may be precipitated by myocardial infarction (may be silent, especially in diabetics), a life-threatening arrhythmia, pulmonary embolism, dissecting aneurysm or a cardiomyopathy (such as viral myocarditis) require early diagnosis and corrective action. Recurrent pulmonary embolism may present a diagnostic problem. There may be a history of deep venous thrombosis, pregnancy, malignancy or taking the contraceptive pill.[4]

Table 49.3 Comparison of asthma and COPD

	Asthma	COPD
Symptoms < 35 years	Common	Unusual
Smoking history	Possible	Invariable
Chronic cough	Uncommon	Common
Dyspnoea	Diurnal and variable	Constant and progressive
Response to inhaled bronchodilator	Good	Poor
Nocturnal waking with symptoms	Common	Uncommon

Table 49.4 Dyspnoea: diagnostic strategy model

Q. Probability diagnosis
A. Bronchial asthma Bronchiolitis (children) Left heart failuer COPD Obesity Lack of fitness
Q. Serious disorders not to be missed
A. Cardiovascular • acute heart failure (e.g. AMI) • arrhythmia • pulmonary embolism • pulmonary hypertension • dissecting aneurysm • cardiomyopathy

• pericardial tamponade • anaphylaxis Neoplasia • bronchial carcinoma Severe infections • SARS • avian influenza • pneumonia • acute epiglottitis (children) Respiratory disorders • inhaled foreign body • upper airways obstruction • pneumothorax • atelectasis • pleural effusion • tuberculosis Acute respiratory distress syndrome Neuromuscular disease • infective polyneuritis • poliomyelitis
Q. Pitfalls (often missed)
A. Interstitial lung disorder • fibrosing alveolitis • extrinsic allergic alveolitis Chemical pneumonitis Metabolic acidosis Radiotherapy Kidney failure (uraemia) Multiple small pulmonary emboli
Q. Seven masquerades checklist
A. Depression ✔ Diabetes ✔ Drugs ✔ Anaemia ✔ Thyroid disorder (thyrotoxicosis) ✔ Spinal dysfunction (ankylosing spondylitis) UTI -
Q. Is the patient trying to tell me something?
A. Consider functional hyperventilation (anxiety and panic attacks).

Severe infections such as lobar pneumonia, tuberculosis and myocarditis must be considered. In children acute epiglottitis, croup, bronchiolitis, pneumonia and bronchitis are serious infections responsible for respiratory distress.

Primary carcinoma is an important consideration, especially in dyspnoea of gradual onset. Other malignant conditions to consider are metastases, lymphangitis carcinomatosis, lymphomas and pleural mesothelioma. Pleural effusion may be the mode of presentation of some of these serious disorders.

Pitfalls

Interstitial pulmonary disease can be a diagnostic dilemma because the physical signs and X-ray appearances can be minimal in the early stages despite the presence of significant dyspnoea. Allergic alveolitis, such as that caused by birds (e.g. hypersensitivity to their droppings) can be a pitfall. The diagnosis is easier if a known disease associated with pulmonary infiltration, such as sarcoidosis, is present. Measuring the diff using capacity will help with diagnosis.

Pericardial tamponade may cause difficulty in diagnosis either with an acute onset, such as malignancy involving pericardium or insidiously. The patient usually has a weak pulse with pulsus paradoxus, hypotension and a raised jugular venous pressure.

It is important to be careful not to attribute dyspnoea simply to obesity or lack of fitness when it could have a true organic disorder such as heart failure.

Seven masquerades checklist

Most of the masquerades have to be considered as underlying causes. Depression can be associated with dyspnoea, anaemia is an important cause of dyspnoea, thyrotoxicosis can rarely present with dyspnoea, and diabetic ketoacidosis can cause rapid deep breathing.

Drugs must also be considered, especially as a cause of interstitial pulmonary fibrosis that presents with dyspnoea, cough and fever. Drugs that cause this disorder include several cytotoxic agents (especially bleomycin, cyclophosphamide, methotrexate), amiodarone, sulphasalazine, penicillamine, nitrofurantoin, gold salts and adrenergic nasal sprays.[3] Poisons that may cause hyperventilation are salicylate, methyl alcohol, theophylline overdosage and ethylene glycol. Anaemia must be considered especially in those at risk. Dyspnoea is unlikely to be caused solely by chronic anaemia unless the haemoglobin level is less than 8 g/dL.[4] It is more likely to occur if another predisposing cause, such as ischaemic heart disease, is present.

Psychogenic considerations

Functional dyspnoea or hyperventilation is common. However, it is important to exclude organic causation such as asthma, drugs and thyrotoxicosis before settling with the psychogenic label and to reassure the patient strongly if there is no organic cause. Any uncomfortable sensation in the chest may be interpreted as dyspnoea by anxious patients. Depression, anxiety and panic attacks may be underlying the problem. Characteristic associated features of hyperventilation with anxiety include dizziness, faintness, palpitations, yawning, paraesthesia of the hands and legs, inability to take a deep breath or a sensation of smothering. These patients may exhibit sighing and irregular breathing on examination. In true psychogenic dyspnoea, chest X-rays and pulmonary function tests are normal but symptoms are often reproduced after 15-30 seconds of voluntary hyperventilation. It is important to remember that it may be present in a patient who has organic disease of a mild degree such as asthma.

THE CLINICAL APPROACH

History

Special attention should be paid to evaluating exactly what the patient means by breathlessness or restriction of breathing. The analysis should then include provoking factors and associated symptoms with a view to differentiating between pulmonary causes such as asthma and COPD. Wheeze is often (but not always) present in asthma and chronic airflow obstruction. Most respiratory causes of dyspnoea also produce cough. The rate of development of dyspnoea gives an indication of the possible cause (Table 49.5).[5] The sudden onset of dyspnoea at rest is suggestive of pulmonary embolism or pneumothorax. Severe dyspnoea developing over 1 or 2 hours is most likely due to left heart failure or bronchial asthma. Bronchial asthma is usually easily distinguished from left heart failure by the history of previous attacks, by the absence of chest pain and the absence of cardiac murmurs. ‘My breathing feels tight’ indicates asthma. A complaint of ‘suffocation or feeling smothered’ or ‘just not getting enough air’ may be a pointer to functional dyspnoea.

Table 49.5 Typical causes of dyspnoea related to time of onset

Sudden • Lung collapse • Inhaled foreign body/other choking
• Spontaneous pneumothorax
• Arrhythmia
• Anaphylaxis
• Myocardial infarction
• Pulmonary embolism
Rapid (over a few hours)
• Asthma • Hyperventilation (can be sudden) • Acute exacerbations of COPD • Pneumonia • Diabetic ketoacidosis • Extrinsic allergic alveolitis • High altitude • Left heart failure (acute pulmonary oedema) • Pericardial tamponade • Poisons **Over days or weeks**
• Congestive heart failure • Pleural effusion • Carcinoma of the bronchus/trachea **Over months or years** • COPD • Tuberculosis • Fibrosing alveolitis • Pneumoconiosis **Non-respiratory causes**
• Anaemia
• Hyperthyroidism
• Obesity

The dyspnoea of asthma tends to occur at rest and at night, while that with chronic airflow obstruction occurs with exertion.

Examination

The routine findings from inspection, percussion and auscultation will determine whether the underlying lung disease is localised or generalised. The generalised findings for various disorders of the lungs are summarised in Table 49.6.

Table 49.6 Comparison of examination findings for various lung disorders

	Trachea	Chest wall movement	Percussion note	Breath sounds	Vocal fremitus	Adventitious sounds
Normal	Midline	Equal expansion	Resonant	Vesicular	Normal	Nil: ± few transient inspiratory basal crackles
Asthma	Midline	Decreased (bilateral)	Resonant	Vesicular - prolonged expiration	Normal or decreased	Expiratory wheezes
Emphysema	Midline	Decreased (bilateral)	Resonant to hyperresonant	Vesicular - decreased	Decreased	Nil or the crackles and wheezes of chronic bronchitis
Consolidation (e.g. lobar pneumonia)	Midline	Decreased on affected side	Dull	Bronchial	Increased	Fine late inspiratory crackles
Collapse: major bronchus	Towards affected side	Decreased (unilateral)	Dull	Absent or decreased	Absent or decreased	Nil
Collapse: peripheral bronchus	Towards affected side	Decreased (unilateral)	Dull	Bronchial	Increased	Coarse crackles
Pleural effusion (> 500 mL)	Towards opposite side (if massive)	Decreased (unilateral)	Stony dull	Absent or decreased	Absent or decreased	None
Pneumothorax (large)	Towards opposite side (if tension)	Decreased (unilateral)	Hyperresonant	Absent or decreased	Absent or decreased	None
Fibrosis (generalised)	Midline	Decreased (bilateral)	Normal	Vesicular	Increased	Fine crackles
Bronchiectasis	Midline	Slight decrease	Resonant to dull	Bronchial	Normal or decreased	Coarse crackles ± localised wheeze

Careful inspection is mandatory. The patient should be stripped to the waist and observed for factors such as cyanosis, clubbing, mental alertness, dyspnoea at rest, use of accessory muscles, rib retraction and any other abnormalities of the chest wall. A coarse tremor or flap of the outstretched hands indicates carbon dioxide intoxication.[6] To obtain maximum value from auscultation, request the patient to open his or her mouth and take deep breaths. Adventitious sounds that are not audible during tidal breathing may then be heard. Wheezes are high-pitched continuous sounds heard either in expiration or inspiration, being more pronounced in expiration.

Crackles are short interrupted sounds heard mainly at the end of inspiration, resembling the crackling sound of hair being rubbed between the fingers near the ear. Fine crackles, previously referred to as crepitations, occur typically in lobar pneumonia and diffuse interstitial fibrosis, and are not cleared by coughing. Medium crackles are typical of congestive cardiac failure, and coarse crackles indicate airway mucus and usually clear on coughing.

Causes of pulmonary crackles

- Left ventricular failure
- Fibrosing alveolitis
- Extrinsic allergic alveolitis
- Pneumonia

- Bronchiectasis
- Chronic bronchitis
- Asbestosis
- Pulmonary fibrosis

Investigations

The two most important initial investigations for respiratory disease are chest X-ray and pulmonary function tests.

Pulmonary function tests (PFTs)

These relatively simple tests provide considerable information.

Peak expiratory flow rate

The most practical instrument for office use to detect chronic airway obstruction due to asthma or chronic bronchitis is the mini peak flow meter, which measures peak expiratory flow rate (PEFR).

The interpretation of the tests, which vary according to sex, age and height, requires charts of predicted normal values. A chart for PEFR in normal adult subjects is presented in Appendix V. The value for a particular patient should be the best of three results.

Spirometry

The measurement of the forced vital capacity (FVC) and the forced expiratory volume in one second (FEV_1) provide a very useful guide to the type of ventilatory deficit. Both the FVC and the FEV_1 are related to sex, age and height.

The FEV_1 expressed as a percentage of the FVC is an excellent measure of airflow limitation. In normal subjects it is approximately 70%. A normal spirometry pattern is shown in Figure 49.1 and abnormal patterns in Figure 49.2. Figure 49.3 summarises the relative values for these conditions.


Flow
Expiratory flow
Inspiratory flow
'effort-independent' flow
full inspiration
full expiration
Volume
FRC
tidal breathing
vital capacity

Figure 49.1 Spirometry patterns showing normal flow volume loop. Reprinted with permission.



Figure 49.2 Spirometry patterns for obstructive airway disease (e.g. asthma) and restrictive lung disease (e.g. fibrosis)



Figure 49.3 Spirometry patterns for respiratory disorders

Lung volume

Tidal volume (TV) and vital capacity (VC) can be measured by a simple spirometer but the total lung capacity and the residual volume are measured by the helium dilution method in a respiratory laboratory.

Gas transfer factor

This test measures the carbon monoxide uptake by a single breath analysis for whole lungs. In normal lungs the transfer factor is a true measure of the diff using capacity of the lungs for oxygen and depends on the thickness of the alveolar-capillary membrane.[5] Gas transfer is usually reduced in patients with severe degrees of emphysema and fibrosis, anaemia and congestive cardiac failure.

Histamine challenge test

This test indicates the presence of airway or bronchial hyper-reactivity, which is a fundamental feature with asthma. The test should not be performed on those with poor lung function and only performed by a respiratory technician under medical supervision. The test is potentially dangerous.

Other investigations (to select from)

- Haemoglobin, red cell indices and PCV
- White blood cell count (e.g. eosinophilia of asthma)
- ESR
- Arterial blood gas analysis
- Oximetry: pulse oximeters monitor oxygen saturation
- Cardiological investigations
  - ECG, including exercise
  - echocardiography (technically difficult in emphysema)
  - nuclear gated blood pool scan to assess heart function
  - cardiac enzymes
- Other medical imaging
  - high-resolution CT
  - MRI
  - ventilation and perfusion radionuclide scan (pulmonary embolism)
- Bronchoscopy, especially fibre-optic bronchoscopy
- Thoracocentesis and pleural biopsy
- Open lung biopsy
- Alpha1-antitrypsin measurement (normal range 1.1 - 2.2 g/L)

PLEURAL EFFUSION

Key points

- Normal pleural space has 10 - 20 mL fluid
- Can be detected on X-ray if . 300 mL fluid in pleural space
- Can be detected clinically if . 500 mL fluid
- Can be subpulmonary - simulates a raised diaphragm
- May be asymptomatic
- Dyspnoea common with large effusion
- Chest pain in setting of pleuritis, infection or trauma
- Signs: refer Table 49.6 at p. 529
- The fluid may be transudate or exudate (diagnosed by aspirate)
- If blood stained - malignancy, pulmonary infarction, TB

Transudate

Protein content < 30 g/L; lactic dehydrogenase < 200 IU/L

Causes

- Heart failure (90% of cases)
- Hypoproteinaemia, e.g. nephrotic syndrome
- Liver failure with ascites
- Constrictive pericarditis
- Hypothyroidism
- Ovarian tumour - right-sided effusion (Meig's syndrome)

Exudate

Protein content > 30 g/L; lactic dehydrogenase > 200 IU/L

Causes

- Infection - bacterial pneumonia, pleurisy, empyema, TB, viral
- Malignancy - carcinoma bronchus, mesothelioma, metastatic
- Pulmonary infarction
- Connective tissue diseases, e.g. SLE, RA
- Acute pancreatitis
- Lymphoma
- Sarcoidosis
- HIV with parasitic pneumonia

Management

Aspiration if symptomatic: may require repeats and pleurodesis. Treat the underlying cause.

PULMONARY FIBROSIS

Pulmonary fibrosis is the end result of a collection of ill-defined disorders producing interstitial pneumonitis leading to fibrosis. In many there is a hypersensitivity reaction to various unusual antigens. The fibrosis may be localised as following unresolved pneumonia, bilateral as with tuberculosis or widespread. Causes of widespread fibrosis include fibrosing alveolitis, rheumatoid arthritis, extrinsic allergic alveolitis including various occupational lung disorders, ornithosis, sarcoidosis and drug - induced interstitial disease (see the section on interstitial lung diseases, pp. 532 - 4).

Key features

- Consider possibility of fibrosis of lungs in chronic dyspnoea and a dry cough with normal resonance.
- Cyanosis and/or finger clubbing is commonly present.
- A ‘tricky’ feature is the presence of fine crackles at the end of inspiration upon auscultation, with faint breath sounds.
- Imaging is diagnostic especially high-resolution CT scanning. It may show the ‘honeycomb lung’ appearance.

DYSPNOEA IN CHILDREN

There are numerous causes of dyspnoea in children but the common causes are asthma, bronchiolitis and pulmonary infections. The important infections that can be fatal - croup, epiglottitis and myocarditis - must be kept in mind and intensively managed.

Bronchiolitis is an important cause of respiratory distress in infants under 6 - 12 months. It should not be confused with asthma (refer to p. 937). Sudden breathlessness or stridor may be due to an inhaled foreign body. Signs of lobar collapse may be present but physical examination may be of little help and a chest X-ray is essential.

Cardiovascular disorders, including congenital heart disease, can cause dyspnoea. Extra respiratory causes include anaemia, acidosis, aspiration, poisoning and hyperventilation.

DYSPNOEA IN THE ELDERLY

Dyspnoea in the elderly is common and is caused usually by heart failure and COPD. The other associations with ageing, such as carcinoma of the lungs, pulmonary fibrosis and drugs, are relevant. The classic problem of the aged is acute heart failure that develops typically in the early morning hours. The acute brain syndrome is a common presentation of all these disorders.

Respiratory disease in the elderly

The respiratory system, like most other bodily systems, matures until about the age of 25 years and subsequently slowly loses efficiency due to a variety of factors such as disease, smoking, pollution and ageing. There is a decline in lung function and gas exchange and decreased ventilatory responses to hypoxia and hypercapnia.

HEART FAILURE

Heart failure occurs when the heart is unable to maintain sufficient cardiac output to meet the demands of the body. Dyspnoea is a common early symptom as pulmonary congestion causes hypoxia (increased ventilation) and decreased compliance (increased work). The incidence of congestive cardiac failure (CCF) has been increasing steeply, partly due to the ageing population.

Symptoms

- increasing dyspnoea progressing to (in order)
    - fatigue, especially exertional fatigue
    - paroxysmal nocturnal dyspnoea
    - weight change: gain or loss

It is convenient to divide heart failure into left and right heart failure but they rarely occur in isolation and often occur simultaneously. Right failure is invariably secondary to left failure. Furthermore, some cardiologists stress the importance of differentiating between systolic and diastolic dysfunction. Both present in the same way clinically and hence referral for cardiac studies to obtain measurement of the left ventricular function is required. This permits an accurate diagnosis and guide to treatment and an accurate prognosis.

Refer to Chronic heart failure (Chapter 129).

CHRONIC OBSTRUCTIVE PULMONARY DISEASE

Chronic bronchitis and emphysema should be considered together as both these conditions usually coexist to some degree in each patient. An alternative, and preferable, term - chronic obstructive pulmonary disease (COPD) - is used to cover chronic bronchitis and emphysema with chronic airflow limitation.[6]

Definitions

COPD A chronic, slowly progressive disorder characterised by the presence of airway obstruction which may (or may not) be partially reversible by bronchodilator therapy.7

Chronic bronchitis A clinical condition characterised by a productive cough on most days for at least 3 months of the year for at least 2 consecutive years in the absence of any other respiratory disease that could be responsible for such excessive sputum production (such as tuberculosis or bronchiectasis).

Emphysema This is defined in pathological rather than clinical terms, as permanent dilatation and destruction of lung tissue distal to the terminal bronchioles.

Chronic airflow limitation A physiological process measured as impairment of forced expiratory flow which is the major cause of dyspnoea in these patients.

For more detail on the management of COPD refer to chapter 124, p. 1281.

INTERSTITIAL LUNG DISEASES

Interstitial lung diseases comprise a group of disorders that have the common features of inflammation and fibrosis of the interalveolar septum, representing a non-specific reaction of the lung to injury of various causes.[7]

Causes of pulmonary infiltration include:

- sarcoidosis
- cryptogenic fibrosing alveolitis (interstitial pulmonary fibrosis)
- extrinsic allergic alveolitis (hypersensitivity pneumonitis)
- drug-induced
- lymphangitis carcinomatosis
- acute pulmonary oedema
- immunological e.g. connective tissue disorders, vasculitis

Common clinical features:

- dyspnoea and dry cough (insidious onset)
- fine inspiratory crackles at lung base
- finger clubbing
- PFTs:
  - restrictive ventilatory deficit
  - decrease in gas transfer factor
- characteristic X-ray changes

High-resolution CT scanning has been a major advance in diagnosis.

Sarcoidosis

Sarcoidosis is a multisystemic disorder of unknown aetiology which is characterised by non-caseating granulomatous inflammation that involves the lung in about 90% of affected patients. A characteristic feature is bilateral hilar lymphadenopathy, which is often symptomless and detected on routine chest X-ray (CXR). Radiological lung involvement can be associated with or occur independently of hilar lymphadenopathy.

Clinical features:[7, 8]

- may be asymptomatic (one-third)
- onset usually third or fourth decade (but any age)
- bilateral hilar lymphadenopathy (on CXR)
- cough
- fever, malaise, arthralgia
- erythema nodosum
- ocular lesions (e.g. anterior uveitis)
- other multiple organ lesions (uncommon)
- overall mortality 2-5%

Erythema nodosum with acute fever, malaise and arthralgia in a young adult female is diagnostic of sarcoidosis.

Diagnosis

Histological evidence from biopsy specimen, usually transbronchial biopsy (essential if an alternative diagnosis, e.g. lymphoma, cannot be excluded) or skin biopsy in cases of erythema nodosum. A better modern diagnostic method is biopsy via video assisted thoracoscopy.

Supporting evidence:

- elevated serum ACE (non-specific)
- PFTs: restrictive pattern; impaired gas transfusion in advanced cases
- ±ve Kveim test (not recommended these days)
- serum calcium

Treatment

Sarcoidosis may resolve spontaneously (hilar lymphadenopathy without lung involvement does not require treatment).

Indications for treatment with corticosteroids:

- no spontaneous improvement or worsening after 3-6 months
- symptomatic pulmonary lesions
- eye, CNS and other systems involvement
- hypercalcaemia, hypercalcuria
- erythema nodosum with arthralgia
- persistent cough

Corticosteroid treatment

- Prednisolone 20-40 mg (o) daily for 6-8 weeks, then reduce to lowest dose that maintains improvement.[8] If there is no response, taper the dose to zero. If there is a response, taper the dose to 10-15 mg (o) daily as a maintenance dose for 6-12 months.[8]
- Prednisolone 20-30 mg for 2 weeks for erythema nodosum of sarcoidosis.

Fibrosing alveolitis

Cryptogenic fibrosing alveolitis (idiopathic pulmonary fibrosis) is the most common diagnosis among patients presenting with interstitial lung disease.

Patients usually present in the fifth to seventh decade with the clinical features as outlined under interstitial lung diseases such as slowly progressive dyspnoea over months to years. Chest X-ray abnormalities are variable but include bilateral diffuse nodular or reticulonodular shadowing favouring the lung bases. Open lung biopsy may be needed for diagnosis and staging. The usual prognosis is poor with death occuring about 2 years after diagnosis. The usual treatment is high doses of oral corticosteroids with azathioprine.[8]

Extrinsic allergic alveolitis

Extrinsic allergic alveolitis (hypersensitivity pneumonitis) is characterised by a widespread diffuse inflammatory reaction in both the small airways of the lung and the alveoli, due to the inhalation of allergens, which are usually spores of micro-organisms such as *Thermophilic actinomycetes* in 'farmer's lung' or (more commonly) avian protein from droppings or feathers in 'bird fancier's lung'. Occupational causes of extrinsic alveolitis have been described by Molina [9] (Table 49.7).

Table 49.7 Various causes of extrinsic allergic alveolitis

Occupation/disease	Source of antigen
Farmer's lung	Mouldy hay, grain and straw
Bagassosis	Mouldy sugar cane fibre (bagasse)
Bird fancier's lung	Dropping dust (e.g. pigeons, budgerigars 'bloom' on feathers)
Mushroom workers	Mushroom compost
Cheese washer's lung	Moulds or mites on cheese
Wheat weevil lung	Infested wheat flour (insect)
Ventilator pneumonitis	Humidified hot air system Air-conditioning system
Wood pulp worker's disorder	Contaminated wood dust
Detergent worker's disorder	Proteolytic enzymes
Suberosis	Mouldy cork bark

Rat handler's lung	Rat urine and serum
Malt worker's lung	Mouldy barley
Coffee worker's lung	Coffee dust
Sisal worker's lung	Sisal dust
Sericultural workers	Silkworms
Furrier's lung	Fur dust
Sausage workers	Dust
Prawn workers	Prawn fumes

Illness may present as acute or subacute episodes of pyrexia, chills and malaise with dyspnoea and a peripheral neutrophilia several hours after exposure.[8, 9] Management is based on prevention, namely avoiding exposure to allergens or wearing protective fine-mesh masks. Prednisolone can be used (with caution) to control acute symptoms. It should be pointed out that this allergic disorder is different from the infection psittacosis.

Drug-induced interstitial lung disease [8]

Drugs are an important cause of this disorder and have three main effects:

1. *Alveolitis with or without pulmonary fibrosis*. This is mainly due to cytotoxic drugs, nitrofurantoin and amiodarone. The drug should be removed and consideration given to prescribing prednisolone 50 mg (o) daily for several weeks, depending on response.
2. *Eosinophilic reactions*. This is presumably an immunological reaction, which may present as wheezing, dyspnoea, a maculopapular rash and pyrexia. The many implicated drugs include various antibiotics, NSAIDs, cytotoxic agents, major tranquillisers and antidepressants, and antiepileptics. Treatment is drug removal and a short course of prednisolone 20 - 40 mg (o) daily for 2 weeks.
3. *Non-cardiogenic acute pulmonary oedema.* This is rare and has been reported to occur with opioids, aspirin, hydrochlorothiazide, β_2-adrenoceptor agonists (given IV to suppress premature labour), cytotoxics, interleukin-2, heroin.

OCCUPATIONAL PULMONARY DISEASE

Various types of acute and chronic pulmonary diseases are related to exposure to noxious substances such as dusts, gases and vapours in the workplace. Common chemical causes include formaldehyde used in processed woods, e.g. chipboard and medium-density fibre. GPs have a crucial role in the identification of the possible work-relatedness of lung disease.

Disorders due to chemical agents include:

- obstructive airways disorders, such as occupational asthma, acute bronchitis, (chronic) industrial bronchitis, byssinosis (asthma-like condition due to cotton dust)
- extrinsic allergic alveolitis
- pulmonary fibrosis (pneumoconiosis) due to mineral dust
- lung cancer due to industrial agents such as asbestos, various hydrocarbons
- pleural disorders, usually associated with asbestosis

Pneumoconiosis

The term *pneumoconiosis* refers to the accumulation of dust in the lungs and the reaction of tissue to its presence, namely chronic fibrosis. The main cause worldwide is inhalation of coal dust, a specific severe variety being progressive massive fibrosis (complicated coal worker's pneumoconiosis) in which the patient suffers severe dyspnoea of effort and cough often productive of black sputum. Table 49.8 summarises the important causes.

Table 49.8 Selected pneumoconioses

Fibrotic lung disease	Agent	Typical occupations
Coal dust		
Coal worker's pneumoconiosis	Coal dust	• Coal mining
Metal dust		
Siderosis	Metallic iron or iron oxide	• Mining • Welding • Foundry work
Inorganic dusts		
Silicosis	Silica (silicon dioxide)	• Quarrying • Rock mining • Stone cutting • Sandblasting
Silicate dusts		
Asbestosis	Asbestos	• Mining • Shipbuilding • Insulation • Power stations
		• Wharf labouring

Of particular concern are diseases caused by inhalation of fibres of asbestos, which is a mixture of silicates of iron, magnesium, cadmium, nickel and aluminium. The diseases include asbestosis, diffuse pleural thickening, pleural plaques, mesothelioma and increased bronchial carcinoma in smokers. Pulmonary asbestosis has classic X-ray changes but high resolution CT scans may be required to confirm the presence of calcifiedpleural plaques. It usually takes 10 - 20 years from exposure for asbestosis to develop and 20 - 40 years for mesothelioma to develop,[7] while bronchial carcinoma is caused by the synergistic effects of asbestosis and cigarette smoking.

ACUTE RESPIRATORY DISTRESS SYNDROME (ARDS)

ARDS, also known as acute lung injury and formerly called ‘adult respiratory stress syndrome’, refers to acute hypoxaemic respiratory failure following a pulmonary or systemic insult with no apparent cardiogenic cause of pulmonary oedema. This occurs about 12-48 hours after the event.[10] The most common cause is sepsis which accounts for about onethird of ARDS patients. The mortality rate is 30-40% increasing if accompanied by sepsis. Management is based on early diagnosis, early referral, identification and treatment of the underlying condition and then optimal intensive care.[11]

Key features

- Sudden onset of respiratory distress
- Stiff lungs - reduced lung compliance
- Bilateral pulmonary infiltrates on X-ray
- No apparent evidence of heart failure
- Absence of elevated left atrial pressure
- Specific gas exchange abnormalities
- Signs: tachypnoea, laboured breathing, rib retraction, central cyanosis, fine crackles on auscultation

The differential diagnoses are pneumonia and acute heart failure. Common risk factors/associations for ARDS include (indirectly - systemic) - sepsis, shock, trauma, burns, multiple transfusions, drug overdose, e.g. heroin, multiple transfusions, obstetric complications, e.g. eclampsia, amniotic fluid embolism and many direct causes such as pulmonary aspiration, pneumonia, toxic gas inhalation, blast injury and pneumonia (e.g. SARS).

SEVERE ACUTE RESPIRATORY SYNDROME (SARS)

SARS is a recently recognised respiratory illness of varying severity (mild to severe) with a known mortality rate of about 10% of clinically established cases. All cases to date exhibit a high fever of > 38°C. It is considered to be an atypical pneumonia caused by a quite unique coronavirus. Cases have acquired SARS following exposure to endemic areas of South-East Asia.

Suspected case ° fever > 38°C + cough, breathing difficulty or dyspnoea + contact with SARS person/area

Key features

- Symptoms of atypical pneumonia: fever, cough, dyspnoea
- Associated may be myalgia, diarrhoea, headache, sore throat, rhinorrhoea, confusion, malaise, rash
- Virulent virus - droplet spread
- Incubation period 2 - 7 days
- Crackles and wheezes may be present on auscultation but no hallmark sign
- Non-specific X-ray signs
- Prime CT scans may show typical changes
- Diagnosis confirmed by PCR studies or isolation of the virus on culture
- Complications appear to be confined to the lung

Optimal treatment for SARS by drugs is still being evaluated. Preventive measures, including wearing of face masks (NIOSH standard mask best) and appropriate infection control procedures, are important. Wear a mask, gloves, goggles and gown for clinical assessment of suspected cases and switch off air-conditioning (refer to chapter 26, p. 262).

Practice tips

- Remember to order a chest X-ray and pulmonary function tests in all doubtful cases of dyspnoea.
- All heart diseases have dyspnoea as a common early symptom.
- Increasing dyspnoea on exertion may be the earliest symptom of incipient heart failure.
- Several drugs can produce a wide variety of respiratory disorders, particularly pulmonary fibrosis and pulmonary eosinophilia. Amiodarone and cytotoxic drugs, especially bleomycin, are the main causes.
- Dyspnoea in the presence of lung cancer may be caused by many factors, such as pleural effusion, lobar collapse, upper airway obstruction and lymphangitis carcinomatosis.
- The abrupt onset of severe dyspnoea suggests pneumothorax or pulmonary embolism.
- If a patient develops a relapse of dyspnoea while on digoxin therapy, consider the real possibility of digoxin toxicity and/or electrolyte abnormalities leading to left heart failure.
- Recurrent attacks of sudden dyspnoea, especially waking the patient at night, are suggestive of asthma or left heart failure.
- Causes of hyperventilation include drugs, asthma, thyrotoxicosis and panic attacks/anxiety.

BRONCHIAL CARCINOMA

Dyspnoea is associated with about 60% of cases of lung cancer.[3] It is not a common early symptom unless bronchial occlusion causes extrinsic collapse. In advanced cancer, whether primary or secondary, direct spread or metastases may cause dyspnoea. Other factors include pleural effusion, lobar collapse, metastatic infiltration, upper airway obstruction due to superior vena cava (SVC) obstruction and lymphangitis carcinomatosis. A special problem arises with coexisting chronic bronchitis and emphysema.

WHEN TO REFER

- Patients with acute onset of severe dyspnoea
- All patients with heart failure resistant to initial therapy or where the diagnosis is in doubt
- Patients with pulmonary disease of uncertain aetiology, especially those requiring respiratory function tests
- Those in whom carcinoma of the lung is suspected

REFERENCES

1. Sandler G. *Common Medical Problems*. London: ADIS Press, 1984: 31 - 56.
2. Cormack J, Marinker M, Morrell D. *Practice: A Handbook of Primary Health Care*. London: Kluwer-Harrap Handbooks, 1980; 3(29): 3.
3. Walsh TD. *Symptom Control*. Boston: Blackwell Scientific Publications, 1989: 157 - 64.
4. Beck ER, Francis JL, Souhami RL. *Tutorials in Differential Diagnosis* (3rd edn). Edinburgh: Churchill Livingstone, 1993: 37.
5. Kelly DT. Cardiac failure. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 97 - 9.
6. Kumar PJ, Clark ML. *Clinical Medicine* (2nd edn). London: Bailliere Tindall, 1990: 637 - 90.
7. Tierney LM et al. *Current Medical Diagnosis and Treatment* (45th edn). New York: The McGraw-Hill Companies, 2006: 275 - 80.
8. Dowden J (Chair: Writing group). *Therapeutic Guidelines*: *Respiratory* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2005: 123 - 9.
9. Molina C. Occupational extrinsic allergic alveolitis. In: Pepys J ed. *Clinics in Immunology and Allergy*. London: WB Saunders, 1984: 173 - 90.
10. Tierney LM et al. *Current Medical Diagnosis and Treatment* (45th edn). New York: The McGraw-Hill Companies, 2006: 309.
11. Ware LB et al. The acute respiratory distress syndrome. N Engl J Med, 2000; 342: 1334.

CHAPTER 50. THE PAINFUL EAR

> The ears should be kept perfectly clean; but it must never be done in company. It should never be done with a pin, and still less with the fingers, but always with an ear picker.
>
> ST JEAN BAPTISTE DE LA SALLE (1651 - 1719)

THE PAINFUL EAR

Pain in the ear (otalgia) is a common symptom in general practice. It affects all ages, but is most prevalent in children, where otitis media is the commonest cause. Ear pain may be caused by disorders of the ear or may arise from other structures, and in many instances the precise diagnosis is difficult to make. Important causes of ear pain are summarised in Table 50.1.[1]

Table 50.1 Causes of ear pain

1 Ear
External ear
Perichondritis
Otitis externa: ▪ *Candida albicans* ▪ *Aspergillus nigra* ▪ *Pseudomonas* spp. ▪ *Staphylococcus aureus*
Furunculosis
Trauma
Neoplasia
Herpes zoster (Ramsay-Hunt syndrome)
Viral myringitis
Wax-impacted
Middle ear
Eustachian insufficiency
Eustachian tube dysfunction
Barotrauma
Acute otitis media
Chronic otitis media and cholesteatoma
Acute mastoiditis
2 Periotic cause
Dental disorders
Upper cervical spinal dysfunction
TMJ arthralgia
Parotitis
Temporal arteritis
Lymph node inflammation
Other referred causes
Pharyngeal disorders
Tonsillitis

Glossopharyngeal neuralgia

A patient with a painful ear often requests urgent attention, and calls in the middle of the night from anxious parents of a screaming child are commonplace. Infants may present with nothing except malaise, vomiting or screaming attacks.

KEY FACTS AND CHECKPOINTS

- Of patients presenting with earache, 77% can be expected to have acute otitis media and 12% otitis externa.[2]
- Approximately one of every 25 patients in general practice will present with an earache.
- Two-thirds of children will sustain at least one episode of otitis media by their second birthday; one in seven children will have had more than six episodes by this age.[3]
- Otitis media is unlikely to be present if the tympanic membrane (TM) is mobile. Pneumatic otoscopy greatly assists diagnosis since the most valuable sign of otitis media is absent or diminished motility of the TM.
- Bullous myringitis, which causes haemorrhagic blistering of the eardrum or external ear canal, is an uncommon cause of severe pain. It is caused by a virus, probably influenza.[4]
- The antibiotic of first choice for acute otitis media (children and adults) is amoxycillin.
- Otitis externa can be distinguished from otitis media by pain on movement of the pinna.

A DIAGNOSTIC APPROACH

Using the safe diagnostic approach model (Table 50.2) the five self-posed questions can be answered as follows.

Table 50.2 The painful ear: diagnostic strategy model

Q. Probability diagnosis
A. Otitis media (viral or bacterial) Otitis externa TMJ arthralgia Eustachian tube dysfunction
Q. Serious disorders not to be missed
A. Neoplasia of external ear Carcinoma of other sites (e.g. tongue, throat) Herpes zoster (Ramsay-Hunt syndrome) Acute mastoiditis Cholesteatoma Necrotising otitis externa
Q. Pitfalls (often missed)
A. Foreign bodies in ear Hard ear wax Barotrauma Dental causes Referred pain: neck, throat Unerupted wisdom tooth and other dental causes TMJ arthralgia Facial neuralgias, esp. glossopharyngeal Post tonsillectomy

- from the wound
- from TMJ due to mouth gag

Q. Seven masquerades checklist

A. Depression

Diabetes -

Drugs -

Anaemia -

Thyroid disorder -

Spinal dysfunction

UTI -

Q. Is the patient trying to tell me something?

A. Unlikely, but always possible with pain.

More likely in children. Consider factitious pain.

Probability diagnosis

The commonest cause of ear pain is acute otitis media. Chronic otitis media and otitis externa are also common. In the tropics, 'tropical ear' due to acute bacterial otitis is a particular problem. Temporomandibular joint (TMJ) arthralgia, which may be acute or chronic, is also common and must be considered, especially when otitis media and otitis externa are excluded.

Serious disorders not to be missed

As always, it is important not to overlook malignant diseases, especially the obscure ones such as carcinoma of the tongue, palate or tonsils that cause referred pain.

Locally destructive cholesteatoma associated with chronic otitis media must be searched for. It signifies the 'unsafe' ear (Figure 50.1) that must be distinguished from the so-called 'safe' ear (Figure 50.2).


superior margin of
tympanic membrane
attic perforation
lateral process
of malleus

Figure 50.1 Infected ear: unsafe perforation



Figure 50.2 Infected ear: safe perforation

Herpes zoster should be considered, especially if it does not erupt on the pinna and is confined to the ear canal (usually the posterior wall), and especially in the older person.

Pitfalls

The medical aphorism ‘more things are missed by not looking than by not knowing’ applies particularly to the painful ear - good illumination and focusing the auroscope are mandatory. Particular attention should be paid to the external canal - look for hard wax, otitis externa, furuncles and foreign objects such as insects.

It may not be possible to visualise the TMs so it is important to clean the canal to permit this (if possible, on the first visit). Otitis media may coexist with otitis externa. Barotrauma should be considered, especially if pain follows air travel or diving.

General pitfalls

- Failing to visualise the TM before diagnosis and treatment
- Not checking out possible referral sites such as the oropharynx and teeth
- Overlooking common musculoskeletal causes such as TMJ arthralgia and cervical spondylosis
- Failing to recognise the unsafe ear

Seven masquerades checklist

Of the conditions in the checklist, depression and dysfunction of the upper cervical spine have to be considered. Depressive illnesses should be considered in any patient complaining of chronic pain.

Disorders of the upper cervical spine are a commonly overlooked cause of periotic pain. Pain from the C2 and C3 levels are referred to the posterior region of the ear.

Psychogenic considerations

Such factors are unlikely, unless the pain causes discomfort in the periotic region, which is likely to be magnified by a depressive state.

THE CLINICAL APPROACH

History

In assessing the painful ear the relevant features are:

- site of pain and radiation
- details of the onset of pain
- nature of the pain
- aggravating or relieving factors, especially swimming
- associated features such as deafness, discharge, vertigo, tinnitus and irritation of the external ear, sore throat

Agonising pain may be caused by perichondritis or furunculosis of the external ear and by the rare problem of herpes zoster (Ramsay-Hunt syndrome).[5] Movement of the pinna markedly increases the pain of acute otitis externa and perichondritis, and movement of the jaw usually causes an exacerbation of TMJ arthralgia or severe otitis externa.

Key questions (especially children)

- Where is the pain?
- Is it in the ear, behind or below it?
- Is it in one ear or both ears?
- Have you noticed any other symptoms such as sore throat, fever or vomiting?
- Has anyone hit you over the ear?
- Has there been a discharge from the ear?
- Have you noticed any deafness?
- Are you allergic to penicillin?
- Have you been swimming in a spa, and where?
- Have you been in an aeroplane?

Physical examination

The patient's general state and behaviour is observed during the history taking. Sudden, jabbing pain may indicate neuralgia, particularly glossopharyngeal neuralgia or a severe infection. The external ear is carefully inspected and the pinna manipulated to determine any tenderness.

Palpate the face and neck and include the parotid glands, regional lymph nodes and the skin. Inspect the TMJs - tenderness from dysfunction typically lies immediately in front of the external auditory meatus. Palpate the TMJ over the lateral aspect at the joint disc. Ask the patient to open the mouth fully when tenderness is maximal. The TMJ can be palpated posteriorly by inserting the little finger into the external canal.

Inspect both ear canals and TMs with the auroscope, using the largest earpiece that comfortably fits into the canal. Better visualisation of the TM can be achieved by pulling the pinna back in young children and up and back in older children. Impacted wax may not explain the otalgia. If herpes zoster involves the facial nerve, vesicles may be noted in and around the external auditory meatus (notably the posterior wall).

If the diagnosis is still doubtful look for causes of referred pain; inspect the cervical spine, the nose and postnasal space and the mouth, including the teeth, pharynx and larynx.

Pharyngeal and mandibular causes of periotic pain are summarised in Figures 50.3 and 50.4.



Figure 50.3 Pharangeal causes of otalgia. Courtesy Of B. Black.



Figure 50.4 Mandibular causes of otalgia. Courtesy Of B. Black.

Inspect sites supplied by the nerves V2, IX, X, XI, C1, C2 and C3 to exclude other causes of referred pain.

Investigations

Investigations are seldom necessary. Hearing tests are essential, especially for children. Simple tests such as speech discrimination, hair rubbing and tuning fork tests can be used. Otherwise audiometry can be used. Audiometry combined with tympanometry and physical measurement of the volume of the ear canal can be performed in children, irrespective of age.

Swabs from discharge, especially to determine bacterial causes, such as *Staphylococcus aureus or Pseudomonas* spp. infection, may be necessary. However, swabs are of no value if the TM is intact.

Radiology and CT scanning may be indicated for special conditions such as a suspected extraotic malignancy.

EAR PAIN IN CHILDREN

Important causes of primary otalgia in children include otitis media, otitis externa, external canal furuncle or abscess, chronic eczema with fissuring of the auricle, impacted wax, foreign body, barotrauma, perichondritis, mastoiditis and bullous myringitis. Secondary otalgia includes pharyngeal lesions, dental problems, gingivostomatitis, mumps and postauricular lymphadenopathy. Peritonsillar abscess (quinsy) may cause ear pain.

Foreign bodies

Foreign bodies (FBs) are frequently inserted into the ear canal. They can usually be syringed out or lifted with thin forceps. Various improvised methods can be used to remove FBs in cooperative children. These include a probe to roll out the FB or a rubber catheter used as a form of suction or otherwise a fine sucker.[6]

Probe method

This requires good vision using a head mirror or head light and a thin probe. The probe is inserted under and just beyond the FB. Lever it in such a way that the tip of the probe 'rolls' the foreign body out of the obstructed passage (Figure 50.5).



Figure 50.5 Probe method of removing a foreign body: **(a)** the tip of the probe is lifted by depressing the outer end of the probe; **(b)** continued gentle levering 'rolls' the foreign body out

Rubber catheter suction method

The only equipment required for this relatively simple and painless method is a straight rubber catheter (large type) and perhaps a suction pump. The end of the catheter is cut at right angles, a thin smear of petroleum jelly is applied to the rim and this end is applied to the FB (Figure 50.6). Suction is applied either orally or by a pump. Gentle pump suction is preferred but it is advisable to pinch closed the suction catheter until close to the FB as the hissing noise may frighten the child.



Figure 50.6 Extracting the foreign body using a rubber catheter; **(a)** catheter cut straight across near its extremity; **(b)** application of suction (orally or by pump)

Insects in the ear

Live insects should be immobilised by first instilling Aquaear drops or olive oil, and then syringing the ear with warm water (Figure 50.7).



Figure 50.7 Insect in the ear: **(a)** first aid; **(b)** office procedure

Dead flies that have originally been attracted to pus are best removed by suction.

Note: If simple methods such as syringing fail to dislodge the FB it is important to refer for examination and removal under microscopic vision. Syringing should not be performed if there is a possibility of the FB perforating the TM.

OTITIS MEDIA IN CHILDREN

Otitis media is very common in children and is the most common reason a child is brought in for medical attention. Persistent middle ear effusions may follow and affect the language and cognitive development of young children.

Features

- Two peaks of incidence: 6 - 12 months of age and school entry.
- Seasonal incidence coincides with URTIs.
- Bacteria cause two-thirds of cases.[7]

- The commonest organisms are *Streptococcus pneumoniae, Haemophilus influenzae* and *Moraxella catarrhalis*.
- Fever, irritability, otalgia and otorrhoea may be present.
- The main symptoms in older children are increasing earache and hearing loss.
- Pulling at the ears is a common sign in infants.
- Removal of wax is necessary in about 30% to visualise the TM.

Visualisation of the tympanic membrane

Use the largest ear speculum that will comfortably fit in the child's ear. A good technique to enable the examination of the ears (also nose and throat) in a reluctant child is where the child is held against the parent's chest while the parent's arm embraces the child's arm and trunk.

Note the following features of the TM: translucency, colour, position and motility.

Treatment

Many children with viral URTIs have mild reddening or dullness of the eardrum and antibiotics are not warranted particularly in the absence of systemic features (fever and vomiting).8 In contrast, where the eardrum is red or yellow and bulging, with loss of anatomical landmarks, antibiotic therapy is indicated.

The debate about the role of antibiotics in acute otitis media is controversial. An Australian study revealed that antibiotics provided only modest benefit: up to 20 children must be treated to prevent one child from experiencing pain by 2 - 7 days after presentation.[9, 10] A North American survey concluded ‘treat children who are very ill (approximately 15%) with antibiotics immediately. Treat all others for pain and pain only and follow up according to age. Treat with antibiotics if more serious symptoms develop, or the fever, pain and other symptoms do not resolve within the time frame’.[11]

The antibiotic of choice is:[8]

- amoxycillin 15 mg/kg 8 hourly (o) for 5 - 7 days *or*

amoxycillin 30 mg/kg 12 hourly (o) for 5 - 7 days If β-lactamase producing bacteria are suspected or documented, or initial treatment fails, use:

- cefaclor 10 mg/kg/day (max. 750 mg/day) orally in three divided doses for 5 - 7 days (cefaclor is second choice irrespective of cause) *or*
- (if resistance to amoxycillin is suspected or proven) amoxycillin/potassium clavulanate

With appropriate treatment most children with acute otitis media are significantly improved within 48 hours. Parents should be encouraged to contact their doctor if no improvement occurs within 72 hours. This problem is usually due to a resistant organism or suppuration. The patient should be re-evaluated at 10 days.

It is of interest that some practitioners refer to the ‘Pollyanna’ phenomenon when treating otitis media; that is, all antibiotics seem to work!

Symptomatic treatment

Rest the patient in a warm room with adequate humi dity. Use analgesics such as paracetamol (acetaminophen) in high dosage. Although the use of antihistamines and decongestants has not been verified scientifically, the author has found nasal decongestants (as oxymetazoline nasal drops or sprays) effective in distressed children with an associated URTI. Otherwise, avoid antihistamines and decongestants.

Follow-up: adequate follow-up with hearing assessment is mandatory.

Complications

- *Middle ear effusion*. 70% of children will have an effusion present 2 weeks from the time of diagnosis, 40% at 4 weeks, with 10% having persistent effusions for 3 months or more. If the effusion is still present at 6 - 8 weeks, a second course of antibiotics should be prescribed.[2] If the effusion persists beyond 3 months refer for an ENT opinion.
- *Acute mastoiditis*. This is a major complication that presents with pain, swelling and tenderness developing behind the ear associated with a general deterioration in the condition of the child. Such a complication requires immediate referral.[7]
- *Chronic otitis media*

- *Rare complications.* These include labyrinthitis, petrositis, facial paresis and intracranial abscess.
- *Serous otitis media (glue ear).* This represents incomplete resolution of suppurative otitis media. Signs include loss of drum mobility, hearing loss and abnormal impedance. Most resolve spontaneously but any necessary treatment includes medications such as bromhexine elixir and Demazin syrup, auto-inflations and 'Oto-vent' assisted nasal inflation.

Recurrent acute otitis media

Antibiotic prevention of acute otitis media is indicated (arguably) if it occurs more often than every other month or for three or more episodes in 6 months.

- Chemoprophylaxis (for about 4 months)
  - amoxycillin twice daily (first choice) *or*
  - cefaclor twice daily

Consider *Pneumococcus* vaccine in children over 18 months of age (if not already given) in combination with the antibiotic. Avoid smoke exposure (cigarettes and wood fires) and group child care.

Viral infections

Most children with viral URTIs have mild reddening or dullness of the eardrum and antibiotics are not warranted. If painful bullous otitis media is present, either prick the bulla with a sterile needle for pain relief, or instil dehydrating eardrops such as anhydrous glycerol.

EAR PAIN IN THE ELDERLY

Causes of otalgia that mainly afflict the elderly include herpes zoster (Ramsay-Hunt syndrome), TMJ arthralgia, temporal arteritis and neoplasia. It is especially important to search for evidence of malignancy.

ACUTE OTITIS MEDIA

Acute otitis media causes deep-seated ear pain, deafness and often systemic illness (Figure 50.8). The sequence of symptoms is a blocked ear feeling, pain and fever. Discharge may follow if the TM perforates, with relief of pain and fever.



Figure 50.8 Acute otitis media causing true otalgia. The ear drum bulges laterally due to pus in the middle ear. Perforation and otorrhoea imminent. *Photo courtesy Bruce Black*

The commonest organisms are viruses (adenovirus and enterovirus), and the bacteria *H. influenzae, S. pneumoniae, Branhamella* (previously *Neisseria*) *catarrhalis* and β-haemolytic streptococci.

The two cardinal features of diagnosis are inflammation and middle ear effusion.

Appearance of the tympanic membrane (all ages)

Translucency. If the middle ear structures are clearly visible through the drum, otitis media is unlikely.

Colour. The normal TM is a shiny pale-grey to brown: a yellow colour is suggestive of an effusion.

Diagnosis

The main diagnostic feature is the redness of the TM. The inflammatory process usually begins in the upper posterior quadrant and spreads peripherally and down the handle of the malleus (Figure 50.9). The TM will be seen to be reddened and inflamed with engorgement of the vessels particularly along the handle of the malleus. The loss of light reflex follows and anatomical features then become difficult to recognise as the TM becomes oedematous. Bulging of the drum is a late sign. Blisters are often seen on the TM and this is thought to be due to a viral infection in the epidermal layers of the drum.


• erythema of prominent blood vessels progressing down handle of malleolus
• normal drum
• progressive erythema
• loss of light reflex
• bulging pars flaccida
• red pars tensa
• anatomical structures unidentifiable

Figure 50.9 The appearances of the left tympanic membrane in the progressive development of acute otitis media

Treatment of acute otitis media (adults)

- Analgesics to relieve pain
- Adequate rest in a warm room
- Nasal decongestants for nasal congestion
- Antibiotics until resolution of all signs of infection
- Treat associated conditions (e.g. adenoid hypertrophy)
- Follow-up: review and test hearing audiometrically

Antibiotic treatment[7]

First choice:

- amoxycillin 750 mg (o) bd for 5 days[7] *or* 500 mg (o) tds for 5 days

A longer course (up to 10 days) may be required depending on severity and response to 5-day course.

Alternatives:

- doxycycline 100 mg (o) bd for 5 - 7 days (daily for milder infections)*or*
- cefaclor 250 mg (o) tds for 5 - 7 days *or*
- (if resistance to amoxycillin is suspected or proven) amoxycillin/potassium clavulanate 500/125 mg (o) tds for 5 days (the most effective antibiotic)

Consider surgical intervention for failed therapy.

CHRONIC OTITIS MEDIA

There are two types of chronic suppurative otitis media and they both present with deafness and discharge without pain. The discharge occurs through a perforation in the TM: one is safe (Figure 50.10a), the other unsafe (Figure 50.10b).



Figure 50.10 (a) Chronic otitis media with loss of the tympanic membrane this is ‘safe ear’**(b)** Unsafe ear: chronic otitis media with attic cholesteatoma. *Photo courtesy Bruce Black*

Chronic discharging otitis media (safe)[8]

If aural discharge persists for > 6 weeks after course of antibiotics, treatment can be with topical steroid and antibiotic combination drops, following ear toilet. The toileting can be done at home by dry mopping with rolled tissue spear. If persistent, referral to exclude cholesteatoma or chronic osteitis is advisable.

Recognising the unsafe ear

Examination of an infected ear should include inspection of the attic region, the small area of drum between the lateral process of the malleus, and the roof of the external auditory canal immediately above it. A perforation here renders the ear ‘unsafe’ (Figure 50.1); other perforations, not involving the drum margin (Figure 50.2), are regarded as ‘safe’.[12]

Cholesteatoma[12]

The status of a perforation depends on the presence of accumulated squamous epithelium (termed cholesteatoma) in the middle ear, because this erodes bone. An attic perforation contains such material; safe perforations do not. Cholesteatoma is visible through the hole as white flakes, unless it is obscured by discharge or a persistent overlying scab. Either type of perforation can lead to chronic infective discharge, the nature of which varies with its origin. Mucus admixture is recognised by its stretch and recoil when this discharge is being cleaned from the external auditory canal. The types of discharge are compared in Table 50.3.

Table 50.3 Comparison of types of discharge

	Unsafe	Safe
Source	Cholesteatoma	Mucosa
Odour	Foul	Inoffensive
Amount	Usually scant, never profuse	Can be profuse
Nature	Purulent	Mucopurulent

Management

If an attic perforation is recognised or suspected, specialist referral is essential. Cholesteatoma cannot be eradicated by medical means: surgical removal is necessary to prevent a serious infratemporal or intracranial complication.

OTITIS EXTERNA

Otitis externa (Figure 50.11), also known as ‘swimmer's ear’, ‘surfer's ear’ and ‘tropical ear’, is common in a country whose climate and coastal living leads to extensive water sports. It is more prevalent in hot humid conditions and therefore in the tropics.



Figure 50.11 Otitis externa

Predisposing factors are allergic skin conditions, ear canal trauma, water penetration (swimming, humidity, showering), water and debris retention (wax, dermatitis, exostoses), foreign bodies, contamination from swimming water including spas, and use of Q tips and hearing aids.

Common responsible organisms

- Bacteria:
  - *Pseudomonas* sp.
  - *Escherichia coli*
  - *S. aureus*
  - *Proteus* sp.
  - *Klebsiella* sp.
- Fungi:
  - *Candida albicans*
  - *Aspergillus* sp.

Clinical features

- Itching at first
- Pain (mild to intense)
- Fullness in ear canal
- Scant discharge
- Hearing loss

Signs

- Oedema (mild to extensive)
- Tenderness on moving auricle or jaw
- Erythema
- Discharge (offensive if coliform)
- Pale cream ‘wet blotting paper’ debris - *C. albicans*
- Black spores of *Aspergillus nigra*
- TM granular or dull red

Obtain culture, especially if resistant *Pseudomonas* sp. suspected, by using small ear swab.

Note: ‘Malignant’ otitis externa occurs in diabetics due to *Pseudomonas* infection at base of skull.

Management

Aural toilet

Meticulous aural toilet by gentle suction and dry mopping with a wisp of cotton wool on a fine broach under good lighting is the keystone of management. This enables topical medication to be applied directly to the skin.

Syringing

This is appropriate in some cases but the canal must be dried meticulously afterwards. For most cases it is not recommended.

Dressings

Dressings are essential in all but the mildest forms. After cleaning and drying, insert 10 - 20 cm of 4 mm Nufold gauze impregnated with a steroid and antibiotic cream.

For severe otitis externa a wick is important and will reduce the oedema and pain in 12 to 24 hours (Figure 50.12). The wick can be soaked in an astringent (e.g. aluminium acetate 4% solution or glycerin and 10% ichthammol). The wick needs replacement daily until the swelling has subsided.



Figure 50.12 Insertion of a wick; it is packed gradually by short back-and-forth movements of the forceps. Courtesy of B. Black.

Topical antimicrobials[8]

The most effective, especially when the canal is open, is an antibacterial, antifungal and corticosteroid preparation such as Kenacomb or Sofradex drops (2 - 3 drops tds), Locacorten-Vioform drops (2 - 3 drops bd) or Ciproxin HC (2 - 3 drops bd). The tragus should be pumped for 30 seconds after instillation by pressing on it repeatedly, within the limitation of any pain.

Other measures

- Strong analgesics are essential.
- Antibiotics have little place in treatment unless a spreading cellulitis has developed.
- Prevent scratching and entry of water.
- Use a wick soaked in combination steroid and antibiotic ointiment for more severe cases.

Practice tip for severe ‘tropical ear’

- Prednisolone (o) 15 mg statim then 10 mg 8 hourly for 6 doses followed by
- Merocel ear wick
- Topical Kenacomb or Sofradex drops

Prevention

- Keep the ear dry, especially those involved in water sports.
- Protect the ear with various water-proofing methods:
  - cotton wool coated with petroleum jelly
  - tailor-made ear plugs (e.g. EAR foam plugs)
  - silicone putty or Blu-tack
  - a bathing cap pulled well forward allows these plugs to stay *in situ*
- Avoid poking objects such as hairpins and cotton buds in the ear to clean the canal.
- If water enters, shake it out or use Aquaear drops (spirit drops help dry the canal).

Necrotising otitis externa

This severe complication usually due to *Pseudomonas aeruginosa* can occur in the immunocompromised, diabetic or elderly patient. It involves cartilage and bone and should be considered with treatment failure, severe persistent pain, fever and visible granulation tissue. Urgent referral is advisable.

EAR EXOSTOSES (‘SURFER'S EAR’)

These bony overgrowths are caused by water retention in the ear.

Prevention

- Use plugs or Blu-Tack to waterproof ear.
- Dry thoroughly with hair dryer after swimming.

FURUNCULOSIS

Furunculosis is a staphylococcal infection of the hair follicle in the outer cartilaginous part of the ear canal. It is usually intensely painful. Fever occurs only when the infection spreads in front of the ear as cellulitis. The pinna is tender on movement - a sign that is not a feature of acute otitis media. The furuncle (boil) may be seen in the external auditory meatus (Figure 50.13).



Figure 50.13 Furuncle (boil) in hair-bearing area at opening of the ear canal

Management

- If pointing, it can be incised after a local anaesthetic or freezing spray.
- Warmth (e.g. use hot face washer, hot water bottle).
- If fever with cellulitis - dicloxacillin.

PERICHONDRITIS

Perichondritis is infection of the cartilage of the ear characterised by severe pain of the pinna, which is red, swollen and exquisitely tender. It is rare and follows trauma or surgery to the ear. As the organism is frequently *P. pyocaneus* the appropriate antibiotics must be carefully chosen (e.g. ciprofloxacin).

INFECTED EAR LOBE

The cause is most likely a contact allergy to nickel in an earring, complicated by a *S. aureus* infection.

Management

- Discard the earrings.
- Clean the site to eliminate residual traces of nickel.
- Swab the site and then commence antibiotics (e.g. flucloxacillin or erythromycin).
- Instruct the patient to clean the site daily, and then apply the appropriate ointment.
- Use a 'noble metal' stud to keep the tract patent.
- Advise the use of only gold, silver or platinum studs in future.

EUSTACHIAN TUBE DYSFUNCTION

This is a common cause of discomfort.[13] Symptoms include fullness in the ear, pain of various levels and impairment of hearing. The most common causes of dysfunction are disorders causing oedema of the tubal lining, such as viral URTI and allergy when the tube is only partially blocked; swallowing and yawning may elicit a crackling or popping sound. Examination reveals retraction of the TM and decreased mobility on pneumatic otoscopy. The problem is usually transient after a viral URTI.

Treatment

- Systemic and intranasal decongestants (e.g. pseudoephedrine or corticosteroids in allergic patients).
- Autoinflation by forced exhalation against closed nostrils (avoid in active intranasal infection).
- Avoid air travel, rapid altitude change and underwater diving.

OTIC BAROTRAUMA

Barotrauma is damage caused by undergoing rapid changes in atmospheric pressure in the presence of an occluded Eustachian tube (Figure 50.14). It affects scuba divers and aircraft travellers.



Figure 50.14 Mechanism of barotrauma, with blocking of the Eustachian tube due to increased pressure at the sites indicated. Courtesy of B. Black.

The symptoms include temporary or persisting pain, deafness, vertigo, tinnitus and perhaps discharge.

Inspection of the TM may reveal (in order of seriousness): retraction; erythema; haemorrhage (due to extravasation of blood into the layers of the TM); fluid or blood in the middle ear; perforation. Perform conductive hearing loss tests with tuning fork.

Treatment

Most cases are mild and resolve spontaneously in a few days, so treat with analgesics and reassurance. Menthol inhalations are soothing and effective. Refer if any persistent problems for consideration of the Politzer bag inflation or myringotomy.

Prevention

Flying. Perform repeated Valsalva manoeuvres during descent. Use decongestant drops or sprays before boarding the aircraft, and then 2 hours before descent.

Diving. Those with nasal problems, otitis media or chronic tubal dysfunction should not dive.

PENETRATING INJURY TO TYMPANIC MEMBRANE

A penetrating injury to the TM can occur in children and adults from various causes such as pencils and slivers of wood or glass. Bleeding invariably follows and infection is the danger.

Management

- Remove blood clot by suction toilet or gentle dry mopping.
- Ensure no FB is present.
- Check hearing.
- Prescribe a course of broad spectrum antibiotics (e.g. cotrimoxazole).
- Prescribe analgesics.
- Instruct patient not to let water enter ear.
- Review in 2 days and then regularly.
- At review in 1 month the drum should be virtually healed.
- Check hearing 2 months after injury.

Complete healing can be expected within 8 weeks in 90 - 95% of such cases.[14]

TEMPOROMANDIBULAR JOINT ARTHRALGIA

If rheumatoid arthritis is excluded, a set of special exercises, which may include ‘chewing’ a piece of soft wood over the molars, invariably solves this problem (Chapter 52). If an obvious dental malocclusion is present, referral is necessary.

WHEN TO REFER

Otitis media

- Incomplete resolution of acute otitis media
- Persistent middle ear effusion for 3 months after an attack of acute otitis media
- Persistent apparent or proved deafness
- Evidence or suspicion of acute mastoiditis or other severe complications
- Frequent recurrences (e.g. four attacks a year)
- Presence of craniofacial abnormalities

Other ear problems

- Attic perforation/cholesteatoma
- FBs in ear not removed by simple measures such as syringing
- No response to treatment after 2 weeks for otitis externa
- Suspicion of carcinoma of the ear canal
- Acute TM perforation that has not healed in 6 weeks
- Chronic TM perforation (involving lower two-thirds of TM)

Practice tips

- The pain of acute otitis media may be masked by fever in babies and young children.
- A red TM is not always caused by otitis media. The blood vessels of the drum head may be engorged from crying, sneezing or nose blowing. In crying babies the TM as well as the face may be red.
- In otitis externa, most cases will resolve rapidly if the ear canal is expanded and then cleaned meticulously.
- If an adult presents with ear pain but normal auroscopy, examine possible referral sites, namely TMJ, mouth, throat, teeth and cervical spine.
- Antibiotics have no place in the treatment of otic barotrauma.
- It is good medicine to make relief of distressing ear pain a priority. Adequate analgesics must be given. There is a tendency to give too low a dose of paracetamol in children. The installation of nasal drops in infants with a snuffy nose and acute otitis media can indirectly provide amazing relief of pain.
- Spirit ear drops APF are a cheap and simple agent to use for recurrent otitis externa where wetness of the ear canal is a persistent problem.

REFERENCES

1. Black B. Otalgia. Aust Fam Physician, 1987; 16: 292 - 6.
2. Shires DB, Hennen BK, Rice DI. *Family Medicine* (2nd edn). New York: McGraw-Hill, 1987: 86 - 93.
3. Jarman R. Otitis media. Aust Paediatr Rev, 1991; 4: 1 - 2.
4. Ludman H. *ABC of Otolaryngology* (3rd edn). London: BMJ, 1993.
5. Sandler G, Fry J. *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986: 285 - 7.
6. Murtagh J. *Practice Tips* (4th edn). Sydney: McGraw-Hill, 2004: 121 - 4.
7. Robinson MJ, Roberton DM. *Practical Paediatrics*. (5th edn). Melbourne: Churchill Livingstone, 2003: 744 - 8.
8. Spicer J et al. *Therapeutic Guidelines: Antibiotics* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 234 - 40.
9. Antibiotics, patient education and otitis media. NPS News, 1999; 3: 3.
10. Del Mar C, Glasziou P, Hayem M. Are antibiotics indicated for the initial treatment of acute otitis media in children? A meta analysis. Br Med J, 1997; 314: 1526 - 9.
11. Rosser W, Shafir M. *Evidence-Based Family Medicine*. Hamilton: BC Decker, 1998: 111 - 13.
12. Black B. How to treat: Otitis media. Australian Doctor, 2002; 29 November: I - VIII.
13. Tierney LM et al. *Current Medical Diagnosis and Treatment* (45th edn). New York: The McGraw-Hill Companies, 2006: 183 - 5.
14. Kruger R, Black B. Penetrating injury eardrum. Aust Fam Physician, 1986; 15: 735.

CHAPTER 51. THE RED AND TENDER EYE

Those with sore eyes … find the light painful, while the darkness, which permits them to see nothing, is restful and agreeable.

DIO CHRYOSTOM (40 - 115)

THE RED AND TENDER EYE

A red eye accounts for at least 80% of patients with eye problems encountered in general practice.[1] An accurate history combined with a thorough examination will permit the diagnosis to be made in most cases without recourse to specialist ophthalmic equipment. A summary of the diagnostic strategy model is presented in Table 51.1.

Table 51.1 The red and tender eye: diagnostic strategy model

Q. Probability diagnosis A. Conjunctivitis • bacterial • adenovirus • allergic
Q. Serious disorders not to be missed A. Acute glaucoma Uveitis • acute iritis • choroiditis Corneal ulcer Herpes simplex keratitis Microbiol keratitis, e.g. fungal, amoeba, bacterial Herpes zoster ophthalmicus Penetrating injury Endophthalmitis Orbital cellulitis
Q. Pitfalls (often missed) A. Scleritis/episcleritis Foreign body Trauma Ultraviolet light 'keratitis' Blepharitis Cavernous sinus arteriovenous fistula
Q. Seven masquerades checklist A. Depression - Diabetes - Drugs ✔ hypersensitivity Anaemia - Thyroid disorder ✔ hyperthyroidism Spinal dysfunction - UTI -
Q. Is the patient trying to tell me something? A. Unlikely.

KEY FACTS AND CHECKPOINTS

- Acute conjunctivitis accounts for over 25% of all eye complaints seen in general practice.[2]
- A purulent discharge indicates bacterial conjunctivitis.[3]
- A clear or mucous discharge indicates viral or allergic conjunctivitis.
- Viral conjunctivitis can be slow to resolve and may last for weeks.
- Pain and visual loss suggest a serious condition such as glaucoma, uveitis (including acute iritis) or corneal ulceration.
- Beware of the unilateral red eye - think beyond bacterial or allergic conjunctivitis. It is rarely conjunctivitis and may be a corneal ulcer, keratitis, foreign body, trauma, uveitis or acute glaucoma.[4]
- Keratitis (inflammation of the cornea) is one of the most common causes of an uncomfortable red eye. Apart from the well-known viral causes (herpes simplex, herpes zoster, adenovirus and measles), it can be caused by fungal infection (usually on a damaged cornea), bacterial infection or inflammatory disorder such as ankylosing spondylitis.[5]
- Herpes simplex keratitis (dendritic ulcer) often presents painlessly as the neurotrophic effect grossly diminishes sensation.

THE CLINICAL APPROACH

The five essentials of the history are:

- history of trauma (especially as indicator of intraocular foreign body [IOFB])
- vision
- the degree and type of discomfort
- presence of discharge
- presence of photophobia

The social and occupational history is also very important. This includes a history of exposure to a ‘red eye’ at school, work or home; incidents at work such as injury, welding, foreign bodies or chemicals; and genitourinary symptoms.

When examining the unilateral red eye keep the following diagnoses in mind:

- trauma
- foreign body, including IOFB
- corneal ulcer
- iritis (uveitis)
- viral conjunctivitis (commonest type)
- acute glaucoma

The manner of onset of the irritation often gives an indication of the cause. Conjunctivitis or uveitis generally has a gradual onset of redness, while a small foreign body will produce a very rapid hyperaemia. Photophobia occurs usually with uveitis and keratitis. It is vital to elicit careful information about visual acuity. The wearing of contact lenses is very important as these are prone to cause infection or the ‘overwear syndrome’, which resembles an acute ultraviolet (UV) burn.

The key eye symptoms

The key eye symptoms are:

- itch
- irritation
- pain (with pus or watering)
- loss of vision (red or white eye)

Key questions

- Have you noticed blurring of your vision?
- Have you been in close contact with others with the same condition?
- Have you had a cold or running nose recently?
- Do you wear contact lenses?
- Can you recall scratching or injuring your eye?
- What were you doing at the time you noticed trouble?
- Have you been putting any drops, ointments or cosmetics in or around your eye?
- Do you suffer from hay fever?

- Do you have any problems with your eyelids?
- Had your eyes been watering for some time beforehand?
- Have you had any other problems?
- Have you been exposed to arc welding?

Loss of vision in the red eye

Consider:

- iritis (uveitis)
- scleritis
- acute glaucoma (pain; nausea and vomiting)
- chemical burns

The painful red eye

Causes to consider:

- keratitis
- uveitis (iritis)
- episcleritis
- scleritis
- acute glaucoma
- hypopyon (pus in the anterior chamber)
- endophthalmitis (inflammation of internal structures - may follow surgery)
- corneal abrasion/ulceration

Pain with discharge:

- keratitis

Pain with photophobia:

- uveitis
- episcleritis

Physical examination

The basic equipment:

- eye testing charts at 45 cm (18 in) and 300 cm (10 ft)
- multiple pinholes
- torch (e.g. Cobalt blue)
- magnifying aid (e.g. binocular loupe)
- glass rod or cotton bud to aid eyelid eversion
- fluorescein sterile paper strips
- anaesthetic drops
- Schiotz tonometer
- ophthalmoscope
- Ishihara's colour vision test

The four essentials of the examination are:

- testing and recording vision
- meticulous inspection under magnification
- testing the pupils
- testing ocular tension4

Also

- local anaesthetic test
- fluorescein staining
- subtarsal examination

Inspection

A thorough inspection is essential, noting the nature of the inflammatory injection, whether it is localised (episcleritis) or diffuse, viewing the iris for any irregularity, observing the cornea, and searching for foreign bodies, especially under the eyelids, and for any evidence of penetrating injury. No ocular examination is complete until the eyelid is everted and closely inspected. Both eyes must be examined since many patients presenting with conjunctivitis in one eye will have early signs of conjunctivitis in the other. Use fluorescein to help identify corneal ulceration. Local anaesthetic drops instilled prior to the examination of a painful lesion are recommended. The local anaesthetic test is a sensitive measure of a surface problem - if the pain is unrelieved a deeper problem must be suspected.

Palpate for enlarged preauricular lymph nodes, which are characteristic of viral conjunctivitis.

The nature of the injection is important. In conjunctivitis the vessels are clearly delineated and branch from the corners of the eye towards the cornea, since it involves mainly the tarsal plate. Episcleral and scleral vessels are larger than conjunctival vessels and are concentrated towards the cornea (Figure 51.1).


papillae of allergic (vernal) conjunctivitis
foreign body
lymphoid follicles of viral conjunctivitis
ciliary flush
conjunctival vessels
irregular pupil
dendritic ulcer
episcleral vessels
blepharitis
mucopurulent discharge —bacterial conjunctivitis

Figure 51.1 Physical signs to search for in a patient with a red eye (eyelids everted)

Ciliary injection appears as a red ring around the limbus of the cornea (the ciliary flush), and the individual vessels, which form a parallel arrangement, are not clearly visible. Ciliary injection may indicate a more serious deep-seated inflammatory condition such as anterior uveitis or a deep corneal infection. The presence of fine follicles on the tarsal conjuctivae indicates viral infection while a cobblestone appearance indicates allergic conjunctivitis.

Note: Slit lamp examination is ideal for the examination of the eye.

RED EYE IN CHILDREN

Children can suffer from the various types of conjunctivitis (commonly), uveitis and trauma. Of particular concern is orbital cellulitis, which may present as a unilateral swollen lid and can rapidly lead to blindness if untreated. Bacterial, viral and allergic conjunctivitis are common in all children. Conjunctivitis in infants is a serious disorder because of the immaturity of tissues and defence mechanisms. Serious corneal damage and blindness can result.

Neonatal conjunctivitis (ophthalmia neonatorum)

This is conjunctivitis in an infant less than 1 month old and is a notifiable disease. Chlamydial and gonococcal infections are uncommon but must be considered if a purulent discharge is found in the first few days of life.[6] In both conditions the parents must be investigated for associated venereal infection and treated accordingly.

Chlamydia trachomatis accounts for 50% or more of cases. Its presentation in neonates is acute, usually 1 - 2 weeks after delivery, with moderate mucopurulent discharge. It is a systemic disease and may be associated with pneumonia. The diagnosis is confirmed by serological tests on the conjunctival secretions.

Treatment is with oral erythromycin and local sulfacetamide eye drops.

Neisseria gonorrhoeae conjunctivitis, which usually occurs within 1 - 2 days of delivery, requires vigorous treatment with intravenous cephalosporins or penicillin and local sulfacetamide drops. The discharge is highly infectious and the organism has the potential for severe corneal infection or septicaemia.[6]

Other common bacterial organisms can cause neonatal conjunctivitis, and herpes simplex virus type II can cause conjunctivitis and/or eyelid vesicles or keratitis.[2]

Trachoma

More than 6 million people worldwide have trachoma caused blindness.

Trachoma is a chlamydial conjunctivitis that is prevalent in outback areas and in the Indigenous population. *C. trachomatis* is usually transmitted by human contact between children and mothers and also by flies, especially where hygiene is inadequate. It is the most common cause of blindness in the world. Recurrent and untreated disease leads to lid scarring and inturned lashes (endtropion) with corneal ulceration and visual loss. It is important to commence control of the infection in childhood (see page 556).

Treatment:

- prevention/community education
- antibiotics - azithromycin
- surgical correction (where relevant)

Blocked nasolacrimal duct

Delayed development of the nasolacrimal duct occurs in about 6% of infants,[6] resulting in blocked lacrimal drainage; the lacrimal sac becomes infected, causing a persistent discharge from one or both eyes. In the majority of infants spontaneous resolution of the problem occurs by the age of 6 months.

Management

- Local antibiotics for infective episodes
- Bathing with normal saline
- Frequent massage over the lacrimal sac
- Referral for probing of the lacrimal passage before 6 months if the discharge is profuse and irritating or between 6 and 12 months if the problem has not self-corrected

RED EYE IN THE ELDERLY

In an elderly patient there is an increased possibility of acute glaucoma, uveitis and herpes zoster. Acute angle closure glaucoma should be considered in any patient over the age of 50 presenting with an acutely painful red eye.

Eyelid conditions such as blepharitis, trichiasis, entropion and ectropion are more common in the elderly.

ACUTE CONJUNCTIVITIS

Acute conjunctivitis is defined as an episode of conjunctival inflammation lasting less than 3 weeks.[2] The two major causes are infection (either bacterial or viral) and acute allergic or toxic reactions of the conjunctiva (see Table 51.2).

Table 51.2 Major causes of a red eye

	Site of inflammation	Pain	Discharge	Vision	Photophobia	Pupil	Cornea	Ocular tension
Bacterial conjunctivitis	Conjunctiva,	Irritation -	Purulent, lids	Normal	No	Normal	Normal	Normal

	including lining of lids (usually bilateral)	gritty	stuck in the morning					
Viral conjnctivitis	Conjunctiva, lining of lids often follicular (uni or bilateral)	Girtty	Watery	Normal	No	Normal	Normal	Normal
Allergic (vernal) conjunctivitis	Conjunctiva, papillary swellings on lids linings(bilateral)	Gritty - itching	Watery	Normal	No	Normal	Normal	Normal
Contact hypersensitivity (dermatoconjunctivitis)	Conjunctiva and eyelids Oedema	Itching	Watery	Normal - may be blurred	No	Normal	Normal	Normal
Subconjunctival haemorrhage	Beefy red area fading at edge (unilateral)	No	No	Normal	No	Normal	Normal	Normal
Herpes simplex keratitis	Unilateral - circumcorneal Dendritic ulcer	Yes - gritty	No, reflex lacrimation	Blurred, but variable, depends on site	Yes	Normal	Normal	Normal
Corneal ulcer	Unilateral - circumcorneal (exclude foreign body)	Yes	No, reflex lacrimation	Blurred, but variable, depends on site	Yes	Normal	Abnormal	Normal
Scleritis/episcleritis	Localised deep redness Tender area	Yes	No	Normal	No	Normal	Normal	Normal
Acute uveitis/iritis	Maximum around cornea	Yes - radiates to brow, temple, nose	No, reflex lacrimation	Blurred	Yes	Constricted, may be irregular	Normal	Normal or low
Acute glaucoma	Diffuse but maximum circumcorneal	Yes, severe with nausea and vomiting	No, reflex lacrimation	Haloes around lights	Yes	Dilated Absent light reflex	Hazy	Hard, elevated

Clinical features

- Diffuse hyperaemia of both tarsal or bulbar conjunctivae
- Absence of ocular pain, good vision, clear cornea
- Infectious conjunctivitis is bilateral (usually) or unilateral (depending on the cause), with a discharge, and a gritty or sandy sensation

Bacterial conjunctivitis

Bacterial infection may be primary, secondary to a viral infection or secondary to blepharitis.

History

Purulent discharge with sticking together of eyelashes in the morning is typical. It usually starts in one eye and spreads to the other. There may be a history of contact with a person with similar symptoms. The organisms are usually picked up from contaminated fingers, face cloths or towels.

Key features:

- gritty red eye
- purulent discharge
- clear cornea

Examination

There is usually a bilateral mucopurulent discharge with uniform engorgement of all the conjunctival blood vessels and a non-specific papillary response (Figure 51.2). Fluorescein staining is negative.



Figure 51.2 Acute bacterial conjunctivitis with mucopurulent discharge, no corneal stain

Causative organisms

These include:

- *Streptococcus pneumoniae*
- *Haemophilus influenzae*
- *Staphylococcus aureus*
- *Streptococcus pyogenes*
- *N. gonorrhoeae* (a hyperacute onset)
- *Pseudomonas aeruginosa*

Diagnosis is usually clinical but a swab should be taken for smear and culture with:[2]

- hyperacute or severe purulent conjunctivitis
- prolonged infection
- neonates

Management

Limit the spread by avoiding close contact with others, use of separate towels and good ocular hygiene.

Mild cases

Mild cases may resolve with saline irrigation of the eyelids and conjunctiva but may last up to 14 days if untreated.[7] An antiseptic eye drop such as propamidine isethionate 0.1% (Brolene) 1 - 2 drops 6 - 8 hourly for 5 - 7 days can be used.

More severe cases

- Chloramphenicol 0.5% eye drops, 1 - 2 hourly for 2 days,[1] decrease to 4 times a day for another 7 days (max. 10 days - cases of aplastic anaemia have been reported with long-term use) Use also chloramphenicol 1% eye ointment each night or preferably[7]
- polymyxin B sulphate 5000 units/mL + gramicidin 25 µg/mL + neomycin 2.5 mg/mL (Neosporin), 1 - 2 drops hourly, decreasing to 6 hourly as the infection improves

Specific organisms

- *Pseudomonas* and other coliforms: use topical gentamicin and tobramycin.
- *N. gonorrhoeae*: use appropriate systemic antibiotics.
- *Chlamydia trachomatis* - may be sexually transmitted. Shows a brick red follicular conjunctivitis with a stringy mucus discharge.

Viral conjunctivitis

The most common cause of this very contagious condition is adenovirus.

History

It is commonly associated with URTIs and is the type of conjunctivitis that occurs in epidemics (pink eye).[1] The conjunctivitis usually has a 2 - 3 week course; it is initially one-sided but with cross-infection occurring days later in the other eye. It can be a severe problem with a very irritable, watering eye.

Examination

The examination should be conducted with gloves. It is usually bilateral with diff use conjunctival infection and productive of a scant watery discharge. Viral infections typically but not always produce a follicular response in the conjunctivae (tiny, pale lymphoid follicles) and an associated preauricular lymph node (Figure 51.3). Subconjunctival haemorrhages may occur with adenovirus infection. High magnification, ideally a slit lamp, may be necessary to visualise some of the changes, such as small corneal opacities, follicles and keratitis.



Figure 51.3 Viral conjunctivitis: watery eye, lid swelling, typical eyelid follicles, Associated local lymphadeuopathy

Diagnosis is based on clinical grounds and a history of infected contacts. Viral culture and serology can be performed to identify epidemics.

Treatment

- Limit cross-infection by appropriate rules of hygiene and patient education.
- Treatment is symptomatic - cool compress and topical lubricants (artificial tear preparations), naphazoline (e.g. Albalon), vasoconstrictors (e.g. phenylephrine) or saline bathing.
- Do not pad.
- Watch for secondary bacterial infection. Avoid corticosteroids, which reduce viral shedding and prolong the problem.

Primary herpes simplex infection

This viral infection produces a follicular conjunctivitis. About 50% of patients have associated lid or corneal ulcers/vesicles which are diagnostic.[2] Only a minority (less than 15%) develop corneal involvement with the primary infection.

Dendritic ulceration highlighted by fluorescein staining is diagnostic (Figure 51.4). Antigen detection or culture may allow confirmation.



Figure 51.4 Herpes simplex keratitis - gritty, watery eye with typical dendritic ulcer, stained with fluorescein

Treatment of herpes simplex keratitis

- Attend to eye hygiene
- Aciclovir 3% ointment, 5 times a day for 14 days or for at least 3 days after healing[7]
- Atropine 1% 1 drop, 12 hourly, for the duration of treatment will prevent reflex spasm of the pupil (specialist supervision)
- Debridement by a consultant

Allergic conjunctivitis

Allergic conjunctivitis results from a local response to an allergen. It includes:

- vernal (hay fever) conjunctivitis, and
- contact hypersensitivity reactions, e.g. reaction to preservatives in drops

Vernal (hay fever) conjunctivitis

This is usually seasonal and related to pollen exposure. There is usually associated rhinitis.

Treatment

Tailor treatment to the degree of symptoms. Antihistamines may be required but symptomatic measures usually suffice.

Treatment options:

1. Topical antihistamines/vasoconstrictors
2. Mast cell stabilisers, e.g. sodium cromoglycate 2% drops, 1 - 2 drops per eye 4 times daily
3. Combination of l and 2
4. Topical steroids (severe cases) Artificial tear preparations may give adequate symptomatic relief.

Contact hypersensitivity

Common topical allergens and toxins include topical ophthalmic medications, especially antibiotics, contact lens solutions (oft en the contained preservative) and a wide range of cosmetics, soaps, detergents and chemicals. Clinical features include burning, itching and watering with hyperaemia and oedema of the conjunctiva and eyelids. A skin reaction of the lids usually occurs.

Treatment

- Withdraw the causative agent.
- Apply normal saline compresses.
- Treat with naphazoline or phenylephrine.
- If not responding, refer for possible corticosteroid therapy.

Chlamydial conjunctivitis

Chlamydial conjunctivitis is encountered in three common situations:

- neonatal infection (first 1 - 2 weeks)
- young patient with associated venereal infection
- isolated Aboriginal people with trachoma Take swabs for culture and PCR testing

Systemic antibiotic treatment:[7]

- Neonates: erythromycin for 3 weeks
- Children over 6 kg and adults: azithromycin 1 g (o) as single dose

Note: Partner must be treated in cases of STI.

SUBCONJUNCTIVAL HAEMORRHAGE

Subconjunctival haemorrhage, which appears spontaneously, is a beefy red localised haemorrhage with a definite posterior margin (Figure 51.5). If it follows trauma and extends backwards it may indicate an orbital fracture. It is usually caused by a sudden increase in intrathoracic pressure such as coughing and sneezing. It is not related to hypertension but it is worthwhile measuring the blood pressure to help reassure the patient.



Figure 51.5 Subconjunctival haemorrhage: this is usually localised haemorrhage that appears spontaneously. It is pain free. If traumatic and extends posteriorly it may indicate an orbital fracture

Management

No local therapy is necessary. The haemorrhage absorbs over 2 weeks. Patient explanation and reassurance is necessary. If haemorrhages are recurrent a bleeding tendency should be excluded.

EPISCLERITIS AND SCLERITIS

Episcleritis and scleritis present as a localised area of inflammation (Figure 51.1 and Figure 51.2). The episclera lies just beneath the conjunctiva and adjacent to the sclera. Both may become inflamed but episcleritis (which is more localised) is essentially self-limiting while scleritis (which is rare) is more serious as the eye may perforate.[3] Both conditions may be confused with inflammation associated with a foreign body, pterygium or pinguecula. There are no significant associations with episcleritis, which is usually idiopathic, but scleritis may be associated with connective tissue disease especially rheumatoid arthritis, herpes zoster and rarely sarcoidosis and tuberculosis.

Features

Episcleritis:

- no discharge
- no watering
- vision normal (usually)
- often sectorial

- usually self-limiting

Treat with topical or oral steroids.

Scleritis:

- painful loss of vision
- urgent referral

History

A red and sore eye is the presenting complaint. There is usually no discharge but there may be reflex lacrimation. Scleritis is much more painful than episcleritis3 and the eye becomes intensely red.

Examination

With scleritis there is a localised area of inflammation that is tender to touch, and more extensive than with episcleritis, being uniform across the eye. The inflamed vessels are larger than the conjunctival vessels.

Management

An underlying cause such as an autoimmune condition should be identified. Refer the patient, especially for scleritis. Corticosteroids or NSAIDs may be prescribed.

UVEITIS (IRITIS)

The iris, ciliary body and the choroid form the uveal tract, which is the vascular coat of the eyeball.[6]

Anterior uveitis (acute iritis or iridocyclitis) is inflammation of the iris and ciliary body and this is usually referred to as acute iritis (Figure 51.6). The iris is sticky and sticks to the lens. The pupil may become small because of adhesions, and the vision is blurred.


ocular (bulbar) conjunctiva
eyelid
episclera (episcleritis)
blepharitis
sclera
choroid
cornea
(keratitis)
retina
posterior
uveitis
anterior
uveitis
iris
ciliary body

Figure 51.6 Diagrammatic representation of eye structures involved in inflammatory disorders

Causes include autoimmune related diseases such as the seronegative arthropathies (e.g. ankylosing spondylitis), SLE, IBD, sarcoidosis and some infections (e.g. toxoplasmosis and syphilis).

Features

- Eye redness, esp. around edge of iris
- Eye discomfort or pain
- Increased tearing
- Blurred vision
- Sensitivity to light
- Floaters in the field of vision
- Small pupil

The examination findings are summarised in Table 51.2 at p. 554. The affected eye is red with the injection being particularly pronounced over the area covering the inflamed ciliary body (cilary flush). However, the whole bulbar conjunctivae can be injected. The patient should be referred to a consultant. Slit lamp examination aids diagnosis.

Management involves finding the underlying cause. Treatment includes pupil dilatation with atropine drops and topical steroids to suppress inflammation. Systemic corticosteroids may be necessary. The prognosis of anterior uveitis is good if treatment and follow-up are maintained, but recurrence is likely.

Posterior uveitis (choroiditis) may involve the retina and vitreous. Blurred vision and floating opacities in the visual field may be the only symptoms. Pain is not a feature. Referral to detect the causation and for treatment is essential.

ACUTE GLAUCOMA

Acute glaucoma should always be considered in a patient over 50 years presenting with an acutely painful red eye. Permanent damage will result from misdiagnosis. The attack characteristically strikes in the evening when the pupil becomes semidilated.[3]

Features

- Patient > 50 years
- Pain in one eye
- ±Nausea and vomiting
- Impaired vision
- Haloes around lights
- Hazy cornea
- Fixed semidilated pupil
- Eye feels hard

Management

Urgent ophthalmic referral is essential since emergency treatment is necessary to preserve eyesight. If immediate specialist attention is unavailable, treatment can be initiated with acetazolamide (Diamox) 500 mg IV and pilocarpine 4% drops to constrict the pupil.

KERATOCONJUNCTIVITIS SICCA

Dry eyes are a common problem, especially in elderly women. Lack of lacrimal secretion can be functional (e.g. ageing), or due to systemic disease (e.g. rheumatoid arthritis, SLE, Sjögren's syndrome), drugs (e.g. β-blockers) or other factors, including the menopause. Up to 50% of patients with severe dry eye have Sjögren's syndrome.

Features

- A variety of symptoms
- Dryness, grittiness, stinging and redness
- Sensation of foreign body, e.g. sand
- Photophobia if severe
- Slit light examination diagnostic with special stains

Treatment

- Treat the cause.
- Bathe eyes with clean water.
- Use artificial tears: hypromellose (e.g. Tears Naturale), polyvinyl alcohol (e.g. Tears Plus).
- Be cautious of adverse topical reactions.
- Refer severe cases.

EYELID AND LACRIMAL DISORDERS

There are several inflammatory disorders of the eyelid and lacrimal system that present as a 'red and tender' eye without involving the conjunctiva. Any suspicious lesion should be referred.

Stye (external hordeolum)

A stye is an acute abscess of a lash follicle or associated glands of the anterior lid margin, caused usually by *S. aureus*. The patient complains of a red tender swelling of the lid margin, usually on the medial side (Figure 51.7). A stye may be confused with a chalazion, orbital cellulitis or dacryocystitis.



Figure 51.7 Hordeolum - a stye. This is a focal staphylococcal infection of the root of an eyelash

Management

- Use heat to help it discharge by using direct steam from a thermos (Figure 51.8) onto the enclosed eye or by hot compresses.
- Perform lash epilation to allow drainage of pus (incise with a D_{11} blade if epilation does not work).
- Use chloramphenicol ointment if the infection is spreading locally.[3]



Figure 51.8 Steaming the painful eye: allow steam to rise from a thermos onto the closed eye for 10 - 15 minutes

Chalazion (meibomian cyst)

Also known as internal hordeolum, this granuloma of the meibomian gland in the eyelid may become inflamed and present as a tender irritating lump in the lid. Look for evidence of blepharitis.

Management

Conservative treatment may result in resolution. This involves heat either as steam from a thermos or by applying a hot compress (a hand towel soaked in hot water) and the application of chloramphenicol ointment for 5 days. If the chalazion is very large, persistent or uncomfortable, or is affecting vision, it can be incised and curetted under local anaesthesia. This is best performed through the inner conjunctival surface using a chalazion clamp (blepharostat) (Figure 51.9).



Figure 51.9 Excision of a meibomian cyst, using a chalazion clamp and curette

Meibomianitis is usually a staphlyococcal micro abscess of the gland and oral antistaphylococcal antibiotics (not topical) are recommended, e.g. di/flucloxacillin 500 mg (o) 6 hourly (adult). Surgical incision and curettage may also be necessary.

Blepharitis

This common chronic condition is characterised by inflammation of the lid margins and is commonly associated with secondary ocular effects such as styes, chalazia and conjunctival or corneal ulceration. (Figure 51.10) Blepharitis is frequently associated with seborrhoeic dermatitis (especially) and atopic dermatitis, and less so with rosacea.8 There is a tendency to colonisation of the lid margin with *S. aureus*, which causes an ulcerative infection.



Figure 51.10 Blepharitis: common complicating features

The three main types are:

- seborrhoeic blepharitis
- staphylococcal blepharitis
- blepharitis associated with rosacea

Features[8]

- Persistent sore eyes or eyelids
- Irritation, grittiness, burning, dryness and 'something in the eye' sensation
- Lid or conjunctival swelling and redness
- Crusts or scales around the base of the eyelids
- Discharge or stickiness, especially in morning
- Inflammation and crusting of the lid margins

Management

- Eyelid hygiene is the mainstay of therapy. The crusts and other debris should be gently cleaned with a cotton wool bud dipped in a 1:10 dilution of baby shampoo or a solution of sodium bicarbonate, once or twice daily. Application of a warm water or saline soak with gauze for 20 minutes is also effective.
- For chronic blepharitis short-term use of a corticosteroid ointment (e.g. hydrocortisone 0.5%) can be very effective.
- Ocular lubricants such as artificial tear preparations may greatly relieve symptoms of keratoconjunctivitis sicca (dry eyes).
- Control scalp seborrhoea with regular medicated shampoos.
- Treat infection with an antibiotic ointment smeared on the lid margin (this may be necessary for several months) (e.g. tetracycline hydrochloride 1% or framycetin 0.5% or chloramphenicol 1% ointment to lid margins 3 - 6 hourly).[7]
- Systemic antibiotics such as flucloxacillin may be required for lid abscess.
- Avoid wearing make-up and contact lenses if inflammation is present.

Dacryocystitis

Acute dacryocystitis is infection of the lacrimal sac secondary to obstruction of the nasolacrimal duct at the junction of the lacrimal sac (Figure 51.11). Inflammation is localised over the medial canthus. There is usually a history of a watery eye for months beforehand. The problem may vary from being mild (as in infants) to severe with abscess formation.



Figure 51.11 Acute dacrocystitis with abscess formation. Associated obstruction of the nasolacrimal duct at the junction of the lacrimal sac which has become infected

Management

- Use local heat: steam or a hot moist compress.
- Use analgesics.
- In mild cases, massage the sac and duct, and instil astringent drops (e.g. zinc sulfate + phenylephrine).
- For acute cases systemic antibiotics are best guided by results of Gram's stain and culture but initially use di/flucloxacillin.
- Measures to establish drainage are required eventually. Recurrent attacks or symptomatic watering of the eye are indications for surgery such as dacryocystorhinostomy.

ORBITAL CELLULITIS

Orbital cellulitis includes two basic types - periorbital (or preseptal) and orbital (or postseptal) cellulitis. The latter is a potentially blinding and life-threatening condition. It is especially important in children in whom blindness may develop in hours. The patient, often a child, presents with unilateral swollen eyelids that may be red.

Features to look for in orbital cellulitis include:[3]

- an unwell patient
- tenderness over the sinuses
- restricted and painful eye movements (Figure 51.12)



Figure 51.12 Important signs in the patient presenting with orbital cellulitis

In periorbital cellulitis, which usually follows an abrasion, there is no pain or restriction of eye movement (Figure 51.13).



Figure 51.13 Periorbital cellulitis following an abrasion to the eye. Treat this as an urgent condition

Immediate referral to hospital for specialist treatment is essential for both types. Treatment is with IV cefotaxime until afebrile, then amoxycillin/clavulanate for 7 - 10 days for periorbital cellulitis and for orbital cellulitis, IV cefotaxime + di(flu) cloxacillin together followed by amoxycillin/clavulanate (o) 10 days.[7]

HERPES ZOSTER OPHTHALMICUS

Herpes zoster ophthalmicus (shingles) affects the skin supplied by the ophthalmic division of the trigeminal nerve. The eye may be affected if the nasociliary branch is involved. Ocular problems include conjunctivitis, uveitis, keratitis and glaucoma.

Immediate referral is necessary if the eye is red, vision is blurred or the cornea cannot be examined. Apart from general eye hygiene, treatment usually includes one of the oral anti-herpes virus agents such as oral aciclovir 800 mg, 5 times daily for 10 days or (if sight is threatened) aciclovir 10 mg/kg IV slowly 8 hourly for 10 days (provided this is commenced within 3 days of the rash appearing)[5, 7] and topical aciclovir ointment 4 hourly.

PINGUECULA AND PTERYGIUM[9]

Pinguecula is a yellowish elevated nodular growth on either side of the cornea in the area of the palpebral fissure. It is common in people over 35 years. The growth tends to remain static but can become inflamed - pingueculitis. Usually no treatment is necessary unless they are large, craggy and uncomfortable, when excision is indicated. If irritating, topical astringent drops such as naphazoline compound drops (e.g. Albalon) can give relief.

Pterygium is a fleshy overgrowth of the conjunctiva onto the nasal side of the cornea and usually occurs in adults living in dry, dusty, windy areas. Excision of a pterygium by a specialist is indicated if it is likely to interfere with vision by encroaching on the visual axis, or if it becomes red and uncomfortable or disfiguring.

CORNEAL DISORDERS[10]

Patients with corneal conditions typically suffer from ocular pain or discomfort and reduced vision. The common condition of dry eye may involve the cornea while contact lens disorders, abrasions/ulcers and infection are common serious problems that threaten eyesight. Inflammation of the cornea - **keratitis** - is caused by factors such as ultraviolet light, e.g. 'arc eye', herpes simplex, herpes zoster ophthalmicus and the dangerous microbiol keratitis. Bacterial keratitis is an ophthalmological emergency which should be considered in the contact lens wearer presenting with pain and reduced vision.

Topical corticosteroids should be avoided in the undiagnosed red eye.

Corneal abrasion and ulceration

There are many causes of abrasions, particularly trauma from a foreign body embedded on the corneal surface or 'cul-de-sac' FB, contact lenses, fingernails including 'french nails', and ultraviolet burns. The abrasion may be associated with an ulcer, which is a defect in the epithelial cell layer of the cornea. Common causes of a corneal ulcer are listed in Table 51.3.

Table 51.3 Corneal ulceration: common causes

Trauma
Contact lens wear/injury
Infection - microbiol keratitis
• bacterial, e.g. *Pseudomonas* (contact lens)
• viral, e.g. Herpes simplex (dendritic ulcer), Herpes zoster ophthalmacus
• fungal
• protozoa, e.g. acanthamoeba
Neurotrophic, e.g. trigeminal nerve defect
Immune related, e.g. rheumatoid arthritis
Spontaneous corneal erosion
Chronic blepharitis
Overexposure, e.g. eyelid defects

Symptoms

- Ocular pain
- Foreign body sensation
- Watering of the eye (epiphoria)
- Blepharospasm
- Blurred vision

Diagnosis is best performed with a slit lamp using a cobalt blue filter and flourescein staining.

> **Practice tip**
>
> If a slit lamp is unavailable, the direct ophthalmoscope can be used to provide illumination as well as blue light for corneal examination. Magnifying loupes can then be used for viewing the illuminated cornea.

Basic management of corneal ulcer

- Stain with fluoroscein.
- Check for a foreign body.
- Treat with chloramphenicol 1% ointment ± homatropine 2% (if pain due to ciliary spasm).
- Double eye pad (if not infected).
- Review in 24 hours.
- Consider specialist referral.

Superficial punctate keratitis

Punctuate keratopathy presents as scattered small lesions on the cornea which stain with fluoroscein if they are deep enough. It is a non-specific finding and may be associated with blepharitis, viral conjunctivitis, trachoma, keratitis sicca (dry eyes), UV light exposure (e.g. welding lamps, sunlamps), contact lenses and topical ocular agents. Management involves treating the cause and careful follow-up.

Microbiol keratitis

This is responsible for at least 1.5 million new cases of blindness every year in the developing world and for significant morbidity in developed countries.

Risk factors

- Contact lens wear
- Corneal trauma, especially agriculture trauma
- Corneal surgery
- Post-herpetic corneal lesion
- Dry eye

- Corneal anaesthetic
- Corneal exposure, e.g. VII nerve lesion
- Ocular surface disease including ulceration

Pseudomonas aeruginosa is the most common causative organism in contact lens wearers.

Acanthamoeba is associated with bathing or washing in contaminated water.

Urgent referral to an ophthalmologist or eye clinic is needed to avoid rapid corneal destruction with perforation especially with bacterial keratitis. An appropriate ‘covering’ topical antibiotic is ciprofloxacin 0.3% ointment.

PROBLEMS WITH CONTACT LENSES

Because a contact lens is a foreign body, various complications can develop and a history of the use of contact lenses is important in the management of the red eye.

Infection

Infection is more likely to occur with soft rather than hard lenses. They should not be worn for sleeping since this increases the risk of infection 10-fold.[11] One cause is *Acanthamoeba* keratitis acquired from contaminated water that may be used for cleaning the lenses.

Hard lens trauma

This may cause corneal abrasions with irreversible endothelial changes or ptosis, especially with the older polymethyl methacrylate based lenses. Patients should change to modern gas permeable hard lenses.

Lost lenses

Patients should be reassured that lenses cannot go behind the eye. The edge of the lens can usually be seen by everting the upper lid.

Preventive measures[12]

- Wash hands before handling lenses.
- Do not use tap water or saline.
- Clean lenses with disinfecting solution.
- Store overnight in a clean airtight case with fresh disinfectant.
- Change the lens container solution daily.
- Discard disposable lenses after 2 weeks.
- Do not wear lenses while sleeping.
- Do not wear lenses while swimming in lakes, rivers or swimming pools.

Refer to an ophthalmologist if a painful red eye develops.

FLASH BURNS

A common problem, usually presenting at night, is bilateral painful eyes caused by ultraviolet (UV) ‘flash burns’ to both corneas some 5 - 10 hours previously. The mechanism of injury is ultraviolet rays from a welding machine causing superficial punctate keratitis. Other sources of UV light such as sunlamps and snow reflection can cause a reaction.

Management

- Local anaesthetic (long-acting) drops: once only application (do not allow the patient to take home more drops).
- Instil homatropine 2% drops statim or other short-acting ocular dilating agent (be careful of glaucoma).
- Use analgesics (e.g. codeine plus paracetamol) for 24 hours.
- Use broad spectrum antibiotic eye ointment in lower fornix (to prevent infection).
- Use firm eye padding for 24 hours, when eyes reviewed (avoid light).

The eye usually heals completely in 48 hours. If not, check for a foreign body.

Note: Contact lens ‘overwear syndrome’ gives the same symptoms.

CAVERNOUS SINUS ARTERIOVENOUS FISTULA

Such a fistula produces conjunctival hyperaemia but no inflammation or discharge. The lesion causes raised orbital venous pressure. The fistula may be secondary to head injuries or may arise spontaneously, particularly in postmenopausal women. They need radiological investigation.

The classic symptom is a ‘whooshing’ sound synchronous with the pulse behind the eye, and the sign is a bruit audible with the stethoscope placed over the orbit.

PENETRATING EYE INJURIES

These require urgent referral to an ophthalmologist. Consider:

- X-ray
- tetanus prophylaxis
- transport by land
- injection of anti-emetic, e.g. metoclopramide

Use no ointment or eye drops, including local anaesthetic.

If significant delay is involved give one dose (in adults) of:[7]

- gentamicin 1.5 mg/kg IV plus cefotaxime 1 g or ceftriaxone 1 g IV (can give ceft riaxone IM but with lignocaine 1%) *or*
- vancomycin IV + oral ciprofloxacin

ENDOPHTHALMITIS

This is an intraocular bacterial infection which may complicate any penetrating injury including intra-ocular sugery. It should be considered in patients with such a history presenting with a red painful eye. Pus may be seen in the anterior chamber (hypopyon).

Urgent referral is mandatory.

Practice tips

- Avoid long-term use of any medication, especially antibiotics (e.g. chloramphenicol: course for a maximum of 10 days).[2] *Note:* Beware of allergy or toxicity to topical medications, especially antibiotics, as a cause of persistent symptoms.
- As a general rule avoid using topical corticosteroids or combined corticosteroid/antibiotic preparations.
- Never use corticosteroids in the presence of a dendritic ulcer.
- To achieve effective results from eye ointment or drops, remove debris such as mucopurulent exudate with bacterial conjunctivitis or blepharitis by using a warm solution of saline (dissolve a teaspoon of kitchen salt in 500 mL of boiled water) to bathe away any discharge from conjunctiva, eyelashes and lids.
- A gritty sensation is common in conjunctivitis but the presence of a foreign body must be excluded.[3]
- Beware of the contact lens ‘overwear syndrome’, which is treated in a similar way to flash burns.

WHEN TO REFER

- Uncertainty about the diagnosis
- Patients with uveitis, acute glaucoma, episcleritis/scleritis or corneal ulceration
- Deep central corneal and intraocular foreign bodies
- Prolonged infections, with a poor or absent response to treatment or where therapy may be complicating management
- Infections or severe allergies with possible ocular complications
- Sudden swelling of an eyelid in a child with evidence of infection suggestive of orbital cellulitis - this is an emergency
- Emergency referral is also necessary for hyphaema, hypopyon, penetrating eye injury, acute glaucoma, severe chemical burn
- Herpes zoster ophthalmicus: if the external nose is involved then the internal eye may be involved
- Summary for urgent referral:
    - trauma (significant)/penetrating injury
    - hyphaema > 3 mm

- corneal ulcer
- severe conjunctivitis
- uveitis/acute iritis
- Behçet's syndrome
- acute glaucoma
- giant cell arteritis
- orbital cellulitis (pre- and post-)
- acute dacryocystitis
- keratitis
- episcleritis/scleritis
- endophthalmitis
- herpes zoster ophthalmicus

Note: As a general rule never use corticosteroids or atropine in the eye before referral to an ophthalmologist.

REFERENCES

1. McDonnell P. Red eye: an illustrated guide to eight common causes. Mod Med Aust, 1989; October: 37 - 9.
2. Della NG. Acute conjunctivitis. In: *MIMS Disease Index*. Sydney: IMS Publishing: 113 - 15.
3. Elkington AR, Khaw PT. *ABC of Eyes*. London: British Medical Association, 1990: 6 - 10.
4. Colvin J. Systematic examination of the red eye. Aust Fam Physician, 1976; 5: 153 - 65.
5. Maclean H. Keratitis (viral and fungal). In: *MIMS Disease Index*. Sydney: IMS Publishing, 1991 - 92: 301 - 3.
6. Robinson MJ, Roberton DM. *Practical Paediatrics* (5th edn). Melbourne: Churchill Livingstone, 2003: 759.
7. Spicer J (Chair). *Therapeutic Guidelines: Antibiotics* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 69 - 78.
8. Barras CW. Blepharitis. In: *MIMS Disease Index*. Sydney: IMS Publishing, 1991 - 92: 80 - 2.
9. Colvin J. Painful eye: an emergency call. Aust Fam Physician, 1985; 14: 1258.
10. Watson SL. Common corneal conditions. Medicine Today, 2005; 6(5): 22 - 30.
11. Schein OD, Poggio EC. Ulcerative keratitis in contact lens wearers. Cornea, 1990; 9(1): 55 - 8.
12. Lazarus MG. Complications of contact lenses. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 121 - 3.

CHAPTER 52. PAIN IN THE FACE

It's as though the devil suddenly thrust red hot electric needles through my right cheek towards my ear.

PATIENT (ANONYMOUS), DESCRIBING ' TIC DOULOUREUX '

PAIN IN THE FACE

When a patient complains of pain in the face rather than the head, the physician has to consider foremost the possibilities of dental disorders, sinus disease, especially of the maxillary sinuses, TMJ dysfunction, eye disorders, lesions of the oropharynx or posterior third of the tongue, trigeminal neuralgia and chronic paroxysmal hemicrania.

The key to the diagnosis is the clinical examination because even the most sophisticated investigation may provide no additional information.

A basic list of causes of facial pain is presented in Table 52.1.[1] The causes can vary from the simple, such as aphthous ulcers, herpes simplex and dental caries, to serious causes, such as carcinoma of the tongue, sinuses and nasopharynx or osteomyelitis of the mandible or maxilla.

Table 52.1 Diagnoses to consider in orofacial pain

Positive physical signs
Cervical spinal dysfunction
Dental pathology
Erysipelas
Eye disorders
Herpes zoster
Nasopharyngeal carcinoma
Oropharyngeal disorders - ulceration (aphthous, infective, traumatic, others) - carcinoma - gingivitis/stomatitis - tonsillitis - erosive lichen planus
Paranasal sinus disorders
Parotid gland - mumps - sialectasis - carcinoma - pleomorphic adenoma
TMJ dysfunction
Temporal arteritis

KEY FACTS AND CHECKPOINTS

- Dental disorders are the commonest cause of facial pain, accounting for up to 90% of pain in and about the face.[2]
- The most common dental disorders are dental caries and periodontal diseases.
- Trigeminal neuralgia is relatively uncommon with a prevalence of 155 persons per million of the population.[3]
- The mean age of onset of trigeminal neuralgia is 50 - 52 years.

- There is a similarity in the 'occult' causes of pain in the ear and in the face (refer to Figures 50.3 and 50.4).
- Sinusitis occurs mainly as part of a generalised upper respiratory infection. Swimming is another common predisposing factor.
- Dental root infection must be sought in all cases of maxillary sinusitis.

Absent physical signs

Atypical facial pain

Chronic paroxysmal hemicrania

Depression-associated facial pain

Facial migraine (lower half headache)

Glossopharyngeal neuralgia

Migrainous neuralgia (cluster headache)

Trigeminal neuralgia (tic douloureux)

A diagnostic approach

A summary of the safety diagnostic model is presented in Table 52.2.

Table 52.2 Pain in the face: diagnostic strategy model

Q. Probability diagnosis
A. Dental pain - caries
- periapical abscess
Maxillary sinusitis
Q. Serious disorders not to be missed
A. Cardiovascular
• myocardial ischaemia
• aneurysm of cavernous sinus
• internal carotid aneurysm
• ischaemia of posterior inferior cerebellar artery
Neoplasia
• carcinoma - mouth
- sinuses
- nasopharynx
- tonsils
- tongue
- larynx
• metastases - orbital
- base of brain
- bone
Severe infections
• erysipelas

- periapical abscess →osteomyelitis
- acute sinusitis →spreading infection

Temporal arteritis

Q. Pitfalls

A. TMJ dysfunction

Migraine variants

- facial migraine
- chronic paroxysmal hemicrania

Eye disorders - glaucoma, iritis, optic neuritis Chronic dental neuralgia

Parotid gland - mumps, carcinoma, sialectasis

Acute glaucoma (upper face)

Cranial nerve neuralgias

- trigeminal neuralgia
- glossopharyngeal neuralgia

Q. Seven masquerades checklist

A. Depression

Diabetes -

Drugs -

Anaemia -

Thyroid disorder -

Spinal dysfunction

UTI -

Q. Is the patient trying to tell me something?

A. Quite probably. Atypical facial pain has underlying psychogenic elements.

Probability diagnosis

The commonest cause of facial pain is dental disorders, especially dental caries. Another common cause is sinusitis, particularly maxillary sinusitis.

TMJ dysfunction causing TMJ arthralgia is a very common problem encountered in general practice and it is important to have some simple basic strategies to give the patient.

Serious disorders not to be missed

It is important not to overlook carcinoma of various structures, such as the mouth, sinuses, nasopharynx, tonsils, tongue, larynx and parotid gland, which can present with atypical chronic facial pain.

It is important therefore to inspect these areas, especially in the elderly, but lesions in the relatively inaccessible nasopharynx can be easily missed. Nasopharynx carcinoma spreads upwards to the base of the skull early and patients can present with multiple cranial nerve palsies before either pain or bloody nasal discharge.[1]

Tumours may arise in the bones of the orbit, for example, lymphoma or secondary carcinoma, and may cause facial pain and proptosis. Similarly, any space-occupying lesion or malignancy arising from the region of the orbit or base of the brain can cause facial pain by involvement (often destruction) of trigeminal sensory fibres. This will lead to a depressed ipsilateral corneal reflex.

Also, aneurysms developing in the cavernous sinus[1] can cause pain via pressure on any of the divisions of the trigeminal nerve, while aneurysms from the internal carotid arising from the origin of the pos terior communicating artery can cause pressure on the oculomotor nerve.

Temporal arteritis typically causes pain over the temporal area but can cause ischaemic pain in the jaws when chewing.

Pitfalls

Commonly overlooked causes of facial pain include TMJ arthralgia and dental disorders, especially of the teeth, which are tender to percussion, and oral ulceration. Diagnosing the uncommon migraine variants, particularly facial migraine and chronic paroxysmal hemicrania, often presents difficulties, including differentiating between the neuralgias. Glossopharyngeal neuralgia, which is rare, causes pain in the back of the throat, around the tonsils and adjacent fauces. The lightning quality of the pain of neuralgia gives the clue to diagnosis.

Common pitfalls

- Failing to refer unusual or undiagnosed causes of facial pain
- Overlooking infective dental causes, which can cause complications
- Failing to consider the possibility of malignant disease of ‘hidden’ structures in the older patient

Seven masquerades checklist

Of these, depression and cervical spinal dysfunction must be considered. The upper cervical spine can cause facial pain from lesions of C2 or C3 via the lesser occipital or greater auricular (Figure 52.1) nerves, which may give pain around the ear. It is important to remember that the C2 and C3 nerves share a common pathway with the trigeminal nerve (Chapter 62).



Figure 52.1 Dermatomes of C2 and C3, with the overlap area indicated

Depressive illness can present with a variety of painful syndromes and facial pain is no exception. The features of depression may be apparent and thus antidepressants should be prescribed. Usually the facial pain and the depression subside concomitantly.

Psychogenic considerations

Psychogenic factors have to be considered in every painful condition. They are considered to be high in patients with atypical facial pain.

The clinical approach

History

Diagnosis of nearly all types of facial pain must be based almost entirely on the history. It is often difficult to delineate the exact nature and distribution of the pain. The history should include the typical analysis of pain, especially noting the site and radiation of the pain.

Examination

The patient's general state and behaviour should be noted. Any sudden jabbing pain in the face causing the characteristic ‘tic’ may indicate neuralgia.

Palpate the face and neck to include the parotid glands, eyes, regional lymph nodes and the skin. Inspect the TMJs and cervical spine. Carefully inspect the nose, mouth, pharynx and postnasal space. In particular inspect the teeth, percussing each tooth if dental disorder is suspected. Bimanual palpation of the floor of the mouth is performed to detect induration or submandibular and submental lymph node enlargement.

The sinuses, especially the maxillary sinuses, should be inspected and a torch light should be placed inside the mouth to test transillumination of the maxillary sinuses. It works best when one symptomatic side can be compared with an asymptomatic side.

Perform a neurological examination on the cranial nerves with special emphasis on the trigeminal, oculomotor and glossopharyngeal nerves.

Investigations

If investigations are being contemplated referral may be appropriate. The association of multiple sclerosis and tumours with neuralgias may have to be investigated. Radiological investigations include plain X-rays of the paranasal sinuses, CT scans, MRI and orthopantomograms.

Facial pain in children

Apart from trauma, facial pain in children is invariably due to dental problems, rarely migraine variants and occasionally childhood infections such as mumps and gingivostomatitis. A serious problem sometimes seen in children is orbital cellulitis secondary to ethmoiditis.

Sinusitis occurs in children, especially older children, and it should be suspected with persistent bilateral mucopurulent rhinorrhoea (beyond 10 days).

Facial pain in the elderly

Many of the causes of facial pain have an increased incidence with age, in particular trigeminal neuralgia, herpes zoster, carcinoma, glaucoma, TMJ dysfunction and cervical spondylosis. Glossopharyngeal neuralgia does not seem to have a particular predilection for the elderly. Xerostomia due to decreased secretions of salivary glands may cause abrasion with minor trauma. It may aggravate the pain of glossitis, which is common in the elderly.

Dental disorders

Dental caries, impacted teeth, infected tooth sockets and dental roots can cause pain in the maxillary and mandibular regions. Caries with periapical and apical abscess formation produces pain from infection extending around the apex of the tooth into the alveolar bone. Retention of a fractured root may cause unilateral paroxysmal pain. Impacted third molars (wisdom teeth) may be associated with surrounding softtissue inflammation (pericoronitis), causing pain that may be localised to the mandible or radiate via the auriculotemporal nerve to the ear. *Candida albicans*, which is an oral commensal, may colonise dentures causing hyperaemia and painful superficial ulceration of the denture-bearing mucosa.

Features of dental caries

- Pain is usually confined to the affected tooth but it may be diffuse.
- Pain is almost always aggravated by thermal changes in the mouth:
    - cold - if dental pulp vital
    - hot - if dental pulp is necrotic
- Pain may be felt in more than one tooth.
- Dental pain will not cross the midline.

Treatment of dental pain

- Arrange urgent dental consultation.
- Pain relief:[4]
    - aspirin 600 mg (o) 4 - 6 hourly *or*

  - paracetamol 0.5 - 1 g (o) 4 - 6 hourly
- If pain severe add:
  - codeine 30 mg (o) 4 - 6 hourly

Tooth abscess, inflamed wisdom tooth or root canal infection [5]

Dental treatment will usually alleviate the problem; however, if severe:

- amoxycillin 500 mg (o) tds for 5 days if unresponsive add:
- metronidazole 400 mg (o), 12 hourly for 5 days For patients hypersensitive to penicillin:
- clindamycin 300 mg (o), 8 hourly for 5 days

Gingivitis and periodontitis

Refer to Chapter 72.

Alveolar osteitis (dry socket)

Refer for specialised toileting. This usually heals naturally in 14 days. Antibiotics of no proven use (Figure 52.2). [5]



Figure 52.2 Dry tooth socket: this is a very painful condition mainly in the lower molar unrelieved by analgesics following a toothextraction 1 - 3 days earlier. The socket has few or no blood clots and sensitive bone surfaces covered by a grayish layer of necrotic tissue

Ludwig's angina

This is a rapidly swelling cellulitis occuring in both the sublingual and submaxillary spaces without abscess formation, often arising from a root canal infection. It resembles an abscess and should be treated as one. It is potentially life-threatening as it can compromise the airway.

Management

- Culture and sensitivity testing
- Specialist consultation Empirical treatment:
- Amoxycillin 2 g IV, 6 hourly *plus* metronidazole 500 mg IV 12 hourly

Pain from paranasal sinuses

Infection of the paranasal sinuses may cause localised pain. Localised tenderness and pain may be apparent with frontal or maxillary sinusitis. Sphenoidal or ethmoidal sinusitis causes a constant pain behind the eye or behind the nose, often accompanied by nasal blockage. Chronic infection of the sinuses may be extremely difficult to detect. The commonest organisms are *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Moraxella catarrhalis*.

Expanding lesions of the sinuses, such as mucocoeles and tumours, cause local swelling and displace the contents of the orbit - upwards for maxillary, laterally for the ethmoids and downwards for the frontal.

Maxillary sinusitis

The maxillary sinus is the one most commonly infected.[6] It is important to determine whether the sinusitis is caused by stasis following a URTI or acute rhinitis, or due to dental root infection.

Clinical features of acute sinusitis

- Facial pain and tenderness (over sinuses)
- Toothache
- Headache
- Purulent postnasal drip
- Nasal discharge
- Nasal obstruction
- Rhinorrhoea
- Cough (worse at night)
- Prolonged fever
- Epistaxis

Suspect bacterial cause if high fever and purulent nasal discharge.

Clinical features of chronic sinusitis

- Vague facial pain
- Offensive postnasal drip
- Nasal obstruction
- Toothache
- Malaise
- Halitosis

Some simple office tests

Diagnosing sinus tenderness [7]

To differentiate sinus tenderness from non-sinus bone tenderness palpation is useful. This is best done by palpating a non-sinus area first and last (Figure 52.3), systematically exerting pressure over the temporal bones (T), then the frontal (F), ethmoid (E) and maxillary (M) sinuses, and finally zygomas (Z), or vice versa.



Figure 52.3 Diagnosing sinus tenderness: T (temporal) and Z (zygoma) represent no sinus bony tenderness, for purposes of comparison

Differential tenderness both identifies and localises the main sites of infection (Figure 52.3).

Diagnosis of unilateral sinusitis

A simple way to assess the presence or absence of fluid in the frontal sinus, and in the maxillary sinus (in particular), is the use of transillumination. It works best when one symptomatic side can be compared with an asymptomatic side.

It is necessary to have the patient in a darkened room and to use a small, narrow-beam torch. For the maxillary sinuses remove dentures (if any). Shine the light inside the mouth, on either side of the hard palate, pointed at the base of the orbit. A dull glow seen below the orbit indicates that the antrum is air-filled. Diminished illumination on the symptomatic side indicates sinusitis.

Management of acute bacterial sinusitis

Principles

- Exclude dental root infection.
- Control predisposing factors.
- Use appropriate antibiotic therapy.

- Establish drainage by stimulation of mucociliary flow and relief of obstruction.

Guidelines for antibiotic therapy

Consider therapy for severe cases displaying at least three of the following:

- persistent mucopurulent nasal discharge (> 7 - 10 days)
- facial pain
- poor response to decongestants
- tenderness over the sinuses especially maxillary
- tenderness on percussion of maxillary molar and premolar teeth that cannot be attributed to by a single tooth

Measures

- analgesics
- antibiotics [5, 8]
  - amoxycillin 500 mg (o) tds for 7 days *or* (if sensitive to penicillin)
  - doxycycline 100 mg (o) bd for 7 days *or*
  - cefaclor 500 mg (o) tds for 7 days *or*
  - amoxycillin + clavulanate 875/125 mg (o) tds for 7 - 14 days if poor response to above (indicates resistant *H. influenzae*)
- In complicated or severe disease, use intravenous cephalosporins or flucloxacillin
- Nasal decongestants (ephedrine-containing nasal drops or sprays)6 only if congestion
- Inhalations (a very important adjunct)

Invasive methods

Surgical drainage may be necessary by atrial lavage or frontal sinus trephine.

Inhalations for sinusitis

The old method of towel over the head and inhalation bowl can be used, but it is better to direct the vapour at the nose. Equipment needed is a container, which can be an old disposable bowl, a wide-mouthed bottle or tin, or a plastic container.

For the inhalant, several household over-thecounter preparations are suitable such as friar's balsam (5 mL), Vicks Vaporub (1 teaspoon), or menthol (5 mL).

The cover can be made from a paper bag (with its base cut out), a cone of paper (Figure 52.4) or a small cardboard carton (with the corner cut away).



Figure 52.4 Method of inhalation for sinusitis

Method

1. Add 5 mL or 1 teaspoon of the inhalant to 0.5 L (or 1 pint) of boiled water in the container.
2. Place the paper or carton over the container.
3. Get the patient to apply nose and mouth to the opening and breathe in the vapour deeply and slowly through the nose, and then out slowly through the mouth.
4. This should be performed for 5 - 10 minutes, three times a day, especially before retiring.

After inhalation, upper airway congestion can be relieved by autoinsufflation.

Chronic sinusitis

Chronic sinusitis or recurrent sinusitis may arise from chronic infection or allergy. It may be associated with nosal polyps and vasomotor rhinitis, but is frequently associated with a structural abnormality of the upper airways.

It does not usually cause pain unless an acute infection intervenes. The acute or chronic attack is treated as for the acute attack but with 14 days of antibiotics. Those with an allergic mucous membrane may respond to intranasal corticosteroids. Surgical intervention will benefit chronic recurrence with mechanical blockage.

TMJ dysfunction

This condition is due to abnormal movement of the mandible, especially during chewing. The basic cause is dental malocclusion. The pain is felt over the joint and tends to be localised to the region of the ear and mandibular condyle but may radiate forwards to the cheek and even the neck.

Examination of the TMJ

- Check for pain and limitation of mandibular movements, especially on opening the mouth.
- Palpate about the joint bilaterally for tenderness, which typically lies immediately in front of the external auditory meatus; palpate the temporalis and masseter muscles.
- Palpate the TMJ over the lateral aspect of the joint disc.
- Ask the patient to open the mouth fully when tenderness is maximal. The TMJ can be palpated posteriorly by inserting the little finger into the external canal.
- Check for crepitus in mandibular movement.

Treatment of TMJ dysfunction

If organic disease such as rheumatoid arthritis and obvious dental malocclusion is excluded, a special set of instructions or exercises can alleviate the annoying problem of TMJ arthralgia in about 3 weeks.

Method 1: ‘Chewing’ the piece of softwood

- Obtain a rod of softwood approximately 15 cm long and 1.5 cm wide. An ideal object is a large carpenter's pencil.
- Instruct the patient to position this at the back of the mouth so that the molars grasp the object with the mandible thrust forward.
- The patient then rhythmically bites on the object with a grinding movement for 2 - 3 minutes at least 3 times a day.

Method 2: The ‘six by six’ program

This is a specific program recommended by some dental surgeons. The six exercises should be carried out six times on each occasion, six times a day, taking about 1 - 2 minutes.

Instruct the patient as follows:

1. Hold the front one-third of your tongue to the roof of your mouth and take six deep breaths.
2. Hold the tongue to the roof of your mouth and open your mouth six times. Your jaw should not click.
3. Hold your chin with both hands keeping the chin still. Without letting your chin move, push up, down and to each side. Remember, do not let your chin move.
4. Hold both hands behind your neck and pull chin in.
5. Push on upper lip so as to push head straight back.
6. Pull shoulders back as if to touch shoulder blades together.

These exercises should be pain-free. If they hurt, do not push them to the limit until pain eases.

Method 3: The TMJ ‘rest’ program

This program is reserved for an acutely painful TMJ condition.

- For eating avoid opening your mouth wider than the thickness of your thumb and cut all food into small pieces.
- Do not bite any food with your front teeth - use small bite-size pieces.
- Avoid eating food requiring prolonged chewing (e.g. hard crusts of bread, tough meat, raw vegetables).
- Avoid chewing gum.
- Always try to open your jaw in a hinge or arc motion. Do not protrude your jaw.
- Avoid protruding your jaw (e.g. talking, applying lipstick).
- Avoid clenching your teeth together - keep your lips together and your teeth apart.
- Try to breathe through your nose at all times.

- Do not sleep on your jaw: try to sleep on your back.
- Practise a relaxed lifestyle so that your jaws and face muscles feel relaxed.

Injection into the TMJ

Indications: painful rheumatoid arthritis, osteoarthritis or TMJ dysfunction not responding to conservative measures.

Method

- The patient sits on a chair, facing away from the therapist. The mouth is opened to at least 4 cm.
- The joint line is palpated anterior to the tragus of the ear: this is confirmed by the opening and closing of the jaw. A 25 gauge needle is inserted into the depression above the condyle of the mandible, below the zygomatic arch and one finger breadth (2 cm) anterior to the tragus. The needle is directed inwards and slightly upwards to lie free within the joint cavity. The 1 mL solution containing 0.5 mL of local anaesthetic and 0.5 mL of corticosteroid should flow quite freely.[9]

Other treatments

- Dental management that may be required for malfunction of the bite includes dental occlusal splinting.
- NSAIDs: A trial of NSAIDs, e.g. ibuprofen 400 mg (0) tds for 10 days, for TMJ inflammation may need consideration. Cease if no response after 10 days.

Inflammatory or ulcerative oropharyngeal lesions

A variety of ulcerative conditions and infections of structures such as gingivae, tongue, tonsils, larynx and pharynx can cause facial pain (refer to Chapter 72). Gingivostomatitis, herpes labialis (cold sores) and aphthous ulceration are common examples. Lesions of the posterior third of the tongue, the oropharynx, tonsils and larynx may radiate to the region of the ear via the tympanic branch of the ninth nerve or the auricular branch of the tenth nerve.

Trigeminal neuralgia

Trigeminal neuralgia (tic douloureux) is a condition of often unknown cause that typically occurs in patients over the age of 50, affecting the second and third divisions of the trigeminal nerve and on the same side of the face.[3] Brief paroxysms of pain, often with associated trigger points, are a feature.

Typical clinical features

- Site: sensory branches of the trigeminal nerve (Figure 52.5) almost always unilateral (often right side)



Figure 52.5 Typical cutaneous sensory distribution of the trigeminal nerve and its branches

- Radiation: tends to commence in the mandibular division and spreads to the maxillary division and (rarely) to the ophthalmic division
- Quality: excruciating, searing jabs of pain like a burning knife or electric shock
- Frequency: variable and no regular pattern
- Duration: 1 - 2 minutes (up to 15 minutes)
- Onset: spontaneous or trigger point stimulus
- Offset: spontaneous
- Precipitating factors: talking, chewing, touching trigger areas on face (e.g. washing, shaving, eating), cold weather or wind, turning onto pillow
- Aggravating factors: trigger points usually in the upper and lower lip, nasolabial fold or upper eyelid (Figure 52.6)



Figure 52.6 Trigeminal neuralgia: typical trigger points

- Relieving factors: nil
- Associated features:
  - rarely occurs at night
  - spontaneous remissions for months or years
- Signs: there are no signs, normal corneal reflex
- Causes:
  - unknown
  - local pressure on the nerve root entry zone by tortuous pulsatile dilated small vessels (probably up to 75%)
  - multiple sclerosis
  - neurosyphilis
  - tumours of the posterior fossa

Note: Precise diagnosis is essential.

Treatment

- Patient education, reassurance and empathic support is very important in these patients.

- Medical therapy:
    - carbamazepine (from onset of the attack to resolution)4 50 mg (elderly patient) or 100 mg (o) bd initially, gradually increase the dose to avoid drowsiness every 4 days to 200 mg bd (maintenance); testing serum levels is unnecessary; higher dosage may be necessary
    - alternative drugs if carbamazepine not tolerated or ineffective (but question the diagnosis if lack of response)
        - gabapentin 300 mg daily initially, then increase
        - phenytoin 300 - 500 mg daily
        - clonazepam
        - baclofen
- Surgery:
    - refer to a neurosurgeon if medication ineffective
    - possible procedures include:
        - decompression of the trigeminal nerve root (e.g. gel foam packing between the nerve and blood vessels)
        - thermocoagulation/radiofrequency neurolysis
        - surgical division of peripheral branches

Glossopharyngeal neuralgia [8, 10]

This is a rare condition of the ninth cranial nerve and branches of the vagus nerve with similar clinical features of severe lancinating pains, particularly felt in one ear, the base of the tongue or beneath the angle of the jaw.

- Sites: back of throat around tonsillar fossa and adjacent fauces deep in ear
- Radiation: ear canal, neck
- Triggers: swallowing, coughing, talking, yawning, laughing
- Treatment: as for trigeminal neuralgia

Migrainous neuralgia (cluster headache)

As described in Chapter 56 the pain is unilateral and centred around the eye with associated lacrimation and stuffiness of the nose.

Facial migraine (lower half headache) [8]

Migraine may rarely affect the face below the level of the eyes, causing pain in the area of the cheek and upper jaw. It may spread over the nostril and lower jaw. The pain is dull and throbbing, and nausea and vomiting are commonly present. The treatment is as for other varieties of migraine with simple analgesics or ergotamine for infrequent attacks.

Chronic paroxysmal hemicrania

In the rare condition of chronic or episodic paroxysmal hemicrania there is a unilateral facial pain that can resemble chronic cluster headache but the duration is briefer, about 15 minutes, and it may recur many times a day even for years. It responds dramatically to indomethacin.[11]

Herpes zoster and postherpetic neuralgia

Herpes zoster may present as hyperaesthesia or a burning sensation in any division of the fifth nerve, especially the ophthalmic division.

Atypical facial pain

This is mainly a diagnosis of exclusion whereby patients, usually middle-aged women, complain of diffuse pain in the cheek (unilateral or bilateral) without demonstrable organic disease. The pain does not usually conform to a specific nerve distribution (although in the maxillary area), varies in intensity and duration and is not lancinating as in trigeminal neuralgia. It is usually described as deep-seated and ‘boring’, severe, continuous and throbbing in nature. It is a very confusing and difficult problem to treat. These patients tend to show psychoneurotic tendencies but caution is needed in labelling them as functional.

Treatment

Trial of an antidepressant[4], e.g.

- dothiepin 25 - 150 mg nocte *or*
- amitriptyline 10 - 150 mg nocte

Temporal arteritis

This may produce mild or severe unilateral or bilateral headache. There may be ischaemic pain in the jaws when chewing. There may be marked scalp tenderness over the affected arteries. See p. 291 for management.

Erysipelas

Classical erysipelas is a superficial form of cellulitis involving the face. It usually presents with the sudden onset of butterfly erythema with a well defined edge. (Figure 52.7). It often starts around the nose and there may be underlying sinus or dental infection which should be investigated. There is an associated 'flu like' illness and fever. It is invariably caused by streptococcus pyogenes. Treament is by phenoxymethyl penicillin or di/flucloxacillin for 7 - 10 days.



Figure 52.7 Erysipelas: typical spreading distribution of the infection

When to refer

- Severe trigeminal neuralgia
- Unusual facial pain, especially with a suspicion of malignancy
- Positive neurological signs, such as impaired corneal reflex, impaired sensation in a trigeminal dermatome, slight facial weakness, hearing loss on the side of the neuralgia
- Possible need for surgical drainage of sinusitis - indications for surgery include failure of appropriate medical treatment, anatomical deformity, polyps, uncontrolled sinus pain[6]
- Dental root infection causing maxillary sinusitis
- Other dental disorders

Practice tips

- Malignancy must be excluded in the elderly with facial pain.
- Problems from the molar teeth, especially the third (wisdom), commonly present with periauricular pain without aural disease and pain in the posterior cheek.

- Facial pain never crosses the midline; bilateral pain means bilateral lesions.

REFERENCES

1. Beck ER, Francis JL, Souhami RL. *Tutorials in Differential Diagnosis*. Edinburgh: Longman Cheshire, 1987: 161 - 4.
2. Gerschman JA, Reade PC. Orofacial pain. Aust Fam Physician, 1984; 13: 14 - 24.
3. Selby G. Trigeminal neuralgia. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 531 - 3.
4. Mashford ML (Chair). *Therapeutic Guidelines: Analgesic* (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2002.
5. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 167 - 243.
6. Stevens M. The diagnosis and management of acute and chronic sinusitis. Mod Med Aust, 1991; April: 16 - 26.
7. Bridges-Webb C. Diagnosing sinus tenderness. Practice tip. Aust Fam Physician, 1981; 10: 742.
8. Tiller J (Chair). *Therapeutic Guidelines: Neurology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2002.
9. Corrigan B, Maitland G. *Practical Orthopaedic Medicine*. Sydney: Butterworths, 1986: 220.
10. Mendelsohn M, Lance J, Wheatley D. How to treat: Facial pain. Australian Doctor, 2003; 7 November: 31 - 6.
11. Burns R. Pitfalls in headache management. Aust Fam Physician, 1990; 19: 1825.

CHAPTER 53. FEVER AND CHILLS

Why! the fever itself is nature's instrument.

Thomas Sydenham (1624 - 89), *Medical Observations*

FEVER AND CHILLS

Although fever is a sign of disease and usually occurs in response to infection (mainly viral), its presence is recognised to play an important role in the individual's defence against infection. The infecting pathogen triggers hypothalmic receptors, causing the thermostatic mechanisms to be reset to maintain core temperature at a higher level. The elevation in temperature results from increased heat production (e.g. shivering) or decreased loss (e.g. peripheral vasoconstriction). The elevation in body temperature activates T-cell production, increases the effectiveness of interferons and limits the replication of some common viruses.[1]

KEY FACTS AND CHECK POINTS

- Fever plays an important physiological role in the defence against infection.
- Normal body temperature (measured orally mid morning) is 36 - 37.2°C (average 36.8°C).
- Fever can be defined as an early morning oral temperature > 37.2°C or a temperature > 37.8°C at other times of day.[2]
- Oral temperature is about 0.4°C lower than core body temperature.
- Axillary temperature is 0.5°C lower than oral temperature.
- Rectal, vaginal and ear drum temperatures are 0.5°C higher than oral and reflect core body temperature.
- There can be a normal diurnal variation of 0.5 - 1°C (lowest in early morning and highest in late afternoon).
- Fevers due to infections have an upper limit of 40.5 - 41.1°C (105 - 106°F).
- Hyperthermia (temperature above 41.1°C) and hyperpyrexia appear to have no upper limit.
- Infection remains the most important cause of acute fever.[3]
- Symptoms associated with fever include sweats, chills, rigors and headache.
- General causes of fever include infections, malignant disease, mechanical trauma (e.g. crush injury), vascular accidents (e.g. infarction, cerebral haemorrhage), immunogenic disorders (e.g. drug reactions, SLE), acute metabolic disorders (e.g. gout), and haemopoietic disorders (e.g. acute haemolytic anaemia).[3]
- Drugs can cause fever, presumably because of hypersensitivity.[3] Important examples are allopurinol, antihistamines, barbiturates, cephalosporins, cimetidine, methyldopa, penicillins, isoniazid, quinidine, phenolphthalein (including laxatives), phenytoin, procainamide, salicylates, sulphonamides.
- Drug fever should abate by 48 hours after discontinuation of the drug.[4]
- Infectious diseases at the extremes of age (very young and aged)3 often present with atypical symptoms and signs. Their condition may deteriorate rapidly.
- Overseas travellers or visitors may have special, even exotic infections and require special evaluation (refer Chapter 14).
- Immunologically compromised patients (e.g. AIDS patients) pose a special risk for infections, including opportunistic infections.
- A febrile illness is characteristic of the acute infection of HIV: at least 50% have an illness like glandular fever. Think of it!

CHILLS/RIGORS[2]

The abrupt onset of fever with a chill or rigor is a feature of some diseases. Examples include:

- pneumococcal pneumonia
- pyogenic infection with bacteraemia
- lymphoma
- pyelonephritis
- visceral abscesses (e.g. perinephric, lung)
- malaria
- biliary sepsis (Charcot's triad - jaundice, right hypochrondial pain, fever/rigors)

Features of a true chill are teeth chattering and bed shaking, which is quite different from the chilly sensations that occur in almost all fevers, particularly those in viral infections. The event lasts 10 - 20 minutes.

Other features:

- shaking cannot be stopped voluntarily
- absence of sweating
- cold extremities and pallor (peripheral vascular shutdown)
- dry mouth and pilo-erection
    - lasts 10 - 20 minutes

HYPERTHERMIA

Hyperthermia or hyperpyrexia is a temperature greater than 41.1°C (106°F). A more accurate definition is a state when the body's metabolic heat production or environmental heat load exceeds normal heat loss capacity. Hyperthermia may be observed particularly in the tropics, in malaria and heatstroke. It can occur with CNS tumours, infections or haemorrhages because of its effect on the hypothalamus.

Heatstroke (sunstroke, thermic fever)[5]

This is the sudden onset of hot, dry, flushed skin with a rapid pulse, temperature above 40°C, and confusion or altered conscious state in a person exposed to a very hot environment. The BP is usually not affected initially but circulatory collapse may precede death. It is a life-threatening emergency. Differential diagnoses include severe acute infection, toxic shock, food, chemical and drug poisoning. The elderly and debilitated are susceptible as are children left in cars.

Treatment

- Immediate effective cooling water applied to skin
- Icepacks at critical points (e.g. axillae, neck, head)
- Ice water bath if possible
- Aim to bring down temperature by 1°C every 10 minutes

Malignant hyperthermia

This is a rare hereditary disorder characterised by rapidly developing hyperpyrexia, muscular rigidity and acidosis in patients undergoing major surgery.

SWEATS

Sweating is a heat loss mechanism, and diffuse sweating that may soak clothing and bedclothing permits rapid release of heat by evaporation. In febrile patients the skin is usually hot and dry - sweating occurs in most when the temperature falls. It is characteristic of only some fevers (e.g. septic infections and rheumatic fever).

FACTITIOUS FEVER

Factitious fever is usually encountered in hospitalised patients attempting to malinger. The situation is usually suspected when:

- a series of high temperatures is recorded to form an atypical pattern of fluctuation
- there is excessively high temperature (41.1°C) and above
- a recorded high temperature is unaccompanied by warm skin, tachycardia and other signs of fever such as a flushed face and sweating
- there is an absence of diurnal variation

The patient may have surreptitiously dipped the thermometer in warm water, placed it in contact with a heat source or heated the bulb by friction with bedclothes or even mucous membranes of the mouth.

NEUROLEPTIC MALIGNANT SYNDROME

This is often confused with ‘malignant’ hyperthermia and heat stroke. The syndrome includes high temperature, muscle rigidity, autonomic dysfunction and altered consciousness. It is a rare and potentially lethal reaction in patients taking antipsychotic drugs, particularly occurring with haloperidol alone or with other drugs especially lithium carbonate.

MEASUREMENT OF TEMPERATURE

Temperature can be measured by several methods, including the mercury thermometer, the liquid crystal thermometer and the electronic probe thermometer. The mercury thermometer, however, is probably still the most widely used and effective temperaturemeasuring instrument.

Basic rules of usage

1. Before use, shake down to 35 - 36°C.
2. After use:
    - shake down and store in antiseptic
    - do not run under hot water
    - wipe rectal thermometer with alcohol and store separately
3. Recording time is 3 minutes orally and 1 - 2 minutes rectally.

Oral use

1. Place under the tongue at the junction of the base of the tongue and the floor of the mouth to one side of the frenulum - the ‘heat’ pocket.
2. Ensure that the mouth is kept shut.
3. Remove dentures.

Note: This is unsuitable for children 4 years and under, especially if irritable.

Rectal use

This is an appropriate route for babies and young children under the age of 4 years but should be used with care. The rule is ‘3 cm in for 3 minutes’ and some authorities claim that this method is the gold standard, especially in infants.

Method

1. Lubricate the stub with petroleum or KY jelly.
2. Insert for 3 cm (1 in) past anal verge.
3. Keep the thermometer between the flexed fingers with the hand resting on the buttocks (Figure 53.1).



Figure 53.1 Rectal temperature measurement in infants

Don't :

Axillary use

This is unreliable with poor sensitivity and generally should be avoided but may be practical in young children.[6, 7]

If used, place high in the axilla for 3 minutes. Fever is present if the temperature is above 37.2°C.

Groin use

This route is not ideal but is more reliable than the axilla. It closely approximates oral temperature. In infants, the thigh should be flexed against the abdomen.

Vaginal use

This is mainly used as an adjunct to the assessment of ovulation during the menstrual cycle. It should be placed deeply in the vagina for 5 minutes before leaving bed in the morning.

Infrared eardrum use

Otic thermography is now accepted standard practice. A systematic review in the Cochrane study describes the tympanic temperature as being inaccurate and unreliable.[8] However some Australian authorities believe that in general practice the benefits of convenience outweigh lack of accuracy.[9] The normal range is the same as for rectal temperature.

Skin use

Plastic strip thermometers placed on the forehead are very inaccurate and should not be used.

Accidental breakage in the mouth

If children bite off the end of a mercury thermometer there is no need for alarm, as the small amount of mercury is non-toxic and the piece of glass will usually pass in the stool.

THE CLINICAL APPROACH

The initial approach is to evaluate the severity of the problem and the nature of the illness. Some infections, particularly bacterial infections, are life-threatening and this requires urgent diagnosis and hospital admission.

According to Yung and Stanley[3] it is helpful to consider fever in three categories: less than 3 days duration; between 4 and 14 days duration and protracted fever (more than 14 days).

Fever of less than 3 days duration

This is very commonly encountered in family practice, often due to a self-limiting viral infection of the respiratory tract. It is important, however, to be vigilant for other infections, so evidence of an infectious disease, urinary tract infection, pneumonia or other infection should be sought. A routine urine examination, especially in females, is an important screening investigation. The majority of patients can be managed conservatively.

Fever present for 4 to 14 days

If fever persists beyond 4 - 5 days a less common infection should be suspected since most common viral infections will have resolved by about 4 days.[3] A checklist of causes is presented in Table 53.1. The careful history is mandatory as outlined for FUO. The basic examination and investigations are along similar lines.

Table 53.1 Common causes of fever of 4 - 14 days duration

- Influenza
- Sinusitis
- Epstein-Barr mononucleosis (EBM)
- Enteroviral infection
- Infective endocarditis
- Dental infections
- Hepatobiliary infections: hepatitis, cholecystitis, empyema of gall bladder
- Abscess
- Pelvic inflammatory disease
- Cytomegalovirus infection
- Lyme disease
- Travel-acquired infection: typhoid, dengue, hepatitis, malaria, amoebiasis
- Zoonosis: brucellosis, Q fever, leptospirosis, psittacosis
- Drug fever

Temperature chart

Charting the patterns of fever may be a diagnostic help because some febrile conditions follow a predictable temperature pattern.[10] Examples are presented in Figure 53.2.



Intermittent fever: quartan fever of malaria (a 4 day pattern with fever peaks every third day)



Undulant fever: due to lymphoma



Continuous pattern



Quotidian fever pattern and remittent fever
(e.g. empyema of gall bladder)

Figure 53.2 Examples of fever patterns from temperature charts

Intermittent fever

This is a fever in which the temperature rises for a few hours and then returns to normal (Figure 53.2a). Malaria is the classic example: in quartan fever, caused by *Plasmodium malariae*, the attacks occur every 72 hours (the term *quartan* means every 4th day by inclusive counting). This compares with tertian fever from *Plasmodium vivax* in which paroxysms of malaria occur every 48 hours. Other examples are cytomegalovirus, EBM and various pyogenic infections (e.g. ascending cholangitis).

Remittent fever

This is a fever in which the temperature returns towards normal for a variable period but is always elevated (Figure 53.2d). Common examples are collections of pus (e.g. pelvic abscess, wound infection, empyema and carcinoma). It is a common feature of empyema.

Undulant fever

Undulant fever is characterised by bouts of continuous or remittent fever for several days, followed by afebrile remissions lasting a variable number of days. It is commonly a feature of brucellosis infection but is also seen in the lymphomas, especially Hodgkin's lymphoma (Figure 53.2b). The latter is referred to as Pel-Ebstein fever with fevers lasting 3 to 10 days followed by afebrile periods of 3 to 10 days.

Continuous fever pattern

This is common with viral infections such as influenza (Figure 53.2c).

Quotidian fever

In this pattern the fever recurs daily (Figure 53.2d). Daily fever spikes in the morning are characteristic of *Pseudomonas* infection (e.g. pulmonary superinfection); afternoon spikes are indicative of cytomegalovirus infection; and evening spikes suggest localised collection of pus (e.g. empyema of the gall bladder).

Double quotidian fever (two fever spikes in a day) is caused by adult Still's syndrome, gonococcal endocarditis and visceral leishmaniasis.

If risk factors are present, especially if taking antibiotics, consider the investigations outlined in Table 86.1, p. 929.

FEVER IN CHILDREN

In children most authorities would consider a fever of 38.5°C and above to be significant and warrant close scrutiny.[11]

The fever is usually a response to a viral infection. Fever itself is not harmful until it reaches a level of 41.5°C.[1] Hyperthermia is uncommon in children. Temperatures above 41°C are usually due to CNS infection or the result of human error, for example

- shutting a child in a car on a hot day
- overwrapping a febrile child

Complications include dehydration (usually mild) and febrile convulsions, which occur in 5% of febrile children between 6 months and 5 years. Febrile convulsions are triggered by a rapid rise in temperature rather than its absolute level.

Note: Teething does *not* cause fever.

Approach to the febrile infant

It is important to decide whether the child looks well or seriously ill. Identification of the very ill child is presented in Chapter 86.

If the child is well and has no risk factors (e.g. unreliable caregiver, poor access to treatment, medical risk factors, taking antibiotics) treat expectantly. The only test required is urine microscopy and culture. Educate the caregiver about review if serious signs develop. Treat the fever as outlined in Table 86.1, p. 929.

Management

- Treatment of low-grade fevers should be discouraged.
- Treatment of high-grade fevers includes:
    - treatment of the causes of the fever (if appropriate)
    - adequate fluid intake/increased fluids
    - paracetamol (acetaminophen) is the preferred antipyretic since aspirin is potentially dangerous in young children (use if temperature > 38.5°C). The usual dose of 10 - 15 mg/kg every 4 - 6 hours may represent undertreatment. Use 20 mg/kg as a loading dose and then 15 mg/kg maintenance
    - evidence favours tepid sponging for first 30 minutes combined with paracetamol[12]
    - ibuprofen 5 - 10 mg/kg every 6 hours is a suitable antipyretic

Advice to parents

- Dress the child in light clothing (stripping off is unnecessary).
- Do not overheat with too many clothes, rugs or blankets.
- Give frequent small drinks of light fluids, especially water.
- Sponging with cool water and using fans is not effective.

Febrile convulsions

Refer to p. 933.

FEVER IN THE ELDERLY

The elderly tend to have a problem with impaired thermoregulation and so they may not develop a fever in response to suppurative infection compared with younger people. This can be misleading in the diagnostic process.

Important facts

- Any fever in the elderly is significant.
- Viral infection is a less common cause of fever in the elderly.
- Fever in the elderly is sepsis until proven otherwise (common sites are the lungs and urinary tract).

The elderly are more vulnerable to hyperthermia and hypothermia. Heatstroke classically occurs in epidemic form during a heatwave. The syndrome consists of hyperpyrexia, decreased sweating, delirium and coma. The core temperature is usually over 41°C.

‘ALARM BELL’ SIGNS

In many patients the existence of a life-threatening infective illness is obvious and prompt action is essential. In others the diagnosis is not clear-cut but there are certain warning signs:

- high fever
- repeated rigors
- drenching night sweats
- severe myalgia (? sepsis)
- severe pain anywhere (? sepsis)
- severe sore throat or dysphagia (? *Haemophilus influenzae* epiglottitis)
- altered mental state
- incessant vomiting
- unexplained rash
- jaundice
- marked pallor
- tachycardia
- tachypnoea

These symptoms and signs are obviously super ‘sensitive’. Patients with some of these features may have potentially life-threatening diseases, but this list would include many with viral infections.

FEVER OF UNDETERMINED ORIGIN

FUO, also referred to as pyrexia of unknown origin (PUO), has the following criteria:[13]

- illness for at least 3 weeks
- fevers > 38.3°C (100.9°F)
- undiagnosed after 1 week of intensive study

Most cases represent unusual manifestations of common diseases and not rare or exotic diseases. Examples are tuberculosis, bacterial endocarditis, hepatobiliary disease and carcinoma of the lung.[14]

Keep in mind that the longer the duration of fever, the less likely the diagnosis is to be infectious - fevers that last greater than 6 months are rarely infectious (only 6%). One study showed that 9% are factitious.[15]

Patients with FUO in definite need of further investigation are:

- babies < 3 months of age
- children with fever > 40°C
- adults > 50 years
- diabetics

- the immunocompromised
- travellers

A diagnostic approach

A knowledge of the more common causes of FUO is helpful in planning a diagnostic approach (refer to Table 53.2).

Table 53.2 Common causes of FUO

Common examples of each group selected
Infection (up to 40%) Bacteria • pyogenic abscess (e.g. liver, pelvic) • urinary infection • biliary infection (e.g. cholangitis) • chronic septicaemia • infective endocarditis • Lyme disease • tuberculosis • brucellosis • osteomyelitis • typhoid/paratyphoid fever Viral, rickettsial, *Chlamydia* • Epstein-Barr mononucleosis • cytomegalovirus • HIV virus infection (AIDS, ARC) • Q fever • psittacosis Parasitic • malaria • toxoplasmosis • amoebiasis
Malignancy (up to 30%) Reticuloendothelial • leukaemia • lymphomas Solid (localised) tumours • kidney • liver • pancreas • stomach • lung Disseminated
Immunogenic (up to 20%) Drugs Connective tissue diseases/vasculitides

• rheumatic fever • rheumatoid arthritis • SLE • polyarteritis nodosa/Wegener's granulomatosis • giant cell arteritis/polymyalgia Sarcoidosis Inflammatory bowel disease e.g. Crohn's
Factitious (1 - 5%)
Remain unknown (5 - 9%)

Source: After Kumar and Clark[16, 17]

History

The history should include consideration of past history, occupation, travel history, sexual history, IV drug use (leads to endocarditis and abscesses), animal contact, medication and other relevant factors. Symptoms such as pruritus, a skin rash and fever patterns may provide clues for the diagnosis. The average patient with a difficult FUO needs to have a careful history taken on at least three separate occasions.[16]

Physical examination

A common mistake is the tendency to examine the patient only once and not re-examine. The patient should be examined regularly (as for history taking) as physical signs can develop eventually. HIV infection must be excluded. Special attention should be paid to the following (Figure 53.3):

- skin - look for rashes, vesicles and nodules
- the eyes and ocular fundi
- temporal arteries
- sinuses
- teeth and oral cavity - ? dental abscess, other signs
- heart - murmurs, pericardial rubs
- lungs - abnormalities including consolidation, pleuritic rub
- abdomen - enlarged/tender liver, spleen or kidney
- rectal and pelvic examination (note genitalia)
- lymph nodes, especially cervical (supraclavicular)
- blood vessels, especially of the legs - ? thrombosis
- urine (analysis)


Factitious fever
Drugs
Lung
• abscess
• TB
• sarcoidosis
• carcinoma
Pyogenic abscess
• lung
• subphrenic
• liver
• perinephric
• empyema gall bladder
Malignancy
• liver
• kidney
• stomach
• pancreas
• lymphoma
• leukaemia
Dental abscess
Lymphadenopathy
• lymphoma
• leukaemia
• HIV infection
• sarcoidosis
• EBM
Heart
• Infective endocarditis
Splenomegaly
• myeloproliferative disorder
• bacterial endocarditis
Crohn's disease

Figure 53.3 Sites to consider in FUO (malignancy is indicated by a star)

Investigations

Basic investigations include:

- haemoglobin, red cell indices and blood film
- white cell count
- ESR/C-reactive protein
- chest X-ray and sinus films
- urine examination (analysis and culture)
- routine blood chemistry
- blood cultures

Further possible investigations:

- stool microscopy and culture
- culture of sputum (if any)
- specific tests for typhoid, EBM, Q fever, brucellosis, psittacosis, cytomegalovirus, toxoplasmosis, syphilis and others
- HIV screening

- tests for rheumatic fever
- tuberculin test
- tests for connective tissue disorders (e.g. DNA antibodies, C-reactive protein)
- upper GIT series with small bowel follow-through
- CT and ultrasound scanning for primary and secondary neoplasia:
    - gall bladder functioning
    - occult abscesses
- MRI - best for detecting lesions of the nervous system
- echocardiography - for suspected endocarditis
- isotope scanning for specific causes
- aspiration or needle biopsy
- laparoscopy for suspected pelvic infection
- tissue biopsies (e.g. lymph nodes, skin, liver, bone marrow) as indicated

FUO in children

Fever in children is usually a transient phenomenon and subsides within 4 - 5 days. At least 70% of all infections are viral. Occasionally a child will present with FUO which may be masked from antibiotic administration. Common causes of prolonged fever in children differ from those in adults. Most cases are not due to unusual or esoteric disorders,[18] the majority representing atypical manifestation of common diseases.

A summary of the common causes (with the most common ranked first) is as follows.[18]

Infectious causes (40%)

- Viral syndrome
- Urinary tract infection
- Pneumonia
- Pharyngitis
- Sinusitis
- Meningitis

Collagen - vascular disorders (15%)

- Rheumatic arthritis
- Aystemic lupus erythematosus
- Rheumatic fever
- Henoch-Schönlein syndrome

Neoplastic disorders (7%)

- Leukaemia
- Reticulum cell sarcoma
- Lymphoma

Inflammatory diseases of the bowel (4%)

SEPTICAEMIA

The diagnosis of septicaemia can be easily missed, especially in small children, the elderly and the immunocompromised, and in the absence of classic signs, which are:

- fever
- rash (suggestive of meningococcus)
- tachycardia
- tachypnoea
- cool extremities

Patients with septicaemia require urgent referral. Investigations should include 2 sets of blood cultures and other appropriate cultures e.g. urine, wound, sputum. Empirical initial treatment (after blood cultures) is di/flucloxacillin 2g IV + gentamicin 4 - 6mg/kg IV (statim).

Definitions

Bacteraemia

The transient presence of bacteria in the blood (usually asymptomatic) caused by local infection or trauma.

Septicaemia (sepsis)

The multiplication of bacteria or fungi in the blood, usually causing severe systemic symptoms such as fever, tachycardia and tachypnoea. Septicaemia has a very high mortality and demands urgent attention.

Severe sepsis

Sepsis associated with organ dysfunction, hypoperfusion or hypotension with 2 or more of: fever, tachycardia, tachypnoea, and elevated WCC.

Septic shock

Sepsis with critical tissue perfusion causing acute circulatory failure including hypotension and peripheral shut down - cool extremities, mottled skin, cyanosis.

Pyaemia

A serious manifestation of septicaemia whereby organisms and neutrophils undergo embolisation to many sites, causing abscesses, especially in the lungs, liver and brain.

Primary septicaemia

Septicaemia where the focus of infection is not apparent, while in *secondary septicaemia* a primary focus can be identified. Examples of secondary septicaemia in adults are:

- urinary tract (e.g. *Escherichia coli*)
- respiratory tract (e.g. *Streptococcus pneumoniae*)
- pelvic organs (e.g. *Neisseria gonorrhoea*)
- skin (e.g. *Staphylococcus aureus*)
- gall bladder (e.g. *E. coli*, *Streptococcus faecalis*)

Patients with septicaemia require urgent referral.

REFERENCES

1. Sewell J. Fever in childhood. Problems in clinical medicine. Aust Paediatr Rev, 1990: 2: 2.
2. Yung AP, McDonald MI, Spelman DW et al. *Infectious Diseases: a Clinical Approach*. Melbourne: Self-published, 2001: 13 - 16.
3. Yung A, Stanley P. Problems in infectious diseases. In: Kincaid Smith et al. *Problems in Clinical Medicine*. Sydney: MacLennan & Petty, 1990: 326 - 35.
4. Lipsky BA, Hirshmann JV. Drug fever. JAMA, 1981; 245: 851 - 4.
5. Tierney LM et al. *Current Medical Diagnosis and Treatment* (41st edn). New York: The McGraw-Hill Companies, 2002: 1595 - 6.
6. Keeley D. Taking infant's temperature: forget the axilla - the rectum is better. Br Med J, 1992; 304: 931 - 2.
7. Craig JV et al. Temperature measurement at the axilla compared with rectum in children and young people: systematic review. Br Med J, 2000; 320: 1174 - 8.
8. Ducesj J. Cochrane Library, 1996: 1 - 124: dare - 978207.
9. Nogrady B. Ear temperature readings get green light. Aust Doctor, 2002; 30 August: 3.
10. Beck ER, Francis JL, Souhami RL. *Tutorials in Differential Diagnosis* (3rd edn). Edinburgh: Churchill Livingstone, 1995: 209 - 13.
11. Fitzgerald D. Assessing fever in children. Medical Observer, 2003; 23 May: 36 - 7.
12. Bernath VF, Anderson JN, Silagy CA. Tepid sponging and paracetamol for reduction of body temperature in children: MJA 2002; 176: 30.
13. Roth AR, Basello GM. Approach to the adult patient with fever of unknown origin. Am Fam Physician, 2003; 68: 2223 - 8.

14. Tierney LM et al. *Current Medical Diagnosis and Treatment* (45th edn). New York: The McGraw-Hill Companies, 2006: 1286 - 8.
15. Gelfand JR. Fever of unknown origin. In: Braunwald E et al. *Harrison's Principles of Internal Medicine* (15th edn, Vol. 1). New York: McGraw-Hill, 2001: 805 - 6.
16. Braunwald E et al. *Harrison's Principles of Internal Medicine* (15th edn). New York: McGraw-Hill, 2001: 90 - 106.
17. Kumar PJ, Clark ML. *Clinical Medicine* (6th edn). London: Elsevier Saunders, 2003: 29 - 30.
18. Tunnessen WW Jr. *Signs and Symptoms in Paediatrics* (2nd edn). Philadelphia: JB Lippincott, 1988: 3 - 6.
19. Spicer J (Chair) *Therapeutic Guidelines: Antibiotic* (Version 13) Melbourne: Therapeutic Guidelines Ltd, 2006: 253 - 56.

CHAPTER 54. FAINTS, FITS AND FUNNY TURNS

Persons who have had frequent and severe attacks of swooning, without any manifest cause, die suddenly.

HIPPOCRATES (?460 - 377 BC), *Aphorisms*, 11, 41

FAINTS, FITS AND FUNNY TURNS

When patients present with the complaint of a ‘funny turn’ it is usually possible to determine that they have one of the more recognisable presenting problems, such as fainting, ‘blackouts’, lightheadedness, weakness, palpitations, vertigo or migraine. However, there are patients who do present with confusing problems that warrant the label of ‘funny turn’. The most common problem with funny turns is that of misdiagnosis, so a proper and adequate history taking is of great importance.

It is important to remember that seemingly ‘funny turns’ may be the subjective interpretation of cultural and linguistic communication barriers, especially in an emotional and frustrated patient.[1] Various causes of faints, fits and funny turns are presented in Table 54.1. A useful simple classification is to consider them as:

- syncope
- seizures
- sleep disorders - sleep apnoea/narcolepsy/cataplexy
- labyrinthine

Table 54.1 Faints, fits and funny turns: checklist of causes (excluding tonic - clonic seizure and stroke)

Psychogenic/communication problems
Conversion reactions (hysteria)
Culture/language conflicts
Fugue states
Hyperventilation
Breath holding attack
Malingering
Personality disorders
Phobia/anxiety states
Psychoses/severe depression
Other conditions
Transient ischaemic attacks
Complex partial seizure (temporal lobe epilepsy)
Tonic, clonic or atonic seizures
Primary absence seizure
Migraine variants or equivalents
Cardiovascular disorders
• arrhythmias
• postural hypotension
• aortic stenosis
Vertigo

Drug reaction
Alcohol and other substance abuse
Hypoglycaemia
Anaemia
Amnesic episodes
Metabolic/electrolyte disturbances
Vasovagal/syncope
Carotid sinus sensitivity
Cervical spondylosis
Sleep disorders
• sleep apnoea
• narcolepsy/cataplexy
Autonomic failure

KEY FACTS AND CHECKPOINTS

- The commonest cause of 'funny turns' presenting in general practice is lightheadedness, often related to psychogenic factors such as anxiety, panic and hyperventilation.[2] Patients usually call this 'dizziness'.
- Absence attacks occur with minor forms of epilepsy and with partial seizures such as complex partial seizures.
- The psychomotor attack of complex partial seizure presents as a diagnostic difficulty. The most commonly misdiagnosed seizure disorder is that of complex partial seizures or variants of generalised tonic - clonic seizures (tonic or clonic or atonic).
- The diagnosis of epilepsy is made on the history (or video electroencephalogram [EEG]), rather than on the standard EEG, although a sleep-deprived EEG is more effective.
- The triad - angina + dyspnoea + blackout or lightheadedness - indicates aortic stenosis.
- Severe cervical spondylosis can cause vertebrobasilar ischaemia by causing pressure on the vertebral arteries that pass through the intervertebral foramina, especially with head turning or looking up.

THE DIAGNOSTIC APPROACH

A summary of the diagnostic strategy model is presented in Table 54.2.

Table 54.2 Faints, fits and funny turns: diagnostic strategy model

Q. Probability diagnosis
A. Anxiety related/hyperventilation
Vasovagal syncope
Postural hypotension
Breath-holding attacks (children)
Q. Serious disorders not to be missed
A. Cardiovascular
• arrhythmias
• aortic stenosis
Cerebrovascular

- TIAs

Neoplasia

- space-occupying lesions

Severe infections

- infective endocarditis

Hypoglycaemia

Q. Pitfalls (often missed)

A. Atypical migraine

Cardiac arrhythmias

Simple partial seizures

Complex partial seizures

Atypical tonic - clonic seizures

Drugs/alcohol/marijuana

Electrolyte disturbances (e.g. hypokalaemia)

Sleep disorders

Transient global amnesia

Rarities

- Atrial myxoma

Q. Seven masquerades checklist

A. Depression

Diabetes   hypoglycaemia

Drugs

Anaemia

Thyroid disorder -

Spinal dysfunction   cervical spondylosis

UTI -

Q. Is this patient trying to tell me something?

A. Highly likely. Psychogenic and ‘communication’ disorders quite significant.

History

The clinical history is of paramount importance in unravelling the problem. A reliable eye witness account of the ‘turn’ is invaluable, as is the setting or circumstances in which the ‘episode’ occurred.

It is essential at first to determine exactly what the patient means by ‘funny turn’. In the process of questioning it is appropriate to evaluate the mental state and personal and social factors of the patient. It may be appropriate to confront the patient about feelings of depression, anxiety or detachment from reality.

It is important to break up the history into three components. The first is the lead-up to the episode, the second is an adequate description of what took place during the episode, and thirdly, the events that took place after the episode.

Apart from the events, note the patient's feelings, symptoms, circumstances and provocative factors. Search for possible secondary gain.

Onset

A sudden onset may be due to cardiovascular causes, especially arrhythmias, which may include the more common supraventricular tachycardias in addition to the less common but more dramatic arrhythmias that may cause unconsciousness. Other causes of a sudden onset include the various epilepsies, vasovagal attacks and TIAs.

Precipitating factors[2]

Enquire about precipitating factors such as emotion, stress, pain, heat, fright, exertion, suddenly standing up, coughing, head movement or hypersomnolence:

- emotion and stress suggest hyperventilation
- fright, pain → vasovagal attack
- standing up → postural hypotension
- exertion → aortic stenosis
- head movement → cervical spondylosis with vertebrobasilar insufficiency
- hypersomnolence → narcolepsy

Associated symptoms[2]

Certain associated symptoms give an indication of the underlying disorder:

- breathing problems and hyperventilation suggest an anxiety state
- tingling in extremities or tightening of the hand → anxiety/hyperventilation
- visual problems → migraine, TIA
- fear or panic → anxiety or complex partial seizure
- hallucinations (taste/smell/visual) → complex partial seizure
- speech problems → TIA/anxiety
- sweating, hunger feelings → hypoglycaemia
- related to food → migraine
- first thing in morning → consider ‘ehangover’f

Drug history

This requires careful analysis and includes alcohol intake and illicit drugs such as marijuana, cocaine and amphetamines. Prescribed drugs that can cause lightheadedness or unconsciousness are listed in Table 54.3.

Table 54.3 Typical drugs that may cause lightheadedness or blackouts

Alcohol
Peripheral vasodilators
• ACE/AIIRA inhibitors
• glyceryl trinitrate
• hydralazine
• prazosin
Antiepileptics
Antihypertensives
Barbiturates
Benzodiazepines
Phenothiazines
Phenoxybenzamine
Tricyclic antidepressants

SSRI antidepressants
OTC anticholinergic compounds

Sudden cessation of certain drugs such as phenothiazines can also be responsible for ‘funny turns’.

Past history

The past history may give an indication of the cause of the ‘turn’. Such conditions include hypertension, migraine, epilepsy, rheumatic heart disease, atherosclerosis (e.g. angina, vascular claudication), alcohol or other substance abuse and psychiatric disorders.

Diary of events

If the diagnosis is elusive it may help to get the patient to keep a diary of circumstances in which events take place, keeping in mind the importance of the time period prior, during and post episode.

The examination

Important focal points of the physical examination include:

- evaluation of the mental state, especially for anxiety
- looking for evidence of anaemia, alcohol abuse and infection
- cerebrovascular examination: carotid arteries, ocular fundi, bruits
- cardiovascular examination: pulses, BP, heart (the BP should be taken lying, sitting and standing)
- the cervical spine

Various manoeuvres

Subject the patient to a number of manoeuvres to try to induce various sensations in order to identify the one that affects them. These should include sudden assumption of the erect posture from a squat, spinning the patient and then a sudden stop, head positioning with either ear down (Figure 46.3, p. 507), Valsalva manoeuvre, and hyperventilation for 60 seconds. Children can spin a showbag ‘windmill’ while hyperventilating (blowing). Ask ‘Which one mimics your complaint?’

Investigations

Depending on the clinical findings, investigations can be selected from the following tests:

- full blood count: ? anaemia ? polycythaemia
- blood sugar: ? diabetes ? hypoglycaemia
- urea and electrolytes
- ECG: ? ischaemia ? arrhythmia
- 24-hour ambulatory cardiac (Holter) monitor: ? arrhythmia
- radiology/imaging
  - cervical X-ray
  - chest X-ray
  - carotid duplex Doppler scan: ? carotid artery stenosis
  - CT scan
  - MRI scan
- EEG or video EEG; include those recorded with sleep deprivation, hyperventilation or photic stimulation
- positron emission tomography (PET) or single photon emission computerised tomography (SPECT) may show localised brain dysfunction when others are negative

CHILDREN

The various forms of seizures are also encountered in childhood when most patients with epilepsy are diagnosed. There is a concern, supported by evidence, that children are being misdiagnosed. It is appropriate to take a visual recording of the attack.[3]

Epilepsy syndromes

The following are special age-related epilepsy syndromes seen in children.[4, 5]

Febrile seizures

Tonic - clonic seizures occur in 2 - 5% of children usually aged 6 months to 5 years who have a high fever generally caused by a viral infection. The longterm prognosis is good.

Infantile spasms (hypsarrhythmia)

Also known as West syndrome, these are generalised tonic seizures with sudden flexion of the arms, forward flexion of the trunk and extension of the legs, lasting only a few seconds, with usual age onset between 3 and 7 months. They are usually restricted to the first 3 years of life and are replaced by other forms of attacks. Prognosis for cognitive development is also unfavourable. The most effective therapy is corticotrophin (ACTH) IM injection. Otherwise, oral prednisolone, vigabatrin, benzodiazepines or sodium valproate can be used.

Lennox-Gastaut syndrome (myoclonic epilepsy of infancy)

This uncommon syndrome refers to a triad of severe difficult-to-control seizures (usually tonic with drop attack), mental retardation and characteristic EEG. The seizures usually begin between ages 1 and 6 years with a peak onset at 3 to 5 years. Prognosis is also poor. Sodium valproate is the therapy of choice.

Benign rolandic epilepsy

This disorder usually begins in children aged 2 - 13 years with a peak age of 5 to 8 years. There is a strong family history of epilepsy. The feature is a simple partial motor or somatosensory seizure involving the face and mouth during sleep, producing a typical ‘glugging’ sound. The child usually wakes from sleep, goes to the parents and is unable to speak and has hemifacial contortions. It may progress to a tonic - clonic seizure. There is a characteristic EEG pattern. The prognosis is excellent as remission usually occurs around puberty. Carbamazepine is the therapy of choice.

Juvenile absence syndrome

These children present with frequent absence seizures, often over a hundred daily. Peak age of onset is 6 to 7 years. The absence seizures can be very subtle. Signs include alteration of awareness (usually in the classroom), sudden onset, facial and other automatisms. First-line treatment is sodium valproate.

Juvenile myoclonic epilepsy (myoclonic epilepsy of Janz)

This is a triad of seizures: myoclonic jerks, tonic - clonic seizures and absences. Onset is around puberty but may occur earlier. The myoclonic jerks and tonic - clonic seizures usually occur in the early morning after waking. Mental development is usually normal but the disorder is usually lifelong and is well controlled with sodium valproate.

Medial temporal lobe epilepsy

This syndrome of complex partial seizures, which usually last 1 to 3 minutes, is seen in childhood. Transient post-ictal confusion and speech dysfunction is common. Those with medically intractable seizures respond well to surgery.

Non-epileptic events resembling epileptic seizures [4]

Many normal and abnormal behaviours seen in children resemble seizures but are unrelated to epilepsy. A careful history is very important. The following are examples:

- *Postures of spasticity and movement disorders.* These occur in neurologically handicapped children such as those with cerebral palsy.

- *Syncope*. The child may describe a ‘sinking feeling’, or ‘everything getting louder’ prior to the loss of consciousness.
- *Breath-holding*. This often occurs after a crying spell and clonic movements may be seen at the end of the event.
- *Masturbation*. This behaviour leads to a tonic - like posture of the legs and preoccupation, especially in young girls.[6]
- *Munchausen-by-proxy*.[7] This syndrome of fictitious epilepsy described by a parent is becoming more recognised.
- *Psychogenic seizures (pseudo seizures)*. A diagnostic dilemma exists when these coexist with genuine seizures. These should be suspected when they occur in particular circumstances and the description of the ‘seizure’ is bizarre.
- *Shuddering*. Shuddering or shivering spells can resemble myoclonic jerks.
- *Night terrors*.[8] These episodes, which usually affect 2 - 4 year olds and 6 - 9 year olds, generally develop within 2 hours of sleep onset and last 1 to 2 minutes (sometimes longer). They are alarming and the child usually cannot be reassured or settled. A 6-week trial of phenytoin or imipramine can be used for severe problems.
- *Tics*. Motor tics can be quite complex but are usually brief involuntary movements involving the face and upper limbs.

BLACKOUTS

The important causes of blackout include the various syncopes that are listed in Table 75.4 (p. 820) and the various forms of epilepsy. The classic tonic - clonic seizure is described in Chapter 75 (p. 817) while descriptions of other seizures producing blackouts or funny turns now follow.

Important causes of convulsions (tonic - clonic seizures) are listed in Table 54.4. Refer to Chapter 125 (p. 1289).

Table 54.4 Important causes of convulsive seizures

• Epilepsy
- first presentation
- known patient with recurrence
• Cerebral hypoxia
• Hypoglycaemia
• Poor cerebral perfusion, e.g.
- oedema of eclampsia
• Neurotrauma
• Cerebrovascular accident
• CNS infections, e.g.
- meningitis
- encephalitis
- septicaemia
- septic emboli
- cerebral abscess
• Toxins
• Alcohol excess
• Hyperthermia
• Metabolic disorders
• Drugs, e.g.
- antidepressants
- theophylline
- amphetamine

- cocaine
- local anaesthetics

• Anaphylaxis

• Expanding brain lesion, e.g.

- neoplasm
- haematoma

COMPLEX PARTIAL SEIZURES

In complex partial seizures (known also as temporal lobe epilepsy) the symptomatology varies considerably from patient to patient and is often a diagnostic problem. It is the commonest type of focal epilepsy and the attacks vary in time from momentary to several minutes (usually 1 to 3 minutes).

Possible manifestations[2]

- Commonest: slight disturbance of perception and consciousness
- Hallucinations:
  - visual
  - taste
  - smell
  - sounds
- Absence attacks or vertigo
- Illusions - objects/people shrink or expand
- Affective feelings - fear, anxiety, anger
- Dyscognitive effects:
  - *déjà vu* (familiarity)
  - *jamais vu* (unreality)
  - waves emanating from epigastrium
- Objective signs:
  - lip-smacking
  - swallowing/chewing/sucking
  - unresponsive to commands or questions
  - pacing around a room

Unreal or detached feelings are common in complex partial seizures. There can be permanent short-term memory loss. The sensation of strange smells or tastes is more common than auditory or visual hallucinations.[1] They can progress to tonic - clonic seizures.

Diagnosis

- EEG
  - diagnostic in 50 - 60% of cases
  - a repeat EEG will increase rate to 60 - 80%
- EEG/video telemetry helpful with frequent attacks
- CT or MRI scan - to exclude tumour when diagnosis confirmed

Medication

- carbamazepine (first choice)[9] *or*
- phenytoin (second choice) *or* (others)

- vigabatrin, sodium valproate, phenobarbitone, tiagabine

TONIC - CLONIC SEIZURES

Variants of tonic - clonic seizures are more common than realised. Some patients may simply stiffen or drop to the ground while others may have one or two jerks or shakes only:

- stiffen and fall ° tonic
- floppy and fall ° atonic
- shaking only ° clonic

Refer to Chapter 75 (p. 821).

SIMPLE PARTIAL SEIZURES

In simple partial seizures (Jacksonian epilepsy) there is no loss of consciousness. These include focal seizures, which may proceed to a generalised tonic - clonic seizure or to motor seizures.

Jacksonian (motor seizure)

Typically, jerking movements begin at the angle of the mouth or in the thumb and index finger and 'march' to involve the rest of the body (e.g. thumb → hand → limb → face ± leg on one side and then on to the contralateral side). A tonic - clonic or complex partial seizure may follow.

Medication

- carbamazepine (first choice)[9] *or*
- phenytoin (second choice) *or* (others)
- sodium valproate, vigabatrin, gabapentin

ABSENCE SEIZURE (PREVIOUSLY CALLED PETIT MAL)

This type of generalised epilepsy typically affects children from 4 years up to puberty:[2]

- child ceases activity and stares suddenly
- child is motionless (may blink or nod)
- no warning
- sometimes clonic (jerky) movement of eyelids, face, fingers
- may be lip-smacking or chewing (called complex absence)
- only lasts few seconds - usually 5 - 10 seconds
- child then carries on as though nothing happened
- usually several per day (not just one or two)
- may lead to generalised seizures in adulthood

Diagnosis

Best evoked in the consulting room by hyperventilation and 'windmill'.

- EEG
  - classic 3 Hz wave and spike
  - may be normal
  - always include hyperventilation
  - easier with sleep deprivation

Medication

- ethosuximide (first choice)[10] *or*
- sodium valproate (second choice) *or* (others)
- clonazepam, gabapentin

Note: Beware of hepatoxicity with sodium valproate, especially in those under 2 years.

NARCOLEPSY

Narcolepsy is characterised by brief spells of irresistible sleep during daytime hours, usually at times when the average person simply feels sleepy. Although patients are usually aware of their disorder some may have no insight into the problem and present with the complaint of unusual turns. Narcolepsy can present as 'attacks' in which the patient may crumple and fall without losing consciousness. It can be part of a tetrad syndrome (narcolepsy, cataplexy, hypnogogic hallucinations, sleep paralysis).

Other features:

- onset in teens or twenties
- can have several attacks per day

Refer to Chapter 71 (p. 786).

Diagnosis

- A clinical diagnosis

If doubtful:

- EEG monitoring
- sleep laboratory studies (sleep latency test) - rapid eye movement is a hallmark

Medication

- amphetamines (in slowly increasing doses) *or* methylphenidate (Ritalin)
- tricyclic antidepressants (e.g. clomipramine), for associated cataplexy

AMNESIC EPISODES

Amnesic episodes in which people cannot recall events or their own identity can be psychogenic (commonly), such as fugue states, conversion disorder, severe depression and factitious states. They may be related to an organic problem such as epilepsy, sleep apnoea, cerebrovascular disorder, post-trauma, Wernicke - Korsakoffsyndromes and drugs, e.g. alcohol, cannabis and anaesthetic agents.

Transient global amnesia[11, 12]

This is a benign, self-limited profound amnesic episode of unknown aetiology tending to occur in middle-aged or elderly people. Proposed causation includes transient ischaemia or dysfunction of the temporal lobes, similar to temporal lobe epilepsy or a migraine variant.

Features

- Typically last 4 - 8 hours (up to 24 hours)
- Identity and conscious state preserved
- Agitated, perplexed, anxious patient

- State of bewilderment, e.g. ‘where am I?’
- Frequent repetition of questions
- Able to perform complex motor skills (e.g. driving)
- Usually single episode (20% recurrence)
- Complete resolution: good prognosis
- No abnormal neurological signs
- All investigations unhelpful

Four diagnostic criteria (after Caplan)[11]

- Witnessed onset of attack
- Dysfunction during attack limited to amnesia and repetitive queries
- No other neurological features
- Memory loss should be transient lasting no longer than 24 hours

Management

- Reassuring explanation/education
- Special investigations, including angiography not needed if conform to above criteria
- No active treatment usually needed

CEREBROVASCULAR DISORDERS

Cerebrovascular disease is one of the major causes of mortality and morbidity in developed countries and can cause recurrent attacks of ischaemia in the carotid and vertebrobasilar systems (particularly vertebrobasilar insufficiency), which may present as ‘funny turns’. In particular, brain-stem ischaemia causes ‘funny turns’ such as impaired consciousness, including transient global amnesia, drop attacks and the ‘locked-in’ syndrome.

PSYCHOGENIC OR COMMUNICATION DISORDERS

Psychogenic causes have to be considered. ‘Hysterical fugue’ is one such manifestation. The problem can be a communication disorder, such as an emotional person trying to communicate a problem in a language foreign to them.

Patients with psychiatric disorders such as schizophrenia or depression may experience feelings of depersonalisation or unreality, which can be interpreted as a ‘turn’ or even temporal lobe epilepsy.

Patients who complain of vague and bizarre symptoms, such as ‘queer feelings in the head’, ‘swimming sensation’, ‘unreal feelings’ and ‘walking on air’ are likely to have an anxiety state.

Severe anxiety or panic attacks typically cause lightheadedness that presents as a ‘funny turn’. Other somatic symptoms include palpitations, sweating, inability to swallow, headache, breathlessness and manifestations of hyperventilation.

PITFALLS IN MANAGEMENT[2]

- The main pitfall associated with seizure disorders and epilepsy is misdiagnosis (not all seizures are generalised tonic - clonic in nature).
- Failing to place appropriate emphasis on the history in making the diagnosis.
- Misdiagnosing syncope with some involuntary movements for epilepsy.
- Overlooking cardiac arrhythmias as a cause of funny turns, including recurrent dizziness.
- Failing to consider the possibility of aortic stenosis with syncopal attacks.

- Misdiagnosing vertigo and syncope for TIA.
- Mistaking visual or sensory migraine equivalents in young adults for TIA.
- Overlooking drugs (including self-administered drugs) as a cause of lightheadedness.

Practice tips

- A detailed clinical analysis is more important in the first instance than laboratory tests. The key to accurate diagnosis is a very careful history, taking the patient second by second through the attack and events preceding the turn.
- Talk to as many eyewitnesses as possible in unravelling the cause.
- For ‘undiagnosed turns’ ask the patient to keep a diary with an accurate record of the attack, including preceding events.
- Remember that migraine is a great mimic and can cause confusion in diagnosis.
- Remember that the EEG can be normal in the confirmed epileptic.
- The more bizarre the description of a ‘funny turn’, the more likely a functional problem is the cause.
- Transient hypoglycaemia can mimic a TIA.

WHEN TO REFER

- Transient ischaemic attacks, especially if the diagnosis is in doubt
- Clinical suspicion of or proven cardiac arrhythmias
- Evidence of aortic stenosis
- Seizures
- General uncertainty of the diagnosis

REFERENCES

1. Kincaid-Smith P, Larkins R, Whelan G. Problems in Clinical Medicine. Sydney: MacLennan & Petty, 1990: 159 - 64.
2. Sandler G, Fry J. *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986: 411 - 30.
3. Hindley D, Ali A, Robson C. Diagnoses made in a secondary care ‘fits, faints and funny turns’ clinic. Arch Dis Child, 2006; 91(3): 214 - 8.
4. Stewart I, Bye A. The diagnosis of childhood fits and funny turns. Mod Med Aust, 1994; 37(8): 65 - 72.
5. Tiller JWG (Chair). *Therapeutic Guidelines: Neurology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2002: 91 - 102.
6. Fleisher DR, Morrison A. Masturbation mimicking abdominal pain or seizures in young girls. J Paediatr, 1990; 116: 810 - 14.
7. Meadow R. Fictitious epilepsy. Lancet, 1984; 2: 25 - 8.
8. Rothner AD. Not everything that shakes is epilepsy: the differential diagnosis of paroxysmal nonepileptiform disorders. Cleve Clin J Med, 1989; 56 Suppl part 2: 206S - 213S.
9. Scott AK. Management of epilepsy. In: Central Nervous System. London: British Medical Association, 1995: 1 - 2.
10. Levine M et al. *Drugs of Choice: A Formulary for General Practice*. Ottawa: Canadian Medical Association, 1995: 98 - 9.
11. Caplan LR. Transient global amnesia: criteria and classification. Neurology, 1986; 36: 441.
12. Horne M. Neurology quiz. Aust Fam Physician, 1994; 23: 935.

CHAPTER 55. HAEMATEMESIS AND MELAENA

> The ‘once smelt never forgotten’ sickly smell of melaena can be diagnosed from a distance of 20 metres without trying.
>
> EMERGENCY ROOM SUPERVISOR, BRISBANE 1985

HAEMATEMESIS AND MELAENA

Acute severe upper GI haemorrhage is an important medical emergency. The dramatic symptom of haematemesis follows bleeding from the oesophagus, stomach and duodenum. Over half the patients are over 60 years of age.[1]

Haematemesis is the vomiting of blood appearing as fresh blood or ‘coffee ground’. Melaena is the passage of black tarry stools, with 50 mL or more of blood required to produce melaena stool. Melaena occurs in most patients with upper GI haemorrhage and haematemesis occurs in over 50%.[1]

Although the incidence is declining, the morfality rate of upper 61 Laemorrtage remains high at approximately6 - 8%.[2]

KEY FACTS AND CHECKPOINTS

- Chronic peptic ulceration accounts for most cases of upper GI haemorrhage.
- Haematemesis is almost always associated with some degree of blood in the stools, although melaena may not necessarily accompany it, especially if bleeding occurs from the oesophagus.
- Black stool caused by oral iron therapy or bismuth-containing antacid tablets can cause confusion.
- Always check for a history of drug intake, especially aspirin and NSAIDs.
- Corticosteroids in conventional therapeutic doses are thought to have no influence on GI haemorrhage.
- The volume of the bleeding is best assessed by its haemodynamic effects rather than relying on the patient's estimation, which tends to be excessive.
- Melaena is generally less life-threatening than haematemesis.
- Resuscitation of the patient is the first task.
- A sudden loss of 20% or more circulatory blood volume usually produces signs of shock such as tachycardia, hypotension, faintness and sweating. Younger patients can compensate better and tolerate a larger loss prior to the development of shock.[1] A useful guide is that shock in a previously well 70 kg man indicates an acute blood loss of at least 1000 - 1500 mL.

CAUSES OF UPPER GI BLEEDING

The major cause of bleeding is chronic peptic ulceration of the duodenum and stomach, which accounts for approximately half of all cases.[3] The other major cause is acute gastric ulcers and erosions, which account for at least 20% of cases. Aspirin and NSAIDs are responsible for many of these bleeds. Causes are summarised in Table 55.1 and illustrated in Figure 55.1.

Table 55.1 Causes of upper gastrointestinal bleeding[4, 5]

Common causes
1 Duodenal ulcer
2 Gastric erosion
3 Gastric ulcer
4 Reflux oesophagitis
5 Oesophageal varices

6 Mallory-Weiss emetogenic syndrome
Others
• Gastric or oesophageal carcinoma
• Stomal ulcer
• Blood dyscrasias
• Anticoagulant therapy
• Vascular malformations/angiodysplasia
• Hereditary haemorrhagic telangiectasia (Rendu-Osler-Weber syndrome)


oesophagitis
oesophageal varices
Mallory-Weiss syndrome
gastric ulcer
duodenal ulcer
acute gastric erosions and ulcers
gastric carcinoma

Figure 55.1 Important causes of haematemesis and melaena

Mallory-Weiss syndrome

In this condition a tear occurs at the lower end of the oesophageal mucosa (at the oesophago-gastric junction) because of an episode of severe or protracted vomiting or coughing. Blood appears in the vomitus after a bout of heavy vomiting or dry retching. It is usually seen in alcoholic patients. It is usually a self-limiting lesion. A definite diagnosis can only be made by oesophagoscopy.

Gastro-oesophageal varices

Such varices are caused by portal hypertension, which in turn is usually due to cirrhosis of the liver. There is a raised incidence of peptic ulcer in those with liver cirrhosis, especially in biliary and alcohol-induced cirrhosis, so this should be kept in mind as a possible source of bleeding. Mortality is about 30%, despite advances.[5]

Management includes injection sclerotherapy, and then intravenous octreotide if it fails. Passage of a Sengstaken-Blakemore or Minnesota tube into the oesophagus and stomach to provide tamponade and the radiological procedure of using a transjugular intrahepatic portosystemic stent are possible options.

THE CLINICAL APPROACH

History

It is important to establish the nature of the vomitus and the possibility of bleeding arising from the mouth, nose or pharynx. A coffee grounds vomitus indicates that the blood has been in contact with gastric acid. Oesophageal bleeding tends to lead to vomiting of fresh blood. Questions to help pinpoint the possible aetiology should be asked.

Key questions

- What drugs have you been taking? (Table 55.2)
- Have you been taking aspirin or tablets for arthritis or back pain?
- How much have you vomited?
- What did the vomit look like?
- Do you notice black dots like coffee grounds or any blood clots?
- Have you had any indigestion, heartburn or stomach pains recently?
- Have you opened your bowels and if so what was the colour?
- Have you noticed whether your bowel motions were black or unusual in any way?
- How much alcohol do you drink?
- Have you had any previous operations on your stomach for a peptic ulcer?
- Were you vomiting normal vomit before the blood appeared?

• Drugs Assosiation with gastrointestinal haemorrhage
• Aspirin
• NSAIDS/CO X-2 inhibit
• Prednisolone
• Warfarin
• Clopidogerl
• Other antiplatelet dn
• Iteparin
• SSRI antidepression

Physical examination

The patient's general state, particularly the circulation, should be assessed immediately on presentation. It is critical to assess the palient's haenidynamic status will vital sign of hrart rate, blood pressur and postual change.[2] A careful abdominal examination should be performed including a digital rectal examination. As a rule abdominal findings are not remarkable except when a mass, hepatomegaly or splenomegaly is found. Other evidence of liver disease should be sought.

Investigations

Investigations to determine the source of the bleeding should be carried out in a specialist unit. Upper gastrointestinal endoscopy is the single most useful test and will detect the cause of the bleeding in at least 80% of cases.[3]

The haemoglobin level will not be an appropriate guide to blood loss or the need for transfusion during the early stages, because haemodilution occurs gradually over the 24 hours following a severe bleed. However, a level below 90 g/L during this period is usually regarded as an indication for transfusion.

Management

The immediate objectives are:

1. restore an effective blood volume (if necessary)
2. establish a diagnosis to allow definitive treatment

All patients with a significant bleed should be admitted to hospital and referred to a specialist unit. Urgent resuscitation is required where there has been a large bleed and there are clinical signs of shock. Such patients require insertion of intravenous lines and rapid infusion of isotonic saline followed by a plasma expander (e.g. Haemaccel) followed by transfusion with blood commenced as soon as possible.

In many patients bleeding is insufficient to decompensate the circulatory system and they settle spontaneously. Approximately 85% of patients stop bleeding within 48 hours.[3]

In some instances intervention via endoscopy to achieve hoemostasis of a bleeding point with a heater probe or injection with odrenaline or both or band varices will be employed. Occasionally surgery will be necessary to arrest bleeding but should be avoided if possible in patients with acute gastric erosion.

REFERENCES

1. Tierney LM et al. Current Medical Diagnoses and Treatment (41st edn). Stamford: McGraw-Hill and Lange, 2002: 502 - 22.
2. Worthley DL, Fraser R J. Management of acute bleeding in the upper gastrointestinal trait. Australian Presculer, 2005; 28:62 - 6.
3. Kumar PJ, Clark ML. Clinical Medicine (6th edn). London: Elsevier Saunders 2003: 291 - 3.
4. Beck ER, Francis JL, Souhami RL. Tutorials in Differential Diagnosis (3rd edn). Edinburgh: Churchill Livingstone, 1995: 75 - 9.
5. Talley NJ, Martin CJ. Clinical Gastroenterology. Sydney: MacLennan & Petty, 1996: 150 - 3.

CHAPTER 56. HEADACHE

When the head aches, all the body is out of tune.

CERVANTES 1547 - 1616

HEADACHE

Headache, one of the cardinal symptoms known to human beings, is a very common complaint in general practice. When a patient presents with 'headache' we need to have a sound diagnostic and management strategy as the problem can be confusing. The key to analysing the symptom of headache is to know and understand the cause, for 'one only sees what they know'.

The patient's manner of presentation can confuse us because many tend to influence us with preconceived ideas that they will verbalise - 'I think I need my blood pressure checked' or 'My eyes need testing' - or they may not mention their anxiety about a cerebral tumour or an impending stroke.

Hypertension is such a rare cause of headache that one is tempted to stress the adage 'hypertension does not cause headache', but we do encounter the occasional patient whose headache appears to be caused by hypertension and it is mandatory to measure the blood pressure of patients presenting with headache. Patients expect this routine and reassurance is difficult without the appropriate physical examination. Where headaches and hypertension coexist, assume that the headaches are not due to hypertension.

The diagnosis of serious causes of headache depends on a careful history, a high index of suspicion of the 'different' presentations and the judicious use of CT scanning.

KEY FACTS AND CHECKPOINTS

- 85% of the population will have experienced headache within 1 year and 38% of adults will have had a headache within 2 weeks.[1]
- 40% of children will have experienced one or more headaches by the age of 7 and 75% by the age of 15.[2]
- Migraine affects at least 10% of the adult population and one-quarter of these patients require medical attention for their attacks at some stage.[3] It is underrecognised and poorly managed in the community.[4]
- 5% of children suffer from migraine by the age of 11 years.[3]
- 70% of sufferers have a positive family history of migraine.
- Many headaches previously considered to be tension are secondary to disorders of the neck, eyes, teeth, temporomandibular joints or other structures.[3]
- Drug-induced headaches are common and must be considered in the history.
- In children the triad of symptoms - dizziness, headache and vomiting - indicates medulloblastoma of the posterior fossa until proved otherwise.
- A typical triad of symptoms in an adult with a cerebral tumour (advanced) is headache, vomiting and convulsions.
- Eye strain is not a common cause.
- Bronchial carcinoma is the commonest cause of intracerebral malignancy.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 56.1.

Table 56.1 Headache: diagnostic strategy model

Q. Probability diagnosis
A. Acute: respiratory infection

Chronic: • tension-type headache

- combination headache
- migraine
- transformed migraine

Q. Serious disorders not to be missed

A. Cardiovascular

- subarachnoid haemorrhage
- intracranial haemorrhage
- carotid or vertebral artery dissection
- temporal arteritis
- cerebral venous thrombosis

Neoplasia

- cerebral tumour
- pituitary tumour

Severe infections

- meningitis, esp. fungal
- encephalitis
- intracranial abscess

Haematoma: extradural/subdural

Glaucoma

Benign intracranial hypertension

Q. Pitfalls (often missed)

A. Cervical spondylosis/dysfunction

Dental disorders

Refractive errors of eye

Sinusitis

Ophthalmic herpes zoster (pre-eruption)

Exertional headache

Hypoglycaemia

Post-traumatic headache

Post-spinal procedure (e.g. epidural, lumbar puncture)

Sleep apnoea

Rarities

- Paget's disease
- Post-sexual intercourse
- Cushing's syndrome
- Conn's syndrome
- Addison's disease (p. 224)
- Dysautonomic cephalgia

Q. Seven masquerades checklist

A. Depression
Diabetes
Drugs
Anaemia
Thyroid disorder
Spinal dysfunction   cervicogenic
UTI

Q. Is the patient trying to tell me something?

A. Quite likely if there is an underlying psychogenic disorder.

Probability diagnosis

The commonest cause of headache presenting in general practice is respiratory infection.[1] The most common causes of chronic recurrent headache are so-called transformed migraine, tension-type headache and combination headaches. Combination headaches, typified by relatively constant pain lasting for many days, have a mix of components such as tension, depression, cervical dysfunction, vascular headache and drug dependence. Neurologists may refer to these headaches as ‘tension-vascular headache’. Tension headache is less common than previously promulgated.[3]

Transformed migraine[5]

This describes the progressive increase in frequency of migraine attacks until the headache recurs daily. The typical migraine features become modified so that the pattern resembles that of tension headache but with the unilateral situation of migraine. Analgesic abuse can transform episodic migraine into chronic daily headache.

Serious disorders not to be missed

For the acute onset of headache it is vital not to miss subarachnoid haemorrhage (SAH) or meningitis. Intracranial haemorrhage, especially involving cerebellar, intraventricular and frontal lobe areas, needs to be considered.

Acute ‘thunderclap’ headache[5]

This is a sudden severe headache that can be caused by the following:

- enlarging aneurysm - an enlarging aneurysm or vascular malformation can cause acute headache
- SAH - the pain is typically occipital, localised at first then generalised and may vary in intensity
- meningitis - must be considered if the headache is generalised, especially in the presence of malaise, fever and neck stiffness: the ache, which is constant and severe, may begin abruptly

For chronic headache, space-occupying lesions including subdural haematomas must be considered. Since headaches tend to decrease with age, headaches developing in the elderly should be viewed with suspicion and this includes considering temporal arteritis (TA). Benign intracranial hypertension should be considered, especially in young obese women. The dangerous cryptococcal meningitis can be difficult as the CT scan may be normal.

Tips on sinister causes of headache[4]

- The most important indicator is time course: beware of acute or subacute tempo.
- Be suspicious of any focal symptoms or signs (except for typical migraine aura).
- Beware of fever, confusion, altered mental state or neck stiffness.

Pitfalls

The list (Table 56.1) contains some controversial causes of headache, although some should be obvious if a careful history is elucidated. These include post-traumatic headache, post-procedural headache (e.g. lumbar puncture and spinal anaesthesia) and exertional headache. Sinusitis can be overlooked in the absence of respiratory signs. Refractive errors of the eye, although an uncommon cause of headache, do warrant consideration.

General pitfalls

- Overinvestigating the patient with headache, especially as a substitute for a careful history and examination
- Failing to appreciate that a combination of factors and cervical dysfunction are common causes of headache
- Omitting to measure the blood pressure in the patient complaining of headache
- Rushing in with antibiotics for a patient (especially children) with fever and headache - bacterial meningitis may be masked
- Attributing the early headache of a spaceoccupying lesion to tension or hypertension

Seven masquerades checklist

Of the masquerades, depression and drugs are important causes of headache. Cervical dysfunction is certainly an important cause and tends to be ignored by some doctors. Australian figures are misleading because many of these patients tend to gravitate to alternative health professionals.

A UK study placed headache from cervical spondylosis on almost equal terms with migraine.[1] The explanation for referral of pain from disorders of the upper cervical spine to the head and eye is that some afferent fibres from the upper three cervical nerve roots converge on cells in the posterior horn of the spinal cord (which can also be excited by trigeminal afferent fibres), thus conveying to the patient the impression of head pain through this shared pathway (Figure 56.1).



Figure 56.1 Typical headache referral patterns for dysfunction of the upper cervical spinal segments

Significant drug causes are listed in Table 56.2. Anaemia can cause headache, usually if the haemoglobin level falls below 100 g/L.[5] Hypo- and hyper-thyroidism may also cause headache, and in diabetics hypoglycaemia is often responsible.

Table 56.2 Drugs that can cause headache

Alcohol
Analgesics (rebound) - aspirin - codeine
Antibiotics and antifungals
Antihypertensives - methyldopa - beta-blockers (e.g. atenolol) - hydralazine

- reserpine
- calcium-channel blockers (e.g. nifedipine)

Caffeine

Corticosteriods

Cyclosporin

Dipyridamole

Ergotamine (rebound)

H2-receptor antagonists (e.g. cimetidine, ranitidine)

MAO inhibitors

Nicotine

Nitrazepam

Nitrous Oxide

NSAIDs (e.g. indomethacin)

Oral contraceptives

PDEs inhibitors (e.g. sildanafil, tadelafil)

Retinoids

Sympathomimetics

Theophylline

Vasodilators (e.g. calcium-channel blockers, nitrates)

Psychogenic considerations

Headache, like tiredness, is one of those symptoms that may reflect a ‘hidden agenda’. Of course, the patient may be depressed (overt or masked) or may have a true anxiety state. The most characteristic feature of psychogenic headache is that the headache is present virtually every minute of the day for weeks or months on end. However, it is common for patients to deny that they are anxious, depressed or unduly stressed. For this reason a detailed history is important to identify lifestyle factors and historical events that can be associated with headache.

Some patients are fearful of their headache lest it represents a cerebral tumour, stroke or hypertension and need appropriate reassurance.

Conversion reactions and other aspects of compensation rewards, especially following an accident (e.g. rear-end collision), may make the symptom of headache difficult to manage. Headache, like backache, is one of the prime symptoms perpetuated or exaggerated for secondary gain.

Severe headaches, especially simulated migraine, are common ‘tickets of entry’ for drug addicts seeking narcotics from empathic practitioners. Such patients require very skilled management.

DIURNAL PATTERNS OF PAIN

Plotting the fluctuation of headache during the day provides vital clues to the diagnosis (Figure 56.2). The patient who wakes up with headache could have vascular headache (migraine), cervical spondylosis, depressive illness, hypertension or a space-occupying lesion. It is usual for migraine to last hours, not days, which is more characteristic of tension headache. The pain of frontal sinusitis follows a typical pattern, namely onset around 9 am, building to a maximum by about 1 pm, and then subsiding over the next few hours. In the absence of respiratory symptoms it is likely to be misdiagnosed as tension headache. The pain from combination headache tends to follow a most constant pattern throughout the day and does not usually interrupt sleep.



Figure 56.2 Typical diurnal patterns of various causes of headache; the relative intensity of pain is plotted on the vertical axis

THE CLINICAL APPROACH

History

A full description of the pain including a pain analysis should be obtained. This includes:

- site
- radiation
- quality
- severity
- frequency
- duration
- onset and offset
- precipitating factors
- aggravating and relieving factors
- associated symptoms

It is useful to get the patient to plot on a prepared grid the relative intensity of the pain and the times of day (and night) that the pain is present. The history, especially of the tempo of the condition, should help diagnose headaches secondary to specific pathology.

Key questions[6]

- Can you describe your headaches?
- How often do you get them?
- Can you point to exactly where in the head you get them?
- Do you have any pain in the back of your head or neck?

- What time of the day do you get the pain?
- Do you notice any other symptoms when you have the headache?
- Do you feel nauseated and do you vomit?
- Do you experience any unusual sensations in your eyes, such as flashing lights?
- Do you get dizzy, weak or have any strange sensations?
- Does light hurt your eyes?
- Do you get any blurred vision?
- Do you notice watering or redness of one or both of your eyes?
- Do you get pain or tenderness on combing your hair?
- Are you under a lot of stress or tension?
- Does your nose run when you get the headache?
- What tablets do you take?
- Do you get a high temperature, sweats or shivers?
- Have you had a heavy cold recently?
- Have you ever had trouble with your sinuses?
- Have you had a knock on your head recently?
- What do you think causes the headaches?

Differences between the clinical features of migraine and tension headache are presented in Table 56.3. ‘Red flag’ indicators of a serious cause of headache are outlined in the box.

Table 56.3 A comparison of typical clinical features of migraine and tension headache[7]

	Migraine	Tension headache
Family history		
Onset before 20		
Prodromata		
Bilateral		
Unilateral		
Throbbing		
Constant		
Less than 1 per week		
Continuous daily		
Lasts less than 24 h		
Vomiting		
Aggravated by the pill		
Aggravated by alcohol		
Relieved by alcohol		

Red flag pointers for headache

- Sudden onset
- Severe and debilitating pain
- Fever
- Vomiting
- Disturbed consciousness
- Worse with bending or coughing

- Maximum in morning
- Neurological symptoms/signs
- Young obese female: ? on medication
- ‘New’ in elderly, especially > 50 years

Physical examination

For the physical examination it is appropriate to use the basic tools of trade, namely the thermometer, sphygmomanometer, pen torch, and diagnostic set, including the ophthalmoscope and the stethoscope. Inspect the head, temporal arteries and eyes. Areas to palpate include the temporal arteries, the facial and neck muscles, the cervical spine and sinuses, the teeth and temporomandibular joints. Search especially for signs of meningeal irritation and papilloedema.

A mental state examination is mandatory and includes looking for altered consciousness or cognition and assessment of mood, anxiety - tension - depression, and any mental changes. Neurological examination includes assessment of visual fields and acuity, reactions of the pupils and eye movements in addition to sensation and motor power in the face and limbs and reflexes, including the plantar response. ‘Red flag’ indicators from the examination are given in the box below.

Special signs

- *Upper cervical pain sign*. Palpate over the C2 and C3 areas of the cervical spine, especially two finger breadths out from the spinous process of C2. If this is very tender and even provokes the headache it indicates headache of cervical origin.

Red flag pointers: from physical examination

- Altered consciousness or cognition
- Meningism
- Abnormal vital signs: BP, temperature, respiration
- Focal neurological signs, including pupils, fundi, eye movement
- Tender, poorly pulsatile cranial arteries

- *Ewing's sign for frotal sinusitis*. Press your finger gently upwards and inwards against the orbital roof medial to the supraorbital nerve. Pain on pressure is a positive finding and indicates frontal sinusitis.
- *The invisible pillowsign*. The patient lies on the examination table with head on a pillow. The examiner then supports the head with his or her hands as the pillow is removed. The patient is instructed to relax the neck muscles and the examiner removes the supporting hands. A positive test indicating tension from contracting neck muscles is when the patient's head does not readily change position. This is uncommon.

Investigations

Investigations can be selected from:

- haemoglobin: ? anaemia
- WCC: leucocytosis with bacterial infection
- ESR: ? temporal arteritis
- radiography
  - chest X-ray, if suspected intracerebral malignancy
  - cervical spine
  - skull X-ray, if suspected brain tumour, Paget's disease, deposits in skull
  - sinus X-ray, if suspected sinusitis
  - CT scan
    - detection of brain tumour (most effective)
    - cerebrovascular accidents (valuable)

        - SAH
    - radioisotope scan (technetium-99m) to localise specific tumours and haematoma
    - MRI: very effective for intracerebral pathology but expensive; produces better definition of intracerebral structures than CT scanning but not as sensitive for detecting bleeding; detects intracranial vasculitis in temporal arteries
    - lumbar puncture
        - diagnosis of meningitis
        - suspected SAH (only if CT scan normal)

Note: Dangerous if raised intracranial pressure.

HEADACHE IN CHILDREN

Respiratory infections and febrile illnesses are a common cause of headache in children but there are other causes that reflect the common causes in adults. Many childhood headaches are isolated but are chronic in a significant number. Migraine is relatively common before adolescence, while tension or muscle contraction headache is more common after adolescence.

Consider often overlooked causes such as hair traction, eye strain (measure and record vision) and hypoglycaemia. Children who have long periods without regular eating are prone to headache including exacerbations of migraine. They should not skip breakfast.[8]

Young children rarely experience sinus headache and this should not really be considered until the sinuses develop, around 5 years for the frontal sinuses.

From 1% of 7 year olds to 5% or more of 15 year old children suffer from migraine, with girls developing it at a higher rate2 with increasing age. There is a strong family history. As a rule the prognosis is good as the majority will have no migraines in the long term. The type is mainly common migraine with symptoms such as malaise or nausea: classic migraine with the typical aura is not a feature of childhood migraine. The rather dramatic migraine, such as vertebrobasilar migraine, is frequent in adolescent girls and hemiplegia occurs in infants and children, especially with their first migraine attack.[9] Vomiting is not necessarily an associated symptom in children.

The possibility of cerebral space-occupying lesions requires due consideration, especially if the headaches are progressive. These are present typically in the morning and are associated with symptoms such as vomiting, dizziness, diplopia, ataxia personality changes and deterioration of school performance. Symptoms that indicate a cerebral tumour or other serious problem are outlined in Table 56.4.

Table 56.4 Pointers to serious causes of headache in children

Headache features
Persistent
Present first thing in morning
Wakes child at night
No past history
No family history
Associated poor health
Associated neurological symptoms
Unilateral localisation

Source: After Wright[2]

Neonates and children aged 6 - l2 months are at the greatest risk from meningitis and it is important to keep this in mind.

Management of the non-serious causes of headache includes reassurance (especially of parents), discouragement of excessive emphasis on the symptom and simple medications, such as paracetamol for the younger child and aspirin for the adolescent. Patients with undiagnosed and/or problematic headache should be referred.

HEADACHES IN THE ELDERLY

The recent onset of headache in the elderly has to be treated with caution because it could herald a serious problem, such as a space-occupying lesion (e.g. neoplasm, subdural haematoma), TA, trigeminal neuralgia or vertebrobasilar insufficiency. Cervical spondylosis is age-related and may be an important factor in the ageing patient. Age-related headaches are summarised in Table 56.5.

Table 56.5 Age-related causes of headache

Children	• Intercurrent infections • Psychogenic • Migraine • Meningitis • Post-traumatic
Adults, including middle age	• Migraine • Cluster headache • Tension • Cervical dysfunction • Subarachnoid haemorrhage • Combination
Elderly	• Cervical dysfunction • Cerebral tumour • Temporal arteritis • Neuralgias • Paget's disease • Glaucoma • Cervical spondylosis • Subdural haemorrhage

Late life migraine can be mistaken for cerebrovascular disease, especially in the presence of preceding neurological symptoms. It is the sequence of the visual and sensory symptoms with the spread from face to tongue to hand over some minutes, with clearing in one area as it appears that helps distinguish migraine from transient ischaemic attacks (TIAs). Although some patients experience headache with TIAs it is not a distinguishing feature. Vomiting is suggestive of migraine rather than cerebrovascular disease.[9]

TENSION-TYPE HEADACHE

Tension or muscle contraction headaches are typically a symmetrical (bilateral) tightness. They tend to last for hours and recur each day. They are often associated with cervical dysfunction and stress or tension, although the patient usually does not realise the headaches are associated with tension until it is pointed out. Seventy-five per cent of patients are females.[3]

Typical clinical features

- Site: frontal, over forehead and temples (Figure 56.3)



Figure 56.3 Typical distribution of pain in tension-type headache

- Radiation: occiput
- Quality: dull ache, like a 'tight pressure feeling', 'heavy weight on top of head', 'tight band around head'; may be tightness or vice-like feeling rather than pain
- Frequency: almost daily
- Duration: hours (can last days)
- Onset: after rising, gets worse during day
- Aggravating factors: stress, overwork with skipping meals
- Relieving factors: alcohol
- Associated features: lightheadedness, fatigue, neck ache or stiffness (occiput to shoulders), perfectionist personality, anxiety/depression
- Physical examination: muscle tension (e.g. frowning), scalp often tender to touch, 'invisible pillow' sign may be positive

IHS criteria for tension-type headache

The International Headache Society (IHS) criteria for episodic tension-type headaches involve the following:

1. The patient should have had at least ten of these headaches.
2. The headaches last from 30 minutes to 7 days.
3. The headaches must have at least two of the following four:
    a. non-pulsating quality
    b. mild or moderate intensity
    c. bilateral location
    d. no aggravation with routine physical activity
4. The headaches must have both of the following:
    a. no nausea or vomiting
    b. photophobia and phonophobia are absent, or one but not the other is present
5. There should be less than 15 days of headache per month and less than 180 days per year.
6. Secondary causes are excluded.

Management[10]

- Careful patient education: explain that the scalp muscles get tight like the calf muscles when climbing up stairs.

- Counselling and relevant advice:
    - Learn to relax your mind and body.
    - During an attack, relax by lying down in a hot bath and practise meditation.
    - Be less of a perfectionist: do not be a slave to the clock.
    - Don't bottle things up, stop feeling guilty, approve of yourself, express yourself and your anger.
- Advise and demonstrate massage of the affected area with a soothing analgesic rub.
- Advise stress reduction, relaxation therapy and yoga or meditation classes.
- Medication - use mild analgesics such as aspirin or paracetamol. Discourage stronger analgesics. Avoid tranquillisers and antidepressants if possible, but consider these drugs if symptoms warrant medication (e.g. amitriptyline 10 - 75 mg (o) nocte increasing to 150 mg if necessary). Diazepam (short-term use) appears to be very effective in middle-aged men; it is prone to cause depression in women (beware of habituation).

Special notes:

- The general aim is to direct patients to modify their lifestyle and avoid tranquillisers and analgesics.
- It is unusual to be awoken from sleep.
- Beware of depression.
- Consider muscle energy therapy and/or mobilisation of the neck followed by exercises if there is evidence of cervical dysfunction.
- Recommend a meditation program.

MIGRAINE

Migraine, or the ‘sick headache’, is derived from the Greek word meaning ‘pain involving half the head’. It affects at least 1 person in 10, is more common in females (18% of women, 6% men) and peaks between 20 and 50 years. There are various types of migraine (Table 56.6), with classic migraine (headache, vomiting and aura) and common migraine (without the aura) being the best known. The most common trigger factor is stress.[3]

Table 56.6 Types of vascular headache

Common migraine (aura is vague or absent)
Classic migraine
Complicated migraine
Unusual forms of migraine
• hemiplegic
• basilar
• retinal
• migrainous stupor
• ophthalmoplegic
• migraine equivalents
• status migrainosus
Cluster headache
Chronic paroxysmal hemicrania
Menstrual migraine
Lower half headache
Benign exertional-sex headache (beware of SAH)
Miscellaneous (e.g. icepick pains, ‘ice-cream’ headache)

Source: After Day[11]

Typical clinical features of classic migraine

- Site: temporofrontal region (unilateral) (Figure 56.4); can be bilateral



Figure 56.4 Typical distribution of pain in migraine (right side)

- Radiation: retro-orbital and occipital
- Quality: intense and throbbing
- Frequency: 1 to 2 per month
- Duration: 4 to 72 hours (average 6 - 8 hours)
- Onset: paroxysmal, often wakes with it
- Offset: spontaneous (often after sleep)
- Precipitating factors: tension and stress (commonest); others in Table 56.7

Table 56.7 Migrainous trigger factors

Exogenous
• Foodstuffs - chocolate, oranges, tomatoes, citrus fruits, cheeses, gluten sensitivity (possible)
• Alcohol - especially red wine
• Drugs - vasodilators, oestrogens, monosodium glutamate, nitrites (‘hot dog’ headache), indomethacin, OCP
• Glare or bright light
• Emotional stress
• Head trauma (often minor), e.g. jarring - ‘footballer's migraine’
• Allergen
• Climatic change
• Excessive noise
• Strong perfume
Endogenous
• Tiredness, physical exhaustion, oversleeping
• Stress, relaxation after stress - ‘weekend migraine’
• Exercise
• Hormonal changes - puberty - menstruation
- climacteric
- pregnancy

- Hunger
- Familial tendency
- ? Personality factors

- Aggravating factors: tension, activity
- Relieving factors: sleep, vomiting
- Associated factors: nausea, vomiting (90%) irritability
  aura - visual 25% (scintillation, scotoma, hemianopia, fortification)
  - sensory (unilateral paraesthesia)
- Other pointers: abdominal pain in childhood; family history of migraine, asthma and eczema

DxT: headache + vomiting + visual aura ° migrane with aura (classic)

IHS criteria for common migraine

The IHS criteria for migraine without aura involves the checklist below.

1. The patient should have had at least five of these headaches.
2. The headaches last 4 - 72 hours.
3. The headache must have at least two of the following:
    a. unilateral location
    b. pulsing quality
    c. moderate or severe intensity, inhibiting or prohibiting daily activities
    d. headache worsened by routine physical activity
4. The headache must have at least two of the following:
    a. nausea and/or vomiting
    b. photophobia and phonophobia
5. Secondary causes of headache are excluded (e.g. normal exam and/or imaging study).

IHS criteria for migraine with typical aura (classic)

There should be at least two attacks, including at least three of the following:

1. reversible brain symptoms (cortical or brain stem)
2. gradual development over 4 minutes
3. aura duration less than 60 minutes
4. headache follows aura in less than 1 hour

Note: If the aura lasts longer than 1 hour, it is migraine with prolonged aura. If it lasts longer than 24 hours, it is a migrainous infarction (stroke).

Management

Patient education - provide explanation and reassurance, especially if bizarre visual and neurological symptoms are present. Patients should be reassured about the benign nature of their migraine. For each migraine sufferer, an individual treatment plan including a migraine action plan should be devised.

Counselling and advice

- Tailor the advice to the individual patient.
- Avoid known trigger factors, especially tension, fatigue, hunger and constant physical and mental stress.

- Advise keeping a diary of foodstuffs or drinks that can be identified as trigger factors. Consider a low amine diet: eliminate chocolate, cheese, red wine, walnuts, tuna, vegemite, spinach and liver.
- Practise a healthy lifestyle, relaxation programs, meditation techniques and biofeedback training.
- Be open to non-drug therapies (e.g. trial of acupuncture, hypnotherapy).

Treatment of the acute attack

- Commence treatment at earliest impending sign.
- Mild headaches may require no more than conventional treatment with '2 aspirin (or paracetamol), and a good lie down in a quiet dark room'.[11]
- Rest in a quiet, darkened, cool room.
- Place cold packs on the forehead or neck.
- Avoid drinking coffee, tea or orange juice.
- Avoid moving around too much.
- Do not read or watch television.
- For patients who find relief from simply 'sleeping off' an attack, consider prescribing temazepam 10 mg or diazepam 10 mg in addition to the following measures.[3]
- For moderate attacks use oral ergotamine or sumatriptan and for severe attacks use injection therapy.
- Avoid pethidine and similar drugs of dependence.

Medication (if necessary)[10]

First-line medication

- Aspirin or paracetamol + antiemetic: e.g. soluble aspirin (Dispirin direct) 600 - 900 mg (o) and metoclopramide 10 mg (o)
- Paracetamol (for children)

Consider NSAIDs (e.g. ibuprofen, diclofenac rapid) If nausea and vomiting is a feature:

- metoclopramide 5 - 10 mg IM or IV *or*
- prochlorperazine 12.5 mg IM or 12.5 - 25 mg rectally
- consider nasal sumatriptan

Alternatives

Choose an ergotamine preparation or a triptan preparation.

- Ergotamine (helps about 80% of patients)
  - oral: e.g. ergotamine 1 mg + caffeine 100 mg (Cafergot)
    2 tabs at 1st warning then 60 minutes if necessary (max. 6 per day)
    May need metoclopramide (o), IM or IV *or*
  - suppository: e.g. ergotamine 2 mg + caffeine 100 mg (Cafergot S)
    1 suppository at 1st warning then½ every 60 minutes (max. 3 per day) *or*
  - medihaler: e.g. 1 inhalation statim then every 5 minutes (max. 6 per day) *or*
  - IM injection: e.g. dihydroergotamine 0.5 - 1 mg, preceded by metoclopramide 10 mg IM, 20 minutes beforehand
- Sumatriptan (a serotonin receptor agonist)[10]
  - 50 - 100 mg (o) at the time of prodrome, repeat in 2 hours if necessary to maximum dose 300 mg/24 hours *or*
  - nasal spray 10 - 20 mg per nostril (up to 40 mg/24 hours) *or*
  - 6 mg, SC injection, repeat in 1 or more hours to maximum dose 12 mg/24 hours
- Zolmitriptan 2.5 - 5 mg (o), repeat in 2 hours if necessary (max. 10 mg/24 hours)
- Naratriptan 2.5 mg (o), repeat in 4 hours (max. 5 mg/24 hours)

Avoid triptans in patients with coronary artery disease, Prinzmetal angina, uncontrolled hypertension or during pregnancy. Do not use it with ergotamine simultaneously and cease if chest pain develops, albeit transient in a young patient. Use with caution in patients taking SSRIs, MAOIs and lithium.

The severe attack

(if other preparations ineffective)

Caution: Consider the possibility of underlying cerebral vascular malformation, SAH or pethidine addiction.

- If at home:[12]
    - dihydroergotamine 0.5 - 1 mg (IM) + metoclopramide 10 mg (IM) *or*
    - sumatriptan 6 mg (SC)
- If in surgery or emergency room:
    - metoclopramide 10 mg (IV) slowly over 2 minutes + oral analgesics *or*
    - metoclopramide 10 mg (IV) + dihydroergotamine 0.5 mg IV slowly *or*
    - sumatriptan 6 mg (SC)

Caution: Do not use ergotamine preparations if sumatriptan used in previous 6 hours, and do not use sumatriptan if ergotamine preparations used in previous 24 hours.

Practice tip for severe classic migraine:

IV metoclopramide + 1 litre N saline IV in 30 minutes + oral aspirin or paracetamol Continue high fluid intake

Status migrainosis: IV dihydroergotamine (may have to be given 8 hourly over 3 - 7 days in hospital) or chlorpromazine 0.1 mg/kg IV, repeated every 15 minutes for up to 3 doses (if necessary). Consider corticosteroids (e.g. dexamethasone 10 - 20 mg IV statim and then taper).

Prophylaxis

Consider prohylactic therapy for frequent attacks that cause disruption to the patient's lifestyle and wellbeing, a rule of thumb being two or more migraine attacks per month; certainly consider it for weekly attacks and a poor response to therapy for the acute attack. Do not give ergotamine.[10]

The most commonly used drugs include:

- beta-blockers - propranolol 40 mg (o) bd or tds (max. 320 mg/day), metoprolol, atenolol
- pizotifen 0.5 - 2.0 mg at night
- cyproheptadine (ideal for children)
- tricyclic antidepressants - amitriptyline
- clonidine
- methysergide (reserve for unresponsive severe migraine) 1 mg tds after food - up to 4 months only
- calcium-channel blockers - nifedipine, verapamil
- NSAIDs - naproxen, indomethacin, ibuprofen
- MAOIs - phenelzine, moclobemide
- sumatriptan
- gabapentin
- sodium valproate
- topiramate

Menstrual migraine

Naproxen 550 mg (o) bd, 48 hours before attack for 4 - 10 days

Guidelines[10, 12]

Select the initial drug according to the patient's medical profile:

- if low or normal weight - pizotifen
- if hypertensive - a beta-blocker
- if depressed or anxious - amitriptyline
- if tension - a beta-blocker
- if cervical spondylosis - naproxen
- food-sensitive migraine - pizotifen
- menstrual migraine - naproxen or ibuprofen

Commonly prescribed first-line drugs are propranolol or pizotifen:[10]

- propranolol 40 mg (o) bd or tds (at first) increasing to 320 mg daily (if necessary)
- pizotifen 0.5 - 1 mg (o) nocte (at first) increasing to 3 mg a day (if necessary)

Each drug should be tried for 2 months before it is judged to be ineffective. Amitriptyline 50 mg nocte can be added to propranolol, pizotifen (beware of weight gain) or methysergide and may convert a relatively poor response to very good control.[3]

CLUSTER HEADACHE

Cluster headache is also known as migrainous neuralgia. It occurs in paroxysmal clusters of unilateral headache that typically occur nightly, usually in the small hours of the morning, although patients may have headaches that occur at other times. A hallmark is the pronounced cyclical nature of the attacks. It occurs typically in males (6:1 ratio) and is rare in childhood. There are no visual disturbances or vomiting.

DxT: retro-orbital headache + rhinorrhoea + lacrimation ° cluster headache

Typical clinical features

- Site: over or about one eye (Figure 56.5); always same side



Figure 56.5 Typical distribution of pain in cluster headache

- Radiation: frontal and temporal regions
- Quality: severe
- Frequency: 1 - 3 times a day, at regular times like clockwork
- Duration: 15 minutes to 2 - 3 hours (average 30 minutes); the clusters last 4 - 6 weeks (can last months)
- Onset: suddenly during night (usually), same time about 2 - 3 hours after falling asleep; the 'alarm clock' headache (e.g. 2 - 4 am)
- Off set: spontaneous
- Aggravating factors: alcohol (during cluster)
- Relieving factors: drugs
- Associated features: family history; rhinorrhoea, ipsilateral nose; lacrimation; flushing of forehead and cheek; redness of ipsilateral eye; Horner's syndrome (uncommon) (Figure 56.6)



Figure 56.6 Features of an attack of cluster headache: ptosis, lacrimation and a discharge from the nostril on the side of pain

Management

Acute attack (brief treatment seldom effective):

- consider 100% oxygen 10 L/min for 15 min (usually good response)
- sumatriptan 6 mg SC injection (or 20 mg intranasal) *or*
- ergotamine (e.g medihaler or rectally)
- metoclopramide 10 mg IV + dihydroergotamine 0.5 mg IV slowly or 1 mg IM
- consider local anaesthetic - greater occipital nerve block

Avoid alcohol during cluster.

Prophylaxis (once a cluster starts)

Consider the following:

- ergotamine (take at night during a cluster): oral or dihydroergotamine IM (preferably given 1 hour prior to predicted times)
- methysergide 2 mg (o) tds
- prednisolone 50 mg/day for 10 days then reduce

- lithium 250 mg (o) bd
- verapamil SR 160 mg (o) daily up to 320 mg
- pizotifen
- indomethacin (helps confirm diagnosis)
- sodium valproate

Note: Some of the above can be used long term for frequent clusters.

CERVICAL DYSFUNCTION/SPONDYLOSIS

Headache from neck disorders, often referred to as occipital neuralgia or cervicogenic, is far more common than realised and is very rewarding to treat by physical therapy, including mobilisation and manipulation and exercises in particular.

Headache can be caused by abnormalities in any structure innervated by the upper two cervical nerves C2, C3 (usually the C1 - 2, C2 - 3 facet joints). Pain from cervical structures can be referred retro-orbitally and over one-half of the head. The headache is often incorrectly diagnosed as migraine but clinical examination of the neck helps differentiation.13 The neck may be responsible for so-called ‘tension’ headache but clinical differentiation can be more difficult.

Typical clinical features

The pain is usually sited in the occipital region with possible radiation to the parietal region, vertex of skull and behind the eye. (Figure 56.7) It is usually present on walking and settles during the day. There is usually a history of trauma including a MVA or blow to the head. Associated features include stiffness and grating of the neck. On examination there is usually tenderness to palpation over the C1, C2 and/or C3 levels of the cervical spine, especially on the side of the headache.



Figure 56.7 Typical distribution of pain in cervical dysfunction (right side)

Treatment

- Physiotherapy modalities: hydrotherapy, muscle energy therapy, mobilisation, manipulation (from experts) and neck exercises (very important)
- Supportive neck pillow
- NSAIDs for cervical spondylosis
- For intractable cases consider mobilisation under general anaesthesia, injections of corticosteroids around, or surgical section of, the greater occipital nerve.[13]

COMBINATION HEADACHE

Combined (also known as mixed) headaches are common and often diagnosed as psychogenic headache or a typical migraine. They have a combination of various degrees of:

- tension and/or depression
- cervical dysfunction
- vasospasm (migraine)
- drugs:
    - analgesics (rebound)
    - alcohol
    - nicotine
    - caffeine
    - NSAIDs

The headache, which has many of the features of tension headache, is usually described as a heavy deep ache ‘as though my head is ready to burst’. It tends to be constant, being present throughout every waking moment. It tends to last for days (average 3 - 7) but can last for weeks or months. It is often related to stress and adverse working conditions, and sometimes follows an accident.

Management

An important strategy is to evaluate each possible component of the headache as a stepwise trial by an elimination process:

- drug evaluation and modification
- cervical dysfunction - physical therapy if present
- depression
- tension and stress
- other psychogenic factors (e.g. conversion reaction)
- vasospasm

Treatment includes cognitive therapy, reassurance that the patient does not have a cerebral tumour, and lifestyle modification. The most effective medication is amitriptyline or other antidepressant.

TEMPORAL ARTERITIS

TA is also known as giant cell arteritis or cranial arteritis. There is usually a persistent unilateral throbbing headache in the temporal region and scalp sensitive with localised thickening, with or without loss of pulsation of the temporal artery. It is related to polymyalgia rheumatica - 20% of sufferers will develop TA.

Typical clinical features

Description

TA is a type of collagen disease causing inflammation of extracranial vessels, especially the superficial temporal artery. It usually presents as a unilateral intermittent headache in a person over 50 years.

- Age: over 50 years (mean age 70 years)
- Site: forehead and temporal region (unilateral) (Figure 56.8)



Figure 56.8 Typical distribution of pain in temporal arteritis (right side)

- Radiation: down side of head towards occiput
- Quality: severe burning pain
- Frequency: daily, a constant ache
- Duration: usually constant (getting worse)
- Onset: non-specific, tends to be worse in morning
- Off set: nil
- Aggravating factors: stress and anxiety
- Relieving factors: nil
- Associated features: malaise, vague aches and pains in muscles (especially of neck), weight loss
- Other pointers:
  - intermittent blurred vision
  - tenderness on brushing hair
  - jaw claudication on eating
  - polymyalgia rheumatica
  - hypertension
  - abnormal emotional behaviour

TA may also involve the intracranial vessels, especially the ophthalmic artery or posterior ciliary arteries, causing optic atrophy and blindness. Vision is impaired in about one-half of patients at some stage. Once the patient goes blind it is usually irreversible.

Diagnosis

Diagnosis is by biopsy and histological examination of the superficial temporal artery. The ESR is usually markedly elevated but may be normal. The biopsy may be normal as TA has a focal nature. MRI has a high sensitivity and specificity.

Note: Consider it with any ‘new’ headache.

Treatment

TA is very responsive to corticosteroids; start treatment immediately to prevent permanent blindness. Initial medication is prednisolone 60 mg orally daily in two divided doses initially for 2 - 4 weeks. Dose reduction and progress is monitored by the clinical state and ESR and CRP levels.10 Concomitant use of H2-receptor antagonists may be appropriate initially. Temporal arteritis may take 1 - 2 years to resolve.

FRONTAL SINUSITIS

The headache of frontal sinusitis can be a diagnostic problem especially in the absence of, or a lapse in time since, an obvious upper respiratory infection or vasomotor rhinitis. Some patients do not have a history of a preceding respiratory infection nor have signs of nasal obstruction or fever. Contrary to popular belief, sinusitis is a relatively uncommon source of headache.

Typical clinical features

It presents typically as a frontal or retro-orbital headache (Figure 56.9). A characteristic is its diurnal variation, developing in the morning around 9 am, being most intense in the middle of the day, then subsiding to off set around 6 pm.



Figure 56.9 Typical distribution of pain of frontal sinusitis (right side)

Examination

There is tenderness over the frontal sinus and pain on percussion over the sinus. Ewing's sign may be elicited. Fever and oedema of the upper eyelid may be present.

Management

Principles of treatment

- Drain the sinus conservatively using steam inhalations
- Antibiotics: amoxycillin/clavulanate or cefaclor or doxycycline
- Analgesics

Referral

If resolution cannot be accomplished by conservative means then referral to an ENT specialist is advisable. Acute purulent sinusitis can be treacherous if it persist and spreads, causing collections of pus in the extradural or subdural space, cerebral abscess or blood-borne spread of infection.

Complications

- Orbital cellulitis
- Subdural abscess
- Osteomyelitis
- Cavernous sinus thrombosis

Symptoms indicating spread of infection:

- increase in fever and chills
- vomiting
- oedema of the eyelids and forehead
- visual disturbances
- dulling of the sensorium
- convulsions

RAISED INTRACRANIAL PRESSURE

Important causes of a space-occupying lesion include a cerebral tumour and subdural haematoma. Sometimes it is not possible to differentiate between a subdural and an extradural haematoma, although the latter classically follows an acute injury. Typical features are generalised headache, usually worse in the morning, aggravated by abrupt changes in intra cranial pressure and later associated with vomiting and drowsiness. Headache is an uncommon presenting symptom of a cerebral tumour.

Typical clinical features of the headache

- Site: generalised, often occipital
- Radiation: retro-orbital
- Quality: dull, deep steady ache
- Frequency: daily
- Duration: may be hours in morning
- Onset: worse in mornings, usually intermittent, can awaken from sleep
- Off set: later in day (if at all)
- Aggravating factors: coughing, sneezing, straining at toilet
- Relieving factors: analgesics (e.g. aspirin), sitting, standing
- Associated features: vomiting (without preceding nausea); vertigo/dizziness; drowsiness; confusion (later); neurological signs (depending on side)

DxT: drowsiness + vomiting + seizure ° raised intracranial pressure

Examination

- Focal CNS signs
- Papilloedema (Figure 56.10) (but may be absent)



Figure 56.10 Papilloedema with swollen optic disc of the ocular fundus due to raised intracranial pressure

INTRACEREBRAL TUMOURS

- Incidence is 5 - 10 per 100 000 population
- Two peaks of incidence: children < 10 years[3] 35 - 60 years
- Main types of tumour:
  - children
    - medulloblastoma
    - astrocytoma (posterior fossa)
    - ependymoma
    - glioma (brain stem)
  - adults
    - cerebral glioma
    - meningioma
    - pituitary adenoma
    - cerebral metastases (e.g. lung)

Investigations

- CT scan and MRI

SUBARACHNOID HAEMORRHAGE

SAH is a life-threatening event that should not be overlooked at the primary care level. The incidence is 12 per 100000 population per annum. About 40% of patients die before treatment, while about one-third have a good response to treatment.

Clinical features:

- sudden onset headache (moderate to intense severity)
- occipital location
- localised at first, then generalised
- pain and stiffness of the neck follows
- vomiting and loss of consciousness often follow
- Kernig's sign positive
- neurological deficit may include
  - hemiplegia (if intracerebral bleed)
  - third nerve palsy (partial or complete) (Figure 56.11)



Figure 56.11 Third nerve palsy (right side)

About one-third of patients experience a ‘senital’ headache.

DxT: occipital headache + vomiting + neck stiffness ° SAH

Diagnosis

CT scanning is the investigation of choice and should be performed in the first few hours. Lumbar puncture is not necessary if the diagnosis can be made by CT, but is used if the CT scan is negative (usually 10 - 20% of cases). It may be falsely negative after 7 days. Even blood staining of CSF and xanthochromia is a positive feature on lumbar puncture.

Special notes

- Less severe headaches can cause diagnostic difficulties.
- Consider an angioma rather than an aneurysm as the cause of SAH if previous episodes.

Management

Immediate referral is required. If there is lingering doubt, review the patient within 12 - 24 hours.

MENINGITIS

The headache of meningitis is usually generalised and radiates to the neck. It is constant and severe and occasionally may begin abruptly. It is aggravated by flexion of the neck. Kernig's sign is positive. Fever and neck stiffness is usually present. Urgent referral to hospital is necessary. If meningitis is suspected or if a child or adult has headache with fever and neck stiffness, antibiotics must not be given until a lumbar puncture has been performed.

DRUG REBOUND HEADACHE

Rebound headaches are usually associated with analgesic and ergotamine dependence. A long list of over-the-counter and prescription medications can cause rebound, for example, aspirin, paracetamol, ibuprofen, opioids and caffeine. The headache is present on waking and typically persists throughout the day but fluctuates in intensity. It is a mild to moderate, dull, bilateral ache with a distribution similar to tension headache. Drug rebound headaches should be suspected in any patient who complains of headache 'all day every day'. A careful drug history should be taken. Treatment includes gradual withdrawal of the drugs and the substitution of antiemetics and sedatives or beta-blocker over about 14 days.

CHRONIC PAROXYSMAL HEMICRANIA

This is a rare headache syndrome which overlaps with cluster headache and facial pain. The unilateral pain, which can be excruciating, is located in the area of the temple, forehead, eye and upper face. It can radiate to the ear, neck and shoulder. It differs from cluster headaches in that the patients are invariably female, the paroxysms are short (average 20 - 30 minutes) and more frequent, with attacks occurring up to 14 times a day. The disorder resembles cluster headaches in nature and distribution and associated autonomic features, such as ipsilateral nasal stuffiness or rhinorrhoea, lacrimation, conjunctival injection and ptosis. The aetiology is unknown but the headache often responds dramatically to indomethacin (25 mg (o) tds).[10]

POST-LUMBAR-PUNCTURE HEADACHE[10]

This is common and is usually present when standing or sitting and rapidly improves with lying flat. It is a form of low-pressure headache, possibly due to CSF leakage. It can be severe with nausea and vomiting. In most resolution occurs within 2 - 7 days. Treatment includes bed rest until resolution. If persistent referral for an epidural blood patch is recommended.

TRIGEMINAL NEURALGIA

The pain of trigeminal neuralgia comes in excruciating paroxysms, which last for seconds to minutes only and usually affect the face rather than the head (see p. 572). The lightning-like jabs of searing or burning pain usually last 1 to 2 minutes but can last as long as 15 minutes.

Icepick headache

Icepick headaches are similar sudden stabbing pains lasting a few seconds usually at the temple (often bilateral) and are more common in migraine sufferers. They can occur unpredictably 30 or more times a day. Treatment is with indomethacin 25 mg tds.[10]

HYPERTENSION HEADACHE

It tends to occur only in severe hypertension such as malignant hypertension or hypertensive encephalopathy. The headache is typically occipital, throbbing and worse on waking in the morning.

The headache may be psychogenic in origin, developing after the diagnosis of hypertension is disclosed to the patient. However, the occasional patient has genuine headache related to milder hypertension and this serves as an accurate indicator of their blood pressure level.

BENIGN INTRACRANIAL HYPERTENSION (PSEUDOTUMOUR CEREBRI)

This is a rare but important sinister headache condition that typically occurs in young obese women. Key features are headache, visual blurring and obscurations, nausea and papilloedema. The CT and MRI scans are normal but lumbar puncture reveals increased CSF pressure and normal CSF analysis.

It is sometimes linked to drugs, including tetracyclines (most common), nitrofurantoin, oral contraceptive pill and vitamin A preparations. The main concern is visual deficits from the high intracranial pressure. Medical treatment includes weight reduction, corticosteroids and diuretics. The treatment of choice to alleviate symptoms is repeated lumbar puncture. Surgery, which involves decompression of the optic nerves or lumbo-peritoneal shunting, is sometimes required for failed medical therapy.

DxT: headache + visual obscurations + nausea ° benign intracranial hypertension

HEADACHES RELATED TO SPECIFIC ACTIVITIES

Sex headache

This can manifest as a dull or explosive headache, provoked by sexual arousal and activity, especially with orgasm. Some are clearly a form of exertional headache. Sometimes sex headache is mistaken for SAH but if the severe headache coincided with orgasm, was not associated with vomiting or neck stiffness, or settled within hours, SAH is unlikely. Treatment is with prophylactic beta-blockers or ergotamine 1 mg (0) 1 - 2 hours before activity.

Cough and exertional headache

Some people experience a severe transient pain with factors such as coughing, sneezing, stooping, straining, lifting and various sporting activities. It is usually benign and examination is normal. A CT scan is indicated if there are focal signs or if the symptoms do not settle.

Treatment is indomethacin 25 mg (0) 2 - 3 times daily for cough headache and 1 - 2 hours before exertional activity.

Gravitational headache

Occipital headache, coming on when standing upright and relieved by lying down, is characteristic of a post-lumbar puncture, an epidural block or low pressure headache. It can last for several weeks after the procedure.

‘Ice-cream’ headache

Frontal or global headache can be provoked by the rapid ingestion of very cold food and drink. It is a form of vascular headache.

WHEN TO REFER

- Evidence or suspicion of SAH or intracerebral haematoma
- Complicated migraine
- Uncertain diagnosis
- Positive neurological signs despite typical headaches

Practice tips

- A patient >55 years old presenting with unaccustomed headache has an organic disorder such as TA, intracerebral tumour or subdural haematoma until proved otherwise.
- The ESR is an excellent screening test to diagnose TA but occasionally can be normal in the presence of active TA.
- If a patient presents twice within 24 hours to the same practice or hospital with headache and vomiting, consider other causes apart from migraine before discharging the patient.[8]
- Treat an unusual or unaccustomed headache with a lot of respect.
- If migraine attacks are severe and unusual (e.g. always on the same side) consider the possibility of cerebral vascular malformation.
- CT scans and MRI have superseded other investigations in the diagnosis of cerebral tumours and intracranial haemorrhage but should be ordered sparingly and judiciously.
- If a headache is occipital in origin or accompanied by neck pain, consider the likely possibility of cervical dysfunction and refer to the appropriate therapist once the diagnosis is established.
- For recurrent migraine sufferers emphasise the importance of trigger factor avoidance and of taking aspirin and metoclopramide medication at the earliest warning of an attack.
- A severe headache of sudden onset is SAH until proved otherwise.
- SAH is overlooked sometimes, mainly because it is not considered in the differential diagnosis. Suspect with very severe and protracted headache, drowsiness and neck stiffness.
- Medical evidence indicates that most headaches are related to fatigue, stress or migraine triggers and respond to application of heat or cold, exercise and common analgesics, including aspirin and ibuprofen.[14]
- If women with migraine demand the oral contraceptive, use a low-dose oestrogen preparation and monitor progress.
- The use of narcotics for migraine treatment (such as pethidine and codeine) is to be avoided whenever possible - the frequent use of ergotamine, analgesics or narcotics can transform episodic migraine into chronic daily headache.[4]

- Headaches increasing in frequency, despite prophylaxis
- Danger signals with headache:
  - sudden onset without previous history
  - recent onset for first time in an older person
  - recurrent in children
  - progressive
  - wakes the patient at night
  - localised pain in definite area or structure (e.g. ear, eye)

- precipitated by raised intracranial pressure (e.g. coughing)
- associated neurological symptoms or signs:
  - convulsions
  - fever
  - confusion
  - impaired consciousness
  - neck stiffness
  - dizziness/vertigo
  - personality change

REFERENCES

1. Cormack J, Marinker M, Morrell D. The patient complaining of headache. In: *Practice*. London: Kluwer Medical, 1982: 3.12.
2. Wright M. Recurrent headaches in children. Aust Paediatr Revi, 1991; 1(6): 1 - 2.
3. Anthony M. Migraine and tension headache. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 313 - 16.
4. Stark R. Management of headache. Proceedings of 25th update course for GPs. Monash University, 2003.
5. Lance JW. Headache and facial pain. Med J Aust, 2000; 172: 450 - 5.
6. Davis A, Bolin T, Ham J. *Symptom Analysis and Physical Diagnosis*. Sydney: Pergamon, 1990.
7. Lance JW. *Mechanism and Management of Headache* (3rd edn). London: Butterworths, 1978: 109 - 12.
8. Smith L. Childhood headache. In: Australian Doctor Education, GP Paediatrics, 2005.
9. Burns R. Pitfalls in headache management. Aust Fam Physician, 1990; 19: 1821 - 6.
10. Tiller J (Chair). *Therapeutic guidelines: Neurology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2002: 25 - 53.
11. Day TJ. Migraine and other vascular headaches. Aust Fam Physician, 1990; 19: 1797 - 1804.
12. Heywood J, Zagami A. Treating acute migraine attack. Curr Ther, 1997; 37(12): 33 - 7.
13. Anthony M. The treatment of migraine - old methods, new ideas. Aust Fam Physician, 1993; 22: 1401 - 05.
14. Rosser W, Shafir MS. Evidence-based family medicine. Hamilton: BC Decker Inc., 1998: 164 - 6.

CHAPTER 57. HOARSENESS

Hoarseness results from imperfect phonation due to impairment of normal vocal cord mobility or vibration. It is an important symptom as it may signal a serious cause such as malignancy or a disease with potential for airway obstruction.[1]

RAYMOND L. CARROLL 1996

HOARSENESS

Hoarseness, which is an important symptom of laryngeal disease presenting in general practice, ranges from the very common, trivial, self-limiting condition of viral upper respiratory tract infection to a life-threatening disorder (Table 57.1). It may be of sudden presentation lasting only a few days or develop gradually and persist for weeks or months. The cut-off point between acute and chronic hoarseness is three weeks duration, by which time most self-limiting conditions have resolved. Hoarseness pertains to harsh, husky or rough tones of voice rather than pitch or volume. Rarely, hoarseness can be a functional or deliberate symptom referred to as ‘hysterical aphonia’.[1] In this condition, patients purposely hold the cords apart while speaking.

Table 57.1 Hoarseness: diagnostic strategy model

Q. Probability diagnosis
A. Viral URTI: acute laryngitis
Non-specific irritative laryngitis
Vocal abuse (shouting, screaming etc.)
Nodules and polyps of cords
Phonaesthenia in elderly - ‘tired’ voice
Acute tonsillitis
Q. Serious disorders not to be missed
A. Carcinoma: larynx, lung including laryngeal nerve palsy
Imminent airway obstruction (e.g. acute epiglottitis, croup)
Other rare severe infections (e.g. tuberculosis, diphtheria)
Foreign body
Motor neurone disease
Myaesthenia gravis
Q. Pitfalls
A. Toxic fumes
Vocal abuse
Benign tumours of vocal cords (e.g. polyps, ‘singers’ nodules, papillomas)
Gastro-oesophageal reflux →
pharyngolaryngitis
Goitre
Dystonia
Physical trauma (e.g. post-intubation), haematoma
Fungal infection (e.g. *Candida* with steroid inhalation, immunocompromised)
Allergy (e.g. angioedema)

Leucoplakia

Q. Masquerades

A. Consider:

- drugs: antipsychotics, anabolic steroids
- smoking →non-specific laryngitis
- hypothyroidism, acromegaly

Q. Is the patient trying to tell me something?

A. Consider:

- functional aphonia
- functional stridor

KEY FACTS AND CHECKPOINTS

- In acute hoarseness the diagnosis is usually obvious from the history alone. Examples include acute upper respiratory tract infection (URTI) or vocal overuse.
- Think ‘hypothyroidism’ if unusual hoarseness develops.
- Carcinoma of the larynx must be excluded if hoarseness persists for longer than 3 weeks in an adult. It can arise intrinsic or extrinsic to the vocal cords.
- Non-malignant vocal cord lesions include polyps, vocal nodules, contact ulcers, granulomas, other benign tumours and leucoplakia.
- In cases of chronic hoarseness the larynx must be visualised for diagnosis but the following are common:
  - children - ‘screamer's nodules’
  - adults - non-specific irritant laryngitis
- Acute laryngeal oedema may develop as a component of the life-threatening acute angioedemic allergic response.
- Elderly or debilitated patients may exhibit a shaky or soft ‘pseudohoarse’ voice due to a weakened respiratory effort. This is termed phonaesthenia.
- Contact ulcers of the larynx occur on the posterior third of the vocal cords where the mucosa is thin. The resultant weak hoarse voice may be accompanied by painful phonation. The ulcers may develop into granulomas. Apart from intubation, the condition is usually found in forceful orators who misuse their larynx when attempting to lower the pitch of their voice.[1]

THE CLINICAL APPROACH

History

Note the nature and duration of the voice change. Inquire about corticosteroid inhalations, excessive or unaccustomed voice straining, especially singing, recent surgery, possible reflux, smoking or exposure to environmental pollutants. Elicit associated respiratory or general symptoms such as cough and weight loss. Consider symptoms of hypothyroidism.

Physical examination

Palpate the neck for enlargement of the thyroid gland or cervical nodes. Perform a simple oropharyngeal examination except if epiglottitis is suspected. Check for signs of hypothyroidism, such as coarse dry hair and skin, slow pulse and mental slowing. Perform indirect laryngoscopy if skilled in the procedure.

Investigations

The following need to be considered:

- Thyroid function tests.
- Chest X-ray if it is possibly due to lung carcinoma with recurrent laryngeal nerve palsy.
- Indirect laryngoscopy (the gag reflex may preclude this).
- Direct laryngoscopy with a flexible fibreoptic endoscope with possible biopsy.
- The choice of imaging to detect suspected neoplasia or laryngeal trauma is special CT scan.

MANAGEMENT PRINCIPLES

Acute hoarseness

- Treat according to cause.
- Advise vocal rest or minimal usage at normal conversation.
- Avoid irritants (e.g. dust, tobacco, alcohol).
- Consider inhalations and cough suppressants in cases of acute URTI and coughing paroxysms.

Chronic hoarseness

- Establish the diagnosis.
- Consider referral to ENT specialist.

HOARSENESS IN CHILDREN

- It is worth bearing in mind that stridor in infants can be caused by a congenital abnormality of the larynx, including laryngomalacia (congenital laryngeal stridor), which is particularly noticeable when the child is asleep, laryngeal stenosis (congenital laryngeal narrowing) and laryngeal paralysis due to birth trauma of the vagus nerve. Vocal cord paralysis/palsy is the most common laryngeal abnormality in children (20% of cases) after laryngomalacia.[1]
- In children exclude the acute infections - laryngotracheobronchitis (croup), tonsillitis and epiglottitis.
- Persistent hoarseness in children is due commonly to vocal cord nodules related to vocal abuse, such as screaming and yelling, often due to noisy children's games.
- It is important to exclude a juvenile papilloma in a hoarse child.[3]

SPECIFIC CONDITIONS

Acute laryngitis

Most cases are caused by the respiratory viruses - rhinovirus, influenza, parainfluenza, coxsackie, adenovirus and respiratory synctial virus, resulting in vocal cord oedema (be cautious of group A *Streptococcus*). The main symptom is hoarseness, which usually persists for 3 - 14 days and leads to loss of voice. Even speaking can be painful. Aggravating factors include smoking, excessive alcohol drinking, and exposure to irritants and pollutants, air-conditioning systems and very cold weather.

Management

- Rest at home, including voice rest (the best treatment).
- Use the voice sparingly, avoid whispering.
- Use a warm sialagogue (e.g. hot lemon drinks).
- Drink ample fluids, especially water.
- Avoid smoking, passive smoke and alcohol.
- Have hot steamy showers as humidity helps.
- Use steam inhalations (e.g. 5 min, 3 times a day).

- Use cough suppressants, especially mucolytic agents.
- Use simple analgesics, such as paracetamol or aspirin, for discomfort.
- Antibiotics are of no proven use unless there is evidence of bacterial infection.

Chronic laryngitis: 'barmaid syndrome'

This typically occurs in a heavy smoker who works in a heavy smoking environment, who is a heavy drinker and continually talks or sings. It is a combination of vocal abuse and chemical irritation. Treatment involves modification of these factors and screening for vocal cord tumours.

Benign tumours of the vocal cords

These include nodules, polyps and papules. Vocal cord nodules, including 'singer's nodules', may respond well to conservative measures such as a voice rest and vocal therapy. If not, they can be removed by microlaryngeal surgery or laser therapy. Dependent polyps and papillomas are removed by microsurgery.

Carcinoma of the larynx

This usually occurs in patients with a history of chronic laryngitis and smoking. It may be preceded by leucoplakia which is treated by vocal cord stripping under microsurgery. The diagnosis based on persistent hoarseness is made after fibre-optic laryngoscopy and biopsy by a specialist. The patient may present with an unexplained cervical lymph node. The condition is curable if detected early. Small local tumours can be treated by radiotherapy or laser therapy. Larger tumours usually require laryngectomy and perhaps dissection of the cervical lymph nodes (Commando operation). Such radical surgery demands considerable patient support, including education about speech, eating and tracheostomy care.

Practice tips

- Consider intubation as a possible cause of transient hoarseness, especially in these times of day surgery.
- Consider gastro-oesophageal reflux disease in the elderly but avoid such a diagnosis without specialist investigation for other causes.
- If stridor is present with acute hoarseness, the airway is compromised. Be on stand-by for possible emergency intervention.
- Prevention is the best treatment for laryngeal carcinoma (i.e. quit smoking).

WHEN TO REFER[1]

- Acute cases that are unexplained, fail to respond by 3 - 4 weeks or recur.
- All chronic cases.
- Any case with stridor or non-tender cervical lymphadenopathy.
- Patients requiring voice therapy when vocal abuse is identified.

REFERENCES

1. Carroll RL. Hoarseness. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 239 - 40.
2. Havas TE. How to treat: Hoarseness and voice dysfunction. Australian Doctor, 2006; 26 May: 29 - 36.
3. Birman C, Fitzsimons, Quayle S. Little voices: therapy update. Australian Doctor, 2004; 7 May: 49 - 50.

CHAPTER 58. JAUNDICE

> The disease is produced by black bile when it flows into the liver. The symptoms are these: ‘an acute pain in the liver, also below the breast, a feeling of suffocation is strong during these days and becomes less strong later’. The liver is tender to palpation and the complexion of the patient is somewhat livid. These are symptoms that occur in the beginning but as the disease progresses, the fever diminishes in strength and the patient feels sated after ingesting a little amount of food. He must drink melikration [a mixture of water and honey].
>
> HIPPOCRATES ON HEPATITIS

JAUNDICE

Jaundice is a yellow discolouration of the skin and mucosal surfaces caused by the accumulation of excessive bilirubin.[1] It is a cardinal symptom of hepatobiliary disease and haemolysis. Important common causes include gallstones, hepatitis A, hepatitis B, hepatitis C, drugs, alcohol and Gilbert's syndrome. The commonest clinical encounter with jaundice, especially physiological jaundice, is in the newborn. As for all patients, the history and examination are paramount, but investigations are essential to clinch the diagnosis of jaundice.

The three major categories of jaundice are (Figure 58.1):


red blood cells
liver
unconjugated bilirubin
spleen
conjugated bilirubin
urobilinogen
kidneys
gut
bilirubin oxidised by bacteria in gut to urobilinogen
urobilinogen excreted in faeces as sterocobilin
urinary urobilin

Figure 58.1 The jaundice pathway

- obstructive
    - extrahepatic
    - intrahepatic
- hepatocellular
- haemolysis

KEY FACTS AND CHECKPOINTS

- Jaundice is defined as a serum bilirubin level exceeding 17 μmol/L.[2]
- Clinical jaundice manifests only when the bilirubin level exceeds 50 μmol/L.[1]
- However, jaundice is difficult to detect visually below 85 μmol/L if lighting is poor.
- It can be distinguished from yellow skin due to hypercarotenaemia (due to dietary excess of carrots, pumpkin, mangoes or pawpaw) and hypothyroidism by involving the sclera.

- The most common causes of jaundice recorded in a general practice population are (in order) viral hepatitis, gallstones, carcinoma of pancreas, cirrhosis, pancreatitis and drugs.[3]
- Always take a full travel, drug and hepatitis contact history in any patient presenting with jaundice.
- A fatty liver (steatosis) can occur not only with alcohol excess but also with obesity, diabetes and starvation. There is usually no liver damage and thus no jaundice.

Table 58.1 Abbreviations used in this chapter

Hepatitis A virus	HAV
Hepatitis A antibody	anti-HAV
Immunoglobulin M	IgM
Immunoglobulin G	IgG
Hepatitis B virus	HBV
Hepatitis B surface antigen	HBsAg
Hepatitis surface antibody	anti-HBs
Hepatitis B core antibody	anti-HBc
Hepatitis Be antigen	HBeAg
Hepatitis C virus	HCV
Hepatitis C virus antibody	anti-HCV
Hepatitis D (Delta) virus	HDV
Hepatitis E virus	HEV
Hepatitis F virus	HFV
Hepatitis G virus	HGV

A DIAGNOSTIC APPROACH

A summary of the diagnostic safety model is presented in Table 58.2.

Table 58.2 Jaundice (adults): diagnostic strategy model

Q. Probability diagnosis
A. Hepatitis A, B, C (mainly B, C) Gallstones Alcoholic hepatitis/cirrhosis
Q. Serious disorders not to be missed
A. Malignancy • pancreas • biliary tract • hepatocellular (hepatoma) • metastases Severe infections • septicaemia • ascending cholangitis

- fulminant hepatitis
- HIV/AIDS

Rarities

- Wilson's syndrome
- Reye's syndrome
- Acute fatty liver of pregnancy

Q. Pitfalls (often missed)

A. Gallstones

Gilbert's syndrome

Cardiac failure

Primary biliary cirrhosis

Autoimmune chronic active hepatitis

Chronic viral hepatitis

Haemochromatosis

Viral infections (e.g. CMV, EBV)

Q. Seven masquerades checklist

A. Depression -

Diabetes -

Drugs

Anaemia

Thyroid disorder -

Spinal dysfunction -

UTI -

Q. Is this patient trying to tell me something?

A. Not usually applicable.

Probability diagnosis

The answer depends on the age and social grouping of the patient, especially if the patient indulges in risk-taking behaviour or has travelled overseas.

Viral hepatitis A, B or C accounts for the majority of cases of jaundice.

In the middle-aged and elderly group, a common cause is obstruction from gallstones or cancer. It is common for older people to have painless obstructive jaundice; bear in mind that the chances of malignancy increase with age.

Alcoholic liver disease is common and may present as chronic alcoholic cirrhosis with liver failure or as acute alcoholic hepatitis. It is worThemphasising that such patients can make a dramatic recovery when they cease drinking alcohol.

In family practice we encounter many cases of drug-induced jaundice, especially in the elderly. These drugs are outlined later in the chapter, under the ‘seven masquerades’.

Serious disorders not to be missed

Malignancy must always be suspected, especially in the elderly patient and those with a history of chronic active hepatitis (e.g. post hepatitis B or C infection). The former is more likely to have carcinoma of the head of the pancreas and the latter, hepatocellular carcinoma (hepatoma).

Metastatic carcinoma must be kept in mind, especially in those with a history of surgery, such as large bowel cancer, melanoma and stomach cancer.

Hepatic failure can be associated with severe systemic infection (e.g. septicaemia and pneumonia), and after surgery in critically ill patients. A patient who has the classic Charcot's triad of upper abdominal pain, fever (and chills) and jaundice should be regarded as having ascending cholangitis until proved otherwise. Wilson's syndrome, although rare, must be considered in all young patients with acute hepatitis. A history of neurological symptoms, such as a tremor or a clumsy gait, and a family history is important. If Wilson's syndrome is suspected an ocular slit lamp examination, serum ceruloplasmin levels (low in 95% of patients) and a liver biopsy should be performed. Early diagnosis and treatment mean a better prognosis.

Reye's syndrome is a rare and severe complication of influenza and some other viral diseases, especially in children when given aspirin. There is rapid development of hepatic failure and encephalopathy.

Pitfalls

Gallstones, especially in the absence of upper abdominal pain, can be overlooked, so this possibility should be kept in mind in the elderly.

Gilbert's syndrome is worth considering, especially as it is the commonest form of unconjugated hyperbilirubinaemia. It affects at least 3% of the population. Like the more severe but rarer Crigler-Najjar syndrome, there is a deficiency of glucuronyl transferase. In Gilbert's syndrome the serum bilirubin level, which may rise to 50 μmol/L but seldom higher,2 tends to fluctuate and to rise during intercurrent infections, such as influenza and in episodes of fasting. All other liver function tests are normal, as is liver serology, but a history of intermittent mild jaundice, a family history or vague right upper quadrant pain may be useful pointers. Patients diagnosed by the author appeared to have a consistently coloured skin resembling a 'suntan' despite living in a cool climate. Gilbert's syndrome is benign with an excellent prognosis and no treatment is required.

Cardiac failure can present as jaundice with widespread tenderness under the right costal margin. It can be insidious in onset or manifest with gross acute failure. It can be confused with acute cholecystitis. The biochemical abnormalities seen are very variable. Usually there is a moderate rise in bilirubin and alkaline phosphatase and sometimes, in acute failure, a marked elevation of transaminase may occur, suggesting some hepatocellular necrosis.

There are many other pitfalls for a family doctor, who may encounter the conditions very rarely, if at all. Such disorders include:

- inherited conjugated hyperbilirubinaemias (Dubin-Johnson and Rotor syndromes) caused by faulty excretion by liver cells
- haemochromatosis (associated pigmentation and diabetes)
- chronic active hepatitis
- primary biliary cirrhosis
- sclerosing cholangitis (associated with ulcerative colitis)

General pitfalls

- Excluding jaundice by examining the sclera in artificial light
- Not realising that the sclera in elderly patients often have an icteric appearance (without jaundice)
- Omitting to take a careful history, including illicit drugs
- A liver biopsy is essential in all patients with chronic hepatitis

Seven masquerades checklist

Of this group the haemolytic anaemias and drugs have to be considered.

Drug-related jaundice

Drug-induced jaundice is common and many drugs are implicated. The patterns of drug-related liver damage include cholestasis, necrosis (‘hepatitis’), granulomas, chronic active hepatitis, cirrhosis, hepatocellular tumours and veno-occlusive disease.[4, 5] Some drugs, such as methyldopa, can initiate haemolysis.

The important drugs to consider are presented in Table 58.3. Antibiotics, especially flucloxacillin, amoxycillin + clavulanate and erythromycin, are commonly implicated.

Table 58.3 Drugs that can cause jaundice

Haemolysis • methyldopa
Hepatocellular damage ***Dose-dependent*** • Paracetamol (can cause acute hepatic necrosis) • Salicylates • Tetracycline ***Dose-independent*** • Anaesthetics (e.g. halothane) • Antidepressants (e.g. MAOIs) • Antiepileptics (e.g. phenytoin, sodium valproate, carbamazepine) • Antibiotics (e.g. penicillins, sulphonamides) • Antimalarials (e.g. Fansidar) • Antituberculosis (e.g. isoniazid) • Anti-inflammatories (e.g. NSAIDs, various) • Carbon tetrachloride • Cardiovascular (e.g. amiodarone, methyldopa, perhexilene) • Statins (e.g. simvastatin)
Cholestasis • Antithyroid drugs • Chlorpromazine • Erythromycin estolate • Penicillins, esp. flucloxacillin • Gold salts • Oral contraceptives/oestrogens • Synthetic anabolic steroids (e.g. methyltestosterone) • Hypoglycaemic drugs (e.g. chlorpropamide) • Amitriptyline
Others • Allopurinol • Cimetidine (aggravated by alcohol) • Cytotoxics (e.g. methotrexate, azathioprine) • Etretinate

- Hydralazine
- Nitrofurantoin
- Vitamin A (mega dosage)
- Various complementary medicines, e.g. herbal agents

Haemolysis

The patient may present with the symptoms of underlying anaemia and jaundice with no noticeable change in the appearance of the urine and stool. The degree of haemolysis may vary from the lemon yellow tinge of pernicious anaemia in an elderly patient to a severe haemolytic crisis precipitated by drugs or broad beans (favism) in a patient with an inherited red cell deficiency of glucose-6-phosphate dehydrogenase (G-6-PD). More common causes include the hereditary haemolytic anaemias, such as congenital spherocytosis and thalassaemia major. Acquired causes include incompatible blood transfusions, malignancies, such as lymphoma, severe sepsis and some drugs.

Splenomegaly occurs in most patients with haemolytic anaemia, and decreased red cell survival can be measured.

Psychogenic considerations

This is not really applicable for an organic problem such as jaundice. Nevertheless, the cause may be related to factors in the patient's lifestyle, such as homosexuality, sexual promiscuity or intravenous drug abuse, and the patient may be reluctant to offer this information. Discreet, concerned probing will be necessary.

THE CLINICAL APPROACH

History

The history should include questioning about the following:

- any episodes of jaundice
- change in colour of faeces and urine
- anorexia, sore throat, weight loss, pruritus
- abdominal pain
- residence and members of household
- contact with patients with hepatitis or jaundice
- recent overseas travel
- exposure to blood or blood products
- needle-stick injuries or exposure to needles, such as acupuncture, tattooing and intravenous drugs
- dietary history - shellfish, drinking water
- sexual history - evidence of promiscuity
- drug history, including alcohol
- recent medical history, including surgery
- family history - family contacts who have had jaundice, haemolytic disease and other genetic liver diseases
- ethnic history - liable to haemolytic disease, contact with hepatitis B
- occupational history - exposure to hazards

Significance of various symptoms

- Pain in the right hypochondrium
    - gallstones
    - acute hepatitis (a constant ache)
    - cholecystitis
- Anorexia, dark urine, fever

  - viral hepatitis probable
  - alcoholic liver disease possible
  - drug-induced hepatitis possible
- Pruritus
  - cholestasis probable
  - possible with all liver diseases
- Arthralgia, rash
  - viral hepatitis
  - autoimmune hepatitis

The examination

The abdominal examination is very important. The liver should be palpated carefully for enlargement, consistency and tenderness under the right costal margin. Search for enlargement of the gall bladder and the spleen. The gall bladder lies in the transpyloric line. A palpable gall bladder indicates extrahepatic biliary obstruction, and splenomegaly may indicate haemolytic anaemia, portal hypertension or viral hepatitis. Test for ascites.

Skin excoriation may indicate pruritus, which is associated with cholestatic jaundice. Look for evidence of chronic liver disease, such as palmar erythema, easy bruising, spider naevi and muscle wasting, and testicular atrophy and gynaecomastia. Test for hepatic flap (asterixis) and fetor, which indicate liver failure. Search for lymphadenopathy, which may be indicative of malignancy.

A summary of the possible findings is presented in Figure 58.2.


General
• jaundice
• loss of body hair
• fever
Epstein-Barr mononucleosis
• lymphadenopathy
• tonsillitis
Liver
• large or small
enlarged gall bladder
(carcinoma head pancreas)
Tremor
• Wilson's syndrome
• hepatic flap
alcoholic facies
parotid enlargement
spider naevi
scratch marks
(if obstructive jaundice)
gynaecomastia
? splenomegaly
needle marks
dilated abdominal veins
(if cirrhosis)
liver palms
leuconychia
testicular atrophy

Figure 58.2 Possible findings on examining the jaundiced patient

Investigations

The main investigations are the standard LFTs and viral serology for the infective causes, particularly hepatitis A, B and C virus (also EBV and CMV).

A summary of the general findings for liver function tests is shown in Table 58.4. Consideration should be given to ordering fractionalisation of bilirubin to determine whether it is conjugated or unconjugated (important in diagnosis of Gilbert's syndrome).

Table 58.4 Characteristic liver function tests for selected types of liver disease

Liver function tests (serological)	Hepatocellular (viral) hepatitis	Haemolytic jaundice	Obstruction	Gilbert's syndrome	Liver metastases/abscess	Alcoholic liver disease
Bilirubin	↑to ↑ ↑	↑ unconjugated	↑to ↑ ↑ ↑	↑up to 50 unconjugated	↑to N	↑to N
Alkaline phosphatase	↑ < 2N	N	↑ ↑ ↑ > 2N	N	↑ ↑to ↑ ↑ ↑	↑
Alanine transferase (ALT)	↑ ↑ ↑ > 5N	N	N or ↑	N	↑	↑
Gamma glutamyl transferase	N or ↑	N	↑ ↑	N	↑	↑ ↑ ↑
Albumin	N or ↑	N	N	N	N to ↓	N to ↓ ↓
Globulin	N or ↑	N	N	N	N	N to ↑

N: is within normal limits

Markers for hepatitis

Serology includes the test for hepatitis B virus (HBV) surface antigen (HBsAg), which is diagnostic of infection. Other markers of HBV are HBeAg antigen and the antibodies (anti-HBs and anti-HBc).

Hepatitis A virus is diagnosed by the presence of IgM antibody for recent infection and IgG antibody which indicates past infection and lifelong immunity.

Specific serological tests for the previously designated non-A, non-B hepatitis are more complex, but tests to identify two of these causative agents (C and D) are now available.

Hepatobiliary imaging

Tests to identify causes such as malignancy or gallstones are now sophisticated and should be chosen with care.

- X-ray: a plain abdominal X-ray shows up to 10% of gallstones
- Ultrasound (US): the most useful investigation for detecting gallstones and dilatation of the common bile duct; also detects liver metastases and other diffuse liver diseases
- HIDA scintiscan: useful in diagnosis of acute cholecystitis
- CT scan: for diagnosis of enlargement of the head of the pancreas and other pathology; indicated if US unsatisfactory
- PTC: percutaneous transhepatic cholangiography
- ERCP: endoscopic retrograde cholangiopancreatography; PTC and ERCP (best) determine the cause of the obstruction
- MRCP: magnetic resonance cholangiography provides non-invasive planning for obstructive jaundice
- Liver isotopic scan: useful for liver cirrhosis, especially of the left lobe

Specific tests

Some Specific tests include:

- autoantibodies for autoimmune chronic active hepatitis and primary biliary cirrhosis
- carcinoembryonic antigen to detect liver secondaries, especially colorectal
- serum iron studies, especially transferrin saturation - elevated in haemochromatosis

- alpha-fetoprotein - elevated in hepatocellular carcinoma; mild elevation with acute or chronic liver disease (e.g. cirrhosis)
- serum ceruloplasmin level - low in Wilson's syndrome
- liver biopsy
- EBV/cytomegalovirus serology (consider if hepatitis serology negative)

JAUNDICE IN CHILDREN

Jaundice in the infant

Jaundice in the newborn is clinically apparent in 50% of term babies and more than 80% of preterm.6 Icterus is therefore common and invariably physiologically benign. However, it is a cause for concern as there are many other causes and investigation is needed to determine whether the bilirubin is conjugated (always pathological) or unconjugated. If conjugated, consider the serious biliary atresia (stools are white); also a cyst obstructing the bile duct or neonatal hepatitis. Prompt referral is essential.

Jaundice occuring in the first 24 hours after birThis not due to immature liver function but is pathological and usually due to haemolysis consequent on blood group incompatibility. In primigravidas it is usually due to ABO incompatibility.

Bilirubin encephalopathy

Unconjugated bilirubin can be regarded as a neurological poison. With increasing serum levels an encephalopathy (which may be transient) can develop, but if persistent can lead to the irreversible brain damage known as kernicterus. The level of bilirubin causing kernicterus is totally unpredictable, but a guideline as a cause for concern in babies with Rh disease is a serum unconjugated bilirubin of 340 μmol/L (20 mg/dL).

Guidelines for treatment for hyperbilirubinaemia (at 24 - 36 hrs) - an example

- > 285 μmol/L - phototherapy
- > 360 μmol/L - consider exchange transfusion

An example of a normogram is presented in Figure 58.3.



Figure 58.3 Typical normogram for decision making in healthy infants with jaundice

Physiological jaundice

This mild form of jaundice, which is very common in infants, is really a diagnosis of exclusion. In a term infant the serum bilirubin rises quickly after birth to reach a maximum by day 3 - 5, then declines rapidly over the next 2 - 3 days before fading more slowly for the next 1 - 2 weeks. Management includes phototherapy.

Pathological jaundice

There are many causes of pathological jaundice, including:

- haemolysis (e.g. blood grouping incompatibilities, ABO or Rhesus, G-6-PD deficiency, hereditary spherocytosis)
- polycythaemia (e.g. intrauterine growth retardation)
- inherited conjugation defects (e.g. uridyl diphosphate glucuronyl transferase deficiency)
- breast milk jaundice
- drugs
- sepsis
- hypothyroidism
- biliary atresia

Such cases require referral for evaluation and management.

ABO blood group incompatibility

This is antibody-mediated haemolysis (Coomb's test positive):

- Mother is O
- Child is A or B

Jaundice develops within first 24 hours.

Perform a direct Coomb's test on infant.

Phototherapy is required immediately.

These children require follow up developmental assessment including audiometry.

Breast milk jaundice

If the secondary causes of prolonged jaundice are excluded, the baby is well and breastfeeding, the likely cause of unconjugated elevated bilirubin is breast milk jaundice. It occurs in 2 - 4% of breastfed infants. It usually begins late in the first week and peaks at 2 - 3 weeks. Diagnosis is confirmed by suspending (not stopping) breastfeeding for 24 - 48 hours. The serum bilirubin falls and then breastfeeding can continue. The mother, who can express milk for this short period, must be reassured that there is nothing wrong with the milk and advised to resume. Some doctors recommend continuation of breastfeeding.

Jaundice in older children

Viral infection is the commonest cause of jaundice in the older child, especially hepatitis A and hepatitis B. It is uncommon for viral hepatitis to become chronic in childhood.

JAUNDICE IN THE ELDERLY

If an elderly person presents with jaundice the usual causes and investigations have to be considered. Obstructive jaundice is the commonest form of jaundice in the elderly and may be caused by gallstones blocking the common bile duct (may be painless) and carcinoma of the head of the pancreas, the biliary tract itself, the stomach or multiple secondaries for other sites. While it is not uncommon for a gallstone to produce marked obstructive jaundice and yet be painless, it is appropriate to adhere to the old adage that painless obstructive jaundice is due to neoplasm - particularly if the gall bladder is palpable (Courvoisier's law).

Alcoholic liver disease, although most frequently affecting patients between 40 and 60 years, can present for the first time over age 60 years. The commonest cause of hepatocellular jaundice in the elderly is probably alcoholic cirrhosis; hepatitis A is still relatively uncommon in old persons.

Drugs do not cause jaundice in the elderly as frequently as they once did, particularly as phenothiazines, especially chlorpromazine, are not prescribed as often as previously. However, drugs should be considered as a potential cause and a careful check of the drug history is important.

INFECTIVE CAUSES OF JAUNDICE

A generation ago hepatitis A (infectious hepatitis or yellow jaundice) was the commonest recognised form of viral hepatitis, presenting usually with an abrupt onset of fever, anorexia, nausea and vomiting. It usually occurred in epidemics and hence was common in overcrowded institutions and camps. Now hepatitis B and C are the most commonly reported types of viral hepatitis with an onset that is more insidious and with a longer incubation period.7, 8 Symptoms include malaise, anorexia, nausea and polyarthritis.

The various forms of hepatitis are summarised in Table 58.5. All forms of hepatitis are common in developing countries and travellers are at risk of contracting these infections: hepatitis A and E from faeco-oral transmission; and hepatitis B, C, D and G from intravenous drugs and bodily fluids (from sexual transmission, in particular, for hepatitis B).

Table 58.5 Characteristic profiles of viral hepatitis A - E

Characteristic	Hepatitis A	Hepatitis B	Hepatitis C	Hepatitis D	Hepatitis E
Pseudonyms	Infectious hepatitis	Serum hepatitis	Parenterally transmitted non-A, non-B	Delta hepatitis	Enterically transmitted non-A, non-B
Agent (virus)	27 nm RNA	42 nm DNA	50 nm RNA	35 nm RNA	30 nm RNA
Transmission	Faecal-oral	Blood and other body fluids	Blood ? other body fluids	Blood and otherbody fluids	Faecal - oral
Incubation period	15 - 45 days	40 - 180 days	14 - 180 days	30 - 50 days	15 - 45 days
Severity of acute illness	Mild to moderate; often subclinical - no jaundice	Mild to severe; jaundice common; arthralgia and rash common	Mild to moderate; often subclinical	Moderate to severe; high mortality; usually jaundice	Mild to moderate; often subclinical
Chronic liver disease	No	Yes 5 - 10%	Yes 20 - 50%	Yes Potentially worst	No
Mortality	0.1 - 0.2%	1 - 3%	1 - 2%	Variable	Variable; high (10 - 20%) in

					pregnant women
Carrier state	No	Yes	Yes	Yes	Uncertain
Risk in travellers	Yes, applies to all A - E: East and South-East Asia, Asian subcontinent (e.g. India), South Pacific Islands (e.g. Fiji), sub-Saharan Africa, Mexico, Russia, other developing countries. A and E with poor sanitation; B, C, D also with IV drug use; B, D sexual contact.				
Antigens	HAV Ag	HBsAg, HBcAg, HBeAg	HCV Ag	HDV Ag	?
Serology	IgM anti-HAV diagnosis	HBsAg diagnosis anti-HBs - exposure immunity	anti-HCV (antibody) HCV-RNA (PCR) HCV genotype	HBsAg + ve HDsAg + ve anti-HDV (antibody)	Being developed
Immunoprophylaxis	Normal Ig	HB Ig	? Ig effective	None	None
Vaccine	Hepatitis A vaccine	Hepatitis B vaccine	None	Hepatitis B vaccine	None

Evidence points to more viruses causing non-ABC hepatitis.9 Hepatitis F virus has been claimed to be transmitted enterically while the newly designated hepatitis G virus (HGV) is transmitted parenterally. It does not appear to cause a severe illness in recipients. It can be predicted that the hepatitis alphabet will continue to expand.

In hepatitis A liver damage is directly due to the virus, but in hepatitis B and C it is due to an immunologic reaction to the virus.

Other infections that can present with jaundice as part of a systemic disease are malaria, Epstein-Barr mononucleosis, cytomegalovirus, Q fever, toxoplasmosis, leptospirosis and, rarely, measles, varicella, yellow fever, rubella, herpes simplex, dengue fever, Lassa fever and Marburg and Ebola virus.

Hepatitis A

Hepatitis A is becoming relatively less prevalent in first world countries. It is enterically transmitted and arises from the ingestion of contaminated food, such as shellfish, or water. There is no carrier state and it does not cause chronic liver disease. Hepatitis A most often causes a subclinical or self-limited clinical illness.

Clinical features

Preicteric (prodromal) phase:

- anorexia, nausea ±vomiting
- malaise
- headache
- distaste for cigarettes in smokers
- mild fever
- ±diarrhoea
- ±upper abdominal discomfort

Icteric phase (many patients do not develop jaundice):

- dark urine
- pale stools

- hepatomegaly
- splenomegaly (palpable in 10%)

Recovery usually in 3 - 6 weeks.

Fulminant hepatitis with liver coma and death may occur but is rare.

Investigations

LFTs and viral markers confirm the diagnosis. The antibodies to HAV are IgM, which indicates active infection, and IgG antibodies, which means immunity and which is common in the general population. Ultrasound is useful to exclude bile duct obstruction, especially in an older patient.

Outcome and treatment

Hepatitis A has an excellent prognosis with most patients making a complete recovery, and patients should be reassured. The mortality is less than 0.5%. Admission to hospital is not usually necessary. There is no Specific treatment, so management is as follows.

- Provide appropriate reassurance and patient education.
- Rest as appropriate.
- Follow a fat-free diet.
- Avoid alcohol, smoking and hepatotoxic drugs (until recovery).
- Advise on hygiene at home to prevent spread to close contacts and family members.
- Wash hands carefully after using the toilet and disinfect them with antiseptic.
- Do not handle food for others with your fingers.
- Do not share cutlery and crockery during meals.
- Do not use tea-towels to dry dishes.

Prevention

Simple health measures such as good sanitation, effective garbage disposal and hand washing are probably responsible for the major decrease in the disease. Immune serum globulin (0.03 - 0.06 mL/kg IM) confers satisfactory passive immunity for close contacts (within 2 weeks of contact) and for travellers to endemic areas for up to 3 months. An active vaccine consisting of a two-dose primary course is the best means of prevention.

Hepatitis B

Hepatitis B has protean clinical manifestations. Transmission is by blood spread, sexual transmission, perinatal spread or by close prolonged family contact. Infection may be subclinical or self-limited acute hepatitis. Fulminant hepatitis is rare. Five per cent of subjects go on to become chronic carriers of the virus. Most are ‘healthy carriers’ but some may develop chronic active hepatitis, cirrhosis and hepatoma. The serology of hepatitis B involves antibody responses to the four main antigens of the virus (core, DNA polymerase, protein X and surface antigens). Passive and active vaccines are available, and should be used freely in groups at risk, including babies of infected mothers. High-risk groups are presented in Table 58.6. The clinical features are the same as those found in hepatitis A infection but may be less abrupt in onset but more severe in the long term.7 A serum sickness-like immunological syndrome may be seen with transient rashes (e.g. urticaria or a maculopapular rash), and polyarthritis affecting small joints in up to 25% of cases in the prodromal period.

Table 58.6 Higher risk groups for contracting hepatitis B (vaccination advisable)[7]

Babies born to hepatitis B positive (carrier) mothers
Sexual partners of hepatitis B carriers (especially acute HBV)
Household contacts of hepatitis B carriers
Intravenous drug users

Recipients of blood or blood products (prior to testing)

Male homosexuals

Kidney dialysis patients

Sex industry workers

Health care workers

Garbage collectors

Institutionalised intellectually disabled patients

Prisoners

Travellers to endemic areas

Investigations[10, 11]

The main viral investigation for HBV is HBsAg (surface antigen), which is searched for routinely. If detected, a full viral profile is then formed.

HBsAg may disappear or persist. Its presence indicates a current or chronic infection as well as a carrier state (Figure 58.4). Hepatitis B carriage is the presence of HBsAg for at least 6 months.



Figure 58.4 Time course of clinical events and serological changes following infection with hepatitis B

Monitoring and outcome[10, 11]

The possible course of events is shown in Figure 58.5. The majority of patients recover completely with the outcome depending on several factors, including the virulence of the virus and the immune state and age of the patient. Some will develop chronic hepatitis, some will develop a fulminant course, and others will become asymptomatic carriers and present a health risk to others.


(a)
Acute hepatitis B infection
95% of infant-acquired infections
3–6% of adult-acquired infections
chronic HBV infection
chronic hepatitis
15–40% lifetime risk
1.3–2.5% per annum
cirrhosis
6–15% in 5 years
20–23% in 5 years
hepatocellular carcinoma
liver failure
liver transplant
(b)
Acute hepatitis C infection
60–80% are chronically infected
20–40% spontaneously clear infection
chronic hepatitis—minimal to severe inflammation and fibrosis
bridging fibrosis
cirrhosis in 10–25% develops over 20–30 years
hepatocellular carcinoma
liver transplantation
liver failure
death

Figure 58.5 Natural history of (a) hepatitis B infection; (b) hepatitis C infection. Source: W. Sievert, B. Katz, Department of Gastroenterology, Monash Medical Centre

Monitor progress with 6 - 12 monthly LFTs, HBeAg and HBV DNA.

- Negative HBeAg and HBV DNA (with anti-HBe) ° resolving, with anti-HBS ° full recovery.
- Positive HBeAg and HBV DNA ° replicating and infective - refer.
- Monitor LFTs every 6 months. Refer if ALT elevated.

Treatment[5]

There is no Specific treatment, initially and appropriate reassurance and patient education is necessary. Advise avoidance of alcohol. Avoid certain drugs, e.g. sedatives, NSAIDs, OCP, until recovery (normal LFTs). Advise about prevention of transmission especially safe sex and sharing needles. Treatment of chronic hepatitis B infection (abnormal LFTs) is with the immunomodulatory and antiviral agents alpha-interferon and lamivudine. New approved anti-viral agents are adefovir and entecavir. This is expensive but it achieves permanent remission in 25% of patients, and temporary remission in a further 25%.7 Liver transplantation has been performed, but is often followed by recurrence of hepatitis B in the graft ed liver. Follow up with regular LFTs and alphafetoprotein screening.

Serology guidelines

HBsAg ° persistent infection

anti-HBs ° past infection and immunity

HBeAg ° highly infectious

HBV DNA ° circulating virus

anti-HBc IgM ° recent infection

anti-HBc IgG ° past infection

Prevention

Active immunisation through hepatitis B vaccination has been a major breakthrough in the management of this serious illness. There is a course of three injections. If there is a negative antibody response after 3 months, revaccinate with a double dose. If the response is positive, consider a test in 5 years with a view to a booster injection.

For non-immune patients at risk (e.g. after a needle-stick injury), hepatitis B immunoglobin (HBIg), which contains a high level of HBV surface antibody, is appropriate.

Prenatal screening of pregnant women and appropriate use of HBIg and HB vaccine is useful in preventing perinatal vertical transmission of HBV.

HEPATITIS C [10, 12]

Hepatitis C virus is responsible for most cases of viral hepatitis in Australia. It is primarily contracted from intravenous drug use or tattooing. It does not seem to be spread very readily by sexual contact although there is a small risk during heterosexual and homosexual intercourse. It is also not readily spread perinatally.

Clinical symptoms of hepatitis C are usually minimal (often asymptomatic), and the diagnosis is often made after LFTs are found to be abnormal. An important feature is that there are at least six major genotypes of HCV and treatment decisions are based on the genotype.

Hepatitis C infection may be self-limiting, but more commonly (in about 70% of cases) causes a slow, relentless progression to chronic hepatitis, cirrhosis (20%) and also hepatoma.8 A liver biopsy is the most reliable way to assess the severity of hepatitis C. A raised ALT level which is tested three times over the next 6 months implies disease activity. HCV RNA (a PCR test) is present when the ALT becomes abnormal while the anti-HCV rises more slowly and may not be detectable for several weeks.

Diagnosis and progress

This is by serology:

- HCV Ab +ve (anti-HCV) ° exposure (current or past)
- HCV-RNA +ve ° chronic viraemia - ve ° spontaneous clearance
- CD_4/HCV ° viral load
- ALTs on LFTs indicate disease activity (tested 3 times over next 6 months) ALT persistently normal ° good prognosis ALT ↑ ↑ ° requires treatment
- If PCR +ve + significant viral load + ALT ↑ perform HCV genotype - determines treatment

Treatment[5]

The current standard treatment is ribavirin and pegylated alpha-interferon. The combination therapy, which can cure many cases of hepatitis C has considerable side-effects, ranging from flu-like symptoms to depression to significant anaemia. At present the determination of the genotype and the viral load will identify those groups most likely to respond to therapy, for example, genotype I infected patients will have a good response while an excellent response including cure can be expected in genotypes 2 and 3. There is no vaccine yet available. These patients should be tested for hepatitis A and B and immunised if not immune. They should avoid alcohol.

Those at increased risk of having hepatitis B and C

- Blood transfusion recipients (prior to HBV and HCV testing)
- Intravenous drug users (past or present)
- Male homosexuals who have practised unsafe sex
- Kidney dialysis patients
- Sex industry workers
- Those with abnormal LFTs with no obvious cause
- Tattooed people

Prevention of transmission of hepatitis B and C viruses

Advice to those who are positive for HCV:

- Do not donate blood or any body organs.
- Do not share needles.

- Advise health care workers, including your dentist.
- Do not share intimate equipment such as toothbrushes, razors, nail files and nail scissors.
- Wipe up blood spills in the home with household bleach.
- Cover up cuts or wounds with an adequate dressing.
- Dispose of bloodstained tissues, sanitary napkins and other dressings safely.
- Use safe sex practices such as condoms.

HEPATITIS D

Hepatitis D is a small defective virus that lacks a surface coat. This is provided by hepatitis B virus, and so hepatitis D infection occurs only in patients with concomitant hepatitis B.

It is usually spread parenterally and if chronic is usually associated with progressive disease with a poor prognosis. Treatment with interferon has a poor success rate. Antibodies to the delta virus, both anti-HDV and anti-HDV IgM (indicating a recent infection) as well as HDV Ag can be measured.[13]

HEPATITIS E

Hepatitis E is an enterically transmitted virus that occurs in outbreaks in certain countries with a poor water supply, such as some Asian subcontinent countries. Epidemiologically, HEV behaves like HAV, with well-documented water-borne epidemics in areas of poor sanitation. There is a high case fatality rate (up to 20%) in pregnant females.

HEPATITIS F

Researchers claim to have identified HGF virus, which is spread enterically.[14]

HEPATITIS G

HGV has been identified as a transfusion-spread virus. It has subsequently been found to be prevalent among Queensland blood donors.[9, 15]

CHOLESTATIC JAUNDICE

Cholestasis refers to the syndrome of biliary obstructive jaundice whereby there is obstruction to the flow of bile from the hepatocyte to the duodenum, thus causing bilirubin to accumulate in the blood. It is classified into two main groups:

- intrahepatic cholestasis - at the hepatocyte or intrahepatic biliary tree level
- extrahepatic cholestasis - obstruction in the large bile ducts

The significant causes are listed in Table 58.7.

Table 58.7 Significant causes of cholestasis in adults

Intrahepatic
• Viral hepatitis
• Alcoholic hepatitis/cirrhosis
• Drugs
• Primary biliary cirrhosis

Extrahepatic

- Common bile duct gallstones
- Cancer of pancreas
- Cancer of bile ducts
- Other cancer: primary or secondary spread
- Cholangitis
- Primary sclerosing cholangitis (? autoimmune)
- Pancreatitis
- Postsurgical biliary stricture or oedema

Symptoms

- Jaundice (greenish tinge)
- Dark urine and pale stools
- Pruritus - worse on palms and soles
- Pain varies from nil to severe

Gallstones and jaundice

Gallstones can be found in the following (Figure 58.6):


common hepatic duct
stone impacted in cystic duct
or gall bladder neck
• biliary pain or
• acute cholecystitis
stone in common bile duct
• biliary pain
• obstructive jaundice
• cholangitis
asymptomatic
stones or
symptomatic
ampulla of Vater
duodenum

Figure 58.6 Clinical presentation of gallstones

- gall bladder (asymptomatic up to 75%) - the majority remain here
- neck of gall bladder (biliary ‘colic’ or acute cholecystitis)
- cystic duct (biliary ‘colic’ or acute cholecystitis)
- common bile duct - may cause severe biliary ‘colic’, cholestatic jaundice or cholangitis

Acute cholecystitis is accompanied by mild jaundice in 20% of cases, due to accompanying common duct stones.[13]

Common bile duct stones may be asymptomatic or may present with any one or all of the triad of abdominal pain, jaundice and fever. The jaundice varies, depending on the amount of obstruction. The liver is moderately enlarged if the obstruction lasts for more than a few hours.

The investigations of choice for cholestatic jaundice are ultrasound and ERCP.

ACUTE CHOLANGITIS

This is due to bacterial infection of the bile ducts secondary to abnormalities of the bile duct, especially gallstones in the common duct. Other causes are neoplasms and biliary strictures.

Charcot's triad (present in 70%) is shown below:

DxT: fever (often with rigor) + upper abdominal pain + jaundice ° acute cholangitis

Older patients can present with circulatory collapse and Gram-negative septicaemia. Urgent referral is necessary.

CARCINOMA OF HEAD OF THE PANCREAS

Pancreatic carcinoma is the fourth commonest cause of cancer death in the United Kingdom and United States.[13]

Typical clinical features

- M > F
- Mainly > 60 years of age
- Obstructive jaundice
- Pain (over 75%) - epigastric and back
- Enlarged gall bladder (50 - 75%)

Possible features

- Weight loss, malaise, diarrhoea
- Migratory thrombophlebitis
- Palpable hard, fixed mass
- Metastases (e.g. left supraclavicular gland of Virchow)
- Occult blood in stool
- Glycosuria

Diagnosis

- Scanning by ultrasound or CT scan may show mass
- ERCP

Prognosis

Very poor: 5 year survival 1 - 2%.

CIRRHOSIS OF THE LIVER

Cirrhosis is accompanied by jaundice as a late and serious manifestation with the exception of primary biliary cirrhosis, where jaundice appears before advanced liver failure. The development of jaundice usually indicates that there is minimal hepatic reserve and is therefore found in conjunction with other signs of liver failure (Figure 58.7).


alcohol facies
• telangiectasia
• plethora
• rhinophyma
• bloatedness
• angular chelitis
• suffused conjunctiva
fetor, breath
jaundice
hepatomegaly
sparse pubic hair
soft atrophic testes
flapping tremor (asterixis)
leuconychia (white nails)
? mental dysfunction
parotid enlargement
spider naevi
gynaecomastia
splenomegaly (possible)
dilated veins (caput medusae)
ascites (possible)
palmar erythema
Dupuytren's contracture
clubbing
bruising
? peripheral neuropathy
? peripheral oedema

Figure 58.7 Possible features of chronic alcoholic liver disease

Causes

Common:

- Alcohol excess
- Chronic viral hepatitis (esp. HBV, HCV)

Others:

- autoimmune chronic active hepatitis
- primary biliary cirrhosis (autoimmune)
- haemochromatosis
- Wilson's syndrome
- drugs (e.g. methotrexate)
- cryptogenic (no cause found)

Clinical features

- Anorexia, nausea ± vomiting
- Swelling of legs
- Abdominal distension
- Bleeding tendency
- Drowsiness, confusion or coma (if liver failure)

Signs

- Spider naevi (distribution of superior vena cava)

- Palmar erythema of hands
- Peripheral oedema and ascites
- Jaundice (obstructive or hepatocellular)
- Enlarged tender liver (small liver in long-term cirrhosis)
- Ascites
- Gynaecomastia
- ± Splenomegaly (portal hypertension)

Complications

- Ascites
- Portal hypertension and GIT haemorrhage
- Portosystemic encephalopathy
- Hepatoma
- Kidney failure

AUTOIMMUNE CHRONIC ACTIVE HEPATITIS (ACAH)[5]

Also termed idiopathic ACAH, it usually affects young females (10 - 40 years) who present insidiously with progressive fatigue, anorexia and jaundice. Diagnosis is made by abnormal LFTs, positive smooth muscle antibodies, a variety of other autoantibodies and a typical liver biopsy. If untreated most patients die within 3 - 5 years. Treatment is with prednisolone orally, monitored according to serum alanine aminotransferase levels, and supplemented with azathioprine.

ALCOHOLIC LIVER DISEASE

The main effects of alcohol excess on the liver are:

- fatty liver
- alcoholic hepatitis (progresses to cirrhosis if alcohol consumption continues)
- alcoholic cirrhosis

If diagnosed, patients are advised to stop drinking alcohol for life except for fatty liver when small amounts can be drunk later.

Fatty liver

Alcohol can cause hepatic steatosis (fatty liver), which is almost universal in obese alcoholics. Non-alcoholic causes include obesity, diabetes mellitus, hypertriglyceridaemia and corticosteroids. A significant number with this very common condition (one in five Australians) will develop cirrhosis. Fatty liver is usually asymptomatic but some complain of malaise and tiredness. Serology is unhelpful. Diagnosis is by liver biopsy and perhaps CT scan. The treatment is weight loss through diet, which improves liver function and reduces fatty deposits.

SPECIAL PATIENT GROUPS

The returned overseas traveller

The overseas traveller presenting with jaundice may have been infected by any one of the viruses - hepatitis A, B, C, D or E. All are prevalent in developing countries, especially in south-eastern and eastern Asia, some Pacific islands and Africa.

Other causes to consider are malaria, ascending cholangitis and drug-induced hepatic damage due to, for example, the antimalarials, including mefloquine (Lariam) and Fansidar.

The pregnant patient

Important hepatic disorders in pregnancy leading to jaundice are cholestasis of pregnancy, acute fatty liver of pregnancy and severe pre-eclampsia. Refer to Chapter 101.

Postoperative jaundice

There are many possible causes of postoperative jaundice either in the immediate or the long-term postoperative phase. Hypoxia associated with shock in a severely ill patient or in a patient with cardiopulmonary disease may lead to transient abnormalities in liver function. Other causes include:

- post-transfusion hepatitis
- coincident viral hepatitis
- drugs, including anaesthetics
- transfusion overload (haemolysis)
- sepsis
- unmasked chronic liver disease and biliary tract disease
- cholestasis: post major abdominal surgery

Practice tips

- All drugs should be suspected as potential hepatotoxins.
- With hepatitis A the presence of IgM antibodies reflects recent infection, and IgG antibody indicates past infection and lifelong immunity.
- There is no chronic carrier state of hepatitis A and E.
- All patients with jaundice should be tested for hepatitis B surface antigen (HBsAg).
- Hepatitis B infection is usually benign and short-lived, but it can be fatal if chronic hepatitis develops, which may lead later to cirrhosis and hepatocellular carcinoma.
- Up to 5% of patients with hepatitis B will become chronic carriers (especially drug addicts).
- Such carriers are identified by persistent titres of HBsAg and possibly HBeAg, the latter indicating the presence of the whole virus, and active replication and high infectivity.
- A raised gamma glutamyl transferase accompanied by a raised MCV is a good screening test for alcohol abuse.
- A systolic murmur may be heard over the liver in alcoholic hepatitis and hepatoma.
- A distaste for smoking (with jaundice) suggests acute viral hepatitis.

Neonates of HBeAg positive mothers

The neonates should have the following:

- hepatitis B immunoglobulin IM within 24 hours of birth
- hepatitis B vaccine at birth, 1 month and 6 months

This is not 100% effective because some infants can be infected in utero.

WHEN TO REFER

- All patients with fulminant hepatitis
- All patients with chronic liver disease
- Painless obstructive jaundice
- Evidence of malignancy
- Symptomatic gallstones

- Patients with cirrhosis
- Acute fatty liver of pregnancy (very urgent)
- Suspected rare conditions (e.g. Wilson's syndrome)

REFERENCES

1. Kincaid-Smith R, Larkins R, Whelan G. *Problems in Clinical Medicine*. Sydney: McLennan & Petty, 1989: 251.
2. Coffman D, Chalstrey J, Smith-Laing G. *Gastrointestinal Disorders*. Edinburgh: Churchill Livingstone, 1986: 106.
3. Sandler G. Fry J. *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986: 468 - 90.
4. Ryan M, Desmond P. Liver toxicity. Aust Fam Physician, 2001; 30 (5): 427 - 31.
5. Birkett D (Chair). *Therapeutic Guidelines: Gastrointestinal* (Version 3). Melbourne: Therapeutic Guidelines Ltd; 2002: 58.
6. Smart J. *Paediatric Handbook* (6Thedn). Melbourne: Blackwell Science, 2000: 421 - 3.
7. Smith BC, Watson KR. Management of the chronic hepatitis B carrier. Aust Fam Physicians, 1995; 24: 331 - 9.
8. Ruff TA, Gust I. Hepatitis, viral (acute and chronic). In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 226 - 30.
9. Bowden DS, Moaven LD, Locarnini SA. New hepatitis viruses: are there enough letters in the alphabet? Med J Aust, 1996; 164: 87 - 9.
10. Cossart Y. Recent advances in diagnosis and management of viral hepatitis. Common sense pathology. RCPA +Australian Doctor, 2006: 2 - 8.
11. McCaughan G, Levy M. How to treat: Hepatitis B infection. Australian Doctor, 2004; 16 January: 28 - 32.
12. Singal DK, George J. Chronic hepatitis C. Aust Doctor, 2001; 15 February: i - viii.
13. Tierney LM et al. *Current Medical Diagnosis and Treatment* (45Thedn). New York: The McGraw-Hill Companies, 2006: 675 - 82.
14. Deka N, Sharma MD, Mukerjee R. Isolation of the novel agent from human stool that is associated with sporadic human hepatitis. J Virol, 1994; 68: 7810 - 15.
15. Moaven LD et al. Prevalence of hepatitis G virus in Queensland blood donors. Med J Aust, 1996; 165: 369 - 71.

CHAPTER 59. NASAL DISORDERS

The face of Mrs Gamp - the nose in particular - was somewhat red and swollen, and it was difficult to enjoy her society without becoming conscious of the smell of spirits.

CHARLES DICKENS (1812 - 1870), *Martin Chuzzlewit*

NASAL DISORDERS

Disorders of the nose, which include the everyday problems of rhinitis, postnasal drip, epistaxis, folliculitis and disorders of smell, are very common in everyday general practice.

The main functions of the nose are:

- filtration - of dust, organisms and other air borne particles
- olfaction (smell)
- self-cleansing and moisturising of the mucous membrane
- humidification and warming of air in its passage to the lungs
- vocal resonance

The main symptoms of nasal disorders are discharge, blockage, sneezing, anosmia, itching, postnasal drip, bleeding and snoring (Table 59.1). Nasal discharge is a common and important symptom to evaluate. The characteristics of nasal discharge are summarised in Table 59.2.

Table 59.1 Typical symptoms for nasal disorders[2]

Foreign body	Unilateral discharge, unilateral blockage
Acute sinusitis	Facial pain, toothache, nasal discharge, postnasal drip
Allergic rhinitis	Sneezing, rhinorrhoea, itch, eye irritation
Infective rhinitis	Blockage, purulent discharge, postnasal drip
Deviated septum	Blockage, postnasal drip
Nasal polyps	Blockage, reduced smell
Nasal tumour	Blockage, unilateral discharge, epistaxis
Adenoidal hypertrophy	Bilateral blockage, snoring, halitosis
Nasal vestibulitis	Local pain, crusting, malodour

Table 59.2 Characteristics of nasal discharge

Nature of discharge	Think of:
Mucopurulent	bacterial rhinitis, foreign body
Watery/mucoid	viral rhinitis, allergic rhinitis, vasomotor rhinitis, CSF
Blood	neoplasia, trauma, bleeding disorder, rhinitis, infection, hypertension
Serosanguineous	neoplasia, foreign body

DISORDERS OF SMELL

The basic sense of smell is detected in the olfactory region by the olfactory nerve (cranial nerve I) while irritant sensors in the nose, mediated by the maxillary branch of the trigeminal nerve (cranial nerve V), detect some noxious odours. The disorders can be classified as:[1]

- anosmia - no smell

- hyposmia - reduced smell
- hyperosmia - increased sensitivity to odours
- dysosmia - distortion of smell perception
    - cacosmia - normal odours seem foul or unpleasant
    - parosmia - a perverse sense of smell

Disorders of smell can be caused by conductive or sensorineural disturbances or considered as idiopathic (see Table 59.3). Conductive disorders present as anosmia or hyposmia, while sensorineural disorders can present with all of the above disorders.[2]

Table 59.3 Causes of reduced sense of smell

Conductive defects
Head trauma
Septal deviation
Rhinitis and sinusitis
Nasal polyps
Rare (not to be missed)
Nasal tumour
Wegener's granulomatosis
Central/sensorineural defects
Head trauma
Viral infections
Ageing
Cigarette and other smoking/inhalation
Drugs
Chemicals, e.g. benzene, chlorine, formaldehyde, cement dust
Parkinson's disease
Endocrine disorders, e.g. diabetes, hypothyroidism
Nutritional deficiencies
Frontal lobe tumour
Kallman syndrome (anosmia + hypogonadism)

Most cases of idiopathic anosmia are considered to be viral neuropathies and may last from a few days to several months. Head trauma, which can cause conductive or sensorineural disturbances, is considered to be caused either from a fracture of the skull involving the cribriform plate or, more commonly, by posterior head trauma. Some patients will never recover their sense of smell. Patients with anosmia lack flavour discrimination and often have accompanying loss of sense of taste. They are also vulnerable to an awareness of smoke, gas, dangerous chemicals and unhealthy food.

The clinical approach

- History: head injury or surgery, recent URTI, drugs, occupation including chemical exposure
- Physical examination, including inspection via Thudicum nasal speculum
- Snifftest - qualitative and quantitative odours, e.g. coffee, cloves, lemon, peppermint, water placebo. Ammonia (for irritant sensation)
- Investigations, e.g. CT scan for sinus disease, nasal polyps

Treatment [2]

- Explanation and reassurance
- Education about smoke detectors, caution about chemicals including gas, excessive perfume, food safety including milk and meat contamination
- Consider dietary supplement with daily zinc sulphate, vitamin A and thiamine

For chronic anosmia following an URTI:

- prescribe a nasal decongestant such as Spray-Tish Menthol for 5 - 7 days

RHINITIS

Rhinitis is inflammation of the nose causing sneezing, nasal discharge or blockage for more than an hour during the day. Rhinitis is subdivided into various types:

- According to time span:
  - seasonal rhinitis: occurs only during a limited period, usually springtime
  - perennial rhinitis: present throughout the year
- According to pathophysiology:
  - allergic rhinitis: an IgE-mediated atopic disorder
  - vasomotor rhinitis: due to parasympathetic overactivity

Both allergic and vasomotor rhinitis have a strong association with asthma.

The classification can be summarised as:

- seasonal allergic rhinoconjunctivitis ° hay fever
- perennial rhinitis
  - allergic (usually due to house dust mites)
  - non-allergic ° vasomotor
    - eosinophilic
    - non-eosinophilic

NB. allergic rhinitis (hay fever) is presented in detail in Chapter 121.

Clinical features of rhinitis

Nasal symptoms:

- sneezing
- nasal obstruction and congestion
- hypersecretion - watery rhinorrhoea, postnasal drip
- reduced sense of smell
- itching nose (usually allergic)

Throat symptoms:

- dry and sore throat
- itching throat

Irritated eyes (allergic)

Abnormal nasal mucous membrane - pale, boggy, mucoid discharge. A transverse nasal crease indicates nasal allergy, especially in a child.

Allergens

- Pollens from trees (spring) and grass (in summer)
- Moulds
- House dust mites (perennial rhinitis)
- Hair, fur, feathers (from cats, dogs, horses, birds)
- Some foods (e.g. cow's milk, eggs, peanuts, peanut butter)

Diagnosis

Allergic rhinitis - nasal allergy:

- detection of allergen-specific IgE antibodies (not specific)
- RAST test or skin testing for specific allergens (can get false negatives)

Vasomotor rhinitis - a diagnosis of exclusion.

Other causes of rhinitis

- Chronic infection (viral, bacterial, fungal)
- Rhinitis of pregnancy
- Rhinitis medicamentosa - following overuse of OTC decongestant nasal drops or sprays
- Drug-induced rhinitis
    - various antihypertensives
    - aspirin
    - phenothiazines
    - oral contraceptives
    - cocaine, marijuana
- Chemical or environmental irritants (vasomotor rhinitis)
    - smoke and other noxious fumes
    - paints and sprays
    - cosmetics

Factors aggravating rhinitis (vasomotor)

- Emotional upsets
- Fatigue
- Alcohol
- Chilly damp weather
- Air-conditioning
- Sudden changes in temperature and humidity

RHINOSINUSITIS

Acute sinusitis

Acute sinusitis is acute inflammation in the mucous membranes of the paranasal sinuses. About 5% of URTIs are complicated by an acute sinusitis,3 which is mainly viral initially while secondary bacterial infection commonly follows. Any factor that narrows the sinus openings into the nasal cavity (the ostia) will predispose to acute sinusitis.

The two prime clinical presentations are:

1. an URTI persisting for longer than 10 daysu
2. an URTI that is unusually severe with pyrexia and a purulent nasal discharge

Refer to Chapter 52, ‘Pain in the face’ (pp. 569 - 70) for features of acute maxillary sinusitis.

Chronic sinusitis

Chronic sinusitis is the most common complication of acute sinusitis. In chronic sinusitis the symptoms and signs of inflammation persist for more than 8 - 12 weeks and are more likely to be associated with factors that impair drainage via the osteomeatal complex, including nasal polyps.

Treatment

- Amoxycillin 500 mg (o) 8 hourly for 10 - 14 days, possibly for longer periods of 3 - 6 weeks[4]
- Consider decongestant spray, e.g. xylometazoline, for maximum of 5 days and intranasal steroids
- Steam inhalations three times daily (see p. 570)

Refer urgently:

- for surgical drainage if there is no response to the above regimen
- those with orbital or facial cellulitis

NASAL POLYPS

Nasal polyps are round, soft, pale, pedunculated outgrowths arising from the nasal or sinus mucosa. They are basically prolapsed, congested, oedematous mucosa, described by some as ‘bags of water’. They occur in patients with all types of rhinitis, but especially in allergic rhinitis (Figure 59.1). Polyps usually arise from the middle meatus and turbinates.



Figure 59.1 Cross-section of nose, demonstrating origin of nasal polyps

Symptoms include nasal obstruction, watery discharge, postnasal drip and loss of smell.

Note:

- Nasal polyps may be associated with asthma and aspirin sensitivity.
- Cystic fibrosis should be considered in any child with nasal polyps.
- A polyp that does not have the typical smooth pale appearance may be malignant.
- If there is a purulent discharge, swab and give antibiotics.

Treatment

The initial treatment should be medical.[5] A medical ‘polypectomy’ can be achieved with oral steroids, e.g. prednisolone 50 mg daily for 7 days. Supplement this with a corticosteroid spray such as betamethasone, starting simultaneously and continuing for at least 3 months. Give antibiotics for any purulent nasal discharge.

Simple polyps can be readily snared and removed, but referral to a specialist surgeon is advisable for surgical intervention since the aim is to remove the polyp with the mucosa of the sinuses (often ethmoidal cells) from which it arises. This complex procedure reduces the incidence of recurrence.

EPISTAXIS

This common emergency should, in some instances, be treated as a life-threatening problem. The common situation is anterior epistaxis seen in children and the young adult while posterior epistaxis is more common in the older hypertensive patient. It has a strong association with higher URTI (rhinitis, sinusitis), hot dry climates and trauma. Neoplasmas should be kept in mind. Systemic factors include hypertension, atherosclerotic vascular disease, bleeding disorders and the rare hereditary haemorrhagic telangiectasia (Table 59.4). The secret of good management is to have the right equipment, good lighting and effective local anaesthesia.

Table 59.4 Causes of epistaxis

Local causes

- Idiopathic
- Trauma including nose picking
- URTIs
  - common cold
  - influenza
  - sinusitis
- Rhinitis
- Intracranial tumours

Systemic causes

- Cardiovascular disorders
  - arteriosclerosis
  - hypertension
- Systemic febrile infections, e.g. malaria
- Drugs: anticoagulants, aspirin, others
- Toxic agents
- Blood disorders, e.g. leukaemia, thrombocytopenia
- Hereditary haemorrhagic telangiectasia

Ideal equipment

Head light, Thudicum nasal speculum, Tilley nasal packing forceps, suction cannula and tubing, Cophenylcaine forte spray ± 5% cocaine solution.

Tamponade options (for difficult bleeding)

Merocel expandable pack, Kaltostat, BIPP (bismuth iodoform paraffin paste) with ribbon gauze, Foley's catheter (no. 12, 14 or 16) with a 30 mL balloon and self-sealing rubber stopper, anterior/posterior balloon, Epistat catheter with or without Kaltostat.

Treatment options

Simple tamponade:

- Pinch 'soft ' part of nose between thumb and finger for 5 minutes
- Apply ice packs to bridge of nose

Simple cautery of Little's area (Figure 59.3) (under local anaesthetic e.g. Cophenylcaine forte nasal spray ± 5% cocaine solution):

- use one of three methods: electrocautery trichloracetic acid or silver nitrate stick (preferred)


anterior ethmoid artery
Little's area
branch from superior labial
greater palatine artery
septal branch of the sphenopalatine artery

Figure 59.3 Little's area on the nasal septum where several blood vessels anastomose. Bleeding is common here, especially in young people

Persistent anterior bleed

Merocel (surgical sponge) nasal tampon or Kaltostat pack

'Trick of the trade' for a recurrent anterior epistaxis:

- apply topical antibiotic (e.g. Aureomycin ointment) bd or tds for 10 days or (better option) Nasalate nasal cream tds for 7 - 10 days or Rectinol ointment

Severe posterior epistaxis

Use a Foley's catheter or an Epistat catheter.

NASAL VESTIBULITIS

Infection of the nasal vestibule can cause a tender, irritating, crusty problem. Low-grade infections and *folliculitis*, which are evident on inspection, cause localised pain, crusts and bleeding, especially if picked from habit. Treatment is with bacitracin or preferably mupirocin (intranasal) ointment topically for 5 - 7 days.

Furunculosis of the nasal vestibule is usually due to *Staphylococcus aureus*. It starts as a small superficial abscess in the skin or the mucus membrane and may develop into a spreading cellulitis of the tip of the nose. The affected area becomes tender, red and swollen. It is best treated by avoidance of touching, hot soaks and systemic antibiotics such as dicloxacillin or as determined by culture from swabs of the vestibule.

Tip: *Staphylococcus aureus* colonises the nose of 20 - 30% of the population.[6] Carriers are prone to transmit nosocomial infection and have an increased risk of serious infections in the presence of serious medical disorders. Treatment includes strict hygiene and eradication with an agent such as mupirocin ointment (match-head size) 2 - 3 times daily for 5 - 7 days (max. 10 days).[6]

Fissure: Painful fissures often develop at the mucocutaneous junction. They may become crusted and chronic. Fissures can be treated by keeping the area moist with petroleum jelly (Vaseline) or saline gel, using hot compresses and the use of an antibiotic or antiseptic ointment if necessary.

Offensive smell from nose

This may be caused by vestibulitis but ensure no foreign body is present.

Treatment

- Take nasal swab for culture.
- Consider mupirocin 2% nasal ointment, instilled 2 to 3 times a day for 10 days or Kenacomb ointment, instil 2 to 3 times a day.

RHINOPHYMA

This disfiguring swelling of the nose is due to hypertrophy of the nasal sebaceous glands. There is no specific association with alcohol. It is almost exclusive to men over the age of 40 years. Rhinophyma may be associated with rosacea.

Treatment

- Good control of rosacea may reduce the risk.
- If surgical correction is warranted refer to a specialist.
- Carbon dioxide laser therapy is the treatment of choice.
- Shave excision is another effective therapy.

NASAL SEPTAL DEVIATION

This causes blockage as a solitary symptom. Mild septal deviation tends to cause alternating blockage while severe deviation causes persistent blockage on one side.

The septum can be divided into anterior and posterior segments. The anterior portion is necessary to support the cartilaginous pyramid of the nose whereas the posterior portion has no supporting role and can be removed without disturbing the support of the nose. The classic submucous resection operation is therefore suitable for posterior septal deviations. Repair of anterior septal deviations is more complex.

Nasal cosmetic surgery

Rhinoplasty is undertaken to improve the function of an obstructed nasal airway or for cosmetic reasons. In counselling for rhinoplasty it is important to undertake careful planning with realistic anticipated outcomes. The GP should provide non-judgmental support for the patient's decision for cosmetic surgery before referral to an expert in rhinoplasty. Each case has to be assessed individually and the surgery tailored to the deformity. Attention to surgery to the airway is important, otherwise the nose may become partially obstructed and stuffy after cosmetic surgery alone.

SEPTAL PERFORATION

A hole in the nasal septum is caused commonly by chronic infection including tuberculosis, repeated trauma such as vigorous nose ‘picking’ or following nasal surgery. It is a known occupational hazard particularly among chrome workers and is seen in drug users who sniffcocaine. In about 5 - 10% of cases perforation is a result of malignant disease.[3] The condition may be asymptomatic depending on the cause, but there is often an irritating nasal crust and a whistling sound on nasal inspiration. It can be demonstrated by looking in one nares while a light is shone in the opposite one. The cartilaginous part is usually involved.

If not due to a serious cause, treat with Vaseline or saline gel and topical antibiotics for any infection. Refer if malignancy is suspected, otherwise treat symptomatically.

NASAL FRACTURES[7]

Fractures of the nose can occur in isolation or combined with fractures of the maxilla or zygomatic arch. Always check for a compound fracture or head injury and, if present, leave alone and refer. If the patient is seen immediately (such as on a sport's field) with a straight forward lateral displacement, reduction may be attempted ‘on the spot’ with digital manipulation before distortion from soft tissue swelling. This involves simply using the fingers to push laterally on the outside of the nose towards the injured side.[2]

Tips:

- If a deformity is present refer the patient within 7 days, ideally from days 3 - 5.
- Skin lacerations, i.e. compound fracture, usually require early repair.
- The optimal time to reduce a fractured nose is about 10 days after injury.
- Closed reduction under local or general anaesthetic is the preferred treatment.
- Open reduction is more suitable for bilateral fractures with significant septal deviation, bilateral fractures with major dislocations or fractures of the cartilaginous pyramid.

Haematoma of nasal septum

Septal haematoma following injury to the nose can cause total nasal obstruction. It is easily diagnosed as a marked swelling on both sides of the septum when inspected through the nose (Figure 59.2).



Figure 59.2 Inferior view of nasal cavity showing bilateral swelling of septal haematoma

It results from haemorrhage between the two sheets of mucoperiosteum covering the septum. It may be associated with a fracture of the nasal septum.

Note: This is a most serious problem as it can develop into a septal abscess. The infection can pass readily to the orbit or the cavernous sinus through thrombosing veins and may prove fatal, especially in children. Otherwise it may lead to necrosis of nasal septal cartilage followed by collapse and nasal deformity.

Treatment

- Remove blood clot through an incision, under local anaesthetic.
- Prescribe systemic (oral) antibiotics (e.g. penicillin or erythromycin).
- Treat as a compound fracture if X-ray reveals a fracture.

STUFFY, RUNNING NOSE IN ADULTS

For simple post-URTI rhinitis, blow the nose hard into disposable paper tissue or a handkerchief until clear, instil a nasal decongestant for 2 to 3 days and also have steam inhalations with Friar's balsam or menthol preparations.

SENILE RHINORRHOEA

This is a common, distressing problem in the elderly, caused by failure of the vasomotor control of the mucosa. It may be associated with a deviated septum and dryness of the mucosa. There are few physical signs apart from the nasal drip. The treatment is to keep the nasal passages lubricated with an oily based preparation e.g. insufflation with an oily mixture or petroleum jelly. Topical decongestants, cause serious side-effects in the elderly.

CSF rhinorrhoea

Following head injury, clear dripping fluid (+ve for glucose) may indicate a fracture of the roof of the ethmoid. Refer for assessment although spontaneous healing can occur.

NEOPLASIA

Malignant nasal disease, which is uncommon, may cause nasal discharge which may be clear at first, becoming thick and offensive. Malignancy should be suspected in the presence of blood. The growth may be in the nasal fossa, sinus or nasopharynx.

Benign tumours include papilloma, fibroma, osteoma, fibroangioma of puberty and nasal polyps. Fibroangiomas occur exclusively in males between the ages of 9 and 24. Patients present with unilateral nasal obstruction and recurrent epistaxis.

Malignant tumours include nasopharyngeal carcinoma, with the maxillary sinus being the most common site. Squamous cell carcinoma is the most common, followed by adenocarcinoma melanoma and lymphoma. Malignant or non-healing granuloma, sometimes called ‘midline granuloma’, is a slowly progressing ulceration of the face starting in the region of the nose.[3] It may represent a form of T cell malignant lymphoma, which responds to radiotherapy. The differential diagnosis is Wegener's granuloma (see p. 291). Diagnosis is by CT scan and biopsy. Treatment of nasopharyngeal and sinonasal carcinoma depends on the site, size and histology, but usually involves a combination of surgery and postoperative radiotherapy.

NASAL DISORDERS IN CHILDREN

Nasal problems, especially nasal discharge (rhinorrhoea), are very common in children but the pattern of presentation is usually different from that of adults. Sinusitis is uncommon in children under the age of 10 and allergic nasal polyps are relatively rare. If a child presents with polyps consider the possibility of cystic fibrosis or neoplasia. Rhinitis, epistaxis and nasal foreign bodies are common.

Rhinorrhoea

This can be normal or abnormal. There is a ‘nasal cycle’ in which there is nasal congestion and decongestion that alternates from side to side and leads to rhinorrhoea. Other causes of normal discharge include vasomotor reactions to external environmental stimuli such as cold wind and irritants and postnasal drip (2 litres of mucus pass down the back of the nose each day)

Abnormal causes

- Adenoid hypertrophy causing post-nasal space obstruction
- Foreign body in nose - usually unilateral discharge
- Allergic rhinitis
- Unilateral choanal atresia
- Sinusitis (possible but rare)
- Tumour (also rare - consider fibroangioma)

Diagnosis may be enhanced by spraying with a vasoconstricting agent and getting the child to blow the nose. A tumour, foreign body or polyp may become visible.

Coanal atresia

Acute bilateral nasal obstruction may occur in newborns with congenital bilateral choanal atresia. This leads to anterior nasal discharge and to acute respiratory distress. Immediate investigation and relief are essential and a finger in the corner of the mouth can be life-saving as can passing a nasal probe down one nostril and perforating the membrane.

Sinusitis

Although rare, sinusitis can represent a serious emergency. Red flags requiring consideration include a sick child, pyrexia, rapid onset, unilateral and deteriorating airway obstruction.

Blocked nose and snoring

The above causes of nasal blockage may lead to snoring, mouth breathing, reduced sense of smell, dribbling and possibly obstructive sleep apnoea.

Nasal trauma and fractures[8]

Areas of concern associated with nasal fractures, which are uncommon, are possible child abuse, open fracture, septal haematoma or abscess and eye or facial changes. If a fracture is undisplaced the treatment is pain relief, ice compresses and rest. If displaced refer for closed reduction under general anaesthetic within 1 - 2 weeks (ideally at 10 days).[8] If associated epistaxis does not settle with pressure, temporary packing may be required.

Epistaxis

Epistaxis is usually intermittent anterior bleeding from Little's area and may follow trauma including nose picking. Bleeding often occurs at night due to vascular vasodilation. At first, try correction with simple measures (pp. 636 - 7) such as pinching below the nasal septum for 5 minutes, supplemented by cold packs. Vaseline applied in the nose at night tends to prevent bleeding while an antibiotic ointment twice daily for 7 - 10 days may help.

If problematic, refer for an ENT appointment.

Tip: Think of a bleeding disorder or a tumour, e.g. juvenile angiofibroma.

Snoring and obstructive sleep apnoea

In normal children these problems are almost always due to adenotonsillar hypertrophy and most cases are relieved by surgery; CPAP is rarely necessary. Sleep studies are performed to confirm clinical features and allay parental concerns.

The snuffling infant

Snuffling in infants is usually caused by rhinitis due to an intercurrent viral infection. The presence of yellow or green mucus should not usually be cause for concern.

Treatment

Reassure the parents.

- Paracetamol mixture or drops for significant discomfort.
- Get the parents to perform nasal toilet with a salt solution (1 teaspoon of salt dissolved in some boiled water); using a cotton bud, gently clear out nasal secretions every 2 waking hours.
- Once the nose is clean, saline nose drops or spray (e.g. Narium nasal mist) can be instilled.
- Stronger decongestant preparations are not advised unless the obstruction is causing a significant feeding problem, when they can be used for up to 4 - 5 days.

Foreign bodies in the nose

The golden rule is ‘a child with unilateral nasal discharge has a foreign body (FB) until proved otherwise’. Such foreign bodies usually consist of beads, pebbles, peas, pieces of rubber, plastic and paper or other small objects handled by the child. A rhinolith may develop in time on the foreign body. In adults, foreign bodies are often rhinoliths, which are sometimes calcium deposits on pieces of gauze or other material that has been used to pack the nose.

Removal of foreign bodies

Removal of FBs from the nose in children is a relatively urgent procedure because of the risks of aspiration. A disc/button battery such as a hearing aid battery in the nose is a medical emergency requiring urgent removal under anaesthetic.[8]

The nose should be examined using a nasal speculum under good illumination. The tip of the nose should be raised and pressed with the tip of a thumb. At first spray a topical decongestant into the nose and see if the child can blow it out after waiting 10 minutes. Do not attempt to remove FBs from the nose by grasping with ‘ordinary forceps’.

Methods of removal

1. Leverage from behind the FB It is best to pass an instrument behind the FB and pull or lever it forward. Examples of instruments are:
    - a Eustachian catheter
    - a probe to rollout the FB
    - a bent hairpin
    - a bent paperclip
2. Snaring the FB This is the appropriate method for soft irregular FBs such as paper, foam rubber and cottonwool that is clearly visible. Examples of instruments are:
    - a foreign-body remover
    - crocodile forceps
    - fine nasal forceps
3. Glue on a stick Apply Superglue to the plastic end of a swab stick. Apply it to the FB, wait about 1 minute and then gently extract the FB.
4. Rubber catheter suction technique The only equipment required is a straight rubber catheter (large type) and perhaps a suction pump. The method involves cutting the end of the catheter at right angles, smearing the rim of the cut end with petroleum jelly and applying this end to the FB, then providing suction. Oral suction may be applied for a recently placed or ‘clean’ object, but gentle pump suction, if available, is preferred.

5. Irritation of the nose Some practitioners sprinkle white pepper into the nose to induce sneezing.
6. The ‘kiss and blow’ technique This mouth-to-mouth method is used for a cooperative child with a firm, round foreign body such as a bead impacted in the anterior nares. It is best to supervise the child's mother to perform the technique, but the practitioner or practice nurse can perform it

Method

- Use a nasal decongestant spray.
- After 20 minutes lay the child on an examination couch with a pillow.
- Obstruct the normal nostril with a finger from the side.
- Place the mouth over the child's mouth, blowing into it until a slight resistance is felt (this indicates that the glottis is closed).
- Then blow hard with a high velocity puff to cause the FB to ‘pop out’.

To encourage cooperation with the technique the child can be asked to give mother (or other) a ‘kiss’. More than one attempt may be needed but it is usually very successful and avoids the necessity for a general anaesthetic.

REFERENCES

1. Beers MH, Berkin R. *The Merck Manual of Diagnoses and Therapy* (17th edn). New Jersey: Merck Research Laboratories, 1999: 687.
2. Mendelsohn M, Ruhno J. The nose - form and function. Aust Doctor 2004; 2 October: 31.
3. Burton M (ed). *Hall and Coleman's Diseases of the Ear, Nose and Throat* (15th edn). Edinburgh: Churchill Livingstone, 2000; 107 - 17.
4. Dowden J, Moulds R. *Therapeutic Guidelines: Respiratory* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2000: 21 - 3.
5. Lund JL. Diagnosis and treatment of nasal polyps. BMJ, 1995; 311: 1411 - 4.
6. Bochner F (Chair). *Australian Medicines Handbook*. Adelaide: Australian Medicines Handbook Pty. Ltd, 2006: 366.
7. Hansen G. Practice Tips. Aust Fam Physician, 1982; 11: 867.
8. Oates K, Currow K, Hu W. Child Health: *A Practical Manual for General Practice*. Sydney: MacLennan+Petty, 2001: 328 - 30.

CHAPTER 60. NAUSEA AND VOMITING

> Nausea, retching and hypersalivation frequently precede the act of vomiting, which is a highly integrated sequence of involuntary visceral and somatic motor events.
>
> Harrison's Principles of Internal Medicine 1994

NAUSEA AND VOMITING

Vomiting or emesis is a rather dramatic event with a diverse number of causes. It is usually preceded by nausea.

Definitions

Nausea The unpleasant sickly sensation that can herald the onset of vomiting or can be present without vomiting.

Vomiting The forceful expulsion of gastric contents through the mouth.

Regurgitation The expulsion of food in the absence of nausea and without diaphragmatic muscular contractions.

Retching An involuntary act with all the movements of vomiting without the expulsion of gastric contents because the cardiac orifice remains closed.

Haematemesis Vomiting of blood. It is presented on p. 593.

KEY FACTS AND CHECKPOINTS

- Nausea and vomiting have a wide range of potential causes emanating from every body system.
- The common cause of acute nausea and vomiting in most age groups is gastroenteritis.
- The most common causes of vomiting in children are infections - viral (especially) and bacterial - including otitis media and urinary infection.
- Drug ingestion is a common cause of nausea and vomiting; thus, a drug history is vital in assessment.
- Vomiting is commonly associated with migraine and may be the only symptom of a variety of migraine. Children with cyclical vomiting syndrome may have a genetic association with migraine.
- The nature of the vomitus provides a clue:
    - faecalent ° intestinal obstruction
    - blood ° bleeding from oesophagus, stomach or duodenum (mostly)
    - coffee-grounds ° bleeding from stomach or duodenum

THE CLINICAL APPROACH

History

A careful history is essential with an emphasis on drug intake, possible psychogenic factors, including self-induced emesis, weight loss, other GIT symptoms or symptoms suggestive of systemic disease.

Diagnostic guidelines

- Surgical GIT causes are unlikely in the absence of abdominal pain.
- Vomiting without bile-stained vomitus ° pyloric obstruction.
- Vomiting of bile ° obstruction below duodenal ampulla.
- Vomiting of ingested food ° oesophageal obstruction.
- Vomiting without nausea and possibly projectile ° ↑ intracranial pressure.

A summary of the diagnostic strategy model is presented in Table 60.1.

Table 60.1 Vomiting: diagnostic strategy model

Q. Probability diagnosis
A. All ages: acute gastroenteritis motion sickness drugs various infections
Neonates: feeding problems
Children: viral infections/fever otitis media UTI
Adults: gastritis alcohol intoxication pregnancy migraine
Q. Serious disorders not to be missed
A. Bowel obstruction
• oesophageal atresia (neonates)
• pyloric obstruction < 3 months
• intestinal malrotation
• intussusception
• malignancy (e.g. oesophagus, stomach)
Severe infection
• botulinum poisoning
• septicaemia
• meningitis/encephalitis
• infective endocarditis
• others (e.g. acute viral hepatitis)
Malignancy
Intracranial disorders
• malignancy
• cerebellar haemorrhage
Acute appendicitis
Acute pancreatitis
Acute myocardial infarction (e.g. painless)
Q. Pitfalls (mainly adults)
A. Pregnancy (early)
Organic failure
• liver, kidney (uraemia), heart, respiratory
Labyrinthine disorders
• Meniere's syndrome
• labyrinthitis
Poisoning
• food
• chemicals
Gut motility disorders
• achalasia

Paralytic ileus

Substance abuse

Radiation therapy

Hypercalcaemia

Functional obstruction

- diabetic gastroparesis
- idiopathic gastroparesis

Q. Seven masquerades checklist

A. Depression possible

Diabetes ketoacidosis

Drugs

Anaemia -

Thyroid and other endocrine disorders

Spinal dysfunction -

UTI

Q. Is this patient trying to tell me something?

A. Possible: extreme stress (e.g. panic attacks)

Consider bulimia (self-induced vomiting)

Functional (psychogenic)

Examination

If fever is present possible sources of infection (e.g. middle ear, meninges and urinary tract) should be checked.

A careful abdominal examination is appropriate in most instances and this includes urinalysis. Look particularly for scars indicating previous surgery. Look for a succussion splash - this indicates pyloric obstruction.

A neurological examination needs to be considered, including ophthalmoloscopy. Consider raised intracranial pressure.

No examination is complete without assessment of the patient's physical fitness, including the level of hydration, especially in infants and the very old. In these age groups the history may be difficult to obtain and the consequences of fluid loss are more complicated. Always be mindful of the possibility of pregnancy in the female patient.

Investigations

These should consider the underlying cause and also biochemical abnormalities resulting from fluid and electrolyte loss.

The following need to be considered:

- pregnancy test
- microscopy and culture of stools
- radiology of GIT
- oesophageal motility studies
- neurological investigation for suspected intracranial pressure
- drug toxicity studies
- biochemistry

VOMITING IN INFANCY

Is the vomiting bile-stained?

- Green vomiting ° urgent surgical referral for possible intestinal malrotation with volvulus (6 hours leeway before gangrene of bowel)[1]

Other causes: meconium ileus, small bowel atresia

- Non bile-stained vomitus: consider pyloric stenosis, GORD, feeding problems, concealed infection (e.g. UTI, meningitis). Both pyloric stenosis and GORD cause projectile vomiting.

Important warning signs in neonates

- Excessive drooling of frothy secretions from mouth
- Bile-stained vomitus - always abnormal
- Delayed passage of meconium (beyond 24 hours)
- Inguinal hernias

Specific conditions

Oesophageal atresia

- Vomiting occurs with the first feeding.
- There is excessive drooling of frothy secretions from the mouth.
- Pass a 10 French gauge catheter through the mouth to aid diagnosis.

Congenital hypertrophic pyloric stenosis

- Usually sudden onset 3rd - 6th week
- Projectile vomitus
- Failure to thrive
- Male:female ° 5:1
- Gastric peristalsis during test feeding (L → R)
    - feel for pyloric tumour either during test feeding or immediately after vomiting (deep in right epigastrium) - Figure 60.1; once felt, further investigation is not necessary



Figure 60.1 Signs of pyloric stenosis

- biochemistry
    - metabolic alkalosis: sodium usually < 130 mmol/L, chloride < 100 mmol/L
- special investigations (if necessary)

  - barium meal (string sign) - concern about aspiration
  - abdominal ultrasound
- treatment
  - correct fluid and electrolyte deficiency (hypochloraemic alkalosis) before surgery
  - surgical management (longitudinal pyloromyotomy)

Acute gastroenteritis

See pp. 475 and 478.

GASTROPARESIS (ADULTS)[2]

Gastroparesis (gastropathy) or severely delayed gastric emptying is a moderately common condition, which is a cause of nausea and vomiting.

Causes include:

- diabetic gastroparesis
- postsurgical gastroparesis, e.g. vagotomy (complete or partial), fundoplication
- idiopathic

Less common causes include:

- connective tissue disorders, e.g. scleroderma
- vasculitides
- myopathic disorders, e.g. muscular dystrophy
- thyroid dysfunction

Symptoms

- Upper abdominal discomfort
- Early satiety
- Nausea
- Postprandial vomiting (1 - 3 hours after meals)
- Abdominal pain

Diagnosis

- Endoscopy → significant gastric residue
- Barium swallow with follow through
- Nuclear medicine gastric emptying test (gastric retention of 60% after 2 hours is abnormal)

Special problems

- Malnutrition
- Dehydration

Management

- Advise patients to eat small, frequent meals, with careful chewing of food.
- Avoid large pieces of bread especially doughy bread (encourage toasting).
- Avoid fat, especially french fries, raw fruit and vegetables.
- Refer diabetic patients to a dietitian for advice.

Medications[3,4]

- motilium 10 - 20 mg (o) tds, 15 - 30 minutes before meals *or*
- metoclopramide 5 - 10 mg (o) tds, 30 minutes before meals *or*
- erythromycin (has prokinetic properties) 125 mg (o) tds, 15 minutes before meals (rapidly develops tolerance)

Other measures

- Injection of botulinum toxin into the pylorus
- Gastric pacing with internally implanted neurostimulators[5]

SYMPTOMATIC RELIEF OF VOMITING

The first-line management is to ensure that any fluid and electrolyte imbalance is corrected and that any underlying cause is identified and treated. Various antiemetics can give symptomatic relief.

Note: Avoid the use of the dopamine antagonist drugs (e.g. metoclopramide and prochlorperazine) in children because of risk of extrapyramidal side-effects.

Drug-induced nausea and vomiting[3]

- metoclopramide 10 mg (o) or IM 8 hourly prn

For cytotoxic drugs (e.g. cisplatin) and radiotherapy:

- metoclopramide 10 mg (o) or IM 1 to 2 hours prior to therapy then 8 hourly (if mild)

For severe cases:

- ondansetron 8 mg (o) or IV prior to therapy then 2 doses 6 hourly plus
- dexamethasone 8 mg IV 30 minutes prior to therapy, then 2 doses 6 hourly

Note: Restrict ondansetron to 8 mg daily for those with hepatic dysfunction

Motion sickness

- promethazine theoclate 25 mg (o) 60 minutes prior to travel *or*
- dimenhydrinate 50 mg (o) 60 minutes prior to travel *or*
- hyoscine 300 - 600 ìg (o) 30 minutes prior to travel *or*
- hyoscine 1.5 mg dermal disc: applied to dry hairless skin behind the ear 5 - 6 hours before travel (effective for 72 hours)[3]

For treatment: repeat oral presentations 4 - 6 hourly during trip (max. 4 doses in 24 hours)

Vestibular disturbances[3]

The phenothiazine derivatives are the most effective, while the dopamine D_2-receptor antagonists are ineffective.

- prochlorperazine 5 - 10 mg (o) or 10 mg rectally, SC or IM 4 times daily prn *or*

Practice tips

- Consider the possibility of anorexia nervosa and bulimia in adolescent females with a history of vomiting immediately after meals, especially after binge eating.
- If weight loss accompanies nausea and vomiting, consider GIT malignancy and obstruction as well as the above psychogenic conditions.
- Early morning nausea and vomiting can be caused typically by alcohol, pregnancy, kidney failure and raised intracranial pressure.
- Intracranial space-occupying lesions can cause vomiting without associated anorexia or nausea.
- Gastroparesis commonly occurs in longstanding diabetes, following surgery or may be idiopathic. Intense

nausea and anorexia are a feature.

- Antiemetic drug therapy should not be used in infants and children with gastroenteritis.
- Antiemetic treatment must be tailored to the specific cause of the problem.
- The major complications of severe vomiting include trauma of the distal oesophagus, such as a Mallory-Weiss tear, and severe fluid and electrolyte disturbances.

- promethazine theoclate 25 mg (o) or IM 4 hourly prn (max. 100 - 150 mg per 24 hours)

Note: Beware of tardive dyskinesia with prolonged use.

Gastroenteritis

For severe cases in adults:

- metoclopramide 10 mg (o) or IM 8 hourly prn

Pregnancy

- pyridoxine hydrochloride 25 - 50 mg tds if still ineffective add
- metoclopramide 10 mg (o) tds or IM (if oral intolerance)

Postoperative vomiting[3]

- metoclopramide 10 mg IM or IV (slowly), 8 hourly prn *or*
- prochlorperazine 12.5 mg IM, 8 hourly prn

Note: A list of some drugs that can cause nausea and vomiting is presented in Table 60.2.

Table 60.2 Some drugs that can cause nausea and vomiting

• Alcohol (including binge drinking)
• Antibiotics (various) esp. erythromycin
• Antidepressants (e.g. serotonin reuptake inhibitors)
• Antihypertensives
• Bromocriptine
• Codeine
• Corticosteroids
• Cytotoxic agents
• Digoxin
• Levodopa
• Iron preparations
• Nicotine and nicotine gum
• NSAIDs (e.g. indomethacin)
• Opioids (e.g. morphine, codeine)
• Oral contraceptives
• Salicylates
• Theophylline

REFERENCES

1. Smart J. *Paediatric Handbook* (6th edn). Melbourne: Blackwell Science, 2000: 523 - 4.
2. Hebbard G. Gastroparesis. Diabetes Management Journal, 2005; 10: 6 - 7.
3. Birkett D (Chair). *Therapeutic Guidelines: Gastrointestinal* (Version 3). Melbourne: Therapeutic Guidelines Ltd, 2002:33 - 8.
4. Talley NJ. Diabetes gastropathy and prokinetics. Am J Gastroenterology, 2003; 98: 264.
5. Abell T et al. Gastric electrical stimulation for medically refractory gastroparesis. Gastroenterology, 2003: 125 - 421.

CHAPTER 61. NECK LUMPS

There are approximately 800 lymph nodes in the body; no fewer than 300 of them lie in the neck and inflammation of these is exceedingly common.

MCNEILL LOVE (EDITOR OF BAILEY & LOVE) 1965

NECK LUMPS

In the management of lumps in the neck it is important to distinguish between the various midline and lateral causes, especially cervical lymphadenopathy, which may be caused by occult malignancy, such as in the aerodigestive tract. With increasing ageing in the population the number of patients presenting with a malignant neck lump is also increasing. The neck is divided into anterior and posterior triangles by the sternomastoid muscle and the anatomical areas are helpful in identifying the origin of the primary lesion (Figure 61.1).


preauricular
(ear, scalp)
postauricular
(ear, scalp)
preparotid
(parotid, ear)
tonsillar
(tonsils)
occipital
(scalp)
submandibular
(tongue, floor of mouth)
Posterior triangle
submental
(lip, mouth)
anterior cervical (thyroid, larynx)
posterior
cervical
Anterior triangle
supraclavicular
(stomach)
(lung
breast)

Figure 61.1 Lymph glands of the neck including common sources of adenopathy (excluding lymphomas)

KEY FACTS AND CHECKPOINTS

- Most nack lumps are reactire lymph nodes-to concurrent infection
- Lymph nodes are normally palpable in children between 3 and 8 years of age; soft mobile nodes up to 1 cm in diameter are commonly felt in the anterior and posterior triangle. A node > 2 cm is considered to be enlarged. Some cervical glands are very prominent, especially tonsillar nodes.
- These prominent nodes often enlarge during intercurrent viral infection.
- Causes of neck swellings are lymph nodes 85%, goitres 8%, others 7%.[1]
- Solitary nodules in the thyroid move on swallowing.
- Consider the possibility of tuberculosis.
- A knowledge of the areas drained by lymph nodes is important (Figure 61.1).
- Examination must extend beyond the neck for lymphadenopathy.

- To examine cervical glands slightly rotate head and palpate with palmar aspect of fingers.
- Palpate submental area with head slightly flexed.
- Biopsy of a complete lymph node is necessary to establish diagnosis for unknown or suspicious causes but do not consider it as the first step in diagnosis.[2]
- Other investigations are chest X-ray and FBE. Bone marrow biopsy or fine-needle aspiration of thyroid nodule or other masses may be considered. Fine-needle aspiration biopsy (FNAB), which is a relatively simple procedure, is the single most helpful investigation for diagnosing the cause.[3]If cytology reveals malignant squameus cells the primary lesion may lie on the skin, long larynx, pharynx, ear or ocsophapus.

The 20:40 and 80:20 rules[3]

- The age of the patient is a helpful guide, as causes of neck lumps can be roughly categorised by the ‘20:40 rule’:
  - 0 - 20 years: congenital, inflammatory, lymphoma
  - 20 - 40 years: inflammatory, salivary, thyroid, lymphoma
  - > 40 years: lymphoma, metastases
- Most neck lumps (80%) are benign in children while the reverse applies to adults.
- Imaging techniques that may assist diagnosis include axial CT scan (especially in fat necks), MRI scan (distinguishes a malignant swelling from scar tissue or oedema), tomogram of larynx (laryngocele or malignancy), barium swallow (pharyngeal pouch), sialogram and carotid angiogram.

A basic suggested approach for the patient presenting with a neck lump is summarised in Figure 61.2.



Figure 61.2 A basic approach for patients presenting with a next lump[4]

CERVICAL LYMPHADENOPATHY

- There are many causes varying from local infections to lymphoproliferative disorders.
- Most malignant nodes in the supraclavicular area have their primary tumour below the clavicle.
- Eighty-five per cent of malignant nodes in the anterior triangle have their primary tumour in the head and neck.[2]

- Always search for:
  - other nodes at distant sites
  - possible primary source of infections or neoplasia
  - hepatosplenomegaly
- Hodgkin's lymphoma usually presents with rubbery, painless nodes in the neck.
- Most swellings are lateral.

Consistency of enlarged nodes

Rules of thumb are:[5]

- hard: secondary carcinoma
- rubbery: lymphoma
- soft: sarcoidosis or infection
- tender and multiple: infection

Causes of cervical lymph node enlargement (lateral cervical swelling)

Acute cervical lymphadenitis

- Acute viral lymphadenitis
- Acute bacterial lymphadenitis - coccal infection

Chronic lymph node infection

- MAIS lymphadenitis (atypical tuberculosis)
- Tuberculosis
- Viral infection (e.g. EBM (Figure 61.3), rubella, cytomegalovirus, HIV)



Figure 61.3 Cervical lymphadenopathy associated with Epstein-Barr mononucleosis

- Toxoplasma gondii infection
- Lat scratch disease-bartonella henselae infection

Neoplastic lymphadenopathy

- Lymphomas, esp. Hodgkin's lymphoma
- Leukaemia

Metastatic

- Check mouth, pharynx, sinuses, larynx, scalp, oesophagus, stomach, breast, lungs, thyroid, skin. A working rule is upper neck - from skin to upper aerodigestive tract; lower neck - from below clavicles (e.g. lung, stomach, breast, colon).
- Examples:
    - occipital or preauricular - check scalp
    - submental - check mouth, tongue, teeth
    - submandibular - check floor of mouth
    - left supraclavicular (under sternomastoid) - consider stomach (Troisier's sign)
    - deep anterior cervical - consider larynx, thyroid, oesophagus, lungs

NON-LYMPH NODE SWELLINGS

Types and causes

Widespread

- Sebaceous cysts
- Lipomas

Midline

- Thyroid nodule (moves upon swallowing)
- Thyroglossal cysts (move up on tongue protrusion)
- Dermoid cyst (beneath chin)
- Midline cervical lymph node swelling

Anterior triangle

- Branchial cyst (in upper part)
    - usually adulthood (20 - 25 years)
- Carotid body tumour
    - opposite thyroid cartilage
    - smooth and pulsatile
    - can be moved laterally but not vertically
    - usually 40 - 60 years
    - requires excision (with care)
- Carotid aneurysm
- Lateral thyroid tumours

Posterior triangle

- Developmental remnants
    - cystic hygroma
    - bronchial sinuses and cysts
- Pancoast tumour (from apex lung)
- Cervical rib

Submandibular swellings

- Submandibular salivary gland

- Cervicofacial actinomycosis (lumpy jaw syndrome)
    - chronic granulomatosis infection due to Gram-positive Actinomyces israeli
    - forms a multilocular abscess (pus has 'sulphur granules')
    - infection follows dental extraction or poor dental hygiene, esp. severe caries
    - treat with high-dose penicillin G, 4 months

Sternomastoid tumour

Refer p. 891.

Pharyngeal pouch

- A soft, squelchy indefinite mass
- Base of left neck
- History of difficulty in swallowing

Thyroid nodule

The most likely cause of a solitary thyroid nodule is the dominant nodule in a multinodular goitre.

Other causes include a true solitary nodule-adenoma, follicular carcinoma or solitary corcinoma-and a colloid cyst. Malignancy must be excluded

Investigations

- Ultrasound
- FNAB (may resolvc a cystic lesion)
- TFTs

NECK LUMPS IN CHILDREN

Eighty per cent of neck lumps are benign while 20% are malignant. Benign lumps usually occur in the anterior triangle, while malignant lumps are more likely in the posterior triangle. The common midline lump in children is the thyroglossal cyst.[3]

Lymphadenopathy

- Most enlarged lymph nodes are either 'normal' or local infections (mainly viral), especially if < 2 cm diameter, and not hard or fixed.
- Inflammatory nodes may be caused by infection in the tonsils, the teeth or other oral or nasopharyngeal cavities.
- They are of concern if supraclavicular node enlargement and fever < 1 week.
- Suspicious nodes are > 2.5 cm, with firmer consistency than normal and less mobility (investigate especially with biopsy).

MAIS lymphadenitis[4,6]

- Child usually 2 - 3 years of age
- Caused by Mycobacterium avium intracellulare, scrofulaceum (MAIS) infection
- Produces chronic cervical lymphadenitis and callar stud abscesses
- A relatively common infection of cervical nodes yet often unrecognised
- Painless swelling due to development of a cold abscess in healthy child
- Nodes enlarge over 4 - 6 weeks prior to erupting into a 'cold' abscess
- Common sites are submandibular, tonsillar and preauricular nodes
- Invariably unilateral, confined to one lymph node group
- No pulmonary involvement

- Unresponsive to antimicrobials: treatment is by surgical excision of abscess and underlying lymph nodes

Acute bacterial lymphadenitis

- Usually coccal infections - Staphylococcus, Streptococcus
- Can progress to abscess formation (fluctuant): requires drainage

WHEN TO REFER

- A persisting lump, depending on its location and size
- A lymph node or group of nodes that are abnormally enlarged and fail to respond to antibiotics

Red flags for neck lumps

- > 40 years, esp. > 70 years
- Nodes > 2.5 cm
- Nodes > 3 - 4 ? malignancy
- Tender mass
- Purple discolouration (collar-stud abscess)
- Single, gradually enlarging node
- Fixed to skin without punctum
- Associated dysphagia
- Hard midline thyroid lump
- Patient at risk of malignancy and HIV

REFERENCES

1. Fry J, Berry H. Surgical Problems in Clinical Practice. London: Edward Arnold, 1987: 38.
2. Coman WB. Neck lumps in adults. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 340 - 1.
3. Cole IE, Turner J. Neck lumps: clues to the diagnosis and management. Mod Med Aust, 1997; April: 37 - 55.
4. Hughes C, O'Brien C. How to treat: Neck lumps. Australian Doctor, 2005; 5 August: 31-38
5. Larkins R, Smallwood R. Clinical Skills. Melbourne: Melbourne University Press, 1993: 133 - 4.
6. Stokes K. Lumps in the Neck in Children. Proceedings notes. Box Hill Hospital Seminar, 1995: 1 - 2.

CHAPTER 62. NECK PAIN

We have all heard of the courtiers who mimicked the wry neck of Alexander the Great.

WILLIAM HEBERDEN (1710 - 1801)

NECK PAIN

Neck pain is a very common symptom in both sexes at all ages and although most pain is experienced in the posterior aspect of the neck, anterior neck pain can occur from causes that overlap between front and back. The main cause of neck pain is a disorder of the cervical spine, which usually manifests as neck pain but can refer pain to the head, shoulders and chest. Such pain usually originates from the facet (apophyseal) joints but can arise from other musculoskeletal structures, such as the intervertebral discs and the muscles or ligaments (Figure 62.1). The other major symptom is limited movement or stiffness.


spinal cord
facet joint
posterior primary ramus
nerve root
sinuvertebral nerve
vertebral artery
sympathetic chain
nucleus pulposus
annulus fibrosus
intervertebral disc

Figure 62.1 Transverse section illustrating the functional unit and nervous network of the cervical spine

General causes of neck pain are presented in Table 62.1.

Table 62.1 Causes of neck pain (a pathological classification)

Musculoskeletal
Joint dysfunction
• apophyseal
• intervertebral disc
Muscular/ligamentous strains or sprains
Trauma
• 'whiplash'
• fracture
• other disorders
Inflammation
• osteoarthritis *
• rheumatoid arthritis
• ankylosing spondylitis
• psoriasis
• inflammatory bowel disorders

• Reiter's syndrome/reactive arthritis • polymyalgia rheumatica • thyroiditis
Infective
Spinal • osteomyelitis • tuberculosis • herpes zoster Extraspinal • cervical adenitis • poliomyelitis • tetanus Extracervical • meningitis • febrile states: - meningism - malaria
Degenerative
• spondylosis*
Neoplasia
• benign • malignant
Fibromyalgia syndrome
Psychogenic
Referred visceral
• heart - IHD - pericarditis • oesophagus • carcinoma lung
Referred cranial
• haemorrhage (e.g. subarachnoid) • tumour • abscess

* Osteoarthritis, or spondylosis, is inflammatory and degenerative.

KEY FACTS AND CHECKPOINTS

- At any time approximately 10% of the adult population are experiencing[1] an episode of neck pain.

- The commonest cause of neck pain is idiopathic dysfunction of the facet joints without a history of injury.
- Disorders of the intervertebral discs are common, especially in the lower cervical spine, and may cause unilateral pain, paraesthesia or anaesthesia in the arm.
- In a UK study radiological cervical disc degeneration was present in 40% of males and 28% of females[1] between 55 and 64 years.
- Strains, sprains and fractures of the facet joints, especially after a ‘whiplash’ injury, are difficult to detect and are often overlooked as a cause of persistent neck pain.
- Cervical spondylosis is a disorder of ageing: radiological signs occur in 50% of people over the age of 50 and in 75% over the age of 65 years.[2]
- In cervical spondylosis, osteophytic projections may produce nerve root and spinal cord compression, resulting in radiculopathy and myelopathy respectively.
- Radiculopathy can be caused by a soft disc protrusion (usually unilateral), a hard calcified lump and osteophytes (may be bilateral).
- Cervical disorders are aggravated by vibration (e.g. riding in a motor vehicle).
- Always determine the C2, C6 and C7 levels by finding the relevant spinous processes (easily palpable landmarks) prior to palpation.
- Palpation of the neck is the cornerstone of cervical management. Palpate gently - the more one presses the less one feels.
- Most episodes of neck pain, including acute torticollis, are transient, lasting from about 2 to 10 days.
- In one study 70% of people with neck pain who sought medical attention had recovered or were recovering within one month.[1]
- Effective management of neck pain is based on the theoretical principle that stiff dysfunctional joints are painful and restoration of normal movement may be associated with resolution of pain.
- The optimal treatment for dysfunctional joints (without organic disease or radiculopathy) is active and passive mobilisation, especially as exercises.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 62.2.

Table 62.2 Neck pain: diagnostic strategy model

Q. Probability diagnosis
A. Vertebral dysfunction Traumatic ‘strain’ or ‘sprain’ Cervical spondylosis
Q. Serious disorders not to be missed
A. Cardiovascular • angina • subarachnoid haemorrhage • arterial dissection Neoplasia • primary • metastasis • Pancoast tumour Severe infections • osteomyelitis • meningitis Vertebral fractures or dislocation

Q. Pitfalls (often missed)

A. Disc prolapse
Myelopathy
Cervical lymphadenitis
Fibromyalgia syndrome
Outlet compression syndrome (e.g. cervical rib)
Polymyalgia rheumatica
Ankylosing spondylitis
Rheumatoid arthritis
Oesophageal foreign bodies and tumours
Paget's disease

Q. Seven masquerades checklist

A. Depression
Diabetes -
Drugs -
Anaemia -
Thyroid disorder Thyroiditis
Spinal dysfunction
UTI -

Q. Is the patient trying to tell me something?

A. Highly probable. Stress and adverse occupational factors relevant.

Probability diagnosis

The main causes of neck pain are vertebral dysfunction, especially of the facet joints, and traumatic strains or sprains affecting the musculoligamentous structures of the neck. The so-called myofascial syndrome is mainly a manifestation of dysfunction of the facet joints. Acute wry neck (torticollis), which is quite common, is yet another likely manifestation of apophyseal joint dysfunction. Spondylosis, known also as degenerative osteoarthrosis and osteoarthritis, is also a common cause, especially in the elderly patient.

Intervertebral disc disruption is also a relatively common phenomenon in the cervical spine, especially at the lower levels C5 - 6 and C6 - 7.[3]

Serious disorders not to be missed

Conditions causing neck pain and stiffness may be a sign of meningitis or of cerebral haemorrhage, particularly subarachnoid haemorrhage, or of a cerebral tumour or retropharyngeal abscess.

Angina and myocardial infarction should be considered in anterior neck pain. Other visceral disorders can refer pain to the neck.

Arterial dissection of the internal carotid artery or vertebral artery should be kept in mind in patients presenting with acute neck pain, especially without musculoskeletal symptoms or signs.

Tumours are relatively rare in the cervical spine but metastases do occur and should be kept in mind, especially with persistent neck pain present day and night.

Metastasis to the spine occurs in 5 - 10% of patients with systemic cancer, making it the second most common neurological complication of cancer. The cervical spine accounts for some 15% of spinal metastases.[2]

The commonest primary tumours are the breast, prostate or lung. Other primaries include the kidney, thyroid and melanoma.

Pitfalls

There are many pitfalls in the clinical assessment of causes of neck pain and many of them are inflammatory.

Rheumatoid arthritis is the prime severe inflammatory arthropathy that involves the neck but the neck can be affected by the seronegative spondyloarthropathies, particularly ankylosing spondylitis, psoriasis and the inflammatory bowel disorders.

While polymyalgia rheumatica affects mainly the shoulder girdle, pain in the lower neck, which is part of the symptom complex, is often overlooked. Diffuse neck pain in myofascial soft tissue with tender trigger areas is part of the uncommon but refractory fibro-myalgia syndrome.

General pitfalls

- Failing to appreciate how often the benign problem of facet joint dysfunction occurs in the neck, causing pain and limited movement. This involves failure to appreciate the value of physical therapy, especially exercise programs, in alleviating the problem.
- Failing to adhere to the idiom: one disc - one nerve root. Involvement of more than one nerve root in the upper limb may mean a neoplastic disorder such as metastatic disease, lymphoma in the thoracic outlet and similar serious diseases.
- Missing the insidious onset of myelopathy, especially the spasticity component, caused by rheumatoid arthritis, osteophytic overgrowth or, rarely, a soft disc prolapse.

Seven masquerades checklist

Cervical spinal dysfunction is the obvious outstanding cause. Thyroiditis may cause neck pain, as in the extremely rare cases of acute specific infection in the thyroid (e.g. syphilis, pyogenic infections), which cause severe pain; non-specific thyroiditis (de Quervain's thyroiditis) produces painful swelling with dysphagia. The association between depression and neck pain is well documented.

Psychogenic considerations

The neck is one of the commonest areas for psychological fixation following injury. This may involve perpetuation or exaggeration of pain because of factors such as anxiety and depression, conversion reaction and secondary gain.

The psychological sequelae that can follow a whiplash injury and chronic neck problems such as spondylosis serve as a reminder that the state of the patient's cervical spine can profoundly affect his or her life and that we should always be aware of the whole person. A feeling of depression is a very common sequel to such an injury and these patients demand our dutiful care and understanding.

THE CLINICAL APPROACH

History

It is important to analyse the pain into its various components, especially the nature of its onset, its site and radiation, and associated features. The diurnal pattern of the pain will provide a lead to the diagnosis (refer to Figure 37.3: the patterns are similar to low back pain).

Key questions

- Can you point to exactly where in your neck you get the pain?

- Do you wake up with pain in the morning?
- Does the pain come on when you have to look up for a while?
- Do you have trouble reversing your car?
- Can you recall an injury to your head or neck such as hitting your head on an overhead bar?
- Does your neck grate or get stiff?
- Do you get headaches or feel dizzy?
- Is the pain present day and night?
- Do you get pain or pins and needles or numbness in your arms?
- Does the pain come on with activity?
- Does the pain wake you at night?
- Do you feel pain on both sides of your neck and over your shoulders?
- Do your hands or arms feel weak or clumsy?

Physical examination

It is appropriate to follow the traditional rule for examination of any joint or complex of joints: look, feel, move, measure, test function and x-ray. Careful examination of the cervical spine is essential for the correct diagnosis and for specific treatment at the painful level.

Three objectives of the examination are to:

- reproduce the patient's symptoms
- identify the level of lesion or lesions
- determine the cause (if possible)

A neurological examination is essential if radicular pain is present, or weakness or other upper limb symptoms, including any pain or paraesthesia that extends below the elbow.

Inspection

The patient should be examined sitting on a couch, rather than on a chair. The body should be fully supported with the hands resting on the thighs. The following should be noted:

- willingness to move the head and neck
- level of the shoulders
- any lateral flexion
- contour of the neck from the side

In the patient with torticollis the head is held laterally flexed with, perhaps, slight rotation to one side - usually away from the painful side. Patients suffering from whiplash injury and severe spondylosis tend to hold the neck stiff and the head forward, and tend to turn the trunk rather than rotate the neck.

Palpation

For this vital component of the examination it is essential to know the surface anatomy of the neck so that the affected level can be determined.

Method

The patient lies prone on the examination couch with the forehead resting on the hands (palms up). The neck should be flexed forward and the shoulders relaxed.

1. Central digital palpation

Systematically palpate the first spinous processes of the cervical vertebrae.

- C2 (axis) is the first spinous process palpable beneath the occiput.
- C7 is the largest ‘fixed’ and most prominent process - situated at the base of the neck.

- C6 is also prominent but usually ‘disappears’ under the palpating finger with extension of the neck.
- The spinous processes of C3, C4 and C5 are difficult to palpate because of cervical lordosis but their level can be estimated (Figure 62.2).



Figure 62.2 Relative sizes of spinous processes of the cervical spine

Standing at the patient's head, place opposed pulps of the thumbs on the spinous processes (starting at C2) and then move down the middle line to C7. Press firmly over each and with arms straight oscillate with moderate firmness three or four times to assess pain, stiffness or muscle spasm.

2 Lateral digital palpation

The facet joints lie in sequence (called the articular pillar) about 2 to 3 centimetres from the midline. Press with opposed thumbs against this pillar in a systematic manner on either side of the midline (top to base) to determine any painful area.

Palpation should be extended to include the anterior neck, searching for evidence of lymphadenitis, muscle spasm, thyroid disorder and other problems.

Movement

Active movements are observed with the patient sitting on the couch. The movements are as follows with normal range indicated:

- flexion - 45°
- extension - 50°
- lateral flexion (R and L) - 45°
- rotation (R and L) - 75°

If there is a full range of pain-free movement, apply overpressure slowly at the end range and note any pain.

The range of movements can be plotted on a special grid called a direction of movement (DOM) diagram (Figure 62.3). This provides a ready reference for serial assessments.



Figure 62.3 Direction of movement diagram to record movements of the neck. This record shows restricted and painful movements (indicated by ll) in right lateral flexion and right rotation; the other movements are free

Neurological examination

A neurological examination for nerve root lesions (C5 to T1) is indicated if the clinical assessment identifies the presence of neurological symptoms and signs such as pain, paraesthesia or anaesthesia in the arm. Nerve root pressure is indicated by:

- pain and paraesthesia along the distribution of the dermatome
- localised sensory loss
- reduced muscular power (weakness or fatigue or both)
- hyporeflexia (reduced amplitude or fatigue or both)

It is necessary to know the sensory distribution for each nerve root and the motor changes. This is summarised in Table 62.3. The dermatomes are illustrated in Figure 62.4.

Table 62.3 Cervical nerve root syndromes

Nerve root	Sensory change	Muscle power	Power loss	Reflex
C5	Outer arm	Deltoid	Abduction arm	Biceps jerk
C6	Outer forearm/thumb/index finger	Biceps	Elbow flexion Extension wrist	Biceps + brachioradialis
C7	Hand/middle and ring fingers	Triceps	Elbow extension	Triceps
C8	Inner forearm/little finger	Long flexors finger, long extensors thumb	Grip	
T1	Inner arm	Interossei	Finger spread	



Figure 62.4 Dermatomes of the upper limb, head and neck

Investigations

The investigations are directed to diagnosing the painful condition and determining if suspected or true organic disease is present in the spine. It is inappropriate to perform sophisticated investigations such as CT scans on most patients. Scanning should be reserved where surgery is contemplated and serious disease is suspected but not confirmed by plain X-ray.

Investigations include:

- haemoglobin, film and WCC
- ESR
- rheumatoid arthritis factor
- HLA-B_{27} antigen
- radiology
  - plain X-ray
  - plain CT scan
  - CT scan and myelogram (if cervical disc surgery contemplated)
  - radionuclide bone scan (for suspected metastatic disease)
  - MRI: the investigation of choice for cervical radiculopathy, myelopathy, suspected spinal infection and tumour

These should be selected conservatively.

NECK PAIN IN CHILDREN

In children and adolescents neck pain, often with stiffness, may be a manifestation of infection or inflammation of cervical lymph nodes, usually secondary to an infected throat - for example, tonsillitis or pharyngitis. However, it is vital to consider the possibility of meningitis. Sometimes a high fever associated with a systemic infection or pneumonia can cause meningism. In the presence of fever the rare possibility of poliomyelitis should be kept in mind. In both children and adults the presence of cerebral pathology, such as haemorrhage, abscess or tumour are un-common possibilities.[4] Acute torticollis is quite common in this age group and the neck may be involved in chronic juvenile arthritis.

NECK PAIN IN THE ELDERLY

In adults the outstanding causes are dysfunction of the joints and spondylosis, with the acute febrile causes encountered in children being rare. However, cerebral and meningeal disorders may cause pain and stiffness in the neck.[4]

Rheumatoid arthritis is the prime severe inflammatory arthropathy that involves the neck, but the neck can be affected by the spondyloarthropathies (e.g. ankylosing spon-dylitis). The painful acute wry neck can affect all ages and is considered to be caused mainly by acute disorders of the apophyseal joints rather than disc prolapse. However, disc lesions do occur and can cause referred pain or radicular pain. In the elderly radicular pain can be caused also by impingement of the nerve root in the intervertebral foramen that has become narrowed from the degenerative changes of longstanding spondylosis.

Problems with a higher probability with increasing age include:

- cervical spondylosis with radiculopathy or myelopathy
- atlantoaxial subluxation complicating rheumatoid arthritis
- polymyalgia rheumatica
- metastatic cancer
- Pancoast tumour of the lung
- angina and myocardial infarction
- pharyngeal and retropharyngeal infection and tumour

CLINICAL PROBLEMS OF CERVICAL SPINAL ORIGIN

Pain originating from disorders of the cervical spine is usually, although not always, experienced in the neck. The patient may experience headache, or pain around the ear, face, arm, shoulder, upper anterior or posterior chest.[5]

Possible symptoms:

- neck pain
- neck stiffness
- headache
- ‘migraine’-like headache
- facial pain
- arm pain (referred or radicular)
- myelopathy (sensory and motor changes in arms and legs)
- ipsilateral sensory changes of scalp
- ear pain (periauricular)
- scapular pain
- anterior chest pain
- torticollis
- dizziness/vertigo
- visual dysfunction

Figure 62.5 indicates typical directions of referred pain from the cervical spine. Pain in the arm (brachialgia) is common and tends to cover the shoulder and upper arm as indicated.



Figure 62.5 Possible directions of referred pain from the cervical spine

CERVICAL DYSFUNCTION

Dysfunction of the 35 intervertebral joints that comprise the cervical spine complex is responsible for most cases of neck pain. The problem can occur at all ages and appears to be caused by disorder (including malalignment) of the many facet joints, which are pain-sensitive. Dysfunction of these joints, which may also be secondary to intervertebral disc disruption, initiates a reflex response of adjacent muscle spasm and myofascial tenderness.

Acute neck pain (ANP)

ANP is most commonly idiopathic or due to a whiplash accident. Serious cases are Mare.[6] Dysfunction can follow obvious trauma such as a blow to the head or a sharp jerk to the neck, but can be caused by repeated trivial trauma or activity such as painting a ceiling or gentle wrestling. People often wake up with severe neck pain and blame it on a 'chill' from a draught on the neck during the night. This is incorrect because it is usually caused by an unusual twist on the flexed neck for a long period during sleep.

Typical clinical features:

- typical age range 12 - 50 years
- dull ache (may be sharp) in neck
- may radiate to occiput, ear, face and temporal area (upper cervical)
- may radiate to shoulder region, especially suprascapular area (lower cervical)
- rarely refers pain below the level of the shoulder
- pain aggravated by activity, improved with rest
- various degrees of stiffness
- neck tends to lock with specific movements, usually rotation
- localised unilateral tenderness over affected joints
- variable restriction of movement but may be normal
- X-rays usually normal: Plain X-rays are not indicated for the investigation of ANP in the absence of 'red flags' and a history of trauma[6]

Management

The aim of treatment is to reduce pain, maintain function and minimise the risk of chronicity.

- Provide appropriate reassurance, information and support.
- Give advice to the patient about rules of living including the following:
  Do:
  - Stay active and resume normal activities.
  - Keep your neck upright in a vertical position for reading, typing and so on.
  - Keep a good posture - keep the chin tucked in.
  - Sleep on a low firm pillow or a special conforming pillow.
  - Sleep with your painful side on the pillow.
  - Use heat and massage: massage your neck firmly three times a day using an analgesic ointment.

  Don't:
  - Look up in a strained position for long periods.
  - Twist your head often towards the painful side (e.g. when reversing a car).
  - Lift or tug with your neck bent forwards.
  - Work, read or study with your neck bent for long periods.
  - Become too dependent on ‘collars’.
  - Sleep on too many pillows.
- Monitor the patient's progress without overtreatment.
- Use basic analgesics (e.g. paracetamol).
- Prescribe an exercise program as early as possible; start with gentle exercises and maintain them at home.
- Refer to an appropriate therapist for cervical mobilisation for persisting pain. Mobilisation combined with exercises is very effective treatment. Occasionally, manipulation may help with a stubborn ‘locked’ neck but should be left to an expert. If manipulation, which carries the rare but real risk of vertebral artery dissection and stroke, is to be performed, informed consent is required.[7]

Evidence of benefit (in summary)[7]

- Staying active: resuming normal activities
- Exercises
- Combined cervical panic mobilisation/exercises
- Pulsed electromagnetic therapy (up to 12 weeks)

Approximately 40% patients recover fully from acute idiopathic ANP, about 30% continue to have wild symptoms include 30% continue to have moderate or several symptoms.[6]

Chronic pain

Additional treatment modalities to consider include:

- a course of antidepressants
- TENS, especially when drugs are not tolerated
- hydrotherapy
- acupuncture
- corticosteroid facet injections (ideally under image intensification)
- facet joint denervation

CERVICAL SPONDYLOSIS

Cervical spondylosis following disc de-generation and apophyseal joint degeneration is far more common than lumbar spondylosis and mainly involves the C5 - 6 and C6 - 7 segments. The consequence is narrowing of the intervertebral foramen with the nerve roots of C6 and C7 being at risk of compression.

Cervical spondylosis is generally a chronic problem but it may be asymptomatic. In some patients the pain may lessen with age, while stiffness increases.

The main clinical features are as follows:

- dull, aching suboccipital neck pain (Figure 62.6)



Figure 62.6 Cervical spondylosis: typical pain distribution with DOM diagram indicating painful and restricted movements

- stiffness
- worse in morning on arising and lifting head
- improves with gentle activity and warmth (e.g. warm showers)
- deteriorates with heavy activity (e.g. working under car, painting ceiling)
- usually unilateral pain - may be bilateral
- pain may be referred to head, arms and scapulae
- may wake patient at night with paraesthesia in arms
- C6 nerve root most commonly involved
- acute attacks on chronic background
- aggravated by flexion (reading) and extension
- associated vertigo or unsteadiness
- restricted tender movements, especially rotation/lateral flexion
- joints tender to palpation
- X-ray changes invariable

Treatment

- Provide appropriate reassurance, information and support.
- Refer for physiotherapy, including warm hydrotherapy.
- Use regular mild analgesics (e.g. paracetamol).
- Use NSAIDs: a trial for 2 weeks and then review.

- Prescribe gentle mobilising exercises as early as possible.
- Give passive mobilising techniques.
- Outline general rules to live by, including advice regarding sleeping and pillows, and day-to-day activities.

Complications

- Radiculopathy (unilateral or bilateral)
- Myelopathy - pressure on spinal cord

ACUTE TORTICOLLIS

Torticollis (acute wry neck) means a lateral deformity of the neck. This is usually a transient self-limiting acutely painful disorder with associated muscle spasm of variable intensity.

The typical features include:

- age of patient between 12 and 30 years
- patient usually awakes with the problem
- pain usually confined to neck but may radiate
- deformity of lateral flexion and slight flexion/rotation
- deformity usually away from the painful side
- loss of extension
- mid-cervical spine (C2 - 3, C3 - 4, C4 - 5)
- any segment between C2 and C7 can cause torticollis
- usually no neurological symptoms or signs

The exact cause of this condition is uncertain, but both an acute disc lesion and apophyseal joint lesion are implicated, with the latter the more likely cause. Acute torticollis is usually a transient and self-limiting condition that can recover within 48 hours. Sometimes it can last for about a week. Encourage heat manage and early mobility. Avoid cervical collars. Management by mobilisation and muscle energy therapy is very effective.

Muscle energy therapy

This amazingly effective therapy relies on the basic physiological principle that the contracting and stretching of muscles leads to automatic relaxation of agonist and antagonist muscles.[8] Lateral flexion or rotation or a combination of movements can be used but treatment in rotation is preferred. The direction of contraction can be away from the painful side (preferred) or towards the painful side, whichever is most comfortable for the patient.

Method

1. Explain the method to the patient, with reassurance that it is not painful.
2. Rotate the patient's head passively and gently towards the painful side to the limit of pain (the motion barrier).
3. Place your hand against the head on the side opposite the painful one. The other (free) hand can be used to steady the painful level - usually C3 - 4.
4. Request the patient to push the head (in rotation) as firmly as possible against the resistance of your hand. The patient should therefore be producing a strong isometric contraction of the neck in rotation away from the painful side (Figure 62.7a). Your counterforce (towards the painful side) should be firm and moderate (never forceful) and should not ‘break’ through the patient's resistance.



Figure 62.7 Muscle energy therapy for acute torticollis: (a) isometric contraction phase for problem on the left side; (b) relaxation phase towards the affected (left) side

5. After 5 - 10 seconds (average 7 seconds) ask the patient to relax; then passively stretch the neck gently towards the patient's painful side (Figure 62.7b).
6. The patient will now be able to turn the head a little further towards the painful side.
7. This sequence is repeated at the new improved motion barrier. Repeat three to five times until the full range of movement returns.
8. Ask the patient to return the following day for treatment although the neck may be almost normal.

The patient can be taught self-treatment at home using this method.

ACCELERATION HYPEREXTENSION WHIPLASH INJURY

Patients with the whiplash syndrome preferably referred to as an acceleration hyperextension injury, present typically with varying degrees of pain-related loss of mobility of the cervical spine, headache and emotional disturbance in the form of anxiety and depression. The problem can vary from mild temporary disability to a severe and protracted course.

The injury occurs as a consequence of hyperextension of the neck followed by recoil hyperflexion, typically following a rear-end collision between motor vehicles. There is reversal of sequence of these movements in a head-on collision. In addition to hyper-extension, there is prolongation or anterior stretching plus longitudinal extension of the neck.[7] It can also occur with other vehicle accidents and in contact sports such as football.

Whiplash causes injury to soft tissue structures including muscle, nerve roots, the cervical sympathetic chain, ligaments, apophyseal joints and their synovial capsules and intervertebral discs. Damage to the apophyseal joints appears to be severe, with possible microfractures (not detectable on plain X-ray) and long-term dysfunction.

Pain and stiffness of the neck are the most common symptoms. The pain is usually experienced in the neck and upper shoulders but may radiate to the suboccipital region, the interscapular region and down the arms. The stiffness felt initially in the anterior neck muscles shifts to the posterior neck.

Headache is a common and disabling symptom that may persist for many months. It is typically occipital but can be referred to the temporal region and the eyes.

Nerve root pain can be caused by a traction injury of the cervical nerve roots or by inflammatory changes or direct pressure subsequent to herniation of a disc.

Paraesthesia of the ulnar border of the hand, nausea and dizziness are all relatively common symptoms.

Delayed symptoms are common. A patient may feel no pain until 24 (sometimes up to 96) hours later; most experience symptoms within 6 hours. Complications of whiplash are summarised in Table 62.4.

Table 62.4 Complications of whiplash

Referred pain (headache, arm pain)
Visual problems
Vertigo
Dysphagia
Depression
Compensation neurosis
Disc rupture increasing to nerve root pain
Osteoarthritis becomes symptomatic

The Canadian guidelines (1995) for whiplash are:

- Grade I - neck pain, stiffness or tenderness
- Grade II - neck symptoms + musculoskeletal signs (e.g. decreased range of motion, point tenderness)
- Grade III - neck symptoms + neurological signs
- Grade IV - neck symptoms + fracture or dislocation

Management principles

The objective of treatment is to obtain a full range of free movement of the neck without pain by attending to both the physical and the psychological components of the problem. Other objectives include an early return to work and discouragement of unnecessary and excessive reliance on cervical collars and legal action.

Treatment

- Establish an appropriate empathy and instil patient confidence with a positive professional approach. Discourage multiple therapists.
- Provide appropriate reassurance and patient education.
- Encourage normalisation of activities ASAP.
- Compare the problem with a sprained ankle, which is a similar injury.
- Inform that an emotional reaction of anger, frustration and temporary depression is common (lasts about 2

weeks).

- X-ray is required.
- Prescribe rest only for grades II and III (max. 4 days).
- Use a cervical collar (limit to 2 days) for grades II and III.
- Use analgesics (e.g. paracetamol) - avoid narcotics.
- Use NSAIDs for 14 days.
- Use tranquillisers, mild - up to 2 weeks.
- Refer for physiotherapy.
- Provide neck exercises (as early as possible).
- Use heat and massage - 'spray and stretch'.
- Give passive mobilisation (not manipulation).

Recovery can take any time from 1 to 2 weeks up to about 3 months. A valuable reference is *Update Quebec Task Force Guidelines for the Management of Whiplash - Associated Disorder* at < http://www.maa.nsw.gov.au >.

CERVICAL DISC DISRUPTION

Disruption of a cervical disc can result in several different syndromes.

1. Referred pain over a widespread area due to pressure on adjacent dura mater.
   Note: A disc disruption is capable of referring pain over such a diffuse area (Figure 62.8) that the patient is sometimes diagnosed as functional (e.g. hysterical).



Figure 62.8 Zone of possible referred pain distribution caused by a cervical disc lesion on the right side

2. Nerve root or radicular pain (radiculopathy). The pain follows the dermatomal distribution of the nerve root in the arm.
3. Spinal cord compression (myelopathy).

RADICULOPATHY

Apart from protrusion from an intervertebral disc, nerve root pressure causing arm pain can be caused by osteophytes associated with cervical spondylosis. Uncommon causes include various tumours involving the vetebral segment, the meninges and nerves or their sheaths. The pain follows neurological patterns down the arm, being easier to localise with lower cervical roots, especially C6, C7 and C8.

1. The cervical roots exit above their respective vertebral bodies. For example, the C6 root exits between C5 and C6 so that a prolapse of C5 - 6 intervertebral disc or spondylosis of the C5 - 6 junction affects primarily the C6 root (Figure 62.9).



Figure 62.9 Typical C6 nerve root (radicular) pain

2. One disc - one nerve root is the rule.
3. Spondylosis and tumours tend to cause bilateral pain (i.e. more than one nerve root).

Clinical presentation

- A sharp aching pain in the neck, radiating down one or both arms
- Onset of pain may be abrupt, often precipitated by a sudden neck movement on awakening
- Paraesthesiae in the forearm and hand (in particular) - in 90% with proven disc prolapse[9]
- Stiffness of neck with limitation of movement
- Nocturnal pain, waking patient during night
- Pain localised to upper trapezius and possible muscle spasm

Investigations

- Plain X-ray (AP, lateral extension and flexion, oblique views to visualise foramina); not good for diagnosis or for surgery
- Plain CT scan
- CT scan and myelogram - excellent visualisation of structures but invasive
- MRI - excellent but expensive, sometimes difficult to distinguish soft disc from osteophytes
- Electromyography - may help delineate lesions requiring surgery

Treatment

Many patients respond to conservative treatment, especially from a disc prolapse. It is basically a self-limiting disorder - about 10% remain severely disabled.[9]

- bed rest
- soft cervical collar
- analgesics
- tranquillisers, especially at night
- traction (with care)
- careful mobilisation (manipulation is contraindicated)

MYELOPATHY

Sometimes the presence of large or multiple osteophytes or in the presence of a narrowed symptoms of spinal-cord involvement may develop.[10, 11] The common cause is a hard mass of maternal projecting from the posterior aspect of the vertebral body to indent the spinal cord. This resultant spinal cord compression may result in several different clinical presculations, notally myelopathy in particular but also central cord and anterior cord syndrome. A full neurological assessment is necessary.

Clinical features of myelopathy:

- older patients, typically men > 50 years
- insidious onset - symptoms over 1 - 2 years
- numbness and tingling in fingers
- leg stiffness
- numb clumsy hands, especially with a high cervical lesion
- signs of UMN: spastic weakness, increased tone and hyper-reflexia (arms > legs)
- neurological deficit, which predicts the level with reasonable accuracy
- bowel and bladder function usually spared

Note: LMN signs occur at the level of the lesion, and UMN signs and sensory changes occur below this level.

Causes

- Cervical spondylosis
- Atlantoaxial subluxation
    - rheumatoid arthritis
    - Down syndrome
- Primary spinal cord tumours (e.g. meningiomas)
- Metastasis to cervical spine → epidural spinal cord compression

Investigations

- MRI scan
- CT scan with myelogram (most accurate)

Central cord syndrome[11]

This rather bizarre condition occurs classically in a patient with a degenerative cervical spine following a hyperextension injury that causes osteophytes to compress the cord anteriorly and posteriorly simultaneously.

The maximum damage occurs in the central part of the cord leading to sensory and motor changes in the upper limbs with relative sparing of the lower limbs due to the arrangements of the long tracts in the cord.

Fortunately, the prognosis is good with most patients achieving a good neurological recovery.

Anterior cord syndrome

The anterior cord syndrome occurs with hyperflexion injuries that produce 'tear drop' fractures of the vertebral bodies or extrusion of disc material. The syndrome can also be produced by comminuted vertebral body fractures.

It is characterised by complete motor loss and the loss of pain and temperature discrimination below the level of the injury, but deep touch, position and vibration sensation remain intact.

Because it is probably associated with obstruction of the anterior spinal artery, early surgical intervention to relieve pressure on the front of the cord may enhance recovery. Otherwise the prognosis for recovery is poor.

Down syndrome

One of the more sinister problems with trisomy 21 syndrome is hypoplasia of the odontoid process, leading to C1 - 2 subluxation and dislocation. If unrecognised in the early stages, sudden death can occur in these children. If suspected, flexion - extension lateral views of the cervical spine will highlight the developing instability and the need for early specialist opinion.

Rheumatoid arthritis

Involvement of the cervical spine is usually a late manifestation of RA. It is important to be aware of the potentially lethal problem of C1 - 2 instability due to erosion of the major odontoid ligaments in the rheumatoid patient. These patients are especially vulnerable to disasters when under general anaesthesia and when involved in motor vehicle accidents. Early cervical fusion can prevent tragedies, especially with inappropriate procedures such as cervical manipulation. It is imperative to X-ray the cervical spine of all patients with severe RA before major surgery to search for C1 - 2 instability.

Treatment of spondylotic myelopathy

Conservative (may help up to 50%):[1]

- soft cervical collar
- physiotherapy for muscle weakness
- analgesics and/or NSAIDs

Surgery is indicated when the myelopathy interferes with daily activities. One procedure is the 'Cloward' method, which is anterior decompression with discectomy and fusion.

WHEN TO REFER

- Persisting radicular pain in an arm despite conservative treatment

Practice tips

- 'One disc - one nerve root' is a working rule for the cervical spine.
- The patient should sit on the couch with the thighs fully supported for inspection and movements of the neck.
- Beware of patients with RA and Down syndrome who have cervical instability. Physical treatments such as cervical manipulation may easily cause quadriplegia.
- All acutely painful conditions of the cervical spine following trauma should be investigated with a careful neurological examination of the limbs, sphincter tone and reflexes. Plain film radiology is mandatory.
- In conscious patients, flexion and extension lateral cervical spinal plain films are useful for diagnosing instability of spinal segments with or without associated spinal fractures.
- The so-called 'whiplash' syndrome is a diagnosis of exclusion of spinal fractures or severe ligamentous disruption causing instability, and even then, for medicolegal and psychological reasons, would best be termed a 'soft tissue injury of the cervical spine'.
- Most 'soft tissue cervical spine injuries' heal within 3 months with conservative treatment. If severe pain

persists, follow-up investigations may be required.
- Dysfunction of the cervical spine is an underestimated cause of headache.
- Always consider dysfunction of the cervical spine as a possible cause of shoulder pain.
- Strains and fractures of the apophyseal joints, especially after a whiplash injury, are difficult to detect, and are often overlooked causes of neck and referred pain.

- Evidence of involvement of more than one nerve root lesion in the arm
- Evidence of myelopathy, such as weakness, numbness or clumsiness of the upper limbs
- Evidence, clinical or radiological, of cervical instability in post-accident victims, Down syndrome or rheumatoid arthritis

REFERENCES

1. Cohen ML. Neck pain. Mod Med Aust, 1989; November: 44 - 53.
2. Payne R. Neck pain in the elderly: a management review. Mod Med Aust, 1988; July: 56 - 67.
3. Bogduk N. Neck pain. Aust Fam Physician, 1984; 13: 26 - 9.
4. Hart FD. *Practical Problems in Rheumatology*. London: Dunitz, 1985: 10 - 14.
5. Sloane PD, Slatt LM, Baker RM. *Essentials of Family Medicine*. Baltimore: Williams & Wilkins, 1988: 236 - 40.
6. NH & MRC. Evidence-based management of acute musculoskeletal pain: a guide for clinicans: Australian Acute Musculoskeletal Pain Guidelines Group. Canberra: Australian Government, 2004: 36 - 43.
7. Beran RG et al. Serious complications with neck manipulation and informed consent. Med J Aust, 2000; 173: 213 - 14.
8. Kenna C, Murtagh J. *Back Pain and Spinal Manipulation* (2nd edn). Oxford: Butterworth-Heinemann, 1997: 83 - 99.
9. Bogduk N. *Medical Management of Acute Cervical Radicular Pain - an Evidence Based Approach*. Newcastle: Newcastle Bone and Joint Institute, 1999: 5 - 59.
10. Corrigan B, Maitland GP. *Practical Orthopaedic Medicine*. Sydney: Butterworths, 1986:352.
11. Young D, Murtagh J. Pitfalls in orthopaedics. Aust Fam Physician, 1989; 18: 645 - 6.

CHAPTER 63. SHOULDER PAIN

It is important to realise that patients' perception of pain in the 'shoulder' may be amazing. For example, pain in the lower border of the scapula may be referred to as shoulder pain.

TEXT, PAGE 667

SHOULDER PAIN

The painful shoulder is a relatively common andsometimes complex problem encountered in general practice. The diagnostic approach involves determining whether the disorder causing the pain arises from within the shoulder structures or from other sources such as the cervical spine (Figure 63.1), the acromio-clavicular (AC) joint or diseased viscera, especially the heart, lungs and sub-diaphragmaticstructures.[1]



Figure 63.1 Typical pain zone arising from disorders of the shoulder joint and the lower cervical spine (C5 level)

KEY FACTS AND CHECKPOINTS

- Virtually all shoulder structures are innervated by the fifth cervical vertebra (C5) nerve root. Pain present in the distribution of the C5 nerve can arise from the:
  - cervical spine
  - upper roots of brachial plexus
  - glenohumeral joint
  - rotator cuff tendons, especially supraspinatus
  - biceps tendon
  - soft tissue (e.g. polymyalgia rheumatica)
  - viscera, especially those innervated by thephrenic nerve (C3, C4, C5).
- The visceral diseases causing a painful shoulder include cardiac disorders, such as angina and pericarditis; lung diseases, especially Pancoast tumour; mediastinal disorders; and diaphragmatic irritation, as from intraabdominal bleeding or a subphrenic abscess.

- A careful history should generally indicatewhether the neck or the shoulder is responsible for the patient's pain.
- By the age of 50 about 25% of people have some wear and tear of the rotator cuff, making it more injury-prone.[2]
- Disorders of the rotator cuff are common, especially supraspinatus tendonopathy.
  The most effective tests to diagnose these problems are the resisted movement tests.[3]
- Injections of local anaesthetic and long-acting corticosteroid produce excellent results for inflammatory disorders around the shoulder joint, especially for supraspinatus tendonopathy.

*Note:*The term tendonopathy or tendonosis is prefered to tendonitis since it has been shown that overuse tendon conditions generallyhave a non-inflammatory pathology.

FUNCTIONAL ANATOMY OF THE SHOULDER

A working knowledge of the anatomical features of the shoulder is essential for understanding the various disorders causing pain or dysfunction of the shoulder. Apart from the AC joint there are two most significant functional joints - the glenohumeral (the primary joint) and the subacromial complex (the secondary joint) (Figure 63.2). The glenohumeral joint is a ball and socket joint enveloped by a loose capsule. It is prone to injury fromtraumatic forces and develops osteoarthritis more often than appreciated. Two other relevantfunctional joints are the scapulothoracic andstenoclavicular joints.



Figure 63.2 The basic anatomical structures of the shoulder joint

The clinically important perihumeral space lies above the glenohumeral joint between the head of the humerus and an arch formed by the bony acromion, the thick coracoacromial ligament and the coracoid process. This relatively tight compartment houses the subacromial bursa and the rotator cuff, particularly the vulnerable supraspinatus tendon.[4] Excessive friction and pinching in this space renders these structures prone to injury.

There is a critical zone of relative ischaemia that appears to affect the rotator cuff about 1 cm medial to the attachment of the supraspinatus tendon,[5] and this area is compromised during adduction and abduction of the arm due to pressure on the rotator cuff tendons from the head of the humerus.

Such factors are largely responsible for the many rotator cuff syndromes, bicipital tendonopathy, subacromial bursitis and lesions of the supraspinatus tendon.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 63.1.

Table 63.1 Shoulder pain: diagnostic strategy model

Q. Probability diagnosis

A. Cervical spine dysfunction

Rotator cuff tendonopathy ±a tear

Adhesive capsulitis

Bicipital tendonopathy

Q. Serious disorders not to be missed

A. Cardiovascular

- angina
- myocardial infarction

Neoplasia

- Pancoast tumour
- primary or secondary in humerus

Severe infections

- septic arthritis (especially children)
- osteomyelitis

Rheumatoid arthritis

Q. Pitfalls (often missed)

A. Polymyalgia rheumatica

Cervical dysfunction

Gout/pseudogout

Osteoarthritis of acromioclavicular joint

Winged scapula - muscular fatigue pain

Q. Seven masquerades checklist

A. Depression ✔

Diabetes ✔

Drugs ✔

Anaemia -

Thyroid disorder rarely

Spinal dysfunction ✔

UTI -

Q. Is the patient trying to tell me something?

A. Shoulder is prone to (uncommonly) psychological fixation for secondary gains, depression and conversion reaction.

Probability diagnosis

The commonest causes of pain in the shoulder zone (Figure 63.1) are cervical disorders and periarthritis (i.e. soft tissue lesions involving the tendons around the glenohumeral joint). The outstanding common disorders of the shoulder joint are the various disorders of the tendons comprising the rotator cuff and biceps tendon. Of these, supraspinatus tendon disorders, which include wearing, calcific degeneration and tearing, are the commonest. It is obvious that the supraspinatus tendon issubjected to considerable friction and wear and tear.

Serious disorders not to be missed

As usual it is important to exclude any malignancy or septic infection, be it septic arthritis or osteomyelitis. Carcinoma of the lung (Pancoast's syndrome) should be kept in mind. For pain in the region of the left shoulder the possibility of myocardial ischaemia has to be considered. Referred pain to the right shoulder from myocardial ischaemia is rare, occurring about once for every 20 episodes of left shoulder referral.

With an acute onset of painful capsulitis the possibility of rheumatoid arthritis (or even gout) is worth considering.

Pitfalls

The shoulder is notorious for diagnostic traps, especially for referred pain from visceral structures, but polymyalgia rheumatica is the real pitfall. A good rule is to consider it foremost in any older person (over 60) presenting with bilateral shoulder girdle pain that is worse in the morning.

Specific pitfalls include:

- misdiagnosing posterior dislocation of the shoulder joint
- misdiagnosing recurrent subluxation of the shoulder joint
- overlooking an avascular humeral head (post fracture)
- misdiagnosing rotator cuff tear or degeneration

Seven masquerades checklist

Of the seven primary masquerades spinal dysfunction and depression are those most likely to be associated with shoulder pain. The degree to which cervical spondylosis is associated with shoulder pain is not always appreciated.

It is important to realise that patients' perception of pain in the 'shoulder' may be amazing. For example, pain in the lower border of the scapula may be referred to as shoulder pain.

Diabetics have a higher incidence of adhesive capsulitis. Drugs are relevant as cortiocosteroids can cause avascular necrosis of the humeral head and anabolic steroids (weight-lift ers) can cause osteolysis of the AC joint.

A summary of common shoulder conditions is presented in Table 63.2.

Table 63.2 Common shoulder conditions (after Murrell)[7]

Problem	Structure affected	Typical Age group	Symptoms	Diagnostic Pointers
Instability	Labrum/capsule	15 - 30	Dislocations	History of dislocation, apprehension sign
Stiff ness	Capsule	40 - 60	Pain, night pain, loss of movement	Loss of external rotation
Impingement	Rotator cuff (fatigue)	30 - 60	Night pain, pain with overhead activities	Impingement signs
Rotator cuff tear	Rotator cuff esp. supraspinatus	50+	As above	Impingement signs, weakness external rotation, weakness supraspinatus
AC joint pain	AC joint cartilage	25 - 45	Localised AC joint pain	Paxinos sign
Arthritis	Glenohumeral joint cartilage	65+	Pain, loss of movement	Crepitus

THE CLINICAL APPROACH

History

In analysing the pain pattern it is appropriate to keep the various causes of shoulder pain in mind (Table 63.3 on p. 669). Many of these conditions, such as rheumatoid arthritis, osteoarthritis and gout, are relatively uncommon.

Table 63.3 Causes of shoulder pain (excluding trauma, fractures and dislocations)

Cervical
• dysfunction
• spondylosis
Cervical radiculopathy
Polymyalgia rheumatica (bilateral)
Acromioclavicular joint
• dysfunction
• osteoarthritis
Shoulder complex
Extracapsular
• subacromial bursitis
• rotator cuff disorders
- supraspinatus tendonopathy
- infraspinatus tendonopathy
- subscapularis tendonopathy
• bicipital tendonitis
Intracapsular (glenohumeral joint)
• adhesive capsulitis
- idiopathic
- blunt trauma
- diabetes
- others
• rheumatoid inflammation
- rheumatoid arthritis
- ankylosing spondylitis
- psoriatic arthropathy
• osteoarthritis
• avascular necrosis
• septic arthritis
Winged scapula - muscular fatigue pain
Malignant disease
• primary or secondary in humerus
• Pancoast (referred from lung)
Referred pain
Cardiac
• ischaemic heart disease
• pericarditis
Lung

Mediastinum, including oesophagus
Diaphragmatic irritation
Herpes zoster

A careful history should generally indicate whether the neck or the shoulder (or both) is responsible for the patient's pain. Enquire about features of movement

- stiffness and restriction
- excessive movement/instability
- weakness
- rough v. smooth

Key questions

- Did you have any injury, even very minor, before your pain started?
- Does the pain keep you awake at night?
- Do you have pain or stiffness in your neck?
- Do you have pain or restriction when clipping or handling your bra or touching your shoulder blades? (indicates painful internal rotation and a problem of capsular restriction or a disorder of the acromioclavicular joint)
- Do you have trouble combing or attending to your hair? (indicates problematic external rotationand also a disorder of the capsule, e.g. adhesive capsulitis)
- Is the pain worse when you wake in the morning? (indicates inflammation)
- Do you have aching in both your shoulders or around your hips?
- Do you get pain associated with sporting activity, including weight training, or with housework, dressing or other activities?
- Do you think you can throw a ball underhand for 10 - 20 metres and/or overhead for 20 - 25 metres with your affected arm?
- Can you lift a full 2 litre container e.g. milk to the level of your shoulder without bending your elbow (or to the top of your head)?
- Can you carry a 20 - 30 kg weight e.g. full suitcase by your side

Physical examination

The diagnosis is based on systematic examination of the cervical spine followed by examination of the shoulder joint. For details of examination of the cervical spine, refer to Chapter 62.

Examination of the shoulder [4, 6]

For the examination of the shoulder it is important to understand thefunctional anatomy of all important tendons.

The tendon disorders are diagnosed by pain on resisted movement (Table 63.4). A knowledge of the anatomical attachments of the rotator cuff tendons to the head of thehumerus (Figure 63.3) provides an understanding of the shoulder movements powered by these muscles.

Table 63.4 Tendon disorders: determining resisted movements

Painful resisted movement at shoulder	Affected tendon
1Abduction	Supraspinatus
2Internal rotation	Subscapularis Teres minor *
3External rotation	Infraspinatus Biceps *

4 Adduction	Pectoralis major Latissimus dorsi*

* Lesser role



Figure 63.3 The attachments of the rotator cuff tendons to the head of the humerus. Reproduced with permission from C. Kenna and J. Murtagh. *Back Pain and Spinal Manipulation*. Sydney: Butterworths, 1989.

With tendon disorders (rotator cuff tendons or biceps) there is painful restriction of movement in one direction, but with capsulitis and subacromial bursitis thereis usually restriction in most directions.

Inspection

Observe the shape and contour of the shoulder joints and compare bothsides. Note the posture and the position of the neck and scapulae. The position of the scapulaeprovides considerable clinical information.

Note any deformity, swelling or muscle wasting.

Palpation

Stand behind the patient and palpate significant structures such as the AC joint, the subacromial space, the supraspinatus tendon and the long head of biceps. The subacromial bursa is one area where it is possible to localise tenderness with inflammation. Feel also over the supraspinatus and infraspinatus muscles for muscle spasm and trigger points. The axilla should be palpated for lymphadenopathy.

Movements

The movements of the shoulder joint are complex and involve the scapulothoracic joint as well as the glenohumeral joint, with each joint accounting for about half the total range. Significant signs of a painful capsular pattern can be gained by determining the movements of flexion, abduction, external rotation and internal rotation.

For each movement, note:

- the range of movement
- any pain reproduction
- any trick movement by the patient
- scapulothoracic rotation

Movements should be tested bilaterally and simultaneously wherever possible.

1 Active movements

- Flexion (anterior elevation) 180°

- Extension (posterior elevation) 45°

With the palm facing medially the patient moves the arm upwards through 180??to a vertical position above the head and then backwards through this plane.

- Abduction - 180°
- Adduction - 80°(from neutral position)

Abduction is possible only if the arm is fully externally rotated. Itis a key combined glenohumeral and scapulothoracic movement, which shouldreach 180°, and thesecomponents should be differentiated if the movement is limited. This is done by fixing the scapula with one hand holding the scapula at its inferior angle and noting the degree of movementof each component (initial glenohumeral range 85 - 100°). Look for the presence of a painful arc, which occurs usually between 60°and 120°of abduction (Figure 63.4). The commonestcause is supraspinatustendonopathy. Other causes include infraspinatus tendonopathy and subacromial bursitis (milder degree).



Figure 63.4 The painful arc syndrome

- Internal rotation - 90?
- External rotation - 90?

These movements are tested with the arm by the side and the elbow flexed to 90°with the palm facing medially. The hand is carried outwards to test externalrotation and inwards towards the abdomen for internal rotation.

2 Resisted movements[6]

Resisted movements (isometric contractions of a muscle) are importantways of testing capsulitis and for pinpointing tenderness of muscle insertions around the shoulder joint, and no examination of the shoulder is complete without them (seeTable 63.4).

Abduction(*supraspinatus test*). With the arm abducted tono more than 15°the patient pushes the elbow away from the side while the examiner's hands resist and prevent the movement, holding for 5 seconds. Compare both sides and note any reproduction of the patient's pain.

A better and more specific test for supraspinatus impingement is testing resisted elevation in the ‘emptying the can’ position (90??of abduction, 30?horizontal flexion and full internal rotation).

Internal rotation (subscapularis test). The examiner stands behind the patient and grasps the palmar surface of the patient's wrists (with the arm bythe side and elbow at 90°). The patient attempts to move the forearm internally (medially) against resistance.

External rotation (infraspinatus test). With the examiner and patient adopting a similar position to that for internal rotation, the examiner grasps the dorsal surface of the forearm near the wrist and asks the patient to press outwards, using the forearm as a lever to produce external rotation. This test is also positive for a C5 nerve root lesion.

3 Special tests

Supraspinatus/infraspinatus rapid differentiation test. A quick test that helps to differentiate between a lesion of either of these tendons causing a painful arc syndrome is the 'thumbs up/thumbs down' abduction test. To test the supraspinatus, perform abduction with thumbs pointing upwards, and then with the thumbs pointing downwards to test the infraspinatus.

Long head of biceps test. The best test is opposed forward elevation of the arm with the elbow at right angles. A positive test is reproduction of pain in the bicipital groove. Another useful test is resisted supination at the wrist (Yergason'stest).

The brachial plexus tension test. This test, devised by Elvey,8 tests the nerve roots and sheaths of the brachial plexus without implicating the cervical spine and the gleno-humeral joint. The upper cervical roots of the plexus are sometimes injured in accidents, so this test is an effective differentiation test.

Impingement test for supraspinatus lesions. Refer to p. 672.

Investigations

Appropriate investigations for shoulder pain include:

- ESR (especially for polymyalgia rheumatica)
- rheumatoid factor
- serum uric acid (acute pain)
- ECG
- radiology
  - X-ray of a specific part of the shoulder - AC joint, axillary view of glenohumeral joint (best view to show osteoarthritis)
  - X-ray of cervical spine and chest (if relevant)
  - radionuclide bone scan - to assess bone tumours
  - shoot-through axillary views (posterior dislocation)
  - high-resolution ultrasound - modern techniques make this an appropriate test to assess shoulder pain due to rotator cuff lesions, especially tears and capsulitis, especially if surgery is contemplated.
    However this test as sometimes reported can be misleading.
  - arthrogram of shoulder (beware of false negatives)
  - CT scan (limited use)
  - MRI - a useful imaging method but not routinely required except for the unstable joint
  - arthroscopy

SHOULDER TIP PAIN

Pain at the shoulder tip may be caused by local musculoskeletal trauma or inflammation or can be referred. Referred causes include:

- peptic ulceration
- diaphragmatic irritation
- ruptured viscus (e.g. perforated ulcer)
- intraperitoneal bleeding (e.g. ruptured spleen)
- pneumothorax
- myocardial infarction

SHOULDER PAIN IN CHILDREN

Shoulder pain in children is not a common presenting problem but the following require consideration:

- septic arthritis/osteomyelitis
- swimmer's shoulder

Swimmer's shoulder

Although it occurs in adults, shoulder pain is the most common complaint in swimmers in the teenage years (over 12 years of age). American studies of college and national competition swimmers showed 40 - 60% had suffered significant pain.[9]

The problem, which is considered to be associated with abnormal scapular positioning and cervicothoracic dysfunction, occurs in the supraspinatus tendon where an avascular zone is compressed by the greater tuberosity when the arm is adducted and relieved whenabducted. Swimmers' shoulders are forced through thousands of revolutions each day, so the susceptible area tends to impinge on the coracoacromial arch, leading to the impingement syndrome, which can progress with continued stress and age.[10]

Symptoms

- Stage 1: pain only after activity
- Stage 2: pain at beginning only, then after activity
- Stage 3: pain during and after activity, affects performance

Management

- Early recognition is important.
- Discuss training program with coach.
- Consider alteration of technique.
- Application of ICE after each swim.
- Use NSAIDs.
- Avoid corticosteroid injections.
- Refer for physiotherapy for scapular stabilisation and cervicothoracic mobilisation.

SHOULDER PAIN IN THE ELDERLY

As a rule most of the shoulder problems increase with age. Special features in the elderly are:

- polymyalgia rheumatica (increased incidence with age)
- supraspinatus tears and persistent 'tendonitis'
- other rotator cuff disorders
- stiff shoulder due to adhesive capsulitis
- osteoarthritis of AC and glenohumeral joints
- cervical dysfunction with referred pain
- the avascular humeral head

Since the rotator cuff is prone to degeneration with age there is a high incidence of rotator cuff tears in the elderly that are mostly asymptomatic.

The avascular humeral head

The humeral head may become avascular after major proximal humeral fractures. With experience, it is usually possible to predict the fractures at special risk. Early humeral head replacement with a prosthesis can lead to excellent pain relief and to a return of good function. Once the head has collapsed, there is secondary capsular contracture. Prosthetic replacement of the head is then rarely associated with an adequate return of joint movement. Thus, early referral of comminuted proximal humeral fractures for an expert opinion in allage groups is good practice.

Early replacement can improve the functional outcome.[11]

ROTATOR CUFF TENDONOPATHY[12]

Rotator cuff tendonopathy also referred to as ‘impingement syndrome', is the commonest cause of shoulder pain. It can be associated with inflammation (tendonitis), a tear in a tendon or impingement under the acromion. It may involve one tendon, usually the supraspinatus, or more of the rotator cuff tendons. It is most frequently encountered in young people engaged in sport involved in overhead activities and people over 50 years, in whomrotator cuff tears occur most often.

Supraspinatus tendonopathy can vary in intensity from mild to extremely severe. The severe cases usually involve calcification (calcific periarthritis) of the tendon and spread to the subacromial bursa (subacromial bursitis).

Supraspinatus tendonopathy

Typical pain profile

• Site:	The shoulder and outer border of arm; maximal over deltoid insertion
• Radiation:	to elbow
• Quality:	throbbing pain, can be severe
• Frequency:	constant, day and night
• Duration:	constant
• Onset:	straining the shoulder (e.g. dog on leash, working under car, fall onto outstretched arm)
• Off set:	nil
• Aggravation:	heat, putting on shirt, toilet activity, lying on shoulder
• Relief:	analgesics only
• Associated features:	trigger point oversupraspinatus origin
• Examination (typical features):	- painful resisted abduction - painful arc - painful resisted external rotation - positive impingement test - positive ‘emptying the can’ sign
• Diagnosis	high-resolution ultrasound

The impingement tests[7]

These are eff ective tests for supraspinatus lesions as they force impingement of the rotator cuff and bursa under the acromion. One of these tests is the ‘emptyingthe can’ resistance test. The arm is placed in the ‘emptying the can’ position (90°of abduction, 30°of horizontal flexion and full internal rotation). Elevation of the arm is resisted against the therapist's downward push. This also tests the strength of supraspinatus. Impingement can also be tested in external rotation when the arm is abducted to 90° and externally rotated.

Treatment[12]

Systematic reviews to date have a lack of sufficient information to provide conclusive evidence-based recommendations for treatment.[13] NSAIDs might provide some relief from pain while corticosteroid injections and physiotherapy could improve range of movement. Experienced therapists believe that peritendon and subacromial corticosteroid injections are efficacious in selected patients.

- Rest during the acute painful phase
- Analgesics and NSAIDs (up to 4 weeks)
- Peritendon or subacromial injection (if no tears on ultrasound)
- Physiotherapy - an active program
- Exercises when acute phase subsides

Injection technique

The ideal injection is a specific injection onto the tendon rather than general infiltration into the subacromial space. As a rule the therapeutic result is quite dramatic after one or two days of initial discomfort (often severe). The tendon can be readily palpated as a tender cord anterolaterally as it emerges from beneath the acromion to attach to the greater tuberosity of the humerus. This identification is assisted by depressing theshoulder via a downward pull on the arm and then externally and internally rotating the humerus. This manoeuvre allows the examiner to locate the tendon readily.

Method

- Identify and mark the tendon.
- Place the patient's arm behind the back, with the back of the hand touching the far waistline. This locates the arm in the desired internal rotation and forces the humeral head anteriorly.
- Insert a 23-gauge 32-mm needle under the acromion along the line of the tendon, and inject around the tendon just under the acromion (Figure 63.5). If the gritty resistance of the tendon is encountered, slightly withdraw the needle to ensure that it lies in the tendon sheath.



XX Palpate for
(1) tenderness
(2) site of injections

Figure 63.5 Injection placement for supraspinatus tendonopathy

- The recommended injection is 1 mL of a soluble or long-acting corticosteroid with 5 mL of 1% lignocaine.

Persistent supraspinatus tendonopathy

There are three factors to consider with this problem:

1. *A very tight subacromial space*. Refer for subacromial decompression by division of the thickened coracoacromial ligament. Even in younger patients this procedure (with or without acromioplasty) may be indicated for those with pain persisting beyond 12 months.
2. *Rotator cuff tear or degeneration*. In middleaged and elderly patients persisting tendonitis is usually due to rotator cuff tear and degeneration, an underdiagnosed condition. Excellent clinical and functional results can be achieved if surgery is performed when the tear is small.
3. *Calcification of the tendon*. This problem usually settles but occasionally surgical intervention is necessary.

Other rotator cuff lesions

The patient may present with dominant signs of subscapularis or infraspinatus lesions, or a combination of two or three tendinous lesions, including the supraspinatus. This problem could be confused with milder adhesive capsulitis, hence the value of investigations such as ultrasound.

Management

A subacromial space injection of 1 mL of corticosteroid and 2 - 3 mL of 1% local anaesthetic, using the posterior approach, generally achieves a good result for multiple affected rotator cuff tendons with or without subacromial bursitis.

Method

With the patient sitting upright the large posterior gap between the medial acromial ridge and the humeral head is identified by palpation from behind. The needle (23 gauge, 32 or 38-mm long) is inserted into this gap just inferior to the acromion. The solution should flow into this space without resistance.

Rotator cuff tears

Asymptomatic rotator cuff tears are common, being present in 4% of people < 40 years old and in more than 50% of these over 60 years, but a significant number willbecome symptomatic over time.[14] Explain to the patient that ‘the rotator cuffis worn not torn’ rather like the frayed heel of a sock that may have a split in it.

Diagnostic tip: 98% specificity for all three signs:[15]

- supraspinatus weakness
- weakness in external rotation
- impingement (in external or internal rotation or both)

If two of these three tests are positive in a patient over 60 there is a 98% chance of a rotator cuff tear.

SUBACROMIAL BURSITIS

Subacromial (subdeltoid) bursitis is the more severe association of the frozen shoulder and mayrequire hospital admission for pain control. It is the only inflammatory disorder around the shoulder joint where localised tenderness is a reliable sign.

Typical pain profile

• Site:	outer shoulder, outer arm
• Radiation:	to outer elbow and upper forearm
• Quality:	intense pain
• Frequency:	constant
• Duration:	constant
• Onset:	either spontaneous or following unaccustomed work
• Off set:	nil
• Aggravation:	heat, brushing hair, most activities
• Relief:	very strong analgesics only
• Examination (typical features):	- ‘frozen’ shoulder - difficult to undress and dress

	- marked tenderness below acromion over deltoid
	- all active movements limited and painful

Management

- Strong analgesics (e.g. paracetamol and codeine)
- Large local injection of 5 - 8 mL of local anaesthetic into and around the bursa just beneath the acromion, followed immediately by 1 mL of corticosteroid (long-acting) into the focus of the lesion

ADHESIVE CAPSULITIS

Adhesive capsulitis or capulosis is an acute inflammation affecting the glenohumeral joint, which becomes fibrotic and contracted. It can arise spontaneously or post injury and may be partial or global which is the classic cause of the 'frozen shoulder'. Differential diagnoses include monoarticular rheumatoid arthritis, a crystal arthropathy such as gout, and septic arthritis. It is worse in diabetics. It is common and estimated to affect 2 - 5% of the general population and 10 - 20% of diabetes.[14]

It generally occurs in three stages:[14]

1. 'freezing, frozen and thawing' - an inflammatory painful phase of 2 - 9 months
2. a fibrotic contracted phase of 4 - 12 months
3. partial or complete resolution of 5 - 26 months

Typical pain profile

Usually affects people in their 40s and 50s.

• Site:	around the shoulder and outer border of arm
• Radiation:	to elbow
• Quality:	deep throbbing pain
• Frequency:	constant, day and night
• Duration:	Constant
• Onset:	Spontaneous, wakes the patient from sleep
• Offset:	nil
• Aggravation:	activity, dressing, combing hair, heat
• Relief:	analgesics only (partial relief)
• Associated features:	stiffness of arm, may be frozen
• Examination (typical features):	- 'frozen' shoulder (some cases)
	- various active and passive movements painful and restricted, especially extension
	- resisted movements pain-free (Patient compensates with scapulo-humeral movements)
• Diagnosis:	high-resolution ultrasound

Treatment

This problem, which can persist for at least 18 - 24 months (average time to restore motion is 30 months) and is usually self-limiting, can be treated with an intra-articular injection of corticosteroid but it is often unsuccessful. The modern treatment is hydrodilation of the joint with a large quantity of sterile solution (to stretch the capsule) ±

corticosteroid. This procedure should be performed slowly to produce an audible'pop' as fluid distends into the subacromial and subcoracoid bursae. Another important treatment is severing adhesions under arthroscopic control. The rule is: if very stiff use arthroscopy; if more mobile use a distension procedure. Active exercises are important to restore function. Fifty per cent of people with adhesive capsulitis do not regain full normal movement if untreated.

Current evidence from systemic reviews indicates that both hydrodistension and intra-articular injections are likely to be beneficial.[13]

BICIPITAL TENDONOPATHY

Bicipital tendonopathy is a lesion such as fraying or tearing of the long head of the biceps, which causes pain in front of the shoulder. Important signs include pain on resisted flexion of the elbow joint and on resisted supination with the elbow flexed to90° and forearm pronated (Yergason's test). A painful arc may be present when theintrascapular part is affected. Hence it is often confused with one of the rotator cuff lesions. Sometimes it is possible to elicit local tenderness by palpation along the course of the tendon in the bicipital groove. This is best done when the arm is externally rotated. Most active shoulder movements, especially external rotation, bring on the pain.

Bicipital tendonopathy is a common problem. It usually follows chronic repetitive strains in young to middle-aged adults (e.g. home decorating, weight training, tennis, swimming freestyle, cricket and baseball pitching). Two complications are complete ruptureand subluxation of the tendon out of its groove.

One treatment to consider is a corticosteroid and local anaesthetic injection at the site of maximal tenderness in the bicipital groove (Figure 63.6). As a rule it is best to refer a significant lesion.



Figure 63.6 Injection placement for bicipital tendonopathy

Rupture of the biceps tendon

Rupture of the long head of biceps usually occurs in the older person. It may be spontaneous or occur after lifting or falling on the outstretched hand. The patient usually feels a tearing or snapping sensation in the shoulder. The shoulder may be painful and difficult to move. The upper arm looks bruised and a lump due to rolled up belly of biceps is obvious on flexion of the elbow. Active treatment is not usually indicated but surgical intervention is appropriate for the young active person especially those in power sports.

POLYMYALGIA RHEUMATICA

It is very important not to misdiagnose polymyalgia rheumatica in the older person (over 50 years) presenting with bilateral pain and stiffness in the shoulder girdle. It may or may not be associated with hip girdle pain. Polymyalgia rheumatica sometimes follows an influenza-like illness. The patients seem to complain bitterly about their pain and seem flat and miserable. In the presence of a normal physical examination they are sometimes misdiagnosed as ‘rheumatics’ or ‘fibrositis’.

Typical pain profile

• Site:	shoulders and upper arms (Figure63.7)
• Radiation:	towards lower neck
• Quality:	a deep intense ache
• Frequency:	daily
• Duration:	constant but easier in afternoon and evening
• Onset:	wakes with pain at greatest intensity
• Offset:	nil
• Aggravating factors:	staying in bed, inactivity
• Relieving factors:	activity (slight relief)
• Associated features:	severe morning stiffness ‘in muscles’ malaise; ±weight loss, depression
• Diagnosis:	greatly elevated ESR (can be normal)
• Treatment:	corticosteroids give dramatic relief but long-term management can be problematic; regular review and support is essential (refer pp. 290 - 1)



Figure 63.7 Polymyalgia rheumatica: typical area of painaround the shoulder girdle

POSTERIOR DISLOCATION OF THE SHOULDER

This is a rare form of shoulder instability, which is oft en misdiagnosed. On first inspection there may not be an obvious abnormality of the shoulder contour. Consider this condition if there is a history of electric shock or a tonic-clonic convulsion. The major clinical sign is painful restriction of external rotation, which is usually completely blocked. Routine shoulder X-rays following trauma should always include the 'axillary shoot-through' view and thenthe diagnosis becomes obvious. Early diagnosis and management can prevent a poor outcome and perhaps litigation.[11]

RECURRENT SUBLUXATION

Recurrent anterior or inferior subluxations, or both, are probably more common than recurrent dislocations, yet frequently are not diagnosed. Patients complaining of attacks of sudden weakness and even a 'dead arm feeling' lasting for a few minutes with overhead activities of the arm should be investigated for this condition.

The disorder is usually apparent on careful stress testing of the shoulder. Air-contrast CT arthrography is considered the best investigation. Surgery is usually curative while conservative treatment oft en fails for younger patients.

GLENOID LABRUM INJURIES [16]

The glenoid labrum is the ring of fibrous tissue attached to the rim of the glenoid and provides volume to the cavity and stability to the glenohumeral joint. Injuries to the labrum are divided into superior labrum anterior to posterior (SLAP) or non-SLAP lesions and further into stable and unstable lesions.

Non-SLAP lesions include degenerative, flap and vertical labral tears as well as unstable lesions such as the classic Bankart lesion, where the labrum and capsule is detached from the rim.

SHOULDER INSTABILITY [7, 16]

Recurrent shoulder instability can be divided into three main types:

1. those with a tendency to generalised laxity of multiple joints including the shoulder and which tend to dislocate with minor injuries. Surgery is less effective and treatment is based on improving muscular stability with physiotherapy rehabilitation.
2. those following trauma, which includes avulsion of the anterior labrum (Bankart lesion). Physiotherapy tends to be less effective and the patients often require surgical repair.
3. those with chronic rotator cuff tendonopathy/impingement who develop subtle instability. Refer first to a sports physician or physiotherapist for assessment and management, preferably conservative initially.

The 'apprehension' test is useful to confirm the diagnosis of traumatic anterior instability. In this test the patient lies supine while the arm is externally rotated with the elbow flexed to 90???The test is more reliable when the patient expresses apprehension that the shoulder will 'come out', rather than pain.

OSTEOARTHRITIS OF THE GLENOHUMERAL JOINT

This is usually secondary to local trauma, long standing rotator cuff lesions and multiple surgical interventions. Shoulder movements are stiff and usually restricted in all directions. Plain X-rays show typical osteoarthritic changes. Treatment includes basic analgesics and short courses of NSAIDs plus exercises to improve mobility. Patients usually manage to cope with osteoarthritis of the shoulder, but for severe pain and stiffness arthroplasty or joint replacement should be considered.

ACROMIOCLAVICULAR OSTEOARTHRITIS

This condition is usually traumatic or degenerative and is relatively common in builders and sportspeople, especially rowers, and the elderly. A key test for AC joint pain is the Paxinos sign, which is positive when pain is elicited on compression of the joint by placing one hand on the back of the acromion and one on the clavicle. It is treated with rest and support and analgesics. Intra-articular injections of corticosteroids can be used for resistant or severe cases. If these measures are ineffectual, pain may be relieved by excision of the lateral end of the clavicle.

Practice tips

- Consider dysfunction of the cervical spine, especially C4 - 5 and C5 - 6 levels, as a cause of shoulder pain.
- Tendonitis and bursitis are very refractory to treatment and tend to last for several months. One well-placed injection of local anaesthetic and corticosteroid may give rapid and lasting relief.
- Test for supraspinatus disorders (including swimmer's shoulder) with the impingement tests, including the 'emptying the can' test.
- Modern ultrasound is the investigation of choice for painful disorders of the rotator cuff, especially to investigate tears in tendons.
- An elderly person presenting with bilateral shoulder girdle pain has polymyalgia rheumatica until proved otherwise. Relief from corticosteroids is dramatic. Although bilateral, it may start as unilateral discomfort.
- Dysfunction of the cervical spine can coexist with dysfunction of the shoulder joints.
- Correlation between clinical symptoms and the degree of tendon injury or failure is not reliable.16

WHEN TO REFER

- Persisting night pain with shoulder joint stiff ness
- Persisting supraspinatus tendonopathy; consider possibility of rotator cuff tear or degeneration, especially in the elderly
- Persisting restriction of movement, such as restricted cross-body flexion (indicates capsular constriction)
- Persisting supraspinatus tendonopathy or other rotator cuff problem, because decompression of the subacromial space with division of the coracoacromial ligament ??acromioplasty gives excellent results
- Confirmed or suspected posterior dislocation of the shoulder - the most commonly missed major joint dislocation
- Confirmed or suspected recurrent subluxation or avascular humeral head
- Children with shoulder joint instability
- Swimmer's shoulder refractory to changes in technique and training schedule
- Severe osteoarthritis of the glenohumeral joint (which usually follows major trauma) for consideration of prosthetic replacement
- Severe osteoarthritis of the AC or glenohumeral joint

REFERENCES

1. Corrigan B. Painful shoulder. Patient Management, 1978; 2: 23.
2. Sloane PD, Slatt LM, Baker RM. *Essentials of Family Medicine*. Baltimore: Williams & Wilkins, 1988: 242.
3. Cormack J, Marinker M, Morrell D. *Practice: A Handbook of Primary Health Care*. London: Kluwer-Harrap, 1980; 3.62: 1 - 4.
4. Kenna C, Murtagh J. *Back Pain and Spinal Manipulation* (2nd edn). Oxford: Butterworth-Heinemann, 1997: 109 - 33.
5. Rathburn JB, Macnab I. The microvascular pattern of the rotator cuff. J Bone Joint Surg, 1970; 52B: 540.
6. Cyriax J. *Textbook of Orthopaedic Medicine*, Vol. 1 (6th edn). London: Bailliere Tindall, 1976: 202.
7. Murrell G. Shoulder dysfunction: how to treat. Australian Doctor, 2004; 17 December: 23 - 30.
8. Elvey R. The investigation of arm pain. In: Grieve GP. *Modern Manual Therapy of the Vertebral Column*. London: Churchill Livingstone, 1986: 530 - 5.
9. Dominguez RH. Shoulder pain in swimmers. Physician Sportsmed, 1980; 8: 36.
10. McLean ID. Swimmers' injuries. Aust Fam Physician, 1984; 13: 499 - 500.
11. Young D, Murtagh J. Pitfalls in orthopaedics. Aust Fam Physician, 1989; 18: 645 - 8.
12. Mashford ML (Chair). *Therapeutic Guidelines: Analgesic* (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2002: 140 - 3.
13. Barton S ed. *Clinical Evidence*. London: BMJ Publishing Group, 2001; 850 - 63.
14. Mashford ML (Chair). *Therapeutic Guidelines: Rheumatology* (Version 1). Melbourne: Therapeutic Guidelines Ltd, 2006: 127 - 131.
15. Murrell GAC, Walton JR. Diagnosis of rotator cuff tears. Lancet, 2001; 357: 769 - 70.
16. Brukner P, Khan K. *Clinical Sports Medicine* (3rd edn). Sydney: McGraw-Hill, 2007: 261 - 6.
17. Sher JS et al. Abnormal findings on magnetic resonance images of asymptomatic shoulders. J Bone Joint Surg Am, 1995; 77: 10 - 15.

CHAPTER 64. PAIN IN THE ARM AND HAND

A pain in the hand is worth a look at the neck. By heck don't forget the neck!

ORTHOPAEDIC SURGEON TO STUDENTS (1965)

PAIN IN THE ARM AND HAND

Pain in the arm and hand is a common problem in general practice, tending to affect the middle aged and elderly in particular.

OVERVIEW OF CAUSES OF A PAINFUL ARM AND HAND

Like pain in the shoulder, pain originating from the cervical spine and shoulder disorders can extend down the arm. While pain from dis-orders of the shoulder joint (because of its C5 innervation) does not usually extend below the elbow, radiculopathies originating in the cervical spine can transmit to distal parts of the arm (Figure 64.4 at p. 683).



Figure 64.4 Lateral tennis elbow test: reproducing pain on resisted extension of the wrist

Important causes are illustrated in Figure 64.1. Myocardial ischaemia must be considered, especially for pain experienced down the inner left arm.


pain of shoulder origin
pain of cervical origin
disease of bone
• osteomyelitis
• neoplasia
myocardial ischaemia
lateral epicondylitis
medial epicondylitis
de Quervain's tenosynovitis
carpal tunnel syndrome
inflammation of tendons around wrist
median nerve neuropathy
ulnar nerve neuropathy

Figure 64.1 Important causes of arm pain (excluding trauma and arthritis)

Soft tissue disorders of the elbow are extremely common, especially tennis elbow. Two types of tennis elbow are identifiable: 'backhand' tennis elbow, or lateral epicondylitis, and 'forehand' tennis elbow, or medial epicondylitis, which is known also as golfer's or pitcher's elbow.

Other significant elbow disorders include inflammatory disorders of the elbow joint, such as rheumatoid arthritis, osteoarthritis and olecranon bursitis, which may follow recurrent trauma, gout, rheumatoid arthritis or infection.

Another important group of disorders are the various regional pain syndromes around the wrists, including the common de Quervain's tenosynovitis (affecting the tendons of extensor pollicis brevis and abductor pollicis longus) and to a lesser extent the extensor tendons to the fingers. Pain from these overuse syndromes can be referred in a retrograde manner into the forearm.

A fascinating and poorly understood syndrome is that related to dysfunction of the upper four vertebral segments of the thoracic spine, which can cause referred pain in the arm that does not correspond to the dermatomes. This syndrome is often confused with the more common regional pain disorders such as tenosynovitis and tennis elbow.

The various causes of the painful arm can be considered with the diagnostic model (Table 64.1).

Table 64.1 Pain in the arm and hand: diagnostic strategy model

Q. Probability diagnosis
A. Dysfunction of cervical spine (lower) Disorders of the shoulder
Medial or lateral epicondylitis

Overuse tendinitis of the wrist

Carpal tunnel syndrome

Osteoarthritis of thumb and DIP joints

Q. Serious disorders not to be missed

A. Cardiovascular

- angina (referred)
- myocardial infarction
- axillary vein thrombosis

Neoplasia

- Pancoast tumour
- bone tumours (rare)

Severe infections

- septic arthritis (shoulder/elbow)
- osteomyelitis
- infections of tendon sheath and fascial spaces of hand

Q. Pitfalls (often missed)

A. Entrapment neuropathies (e.g. median nerve, ulnar nerve)

Pulled elbow - children

Foreign body (e.g. elbow)

Rarities

- Polymyalgia rheumatica (for arm pain)
- Reflex sympathetic dystrophy
- Thoracic outlet syndrome
- Arm claudication (left arm)
- Kienböck's disorder

Q. Seven masquerades checklist

A. Depression

Diabetes

Drugs -

Anaemia -

Thyroid disorder -

Spinal dysfunction

UTI -

Q. Is the patient trying to tell me something?

A. Highly likely, especially with so-called RSI syndromes.

A DIAGNOSTIC APPROACH

Probability diagnosis

The commonest causes of arm pain are referred pain and radiculopathies caused by disorders of the cervical spine, the tennis elbows (lateral and, to a lesser extent, medial epicondylitis), carpal tunnel syndrome and regional pain syndromes caused by inflammation of the tendons around the wrist and thumb.

Disorders of the shoulder, particularly supraspinatus tendinitis, should be considered if the pain is present in the C5 dermatome distribution. Pain in the hand is commonly caused by osteoarthritis of the carpometacarpal joint of the thumb and the distal inter-phalangeal (DIP) joints, and also by the carpal tunnel syndrome (CTS).

Serious disorders not to be missed

Like any other presenting problem it is vital not to overlook malignant disease or severe infection. In the case of the arm, possible malignant disease includes tumours in bones, lymphoma involving axillary glands and Pancoast's syndrome.

Neoplastic tumours of the hand are uncommon and usually benign. Benign tumours include giant cell tumour of the tendon sheath, pigmented villonodular synovitis, neuri-lemmoma and neurofibroma. Malignant tumours are exceptionally rare but can include synovioma and rhabdomyosarcoma.

In addition, myocardial ischaemia, especially infarction in the case of pain of sudden onset, should be considered for left arm pain.

Sepsis can involve joints, the olecranon bursa and the deeper compartments of the hand, the latter leading to serious sequelae if not rapidly diagnosed and treated.

Subclavian or axillary vein thrombosis, known as ‘effort thrombosis’, causes swelling in the arm with pain high in the axilla. It is seen in people working constantly above their head, such as painters and basketballers. It is an emergency requiring antithrombotic therapy.

Pitfalls

Such conditions may include entrapment syndromes for peripheral nerves. If in doubt the patient should be referred for electromyography. Variations of peripheral nerve entrapments include the pronator syndrome (compression of the median nerve by the pronator teres or a fibrous band near the origin of the deep flexor muscles) and ulnar nerve entrapment at the elbow in the cubital fossa and, rarely, in Guyon's canal in the wrist.

Lesions of the nerve roots comprising the brachial plexus can also cause arm pain, especially in the C5 and C6 distribution. These can be detected by the brachial plexus tension tests.

Rarer causes of arm pain

These include polymyalgia rheumatica, although the pain typically involves the shoulder girdle, regional pain syndrome (Sudeck's atrophy) and the thoracic outlet syndromes.

The thoracic outlet syndromes include problems arising from compression or intermittent obstruction of the neurovascular bundle supplying the upper extremity, for example, cervical rib syndrome, costoclavicular syndrome, scalenus anterior and medius syndrome, ‘effort thrombosis’ of axillary and subclavian veins and the subclavian steal syndrome.

The commonest cause of the thoracic outlet syndrome is sagging musculature related to ageing, obesity, and heavy breasts and arms, aptly described by Swift and Nichols as ‘the droopy shoulder syndrome’.[1]

Cervical ribs are relatively common and may, or may not, contribute to the thoracic outlet syndrome. Often the cause is a functional change in the thoracic outlet due to the ‘droopy shoulder syndrome’ with no significant anatomical fault.[2]

Arm claudication is also rare. It can occur with arterial obstruction due to occlusion of the proximal left subclavian artery or the innominate artery. Exercise of the arm may be associated with central nervous system symptoms as well as claudication.

Seven masquerades checklist

Of the seven primary masquerades, spinal dysfunction and depression are those most likely to be associated with arm pain. Nerve root pain arising from entrapment in intervertebral foramina of the cervical spine or from a disc prolapse frequently leads to pain and/or paraesthesia in the arm.

Although diabetic neuropathy primarily manifests in the lower limbs it may be associated with neuropathies in the hands, including erythermalgia (redness and burning related to heat). Hypothyroidism may cause a carpal tunnel syndrome.

Psychogenic considerations

The hand can be regarded as a highly emotive 'organ' that is frequently used to give outward expression to inner feelings. These can range from grossly disturbed psychiatric behaviour, manifested as a hysterical conversion disorder by a non-functioning hand, to occupational neuroses such as repetition strain injury (RSI) and malingering.[3] Experienced occupational physicians and surgeons[3] find the hand and arm a source of functional disability most often as a result of industrial injury. Of great concern are the various so-called RSI disorders, which in some people may be a means of work avoidance or a 'ticket' for compensation or both.

THE CLINICAL APPROACH

History

The painful arm represents a real diagnostic challenge, so the history is very relevant.

It is common for arm pain to cause sleep disturbances and three causes are cervical disorders, CTS and the thoracic outlet syndrome. The working rule is:

- thoracic outlet syndrome - patients cannot fall asleep
- CTS - patients wake in the middle of the night
- cervical spondylosis - wakes the patient with pain and stiffness that persists well into the day[4]

The history should include an analysis of the pain and a history of trauma, particularly unaccustomed activity. In children evidence should be obtained about the nature of any injury, especially pulling the child up by the arms or a fall on an outstretched hand, which can cause potentially serious fractures around the elbow.

Physical examination

As part of the physical examination of the painful arm it may be necessary to examine a variety of joints, including the cervical spine (Chapter 62), shoulder (Chapter 63), elbow, wrist and the various joints of the hand. The arms should be inspected as a whole and it is very important to have both arms free of clothing and compare both sides.

Elbow joint

Inspection (from anterior, lateral and posterior aspects). Hold elbow in an anatomical position to measure the carrying angle of forearm - elbow fully extended, forearm supinated (palm facing forwards) normal 5 - 15? (greater in females). Note any swellings:

- olecranon bursitis (bursa over olecranon)
- nodules
    - RA (subcutaneous border ulna)
    - gout
    - SLE (rare) and rheumatic fever (very rare)
    - granulomas (e.g. sarcoid)

Palpation. Perform with patient supine and elbow held in approximately 70? flexion. Palpate bony landmarks and soft tissue. Note especially any tenderness over lateral epicondyle (tennis elbow) and medial epicondyle (golfer's elbow).

Movement (test active and passive). Hinge joint:

- extension - flexion (0° to 150°)
    - the arc for daily living is 30 - 130°
    - limitation of extension is an early sign of synovitis
- pronation - supination (rotation)
    - occurs at radiohumeral joint
    - test in two positions:
        - 90° flexion (held to side of body)
        - at full extension
    - supination 85? plus
    - pronation 75? plus

Resisted movements

- Painful resisted flexion at wrist ° medial epicondylitis.
- Painful resisted extension at wrist ° lateral epicondylitis.

Wrist joint

Follow the usual rules: LOOK, FEEL, MOVE, TEST FUNCTION, MEASURE and X-RAY. Note swellings or deformities, including the anatomical snuff box and distal end of radius. Feel for heat, tenderness and swelling, especially over the radial aspect of the wrist.

Movements. With elbow fixed at 90? and held into the waist:

1. compare dorsiflexion and palmar flexion on both sides (normal range 80 - 90°)
2. compare ulnar deviation (normal to 45°) and radial deviation (30°)
3. compare pronation and supination (normal to 90° for both)

Neurological examination

Test sensation, motor power and reflexes where indicated.

Summary of tests for motor power:

- C5 - test resisted movement deltoid
- C6 - test resisted movement biceps
- C7 - test resisted movement triceps
- C8 - test resisted EPL and FDL
- T1 - test resisted interossei

Sensory patterns are presented in Figure 62.4.

Investigations

Pain in the arm and hand can be difficult to diagnose but the rule to follow is: ‘If in doubt, X-ray and compare both sides’. This applies particularly to elbow injuries in children. The presence of a foreign body in the hand or arm also requires consideration.

Investigations to consider include:

- blood film and WCC
- ESR
- ECG
- X-rays
    - cervical spine
    - upper thoracic spine
    - elbow/forearm/shoulder

  - wrist and hand
  - ultrasound
  - arthrograms (shoulder, elbow, wrist)
  - CT scanning
  - technetium bone scan
- nerve conduction studies
- electromyography

Note: Modern sophisticated ultrasound exam-ination is becoming a vital diagnostic modality for soft tissue disorders.

ARM PAIN IN CHILDREN

The main concerns with children are the effects of trauma, especially around the elbow. Considerable awareness of potential problems and skilful management are required with children's elbow fractures. Foreign bodies in the arm also have to be considered.

Pulled elbow

This typically occurs in children under 8 years of age, usually at 2 - 5 years, when an adult applies sudden traction to the child's extended and pronated arm (Figure 64.2a): the head of the radius can be pulled distally through the annular radioulnar ligament (Figure 64.2b).[5]



Figure 64.2 Pulled elbow: (a) mechanism of injury; (b) annular ligament displaced over head of radius; (c) reduction by supination

Symptoms and signs

- The crying child refuses to use the arm.
- The arm is limp by the side or supported in the child's lap.
- The elbow is flexed slightly (any flexion will be strenuously resisted).
- The forearm is pronated or held in mid-position.
- The arm is tender around the elbow (without bruising or deformity).

Note: An X-ray is not necessary.

Treatment method

1. Gain the child's confidence.
2. The child stands facing the doctor with the parent holding the non-affected arm.
3. Place one hand around the child's elbow to give support, pressing the thumb over the head of the radius.
4. With the other hand, firmly and smoothly flex the elbow and suddenly and firmly twist the forearm into full supination (Figure 64.2c). A faint click (which will be painful) will be heard. After a few minutes the child will settle and resume full pain-free movement. Warn parents that recurrences are possible up to 6 years.

Alternative method

An easier method for the child is to very gently alternate prouation and supination through a small arc as you flex the elbow.

Note: Spontaneous resolution can occur event-usually. Place the arm in a sling if necessary. If you cannot get child's co-operation send them home in a ‘high’ sling.

Fractures and avulsion injuries around the elbow joint, which are a major problem in children, are discussed in more detail in Chapter 135.[6]

ARM PAIN IN THE ELDERLY

Elderly patients are more likely to be affected by problems such as referred pain, radiculopathy or myelopathy from cervical spondylosis, tumours, polymyalgia rheumatica, entrapment neuropathies such as CTS and ulnar nerve entrapment. The latter can be related to trauma such as Colles' fractures. In addition the elderly are more prone to suffer from the thoracic outlet syndrome as previously described under ‘Pitfalls’. Osteoarthritis of the hand and tenosynovitis, such as trigger thumb or finger, are more common with advancing age.

TENNIS ELBOW

Tennis elbow is caused by overuse or overload of the muscles of the forearm, especially in the middle aged. Two types are identifiable: ‘backhand’ tennis elbow, or lateral epi-condylitis, and ‘forehand’ tennis elbow, or medial epicondylitis, which is known also as golfer's or pitcher's elbow. ‘Backhand’ tennis elbow, which will be termed lateral tennis elbow, is the common classic variety. It is an overload injury caused by excessive strain on the extensor muscles of the forearm resulting from wrist extension. Both conditions are generally self limiting, but symptoms can persist for up to 2 years, or even much longer.

Lateral tennis elbow (lateral epicondylitis)

The patient who presents with this common and refractory problem is usually middle aged and only about one in 20 plays tennis. A typical clinical profile is presented in Table 64.2.

Table 64.2 Lateral tennis elbow: typical clinical profile

Age:	40 - 60 years
Occupation:	Carpenter, bricklayer, housewife, gardener, dentist, violinist
Sport:	Tennis, squash
Symptoms:	• pain at outer elbow, referred down back of forearm • rest pain and night pain (severe cases) • pain in the elbow during hand movements (e.g. turning on taps, turning door handles, picking up objects with grasping action, carrying buckets, pouring tea, shaking hands)

Signs:	• no visible swelling • localised tenderness over lateral epicondyle, anteriorly • pain on passive stretching wrist • pain on resisted extension wrist • normal elbow movement
Course:	6 to 24 months
Management:	Basic • rest from offending activity • RICE[*] and oral NSAIDs if acute • exercises - stretching and strengthening Additional (if refractory) • corticosteroid/LA injection (max. two) • manipulation • surgery

[*] RICE: rest, ice, compression, elevation

Signs

On examination the elbow looks normal, and flexion and extension are painless.

There are three important positive physical signs:

1. localised tenderness to palpation over the anterior aspect of the lateral epicondyle
2. pain on passive stretching at the wrist with the elbow held in extension and the forearm prone (Figure 64.3)



Figure 64.3 Lateral tennis elbow test: reproducing pain on passive stretching at the wrist

3. pain on resisted extension of the wrist with the elbow held in extension and the forearm prone (Figure 64.4).

Management

Although there are a myriad of treatments the cornerstones of therapy are rest from the offending activity and exercises to strengthen the extensors of the wrist. The application of ice may help relieve discomfort of acute pain. Three systematic reviews have found little evidence for efficacy of any one specific intervention but short-term use of NSAIDs and progressive strengthening and stretching exercises were better than placebo.[7] A trial of NSAIDs oral or topical NSAID applied this may be worthwhile.[8]

Exercises

Stretching and strengthening exercises for the forearm muscles represent the best management for tennis elbow. Three options are presented.

1. The wringing exercise. Chronic tennis elbow can be cured by a simple wringing exercise using a small hand towel.[9]

   Method

   - Roll up the hand towel.
   - With the arm extended, grasp the towel with the affected side placed in neutral.
   - Then exert maximum wring pressure:
     - first flexing the wrist for 10 seconds
     - then extending the wrist for 10 seconds

   This is an isometric 'hold' contraction.

   This exercise should be performed only twice a day, initially for 10 seconds in each direction. After each week increase the time by 5 seconds in each twisting direction until 60 seconds is reached (week 11). This level is maintained indefinitely.

   Note: Despite severe initial pain, the patient must persist, using as much force as possible. Review at 6 weeks to check progress and method.

2. 'Weights' exercise. The muscles are strengthened by the use of hand-held weights or dumbbells. A suitable starting weight is 0.5 kg, building up gradually (increasing by 0.5 kg) to 5 kg, depending on the patient.

   Method

   - To perform this exercise the patient sits in a chair beside a table.
   - The arm is rested on the table so that the wrist extends over the edge.
   - The weight is grasped with the palm facing downwards (Figure 64.5).
   - The weight is slowly raised and lowered by flexing and extending the wrist.
   - The flexion/extension wrist movement is repeated 10 times, with a rest for 1 minute, and the program is repeated twice.

3. The pronating exercise.[10] A suitable stretching exercise is to rhythmically rotate the hand and wrist inwards with the elbow extended and the forearm pronated (Figure 64.6). Another proven exercise program is that outlined by Nirschl[11] and this can be provided by referral to a physiotherapist familiar with the program.



Figure 64.5 Lateral tennis elbow: the dumbbell exercise with the palm facing down



Figure 64.6 Tennis elbow stretching exercise: the hand and wrist are rhythmically rotated inwards until the painful point is reduced

Injection therapy

The injection of 1 mL of corticosteroid and 1 mL of local anaesthetic should be reserved for those severe cases when pain restricts simple daily activities, and not used initially for those patients with only intermittent pain. The key to a successful injection is to have the tender lesions pinpointed precisely. The point of maximal tenderness is usually on or just distal to the lateral epicondyle. Relief seems to last for 2 to 6 weeks (Figure 64.7). Some therapists use local anaesthetic only.


vertical approach
for injection
supinated forearm
tender lateral
epicondyle
thumb stretches skin
around epicondyle

Figure 64.7 Lateral tennis elbow: injection technique

A Netherlands study showed that corti-costeroid injections are the best short-term treatment for tennis elbow. Over the longer term physiotherapy offers better results than injection but is on a par with a wait and see approach.[12]

Surgery

Severe and refractory cases can be referred for surgery but this is rarely indicated and there is no evidence to date on its efficacy. The usual procedure is the stripping of the common extensor origin combined with debridement of any granulation tissue.[3]

Medial tennis elbow (medial epicondylitis)

In 'forehand' tennis elbow, or golfer's elbow, the lesion is the common flexor tendon at the medial epicondyle. The pain is felt on the inner side of the elbow and does not radiate far. The main signs are localised tenderness to palpation and pain or resisted flexion of the wrist.

In tennis players it is caused by stroking the ball with a bent forearm action or using a lot of top spin, rather than stroking the ball with the arm extended.

The treatment is similar to that for lateral epicondylitis except that in a dumbbell exercise program that palm must face upwards.

A similar injection method is used to that for lateral epicondylitis. The elbow is flexed and supinated with full external rotation of the shoulder of the affected arm. The anterior approach is used, and the tender area of the medial epicondyle injected as for lateral epicondylitis.

After-care and prevention (both types)

Tennis should be resumed gradually. Players recovering from tennis elbow should start quietly with a warm-up period and obtain advice on style, including smooth stroke play. During a game they should avoid elbow bending and 'wristy' shots. A change to a good quality racket (wooden or graphite frame) with a medium-sized head and suitable grip size may be appropriate.[11] The patient should be advised not to use a tightly strung, heavy racket or heavy tennis balls. It may be worthwhile to advise the use of a non-stretch band or brace situated about 7.5 cm (3 in) below the elbow.

OLECRANON BURSITIS

Olecranon bursitis presents as a swelling localised to the bursa (which has a synovial membrane) over the olecranon process. The condition may be caused by trauma, arthritic conditions (rheumatoid arthritis and gout) or infection.

Traumatic bursitis may be caused by a direct injury to the elbow or by chronic friction and pressure as occurs in miners (beat elbow), truck drivers or carpet layers. Acute olecranon bursitis with redness and warmth can occur in rheumatoid arthritis, gout, pseudogout, haemorrhage and infection (sepsis).[13] Septic bursitis must be considered where the problem is acute or subacute in onset, and hence aspiration of the bursa contents with appropriate laboratory examination is necessary (smear, Gram stain, culture and crystal examination). Treatment depends on the cause.

Simple aspiration/injection technique

Chronic recurrent traumatic olecranon bursitis with a synovial effusion may require surgery but most cases can resolve with partial aspiration of the fluid and then injection of corticosteroid through the same needle. Sepsis must be ruled out.

OVERUSE SYNDROMES OF FOREARM MUSCLES[8]

Pain is often experienced in the belly of a muscle, such as the flexors and extensors, following unaccustomed use of the wrists and elbows. There is pain on contraction and stretching of the muscles and tenderness on palpation. This problem can be limiting for a significant period. Early treatment includes relative rest, ice packs, analgesics (paracetamol) and gradual return to activity. Referral for physiotherapy to supervise rehabilitation is important.

CARPAL TUNNEL SYNDROME

Patients with CTS complain of ‘pins and needles’ affecting the pulps of the thumb, and index, middle and half of the ring finger (Figure 64.8). They usually notice these symptoms after, rather than during, rapid use of the hands. They may also complain of pain, which may even radiate proximally as far as the shoulder, from the volar aspect of the wrist. Causes or associations of CTS are presented in Table 64.3.

Table 64.3 Carpal tunnel syndrome: causes or associations

Idiopathic
Trauma
Fibrosis
Granulomatous disorders (TB, etc.)
Rheumatoid arthritis
Acromegaly
Amyloidosis
Pregnancy
Premenstrual oedema
Hypothyroidism
Paget's disease
Diabetes mellitus
Tophaceous gout
Consider cranial nerve root pressure

The pathognomonic symptom

Patients complain of awakening from their sleep at night with ‘pins and needles’ affecting the fingers. They get out of bed, shake their hands, the ‘pins and needles’ subside and they return to sleep. In severe cases, the patient may awaken two or three times a night and go through the same routine.



Figure 64.8 Carpal tunnel syndrome (median nerve compression syndrome)

Work-related CTS

CTS is seen in many work situations requiring rapid finger and wrist motion under load, such as meat workers and process workers. A type of flexor tenosynovitis develops and thus nerve compression in the tight tunnel. It is advisable to arrange confirmatory investigations by nerve conduction studies and electromyography for this work-induced overuse disorder. This testing is also indicated where the diagnosis is uncertain or if the condition persists and numbness or weakness develops.

Simple clinical tests

In the physical examination a couple of simple tests can assist with confirming the diagnosis. These are Tinel's test and Phalen's test. However, they are 'soft' signs with a relatively low sensitivity and specificity.[14]

The Tinel test

- Hold the wrist in a neutral or flexed position and tap over the median nerve at the flexor surface of the wrist. This should be over the retinaculum just lateral to the palmaris longus tendon (if present) and the tendons of flexor digitorium superficialis (Figure 64.9).



Figure 64.9 Carpal tunnel syndrome: Tinel sign

- A positive Tinel's sign produces a tingling sensation (usually without pain) in the distribution of the median nerve.

The Phalen test

- The patient approximates the dorsum of both hands, one to the other, with wrists maximally flexed and fingers pointing downwards.
- This position is held for 60 seconds.
- A positive test reproduces tingling and numbness along the distribution of the median nerve.

Two point discrimination

The test that has the highest specificity of all basic clinical tests is two point discrimination, but it has low sensitivity for CTS.[14]

Treatment

The treatment is determined by the severity. For mild cases simple rest and splinting (particularly at night) is sufficient. Carpal tunnel corticosteroid infiltration is frequently of diagnostic as well as therapeutic value. Ultrasound therapy has been used with some success. Surgical release (flexor retinaculotomy) is necessary for patients with sensory or motor deficits and those with recalcitrant CTS.

Systematic evidence-based reviews indicate the benefit of short-term oral corticosteroids[7] and local corticosteroid injection (short-term). NSAIDs, diuretics and wrist splinting are unlikely to be beneficial.

In reference to surgery, one review found similar clinical outcomes between open carpal tunnel release and endoscopic release but the latter had more complications.[7]

Injection into the carpal tunnel

Injections may relieve symptoms permanently or, more commonly, temporarily. The injections may be repeated. Do not use local anaesthetic in the injection.

Method

- The patient sits by the side of the doctor with the hand palm upwards, the wrist slightly extended.
- Identify the palmaris longus tendon and ulnar artery.
- Insert the needle (23 gauge) at a point about 2 - 2.5 cm proximal to the main transverse crease of the wrist and between the palmaris longus tendon and the artery (Figure 64.10). Take care to avoid the superficial veins.



Figure 64.10 Injection technique for carpal tunnel syndrome

- Advance the needle distally, parallel to the tendons and nerve at about 5? to the horizontal. It should pass under the transverse carpal ligament (flexor retinaculum) and come to lie in the carpal tunnel.
- Inject 1 mL of corticosteroid. This is usually painless and runs freely. Ensure the patient feels no severe pain or paraesthesia during the injection.
- Withdraw the needle and ask the patient to flex and extend the fingers for 2 minutes.

TRIGGER FINGER/THUMB (FIEXOR TENOSYNOVITIS)

In the fingers the common work-induced condition is stenosing flexor tenosynovitis, also known as trigger thumb and finger. Trigger finger of thumb has a reported lifetime risk of 2.6% in the population and is more common 15 in the 5th and 6th decades of life. It is associated with types diabetes, rheumatoid arthritis, gout, hypothyroidism and amyloidosis. It is caused by the same mechanism as de Quervain's stenosing tenosynovitis. In middle age these tendons, which are rapidly and constantly being flexed and extended, can undergo attrition wear and tear, and fibrillate and fragment; this causes swelling, oedema and painful inflammation and the formation of a nodule on the tendon that triggers back and forth across the thick, sharp edge of the ‘pulley’ (of the fibrosseous tunnel in the finger) (Figure 64.11).



Figure 64.11 Trigger thumb

These patients may present with a finger locked in the palm of the hand; the finger can only be extended passively (manually) with the other hand. It is easily diagnosed by triggering. If the pulp of the finger is placed over the 'pulley' crepitus can be felt and tenderness elicited. The thumb and fourth (ring) finger are commonly affected, at the level of the metacarpal head.

Treatment

Although surgery is simple and effective, treatment by injection is often very successful. The injection is made under the tendon sheath and not into the tendon or its nodular swelling. Controlled trials report a success rate of up to 70%.

Method

- The patient sits facing the doctor with the palm of the affected hand facing upwards.
- Draw 1 mL of long-acting corticosteroid solution into a syringe and attach a 25-gauge needle for the injection.
- Insert the needle at an angle distal to the nodule and direct it proximally within the tendon sheath (Figure 64.12). This requires tension on the skin with free fingers.



Figure 64.12 Injection site for trigger finger

- By palpating the tendon sheath, you can (usually) feel when the fluid has entered the tendon sheath.
- Inject 0.5 - 1 mL of the solution, withdraw the needle and ask the patient to exercise the fingers for 1 minute.

Postinjection

Improvement usually occurs after 48 hours and may be permanent. The injection can be repeated after 3 weeks if the triggering is not completely relieved. If triggering recurs, surgery is indicated. This involves division of the thickened tendon sheath only.

DE QUERVAIN'S TENOSYNOVITIS (WASHERWOMAN'S SPRAIN)

At the wrist a not uncommon work-induced condition is de Quervain's stenosing tenosynovitis of the first dorsal extensor compartment tendons (extensor pollicis brevis and abductor pollicis longus), which pass along the radial border of the wrist to the base of the thumb. It is usually seen when the patient is required to engage in rapid, repetitious movements of the thumb and the wrist, especially for the first time, and thus is common in assembly line workers, such as staple gun operators.

Clinical features

- Typical age 40 - 50 years
- Pain at and proximal to wrist on radial border
- Pain during pinch grasping
- Pain on thumb and wrist movement
- Dull ache or severe pain (acute flare-up)
- Can be disabling with inability to use hand (e.g. writing)

Triad of diagnostic signs

- Tenderness to palpation over and just proximal to radial styloid
- Firm tender localised swelling in area of radial styloid (may be mistaken for exostosis)

- Positive Finklestein's sign (the pathognomonic test)

Finkelstein's test

- The patient folds the thumb into the palm with the fingers of the involved hand folded over the thumb, thus making a fist.
- Rotate the wrist in an ulnar direction to stretch the involved tendons as you stabilise the forearm with the other hand (Figure 64.13).



Figure 64.13 Finkelstein's test

- A positive test is indicated by reproduction of or increased pain.

Treatment

- Rest and avoid the causative stresses and strains on the thumb abductors.
- Use a custom-made splint that involves the thumb and immobilises the wrist.
- Consider of trial of oral NSAIDs for 14-21 days.
- Local long-acting corticosteroid injection can relieve and may even cure the problem but care should be taken to inject the suspension within the tendon sheath rather than into the tendon.
- Surgical release is required for chronic cases.

Method of tendon sheath injection

- Identify and mark the most tender site of the tendon and the line of the tendon. Identify and avoid the radial artery.
- Thoroughly cleanse the skin with an antiseptic such as povidone iodine 10% solution.
- Insert the tip of the needle (23 gauge) about 1 cm distal to the point of maximal tenderness (Figure 64.14).


maximal tenderness over abductor pollicis longus tendon
injection site 1 cm distal
wrist crease

Figure 64.14 Tendon sheath injection

- Advance the needle almost parallel to the skin along the line of the tendon.
- Inject about 0.5 mL of the corticosteroid suspension within the tendon sheath. If the needle is in the sheath very little resistance to the plunger should be felt, and the injection causes the tendon sheath to billow out.

TENDINITIS/TENOSYNOVITIS

After excluding CTS, trigger thumb/finger, de Quervain's tenosynovitis, rheumatoid and related disease, tendinitis is uncommon in the hand.[16] Tendinitis may occur in other extensor compartments of the wrist and hand with unusual repetitive stressful actions, such as power drills jamming, and in conveyer quality control where an object is picked up with the forearm prone, supinating to examine it and pronating to replace it.

Treatment is rest from the provoking activity, splintage and tendon sheath injection with long-acting corticosteroid in a manner similar to that described for de Quervain's tenosynovitis.

INTERSECTION SYNDROME

Intersection syndrome is caused by a bursitis that develops at the site where the extensor pollicis brevis and abductor pollicis longus tendors cross over the extensor carpi radialis tendons figure 64.15. The bursitis is due to friction at the crossing point or due to tenosynovitis of the extensor tendons on palpation tenderness is found dorsally on the radial side with swelling and refutes. Treatment is based on relative test, a trial of NSAIDs and an infection of local anaesthetic and corticosteroid into the bursa.

POST-TRAUMATIC CHRONIC WRIST PAIN[10]

Patients often present with persistent wrist pain following trauma, such as a fracture, sprain to the wrist or even a seemingly mild strain, such as falling down with the wrist flexed into the hand. An undiagnosed fracture, ischaemic necrosis or unstable ligamentous injury should be investigated by radiology or referral where appropriate. Look for a scapholunate ligament tear with tenderness 2 cm distal to the tubercle on the radial side of the lunate. For persistent tenderness an injection of corticosteroid and local anaesthetic into the tender site is advisable.[8]

ISCHAEMIC NECROSIS

Ischaemic necrosis particularly of the scaphoid can occur following failure to recognise a fracture. Tenderness in the ‘anatomical snuff box’ following trauma should be treated as a scaphved until repeated X-rays prove negative. In children, chronic pain in the region of the lunate suggests avascular necrous-Keinbocks' disease, presenting with dorsal wist pain. (p. 692).

GANGLIA

About 60-70% of these common soft tissue tumour occur on the dorsal aspect of the wrist. The vast majority arise from the dorsal scapholunate ligameal. Pain can result from compression on an adjacent nerve or joint space. If the diagnosis is uncertain an ultrasound scan (or even an MRI) may pinpoint the tumour see page 1212 for treatment.


extensor carpi radialis longus
extensor carpi radialis brevis
abductor pollicis longus
extensor pollicis brevis

Figure 64.15 Intersection syndrome: pain is present over the intersection of tendons

NEUROVASCULAR DISORDERS OF THE HAND

Painful vascular disorders, which are more likely to occur in women in cold weather, include Raynaud's phenomenon, eryth-ermalgia, chilblains and acute blue fingers syndrome. Acrocyanosis is not a painful condition.

Raynaud's phenomenon

The basic feature of Raynaud's phenomenon, which is a vasospastic disorder, is sequential discolouration of the digits from pallor to cyanosis to rubor upon exposure to cold and other factors (a useful mnemonic is WBR, namely white ® blue ® red). (Figure 64.16) The rubor is a reactive hyperaemia when fingers become red and tender. Associated symptoms are pain, tingling and numbness. It is possible to get loss of tissue pulp at the ends of the fingers and subsequent necrotic ulcers. The benign form is the commonest, but may indicate an evolving connective tissue disorder. It is highly significant if it extends to the MCP joints.



Figure 64.16 Raynavd's phenomenon: symptoms and colour changes of fingers with cold

Causes

primary

- Raynaud's disease (idiopathic)

secondary

- Occupational trauma (vibrating machinery)
- Connective tissue disorders
  - rheumatoid arthritis
  - SLE
  - systemic sclerosis, CREST
  - polyarteritis nodosa
- Arterial disease
  - Buerger's disease
- Haematological disorders
  - polycythaemia
  - cold agglutinin disease
  - leukaemia
- Drugs
  - beta-blockers
  - sympathomimetic drugs with α receptor activities
  - ergotamine
  - nasal decongestants

Aggravating factors

- Smoking
- Cold wet weather

- Stress or emotional upset

Differential diagnosis

- Chilblains-itchy, patchy discoloration without pallor
- Diffusely cold mottled hands-recover quickly on warming.

Investigations

- Exclude underlying causes with appropriate tests.

Treatment

- In an attack it is best to warm the extreauties gradually.
- Total body protection from cold wear layered clothing to present heat lm.
- Use an electric blanket at night, as required.
- Use mittens, fleece-lined glooves and thick woollen socks.
- Gloves or mittens should be warm when handling cold surfaces and objects, such as frozen food.
- Avoid smoking.
- Consider sympathectomy.

Vaso dilators (during cold weather)

- topical glyceryl trinale 2% ointment-applied to the base of the affected fingers two or four times daily or applied over the radial artery or dorsum of the hand or
- amlodepine 5-20 mg(o) once daily or
- nifedipine SR 30-60 mg (o) once daily or
- diltiazem SR 180-240 mg (o) once daily

Erythermalgia

This condition is characterised by erythema (redness), a burning sensation and swelling of the hands (and feet) after exposure to heat and exercise. It may be primary or secondary to a disease such as diabetes, haematological disorders[13] and connective tissue disease. Treatment of primary erythermalgia includes trials of aspirin, phenoxybenzamine (Dibenyline), methysergide or sympathectomy.

Acute blue fingers in women syndrome

This unusual syndrome involves the sudden onset of pain and cyanosis of the ventral aspect of the digit initially, and then the entire digit. It lasts for two or three days and the attacks recur one or more times per year. No abnormalities are found on physical or on laboratory examination.

The cause is probably spontaneous rupture of a vein at the base of the finger.

Chilblains (perniosis)

Precautions

- Think Raynaud's phenomenon
- Protect from trauma and secondary infection
- Do not rub or massage injured tissues
- Do not apply heat or ice

Physical treatment

- Elevate affected part
- Warm gradually to room temperature

Drug treatment

- Apply glyceryl trinitrate vasodilator spray or ointment or patch (use plastic gloves and wash hands for ointment)

Other treatment

- Rum at night
- Nifedipine SR 30mg daily

REGIONAL PAIN SYNDROME

The hand can be affected by the complex regional pain syndrome, previously called reflex sympathetic dystrophy (RSD, also in this case Sudeck's atrophy). The patient presents with severe pain, swelling and disability of the hand. It may occur spontaneously or, more usually, it follows trauma that may even be trivial. It can occur after a Colles' fracture, especially with prolonged immobilisation.

Clinical features

- Throbbing burning pain, worse at night
- Paraesthesia
- Initial: red, swollen hand; warm, dry skin
- Later: cold, cyanosed and mottled, moist skin; shiny and stiff fingers
- Wasting of small muscles
- X-rays - patchy decalcification of bone (diagnostic)

The problem eventually settles but may take years. Patients need considerable support, encouragement, basic pain relief, mobility in preference to rest and perhaps referral to a pain clinic.

KIENBÖCK'S DISORDER

Kienböck's disorder is avascular necrosis of the carpal lunate bone (Figure 64.17), which may fragment and collapse, eventually leading to osteoarthritis of the wrist.

It presents usually in young adults over the age of 15 as insidious, progressive wrist pain and stiffness that limits grip strength and hand function. Males are affected more often than females and the right hand more than the left, indicating the relationship to trauma.

ARTHRITIC CONDITIONS OF THE WRIST AND HAND

Arthritis of the hand is an inappropriate diagnosis and specificity is required to highlight the various joints that are the targets of the specific arthritides, which include osteo-arthritis, rheumatoid arthritis, spondylo-arthropathies, gout, haemochromatosis, and connective tissue disorders. Typical target areas in the hand are shown in Figure 64.17.


spondyloarthropathies (e.g. psoriasis) affect mainly DIP joints
Heberden's nodes
Bouchard's nodes
SLE and viral polyarthritis target mainly PIP joints
haemochr omatosis targets 2nd and 3rd MCP joints
H
H
RA tends to spare 4th MCP joint
OA affects 1st CMC joint
avascular necrosis lunate (Kienböck's disorder)
Targets of Rheumatoid arthritis
Osteoarthritis
major
lesser
major
lesser

Figure 64.17 Typical sites of arthritic conditions and osteochondritis in the hand

Osteoarthritis

Osteoarthritis commonly involves the interphalangeal joints of the fingers (especially the DIP joints)[17] and the carpometacarpal joint of the thumb. Degenerative changes produce bony swellings around the margins of the joints - Heberden's nodes of the DIP joints and less commonly Bouchard's nodes of the PIP joints. A patchy distribution occurs in meta-carpophalangeal, intercarpal and wrist joints, usually related to trauma.

Rheumatoid arthritis

In rheumatoid arthritis the DIP joints are often spared (only about 30% involved) but the metacarpophalangeal and proximal inter-phalangeal joints and wrist joints are generally affected symmetrically and bilaterally. Rheumatoid arthritis tends to affect the metacarpophalangeal joints of the fourth finger less commonly.

Gout

Gout may involve normal joints of the hand but is encountered more frequently in osteoarthritic joints of the hand (especially DIP joints) in elderly people taking diuretics. This clinical feature is known as nodular gout.

Seronegative arthropathies

A similar appearance to rheumatoid arthritis occurs except that with psoriatic arthritis the terminal joints are often involved with swelling, giving the appearance of ‘sausage digits’.

INFECTIONS OF THE HAND

Although not encountered as frequently as in the past, serious suppurative infections of the deep fascial spaces of the hand and tendon sheath can still occur, especially with penetrating injuries and web space infection.

Infections of the hand include:

- infected wounds with superficial cellulitis or lymphangitis (Streptococcus pyogenes)

- subcutaneous tissues
    - nail bed (paronychia)
    - pulp (whitlow, e.g. herpes simplex)
- erysipeloid - this is a specific infection in one finger of fishers or meat handlers, caused by Erysipelothrix insidiosa. There is a purplish erythema that gradually extends over days. It is rapidly cured by penicillin.
- tendon sheath infection (suppurative tenosynovitis) - this is a dangerous and painful infection that can cause synovial adhesions with severe residual finger stiffness. The affected finger is hot and swollen and looks like a sausage.
- deep palmar fascial space infection - infection from an infected tendon sheath or web space may spread to one of the two deep palmar spaces: the medial (midpalmar space) or lateral (thenar) space.
- sporotrichosis (gardener's arm) - a chronic fungal infection from contaminated spikes of wood or rose thorns presenting as hard non-tender nodules in the skin of the hand and extending along the lymphatics of the arm. The diagnosis is confirmed by biopsy. Treat with itraconazole.

Management of serious infection

- Early appropriate antibiotic treatment for infection and early surgical referral where necessary.
- Antibiotics (adult doses)[19]
    - Streptococcus pyogenes (mild to moderate cellulitis, lymphangitis) procaine penicillin 1.5 g IM daily or phenoxymethyl penicillin 500 mg (o) 6 hourly
    - If server to cover both *strep: pyogenes* and *Staphylococcus aureus* infection (suspected or proved) flucloxacillin/dicloxacillin 2g IV 6 hourly. until improved, then oral for 10 days.

‘CRACKED’ HANDS AND FINGERS

- Wear protective work gloves: cotton-lined PVC gloves.
- Use soap substitutes (e.g. Cetaphil lotion, Dove).
- Apply 2 - 5% salicylic acid and 10% liq picis carb in white soft paraffin ointment. or
- Corticosteroid ointment: class II - III.

WHEN TO REFER

- Disabling osteoarthritis of carpometacarpal joint for possible surgical repair
- Myelopathy (motor weakness) and persistent radiculotherapy (nerve root pain and sensory changes) in the arm
- Unresolving nerve entrapment problems such as median and ulnar nerves
- Elbow injuries in children with proven or possible supracondylar fracture or avulsion epicondylar fractures
- Evidence or suspicion of suppurative infection of the tendon sheaths or deep palmar fascial spaces
- Septic arthritis and osteomyelitis
- Regional pain syndrome
- Other conditions not responding to conservative measures

Practice tips

- With elbow injuries in children, X-ray both elbows and compare one side with the other; this helps to determine whether there is displacement of fragments or a disturbance in the normal anatomy of the elbow.
- Tendinitis and other entheseal problems of the arm are common and tend to take 1 - 2 years to resolve spontaneously, yet they resolve rapidly with rest, an exercise program or corticosteroid injections. Surgical relief is effective for refractory cases.
- The so-called thoracic outlet syndrome is probably most often caused by ‘the droopy shoulder

syndrome' rather than by a cervical rib.

- Consider corticosteroid injections for the CTS and stenosing tenosynovitis (de Quervain's and trigger finger or thumb). They are very effective and often curative.
- The site of arthritis in the hand provides a reasonable guide as to the cause.
- Always keep regional pain syndrome in mind for persistent burning pain in the hand following injury - trivial or severe.

REFERENCES

1. Swift TR, Nichols FT. The droopy shoulder syndrome. Neurology, 1984; 34: 212 - 15.
2. Bertelsen S. Neurovascular compression syndromes of the neck and shoulder. Acta Chir Scand, 1969; 135: 137 - 48.
3. Ireland D. The hand (part two). Aust Fam Physician, 1986; 15: 1502 - 13.
4. Dan NG. Entrapment syndromes. Med J Aust, 1976; 1: 28 - 31.
5. Corrigan B, Maitland GP. Practical Orthopaedic Medicine. Sydney: Butterworths, 1986: 75 - 7.
6. Young D, Murtagh J. Pitfalls in orthopaedics. Aust Fam Physician, 1989; 18: 645 - 53.
7. Barton S ed. Clinical Evidence. London: BMJ Publishing Group, 2001: 717 - 27.
8. Mashford ML (Chair). Therapeutic Guidelines: Analgesic (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2002: 143 - 6.
9. White ADN. Practice tip. A simple cure for chronic tennis elbow. Aust Fam Physician, 1987; 16: 953.
10. Oakes B, Fuller P, Kenihan M, Sandor S. Sports Injuries. Melbourne: Pitman, 1985: 51 - 5.
11. Brinbaum AJ. Tennis elbow: don't worry, it can be avoided and it can be cured. Tennis, 1978; April: 96 - 103.
12. Smidt N et al. Corticosteroid injections, physiotherapy or a wait and see policy for lateral epicondylitis: a randomised controlled trial. Lancet, 2002; 359: 657 - 62.
13. Sheon R, Moskowitz R, Goldberg V. Soft Tissue Rheumatic Pain. Philadelphia: Lea & Febiger, 1987: 134 - 40.
14. Buch-Jaeger N, Foucher G. Carpal tunnel syndrome, Hand Surgery, 1994; 19B: 72 - 4.
15. Mashford ML (Chair), Therapeutic Guidelines: Rheumatology (version 1) Thelbourne: Therapeutic Guidelines Limited, 2006.
16. Ireland D. The hand (part one). Aust Fam Physician, 1986; 15: 1162 - 71.
17. Corrigan B, Maitland G. Practical Orthopaedic Medicine. Sydney: Butterworths, 1986: 97 - 100.
18. Spicer V. Therapeutic Guidelines: Antibiot ic (version12). Melbourne: Therapeutic Guidelines Limited 2003:218 - 25.
19. Spicer J. *Therapeutic Guidelines: Antibiotic* (Version 12). Melbourne: Therapeutic Guidelines Limited, 2003: 218 - 25.

CHAPTER 65. HIP, BUTTOCK AND GROIN PAIN

Which of your hips has the most profound sciatica?

WILLIAM SHAKESPEARE (1564 - 1616), *MEASURE FOR MEASURE*

HIP, BUTTOCK AND GROIN PAIN

Pain in the hip, buttock, groin and upper thigh tend to be interrelated. Patients often present complaining of pain in the hip but pain to in the buttock or lower back as the site of their pain. Most pain in the buttock has a lumbosacral origin. Pain originating from disorders of the lumbosacral spine (commonly) and the knee (uncommonly) can be referred to the hip region, while pain from the hip joint (L3 innervation) may be referred commonly to the thigh and the knee. Disorders of the abdomen, retroperitoneal region and pelvis may cause hip and groin pain, sometimes mediated by irritation of the psoas muscle.

KEY FACTS AND CHECKPOINTS

- Hip troubles have a significant age relationship (Figure 65.1).



Figure 65.1 Typical ages of presentation of hip disorders

- Children can suffer from a variety of serious disorders of the hip such as developmental dysplasia of the hip (DDH), Perthes' disease, tuberculosis, septic arthritis slipped capital femoral epiphysis (SCFE), and inflammation arthritis all of which demand early recognition and management.
- SCFE typically presents in the obese adolescent (10 - 15 years) with knee pain and a slight limp.
- Every newborn infant should be tested for DDH, which can usually be treated successfully when diagnosed early.
- Limp has an inseparable relationship with painful hip and buttock conditions, especially those of the hip.
- The spine is the most likely cause of pain in the buttock in adults.
- Disorders of the hip joint commonly refer pain to the knee and thigh.
- If a woman, especially one with many children, presents with bilateral buttock or hip pain, consider dysfunction of the sacroiliac joints (SIJs) as the cause.
- If a middle-aged or elderly woman presents with hip pain, always consider the underdiagnosed conditions of trochanteric bursitis or gluteus medius tendinitis.

A DIAGNOSTIC APPROACH

A summary of the diagnostic model is presented in Table 65.1.

Table 65.1 Hip and buttock pain: diagnostic strategy model

Q Probability diagnosis
A. Traumatic muscular strains Referred pain from spine Osteoarthritis of hip

Q. Serious disorders not to be missed

A. Cardiovascular

- buttock claudication

Neoplasia

- metastatic carcinoma

steoid osteoma

Septic infections

- septic arthritis
- osteomyelitis
- tuberculosis
- pelvic and abdominal infections
  - pelvic abscess
  - PID

Childhood disorders

- DDH
- Perthes' disease
- slipped femoral epiphysis
- synovitis (irritable hip)
- juvenile chronic arthritis

Q. Pitfalls (often missed)

A. Polymyalgia rheumatica

Fractures

- stress fractures femoral neck
- subcapital fractures
- sacrum

Pubic rami

Avascular necrosis femoral head

Sacroiliac joint disorders

Inguinal or femoral hernia

Bursitis or tendinitis

- gluteus medius tendinitis
- trochanteric bursitis
- ischial bursitis

Osteitis pubis

Neurogenic claudication

Chilblains

Rarities

- Haemarthrosis (e.g. haemophilia)
- Paget's disease

- Nerve entrapments
  - sciatica ‘hip pocket nerve’
  - obturator
  - lateral cutaneous nerve thigh

Q. Seven masquerades checklist

A. Depression

Diabetes -

Drugs -

Anaemia -

Thyroid disorder -

Spinal dysfunction

UTI -

Q. Is this patient trying to tell me something else?

A. Non-organic pain may be present. Patient with arthritis fearful of being crippled.

Probability diagnosis

The commonest cause of hip and buttock pain presenting in general practice is referred pain from the lumbosacral spine and the sacroiliac joints.[1] The pain is invariably referred to the outer buttock and posterior hip area (Figure 65.2). The origin of the pain can be the facet joints of the lumbar spine, intervertebral disc disruption or, less commonly, the SIJs. Much of this pain is inappropriately referred to as ‘lumbago’, ‘fibrositis’ and ‘rheumatism’.


cutaneous distribution of T12 and L1 (also referred to groin)
course of sciatic nerve
typical zone of referred pain from sacroiliac joint

Figure 65.2 Referral patterns of pain from the lumbosacral spine and the sacroiliac joints

Trauma and overuse injuries from sporting activities are also common causes of muscular and ligamentous strains[2] around the buttock hip and groin.

The hip joint is a common target of osteoarthritis. This usually presents after 50 years but can present earlier if the hip has been affected by another condition.

Serious disorders not to be missed

The major triad of serious disorders - cardiovascular, neoplasia and severe in-fections - are applicable to this area albeit limited in extent.

Aortoiliac occlusion

Ischaemic muscle pain, including buttock claudication secondary to aortoiliac arterial occlusion, is sometimes confused with musculoskeletal pain. An audible bruit over the vessels following exercise is one clue to diagnosis.

Neoplasia

Primary tumours including myeloma I lymphosarcoma sarcoma and, can arise rarely in the upper femur and pelvis (especially the ilium). However, these areas are relatively common targets for metastases, especially from the prostate breast and lung.

Infection

Some very important, at times ‘occult’, infections can develop in and around the hip joint.

Osteomyelitis is prone to develop in the metaphysis of the upper end of the femur and must be considered in the child with intense pain, a severe limp and fever. Tuberculosis may also present in children (usually under 10 years) with a presentation similar to Perthes' disease.

Transient synovitis or ‘irritable hip’ is the most common cause of hip pain and limp in childhood.

Inflammation of the side wall of the pelvis as in a deep pelvic abscess (e.g. from appendicitis), pelvic inflammatory disease (PID) including pyosalpinx, or an ischiorectal abscess can cause deep hipand groin pain and a limp. This pain may be related to irritation of the obturator nerve.

Retroperitoneal haematoma can cause referred pain and femoral nerve palsy.

Childhood disorders that must not be missed include:

- DDH and acetabular dysplasia
- Perthes' disease
- SCFE
- stress fractures of the femoral neck

Inflammatory disorders of the hip joint that should be kept in mind include:

- rheumatoid arthritis
- juvenile chronic arthritis (JCA)
- rheumatic fever (a flitting polyarthritis)
- spondyloarthropathy

‘Red flag’ pointers to potentially serious hip conditions

- Swelling, redness, very limited joint motion
- Pain, fever, systemic features (in absence of trauma)
- Neurological changes eg. loss of power
- Rapid joint swelling after trauma
- Constant localised pain unaffected by movement

Pitfalls

There are many pitfalls associated with hip and buttock pain and these include the various childhood problems. Fractures can be a pitfall, especially subcapital fractures.

SIJ disorders are often missed, whether it be the inflammation of sacroiliitis or mechanical dysfunction of the joint.

Inflammatory conditions around the hip girdle are common and so are often mis-diagnosed. These include the common gluteus medius tendinitis and trochanteric bursitis.

Polymyalgia rheumatica commonly causes shoulder girdle pain in the elderly but pain around the hip girdle can accompany this important problem.

Chilblains around the upper thighs occur in cold climates and are often known as ‘jodhpur’ chilblains because they tend to occur during horse riding in very cold weather.

Nerve entrapment syndromes require consideration. Meralgia paraesthetica is a nerve entrapment causing pain and paraesthesia over the lateral aspect of the hip (Figure 65.3).



Figure 65.3 ‘Hip pocket nerve’ syndrome: location and relations of sciatic nerve in the buttock

An interesting modern phenomenon is the so-called ‘hip pocket nerve’ syndrome. If a man presents with ‘sciatica’, especially confined to the buttock and upper posterior thigh (without local back pain), consider the possibility of pressure on the sciatic nerve from a wallet in the hip pocket. This problem is occasionally encountered in people sitting for long periods in cars (e.g. taxi drivers). It appears to be related to the increased presence of plastic credit cards in wallets (Figure 65.3).

Paget's disease can involve the upper end of the femur and the pelvis. Increased pain may indicate a fracture or malignant change causing osteosarcoma

General pitfalls

- Failure to test the hips of neonates carefully and follow up developmental dysplasia of the hip.
- Misdiagnosis of arthritis and other disorders of the hip joint because of referred pain.
- Overlooking an SCFE or a stress fracture of the femoral neck in teenage boys, especially athletes. If an X-ray shows that the epiphysis is fusing or has fused, arrange a technetium bone scan.

Seven masquerades checklist

The one outstanding masquerade is spinal dysfunction, which is the most likely cause of pain in the buttock. Many dermatomes meet at the buttock and theoretically pain in the buttock can result from any lesion situated in a tissue derived from L1, L2, L3, S2, S3 and S4.[1] Symptoms from L3 can spread from the outer buttock to the front of the thigh and down the leg, over the medial aspect of the knee to the calf. Such a distribution is common to an L3 nerve root lesion and arthritis of the hip.

Furthermore, dysfunction of the facet joints and sacroiliac joints can refer pain to the buttock. The relatively common L1 lesion due to dysfunction at the T12 - L1 spinal level can lead to referred pain over the outer upper quadrant of the buttock (Figure 65.2) and also to the groin (Figure 66.1).

Psychogenic considerations

Cyriax[1] claims that the hip shares with the back and the shoulder ‘an enhanced liability to fixation’ for psychological reasons. This problem is often related to work compensation factors and overpowering stresses at home. A common finding in psychoneurotic patients complaining of buttock and thigh pain is 90? limitation of flexion at the hip joint. The importance of testing passive movements of the hip joint is obvious, for often such limitation of flexion is combined with a full range of rotation. In arthritis of the hip joint internal rotation is invariably affected first.

Such patients often walk into the office with a marked limp and leaning on a thick stick. It requires great skill to evaluate and manage them tactfully and successfully.

On the other hand, patients with genuine osteoarthritis fear being crippled and ending in a wheelchair. They require considerable education and reassurance.

THE CLINICAL APPROACH

History

Pain associated with hip joint pathology is usually described as a deep aching pain, aggravated by movement[2] and felt in the groin and anteromedial aspect of the upper thigh, sometimes exclusively around the knee (Figure 67.1). A limp is a frequent association.

An obstetric history in a woman may be relevant for sacroiliac pain.

Key questions

- Can you tell me how the pain started?
- Could you describe the pain?
- Point to where the pain is exactly.
- Does the pain come on after walking for a while and stops as soon as you rest?
- Is there any stiffness, especially first thing in the morning?
- Do you have any difficulty climbing stairs?
- Do you get any backache?
- Do your movements feel free?
- Do you have a limp?
- Do you have a similar ache around the shoulders?
- Have you had an injury such as a fall?
- Have you lost any weight recently?
- Do you have night pain?
- Do you have trouble with shoes and socks?
- How far can you walk?

Physical examination

Follow the traditional methods of examination of any joint: look, feel, move, measure, test, function, look elsewhere and x-ray. The patient should strip down to the underpants to allow maximum exposure.

Inspection

Ask the patient to point exactly to the area of greatest discomfort. Careful observation of the patient, especially walking, provides useful diagnostic information. Note any antalgic (painful gast) If walking with a limp, the leg adducted and foot somewhat externally rotated, osteoarthritis of the hip joint is the likely diagnosis.

If called to a patient who has suffered an injury such as a fall or vehicle accident, note the position of the leg. If shortened and externally rotated (Figure 65.4a), a fractured neck of femur is the provisional diagnosis; if internally rotated, suspect a posterior dislocation of the hip (Figure 65.4b). With anterior dislocation of the hip, the hip is externally rotated.



Figure 65.4 (a) General configuration of the legs for a fractured neck of femur; **(b)** general configuration for a posterior dislocated hip

Get the patient to lie supine on the couch with the anterior superior iliac spines (ASISs) of the pelvis placed squarely and note the shape and position of the limbs.

Palpation

Feel one to two finger breadths below the midpoint of the inguinal ligament for joint tenderness. Check for trochanteric bursitis, gluteus medius tendinitis and other soft tissue problems over the most lateral bony aspect of the upper thigh.

Movements

- Passive movements with patient supine (normal range is indicated):
  - flexion with patients knees flexed (compare both sides): 140°
  - external rotation (knee and hip extended in adults): 45 - 50°
  - internal rotation (knee and hip extended in adults): 45°
  - abduction (stand on same side - steady pelvis): 45°
  - adduction (should see the patella of the opposite leg): 25°

  In children it is most important to measure rotation and abduction/adduction with the knee and hip flexed to detect early Perthes' disease or SCFE.
- Patient prone:
  - extension (one hand held over SIJ): 25°

  Note: Osteoarthritis of the hip affects internal rotation (IR), extension and abduction first.

Measurements

- True leg length (ASIS to medial malleolus)
- Apparent leg length (umbilicus to medial malleolus)

Note:

- Unequal true leg length ° hip disease on shorter side.
- Unequal apparent leg length ° tilting of pelvis.

Feel the height of the greater trochanters relative to the ASIS to determine if shortening is in the hip or below.

Test function and special tests

Gait:

- Trendelenburg test - tests hip abductors (gluteus medius)
- Thomas test - tests for fixed flexion deformity

Look elsewhere

Examine the lumbosacral spine, SIJs, groin and knee. Consider hernias and possibility of PID. Feel pulses and test for femoral bruits.

Investigations

These can be selected from:

- serological tests
    - RA factor
    - FBE
    - ESR
    - C-reactive protein
- radiological
    - plain AP X-ray of pelvis showing both hip joints
    - lateral X-ray (‘Frog’ lateral best in children)
    - X-ray of lumbosacral spine and SIJs
    - CT scan: hip joint, pelvis, lumbosacral spine
    - MRI scan
        - stress fracture
        - early avascular necrosis
        - early osteomyelitis
        - labial tears of hip joint
        - metastases
        - soft tissue tumours
    - isotope bone scan; useful for whole body bony metastaasses
- needle aspiration of joint: if septic arthritissuspected.

Role of ultrasound. Ultrasound diagnosis is now sensitive in children in detecting fluid in the hip joint, and can diagnose septic arthritis and also localise the site of an osteomyelitic abscess around a swollen joint. It can accurately assess the neonatal hip joint and confirm the position of the femoral head in children < 6 months. US used less since ↑

HIP PAIN IN CHILDREN

Hip disorders have an important place in childhood and may present with a limp when the child is walking. These important disorders include:

- DDH
- congenital subluxation of hip and acetabular dysplasia
- transient synovitis
- Perthes' disease

- septic arthritis/osteomyditis
- SCFE
- pathological fractures through bone cyst

The important features of hip pain in children are summarised in Table 65.2.

Table 65.2 Comparison of important causes of hip pain in children

	DDH	Transient synovitis	Perthes'	SCFE	Septic arthritis
Age (years)	0 - 4	4 - 8	4 - 8	10 - 15	Any
Limp	+	+	+	+	Won't walk
Pain	-	+	+	+	+ + +
Limited movement	Abduction	All, especially abduction and IR	Abduction and IR	All, especially IR	All
Plain X-ray	• No diagnostic value in neonatal period (use ultrasound)	• Normal	• Subchondral fracture • Dense head • Pebble stone epiphysis	• AP may be normal • Frog lateral view shows slip	Normal Use ultrasound

Developmental dysplasia of the hip

In DDH, previously known as congenital dislocation of the hip, the underdeveloped femoral head dislocates posteriorly and superiorly. DDH is described as transiently unstable or as a mild subluxation (1 in 80 hips at birth, which stabilise in a few days) and frankly dislocated (1 in 800 hips).[3]

Clinical features

- Females:males ° 6:1
- Asymmetry in 40%
- Bilateral in one-third
- Tight adductors and short leg evident
- Diagnosed early by Ortolani and Barlow tests (abnormal thud or clunk on abduction); test usually negative after 2 months
- Ultrasound excellent (especially up to 3 - 4 months) and more sensitive than clinical examination
- X-ray difficult to interpret up to 3 months, then helppful

Note:

1. When diagnosed and treated from birth it is possible to produce a normal joint after a few months in an abduction splint.
2. Every baby should be examined for DDH during the first day of life and before discharge from hospital and at the 6-weeks check.[3] The Ortolani and Barlow tests remain important means of detecting an unstable or dislocated hip, but ultrasound is becoming more important and is recommended for high-risk babies (e.g. breech, family history of DDH).

Screening examination

Carry out the examination on a large firm bench with the baby stripped. Relaxation is essential; give the baby a bottle if necessary. Be gentle and have warm hands.

With the legs extended any asymmetry of the legs or skin creases is noted.

- Hold the leg in the hand with knee flexed - thumb over groin (lesser trochanter) and middle finger over greater trochanter - see Figure 65.5.



Figure 65.5 Examination of the infant for DDH: demonstrating Ortolani's sign on left side

- Flex hips to about 90°, abduct to 45° (note any clunk).
- Gently rock the femur backwards and forwards on the pelvis by pressing forward with the middle finger and backwards with the thumb.

Note any jerk or clunk. If the femoral head displaces, there is dislocation. Plain X-ray has little or no place in the diagnosis of DDH in the neonatal period.[4] Early referral for treatment is essential. If not detected early the femoral head stays out of the acetabulum and after the age of 1 year the child may present with delay in walking or a limp. The diagnosis is then detected by X-ray.

Treatment (guidelines)

- DDH must be referred to a specialist[3]
- 0 - 6 months - Pavlik harness or abduction splint
- 3 - 18 months - reduction (closed or open) and cast (pelvic spica)
- > 18 months - open reduction and possible osteotomy

Note: Despite early treatment some cases progress to acetabular dysplasia (underdevelopment of the ‘roof’ of the hip joint) and to premature osteoarthritis. Thus a follow-up X-ray of the pelvis during teenage years should be considered for anyone with a history of DDH.

Perthes' disease

Perthes' disease results in the femoral head become agpartly or totally avascular (i.e. avascular necrosis).

Clinical features

- Males:females ° 4:1
- Usual age 4 - 8 years (rarely 2 - 18 years)
- Sometimes bilateral
- Presents as a limp and aching (hip or groin pain)
- May be knee pain
- ‘Irritable’ hip early
- Limited movement in abduction and IR

X-ray. Joint space appears increased and femoral head too lateral: typical changes of sclerosis, deformity and collapse of the femoral capital epiphysis may be delayed.

Management

- Refer urgently.(provide crutetes)
- Aim is to keep femoral head from becoming flat.
- Choice of treatment depends on severity of the condition and age of the patient.

If untreated, the femoral head usually becomes flat over some months, leading to eventual osteoarthritis. Some untreated cases of Perthes' disease heal and have a normal X-ray.

Transient synovitis

This common condition is also known as ‘irritable hip’ or observation hip[5] and is the consequence of a self-limiting synovial inflammation.

Clinical features

- Child aged 4 - 8 years
- Sudden onset of hip pain and a limp
- Child can usually walk (some may not)
- May be history of trauma or recent LIRTI
- Painful limitation of movements, especially abduction and rotation
- Blood tests and X-rays normal (may be soft tissue swelling); ESR may be mildly elevated
- Ultrasound shows fluid in the joint

Differential diagnosis. This includes septic arthritis, JCA, Perthes' disease.

Outcome. It settles to normal within 7 days, without sequelae.

Treatment. Refer early. Treatment is bed rest or the use of crutches and analgesics. Follow-up X-ray is needed in 4 to 6 months to exclude Perthes' disease. Aspiration under general anaesthetic may be needed to exclude septic arthritis.

Slipped capital femoral epiphysis

One problem of the displaced capital epiphysis of the femoral head is patients when some develop avascular necrosis despite expert treatment. Diagnosis of the condition before major slipping is important. This necessitates early consultation with the teenager ex-periencing hip or knee discomfort and then accurate interpretation of X-rays.

Clinical features

- Adolescent 10 - 15 years, often obese
- Most common in the oversized and undersexed (e.g. the heavy prepubertal boy)
- Bilateral in 20%
- Limp and irritability of hip on movement
- Anterior hip (groin) pain
- Knee pain
- Hip rotating into external rotation on flexion and often lies in external rotation (ER)
- Most movements restricted, especially IR

Any adolescent with a limp or knee pain should have X-rays (AP and frog view) of both hips (Figure 65.6). Otherwise, this important condition will be overlooked.



Figure 65.6 Appearance of a slipped capital femoral epiphysis. Note that, in the normal state, a line drawn along the superior surface of the femoral neck passes through the femoral head, but passes above it with SCFE

Management

- Cease weight-bearing and refer urgently to orthopaedic surgeon. Also refer of the clinical signs indicate SCFE but the X-rays look norn
- If acute slip, gentle reduction via traction is better than manipulation in preventing later avascular necrosis.
- Once reduced, pinning is performed.

Septic arthritis

Septic arthritis of the hip should be suspected in all children with acutely painful or irritable hip problems. These patients may not be obviously sick on presentation particulary in infacts < 2 years. A negative needle aspiration does not exclude septic arthritis. If sepsis is suspected it is better to proceed to an arthrotomy if clinically indicated.

The irritable hip syndrome should be diagnosed only after negative investigations, including plain films and ultrasound, full blood examination, ESR and bone scan. Needle aspiration has to be considered but irritable hip is often diagnosed without it, by observing in hospital in traction. If a deterioration or elevated temperature develops, needle aspiration with or without arthrotomy is indicated.

Avulsion bony injuries

Forceful contraction of insides originating around the pelvis can lead to avulsion at there origin in those with skelctal immaturity. This causes acute pain and muscular aysfunction:

- anterior superior iliac spine (sartorius)
- anterior inferior iliac spine long head rectus femoris
- ischial tuberosity (hamstrings)
- lesser trochanter (psoas)

Management includes X-ray and referral. As a rule surgical reattachment is not required.

The little athlete

The most common problem in the little athlete is pain or discomfort in the region of the iliac crest or anterior or superior iliac spines, usually associated with traction apophysitis or with acute avulsion fractures.[6] There is localised tenderness with pain on stretching and athletes should rest until they can compete without discomfort.

If there are persistent signs, pain in the knee, hip irritability or restricted range of motion, X-rays should be ordered to exclude serious problems such as SCFE or Perthes' disease.

HIP AND BUTTOCK PAIN IN THE ELDERLY

The following conditions are highly significant in the elderly:

- osteoarthritis of the hip
- aortoiliac arterial occlusion → vascular claudication
- spinal dysfunction with nerve root or referred pain
- degenerative spondylosis of lumbosacral spine → neurogenic claudication
- polymyalgia rheumatica
- trochanteric bursitis
- fractured neck of femur
- secondary tumours

Subcapital fractures

The impacted subcapital fractured femoral neck can often permit weight-bearing by an elderly patient. No obvious deformity of the leg is present. Radiographs are therefore essential for the investigation of all painful hips in the elderly. Patients often give a story of two falls - the first[7] very painful, the second with the hip just 'giving way' as the femoral head fell off.

The displaced subcapital fracture has at least a 40% incidence of avascular necrosis and usually requires prosthetic replacement in patients over 70 years. MRI scan is the investigation of choice if the X-ray are normal. Intertrochanteric fractures are also common (see p. 1403).

Avascularr necrosis

Consider this with hip pain in those at risk: corticosteroid use, SLE, sickle cell disease, past hip fracture, pregnancy, alcoholic liver disease Investigate with imaging (as above) and refer.

Osteoarthritis of the hip

Osteoarthritis of the hip is the most common form of hip disorder. It can be caused by primary osteoarthritis, which is related to an intrinsic disorder of articular cartilage, or to secondary osteoarthritis. Predisposing factors to the latter include previous trauma, DDH, septic arthritis acetabular dysplasia, SCFE and past in-flammatory arthritis.

Clinical features

- Equal sex incidence
- Usually after 50 years, increases with age
- May be bilateral: starts in one, other follows
- Insidious onset
- At first, pain worse with activity, relieved by rest, and then nocturnal pain and pain after resting
- Stiffness, especially after rising
- Characteristic deformity
- Stiffness, deformity and limp may dominate (pain mild)
- Pain usually in groin - may be referred to medial aspect of thigh, buttock or knee

Physical examination

- Antalgic gait
- Usually gluteal and quadriceps wasting
- First hip movements lost are IR and extension
- Fixed flexion deformity
- Hip held in flexion and ER (at first)
- Eventually all movements affected
- Order of movement loss is IR, extension, abduction, adduction, flexion, ER

Treatment

- Careful explanation: patients fear OA of hip
- Weight loss if overweight
- Relative rest
- Use crutchess for acute pain
- Analgesics and NSAIDs (judicious use)
- Aids and supports (e.g. walking stick)
- Physiotheropy
- Physical therapy, including isometric exercises
- Hydrotherapy - very useful

Surgery

This is an excellent option for those with severe pain or disability unresponsive to conservative measures. Total hip replacement is the treatment of choice in older patients but a femoral osteotomy[8] may be considered in younger patients in selected cases. > 90% achieve a good result. Most replacements last 10-15 years. Inselected patients is their 30's and 40's with seven disease total hip replacement is being performed successfullay. A type of total hip replacement called hip resurfacing is becoming once popular in certain situation in patients under 60 years of age.

GROIN PAIN

All condition involving the hip joint, especially OA, can present with groin pain (see p. 1412). Consider an unrecognised fracture neck of femur, psoas abscess, Paget's disease' osteitis pulis and hernias. Also consider hip labral or chondral lesions. Hip labral injuries present with inguinal pain or upper anhenor thigh pain and requne investigation and referred.

SACROILIAC PAIN

Pain arising from SIJ disorders is normally experienced as a dull ache in the buttock but can be referred to the groin or posterior aspect of the thigh. It may mimic pain from the lumbosacral spine or the hip joint. The pain may be unilateral or bilateral.

There are no accompanying neurological symptoms such as paraesthesia or numbness but it is common for more severe cases to cause a heavy aching feeling in the upper thigh.

Causes of SIJ disorders

The SIJs are difficult to palpate and examine but there are several tests that provoke them.

Direct pressure. With the patient lying prone a rhythmic springing force is applied directly to the upper and lower sacrum respectively.

Winged compression test. With the patient lying supine and with arms crossed, ‘separate’ the iliac crests with a downwards and outwards pressure. This compresses the SIJs.

Lateral compression test. With hands placed on the iliac crests, thumbs on the ASISs and heels of hand on the rim of the pelvis, compress the pelvis. This distracts the SIJs.

Patrick or Fabere test. This method can provoke the hip as well as the SIJ. The patient lies supine on the table and the foot of the involved side and extremity is placed on the opposite knee (the hip joint is now flexed, externally rotated and abducted). The knee and opposite ASIS are pressed downwards simultaneously (Figure 65.7). If low back or buttock pain is reproduced the cause is likely to be a disorder of the SIJ.



Figure 65.7 The Patrick (Fabere) test for right-sided hip or sacroiliac joint lesion, illustrating directions of pressure from the examiner

Unequal sacral 'rise' test. Squat behind the standing patient and place hands on top of the iliac crests and thumbs on the posterior superior iliac spines (PSIS). Ask the patient to bend slowly forwards and touch the floor. If one side moves higher relative to the other a problem may exist in the SIJs (e.g. a hypomobile lesion in the painful side if that side's PSIS moves higher).

Mechanical disorders of the SIJ

These problems are more common than appreciated and can be caused by hypomobile or hypermobile problems.

Hypomobile SIJ disorders are usually encountered in young people after some traumatic event, especially women following childbirth (notably multiple or difficult childbirth), and in those with structural problems (e.g. shortened leg). Pain tends to follow rotational stresses of the SIJ (e.g. tennis, dancing). Excellent results are obtained by passive mobilisation or manipulation, such as the non-specific rotation technique with the patient lying supine, as described in *Practice Tips* by the author.[9]

Hypermobile SIJ disorders are sometimes seen in athletes with instability of the symphysis pubis, in women after childbirth and in those with a history of severe trauma to the pelvis (e.g. MVAs, horse riders with foot caught in the stirrups after a fall). The patient presents typically with severe aching pain in the lower back, buttocks or upper thigh. Such problems are difficult to treat and manual therapy usually exacerbates the symptoms. Treatment consists of relative rest, analgesics and a sacroiliac supportive belt.

GLUTEUS MEDIUS TENDINITIS AND TROCHANTERIC BURSITIS

Pain around the lateral aspect of the hip is a common disorder, and is usually seen as lateral hip pain radiating down the lateral aspect of the thigh in older people engaged in walking exercises, tennis and similar activities. It is analogous in a way to the shoulder girdle, where supraspinatus tendinitis and subacromial bursitis are common wear-and-tear injuries.

The two common causes are tendinitis of the gluteus medius tendon, where it inserts into the lateral surface of the greater trochanter of the femur, and bursitis of one or both of the trochanteric bursae. Distinction between these two conditions is difficult, and it is possible that, as with the shoulder, they are related. The pain of bursitis tends to occur at night; that of tendinitis occurs with such activity as long walks and gardening.

Typical features of trochanteric bursalgia:

- female > 45 - 50 years
- pain on outside hip referred to as far as foot
- pain on lying on hip at night
- limp

Treatment

A trial of NSAIDs in younger patients is appropriate effective treatment for both conditions is similar.

1. Determine the points of maximal tenderness over the trochanteric region and mark them. (For tendinitis, this point is immediately above the superior aspect of the greater trochanter - see Figure 65.8.)



Figure 65.8 Injection technique for gluteus medius tendinitis (into area of maximal tenderness)

2. Inject aliquots of a mixture of 1 mL of long-acting corticosteroid with 5 - 7 mL of local anaesthetic into the tender area, which usually occupies an area similar to that of a standard marble.

The injection may be very effective. Follow-up management includes sleeping with a small pillow under the involved buttock, sleeping on a sheepskin rug and stretching the gluteal muscles with knee - chest exercises. Advise the patients to walk with the feet turned out - ‘the Charlie Chaplin gait’. One or two repeat in-jections over 6 or 12 months may be required. Surgical intervention is rarely necessary. Local application of ice and massage therapy may provide relief.

FASCIA LATA SYNDROME

Pain in the lateral thigh can be caused by inflammation off the fascia lata. It is often due to over use or weak musculature around the tip. Treatment is relative rest and physiotherapy.

ISCHIAL BURSITIS

This is known as tailor's bottom or 'weaver's bottom', which is occasionally seen is a bursa overlying the ischial tuberosity. Irritation of the sciatic nerve may coexist and the patient may appear to have sciatica.

Features

- severe pain when sitting, especially on a hard chair,
- tenderness at or just above the ischial tuberosity

Treatment

- Infiltration into the tender spot of a mixture of 4 ml of 1% lignocaine and 1 ml of LA corticocosteroid (avoid the sciatic nerve)
- Foam rubber cushion with two holes cut out for the ischial prominences

SNAPPING OR CLICKING HIP

Some patients complain of a clunking, clicking or snapping hip. This represents air annoying, problem which may be painful.

Causes

- A taut iliotibial band (tendon or tensor fascia femoris) slipping backwards and forwards over the prominence of the greater trochanter *or*
- The iliopsoas tendon snapping across the iliopectineal eminence
- The gluteus maximus sliding across the greater trochanter
- Joint laxity

Treatment

The basics of treatment are:

- explanation and reassurance
- exercises to stretch the iliotibial band[10]

Occasionally surgery is necessary to lengthen the ilio fibral band.

Exercises

- The patient lies on the 'normal' side and flexes the affected hip, with the leg straight and a weight around the ankle (Figure 65.9), to a degree that produces a stretching sensation along the lateral aspect of the thigh.



Figure 65.9 Treatment for the clicking hip

- This iliotibial stretch should be performed for 1 - 2 minutes, twice daily.

WHEN TO REFER

- Clinical evidence or suspicion of severe childhood disorders: DDH, Perthes' disease, septic arthritis, SCFE or osteomyelitis
- Undiagnosed pain, especially night pain
- Any fractures or suspicion of fractures such as impacted subcapital fracture or stress fracture of the femoral neck
- Patients with true claudication in buttock whether it is vascular from aortoiliac occlusion or neurogenic from spinal canal stenosis
- Patients with disabling osteoarthritis of the hip not responding to conservative measures; excellent results are obtained from surgery to the hip
- Any mass or lump

Practice tips

- Training on a plastic DDH model should be essential for all neonatal practitioners in order to master the manoeuvres for examining the neonatal hip.
- True hip pain is usually feltis the groin, thigh and medial aspect of the knee.
- The name of the Fabere test is an acronym for Flexion, Abduction, External Rotation and Extension of the hip.
- Night pain adds up to inflammation, bursitis or tumour.
- The hip joint can be the target of infections such as *Staphylococcus aureus* or tuberculosis or inflammatory disorders such as rheumatoid and the spondyloarthropathies, but these are rare numerically compared with osteoarthritis.

REFERENCES

1. Cyriax J. *Textbook of Orthopaedic Medicine*, Vol. 1 (6th edn). London: Balliere Tindall, 1976: 568 - 94.
2. Cormack J, Marinker M, Morell D. Hip problems. *Practice*. London: Kluwer-Harrap Handbooks, 1980; 3.65: 1 - 6.
3. Anonymous. *The Eastern Seal Guide to Children's Orthopaedics*. Toronto: Eastern Seal Society, 1982.
4. Robinson MJ. *Practical Paediatrics*. Melbourne: Churchill Livingstone, 1990: 72 - 4.
5. Corrigan B, Maitland G. *Practical Orthopaedic Medicine*. Sydney: Butterworths, 1986: 103 - 24.
6. Larkins PA. The little athlete. Aust Fam Physician, 1991; 20: 973 - 8.
7. Young D, Murtagh J. Pitfalls in orthopaedics. Aust Fam Physician, 1989; 18: 654 - 5.
8. Corrigan B, Maitland G. *Practical Orthopaedic Medicine*. Sydney: Butterworths, 1986: 324 - 31.
9. Murtagh J. *Practice Tips* (4th edn). Sydney: McGraw-Hill, 2004: 155.
10. Sheon RP, Moskowitz RW, Goldberg VM. Soft Tissue Rheumatic Pain (2nd edn). Philadelphia: Lea & Febiger, ?1987: 211 - 12.
11. Sheon RP, Moskowitz RW, Goldberg VM. *Soft Tissue Rheumatic Pain* (2nd edn). Philadelphia: Lea & Febiger, 1987: 211 - 12.

CHAPTER 66. PAIN IN THE LEG

Thou cold sciatica

Cripple our senators, that their limbs may halt

As lamely as their manners.

WILLIAM SHAKESPEARE (1564 - 1616), TIMON OF ATHENS

PAIN IN THE LEG

Pain in the leg has many causes, varying from a simple cramp to an arterial occlusion. Overuse of the legs in the athlete can lead to a multiplicity of painful leg syndromes, ranging from simple sprains of soft tissue to compartment syndromes. A major cause of leg pain lies in the source of the nervous network to the lower limb, namely the lumbar and sacral nerve roots of the spine. It is important to recognise radicular pain, especially from L5 and S1 nerve roots, and also the patterns of referred pain, such as from apophyseal (facet) joints and SIJs.

KEY FACTS AND CHECKPOINTS

- Always consider the lumbosacral spine, the SIJs and hip joints as important causes of leg pain.
- Hip joint disorders may refer pain around the knee only (without hip pain).
- Nerve root lesions may cause pain in the lower leg and foot only (without back pain).
- Nerve entrapment is suggested by a radiating burning pain, prominent at night and worse at rest.
- Older people may present with claudication in the leg from spinal canal stenosis or arterial obstruction or both.
- Think of the hip pocket wallet as a cause of sciatica from the buttocks down.
- Acute arterial occlusion to the lower limb requires relief within 4 hours (absolute limit of 6 hours).
- The commonest site of acute occlusion is the common femoral artery.
- Varicose veins can cause aching pain in the leg.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 66.1.

Table 66.1 Pain in the leg: diagnostic strategy model

Q. Probability diagnosis A. Cramps Nerve root 'sciatica' Muscular injury (e.g. hamstring) Osteoarthritis (hip, knee) Overuse injury (e.g. Achilles tendinitis)
Q. Serious disorders not to be missed A. Vascular • arterial occlusion (embolism) • thrombosis popliteal aneurysm • deep venous thrombosis • iliofemoral thrombophlebitis Neoplasia • primary (e.g. myeloma)

- metastases (e.g. breast to femur) Infection
- osteomyelitis
- septic arthritis
- erysipelas
- lymphangitis
- gas gangrene

Q. Pitfalls (often missed)

A. Osteoarthritis hip

Osgood-Schlatter disorder

Spinal canal stenosis

Herpes zoster (early)

Trochanteric bursitis/gluteus meduis tendonitis

Nerve entrapment

'Hip pocket nerve'

Iatrogenic: injection into nerve

Sacroiliac disorders

Gluteus medius tendinitis

Sympathetic dystrophy (causalgia)

Peripheral neuropathy

Rarities

- Osteoid osteoma
- Polymyalgia rheumatica (isolated)
- Paget's disease
- Popliteal artery entrapment
- Tabes dorsalis
- Ruptured Baker's cyst

Q. Seven masquerades checklist

A. Depression

Diabetes

Drugs (indirect)

Anaemia (indirect)

Thyroid disorder -

Spinal dysfunction

UTI -

Q. Is this patient trying to tell me something else?

A. Quite possible. Common with work-related injuries.

Probability diagnosis

Many of the causes, such as foot problems, ankle injuries and muscle tears (e.g. hamstrings and quadriceps), are obvious and common. There is a wide range of disorders related to overuse syndromes in athletes.

A very common cause of acute severe leg pain is cramp in the calf musculature, the significance of which escapes some patients as judged by middle-of-the-night calls.

One of the commonest causes is nerve root pain, invariably single, especially affecting the L5 and S1 nerve roots. Tests of their function and of the lumbosacral spine for evidence of disc disruption or other spinal dysfunction will be necessary. Should multiple nerve roots be involved other causes, such as compression from a tumour, should be considered. Remember that a spontaneous retroperitoneal haemorrhage in a patient on anticoagulant therapy can cause nerve root pain and present as intense acute leg pain. The nerve root sensory distribution is presented in Figure 66.1.



Figure 66.1 Dermatomes of the lower limb, representing approximate cutaneous distribution of the nerve roots

Other important causes of referred thigh pain include ischiogluteal bursitis (weaver's bottom) and gluteus medius tendinitis or trochanteric bursitis.

Serious disorders not to be missed

Neoplasia

Malignant disease, although uncommon, should be considered, especially if the patient has a history of one of the primary tumours, such as breast, lung or kidney. Such tumours can metastasise to the femur. Consider also osteogenic sarcoma and multiple myeloma, which is usually seen in the upper half of the femur. The possibility of an osteoid osteoma should be considered with pain in a bone relieved by aspirin.

Infections

Severe infections are not so common, but septic arthritis and osteomyelitis warrant con-sideration. Superficial infections such as erysipelas and lymphangitis occur occasionally.

Vascular problems

Acute severe ischaemia can be due to thrombosis or embolism of the arteries of the lower limb. Such occlusions cause severe pain in the limb and associated signs of severe ischaemia, especially of the lower leg and foot.

Chronic ischaemia due to arterial occlusion can manifest as intermittent claudication or rest pain in the foot due to small vessel disease.[1]

Various pain syndromes are presented in Figure 66.2. It is important to differentiate vascular claudication from neurogenic claudication (Table 66.2).


buttock and thigh claudication
calf claudication
calf claudication
foot claudication or rest pain in foot if small vessels involved
high obstruction (abdominal–pelvic)
obstruction in thigh (superficial femoral)
peripheral obstruction

Figure 66.2 Arterial occlusion and related symptoms according to the level of obstruction

Table 66.2 Comparison of the clinical features of neurogenic and vascular claudication

	Neurogenic claudication	Vascular claudication
Cause	• Spinal canal stenosis	• Aortoiliac arterial occlusive disease
Age	• Over 50 • Long history of backache	• Over 50
Pain site and radiation	• Proximal location, initially lumbar, buttocks and legs • Radiates distally	• Distal location • Buttocks, thighs and calves (especially) • Radiates proximally
Type of pain	• Weakness, burning, numbing or tingling (not cramping)	• Cramping, aching, squeezing
Onset	• Walking (uphill and downhill) • Distance walked varies • Prolonged standing	• Walking a set distance each time, especially uphill
Relief	• Lying down • Flexing spine (e.g. squat position) • May take 20 - 30 minutes	• Standing still - fast relief • Slow walking decreases severity
Associations	• Bowel and bladder symptoms	• Impotence

		• Rarely, paraesthesia or weakness
Physical examination		
• Peripheral pulses	• Present	• Present (usually) • Reduced or absent in some, especially after exercise
• Lumbar extension	• Aggravates	• No change
• Neurological	• Saddle distribution • Ankle jerk may be reduced after exercise	• Note: abdominal bruits after exercise
Diagnosis confirmation	• Radiological studies	• Duplex ultrasound • Ankle brachial index • Arteriography

Venous disorders

The role of uncomplicated varicose veins as a cause of leg pain is controversial. Nevertheless, varicose veins can certainly cause a dull aching ‘heaviness’ and cramping, and can lead to painful ulceration.

Superficial thrombophlebitis is usually obvious, but it is vital not to overlook deep venous thrombosis. These more serious conditions of the veins can cause pain in the thigh or calf.

Pitfalls

There are many traps and pitfalls in the painful leg. Herpes zoster at the pre-eruption phase is an old trap and more so when the patient develops only a few vesicles in obscure parts of the limbs.

In future we can expect to encounter more cases of spinal canal stenosis (secondary to the degenerative changes) in the elderly. The early diagnosis can be difficult, and buttock pain on walking has to be distinguished from vascular claudication due to a high arterial obstruction.

The many disorders of the SIJ and hip region can be traps, especially the poorly diagnosed yet common gluteus medius tendinitis. Another more recent phenomenon is the ‘hip pocket nerve syndrome’, where a heavy wallet crammed with credit cards can cause pressure on the sciatic nerve.

One of the biggest traps, however, is when hip disorders, particularly osteoarthritis, present as leg pain, especially on the medial aspect of the knee.

Nerve entrapments (Figure 66.3) are an interesting cause of leg pain, although not as common as in the upper limb. Some entrapments to consider include:


lateral femoral cutaneous nerve of thigh
obturator nerve
superficial peroneal
posterior tibial at tarsal tunnel (causes pain on sole of foot)
deep peroneal

Figure 66.3 Distribution of pain in the leg from entrapment of specific nerves; the sites of entrapments are indicated by an **X**

- lateral cutaneous nerve of thigh, known as meralgia paraesthetica
- common peroneal nerve
- posterior tibial nerve at ankle (the 'tarsal tunnel' syndrome)
- obturator nerve, in obturator canal
- femoral nerve (in inguinal region or pelvis)

Then there are the rare causes. One overlooked problem is sympathetic dystrophy, which may follow even minor trauma to the limb. This 'causalgia' syndrome manifests as burning or aching pain with vasomotor instability in the limbs. The essential feature is the disparity between the intensity of the pain and the severity of the inciting injury.

General pitfalls

- Overlooking beta-blockers and anaemia as a precipitating factor for vascular claudication
- Overlooking hip disorders as a cause of knee pain
- Mistaking occlusive arterial disease for sciatica
- Confusing nerve root syndromes with entrapment syndromes

Seven masquerades checklist

The outstanding cause of leg pain in this group is spinal dysfunction. Apart from nerve root pressure due to a disc disruption or foraminal entrapment, pain can be referred from the apophyseal (facet joints). Such pain can be referred as far as the mid-calf (Figure 66.4).



Figure 66.4 Possible referred pain patterns from dysfunction of an apophyseal joint, illustrating pain radiation patterns from stimulation by injection of the right L4 - 5 apophyseal joint. Reproduced with permission from C. Kenna and J. Murtagh. *Back Pain and Spinal Manipulation*. Sydney: Butterworths, 1989.

The other checklist conditions - depression, diabetes, drugs and anaemia - can be associated with pain in the leg. Depression can reinforce any painful complex.

Diabetes can cause discomfort through a peripheral neuropathy that can initially cause localised pain before anaesthesia predominates. Drugs such as beta-blockers, and anaemia, can precipitate or aggravate intermittent claudication in a patient with a compromised circulation.

Psychogenic considerations

Pain in the lower leg can be a frequent complaint (maybe a magnified one) of the patient with non-organic pain, such as the malingerer, the conversion reaction patient (hysteria) and the depressed. Sometimes regional pain syndrome (reflex or post-traumatic) is incorrectly diagnosed as functional.

THE CLINICAL APPROACH

Careful attention to basic detail in the history and examination can point the way of the clinical diagnosis.

History

In the history it is important to consider several distinctive aspects, outlined by the following questions.

- Is the pain of acute or chronic onset?
- If acute, did it follow trauma or activity?
    - If not, consider a vascular cause: vein or artery; occlusion or rupture.
- Is the pain ‘mechanical’ (related to movement)?
    - If it is unaltered by movement of the leg or a change in posture, it must arise from a soft tissue lesion, not from bone or joints.
- Is the pain postural?
    - Analyse the postural elements that make it better or worse.

  - If worse on sitting, consider a spinal cause (discogenic) or ischial bursitis.
  - If worse on standing, consider a spinal cause (instability) or a local problem related to weight-bearing (varicose veins).
  - If worse lying down, consider vascular origin, such as small vessel peripheral vascular disease.
  - Pain unaffected by posture is activity-related.
- Is the pain related to walking?
  - *No*: Determine the offending activity (e.g. joint movement with arthritis).
  - *Yes*: If immediate onset, consider local cause at site of pain (e.g. stress fracture). If delayed onset, consider vascular claudication or neurogenic claudication.
- Is the site of pain the same as the site of trauma?
  - If not, the pain in the leg is referred. Important considerations include lesions in the spine, abdomen or hip and entrapment neuropathy.
- Is the pain arising from the bone?
  - If so, the patient will point to the specific site and indicate a ‘deep’ bone pain (consider tumour, fracture or, rarely, infection) compared with the more superficial muscular or fascial pain.
- Is the pain arising from the joint?
  - If so, the clinical examination will determine whether it arises from the joint or juxtaposed tissue.

Physical examination

The first step is to watch the patient walk and assess the nature of any limp.

Note the posture of the back and examine the lumbar spine. Have both legs well exposed for the inspection.

Inspect the patient's stance and note any asymmetry and other abnormalities, such as swellings, bruising, discolouration, or ulcers and rashes. Note the size and symmetry of the legs and the venous pattern. Look for evidence of ischaemic changes, especially of the foot.

Palpate for local causes of pain and if no cause is evident examine the spine, blood vessels (arteries and veins) and bone. Areas to palpate specifically are the ischial tuberosity, trochanteric area, hamstrings and tendon insertions. Palpate the superficial lymph nodes. Note the temperature of the feet and legs. Perform a vascular examination, including the peripheral pulses and the state of the veins if appropriate.

If evidence of peripheral vascular disease (PVD), remember to auscultate the abdomen and adductor hiatus, and the iliac, femoral and popliteal vessels.

A neurological examination may be appropriate, particularly to test nerve root lesions or entrapment neuropathies.

Examination of the joints, especially the hip and SIJs, is very important.

Investigations

A checklist of investigations that may be necessary to make the diagnosis is as follows:

- FBE and ESR
- Radiology:
  - leg X-rays, especially knee, hip
  - plain X-ray of lumbosacral spine
  - CT scan of lumbosacral spine
  - MRI scan of lumbosacral spine
  - bone scan
- electromyography
- vascular
  - arteriography
  - duplex ultrasound scan

  - ankle brachial index
  - venous pool radionuclide scan
  - contrast venography
  - air plethysmograph (varicose veins)
  - D-dimer test

LEG PAIN IN CHILDREN

Aches and pains in the legs are a common complaint in children. The most common cause is soreness and muscular strains due to trauma or unaccustomed exercise. One cause of bilateral leg pain in children is leukeuina.

It is important to consider child abuse, especially if bruising is noted on the backs of the legs.

Growing pains

So-called 'growing pains', or idiopathic leg pain, is thought to be responsible for up to 20% of leg pain in children.[2] Such a diagnosis is vague and often made when a specific cause is excluded. It is usually not due to 'growth' but related to excessive exercise or trauma from sport and recreation, and probably emotional factors.

The pains are typically intermittent and symmetrical and deep in the legs, usually in the anterior thighs or calves. Although they may occur at any time of the day or night, typically they occur at night, usually when the child has settled in bed. The pains usually last for 30 to 60 minutes and tend to respond to attention such as massage with an analgesic balm or simple analgesics.

Serious problems

It is important to exclude fractures (hence the value of X-rays if in doubt), malignancy (such as osteogenic sarcoma, Ewing's tumour or infiltration from leukaemia or lymphoma), osteoid osteoma, osteomyelitis, scurvy and beri-beri (rare disorders in developed countries) and congenital disorders such as sickle-cell anaemia, Gaucher's disorder and Ehlers-Danlos syndrome.

LEG PAIN IN THE ELDERLY

The older the patient the more likely it is that arterial disease with intermittent claudication and neurogenic claudication due to spinal canal stenosis will develop. Other important problems of the elderly include degenerative joint disease, such as osteoarthritis of the hips and knees, muscle cramps, herpes zoster, Paget's disease, polymyalgia rheumatica (affecting the upper thighs) and sciatica.

SPINAL CAUSES OF LEG PAIN

Problems originating from the spine are an important, yet at times complex, cause of pain in the leg.

Important causes are:

- nerve root (radicular) pain from direct pressure
- referred pain from:
  - disc pressure on tissues in front of the spinal cord
  - apophyseal joints
  - SIJs
- spinal canal stenosis causing claudication

Various pain patterns are presented in **Figures** 66.3 and 66.4.

Nerve root pain

Nerve root pain from a prolapsed disc is a common cause of leg pain. A knowledge of the dermatomes of the lower limb (**Figure** 66.1) provides a pointer to the involved nerve root, which is usually L5 or S1 or both. The L5 root is invariably caused by an L4 - 5 disc prolapse and the S1 root by an L5 - S1 disc prolapse. The nerve root syndromes are summarised in Table 66.3.

Table 66.3 Comparison of neurological findings of the neurological levels L4, L5 and S1. Reproduced with permission from S. Hoppenfeld. Physical Examination of The Spine and Extremities. Norwalk, Ct: Appleton & Lange

Nerve root	Pain distribution	Sensory loss	Motor weakness changes	Reflex
L3	Front of thigh, inner aspect of thigh, knee and leg	Anterior aspect of thigh 	Extension of knee	Knee jerk
L4	Anterior thigh to front of knee	Lower outer aspect of thigh and knee, inner great toe 	Flexion, adduction of knee, inversion of foot 	Knee jerk 
L5	Lateral aspect of leg, dorsum of foot and great toe	Dorsum of foot, great toe, 2nd and 3rd toes, anterolateral aspect of lower leg 	Dorsiflexion of great toe 	Tibialis posterior (clinically impractical) 

				
S1	Buttock to back of thigh and leg, central calf, lateral aspect of ankle and sole of foot	Lateral aspect of ankle, foot (4th and 5th toes)  	Plantar flexion of ankle and toes, eversion of foot 	Ankle jerk  peroneus longus + brevis

A summary of the physical examination findings for the most commonly involved nerve roots is presented in Figure 66.5.


(a)
tibialis posterior
flexor digitorum longus
posterior tibial nerve
flexor hallucis longus
flexor retinaculum
tarsal tunnel



Figure 66.5 (a) Anatomy of the tarsal tunnel syndrome (b) showing injection sites

Sciatica

Sciatica is defined as pain in the distribution of the sciatic nerve or its branches (L4, L5, S1, S2, S3) that is caused by nerve pressure or irritation. Most problems are due to entrapment neur-opathy of a nerve root, in either the spinal canal (as outlined above) or the intervertebral foramen.

It should be noted that back pain may be absent and peripheral symptoms only will be present.

Treatment of sciatica

Acute sciatica. A protracted course can be anticipated, in the order of 12 weeks (see page 387). The patient should be reassured that spontaneous recovery can be expected. A trial of conservative treatment would be recommended thus:

- back care education
- relative bed rest if very painful only (2 days is optimal) - a firm base is ideal
- return to activities of daily living ASAP
- analgesics (avoid narcotic analgesics)
- NSAIDs (2 weeks is recommended)
- basic exercise program, including swimming
- traction can help, even intermittent manual

Referral to a therapist of your choice (e.g. physiotherapist) might be advisable. Con-ventional spinal manipulation is usually contraindicated for radicular sciatica. If the patient is not responding or the circumstances demand more active treatment, an epidural anaesthetic injection is appropriate. Surgical intervention may be necessary.

Chronic sciatica. If a trial of NSAIDs, rest and physiotherapy has not brought significant relief, an epidural anaesthetic (lumbar or caudal) using half-strength local anaesthetic (e.g. 0.25% bupivacaine HCl) and a depot corticosteroid (e.g. triamcinolone) is advisable. The lumbar route under image intensification is preferred.

REFERRED PAIN

Referred pain in the leg can arise from disorders of the SIJs or from spondylogenic disorders. It is typically dull, heavy and diffuse. The patient uses the hand to describe its distribution compared with the use of fingers to point to radicular pain.

Spondylogenic pain

Non-radicular or spondylogenic pain is that which originates from any of the components of the vertebrae (spondyles), including joints, the intervertebral disc, ligaments and muscle attachments. An important example is distal referred pain from disorders of the apophyseal joints, where the pain can be referred to any part of the limb as far as the calf and ankle but most commonly to the gluteal region and proximal thigh (Figure 66.4).

Another source of referred pain is that caused by compression of a bulging disc against the posterior longitudinal ligament and dura. The pain is typically dull, deep and poorly localised. The dura has no specific dermatomal localisation, and so the pain is usually experienced in the low back, sacroiliac area and buttocks. Less commonly it can be referred to the coccyx, groin and both legs to the calves. It is not referred to the ankle or the foot.

Sacroiliac dysfunction

This causes typically a dull ache in the buttock but it can be referred to the iliac fossa, groin or posterior aspects of the thighs. It rarely radiates to or below the knee. It may be caused by inflammation (sacroiliitis) or mechanical dysfunction. The latter must be considered in a postpartum woman presenting with severe aching pain present in both buttocks and thighs.

NERVE ENTRAPMENT SYNDROMES

Entrapment neuropathy can result from direct axonal compression or can be secondary to vascular problems, but the main common factor is a nerve passing through a narrow rigid compartment where movement or stretching of that nerve occurs under pressure.

Clinical features of nerve entrapment:

- pain at rest (often worse at night)
- variable effect with activity
- sharp, burning pain
- radiating and retrograde pain
- clearly demarcated distribution of pain
- paraesthesia may be present
- tenderness over nerve
- may be positive Tinel's sign

Meralgia paraesthetica

This is the commonest lower limb entrapment and is due to the lateral femoral cutaneous nerve of the thigh being trapped under the lateral end of the inguinal ligament, 1 cm medial to the ASIS.[3]

The nerve is a sensory nerve from L2 and L3. It occurs mostly in middle-aged people, due mainly to thickening of the fibrous tunnel beneath the inguinal ligament, and is associated with obesity, pregnancy, ascites or local trauma such as belts, trusses and corsets. Its entrapment causes a burning pain with associated numbness and tingling (Figure 66.3 at p. 711).

The distribution of pain is confined to a localised area of the lateral thigh and does not cross the midline of the thigh.

Differential diagnosis

- L2 or L3 nerve root pain (L2 causes buttock pain also)

- Femoral neuropathy (extends medial to mid-line)

Treatment options

- Injection of corticosteroid medial to the ASIS, under the inguinal ligament
- Surgical release (neurolysis) if refractory

Note: Meralgia paraesthetica often resolves spontaneously.

Peroneal nerve entrapment

The common peroneal (lateral popliteal) nerve can be entrapped where it winds around the neck of the fibula or as it divides and passes through the origin of the peroneus longus muscle 2.5 cm below the neck of the fibula. It is usually injured, however, by trauma or pressure at the neck of the fibula.

Symptoms and signs:

- pain in the lateral shin area and dorsum of the foot
- sensory symptoms in the same area
- weakness of eversion and dorsiflexion of the foot (described by patients as ‘a weak ankle’)

Differential diagnosis

- L5 nerve root (similar symptoms)

Treatment

- Shoe wedging or other orthotics to maintain eversion
- Neurolysis is the most effective treatment

Tarsal tunnel syndrome

This is an entrapment neuropathy of the posterior tibial nerve in the tarsal tunnel beneath the flexor retinaculum on the medial side of the ankle (Figure 66.5a). The condition is due to dislocation or fracture around the ankle or tenosynovitis of tendons in the tunnel from injury, rheumatoid arthritis, and other inflammations

Symptoms and signs

- A burning or tingling pain in the toes and sole of the foot, occasionally the heel
- Retrograde radiation to calf, perhaps as high as the buttock
- Numbness is a late symptom
- Discomfort often in bed at night and worse after standing
- Removal of shoe may give relief
- Sensory nerve loss variable, may be no loss
- Tinel test (finger or reflex hammer tap over nerve below and behind medial malleolus) may be positive
- Tourniquet applied above ankle may reproduce symptoms

The diagnosis is confirmed by electro-diagnosis.

Treatment

- Relief of abnormal foot posture with orthotics
- Corticosteroid injection
- Decompression surgery if other measures fail

Injection for tarsal tunnel syndrome

Using a 23-gauge 32 mm needle, a mixture of triamcinolone 10 mg/mL or 40 mg methyl-prednisolone in 1% xylocaine or procaine is injected into the tunnel from either above or below the flexor retinaculum. The sites of injection are shown in Figure 66.5b; care is required not to inject the nerve.

VASCULAR CAUSES OF LEG PAIN

Occlusive arterial disease

Risk factors for peripheral vascular disease (for development and deterioration):

- smoking
- diabetes mellitus
- hypertension
- hypercholesterolaemia
- family history
- atrial fibrillation (embolism)

Aggravating factors:

- beta-blocking drugs
- anaemia

Acute lower limb ischaemia

Sudden occlusion is a dramatic event that requires immediate diagnosis and management to save the limb.

Causes:

- embolism - peripheral arteries
- thrombosis
    - major artery
    - popliteal aneurysm
- traumatic contusion (e.g. postarterial puncture)

The symptoms and signs of acute embolism and thrombosis are similar, although thrombosis of an area of atherosclerosis is often preceded by symptoms of chronic disease (e.g. claudication). The commonest site of acute occlusion is the common femoral artery (Figure 66.6).



Figure 66.6 Common sites of acute arterial occlusion

Signs and symptoms - the 6 P's

- Pain
- Pallor
- Paraesthesia or numbness
- Pulselessness
- Paralysis
- ‘Perishing’ cold

The pain is usually sudden and severe and any improvement may be misleading. Sensory changes initially affect light touch, not pinprick. Paralysis (paresis or weakness) and muscle compartment pain or tenderness is a most important and ominous sign.

Other signs include mottling of the legs, collapsed superficial veins, and no capillary return. If the foot becomes dusky purple and fails to blanch on pressure, irreversible necrosis has occurred.

Note: Look for evidence of atrial fibrillation.

Examination of arterial circulation

This applies to chronic ischaemia and also to acute ischaemia.

Skin and trophic changes

Note colour changes, hair distribution and wasting. Note the temperature of the legs and feet with the backs of your fingers.

Palpation of pulses

It is important to assess four pulses carefully (Figure 66.7). Note that the popliteal and posterior tibial pulses are difficult to feel, especially in obese subjects.



Figure 66.7 Sites of palpation of peripheral pulses in the leg

Femoral artery. Palpate deeply just below the inguinal ligament, midway between the ASIS and the symphysis pubis. If absent or diminished, palpate over abdomen for aortic aneurysm.

Popliteal artery. Flex the leg to relax the hamstrings. Place fingertips of both hands to meet in the midline. Press them deeply into the popliteal fossa to compress artery against the upper end of the tibia (i.e. just below the level of the knee crease). Check for a popliteal aneurysm (very prominent popliteal pulsation).

Posterior tibial artery. Palpate, with curved fingers, just behind and below the tip of the medial malleolus of the ankle.

Dorsalis pedis artery. Feel at the proximal end of the first metatarsal space just lateral to the extensor tendon of the big toe.

Oedema

Look for evidence of oedema: pitting oedema is tested by pressing firmly with your thumb for at least 5 seconds over the dorsum of each foot, behind each medial malleolus and over the shins.

Postural colour changes (Buerger's test)

Raise both legs to about 60? for about 1 minute, when maximal pallor of the feet will develop. Then get the patient to sit up on the couch and hang both legs down.[4] Note, comparing both feet, the time required for return of pinkness to the skin (normally less than 10 seconds) and filling of the veins of the feet and ankles (normally about 15 seconds). Look for any unusual rubor (dusky redness) that takes a minute or more in the dependent foot. A positive Buerger's test is pallor on elevation and rubor on dependency and indicates severe chronic ischaemia.

Auscultation for bruits after exercise

Listen over abdomen and femoral area for bruits.

Note: Neurological examination (motor, sen-sory, reflexes) is normal unless there is associated diabetic peripheral neuropathy.

Management of acute ischaemia

Golden rules. Occlusion is usually reversible if treated within 4 hours (i.e. limb salvage). It is often irreversible if treated after 6 hours (i.e. limb amputation).

Treatment

- Intravenous heparin (immediately) 5000 U
- Emergency embolectomy (ideally within 4 hours)
  - under general or local anaesthesia
  - through an arteriotomy site in the common femoral artery
  - embolus extracted with Fogarty balloon or catheter or
- Stenting of vessels (a good modern option) - discuss this with an interventional cardiovascular physician.
- Arterial bypass if acute thrombosis in chronically diseased artery
- In selected cases thrombolysis with streptokinase or urokinase appropriate
- Amputation (early) if irreversible ischaemic changes
- Lifetime anticoagulation with warfarin will be required

Note: An acutely ischaemic limb is rarely life threatening in the short term. Thus, even in the extremely aged, demented or infirm, a simple embolectomy not only is worthwhile but also is usually the most expedient treatment option.

Chronic lower limb ischaemia

Chronic ischaemia caused by gradual arterial occlusion can manifest as intermittent claudication, rest pain in the foot, or overt tissue loss - ulceration, gangrene (Figure 66.8).



Figure 66.8 Gangrene of the lateral aspect of the foot following attempted amputation of an ischaemic toe. A below knee amputation was eventually required

Intermittent claudication is a pain or tightness in the muscle on exercise (Latin claudicare, to limp), relieved by rest. Rest pain is a constant severe burning-type pain or discomfort in the forefoot at rest, typically occurring at night when the blood flow slows down.

The main features are compared in Table 66.4.

Table 66.4 Comparison between intermittent claudication and ischaemic rest pain

	Intermittent claudication	Ischaemic rest pain
Quality of pain	Tightness/cramping	Constant ache
Timing of pain (typical)	Daytime; walking, other exercise	Night-time; rest

Tissue affected	Muscle	Skin
Site	Calf > thigh > buttock	Forefoot, toes, heels
Aggravation	Walking, exercise	Recumbent, walking
Relief	Rest	Hanging foot out of bed; dependency
Associations	Beta-blockers Anaemia	Night cramps Swelling of feet

Intermittent claudication

The level of obstruction determines which muscle belly is affected (Figure 66.2 and 66.7).

Proximal obstruction (e.g. aortoiliac)

- Pain in the buttock, thigh and calf, especially when walking up hills and stairs
- Persistent fatigue over whole lower limb
- Impotence is possible (Leriche syndrome)

Obstruction in the thigh

- Superficial femoral (the commonest) causes pain in the calf (e.g. 200 - 500 m, depending on collateral circulation
- profunda femoris → claudication at about 100 m
- multiple segment involvement → claudication at 40 - 50 m

Causes:

- atherosclerosis (mainly men over 50, smokers)
- embolisation (with recovery)
- Buerger's disease: affects small arteries, causes rest pain and cyanosis (claudication uncommon)
- popliteal entrapment syndrome (< 40 years of age)

Note: The presence of rest pain implies an immediate threat to limb viability.

Investigations

- FBE: exclude polycythemia and thrombocytosis
- Colour Doppler ultrasound
  - measure resting ankle systolic BP
  - determine ankle/brachial index; normal value 0.9 - 1.1
- Angiography: the gold standard, reserved for proposed intervention
- Digital substraction angiography (developing)

MANAGEMENT OF OCCLUSIVE VASCULAR DISEASE

Prevention (for those at risk)

- Smoking is the risk factor and must be stopped.
- Other risk factors, especially hyperlipidaemia, must be attended to and weight reduction to ideal weight is important.
- Exercise is excellent, especially walking.

Diagnostic plan

- Check if patient is taking beta-blockers.
- General tests: blood examination, random blood sugar, urine examination, ECG.

- Measure blood flow by duplex ultrasound examination or ankle brachial index.
- Arteriography should be performed only if surgery contemplated.

Treatment

- General measures (if applicable): control obesity, diabetes, hypertension, hyperlipidaemia, cardiac failure.
- Achieve ideal weight.
- There must be absolutely no smoking.
- Exercise: daily graduated exercise to the level of pain. About 50% will improve with walking; so advise as much walking as possible.
- Try to keep legs warm and dry.
- Maintain optimal foot care (podiatry).
- Drug therapy: aspirin 150 mg daily.

Note:

- Vasodilators and sympathectomy are of little value.
- About one-third progress, while the rest regress or don't change.[5]

When to refer to a vascular surgeon

- ‘Unstable’ claudication of recent onset; deteriorating
- Severe claudication - unable to maintain lifestyle
- Rest pain
- ‘Tissue loss’ in feet (e.g. heel crack, ulcers on or between toes, dry gangrenous patches, infection)

Surgery. Reconstructive vascular surgery is indicated for progressive obstruction, intolerable claudication and obstruction above the inguinal ligament:

- endarterectomy - for localised iliac stenosis
- bypass graft (iliac or femoral artery to popliteal or anterior or posterior tibial arteries)

Percutaneous transluminal dilation: This angioplasty is performed with a special intra-arterial balloon catheter for localised limited occlusions. An alternative to the balloon is laser angioplasty.

VENOUS DISORDERS

Varicose veins

Varicose veins are dilated, tortuous and elongated superficial veins in the lower extremity.

The veins are dilated because of incompetence of the valves in the superficial veins or in the communicating or perforating veins between the deep and superficial systems (Figure 65.9). The cause is a congenital weakness in the valve and the supporting vein wall but there are several predisposing factors (Table 66.5), the most important being family history, female sex (5:1), pregnancy and multiparity. Previous DVT can also damage valves, especially calf perforators, and cause varicose veins.


incompetent valve
deep vein
varicose superficial vein
muscles that pump up blood
perforating veins
incompetent valve
site of varicose 'eczema'

Figure 66.9 The common sites of varicose veins

Table 66.5 Risk factors for varicose veins

Female sex
Family history
Pregnancy
Multiparity
Age
Occupation
Diet (low fibre)

Dilated superficial veins, which can mimic varicose veins, may be caused by extrinsic compression of the veins by a pelvic or intra-abdominal tumour (e.g. ovarian carcinoma, retroperitoneal fibrosis). Uncommonly, but importantly, superficial veins dilate as they become collaterals following previous DVT, especially if the ilio-femoral segment is involved.

Symptoms

Varicose veins may be symptomless, the main complaint being their unsightly appearance. Symptoms include swelling, fatigue, heaviness in the limb, an aching discomfort and itching.

Varicose veins and pain

They may be painless even if large and tortuous. Pain is a feature where there are incompetent perforating veins running from the posterior tibial vein to the surface through the soleus muscle.

Severe cases lead to the lower leg venous hypertension syndrome[6] characterised by pain that is worse after standing, cramps in the leg at night, irritation and pigmentation of the skin, swelling of the ankles and loss of skin features such as hair.

A careful history will usually determine if the aching is truly due to varicose veins and not to transient or cyclical oedema, which is a common condition in women.[7]

The complications of varicose veins are summarised in Table 66.6.

Table 66.6 Complications of varicose veins

Superficial thrombophlebitis
Skin 'eczema' (10%)
Skin ulceration (20%)
Bleeding
Calcification
Marjolin's ulcer (squamous cell carcinoma)

Examination

The following tests will help determine the site or sites of the incompetent valves.

Venous groin cough impulse. This helps determine long saphenous vein incompetence.

Place the fingers over the line of the vein immediately below the fossa ovalis (4 cm below and 4 cm lateral to the pubic tubercle).[8] Ask the patient to cough - an impulse or thrill will be felt expanding and travelling down the long saphenous vein. A marked dilated long saphenous vein in the fossa ovalis (saphena varix) will confirm incompetence. It disappears when the patient lies down.

Trendelenburg test. In this test for long saphenous vein competence the patient lies down and the leg is elevated to 45? to empty the veins (Figure 66.10a). Apply a tourniquet with sufficient pressure to prevent reflux over the upper thigh just below the fossa ovalis. (Alternatively, this opening can be occluded by firm finger pressure, as originally described by Trendelenburg.)



Figure 66.10a Trendelenburg test: the leg is elevated to 45? to empty the veins and a tourniquet applied

The patient then stands. The long saphenous system will remain collapsed if there are no incompetent veins below the level of the fossa ovalis. When the pressure is released the vein will fill rapidly if the valve at the sapheno-femoral junction is incompetent (Figure 66.10b). This is a positive Trendelenburg test.



Figure 66.10b Trendelenburg test: test for competence of long saphenous venous system (medial aspect of knee)

Note: A doubly positive Trendelenburg test is when the veins fill rapidly before the pressure is released and then with a ‘rush’ when released. This indicates coexisting incompetent perforators and long saphenous vein.

Short saphenous vein incompetence test. A similar test to the Trendelenburg test is performed with the pressure (tourniquet or finger) being applied over the short saphenous vein just below the popliteal fossa (Figure 66.11).



Figure 66.11 Testing for competence of the short saphenous vein

Incompetent perforating vein test. Accurate clinical tests to identify incompetence in the three common sites of perforating veins on the medial aspect of the leg, posterior to the medial border of the tibia, are difficult to perform. The general appearance of the leg and palpation of the sites give some indication of incompetence here.

Note: Venous duplex ultrasound studies will accurately localise sites of incompetence and determine the state of the functionally important deep venous system.

Management of varicose veins

Prevention

- Maintain ideal weight.
- Eat a high-fibre diet.
- Rest and wear supportive stockings if at risk (pregnancy, a standing occupation).

Treatment

- Keep off legs as much as possible.
- Sit with legs on a footstool.
- Use supportive stockings or tights (apply in morning before standing out of bed).
- Avoid scratching itching skin over veins.

Compression sclerotherapy

- Use a small volume of sclerosant (e.g. sodium tetradecyl sulphate - Fibro-vein 3%).
- It is ideal for smaller isolated veins particularly below the knee joint.

Surgical ligation and stripping

- This is the best treatment when a clear association exists between symptoms and obvious varicose veins (i.e. long saphenous vein incompetence).
- Remove obvious varicosities and ligate perforators.

Note: Surgery for varicose veins may not relieve heavy, aching legs.

Superficial thrombophlebitis

- Usually occurs in superficial varicose veins
- Presents as a tender, reddened subcutaneous cord in leg
- Usually localised oedema
- No generalised swelling of the limb or ankle
- Requires symptomatic treatment only (see below) unless there is extension above the level of the knee when there is a risk of pulmonary embolism
- Venous duplex scan is diagnostic and also determines:
    - extent of superficial thrombosis, and
    - if coexisting, unsuspected DVT is present

Treatment

The objective is to prevent propagation of the thrombus by uniform pressure over the vein.

- Cover whole tender cord with a thin foam pad.
- Apply a firm elastic bandage (preferable to crepe) from foot to thigh (well above cord).
- Leave pad and bandage on for 7 - 10 days.
- Bed rest with leg elevated is recommended.
- Prescribe an NSAID (e.g. indomethacin, for 10 days).

Note:

- No anticoagulants are required.
- The traditional glycerin and ichthyol dressings are still useful.
- Consider association between thrombophlebitis and deep-seated carcinoma.
- If the problem is above the knee, ligation of the vein at the saphenofemoral junction is indicated.

Deep venous thrombosis

Refer to Chapter 133-Thrombbbosis and embolism thrambo (page xxx).

Iliofemoral thrombophlebitis (phlegmasia dolens)[19]

This rare but life-threatening condition is when an extensive clot obstructs the iliofemoral veins so completely that subcutaneous oedema and blanching occurs. This initially causes a painful 'milky white leg', previously termed phlegmasia alba dolens (used to be seen in late pregnancy or early puerperium). It may deteriorate and become cyanotic - phlegmasia cerulea dolens - representing incipient venous infarction. Massive iliofemoral occlusion is an emergency as such patients may develop 'shock', gangrene and pulmonary embolus.

MANAGEMENT OF OTHER PAINFUL CONDITIONS

Cellulitis and erysipelas

The causative organisms are *Streptococcus pyogenes* (commonest) and *Staphylococcus aureus*. Others include *Haemophilius influenzae*, Aeromonas and fungal infection (especially in the immunocompromised). Predisposing factors include cuts, abrasions, ulcers, insect bites, foreign matter, IV drug use and skin disorders such as eczema and tinea pedis of toe webs.

- Rest in bed.
- Elevate limb (in and out of bed).
- Use aspirin or paracetamol for pain and fever.
- Wound cleansing and dressing with non-sticking saline dressing.

Streptococcus pyogenes (the common cause)[11]

- Severe
  - flucloxacillin/dicloxacillin 2 g IV 6 hourly
- Less severe
  - fludicloxacillin 500 mg (0) 6 hourly 7 - 10 days.
- If penicillin sensitive
  - cephalexin 0.5 mg (o) 6 hourly

Staphylococcus aureus[10]

- Severe, may be life-threatening
  - flucloxacillin/dicloxacillin 2 g IV 6 hourly
- Less severe
  - flucloxacillin/dicloxacillin 500 mg (o) 6 hourly *or*
  - cephalexin 500 mg (o) 6 hourly

Furuncle (boil) of groin

A painful furuncle caused by S. aureus in the hairy area of the groin is common. The aim is to treat conservatively.

- Localised
  - local antiseptics

  - hot compresses
  - drain when ‘ripe’
- deep/extensive
  - dicloxacillin 500 mg (o) 6 hourly for 5 - 7 days
  - drain when ‘ripe’, not before

Nocturnal cramps

Note: Treat cause (if known) - tetanus, drugs, sodium depletion, hypothyroidism.

Physical measures

- Muscle stretching and relaxation exercises: calf stretching for 3 minutes before retiring,[12] then rest in chair with the feet out horizontal to the floor with cushion under tendoachilles for 10 minutes.
- Massage and apply heat to affected muscles.
- Try to keep bedclothes off feet and lower part of legs - a doubled-up pillow at the foot of the bed can be used.

Medication

- Tonic water before retiring may help.
- Drug treatment:
  - quinine sulphate 300 mg nocte daily or alternate nights (benefit confirmed by one systematic review[12]but remains controversial) *or*
  - biperiden 2 - 4 mg nocte

Roller injuries to legs

A patient who has been injured by a wheel passing over a limb, especially a leg, can present a difficult problem. A freely spinning wheel is not so dangerous, but serious injuries occur when a non-spinning (braked) wheel passes over a limb and these are compounded by the wheel then reversing over it. This leads to a ‘degloving’ injury due to shearing stress. The limb may look satisfactory initially, but skin necrosis may follow.

- Admit to hospital for observation.
- Fasciotomy with open drainage may be an option.
- Surgical decompression with removal of necrotic fat is often essential.

WHEN TO REFER

- The sudden onset of pain, pallor, pulselessness, paralysis, paraesthesia and coldness in the leg
- Worsening intermittent claudication
- Rest pain in foot
- Presence of popliteal aneurysm
- Superficial thrombophlebitis above knee
- Evidence of DVT
- Suspicion of gas gangrene in leg
- Worsening hip pain
- Evidence of disease in bone (e.g. neoplasia, infection, Paget's)
- Severe sciatica with neurological deficit (e.g. floppy foot, absent reflexes)

Practice tips

- Always X-ray the legs (including hips) of a patient complaining of unusual deep leg pain, especially a child.
- Pain that does not fluctuate in intensity with movement, activity or posture has an inflammatory or neoplastic cause.

- Hip disorders such as osteoarthritis and slipped femoral epiphysis can present as pain in the knee (usually medial aspect).
- Consider retroperitoneal haemorrhage as a cause of acute severe nerve root pain, especially in people on anticoagulant therapy.
- Avoidance of amputation with acute lower limb ischaemia depends on early recognition (surgery within 4 hours - too late if over 6 hours).

REFERENCES

1. House AK. The painful limb: is it intermittent claudication? Mod Med Aust, 1990: November; 16 - 26.
2. Tunnessen WW. *Signs and Symptoms in Paediatrics* (2nd edn). Philadelphia: Lippincott, 1988: 483.
3. Hart FD. *Practical Problems in Rheumatology*. London: Dunitz, 1983: 120.
4. Bates B. *A Guide to Physical Examination and History Taking* (5th edn). New York: Lippincott, 1991: 450.
5. Fry J, Berry H. *Surgical Problems in Clinical Practice*. London: Edward Arnold, 1987: 125 - 34.
6. Ryan P. *A Very Short Textbook of Surgery* (2nd edn). Canberra: Dennis & Ryan, 1990: 61.
7. Hunt P, Marshall V. *Clinical Problems in General Surgery*. Sydney: Butterworths, 1991: 172.
8. Davis A, Bolin T, Ham J. *Symptom Analysis and Physical Diagnosis* (2nd edn). Sydney: Pergamon, 1990: 179.
9. Colucciello SA. Evaluation and management of deep venous thrombosis. Primary Care Rep, 1996; 2(12): 105.
10. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 12). Melbourne: Therapeutic Guidelines Ltd, 2003: 218 - 2.
11. Murtagh JE. *Practice Tips* (4th edn). Sydney: McGraw-Hill, 2004: 274.
12. Barton S ed. *Clinical Evidence*. London: BMK Publishing Group, 2001: 768 - 9.

CHAPTER 67. THE PAINFUL KNEE

The human knee is a joint and not a source of entertainment.

PERCY HAMMOND (1912), *Review of a Play*

THE PAINFUL KNEE

The knee, which is a gliding hinge joint, is the largest synovial joint in the body. Its small area of contact of the bone ends at any one time makes it dependent on ligaments for its stability. Although this allows a much increased range of movement it does increase the susceptibility to injury, particularly from sporting activities. Finding the cause of a knee problem is one of the really difficult and challenging features of practice. It is useful to remember that peripheral pain receptors respond to a variety of stimuli. These include inflammation due either to inflammatory disorders or chemical irritation such as crystal synovitis, traction pain (e.g. trapped meniscus stretching the capsule), tension on the synovium capsule (e.g. effusion or haemarthrosis), and impact loading of the subchondral bone.

KEY FACTS AND CHECKPOINTS

- Disorders of the knee account for about one presentation per fifty patients per year.[1]
- The commoner presenting symptoms in order of frequency are pain, stiffness, swelling, clicking and locking.[1]
- The age of presentation of a painful knee has varied significance as many conditions are age-related.
- Excessive strains across the knee, such as a valgus-producing force, are more likely to cause ligament injuries, while twisting injuries tend to cause meniscal tears.
- A ruptured anterior cruciate ligament (ACL) is a commonly missed injury of the knee.[2] It should be suspected with a history of either a valgus strain or a sudden pivoting of the knee, often associated with a cracking or popping sensation. It is often associated with the rapid onset of haemarthrosis or inability to walk or weight bear.
- A rapid onset of painful knee swelling (minutes to 1 - 4 hours) after injury indicates blood in the joint - *haemarthrosis*.
- Swelling over 1 - 2 days after injury indicates synovial fluid - *traumatic synovitis*.
- Any collateral ligament repair should be undertaken early but, if associated with ACL injuries, early surgery may result in knee stiffness. Thus, surgery is often delayed. With isolated ACL ruptures, early reconstruction is appropriate in the high-performance athlete; otherwise, delayed reconstruction is appropriate if there is clinical instability.[3]
- Acute spontaneous inflammation of the knee may be part of a systemic condition such as rheumatoid arthritis, rheumatic fever, gout, pseudogout (chondrocalcinosis), a spondyloarthropathy (psoriasis, ankylosing spondylitis, Reiter's syndrome, bowel inflammation), Lyme disease and sarcoidosis.
- Consider Osgood-Schlatter disorder (OSD) in the prepubertal child (especially a boy aged 10 - 14) presenting with knee pain.
- Disorders of the lumbosacral spine (especially L3 to S1 nerve root problems) and of the hip joint (L3 innervation) refer pain to the region of the knee joint.
- If infection or haemorrhage is suspected the joint should be aspirated.
- The condition known as anterior knee pain is the commonest type of knee pain and accounts for at least 11% of sports-related musculoskeletal problems. The prime cause of this is patellofemoral dysfunction pain. It is a benign condition with a good prognosis.

THE KNEE AND REFERRED PAIN - KEY KNOWLEDGE

Pain from the knee joint

Disorders of the knee joint give rise to pain felt accurately at the knee, often at some particular part of the joint, and invariably in the anterior aspect, very seldom in the posterior part of the knee. An impacted loose body complicating osteoarthritis and a radial tear of the lateral meniscus[4] are the exceptional disorders liable to refer pain proximally and distally in the limb, but the problems obviouslyoriginate from the knee.

Pain referred to the knee

Referred pain to the knee or the surrounding regionis a time-honoured trap in medicine. The two classicproblems are disorders of the hip joint and lumbosacralspine.

- The hip joint is mainly innervated by L3, henceapin is referred from the groin down the frontand medial aspects of the thigh to the knee(Figure 67.1). Sometimes the pain can beexperienced on the anteromedial aspect of theknee only. It is not uncommon for childrenwith a slipped upper femoral epiphysis topresent with a limp and knee pain.



Figure 67.1 Possible area of referred pain from disorders of the hip joint

- Knee pain can be referred from thelumbosacral spine. Patients with disclesions may notice that sitting, coughing orstraining hurts the knee, whereas walkingdoes not.

L3 nerve root pressure from an L2 - 3 disc prolapse(uncommon) and L4 nerve root pain will causeanteromedial knee pain; L5 reference from an L4 - 5disc prolapse can cause anterolateral knee pain, whileS1 reference from an L5 - S1 prolapse can cause painat the back of the knee (Figure 67.1).

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 67.1.

Table 67.1 The painful knee: diagnostic strategy model

Q. Probability diagnosis
A. Ligament strains and sprains
± traumatic synovitis
Osteoarthritis
Patellofemoral syndrome
Prepatellar bursitis

Q. Serious disorders not to be missed

A. Acute cruciate ligament tear

Vascular disorders

- deep venous thrombosis
- superficial thrombophlebitis

Neoplasia

- primary in bone
- metastases

Severe infections

- septic arthritis
- tuberculosis

Rheumatoid arthritis

Juvenile chronic arthritis

Rheumatic fever

Q. Pitfalls (often missed)

A. Referred pain: back or hip

Foreign bodies

Intra-articular loose bodies

Osteochondritis dissecans

Osteonecrosis

Osgood-Schlatter disorder

Meniscal tears

Fractures around knee

Pseudogout (chondrocalcinosis)

Gout → patellar bursitis

Ruptured popliteal cyst

Rarities

Sarcoidosis

Paget's disease

Spondyloarthropathy

Q. Seven masquerades checklist

A. Depression

Diabetes

Drugs - (indirect)

Anaemia -

Thyroid disorder -

Spinal dysfunction

UTI -

Q. Is this patient trying to tell me something?

A. Psychogenic factors relevant, especially with possible injury compensation.

Probability diagnosis

A UK study[1] has highlighted the fact that the commonestcause of knee pain is simple ligamentousstrains and bruises due to overstress of the knee orother minor trauma. Traumatic synovitis may accompanysome of these injuries. Some of these so-calledstrains may include a variety of recently describedsyndromes, such as the synovial plica syndrome, patellar tendonitis and infrapatellar fat-pad inflammation(Figure 67.2).



Figure 67.2 Lateral view of knee showing typical sites of various causes of knee pain

Low-grade trauma of repeated overuse, such asfrequent kneeling, may cause prepatellar bursitisknown variously as ‘housemaid's knee’ or ‘carpetlayer's knee’. Infrapatellar bursitis is referred to as‘clergyman's knee’.

Osteoarthritis of the knee, especially in the elderly, is a very common problem. It may arise spontaneouslyor be secondary to previous trauma with associatedinternal derangement and instability.

The most common overuse problem of the knee isthe patellofemoral joint pain syndrome (often previouslyreferred to as chondromalacia patellae).

Serious disorders not to be missed

Neoplasia in the bones around the knee is relativelyuncommon but still needs consideration. The commonestneoplasias are secondaries from the breast, lung, kidney, thyroid and prostate. Uncommonexamples include osteoid osteoma, osteosarcomaand Ewing's tumour (more likely in younger people). Septic arthritis and infected bursitis are prone tooccur in the knee joint, especially following contaminatedlacerations and abrasions. Septic arthritis fromblood-borne infection can be of the primary type inchildren, where the infection is either staphylococcalor due to Haemophilus influenzae, or gonococcalarthritis in adults. Rheumatic fever should be kept inmind with a fleeting polyarthritis that involves theknees and then affects other joints.

Inflammatory disorders such as spondyloarthropathies, sarcoidosis, chondrocalcinosis (a crystalarthropathy due to calcium pyrophosphate dihydratein the elderly), gout and juvenile chronic arthritis haveto be considered in the differential diagnosis.

Pitfalls

There is a myriad of pitfalls in knee joint disorders, often arising from ignorance, because there is a myriad of problems that are difficult to diagnose. Fortunately, many of these problems can be diagnosed by X-ray. A particular trap is a foreign body, such as a broken needle acquired by kneeling on carpet.

The presence of a spontaneous effusion demands careful attention because it could represent a rheumatic disorder or conditions such as osteochondritis dissecans (more common in the young) or osteonecrosis of the femoral condyle (a necrotic problem in the elderly) and perhaps a subsequent loose body in the joint.

A ruptured Baker's cyst will cause severe pain behind the knee and can be confused with deep venous thrombosis. It is important to bear in mind complications of varicose veins, which can cause pain or discomfort around the knee joint.

General pitfalls

- Overlooking referred pain from the hip or low back as a cause of knee pain
- Failing to realise that meniscal tears can develop due to degeneration of the menisci with only minimal trauma
- Failing to X-ray the knee joint and order special views to detect specific problems, such as a fractured patella or osteochondritis dissecans

Ottawa knee rules for X-ray of an injured knee

- Patient aged 55 years or moret
- Isolated tenderness of the patella
- Tenderness at the head of the fibula
- Inability to flex to 90°
- Immediate inability to weight bear

Seven masquerades checklist

Of these, spinal dysfunction is the prime association. Diabetes may cause pain through a complicating neuropathy and drugs such as diuretics may cause gout in the elderly.

Psychogenic considerations

Patients, young and old, may complain of knee pain, imaginary or exaggerated, to gain attention, especially if compensation for an injury is involved. This requires discreet clinical acumen to help patients work through the problem.

THE CLINICAL APPROACH

History

The history is the key to diagnosis. If any injury is involved careful description of the nature of the injury is necessary. This includes past history. A special problem relates to the elderly who can sustain knee injuries after a ‘drop attack’, but attention can easily be diverted away from the knee with preoccupation with the cerebral pattern.

It is relevant to define whether the pain is acute or chronic, dull or sharp, and continuous or recurring. Determine its severity and position and keep in mind age-related causes.

Key questions

Related to an injury

- Can you explain in detail how the injury happened?
- Did you land awkwardly after a leap in the air?
- Did you get a direct blow? From what direction?

- Did your leg twist during the injury?
- Did you feel a ‘pop’ or hear a ‘snap’?
- Did your knee feel wobbly or unsteady?
- Did the knee feel as if the bones separated momentarily?
- How soon after the injury did the pain develop?
- How soon after the injury did you notice swelling?
- Have you had previous injury or surgery to the knee?
- Were you able to walk after the injury or did you have to be carried off the ground or court?
- Does this involve work care compensation?

No history of injury

- Does the pain come on after walking, jogging or other activity?
- How much kneeling do you do? Scrubbing floors, cleaning carpets?
- Could there be needles or pins in the carpet?
- Does your knee lock or catch?
- Does swelling develop in the knee?
- Does it ‘grate’ when it moves?
- Does the pain come on at rest and is there morning stiffness?
- Do you feel pain when you walk on steps or stairs?

Significance of symptoms

Swelling after injury

The sudden onset of painful swelling (usually within 60 minutes) is typical of haemarthrosis (Figures 67.3 and 67.4). Bleeding occurs from vascular structures such as torn ligaments, torn synovium or fractured bones, while injuries localised to avascular structures such as menisci do not usually bleed. About 75% of cases are due to ACL tears.[5] If a minor injury causes acute haemarthrosis suspect a bleeding diathesis or anticoagulant usage. The causes of haemarthrosis are listed in Table 67.2.



Figure 67.3 Haemarthrosis in a sportsman presenting with an acutely painful swollen knee



Figure 67.4 Haemarthosis: Surgical release of intraarticular blood under pressure in the knee shown in Figure 67.3

Table 67.2 Causes of haemarthrosis

Torn cruciate ligaments, esp. ACL
Capsular tears with collateral ligament tears
Peripheral meniscal tears
Dislocation or subluxation of patella
Osteochondral fractures
Bleeding disorders (e.g. haemophilia), anticoagulants

Swelling of an intermediate rate of onset, stiffness and pain in the order of hours (e.g. 6 - 24 hours) is typical of an effusion of synovial fluid. Causes include meniscal tears and milder ligamentous injuries. Swelling gradually developing over days and con - fined to the anterior knee is typical of bursitis such as ‘housemaid's knee’.

Recurrent or chronic swelling

This indicates intra-articular pathology and includes:

- patellofemoral pain syndrome
- osteochondritis dissecans
- degenerative joint disease including degenerative meniscus tears
- arthritides

Locking

Locking usually means a sudden inability to extend the knee fully (occurs at 10 - 45°, average 30°) but ability to flex fully.[6]

Causes

True locking:

- torn meniscus (bucket handle)
- loose body (e.g. bony fragment from osteochondritis dissecans)
- torn ACL (remnant)
- avulsed anterior tibial spine
- dislocated patella
- synovial osteochondromatosis Pseudo-locking:
- patello-femoral disorders
- first or second degree medial ligament tear
- strain of ACL
- gross effusion
- pain and spasm of hamstrings

Catching

'Catching' of the knee implies that the patient feels that something is 'getting in the way of joint movement' but not locking. Causes include any of the conditions that cause locking, but a subluxing patella and loose bodies in particular must be considered.

Causes of loose bodies

- Osteochondritis dissecans (usually lateral side of medial femoral condyle)
- Retropatellar fragment (e.g. from dislocation of patella)
- Dislodged osteophyte
- Osteochondral fracture - post injury
- Synovial chondromatosis

Clicking

Clicking may be due to an abnormality such as patellofemoral maltracking or subluxation, a loose intra-articular body or a torn meniscus, but can occur in normal joints when people climb stairs or squat.

Anterior knee pain

Common causes include:

- patellofemoral syndrome
- osteoarthritis of the knee
- patellar tendonopathy
- osteonecrosis

Lateral knee pain

Consider:

- osteoarthritis of lateral compartment of knee
- lesions of the lateral meniscus
- patellofemoral syndrome

Medial knee pain

Consider:

- osteoarthritis of medial compartment of knee
- lesions of the medial meniscus
- patellofemoral syndrome

Physical examination

The provisional diagnosis may be evident from a combination of the history and simple inspection of the joint but the process of testing palpation, movements (active and passive) and specific structures of the knee joint helps to pinpoint the disorder.

Inspection

Inspect the knee with the patient walking, standing erect and lying supine. Get the patient to squat to help localise the precise point of pain. Get the patient to sit on the couch with legs hanging over the side and note any abnormality of the patella. Note any deformities, swelling or muscle wasting.

The common knee deformities are genu valgum 'knock knees' (Figure 67.5a), genu recurvatum 'back knee' (Figure 67.5b) and genu varum 'bowed legs' (Figure 67.5c).



Figure 67.5 Knee deformities; (a) genu valgum ('knock knees'): tibia deviates laterally from knee; (b) genu recurvatum ('back knee'); (c) genu varum ('bowed legs')

A useful way of remembering the terminology is to recall that the 'l' in valgus stands for 'l' in lateral.[7] In the normal knee the tibia has a slight valgus angulation in reference to the femur, the angulation being more pronounced in women.

Palpation

Palpate the knee generally concentrating on the patella, patella tendon, joint lines, tibial tubercle, bursae and popliteal fossa.

Palpate for presence of any fluid, warmth, swelling, synovial thickening, crepitus, clicking and tenderness. Feel for a popliteal (Baker's) cyst in the popliteal fossa. Draw the fingers upwards over the suprapatellar pouch: synovial thickening, a hallmark of chronic arthritis, is most marked just above the patella - it feels warm, boggy, rubbery and has no fluid thrill.

Flex the knees to 45° and check for a pseudocyst, especially of the lateral meniscus (Figure 67.6).



Figure 67.6 Pseudocyst of the lateral meniscus: flex the knees to 45° to force lump (if present) to appear

Fluid effusion

The bulge sign: compress the medial side of the joint and evacuate any fluid. The test is positive when the lateral side of the joint is then stroked and the fluid is displaced across the joint, creating a visible bulge or filling of the medial depression (Figure 67.7).



Figure 67.7 The bulge sign with a knee effusion: fluid bulges into the medial compartment

The test will be negative if the effusion is gross and tense, in which case the *patellar tap test* (Figure 67.8) is used by sharply tapping the lower pole of the patella against the femur with the index finger. A positive tap is when the patella can be felt to tap against the femur and then float free.



Figure 67.8 The patellar tap test

Movements

Extension: normal is 0 - 5°. The loss of extension is best measured by lifting the heel off the couch with the knee held down. In the normal knee the heel will lift 2.5 - 4 cm off the couch, that is, into hyperextension.

Flexion (supine or prone): normal to 135°. The normal knee flexes heel to the buttock but in locking due to medial meniscus tears there may be a gap of 5 or more centimetres between the heel and buttock.

Rotation: normal 5 - 10°. Test at 90° with patient sitting over the edge of the couch; rotate the feet with the hand steadying the knee.

Note: Normally, no abduction, adduction or rotation of the tibia on the femur is possible with the leg fully extended.

Ligament stability tests

Collateral ligaments. Adduction (varus) and abduction (valgus) stresses of the tibia on the femur are applied in full extension and then at 30° flexion with the leg over the side of the couch. With ligament strains there is localised pain when stressed. With a complete (third degree) tear the joint will open out. This end-point feel should be carefully noted: firmness indicates stability, 'mushiness' indicates damage (Figure 67.9).



Figure 67.9 Medial and lateral ligament instability: (a) medial instability of knee joint; (b) lateral instability of knee joint

Cruciate ligaments. Stability of the ACL can be tested with the anterior drawer test. This is done with the patient supine and the knee flexed to 90°. The tibia is pulled forwards off the femur and in the presence of a cruciate ligament injury there will be increased gliding of the tibia on the femur. An aberrant positive sign can occur in the presence of posterior cruciate ligament (PCL) insufficiency, in which case the knee is actually brought back to its normal site from a dropped-back position. This gives the appearance of a positive anterior drawer sign. In that situation, a Lachman's test will be negative. In the presence of medial ligament injury, the increased external rotation of the tibia against the femur may add to the positive drawer sign.

Specific provocation tests

The simplest menisci function tests are those outlined in Table 67.4 at p. 737.

Table 67.4 Typical symptoms and signs of meniscal injuries

	Medial meniscus tear	Lateral meniscus tear
Mechanism	• Abduction (valgus) force • External rotation of lower leg on femur	• Adduction (varus) force • Internal rotation of leg on femur

Symptoms		
1 Knee pain during and after activity	Medial side of knee	Lateral side of knee
2 Locking	Yes	Yes
3 Effusion	+ or -	+ or -
Signs		
1 Localised tenderness over joint line (with bucket handle tear)	Medial joint line	Lateral joint line (may be cyst)
2 Pain on hyperextension of knee	Medial joint line	Lateral joint line
3 Pain on hyperflexion of knee joint	Medial joint line	Lateral joint line
4 Pain on rotation of lower leg (knee at 90°)	On external rotation	On internal rotation
5 Weakened or atrophied quadriceps	May be present	May be present

- *McMurray's test*. The patient lies on the couch and the flexed knee is rotated in varying degrees of abduction as it is straightened into extension. A hand over the affected knee feels for ‘clunking’ or tenderness.
- *Apley grind/distraction test*. The patient lies prone, the knee is flexed to 90° and then rotated under a compression force. Reproduction of painful symptoms may indicate meniscal tear. Then repeat the rotation under distraction - tests ligament damage.
- *Patella apprehension test*. At 15 - 20° flexion, attempt to push the patella laterally and note the patient's reaction.
- *Patellar tendonitis*. Palpate patellar tendon (refer to Figure 67.19 at p. 741).



Figure 67.19 Patellar tendonitis: method of palpation

- *Patellofemoral pain test*. Refer to Figure 67.18 at p. 741.



Figure 67.18 Special sign of the patellofemoral pain syndrome

Examine the lumbosacral spine and the hip joint of the affected side.

Measurements

Quadriceps. For suspected quadriceps wasting, measure the circumference of the thighs at equal points above the tibial tuberosity. It is helpful to assess quadriceps function by feeling the tone. *Static Q angle* (Figure 67.10).


Q
long axis of femur
Q angle
midpoint of patella
line from midpoint of tibial tuberosity to midpoint of patella

Figure 67.10 The Q angle of the knee gives a measure of patellar alignment

If the Q angle is > 15° in men and > 19° in women there is a predisposition to patellofemoral pain and instability.[8]

Investigations

Investigation for the diagnosis of knee pain can be selected from:

- blood tests
  - RA factor tests; ANA; HLA B27

  - ESR
  - blood culture (suspected septic arthritis)
- radiology[9]
  - plain X-ray
  - special views
    - intercondylar (osteochondritis dissecans, loose bodies)
    - tangential (or skyline view for suspected patella pathology)
    - oblique (to define condyles and patella)
    - weight-bearing views looking for degenerative arthritis
  - bone scan: for suspected tumour, stress fracture, osteonecrosis, osteochondritis dissecans
  - MRI: excellent for diagnosing cartilage and menisci disorders and ligament damage; the investigation of choice for internal 'derangement'
  - arthrography (generally superseded by arthroscopy) or MRI
  - ultrasound: good for assessment of patellar tendon, soft tissue mass, fluid collection, Baker's cyst and bursae
- CT: useful for complex fractures of tibial plateau and patellofemoral joint special dysfunction
- special
  - examination under anaesthesia
  - arthroscopy
  - knee aspiration: culture or crystal examination

Fractures that may be missed on plain films[9]

- Patellar fracture
- Tibial plateau fracture
- Tibial spine fracture
- Epiphyseal injuries in children
- Osteochondral fracture
  - patella
  - femoral condyle
- Stress fracture upper tibia
- Avulsion fracture (e.g. segond fracture of upper lateral tibia, with ACL tear

KNEE PAIN IN CHILDREN

Children may present with unique conditions that are usually related to growth, including epiphyseal problems. Their tendency towards muscle tightness, especially in the growth spurt, predisposes them to overuse injuries such as patellar tendonitis and patellofemoral pain syndrome.

First decade

A painful knee during the first decade of life (0 - 10 years) in non-athletes is an uncommon presenting symptom, but suppurative infection and juvenile chronic arthritis have to be considered.

Genu valgum or varum is a common presentation but usually not a source of discomfort for the child. However, genu valgum, which is often seen around 4 - 6 years, may predispose to abnormal biomechanical stresses which contribute to overuse-type injuries if the child is involved in sport.

Second decade

Pain in the knee presents most frequently in this decade and is most often due to the patellofemoral syndrome,[10] which is related to the retropatellar and peripatellar regions and usually anterior to the knee. It occurs in the late teenage years of both sexes.

An important problem is subluxation of the patella, typically found in teenage girls. It is caused by maltracking of the patellofemoral mechanism without complete dislocation of the patella (Figure 67.11).



Figure 67.11 Lateral subluxation of the patella

On examination, the patella is usually in a high and lateral position. Surgery may be required if symptoms persist.

OSD is common in pre-pubertal adolescent boys but can occur in those aged 10 - 16 years.

Other conditions found typically in this age group include:

- slipped upper femoral epiphysis - usually in middle teenage years after a growth spurt
- anserinus (‘goose foot’) bursitis
- osteochondritis dissecans

Age-related causes of the painful knee are presented in Table 67.3.[10]

Table 67.3 Age-related causes of painful knee

First decade (0 - 10 years)
Infection
Juvenile chronic arthritis
Second decade (10 - 20 years)
Patellofemoral syndrome
Subluxation/dislocation of patella
Slipped femoral epiphysis (referred)
‘Hamstrung’ knee
Osteochondritis dissecans
Osgood-Schlatter disorder
Anserinus tendonitis

Third decade (20 - 30 years)
Bursitis
Mechanical disorders
Fourth and fifth decades (30 - 50 years)
Cleavage tear of medial meniscus
Radial tear of lateral meniscus
Sixth decade and older (50 years and over)
Osteoarthritis
Osteonecrosis
Paget's disease (femur, tibia or patella)
Anserinus bursitis
Chondrocalcinosis and gout
Osteoarthritis of hip (referred pain)

The little athlete

Children competing in sporting activities, especially running and jumping, are prone to overuse injuries such as the patellofemoral pain syndrome, traumatic synovitis of the knee joint and OSD. Haemarthrosis can occur with injuries, sometimes due to a synovial tear without major joint disruption. If knee pain persists, especially in the presence of an effusion, X-rays should be performed to exclude osteochondritis of the femoral condyle.[11]

The Ottawa knee rules

A knee X-ray series is only required for children with any of these findings:

- isolated tenderness of the patella (no bone tenderness of the knee other than the patella)
- tenderness at the head of the fibula
- inability to flex to 90°
- inability to bear weight both immediately and in the emergency department (four steps; unable to transfer weight twice onto each lower limb regardless of limping)

Osgood-Schlatter disorder

OSD is a traction apophysitis resulting from repetitive traction stresses at the insertion of the patellar tendon into the tibial tubercle, which is vulnerable to repeated traction in early adolescence.

Clinical features

- Commonest ages 10 - 14 years
- Boys:girls ° 3:1
- Bilateral in about one-third of cases
- Common in sports involving running, kicking and jumping
- Localised pain in region of tibial tubercle during and after activity
- Aggravated by kneeling down and going up and downstairs
- Development of lump in area
- Localised swelling and tenderness at affected tubercle
- Pain reproduced by attempts to straighten flexed knee against resistance

X-ray to confirm diagnosis (widening of the apophysis and possible fragmentation of bone) and exclude tumour or fracture (see Figure 67.12).



Figure 67.12 Features of Osgood-Schlatter disorder

Management

This is conservative as it is a self-limiting condition (6 - 18 months: average 12 months).

- If acute, use ice packs and analgesics.
- Main approach is to abstain from or modify active sports.
- Localised treatments such as electrotherapy are unnecessary.
- Corticosteroid injections should be avoided.[12]
- Plaster cast immobilisation should also be avoided.
- Surgery may be used (rarely) if an irritating ossicle persists[13] after ossification.
- Gentle quadriceps stretching.
- Graded return to full activity.

Prevention

- Promote awareness and early recognition of OSD.
- Program of stretching exercises for quadriceps mechanism in children in sport.

OSTEOCHONDRITIS DISSECANS: JUVENILE FORM[6]

This commonly occurs in adolescent boys aged 5 - 15 years whereby a segment of articular cartilage of the femoral condyle (85%) undergoes necrosis and may eventually separate to form an intra-articular loose body (Figure 67.13).



Figure 67.13 Osteochondritis dissecans: on X-ray, sclerosis of the lateral aspect of the medial condyle

It usually presents as pain and effusion and locking.

If the fragment has separated, surgery to reattach it can be contemplated.

KNEE PAIN IN THE ELDERLY

Rheumatic disorders are very common and responsible for considerable pain or discomfort, disability and loss of independence in the elderly.

Osteoarthritis is the most common cause and excellent results are now being obtained using total knee replacement in those severely affected.

The elderly are particularly prone to crystalassociated joint diseases, including monosodium urate (gout), CPPD (pseudogout) and hydroxyapatite (acute calcific periarthritis).

Chondrocalcinosis of knee (pseudogout)

The main target of CPPD is the knee, where it causes chondrocalcinosis. Unlike gout, chondrocalcinosis of the knee is typically a disorder of the elderly with about 50% of the population having evidence of involvement of the knee by the ninth decade.[14] Most cases remain asymptomatic but patients (usually aged 60 or older) can present with an acutely hot, red, swollen joint resembling septic arthritis.

Investigations include aspiration of the knee to search for CPPD crystals, and X-ray. If positive, consider an associated metabolic disorder such as haemochromatosis, hyperparathyroidism or diabetes mellitus. The treatment is similar to acute gout although colchicine is less effective. Acute episodes respond well to NSAIDs or intra-articular corticosteroid injection.

Osteonecrosis [6, 15]

Spontaneous osteonecrosis of the knee (SONK) is more common after the age of 60, especially in females; it can occur in either the femoral (more commonly) or tibial condyles. The aetiology is unknown. The sudden onset of pain in the knee, with a normal joint X-ray, is diagnostic of osteonecrosis. However, the X-ray (especially later) will demonstrate an area of osteonecrosis. The pain is usually persistent, with swelling and stiffness, and worse at night. It can take three months for the necrotic area to show radiologically although a bone scan or MRI may be positive at an early stage (Figure 67.14). The condition may resolve in time with reduction of weight-bearing. Surgery in the form of subchondral drilling may be required for persistent pain in the early stages.



Figure 67.14 Osteonecrosis: necrosis in the medial femoral condyle can take three months to show radiologically

Osteochondritis dissecans: adult form[6]

The adult form occurs more often in males and may be the result of cysts of osteoarthritis fracturing into the joint. Up to 30% are bilateral. Symptoms depend on whether the osteochondral fragment becomes separated. A loose fragment may produce locking or collapse of the knee.

Loose bodies[6]

The large knee joint is a ‘haven’ for intra-articular loose bodies, which may be formed from bone, cartilage or osteochondral fragments following injury (‘chip’ fragment), osteochronditis dissecans, osteoarthritis, synovial chondromatosis or other conditions. They may be asymptomatic but usually cause clicking or locking with swelling. Diagnosis is by X-ray and surgical removal is necessary for recurrent problems.

The knee ‘mouse’

This common complaint is usually a result of a pedunculated fibrous lump in the prepatella bursa, often secondary to trauma, such as falls onto the knee.

ACUTE INJURIES

Meniscal tears

Medial and lateral meniscal tears are usually caused by abduction and adduction forces causing the meniscus to be compressed between the tibial and femoral condyles and then subjected to a twisting force.

The medial meniscus is three times more likely to be torn than the lateral. These injuries are common in contact sports and are often associated with ligamentous injuries. Suspect these injuries when there is a history of injury with a twisting movement with the foot firmly fixed on the ground.

However, pain in the knee can present in the patient aged 30 - 50 years as the menisci degenerate, with resultant cleavage tears from the posterior horn of the medial meniscus and ‘parrot beak’ tears of the mid-section of the lateral meniscus. These problems cause pain because these particular deformities create tension on the joint capsule and stretch the nerve ends. X-rays are not specifically useful but an MRI scan should confirm diagnosis.

Clinical features

- General symptoms[8]
    - joint line pain (49%)
    - locking (17%)
    - swelling (14%)
    - loss of movement
        - restricted flexion
        - loss of last 5 - 10° extension
- Parrot beak tear of lateral meniscus
    - pain in the lateral joint line
    - pain radiating up and down the thigh
    - pain worse with activity
    - a palpable and visible lump when the knee is examined at 45°

Arthroscopic partial meniscectomy offers relief. The peripheral meniscus is vascular and can be repaired within 6 - 12 weeks of injury.[15]

- Cleavage tear of medial meniscus
    - pain in medial joint line
    - pain aggravated by slight twisting of the joint
    - pain provoked by patient lying on the side and pulling the knees together
    - pain worse with activity

Arthroscopic meniscectomy is appropriate treatment, but some do settle with a trial of physiotherapy.

A diagnostic memoire

Table 67.4 is a useful aid in the diagnosis of these injuries. There is a similarity in the clinical signs between the opposite menisci, but the localisation of pain in the medial or lateral joint lines helps to differentiate between the medial and lateral menisci.

Note: The diagnosis of a meniscal injury is made if three or more of the five examination findings (‘signs’ in Table 67.4) are present.

Ligament injuries

Tears of varying degrees may occur in the:

- anterior cruciate ligament
- posterior cruciate ligament
- medial collateral ligament
- lateral collateral ligament

Anterior cruciate ligament rupture

This is a very serious and disabling injury which may result in chronic instability. Chronic instability can result in degenerative joint changes if not dealt with adequately. Early diagnosis is essential but there is a high misdiagnosis rate. Sites of ACL rupture are shown in Figure 67.15.



Figure 67.15 Sites of rupture of the anterior cruciate ligament

Mechanisms

- Sudden change in direction with leg in momentum
- Internal tibial rotation on a flexed knee (commonest)
- Valgus force (e.g. a rugby tackle)
- May be associated with collateral ligament tears and meniscus injuries

Clinical features

- Onset of severe pain after a sporting injury, such as landing from a jump, or a forced valgus rotational strain of the knee when another player falls across the abducted leg
- Immediate effusion of blood, usually within 30 minutes
- common sports: contact sports - rugby, football and soccer, basketball, volleyball, skiing
- Differential diagnosis is a subluxed or dislocated patella
- Subsequent history of pain and ‘giving way’ of the knee

Examination

- Gross effusion
- Diffuse joint line tenderness
- Joint may be locked due to effusion, anterior cruciate tag or associated meniscal (usually medial) tear
- Ligament tests
    - anterior drawer: negative or positive

- pivot shift test: positive (only if instability)
- Lachman test: lacking an end point

Note: It may be necessary to examine the knee under anaesthesia, with or without arthroscopy, to assess the extent of injury.

The Lachman test

This test is emphasised because it is a sensitive and reliable test for the integrity of the ACL. It is an anterior drawer test with the knee at 15 - 20° of flexion. At 90° of flexion, the draw may be negative but the anterior cruciate torn.

Functional instability due to anterior cruciate deficiency is best elicited with the pivot shift test. This is more difficult to perform than the Lachman test.

Method

1. The examiner should be positioned on the same side of the examination couch as the knee to be tested.
2. The knee is held at 15 - 20° of flexion by placing a hand under the distal thigh and lifting the knee into 15 - 20° of flexion.
3. The patient is asked to relax, allowing the knee to 'fall back' into the steadying hand and roll slightly into external rotation.
4. The anterior drawer is performed with the second hand grasping the proximal tibia from the medial side (Figure 67.16) while the thigh is held steady by the other hand. The examiner's knee can be used to steady the thigh.



Figure 67.16 The Lachman test

5. The feel of the end point of the draw is carefully noted. Normally there is an obvious jar felt as the anterior cruciate tightens. In an anterior cruciate deficient knee there is excess movement and no firm end point. The amount of draw is compared with the opposite knee. Movement greater than 5 mm is usually considered abnormal.

Pivot shift test

This is an important test for anterolateral rotatory instability. It is positive when anterior cruciate injuries are sufficient to produce a functional instability.

Method

- The tibia is held in internal rotation by grasping the ankle firmly, with the knee in full extension.
- A valgus force is applied to the knee with the hand placed on the lateral aspect of the knee just below it (this maximises subluxation in the presence of an ACL tear).
- The knee is then flexed from 0 - 90°, listening for a 'clunk' of reduction. The test is positive when there is a sudden change of rhythm during flexion which corresponds to relocation of the subluxed knee. This usually occurs between 30° and 45° of flexion.
- From this flexed position the knee is extended, seeking a click into subluxation. This is called a positive jerk test.

Management[16]

The management depends on the finding by the surgeon. Surgical repair is reserved for complete ligament tears. This usually involves reconstruction of the ligament using patellar or hamstring tendons. Early reconstruction is appropriate in younger patients who participate in high levels of sporting activity in which it can be predicted that functional instability will be a problem. In less active people, a conservative approach is appropriate. The ACL may be trimmed. Cruciate reconstruction can then be undertaken if the knee becomes clinically unstable. The presence of an ACL injury with a significant medial ligament injury will necessitate reconstructive surgery but this is probably best delayed for some weeks as the subsequent incidence of knee stiffness is high.

Posterior cruciate ligament rupture

Mechanisms

- Direct blow to the anterior tibia in flexed knee
- Severe hyperextension injury

Clinical features

- Posterior (popliteal) pain, radiating to calf
- Usually no or minimal swelling
- Minimal disability apart from limitation of running or jumping
- Pain running downhill
- Recurvatum
- Posterior sag or draw

Management

- Usually managed conservatively with immobilisation and protection for 6 weeks
- Graduated weight-bearing and exercises

Medial collateral ligament rupture

Mechanisms

- Direct valgus force to knee - lateral side knee (e.g. rugby tackle from side)
- External tibial rotation (e.g. two soccer players kicking ball simultaneously)

Clinical features

These depend on the degree of tear (1st, 2nd or 3rd degree):

- pain on medial knee
- aggravated by twisting or valgus stress
- localised swelling over medial aspect
- pseudo-locking - hamstring strain
- ± effusion

- no end point on valgus stress testing (3rd degree) (Figure 67.9a at p. 732)

Note: Check lateral meniscus if MCL tear. Pellegrini - Stieda syndrome - calcification in haematoma at upper (femoral) origin of MCL may follow.

Management

If an isolated injury, this common injury responds to conservative treatment with early limited motion bracing to prevent opening of the medial joint line. Six weeks of limited motion brace at 20 - 70° followed by knee rehabilitation usually returns the athlete to full sporting activity within 12 weeks.

Note: The same principles of diagnosis and management apply to the less common rupture of the lateral collateral ligament which is caused by a direct varus force to the medial side of the knee. However, lateral ligament injuries tend to involve the cruciate ligament and reconstruction of both ligaments is usually necessary.[16]

Complex regional pain syndrome I

A localised complex regional pain syndrome I (also known as reflex sympathetic dystrophy) can follow a direct fall onto the knee.

Symptoms

- Hypersensitivity
- Full extension, loss of flexion
- Possible increasing sweating
- Tenderness of the joint

OVERUSE SYNDROMES

The knee is very prone to overuse disorders. The pain develops gradually without swelling, is aggravated by activity and relieved with rest. It can usually be traced back to a change in the sportsperson's training schedule, footwear or technique, or to related factors. It may also be related to biomechanical abnormalities ranging from hip disorders to feet disorders.

Overuse injuries include:

- patellofemoral pain syndrome (‘jogger's knee’, ‘runner's knee’)
- patellar tendonitis (‘jumper's knee’)
- anserinus tendonitis/bursitis
- semimembranous tendonitis/bursitis
- biceps femoris tendonitis
- quadriceps tendonitis/rupture
- popliteus tendonitis
- iliotibial band friction syndrome (‘runner's knee’)
- the hamstrung knee
- synovial plica syndrome
- infrapatellar fat-pad inflammation

It is amazing how often palpation identifies localised areas of inflammation (tendonitis or bursitis) around the knee, especially from overuse in athletes and in the obese elderly (Figure 67.17).


(a)
quadriceps tendonitis or rupture
iliotibial band friction syndrome
anserinus bursitis/ tendonitis
patellar tendonitis
biceps femoris tendonitis
Osgood-Schlatter disorder
(b)
quadriceps tendonitis or rupture
patella
patellofemoral joint pain syndrome
patellar tendonitis
medial collateral ligament
semimembranous tendonitis/bursitis
anserinus tendonitis/bursitis

Figure 67.17 Typical painful areas around the knee for overuse syndromes: (a) anterior aspect; (b) medial aspect

Patellofemoral pain syndrome

This syndrome, also known as chondromalacia patellae and referred to as ‘jogger's knee’, ‘runner's knee’ or ‘cyclist's knee’, is the most common overuse injury of the knee. There is usually no specific history of trauma. It may be related to biomechanical abnormalities and abnormal position and tracking of the patella (e.g. patella alta). It usually presents in 13 - 15 year old females with faulty knee mechanisms or in 50 - 70 year olds with osteoarthritis of the patellofemoral joint.[17]

Clinical features

- Pain behind patella or deep in knee
- Pain aggravated during activities that require flexion of knee under loading:

  - climbing stairs
  - walking down slopes or stairs
  - squatting
  - prolonged sitting
- The ‘movie’ sign: using aisle seat to stretch knee
- Crepitus around patella may be present

Signs of chondromalacia patellae

Patellofemoral crepitation during knee flexion and extension is often palpable, and pain may be reproduced by compression of the patella onto the femur as it is pushed from side to side with the knee straight or flexed (Perkins’ test).

Method for special sign (Figure 67.18)

- Have the patient supine with the knee extended.
- Grasp the superior pole of the patella and displace it inferiorly.
- Maintain this position and apply patellofemoral compression.
- Ask the patient to contract the quadriceps (it is a good idea to get the patient to practise quadriceps contraction before applying the test).
- A positive sign is reproduction of the pain under the patella and hesitancy in contracting the muscle.

Treatment

- Give reassurance and supportive therapy.
- Reduce any aggravating activity.
- Refer to a physiotherapist with expertise.
- Correct any underlying biomechanical abnormalities such as pes planus (flat feet) by use of orthotics and correct footwear.
- Employ quadriceps (especially) and hamstring exercises.
- Consider course (trial) of NSAIDs.

Patellar tendonitis (‘jumper's knee’)

‘Jumper's knee’, or patellar tendonitis (Figure 67.2 at p. 727) (also referred to as patellar tendonopathy), is a common disorder of athletes involved in repetitive jumping sports, such as high jumping, basketball, netball, volleyball and soccer. It probably starts as an inflammatory response around a small tear.

Clinical features

- Gradual onset of anterior pain
- Pain localised to below knee
- Pain eased by rest, returns with activity

The diagnosis is often missed because of the difficulty of localising signs. The condition is best diagnosed by eliciting localised tenderness at the inferior pole of the patella with the patella tilted. There may be localised swelling.

Method

- Lay the patient supine in a relaxed manner with the head on a pillow, arms by the side and quadriceps relaxed (a must).
- The knee should be fully extended.
- Tilt the patella by exerting pressure over its superior pole. This lifts the inferior pole.
- Now palpate the surface under the inferior pole. This allows palpation of the deeper fibres of the patellar tendon (Figure 67.19).

- Compare with the normal side.
- Very sharp pain is usually produced in the patient with patellar tendonitis.

Management

Early conservative treatment including rest from the offending stresses is effective. This includes adequate warm-up and warm-down. Training modification includes calf, hamstring and quadriceps muscle stretching. Modified footwear and a patellar tendon strap may be helpful in some cases. The use of NSAIDs and corticosteroid injections is disappointing. Chronic cases may require surgery.

Anserinus tendonitis/bursitis

Localised tenderness is found over the medial tibial condyle where the tendons of the sartorius, gracilis and semitendinosus insert into the bone. It is a common cause of knee pain in the middle aged or elderly, especially the overweight woman. Pain is aggravated by resisted knee flexion.

Semimembranous tendonitis/bursitis

This inflamed area is sited either at the tendon insertion or in the bursa between the tendon and the medial head of the gastrocnemius. It is an uncommon problem. The bursa occurs on the medial side of the popliteal fossa between the medial head of gastrocnemius and the semimembranous tendon. It often communicates with the knee joint and, if so, treat knee joint pathology. If not, one can give an injection of depot triamcinolone or betamethasone.

Biceps femoris tendonitis/bursitis

The tendon and/or the bursa that lies between the tendon insertion and the fibular collateral ligament at the head of the fibula may become inflamed due to overuse. It is usually encountered in sprinters.

Popliteus tendonitis

Tenosynovitis of the popliteus tendon may cause localised pain in the posterior or the posterolateral aspect of the knee. Tenderness to palpation is elicited with the knee flexed to 90°.

Iliotibial band syndrome

Inflammation develops over the lateral aspect of the knee where the iliotibial band passes over the lateral femoral condyle. An inflammed bursa can occur deep to the band. The problem, which is caused by friction of the iliotibial band on the bone, is common in long-distance runners, especially when running up and down hills, and cyclists. It presents with welllocalised lateral knee pain of gradual onset. Palpation reveals tenderness over the lateral condyle 1 - 2 cm above the joint line.

Treatment of tendonitis and bursitis (small area)

Generally (apart from patellar tendonitis), the treatment is an injection of local anaesthetic and longacting corticosteroids into and deep to the localised area of tenderness. In addition it is important to restrict the offending activity with relative rest and refer for physiotherapy for stretching exercises. Attention to biomechanical factors and footwear is important.

If conservative methods fail for iliotibial tract tendonitis, surgical excision of the affected fibres may cure the problem.

Prepatellar bursitis

Repetitive low-grade direct trauma, such as frequent kneeling, can cause inflammation with swelling of the bursa which lies between the anterior surface of the patella and the skin. ‘Housemaid's knee’, or ‘carpet layer's knee’, can be difficult to treat if rest from the trauma does not allow it to subside. If persistent, drain the fluid with a 23-gauge needle and then introduce 0.5 - 1 mL of long-acting corticosteroid. The presence of a bursa ‘mouse’ and persistent bursitis usually means that surgical intervention is required.

Acute bursitis may also be caused by acute infection, or one of the inflammatory arthropathies (e.g. gout, Reiter's syndrome).

Infrapatellar bursitis

'Clergyman's knee' is produced by the same mechanisms as patellar bursitis and can be involved with inflammatory disorders or infection. Treatment is also the same.

The hamstrung knee

Cross describes this condition in young active sportspeople (second decade)10 as one that causes bilateral knee pain and possibly a limp. It is caused by a failure to warm up properly and stretch the hamstring muscles, which become tender and tight during the growth spurt. A 6 week program of straight leg raising and hamstring stretching will alleviate the pain completely.

Synovial plica syndrome

This syndrome results from a synovial fold (an embryological remnant) being caught between the patella and the femur during walking or running. It causes an acute 'catching' knee pain of the medial patellofemoral joint (Figure 67.2 at p. 727) and sometimes a small effusion. It generally settles without treatment.

Infrapatellar fat-pad inflammation

Acute compression of the fat-pad, which extends across the lower patella deep to the patellar tendon and into the knee joint (Figure 67.2 at p. 727), during a jump or other similar trauma, produces local pain and tenderness similar to the sensation of kneeling on a drawing pin.[18]

The pain usually settles without therapy over a period of days or weeks. There is localised tenderness and it can be confused with patellar tendonitis.

ARTHRITIC CONDITIONS

Osteoarthritis

Osteoarthritis is a very common problem of the knee joint. Symptoms usually appear in middle life or later. It is more common in women, the obese, and in those with knee deformities (e.g. genu varum) or previous trauma, especially meniscal tears. The degenerative changes may involve either the lateral or the medial tibiofemoral compartment, the patellofemoral joint or any combination of these sites.

Clinical features

- Slowly increasing joint pain and stiffness
- Aggravated by activities such as prolonged walking, standing or squatting
- Descending stairs is usually more painful than ascending stairs (suggestive of patellofemoral OA)
- Pain may occur after rest, especially prolonged flexion
- Minimal effusion and variable crepitus
- Restricted flexion but usually full extension
- Often quadriceps wasting and tender over medial joint line
- Diagnosis confirmed by X-ray (weightbearing view)

Management options

- Relative rest
- Weight loss
- Analgesics and/or judicious use of NSAIDs

- Glucosamine: a Cochrane review showed that it is both safe and effective (see Chapter 35)
- Walking aids and other supports
- Physiotherapy (e.g. hydrotherapy, quadriceps exercises, mobilisation and stretching techniques)
- Viscosupplementation: intra-articular injection of hylans
- Intra-articular injections of corticosteroids are generally not recommended but a single injection for severe pain can be very effective
- Surgery is indicated for severe pain and stiffness and includes arthroscopic debridement and wash out, osteotomy, arthrodesis and total joint replacement (Figure 67.20) or hemiarthroplasty, especially for the medial compartment with focal arthritis and varus deformity



Figure 67.20 Total joint replacement of knee

Rheumatoid arthritis

The knee is frequently affected by RA although it rarely presents as monoarticular knee pain. RA shows the typical features of inflammation - pain and stiffness that is worse after resting. Morning stiffness is a feature.

Note: The spondyloarthropathies have a similar clinical pattern to RA.

Synovectomy is a useful option with persistent boggy thickening of synovial membrane but without destruction of the articular cartilage.[2]

Baker's cyst

A popliteal cyst (Baker's cyst) is a herniation of a chronic knee effusion between the heads of the gastrocnemius muscle and usually is associated with osteoarthritis, rheumatoid arthritis or internal derangement of the knee. It presents as a mass behind the knee and may or may not be tender or painful.

It tends to fluctuate in size.

A Baker's cyst indicates intra-articular pathology and indicates a full assessment of the knee joint.

Rupture may result in pain and swelling in the calf, mimicking DVT.

Treat underlying knee inflammation (synovitis).

Surgical removal of the cyst is advisable for persistent problems.

Septic arthritis

Tends to be more common in the knee than other joints. Septic (pyogenic) arthritis should be suspected when the patient complains of intense joint pain, malaise and fever. In the presence of acute pyogenic infection the joint is held rigidly. The differential diagnosis includes gout and pseudogout (chondrocalcinosis).

PRINCIPLES OF MANAGEMENT

Most painful knee conditions are not serious and, providing a firm diagnosis is made and internal knee disruption or other serious illness discounted, a simple management plan as outlined leads to steady relief. For more serious injuries the primary goal is to minimise the adverse consequences of forced tinactivity.

- First aid: RICE (avoid heat in first 48 hours).
- Lose weight if overweight.
- Adequate support for ligament sprains - supportive elastic tubular (tubigrip) bandage or a firm elastic bandage over Velband.
- Simple analgesics - paracetamol (acetaminophen).
- Judicious use of NSAIDs and corticosteroid injections.
- Physiotherapy to achieve strength and stability.
- Attend to biomechanical abnormalities, inappropriate footwear and athletic techniques.
- Orthotics and braces to suit the individual patient.
- Specialised exercise techniques (e.g. the McConnell technique).[2]
- Quadriceps exercises: these simple exercises are amazingly effective.

Quadriceps exercises (examples)

- Instruct the patient to tighten the muscles in front of the thighs (as though about to lift the leg at the hip and bend the foot back but keeping the leg straight). The patient should hold the hand over the lower quadriceps to ensure it is felt to tighten. This tightening and relaxing exercise should be performed at least 6 times every 2 hours or so until it becomes a habit. It can be done sitting, standing or lying (Figure 67.21).



Figure 67.21 A quadriceps exercise: with outstretched legs the quadriceps muscle is slowly and deliberately tightened by straightening the knee to position (a) from the relaxed position (b)

- Sitting on a chair the patient places a weight of 2 - 5 kg around the ankle (e.g. a plastic bag with sand or coins in a sock) and lift s the leg to the horizontal and then gently lowers it (avoid in patellofemoral problems).

WHEN TO REFER

- Early referral is required for knees ‘at risk’ following acute injuries where one or more of the following are present:
  - locked knee
  - haemarthrosis
  - instability
- Clinical evidence of a torn cruciate ligament, third degree tear of the collateral ligaments or torn meniscus
- Undiagnosed acute or chronic knee pain
- Recurrent subluxation or dislocation of the patella
- Suspected septic arthritis
- Presence of troublesome intra-articular loose body

Practice tips

- The absence of an effusion does not rule out the presence of severe knee injury.
- Examine the hip and lumbosacral spine if examination of the knee is normal but knee pain is the complaint.
- Always think of an osteoid osteoma in a young boy with severe bone pain in a leg (especially at night) that responds nicely to aspirin or paracetamol or other NSAID.
- Tears of the meniscus can occur, especially in middle age, without a history of significant preceding trauma.
- If a patient presents with a history of an audible ‘pop’ or ‘crack’ in the knee with an immediate effusion (in association with trauma) he or she has an ACL tear until proved otherwise.
- Haemarthrosis following an injury should be regarded as an anterior cruciate tear until proved otherwise.
- The ‘movie’ sign whereby the patient seeks an aisle seat to stretch the knee is usually due to patellofemoral pain syndrome.
- The ‘bed’ sign, when pain is experienced when the knees touch while in bed, is suggestive of a medial meniscal cleavage tear.
- A positive squat test (medial pain on full squatting) indicates a tear of the posterior horn of the medial meniscus.
- Joint aspiration should not be performed on the young athlete with an acute knee injury.
- If an older female patient presents with the sudden onset of severe knee pain think of osteonecrosis.
- Reserve intra-articular corticosteroid injections for inflammatory conditions such as rheumatoid arthritis or a crystal arthropathy: regular injections for osteoarthritis are to be avoided. Do not give the injections when the inflammation is acute and diffuse or in the early stages of injury.
- Many inflammatory conditions around the knee joint, such as bursitis or tendonitis, respond to a local injection of local anaesthetic and corticosteroid but avoid giving injections into the tendon, especially the patellar tendon.
- Keep in mind the technique of autologous cartilage transplantation: in this technique cartilage cells (chondrocytes) are taken from the patient, multiplied in a laboratory and eventually implanted into the damaged area. It can be used for damage in any major joint, especially the knee, being ideal for osteochondritis dissecans.

REFERENCES

1. Knox JDE. Knee problems. In: *Practice*. London: Kluwer - Harrap Handbooks, 1982; 3.66: 1 - 5.
2. Selecki Y, Helman T. How to treat knee pain. Aust Dr. Weekly, 1993; 22 April: i - viii.
3. McLean I. Assessment of the acute knee injury. Aust Fam Physician, 1984; 13: 575 - 80.
4. Cyriax J. *Textbook of Orthopaedic Medicine*, Vol. 1 (6th edn). London: Bailliere Tindall, 1976: 594.
5. Noyes FR. Arthroscopy in acute traumatic haemarthrosis of the knee. J Bone Joint Surg, 1980: 624 - 87.
6. Corrigan B, Maitland GD. *Practical Orthopaedic Medicine*. Sydney: Butterworths, 1986: 126 - 61.
7. Hoppenfeld S. Physical Examination of the Spine and Extremities. Englewood Cliffs: Prentice-Hall, 1976: 172 - 3.
8. Cross MJ, Crichton KJ. *Clinical Examination of the Injured Knee*. London: Harper & Row, 1987: 21 - 46.

9. Lau L ed. *Imaging Guidelines* (4th edn). Melbourne: RAZNC Radiologists, 2001: 200 - 1.
10. Cross MJ. The painful knee. Aust Fam Physician, 1984; 13: 166 - 8.
11. Larkins P. The little athlete. Aust Fam Physician, 1991; 20: 973 - 8.
12. Rostrom PKM, Calver RF. Subcutaneous atrophy following methyl prednisolone injection in Osgood-Schlatter epiphysitis. J Bone Joint Surg, 1979; 61A: 627 - 8.
13. Mital MA, Matza RA, Cohen J. The so-called unresolved Osgood-Schlatter's lesion. J Bone Joint Surg, 1980; 62A: 732 - 9.
14. Wilkins E et al. Osteoarthritis and articular chondrocalcinosis in the elderly. Ann Rheum Dis, 1983; 42(3): 280 - 4.
15. Rush J. Spontaneous osteonecrosis of the knee. Current Orthopaedics, 1999; 13: 309 - 14.
16. Edwards E, Miller R. *Management of acute knee injuries*. Med Observer, 2000; 17 March: 67 - 9.
17. Mashford ML (Chair). *Therapeutic Guidelines: Analgesic* (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2002: 149 - 52.
18. Fricker P. Anterior knee pain. Aust Fam Physician, 1988; 17: 1055 - 6.

CHAPTER 68. PAIN IN THE FOOT AND ANKLE

> The victim goes to bed and sleeps in good health. About two o'clock in the morning he is awakened by a severe pain in the great toe; more rarely in the heel, ankle, or instep … The part affected cannot bear the weight of the bed clothes nor the jar of a person walking in the room. The night is spent in torture.
>
> Thomas Sydenham (1624 - 89)

PAIN IN THE FOOT AND ANKLE

Pain in the foot (podalgia) and ankle problems are a common occurrence in general practice. Various characteristics of the pain can give an indication of its cause, such as the description of gout by Thomas Sydenham. There are many traumatic causes of podalgia and ankle dysfunction, especially fractures and torn ligaments, but this chapter will focus mainly on everyday problems that develop spontaneously or through overuse. The main causes of foot pain are presented in Table 68.1.[1]

Table 68.1 Causes of foot pain (after Johnson)[1]

General
Arthritis - OA, gout, RA, seronegative spondyloarthropathies. Diabetes - neuropathy [sensory (Charcot), motor, autonomic, single nerve], sepsis, vasculopathy. Peripheral neuritis - alcohol, vitamin B12 deficiency. Vascular - arteriosclerosis (claudication, gangrene), hemiplegia, Raynaud's, regional pain syndrome (Sudeck's). Infections - cellulitis, septic arthritis, TB, actinomyces. Other: Paget's disease of bone, osteoid osteoma, hypermobility syndrome (including Marfan).
Ankle and hindfoot
Tendoachilles (bursitis, tendonitis, tear), posterior tibial tendonitis, rupture or subluxation, plantar fasciitis, sprain, bruised heel, phlebitis, cellulitis.
Midtarsal
Acute or chronic foot strain, synovitis of subtaloid, tarsal coalition, hypomobility of transverse tarsal joints, osteochondritis of navicular (Kohler's), dorsal exostosis, peroneus brevis tendonitis, flexor hallucis longus tendonitis.
Forefoot
Bunion, bunionette, Tailor's bunion, intermetatarsal bursitis, traumatic synovitis of MTP joint, sesamoiditis, march fracture, Freiberg's disorder.
Toes
Hallux valgus, hallux rigidus, varus little toe, mallet toe, clawed toe, corn, wet corn, ingrown toenail, onychogryphosis, subungual exostosis, deep peroneal nerve entrapment, digital nerve entrapment (Morton's neuralgia).
Sole
Callus, plantar wart, epidermoid cyst, foreign body, tarsal tunnel syndrome, Dupuytren's (Ledderhose's) contracture.

KEY FACTS AND CHECKPOINTS

- Foot deformities such as flat feet (pes planus) are often painless.
- Foot strain is probably the commonest cause of podalgia.[2]
- A common deformity of the toes is hallux valgus, with or without bunion formation.
- Osteoarthritis is a common sequel to hallux valgus.
- Osteoarthritis affecting the ankle is relatively rare.

- All the distal joints of the foot may be involved in arthritic disorders.
- Many foot and ankle problems are caused by unsuitable footwear and lack of foot care.
- Ankle sprains are the most common injury in sport, representing about 25% of injuries.
- Severe sprains of the lateral ligaments of the ankle due to an inversion force may be associated with various fractures.
- Bunions and hammer toes are generally best treated by surgery.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 68.2.

Table 68.2 The painful foot and ankle: diagnostic strategy model

Q. Probability diagnosis
A. Acute or chronic foot strain
Sprained ankle
Osteoarthritis, esp. great toe
Plantar fasciitis
Achilles tendonitis
Wart, corn or callus
Ingrowing toenail/paronychia
Q. Serious disorders not to be missed
A. Vascular insufficiency
• small vessel disease
Neoplasia
• osteoid osteoma
• osteosarcoma
• synovial sarcoma
Severe infections (rare)
• septic arthritis
• actinomycosis
• osteomyelitis
Rheumatoid arthritis
Peripheral neuropathy
Regional pain syndrome
Ruptured Achilles tendon
Ruptured tibialis posterior tendon
Q. Pitfalls (often missed)
A. Foreign body (especially children)
Gout
Nerve entrapment
• Morton's neuroma
• tarsal tunnel syndrome

• deep peroneal nerve Chilblains Stress fracture (e.g. navicular) Erythema nodosum *Rarities* Spondyloarthropathies Osteochondritis • navicular (Köhler's) • metatarsal head (Freiberg's) • calcaneum (Sever's) Glomus tumour (under nail) Paget's disease
Q. Seven masquerades checklist
A. Depression? Diabetes Drugs Anaemia ? Thyroid disorder - Spinal dysfunction UTI -
Q. Is the patient trying to tell me something?
A. A non-organic cause warrants consideration with any painful condition.

Probability diagnosis

Common causes include osteoarthritis, especially of the first metatarsophalangeal (MTP) joint, acute or chronic foot strain, plantar fasciitis, plantar skin conditions such as warts, corns and calluses and various toenail problems.

Serious disorders not to be missed

The very important serious disorders to consider include:

- vascular disease - affecting small vessels
- diabetic neuropathy
- osteoid osteoma
- rheumatoid arthritis
- regional pain syndrome

Vascular causes

The main problem is ischaemic pain that occurs only in the foot. The commonest cause is atheroma.

Vascular causes include:

- acute arterial obstruction
- chilblains
- atherosclerosis, especially small vessel disease
- functional vasospasm (Raynaud's) - rare

Symptoms:

- claudication (rare in isolation)
- sensory disturbances, especially numbness at rest or on walking
- rest pain - at night, interfering with sleep, precipitated by elevation, relieved by dependency

For treatment refer to p. 718.

Regional pain syndrome I

Also known as reflex sympathetic dystrophy or Sudeck's atrophy, regional pain syndrome is characterised by severe pain, swelling and disability of the feet. It is a neurovascular disorder resulting in hyperaemia and osteoporosis that may be a sequela of trauma (often trivial) and prolonged immobilisation. Regional pain syndrome usually lasts 2 years and recovery to normality usually follows.

The clinical features include sudden onset in middle-aged patients, pain worse at night, stiff joints and skin warm and red. X-rays that show patchy decalcification of bone are diagnostic. Treatment includes reassurance, analgesics, mobility in preference to rest, and physiotherapy.

Osteoid osteoma

Osteoid osteomas are rare but important little ‘brain teasers’ of benign tumours that typically occur in older children and adolescents. Males are affected twice as often as females. Any bone (except those of the skull) can be affected but the tibia and femur are the main sites. Nocturnal pain is a prominent symptom with pain relief by aspirin being a feature.

Diagnosis is dependent on clinical suspicion and then X-ray, which shows a small sclerotic lesion with a radiolucent centre. Treatment is by surgical excision.

Pitfalls

There are many traps in the diagnosis and management of problems presenting with a painful foot. Common problems require consideration - these include gouty arthritis, chilblains, a stress fracture and a foreign body in the foot, especially in children. Nerve entrapment, as outlined in Chapter 66, is uncommon but Morton's neuroma is reasonably common.

Less common disorders include regional pain syndrome which is often misdiagnosed, the spondyloarthropathies (psoriasis, Reiter's syndrome, ankylosing spondylitis and the inflammatory bowel disorders) and osteochondritis of the calcaneus, navicular bone and metatarsal head. If there is an exquisitely tender small purple-red spot beneath a toenail, a glomus tumour (a benign hamartoma) is the diagnosis. It is worth noting that most of these conditions are diagnosed by X-rays.

General pitfalls

- Failing to order X-rays of the foot.
- Failing to order X-rays of the ankle following injury.
- Failing to appreciate the potential for painful problems caused by diabetes - neuropathy and small vessel disease.
- Neglecting the fact that most of the arthritides can manifest in joints in the foot, especially the forefoot.
- Regarding the sprained ankle in adults and children as an innocuous injury: associated injuries include chondral fractures to the dome of the talus, impaction fractures around the medial recess of the ankle, avulsion fractures of the lateral malleolus and base of fifth metatarsal.
- Misdiagnosing a stress fracture of the navicular which, like the scaphoid fracture, causes delayed union and non-union. Cast immobilisation for 8 weeks initially may prevent the need for surgery.
- Misdiagnosing a complete rupture of the Achilles tendon because the patient can plantar flex the foot.

Seven masquerades checklist

The checklist has four conditions that should be considered, especially diabetes and spinal dysfunction. Diabetes may be responsible for a simple type of atherosclerotic pattern, possibly complicated by infection and ulceration. The neuropathy of diabetes can cause a burning pain with paraesthesia. It has a ‘sock’-type pattern as opposed to the dermatome pattern of nerve root pressure arising from the lumbosacral spine. The common S1 pain is experienced on the outer border of the foot, into the fifth toe and on the outer sole and heel of the foot.

Drugs and anaemia could indirectly cause pain through vascular insufficiency. The drugs that could cause vasospasm include betablockers and ergotamine. An alcoholic neuropathy also has to be considered.

Psychogenic considerations

Any painful condition can be closely associated with psychogenic disorders, including depression.

THE CLINICAL APPROACH

History

This is very important, as always, since various characteristics of the pain can give an indication of its cause. Questions should address the quality of the pain, its distribution, mode of onset, periodicity, relation to weight-bearing, and associated features such as swelling or colour change. It is relevant to enquire about pain in other joints such as the hand and spine, including the sacroiliac joints, which might indicate that the foot pain is part of a polyarthritis. A history of diarrhoea, psoriasis, urethritis or iritis may suggest that one of the spondyloarthropathies has to be excluded.

Key questions

The practitioner should address the following questions:

- Does the pain arise from a local condition or is it part of a generalised disease?
- Is there a history of psoriasis, chronic diarrhoea or colitis, urethritis or iritis?
- Is pain also present in other joints, thus indicating the foot pain is part of a polyarthritis, such as rheumatoid arthritis?
- Is the problem related to unsuitable footwear?
- Does the nature of the pain point to the cause?
    - throbbing pain →inflammation
    - burning pain →nerve entrapment, diabetic neuropathy or regional pain syndrome
    - severe episodic pain →gout
    - pain worse at night →ischaemia (small vessel disease), regional pain syndrome, cramps or osteoid osteoma
    - pain worse at night, relieved by aspirin → osteoid osteoma
    - pain worse on standing after sitting and getting out of bed →plantar fasciitis

For ankle injuries it is important to ask about the nature of the injury:

- Did the foot twist in (invert) or twist out (evert)?
- Was the foot pointing down or up at the time of injury?
- Point with one finger to where it hurts (the finger-pointing sign).
- What happened immediately after the injury?
- Were you able to walk straight away?
- What happened when you cooled off?

If there has been a fall onto the foot from a height, consider the possibility of a fracture of the calcaneus or talus or disruption of the syndesmosis between the tibia and fibula.

PHYSICAL EXAMINATION

Inspection

Inspect the feet with the patient standing, sitting, walking (in shoes and barefooted) and lying down (note plantar surfaces). Inspect the footwear (normally, a shoe wears first on the outer posterior margin of the heel).

Note:

- any gait abnormalities, including limping and abnormal toe in or toe out
- deformities, such as hammer toes, bunions - medial (hallux valgus) and lateral (Tailor's bunion) - and claw toes
- swellings, including callosities
- muscle wasting
- skin changes and signs of ischaemia

Palpation

Systematic palpation is very useful as most structures in the foot are accessible to palpation.

Movements (active and passive)

The joints to test are:

- ankle (talar) joint
- hindfoot (subtalar) joint
- midfoot (midtarsal) joint

Movements

- Plantar flexion (normal - 50?) and dorsiflexion (20?) of foot (Figure 68.1)



Figure 68.1 Dorsi-flexion and plantar-flexion of the ankle joint

- Inversion and eversion of hindfoot (mainly subtalar joint) - hold heel and abduct and adduct (Figure 68.2)



Figure 68.2 Testing inversion and eversion of the hindfoot

- Inversion and eversion of forefoot (midtarsal joint) - hold heel in one hand to fix hindfoot, hold forefoot in the other and abduct and adduct (rotation movement) (Figure 68.3)



Figure 68.3 Testing (**a**) eversion and (**b**) inversion of the forefoot

- test other joints individually (e.g. MTP), midtarsal

Special tests

- Achilles tendon, including calf squeeze (Thompson's or Simmond's test) (Figure 136.17 at pp. 1421 - 2)
- Compress MTP joints from above and below
- Compress metatarsals mediolaterally between thumb and forefinger
- Press upwards from sole of foot just proximal to proximal third and fourth MTP joints - Morton's test
- Check circulation - test dorsalis pedis and posterior tibial pulses
- Neurological examination, including tests for L4, L5 and S1 nerve root function

Investigations

The choice of investigations depends on the clinical features elicited by the history and examination. Select from the following list:

- for systemic diseases:
    - blood glucose
    - RA tests
    - ESR/C-reactive protein
    - HLA-B27
- serum uric acid
- radiology
    - X-ray ± stress and weight-bearing views
    - radionuclide scans (for bone or joint pathology)
    - CT or MRI (especially helpful) scans
    - ultrasound (operator dependent)
- nerve conduction studies

Note: High-resolution ultrasound is used to diagnose disorders of the Achilles tendon and to locate foreign bodies such as splinters of wood and glass.

Radionuclide scanning may detect avascular necrosis in bones, stress fractures, osteoid osteomas, inflammatory osteoarthritis and similar lesions.[3]

FOOT AND ANKLE PAIN IN CHILDREN

Apart from the common problem of trauma, special problems in children include:

- foreign bodies in the foot
- tumours (e.g. osteoid osteoma, osteosarcoma, Ewing's tumour)
- plantar warts
- osteomyelitis/septic arthritis
- ingrowing toenails
- osteochondritis/aseptic necrosis
- osteochondritis dissecans of talus (in adolescents)
- pitted keratolysis and juvenile plantar dermatosis (adolescents)
- stress fractures

Think of osteoid osteoma in children with night pain.

Osteochondritis/aseptic necrosis

Three important bones to keep in mind are:

- the calcaneum - Sever's disorder
- the navicular - Köhler's disorder
- the head of the second metatarsal - Freiberg's disorder

Sever's disorder is traction osteochondritis while the other disorders are a ‘crushing’ osteochondritis with avascular necrosis.

Sever's disorder of the heel

This is calcaneal apophysitis which presents in a child (usually a boy) aged 7 - 15 years (average of 10 years) with a painful tender heel at the insertion of the tendoachilles. It is diagnosed by X-ray. The only treatment is to ensure that the child avoids wearing flat-heeled shoes and wears a slightly raised heel. Strenuous sporting activities should be restricted for 12 weeks and then reviewed.

Köhler's disorder of the navicular

This disorder causes a painful limp (usually mild) with some swelling and tenderness around the navicular in a child (usually a boy) aged 3 - 6 years, although it is seen sometimes in older children. Complete recovery occurs with temporary resting. Sometimes a supportive strapping is helpful.

Freiberg's disorder

This problem affects the head of the second metatarsal (rarely the third) which feels tender and swollen on palpation. It is more common in girls aged 12 - 16 years and can present in young adults. Plain X-ray shows the characteristic collapse of the metatarsal head. The treatment is restriction of activity, protective footwear and protective padding.

Sprained ankle in a child

Children rarely sprain ligaments so it is important to assess apparent strains carefully, including an X-ray.

The little athlete

The ‘little athlete’ can suffer a variety of injuries from accidents and overuse. Diffuse heel pain, which is common, is most often related to Sever's apophysitis of the calcaneum. Occasionally, a juvenile-type plantar fasciitis may occur. Little athletes can develop tendonitis around the ankle, either on the lateral side (peroneals) or medially (tibialis posterior). Occasionally, a stress fracture of the metatarsals or other bones can occur.[4] Special attention must be paid to any developmental structural abnormalities and to footwear.

FOOT AND ANKLE PROBLEMS IN THE ELDERLY

Foot problems are more prevalent in old age. Some are due to a generalised disease, such as diabetes or peripheral vascular disease, while others, such as bunions, hammer toes, calluses and corns, atrophy of the heel fat-pad and Morton's neuroma, increase with ageing. The transverse arch may flatten out and the protective pads under the metatarsals may atrophy, resulting in painful callosities.

Unfortunately, many elderly people regard foot problems as a normal process but these problems actually require considerable care and attention, especially in the presence of peripheral vascular disease, diabetes or rheumatoid arthritis. Deformed toenails (onychogryphosis) is also common albeit not a painful condition.

Flat foot occurring in middle age is usually due to stretching or rupture of the tibialis posterior tendon.[5]

SPRAINED ANKLE

There are two main ankle ligaments that are subject to heavy inversion or eversion stresses, namely the lateral ligaments and the medial ligaments respectively. Most of the ankle ‘sprains’ or tears involve the lateral ligaments (up to 90%) while the stronger, tauter medial (deltoid) ligament is less prone to injury. It is important not to misdiagnose a complete rupture of the lateral ligaments.

Most sprains occur when the ankle is plantar flexed and inverted, such as when landing awkwardly after jumping or stepping on uneven ground. It is a very common sporting injury and is presented in more detail in Chapter 136. Note the Ottawa rules for taking an X-ray of the ankle in Chapter 67 (p. 734).

Clinical features of sprained lateral ligaments

Common features:

- ankle ‘gives way’
- difficulty in weight-bearing
- discomfort varies from mild to severe
- bruising (may take 12 - 24 hours) indicates more severe injury
- may have functional instability: ankle gives way on uneven ground

Physical examination (perform as soon as possible):

- note swelling and bruising
- palpate over bony landmarks and three lateral ligaments (Figure 136.13 at p. 1417)
- test general joint laxity and range of motion
- a common finding is a rounded swelling in front of the lateral malleolus (the ‘signe de la coquille d'oeuf ’)
- test stability in AP plane (anterior draw sign)

Is there an underlying fracture?

For a severe injury the possibility of a fracture - usually of the lateral malleolus or base of fifth metatarsal - must be considered. If the patient is able to walk without much discomfort straight after the injury a fracture is unlikely. However, as a rule, ankle injuries should be X-rayed. See the Ottawa rules (p. 734).

HEEL PAIN

Important causes of heel pain in adults (Figure 68.3)[6] include:

- Achilles tendon disorders
    - tendonitis/peritendonitis (Chapter 136)

    - bursitis
        - postcalcaneal
        - retrocalcaneal
    - tendon tearing (Chapter 136)
        - partial
        - complete
- bruised heel
- tender heel pad
- usually atrophy
- also inflammation
- neuropathies (e.g. diabetic, alcoholic)
- tenosynovitis (FHL, FDL)
- ‘pump bumps’
- plantar fasciitis
- periostitis
- calcaneal apophysitis
- peroneal tendon dislocation
- nerve entrapments
    - tarsal tunnel
    - medial calcaneal nerve
    - nerve to abductor digiti minimi

Ultrasound examination is useful to differentiate the causes of Achilles tendon disorders.

Achilles tendonitis/peritendonitis

The inflammation is a combination of degenerative and inflammatory changes due to overuse and may occur either in the tendon itself or in the surrounding paratendon. It presents as a painful local swelling of the tendon. The latter is called peritendinitis rather than tenosynovitis because there is no synovial sheath.

Management

- Relative rest
- Course of NSAIDs
- Heel padding
- Consider heel ‘raisers’
- Physiotherapy

Achilles tendon bursitis

Bursitis can occur at two sites:

- posterior and superficial - between skin and tendon
- deep (retrocalcaneal) - between calcaneus and tendon (Figure 68.4)


peroneal tendon dislocation or tendonitis
arthritis subtalar joint
plantar fasciitis
disease of calcaneus
peritendonitis
tendonitis
ruptured tendoachilles
partial tear
retrocalcaneal bursitis
postcalcaneal bursitis
calcaneal apophysitis (Sever's disorder)
tender heel pad

Figure 68.4 Important causes of the painful heel

The former occurs mainly in young women from shoe friction and is readily palpated. Tenderness from the deep bursitis is elicited by squeezing in front of the tendon with the thumb and index finger: a swelling may be seen bulging on either side of the tendon.

Treatment

- Avoid shoe pressure (e.g. wear sandals)
- 1 - 2 cm heel raise inside the shoe
- Apply local heat and ultrasound
- NSAIDs
- Inject corticosteroid into bursa with a 25-gauge needle

Fat-pad disorders

A tender heel pad or cushion causes a dull throbbing pain under the heel. It is localised more proximal to that of plantar fasciitis. Once established it is very difficult to treat.

The fat-pad, which consists of globules of fat encapsulated in multiple U-shaped scepti, acts as a hydraulic shock absorber on heel strike. It also contains significant nerve endings.7 It can undergo atrophy, especially in the elderly, and also become inflamed.

Treatment

- Reduction of aggravating activity
- Weight loss (if applicable)
- Simple analgesics
- Orthotic (cushioning heel cup) + or - foam insert
- Good footwear

Problems are treated with an orthotic or an insert and good footwear. Corticosteroids should be avoided as they can accelerate the atrophy.[6]

Plantar fasciitis

This common condition (also known as ‘policeman's heel’) is characterised by pain on the plantar aspect of the heel, especially on the medial side; it usually occurs about 5 cm from the posterior end of the heel although it can be experienced over a wide area beneath the heel. The pain radiates into the sole.

History

- Pain:
    - under the heel

  - first steps out of bed
  - relieved after walking about
  - increasing towards the end of the day
  - worse after sitting
- May be bilateral - usually worse on one side
- Typically over 40 years
- Both sexes
- Sometimes history of injury or overuse
- No constant relationship to footwear

Signs[8]

- Tenderness:
  - localised to medial tuberosity
  - may be more posterior
  - may be lateral
  - may be widespread
  - not altered by tensing fascia (but this action may cause pain)
- Heel pad may bulge or appear atrophic
- Crepitus may be felt
- No abnormality of gait, heel strike, or foot alignment
- Patient often obese

Treatment

Plantar fasciitis tends to heal spontaneously in 12 - 24 months. It has a variable response to treatment with NSAIDs, injections, ultrasound and insoles. Rest from long walks and from running is important. Systematic reviews to date indicate unknown effectiveness from a whole variety of treatments, suggesting a conservative approach based on maximising comfort during the 3 month period of considerable discomfort.[9]

Protection

Symptomatic relief is obtained by protecting the heel with an orthotic pad to include the heel and arch of the foot (e.g. Rose insole). Otherwise, a pad made from sponge or sorbo rubber that raises the heel about 1 cm is suitable. A hole corresponding to the tender area should be cut out of the pad to avoid direct contact with the sole. The aim is to get all of the foot to take the stress.

Injection technique

Disabling plantar fasciitis can be treated by injecting local anaesthetic and long-acting corticosteroid into the site of maximal tenderness in the heel. An alternative is to inject the corticosteroid into the anaesthetised heel.

Method

1. Perform a tibial nerve block. (The area of maximal tenderness should be marked prior to nerve block.)
2. When anaesthesia of the heel is present (about 10 minutes after the tibial nerve block), insert a 23-gauge needle with 1 mL of long-acting corticosteroid (e.g. methylprednisolone acetate) perpendicular to the sole of the foot at the premarked site (Figure 68.5). Insert the needle until a ‘give’ is felt as the plantar fascia is pierced.



Figure 68.5 Injection approach for plantar fasciitis

3. Inject half the steroid against the periosteum in the space between the fascia and calcaneus.
4. Reposition the needle to infiltrate into the fascial attachments over a wider area.

'Cracked' heels

- Soak feet for 30 minutes in warm water containing an oil such as Alpha-Keri or Derma Oil.
- Pat dry, then apply a cream such as Nutraplus (10% urea) or Eulactol heel balm. Hydrocortisone 0.5% cream for resistant cases.
- For severe cases use sorbolene cream with 20% glycerol and 30% urea (test skin sensitivity first).

ARTHRITIC CONDITIONS

Arthritis of the foot or ankle is a rather meaningless diagnosis and specificity is required. Typical sites of arthritic targets are shown in Figure 68.6.



Figure 68.6 Typical sites of arthritic causes of podalgia on skeleton of right foot (plantar aspect)

Osteoarthritis

Osteoarthritis may occur in any of the joints of the foot but it commonly involves the first MTP joint, leading to hallux rigidus. It can affect the subtalar joint, but the ankle joint proper is usually not affected by osteoarthritis.

Hallux rigidus

Osteoarthritis of the first MTP joint can lead to gradual loss of motion of the toe and considerable discomfort. Roomy protective footwear and relative rest is the basis of treatment, coupled with daily selfmobilisation (stretching toe into plantar flexion morning and night). Other measures include manipulation under general anaesthesia or surgery (arthrodesis or arthroplasty) for severe cases.

Rheumatoid arthritis

Rheumatoid arthritis is typically a symmetrical polyarthritis presenting with pain in the MTP joints. It may also affect the ankle, mid-tarsal and tarsometatarsal joints. The interphalangeal joints are seldom affected primarily. It causes pain and stiffness under the balls of the feet, especially first thing in the morning.

Gout

Gout typically affects the first MTP and should be considered with the sudden onset of pain, especially in the presence of redness, swelling and tenderness. It can affect any synovial joint and occasionally may be polyarticular. Gout is often dismissed by the patient as a ‘sprain’. A history of alcohol consumption or diuretic treatment is relevant.

Spondyloarthropathies

This group of arthritic disorders (Reiter's syndrome, ankylosing spondylitis, psoriatic arthritis and arthritides associated with chronic bowel disorders) may involve peripheral joints. Other foot involvement includes plantar fasciitis, Achilles tendonitis and sausage-shaped toes due to tenosynovitis, and arthritis of the proximal interphalangeal joints.

BURNING FEET

It is not uncommon for people, especially the elderly, to present with the complaint of burning feet. A careful history is needed to elicit exactly what they mean by 'burning' - is it real pain, a cold sensation or paraesthesia?

A checklist of causes is as follow:

- vascular
    - ischaemic rest pain from small vessel disease
    - chilblains or other cold reaction
    - functional vasospasm (Raynaud's)
- diabetic neuropathy
- tarsal tunnel syndrome (refer to pp. 715 - 6)
- regional pain syndrome
- Morton's neuroma (localised pain between toes)
- psychogenic, especially anxiety

It is worth considering tarsal tunnel syndrome if there is anterior burning pain in the forefoot with associated aching in the calf. It is usually present in menopausal women and worse at night. Caused by entrapment of the posterior tibial nerve near the medial malleolus, it many be associated with rheumatoid arthritis. Treat with physiotherapy, a medial arch support and a corticosteroid injection before contemplating surgery.

FOOT STRAIN

Foot strain is probably the commonest cause of podalgia. A foot may be strained by abnormal stress, or by normal stress for which it is not prepared. In foot strain the supporting ligaments become stretched, irritated and inflamed. It is commonly encountered in athletes who are relatively unfit or have a disorder such as flat feet, or in obese adults.

Symptoms and signs

- Aching pain in foot and calf during or after prolonged walking or standing
- Initial deep tenderness felt on medial border of plantar fascia (Figure 68.7)


Freiberg's disorder
tendonitis of FHL tendon
Morton's neuroma
bunion
stress (march) fracture
fractured tubercle 5th metatarsal
foot strain
Köhler's disorder
plantar fasciitis
FHL: flexor hallucis longus

Figure 68.7 Typical sites of important causes of podalgia (other than arthritis) - right foot

- Worse with new shoes, especially a change to high heels

Acute foot strain

Acute ligamentous strain, such as occurs to the occasional athlete or to the person taking long unaccustomed walks, is usually self-limiting. It recovers rapidly with rest.

Chronic foot strain

Foot strain will become chronic with repeated excessive stress or with repeated normal stress on a mechanical abnormality. A common consequence is an everted foot, leading to flattening of the longitudinal arch on weight-bearing. It is important to establish whether the symptoms commenced after the patient began wearing a different type of footwear.

Treatment

The treatment is basically the same as that of the adult flat foot. Acute strain is treated with rest and by reducing walking to a minimum. Try the application of cold initially and then heat. The management of chronic strain is based on an exercise program and orthotics, including arch supports, to correct any deformity.

ACHING FEET

- Avoid wearing high heels.
- Wear insoles to support the foot arch.
- Perform foot exercises.
- Soak the feet in a basin of warm water containing therapeutic salts (Epsom salts is suitable).
- Massage feet with baby oil followed by a special ribbed wooden foot massager.

FLAT FEET (PES PLANUS)

Flat feet are normal in young children. No treatment is required in flat feet in which the arch is restored by standing on tiptoe (see p. 897). If painful, treat with exercises and insoles. Refer if concerned. Hind foot fusion can be performed for severe pain.

CLAW FOOT (PES CAVUS)

High foot arch is usually of congenital origin. It may be secondary to polio or various neurological conditions. The foot is inflexible and the toes may be ‘hammer’ or clawed. Treatment includes special orthotics with good shock-absorbing properties, appropriate footwear, foot exercises and padding under the metatarsal heads. Operative treatment involves soft tissue release or arthrodesis to strengthen toes.

DISORDERS OF THE ANKLE TENDONS

Inflammation of a tendon sheath about the ankle may result from repetitive overuse, trauma such as a sprained ankle or unaccustomed stress, including sporting injuries.

Tenosynovitis commonly involves the tibialis posterior tendon over the medial compartment or the peroneal tendons over the lateral compartment. It may also affect the tibialis anterior and extensor digitorium longus tendons. Friction at the point where the tendons become angulated at the ankle causes the inflammation. The patient presents with pain, swelling and restricted movement. On examination there is swelling and tenderness where the tendon bulges out from behind and below the malleolus.

If necessary the diagnosis can be confirmed by ultrasound or MRI imaging. Complications include tenovaginitis, weakness, ganglion formation, subluxation or dislocation and rupture.

Treatment of tendonitis includes partial immobilisation (rarely in a cast) or an orthotic device to support the arch of the foot.

A carefully directed injection of corticosteroid into the tendon sheath can be very effective.

Peroneal tendonitis

This occurs along the course of the tendon from behind the lateral malleolus to the outer side of the foot and is common in athletes and ballet dancers. Pain is reproduced on palpation, on stretching the tendons by passive inversion of the foot or by resisting eversion of the foot.

Peroneal tendon dislocation

It is most commonly a dislocated leg tendon as a result of forceful dorsiflexion. An audible painless snapping sensation may be experienced. Surgical repair is necessary.

Tibialis posterior tendonitis

This is a common problem, especially in middleaged females, in ballet dancers and in those with pes planus with a valgus deformity. Pain is reproduced on palpation usually anterior and inferior to the medial malleolus, on stretching by passive eversion of the foot or resisted active inversion of the foot. Complications include a tarsal tunnel syndrome. First-line treatment is by an injection of corticosteroid with local anaesthetic into the tendon sheath at its tender area. (See Figure 68.8.)



Figure 68.8 Medial aspect of the foot illustrating tendon disorders: tendonitis of tibialis anterior and tibialis posterior

Tibialis posterior tendon dislocation

This can occur with forceful ankle dorsiflexion and inversion. The patient usually experiences pain and cannot weight-bear. The dislocated tendon may be seen overlying the medial malleolus. Immediate surgical repair is recommended.

Tibialis posterior tendon rupture

Rupture of the tibialis posterior tendon after inflammation, degeneration or trauma10 is a relatively common and misdiagnosed disorder, especially in middle-aged females. It causes collapse of the longitudinal arch of the foot, leading to a flat foot.[5]

It is uncommon for patients to feel obvious discomfort at the moment of rupture. They may subsequently present with the sudden appearance of an ‘abnormal’ flat foot. There is gross eversion of the foot.

A simple test is the ‘too many toes’ test whereby more toes are seen on the affected side when the feet are viewed from about 3 metres behind the patient (Figure 68.9).[5]



Figure 68.9 Tibialis posterior tendon rupture (right foot): the 'too many toes' posterior view

The single heel raise test is also diagnostic. The most useful investigation is an ultrasound examination. Minor cases can be treated conservatively with orthotics, but severe problems respond well to surgical correction.

SESAMOIDITIS[11]

The two sesamoids that lie beneath the head of the first metatarsal may develop painful conditions such as chondromalacia, osteoarthritis and stress fractures. A special 'sesamoid' X-ray assists diagnosis. Painful callus can develop over here in the elderly. Welldesigned insoles are usually effective as is surgical excision for persistent problems.

METATARSALGIA

Metatarsalgia is not a disease but refers to pain and tenderness over the plantar heads of the metatarsals (the forefoot). Causes include foot deformities (especially with depression of the transverse arch), arthritis of the MTP joints, trauma, Morton's neuroma, Freiberg's disorder and entrapment neuropathy. However it can occur in normal feet after prolonged standing.

Depression of the transverse arch results in abnormal pressure on the second, third and fourth metatarsal heads with possible callus formation. Repetitive foot strain, pes cavus and high heels may cause a maldistribution of weight to the forefoot.

Treatment involves treating any known cause, advising proper footwear and perhaps a metatarsal bar. Flat-heeled shoes with ample width seldom cause problems in the metatarsal region.

STRESS FRACTURES

Clinical features:

- the aches or pains may be slow in onset or sudden

- common in dancers, especially classical ballet, and in unfit people taking up exercise
- examination is often unhelpful: swelling uncommon[12]
- routine X-rays often unhelpful
- a bone scan is the only way to confirm the suspected diagnosis
- basis of treatment is absolute rest for 6 or more weeks with strong supportive footwear
- a walking plaster is not recommended

Avulsion fracture of base of fifth metatarsal

Known also as a Jones fracture, it is usually a traumatic fracture but can be a stress fracture and associated with severe ankle sprains.

March fracture of metatarsal

Stress or fatigue fracture of the forefoot usually involves the neck of the second metatarsal (sometimes the third).

Tarsals, especially navicular

Stress fracture of the navicular, which is a disorder of athletes involved with running sports, presents as poorly localised midfoot pain during weight-bearing. Examination and plain X-ray are usually normal. It is a recently recognised serious disorder due to the advent of nuclear bone scans and CT scans. A protracted course of treatment can be expected.

Calcaneum

Stress fractures of the os calcis usually have an insidious onset. Osteoporosis is a predisposing factor, as is an increased training program.[6]

MORTON'S NEUROMA

Morton's interdigital neuroma is probably misdiagnosed more often than any other painful condition of the forefoot. It is not a true neuroma but a fibrous enlargement of an interdigital nerve, and its aetiology is still uncertain. The diagnosis is made on clinical grounds. An ultrasound examination may detect a neuroma.

Clinical features

- Usually presents in adults < 50 years
- Four times more common in women
- Bilateral in 15% of cases
- Commonest between third and fourth metatarsal heads (Figure 68.10), and second and third (otherwise uncommon)



Figure 68.10 Morton's neuroma: typical site and pain and paraesthenia distribution

- Severe burning pain (sometimes sharp and shooting) between third and fourth or second and third toes
- Worse on weight-bearing on hard surfaces (standing and walking)
- Aggravated by wearing tight shoes
- Relieved by taking offshoe and squeezing the forefoot
- Localised tenderness between metatarsal heads

Treatment

Early problems are treated conservatively by wearing loose shoes with a low heel and using a sponge rubber metatarsal pad. An orthosis with a dome under the affected interspace helps to spread the metatarsals and thus takes pressure off the nerve. Most e ventually require surgical excision, preferably with a dorsal approach. A corticosteroid injection can be considered.

HALLUX VALGUS

Hallux valgus with associated bunion formation and splaying of the forefoot is common. It may be a consequence of poor-fitting footwear.

A bunionette, also caused by pressure, may form over the fifth metatarsal. Pain, if present, may be due to shoe pressure on an inflamed bunion, a hammer toe, metatarsalgia or secondary arthritis of the first MTP joint.

Hallux valgus with bunions should be treated by correcting footwear prior to any surgical correction. Systematic evidence-based reviews found that preventive orthoses and night splints were unlikely to be beneficial but absorbable pin fixation was likely to be beneficial.[9]

HAMMER TOES

Mainly involve the second toe with extended MTP joint, hyperflexed PIP joint and extended DIP joint. Painful corns will appear over the prominent joint. They respond well to surgery if problematic and are not helped by good footwear.

CLAW TOES

Often follows pdio. The feature is extended MTP joint, flexion PIP and DIP. Refer for surgical opinion.

CALLUS, CORN AND WART

The diagnosis of localised, tender lumps on the sole of the foot can be difficult. The differential diagnosis of callus, corn and wart is aided by an understanding of their morphology and the effect of paring these lumps (Table 68.3).

Table 68.3 Comparison of the main causes of a lump on the sole of the foot

	Typical site	Nature	Effect of paring
Callus	Where skin is normally thick: beneath heads of metatarsals, heels, inframedial side of great toe	Hard, thickened skin	 **Figure 68.13a** normal skin
Corn	Where skin is normally thin: on soles, fifth toe, dorsal projections of hammer toes	White, conical mass of keratin, flattened by pressure	
Wart	Anywhere, mainly over metatarsal heads, base of toes and heels; has bleeding points	Viral infections, with abrupt change from skin at edge	 **Figure 68.13c** exposes bleeding points

A callus (Figure 68.11) is simply a localised area of hyperkeratosis related to some form of pressure and friction. It is very common under the metatarsal heads, especially the second.



Figure 68.11 Callus

A corn (Figure 68.12) is a small, localised, conical thickening. It is a horny plug of keratin in the epidermis. A corn develops in response to chronic irritation, usually over a bony prominence of the foot, e.g. outer distal fifth toe. It may resemble a plantar wart but gives a different appearance on paring.



Figure 68.12 Corn

A wart (Figure 68.13) is more invasive, and paring reveals multiple small, pinpoint bleeding spots.



Figure 68.13 Plantar wart

Treatment

Calluses

No treatment is required if asymptomatic. Remove the cause. Proper footwear is essential - wide enough shoes and cushioned pads over ball of foot. Proper paring gives relief, also filing with callus files. If severe, apply daily applications of 10% salicylic acid in soft paraffin with regular paring.

Corns

Remove cause of friction and use wide shoes to allow the foot to expand to its full width. Soften corn with a few daily applications of 15% salicylic acid in collodion or commercial ‘corn removers’ with salicylic acid and then pare (Figure 68.13b). For soft corns between the toes (usually last toe-web) keep the toewebs separated with lamb's wool or a cigarette filter tip at all times and dust with a foot powder.



Figure 68.13b exposes white, avascular corn with concave surface

Plantar warts (Figure 68.14)



Figure 68.14 Plantar warts on the sole of the foot and toes showing a mosaic pattern

There are many treatments for this common and at times frustrating problem. A good rule is to avoid scalpel excision, diathermy and electrocautery because of the problem of scarring. One of the problems with the removal of plantar warts is the 'iceberg' configuration - not all the wart may be removed.

Methods of removal

- Salicylic acid 17%, lactic acid 17% in collodion
  - apply daily, dry and cover
- Liquid nitrogen
  - pare wart (a 21-gauge blade is recommended)
  - apply liquid nitrogen
  - repeat weekly

Can be painful and the results are often disappointing.

- Topical chemotherapy
  - pare wart (particularly in children)
  - apply Upton's paste to wart each night and cover
  - review if necessary

(Upton's paste comprises trichloracetic acid 1 part, salicylic acid 6 parts, glycerin 2 parts.)

- Topical chemotherapy and liquid nitrogen
  - pare wart
  - apply paste of 70% salicylic acid in raw linseed oil
  - occlude for 1 week
  - pare on review, then apply liquid nitrogen and review
- Curettage under local anaesthetic
  - pare the wart vigorously to reveal its extent
  - thoroughly curette the entire wart with a dermal curette
  - hold the foot dependent over kidney dish until bleeding stops (this always stops spontaneously and avoids a bleed later on the way home)
  - apply 50% trichloracetic acid to the base

Occlusion method

- Occlusion with topical chemotherapy: a method of using salicylic acid in a paste under a special occlusive dressing is described:

Equipment

2.5 cm (width) elastic adhesive tape 30% salicylic acid in Lassar's paste of plasticine consistency

Method

1. Cut two lengths of adhesive tape, one about 5 cm and the other shorter.
2. Fold the shorter length in half, sticky side out (Figure 68.15a).



Figure 68.15 Treatment of plantar wart: **(a)**‘window’ to fit the wart is cut out of shoulder of elastic adhesive tape; **(b)** larger strip covers the wart and shoulder strip

3. Cut a half-circle at the folded edge to accommodate the wart.
4. Press this tape down so that the hole is over the wart.
5. Roll a small ball of the paste in the palm of the hand and then press it into the wart.
6. Cover the tape, paste and wart with the longer strip of tape (Figure 68.15b).

This paste should be reapplied twice weekly for 2 - 3 weeks. The reapplication is achieved by peeling back the longer strip to expose the wart, adding a fresh ball of paste to the wart and then recovering with the upper tape.

The plantar wart invariably crumbles, and vanishes. If the wart is particularly stubborn, 50% salicylic acid can be used.

Complementary therapy

- Apply the inner surface of banana skin daily and cover with tape. Cut a small disc of banana skin to cover the wart. Apply the inner surface to the wart on a daily or second daily basis, cover with tape and keep dry. Continue for a few weeks or as necessary.
- Poultice of aspirin and tea tree oil:
  - - Place a non-effervescent 125 - 300 mg soluble aspirin tablet on the centre of the wart and dampen

it with 15% tea tree (*Melaleuca*) oil in alcohol.

- Cover with a cotton pad and tape firmly with Micropore (or similar tape). Allow it to get wet.
- After one week remove the dressing and debride or curette the friable slough.
- Repeat if necessary.

INGROWN TOENAIL (ONYCHOCRYPTOSIS)

Ingrown toenail is a very common condition, especially in adolescent boys. Although not so common in adults, it may follow injury or deformity of the nail bed. It is typically located along the lateral edges of the great toenail and represents an imbalance between the soft tissues of the nail fold and the growing nail edge. The basic cause is a redundant skin fold. It is exacerbated by faulty nail trimming, constricting shoes and poor hygiene. A skin breach is followed by infection, then oedema and granulation tissue of the nail fold.[5]

Treatment

Prevention

All patients should be instructed on correct foot and nail care. Foot hygiene includes foot baths, avoiding nylon socks, and frequent changes of cotton or wool socks. Cotton wool pledgets can be placed beneath the nail edge to assist separation.

It is important to fashion the toenails so that the corners project beyond the skin (Figure 68.16). The end of the nail (not the corners) should be cut squarely so that the nail can grow out from the nail fold. Then each day, after a shower or bath, use the pads of both thumbs to pull the nail folds as indicated.


cut nail towards centre
corners of nail project beyond skin
stretch nail folds with thumbs daily

Figure 68.16 Method of fashioning toenails

Surgical methods

1. *Excision of ellipse of skin*. This ‘army method’ transposes the skin fold away from the nail. The skin heals, the nail grows normally and the toe retains its normal anatomy.
   Under digital block, an elliptical excision is made such that the skin fold is forced off the nail with a blunt instrument and held there by the wound closure (Figure 68.17). Any granulation tissue and debris should be removed with a curette.



Figure 68.17 Treatment of ingrown toenail: excision of ellipse of skin

2. *Electrocautery*. This is similar in principle to the preceding method but is simple, quick and very effective with minimal after-pain, especially for severe ingrowing with much granulation tissue. Under digital block the electrocautery needle removes a large wedge of skin and granulation tissue so that the ingrown nail stands free of skin (Figure 68.18).



Figure 68.18 Treatment of ingrown toenail: electrocautery of wedge of tissue

3. *Skin wedge excision*. Another similar method under digital block is to dissect away all the skin fold adjacent to the nail, starting from the nail base, extending proximally for about 4 mm and then sweeping around the side of the nail to under its tip, using a 3 - 4 mm margin all the way. Removal includes granulation and subcutaneous tissue. Bleeding points are cauterised and the raw area dressed. Dressings are necessary for the next 4 - 6 weeks.
4. *Wedge of nail excision and phenolisation*. This method uses 80% phenol (concentrated solution) to treat the nail bed following excision with scissors of a wedge for about one quarter of the length (rather than a standard wedge resection) of the ingrown nail. A cotton wool stick soaked in phenol is introduced deep into the space of the nail bed (Figure 68.19). Leave the stick in this site for 3 minutes (by the clock). Then remove and flush this pocket with isotonic saline or alcohol, then dry with a cotton wool stick. Dress with paraffin gauze, then with dry gauze. Re-dress as appropriate. The success rate is almost 100%.



Figure 68.19 Phenolisation method

Warning: Take care not to spill the phenol onto the surrounding skin as it is very corrosive.

Practice tips

- Good-quality X-rays are mandatory in all severely sprained ankle injuries.
- If in doubt about the diagnosis of a painful foot - X-ray.
- Children rarely sprain ligaments. All joint injuries causing pain and swelling in children need to be X-rayed.
- Think of the rare problem of a dislocating peroneal tendon if a sharp click and stab of pain is experienced just behind and below the lateral malleolus.
- Paraesthesia of part or whole of the foot may be caused by peripheral neuropathy, tarsal tunnel syndrome, mononeuritis (e.g. diabetes mellitus), rheumatoid arthritis or a nerve root lesion from the lumbosacral spine.
- Avoid giving injections of corticosteroids into the Achilles tendon.
- Avoid invasive procedures such as surgical excision, diathermy or electrocautery for plantar warts. Be aware of the limitations of liquid nitrogen.
- High-resolution ultrasound can help diagnose Achilles tendon disorders.
- Keep in mind the possibility of pain around the sesamoid bones of the first metatarsal.
- Beware of acral lentiginous melanoma on the sole of the foot, especially if amelanotic.

PARONYCHIA

Initial treatment:

- antiseptic (e.g. Betadine-soaked) dressing
- elevation of nail fold to drain pus
- application of petroleum gauze dressing
- antibiotics if extensive or cellulitis developing

Sometimes the nail requires avulsion to establish free drainage of a periungual abscess. Refer to Chapter

119.

REFERENCES

1. Johnson FL. The painful foot: an overview of podalgia. Aust Fam Physician, 1987; 16: 1086.
2. Cailliet R. *Foot and Ankle Pain*. Philadelphia: FA Davis, 1983: 105 - 15.
3. de Jager JP. Problems with the shoulder, knee, ankle and foot. Med J Aust, 1996; 165: 570 - 1.
4. Larkins PA. The little athlete. Aust Fam Physician, 1991; 20: 973 - 8.
5. Quirk R. Flat foot in middle age: diagnosis and treatment. Mod Med Aust, 1995; 38(11): 44 - 7.
6. Brown CH. A review of subcalcaneal heel pain and plantar fasciitis. Aust Fam Physician, 1996; 25: 875 - 85.
7. Jahss MH et al. Investigations into the fat-pads of the sole of the foot: anatomy and histology. Foot and Ankle, 1992; 13: 233 - 42.
8. Harbison S. Plantar fasciitis. Aust Fam Physician, 1987; 16: 113.
9. Barton S ed. *Clinical Evidence*. London: BMJ Publishing Group, 2001: 742 - 3, 823 - 31.
10. Masterton E et al. The planovalgus rheumatoid foot - is tibialis tendon rupture a factor? Br J Rheumatol, 1995; 34: 645 - 6.
11. Quirk R. Metatarsalgia. Aust Fam Physician, 1996; 25: 863 - 9.
12. Quirk R. Stress fractures of the foot. Aust Fam Physician, 1987; 16: 1101 - 2.

CHAPTER 69. WALKING DIFFICULTY

Would ye not think his cunning to be great that could restore this cripple to his legs again?

William Shakespeare (1564 - 1616), *King Henry VI, Part II*, Act 2, Scene 1

WALKING DIFFICULTY

The clinical evaluation of the patient presenting with difficulty walking can be very complex, especially for abnormal gaits caused by neurological conditions. Not all gaits fall into a single category; gait disturbances may be multifactorial, especially in the elderly.

Non-neurological conditions are the most common cause of walking difficulties. They include various arthritic conditions of the lower limbs, usually presenting as a limp, other mechanical factors, such as swelling of the legs, disorders of circulation such as intermittent claudication, and general debility (e.g. malignancy, anaemia and endocrine disorders such as hyperparathyroidism).

It is important for the general practitioner not to overlook hypokalaemia and drugs or the myopathies as a cause of walking difficulty. The drugs that require special consideration include alcohol, corticosteroids, chloroquine, colchicine, clofibrate, bretylium, HMGCoA reductase inhibitors (the statins), gemfibrozil, penicillamine, diuretics, beta-blockers and general anaesthetic agents.

ABNORMAL GAITS

It is convenient to classify abnormal gaits as painless or painful (antalgic). With antalgic gaits the rhythm is disturbed; with painless abnormal gaits the contour is affected. One type of skeletal mechanical abnormality is described as arthrogenic (due particularly to hip disorders) and a second type as osteogenic (due to a shortened limb).

Neurogenic gaits and myogenic gaits are considered together below, under the heading ‘Neurological disorders of gait’.

Psychogenic or ‘hysterical’ gait may have to be considered if the gait is bizarre or seems greatly exaggerated. On the other hand, loss of confidence, especially in the elderly, is an important cause of gait disturbance. However, many abnormal gaits that are caused by neurological disease may also appear bizarre, and caution is advised. Doubtful cases should be referred for an expert opinion.

EXAMINATION OF GAIT AND POSTURE[1]

Disorders of gait and posture go hand in hand because of a common physiological process.

The source of the abnormality is indicated.

1. Ask the patient to stand. Note any difficulty in reaching a standing position. Difficulty ° proximal muscle weakness.
2. Ask the patient to stand with eyes closed (Romberg's test). If positive (sways or falls) ° loss of proprioception (e.g. peripheral neuropathy).
3. Ask the patient to walk (ensure sufficient testing length).
    - Gait initiation hesitancy ° basal ganglia or frontal cortex
    - Stride length very short ° basal ganglia or frontal cortex irregular ° cerebellar
    - Narrow or broad base narrow ° UMN, muscle weakness, basal ganglia broad ° cerebellum, proprioception, vestibular
    - Stiff or ‘sloppy’ stiff ° UMN, basal ganglia sloppy ° LMN, muscle weakness
    - Heel strike loss of normal strike ° UMN, LMN, myopathy
    - High stepping positive ° LMN (distal), proprioception, muscle weakness

- Arm swing decreased ° basal ganglia, UMN (frontal lobes)
- Pelvis control Trendelenburg gait ° proximal muscle weakness

4. Ask the patient to walk using provocation tests.
    - ‘Tightrope’ walk tests proprioception
    - Stand on tiptoes and back on the heels tests distal muscle weakness

Note: Peripheral neuropathy is a common cause of an LMN lesion.

NEUROLOGICAL DISORDERS OF GAIT

Cerebellar gait

The patient has difficulty standing, has a broad-based stance and an unsteady gait, with swaying from side to side. It has been described as the ‘dirty diaper’ gait. The patient may stagger or bump into walls and may be accused of being drunk. When asked to walk in a straight line the patient tends to veer towards the side of the lesion. Ask the patient to walk ‘heel to toe’ in a straight line and perform a heel - knee - shin test to demonstrate cerebellar ataxia. Significant causes are multiple sclerosis, alcoholic cerebellar degeneration and space-occupying lesions.

Basal ganglia gait (Parkinson's type)

Identifying this disorder in its early stages can be difficult as the first sign may be a limp with one leg being described as weak or stiff or slow.[2] However, the typical gait is a shuffling of the feet with small steps in a forward flexed posture, rather like a person walking with a belt around the ankles. This leads to a hurrying (festinant) gait as though there is an impending feeling of falling forward.

Spastic gait

Spastic gait may be regarded as the typical bilateral or paraplegic gait, or as a hemiplegic gait. With the former the gait affects both legs - they are stiff or weak, leading to slow, jerking walking, dragging of the feet and scraping of the toes. This scuffing can be heard as the front of the shoe drags along the ground. Every step can be a struggle and the patient may appear as though walking through glue on the floor.

A scissor-type gait will develop with bilateral hip adduction. Spasticity is caused by a UMN lesion, including multiple sclerosis and spinal cord compression. The typical UMN posture is of a flexed upper limb and an extended lower limb.

With hemiplegia the patient drags the affected leg stiffly with the hips adducted, the knee extended and the foot plantar flexed, leading to scraping of the toes. Mounting stairs can be very difficult, especially if clonus is induced with dorsiflexion of the foot.

Foot drop gait

The patient cannot dorsiflex the foot, leading to a high-stepping gait with extra flexion of the hip and knees to lift the foot off the ground. The foot then slaps down on the ground.

Vestibular gait

If unilateral the patient tends to veer off to the side of the lesion.

Apraxia

With apraxia of gait (due to a prefrontal lobe lesion) there is a failure of control of the legs. The patient may stand up and try to walk but looks with bewilderment at the legs and moves them in an inappropriate manner with a broad-based, smallstepping unsteady gait. Turning is difficult. Apraxia is caused by bilateral cortical involvement, such as hydrocephalus, multi-infarct states and tumours of the corpus callosum.[2]

Neurogenic claudication

With intermittent claudication of the cauda equina, due to spinal canal stenosis, the patient develops pain in the leg after walking a certain distance. However, weakness and numbness are usually more prominent than pain.

Drop attacks

In a drop attack the patient suddenly falls to the ground, without other symptoms, and gets up almost immediately. There is no loss of consciousness. Drop attacks can be caused by disorders such as epilepsy, Parkinson's disease and vertebrobasilar insufficiency. However, in most cases, particularly middle-aged and elderly women, there is no obvious cause.

Waddling gait

A waddling gait is usually caused by muscular dysfunction affecting the pelvic girdle muscles and trunk. There is a wide-based gait with a marked 'rocking and rolling' body swing from side to side and related compensatory movements of the pelvis, that is, a bilateral Trendelenburg gait.

Proximal muscle weakness

The patient may complain when getting out of a low chair or going up or downstairs. The weakness can be demonstrated by asking the patient to squat down and, after a second, rise from the squatting position.[2] Waddling of gait reflects extreme cases. Causes include myopathies, motor neurone disease and Guillain-Barré syndrome.

Distal muscle weakness

This causes a high-stepping gait as the foot is floppy and tends to flap, with walking similar to foot drop gait. Causes include peripheral neuropathy, myotonic dystrophy and peroneal muscular atrophy.

LIMP

Limp is a symptom commonly associated with painful disorders of the lower limb, especially of the hip and knee joints. A limp implies an asymmetrical gait pattern caused by one of four general factors:

1. unequal leg length
2. antalgic (painful) gait (e.g. hip disorder)
3. restricted joint movement (e.g. ankylosed knee)
4. neuromuscular weakness (e.g. poliomyelitis)

Limp has an inseparable relationship with painful hip and buttock conditions, especially those of the hip. Painful hip and pelvic conditions that cause limp are presented on p. 696.

Limp in adults

In adults the cause of limp is usually more obvious than in children and is commonly due to degenerative osteoarthritis of the hip or knee, to spinal disorders, especially sciatica caused by a disc prolapse, or to overuse disorders of the knee, ankle or foot.

Limp in children

The child who limps presents an interesting diagnostic dilemma. The limp must be considered to be due to a definite organic cause, although conversion reactions can be a factor.[3] It is appropriate to focus initially on the hip. The diagnostic strategy is presented in Table 69.1.

Table 69.1 Limp in children: diagnostic strategy (modified)

Q. Probability diagnosis

A. Post trauma/intense exercise causing strain syndromes

Ill-fitting shoes

Hip disorders, esp. transient synovitis

Heel disorders (12 - 14 years)

Q. Must not be missed

A. Toddlers: DDH

child abuse

septic arthritis

foreign body (e.g. needle in foot)

4 - 8 years: Perthes' disorder

transient synovitis

Adolescents: SCFE

avulsion injuries (e.g. ischial tuberosity)

osteochondritis dissecans of knee

Duchenne's muscular dystrophy

All groups: septic infections

- septic arthritis
- osteomyelitis
- tuberculosis tumour (e.g. osteosarcoma)

juvenile chronic arthritis spinal disorders

- discitis
- fracture

Q. Pitfalls (often missed)

A. Foreign body (e.g. in foot)

Osteochondritis (aseptic necrosis)

- femoral head - Perthes' disorder
- knee - Osgood-Schlatter disorder
- calcaneum - Sever's disorder
- navicular - Köhler's disorder

Myalgia ° ‘growing pains’

Overuse syndrome (esp. adolescent)

- patellar tendonitis (jumper's knee)

Stress fractures (e.g. tibia, femoral neck, navicular)

Limp can be considered as acute, subacute or chronic. An acute limp may be due to injury, infection (osteomyelitis, septic arthritis), spinal injuries, a fracture or an irritable hip (transient synovitis). Subacute causes include juvenile rheumatoid arthritis and tumour or leukaemia. Chronic causes include cerebral palsy, DDH, Perthes' disorder and chronic SCFE.

Key checkpoints

- Trauma, sepsis and DDH are perhaps the most common reasons for an infant to limp and refuse to walk. However, a painless waddling gait suggests DDH or Perthes' disorder, which usually begins with a painless limp.
- Multiple fractures and epiphyseal separations in toddlers are highly suggestive of child battering; a skeletal survey should be ordered if this is suspected.
- Perthes' disorder can present from ages 4 to 12 years but is usually found from 4 - 8 years with a peak age of 5 - 7 years.
- Infections of and around the hip joints are most common in infancy. Classically, the hip is held immobile in about 30° of flexion with slight abduction and exter nal rotation. The commonest organism is *Staphylococcus aureus*, followed by *Haemophilus influenzae*.
- Tuberculosis may also occur in children (usually under 10 years) with a presentation similar to Perthes' disorder.
- SCFE typically presents in the obese adolescent (10 - 15 years) with knee pain and a slight limp.
- Growing pains are a controversial issue but do appear to exist as an aching myalgia usually manifest in the leg muscles (anterior thigh, calf, posterior knee). The pain is bilateral, non-articular and usually unrelated to activity.

Diagnostic approach

History

The age of the patient gives a diagnostic pointer. A careful history, especially of trauma, may lead to the diagnosis. A history of injury is usually but not always available. The relationship of the limp to exercise and footwear is significant. The location of any associated pain is relevant: low back pathology can refer to the buttocks and hip pathology can cause knee pain.

Examination

The hip and knee joints should be carefully examined if the source of limp has no specific localisation. Get the child to walk and run on the toes and heels (if appropriate). Note the gait and check whether it is antalgic (painful), hemiplegic (the arm is held out in a balancing action) or Trendelenburg (classic for DDH). Look for evidence of muscular dystrophy.

Investigations

The following have to be considered:

- FBE and ESR
- blood culture
- needle aspiration of joint
- radiological
    - plain X-ray
    - ultrasound
    - bone scan
    - CT or MRI scan

Management

Management is based on the cause. Surgical drainage supplemented with antibiotics is essential for septic arthritis. If the child initially has a limp and cannot walk, admit them to hospital for:

- skin traction
- FBE and ESR
- ultrasound of hip
- blood culture

Specific conditions

Osteomyelitis

Osteomyelitis should be suspected in a child with an acute febrile illness and metaphyseal tenderness and the child admitted to hospital. Blood should be collected for FBE, ESR and culture. A plain X-ray and nuclear scan are valuable but may not provide immediate confirmation of the diagnosis. The child should be admitted to hospital, an IV line should be inserted and IV antibiotics commenced to cover the organisms involved:

- > 5 years: flucloxacillin IV for 4 - 6 days then oral
- < 5 years: di(flu)cloxacillin for at least 21 days[4]

Septic arthritis

Septic arthritis should be suspected in a child with pyrexia and an acute arthritis with limited motion. Manage as for osteomyelitis.

Bone tumour

Chronic limp is a common presentation of malignant bone tumours. Radiological investigation is mandatory.

Irritable hip syndrome (transient synovitis)

Typical age is 3 - 8 years and the child presents with an acute limp with restricted hip motion. Plain X-ray is normal. Orthopaedic assessment is recommended.

Perthes' disorder, SCFE and DDH

Refer to Chapter 65.

PAGET'S DISEASE

Paget's disease of bone (osteitis deformans) is a chronic disorder of the adult skeleton in which new soft bone replaces localised areas of normal bone. The cause is unknown but a viral aetiology is suspected. There is a great increase in bone turnover with osteoclastic resorption followed by increased osteoblastic activity. The disorder is quite common:

- 1 in 200 of the population at 40 years
- 1 in 10 of population at 90 years

Paget's disease is usually asymptomatic but some patients may present with deep aching pain in the lower back and lower limbs. They may also present with a disturbance of gait due to unequal leg length, osteoarthritis of associated joints such as the knee or hip, or a change in the distribution of mechanical forces in the lower extremities (Figure 69.1).



Figure 69.1 Paget's disease of the left leg showing deformity of the tibia with a ‘sabre’ tibia due to its enlargement of length and bulk

Features

- Male:female ratio ° 2:1.
- 95% asymptomatic (discovered by X-ray or raised serum alkaline phosphatase [ALP] level).
- Symptoms may include joint pain and stiffness (e.g. hips, knees), bone pain (usually spine), deformity, headache and deafness.
- Bone pain is typically deep and aching; it occurs at rest, particularly at night.
- Signs may include deformity, enlarged skull (‘hats don’t fit any more’), bowing of tibia, waddling gait, hyperdynamic circulation (Figure 69.2).


skull enlargement
cranial nerve compression
II, V, VII
VIII – deafness
cardiac hypertrophy
? high output failure
arthritis
Bone problems
• pain
• deformities
• pathological fractures
• osteogenic sarcoma
bowed (legs) tibia

Figure 69.2 Paget's disease: possible clinical features

- Bones most commonly affected, in decreasing order, are the pelvis, femur, skull, tibia, vertebrae, clavicle and humerus.

Diagnosis

- Raised serum ALP level (often very high > 1000 U/L).

Note: calcium and phosphate normal.

- Plain X-ray: dense expanded bone - best seen in skull and pelvis.

Note: Can mimic prostatic secondaries so every male Pagetic patient should have a DRE and serum PSA test.

- Bone isotopic scans: useful in locating specific areas.
- Watch for the uncommon complication of osteogenic sarcoma.

Treatment [5]

The two major goals are relief of pain and prevention of long-term complication (e.g. deafness, deformities).

Localised and asymptomatic disease requires no treatment.

Three groups of drugs are currently available:

- the calcitonins
- the biphosphonates - etidronate, pamidronate disodium (APD), alendronate, risedronate
- various antineoplastic agents (e.g. mithramycin)

Biphosphonates have become the preferred drugs for first-line therapy and include:[5]

- alendronate 40 mg (o) daily for 6 months (oesophagitis can be problematic)
- pamidronate disodium 30 - 60 mg IV infused over 2 - 4 hours (usually the preferred option)
- risedronate 30 mg (o) daily for 2 months
- tiludronate 400 mg (o) daily for 3 months

All oral agents should be taken on an empty stomach.

Repeated doses may rarely be required in severe cases as judged by symptoms and disease activity (e.g. monitoring with serum ALP).

LEG SWELLING

Diagnostic features and pitfalls

- Not all swollen legs require investigation and treatment.
- The significance of leg swelling varies according to the age group, to whether it is bilateral or unilateral, and whether the onset is sudden or gradual (Table 69.2).

Table 69.2 Causes of swollen legs

Physiological
• prolonged standing or walking
• prolonged sitting (e.g. elderly on long journey)
• pregnancy
• hot weather

- mechanical factors (e.g. constricting garters/pantyhose)

Local disorders

- skin (e.g. allergy)
- arthritis with particular oedema
- infection (e.g. cellulitis, filariasis)
- trauma
- thrombophlebitis
- vascular obstruction

  -venous (e.g. DVT, varicose veins)

  -lymphatic ??lymphoedema

Generalised disease

- cardiac (e.g. CCF)
- kidney (e.g. nephrotic syndrome)
- hepatic (e.g. cirrhosis)

Drugs

- NSAIDs, antihypertensives (e.g. nifedipine), oestrogens, others

Lipoedema

Lymphoedema: primary or secondary

Idiopathic (periodic) oedema

- DVT must be considered in all unilateral cases and ultrasound examination performed if appropriate (Table 69.3).

Table 69.3 Swollen legs: diagnostic perspective

Q.	A.
Q. Probability diagnosis	A. Chronic venous insufficiency (varicose veins) Physiological (e.g. dependency)
Q. Must not be missed	A. Deep venous thrombosis Thrombophlebitis Obstruction from pelvic cancer
Q. Pitfalls	A. Drugs (sodium retention) Idiopathic (cyclical) oedema

- If a DVT is present, consider occult malignancy (e.g. carcinoma pancreas).
- Consider pelvic cancer causing lymphatic obstruction in a woman > 40 years presenting with painless unilateral leg oedema.
- A drug history is essential as several drugs can cause oedema.
- Pitting oedema is a feature of venous thrombosis or insufficiency, not lymphatic obstruction.

Investigations

Select from these first-line tests:

- urinalysis (? albumin)
- FBE and ESR

- serum urea/creatinine
- serum albumin/LFTs
- ultrasound (DVT screen)
- other radiographs (e.g. CT scan, venogram)

Calf swelling of sudden onset

Causes to consider:

- acute arterial occlusion
- ruptured Baker's cyst
- ruptured medial head of gastrocnemius
- DVT (usually gradual)
- cellulitis/erysipelas
- compartment syndrome

Pain accompanies most of these conditions but the absence of pain does not exclude DVT or thrombophlebitis. Refer to Chapter 66.

Lipoedema

Lipoedema is the development of bilateral leg swelling that does not involve the feet (in lymphoedema the swelling develops in the most distal part of the foot).

Features

- Exclusive to obese women
- Spares the feet
- Bilateral and symmetrical distribution of fat

REFERENCES

1. Horne M. Gait and postural disorders. Monash University Neurology Notes 1996: 1 - 4.
2. Kincaid-Smith P, Larkins R, Whelan G. *Problems in Clinical Medicine*. Sydney: MacLennan & Petty, 1990: 190 - 4.
3. Paxton G, Munro J eds. *Paediatric Handbook* (7th edn). Melbourne: Blackwell Science, 2003: 550 - 4.
4. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 11). Melbourne: Therapeutic Guidelines Ltd, 2000: 178 - 80.
5. Shenfield G (Chair). *Therapeutic Guidelines: Endocrinology* (Version 3). Melbourne: Therapeutic Guidelines Ltd, 2004: 121 - 6.

CHAPTER 70. PALPITATIONS

> The most important requirement of the art of healing is that no mistakes or neglect occur. There should be no doubt or confusion as to the application of the meaning of complexion and pulse. These are the maxims of the art of healing.
>
> HUANG TI (THE YELLOW EMPEROR) 2697 - 2597 BC

PALPITATIONS

Palpitations are an unpleasant awareness of the beating of the heart. By definition it does not always imply 'racing' of the heart but any sensation in the chest, such as 'pounding', 'flopping', 'skipping', 'jumping', 'thumping' or 'fluttering' of the heart. The problem requires careful attention and reassurance (if appropriate) because heartbeat is regarded as synonymous with life. To the practitioner it may simply represent anxiety or it could be a prelude to a cardiac arrest.

KEY FACTS AND CHECKPOINTS

- The symptom of palpitations is suggestive of cardiac arrhythmia but may have a non-cardiac cause.
- Palpitations not related to emotion, fever or exercise suggest an arrhythmia.
- Perhaps the commonest arrhythmia causing a patient to visit the family doctor is the symptomatic premature ventricular beat/complex (ventricular ectopic).
- The commonest cause of an apparent pause on the ECG is a blocked premature atrial beat/complex (atrial ectopic).
- A 12-lead electrocardiographic diagnosis is mandatory. If the cause is not documented, an ambulatory electrographic monitor (e.g. Holter) may be used.
- Consider myocardial ischaemia as a cause of the arrhythmia.
- Consider drugs as a cause, including prescribed drugs and non-prescribed such as alcohol, caffeine and cigarettes.
- Common triggers of paroxysmal supraventricular tachycardia (PSVT) include anxiety and cigarette smoking.
- The commonest mechanism of any arrhythmia is re-entry.
- Get patients to tap out the rate and rhythm of their abnormal beat.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 70.1, which includes significant causes of palpitations.

Table 70.1 Palpitations: diagnostic strategy model

Q. Probability diagnosis
A. Anxiety
Premature beats (ectopics)
Sinus tachycardia
Drugs (e.g. stimulants)
Q. Serious disorders not to be missed
A. Myocardial infarction/angina
Arrhythmias
• ventricular tachycardia
• bradycardia
• sick sinus syndrome
• torsade de pointes

WPW syndrome Electrolyte disturbances • hypokalaemia • hypomagnesaemia • hypoglycaemia (type 1 diabetes)
Q. Pitfalls (often missed)
A. Fever/infection Pregnancy Menopause Drugs (e.g. caffeine, cocaine) Mitral valve disease Aortic incompetence Hypoxia/hypercapnia *Rarities* Tick bites (T1 - 5) Phaeochromocytoma
Q. Seven masquerades checklist
A. Depression Diabetes indirect Drugs Anaemia Thyroid disorder Spinal dysfunction UTI possible
Q. Is the patient trying to tell me something?
A. Quite likely. Consider cardiac neurosis, anxiety.

Probability diagnosis

If the palpitations are not caused by anxiety or fever, the common causes are sinus tachycardia and premature complexes (atrial or ventricular). Sinus tachycardia, which by definition is a rate of 100 - 160/minute, may be precipitated by emotion, stress, fever or exercise.

PSVT and atrial fibrillation are also quite common arrhythmias. Some cardiologists claim that the commonest arrhythmia causing a patient to visit the family doctor is the symptomatic ventricular ectopic beat.[1]

Sinus tachycardia can be differentiated clinically from PSVT in that it starts and stops more gradually than PSVT (abrupt) and has a lower rate of 100 - 150 compared with 160 - 220.

Important causes of tachyarrhythmias are:

- ischaemic heart disease, especially acute coronary syndrome
- hypertension
- heart failure
- mitral disease

- thyrotoxicosis
- atrial septal deficit

Serious disorders not to be missed

It is vital not to overlook acute coronary syndromes, as a cause of the arrhythmia manifesting as palpitations. About 25% of infarcts are either silent or unrecognised.

Sinister life-threatening arrhythmias are:

- ventricular tachycardia
- atypical ventricular tachycardia (torsade de pointes)
- sick sinus syndrome (SSS)
- complete heart block

It is also important not to miss:

- hypokalaemia
- hypomagnesaemia

Pitfalls

There are many pitfalls in the diagnosis and management of arrhythmias, especially in the elderly where symptoms of infection may be masked. Palpitations associated with the menopause can be overlooked. Valvular lesions, usually associated with rheumatic heart disease, such as mitral stenosis, and aortic incompetence may cause palpitations. The rare tumour, phaeochromocytoma, presents with palpitations and the interesting characteristic of postural tachycardia (a change of more than 20 beats/minute). The toxin from tick bites in dermatomes T1 - 5 can cause palpitations.

General pitfalls

- Misdiagnosing PSVT as an anxiety state
- Overlooking a cardiac arrhythmia as a cause of syncope or dizziness
- Overlooking atrial fibrillation (AF) in the presence of a slow heartbeat
- Overlooking mitral valve prolapse in a patient, especially a middle-aged woman, presenting with unusual chest pains and palpitations (auscultate in standing position to accentuate click(s) ± murmurs)

Seven masquerades checklist

Surprisingly, all the masquerades have to be considered, either as direct or indirect causes: depression, especially with anxiety and in the postpartum period; diabetes, perhaps as an arrhythmia associated with a silent myocardial infarction or with hypoglycaemia; drugs as a very common cause (Table 70.2); anaemia, causing a haemodynamic effect; hyperthyroidism; spinal dysfunction of the upper thoracic vertebrae T1 - 5; and urinary tract infection, especially in the elderly.

Table 70.2 Some drugs that cause palpitations

Alcohol
Aminophylline
Amphetamines
Anti-arrhythmic drugs
Antidepressants
• tricyclics
• MAO inhibitors
caffeine

Cocaine
Class 1_A and 1_C drugs
Digitalis
Diuretics → K ↓, Mg ↓
Glyceryl trinitrate
Sympathomimetics
• in decongestants
• salbutamol
• terbutaline
Thyroxine

PSVT has been described as resulting from injury or dysfunction of the upper thoracic spine (especially T4 and T5) in the absence of organic heart disease.[2] The author has personally encountered several cases of PSVT alleviated by normalising function of the spine.

Psychogenic considerations

Emotional factors can precipitate a tachycardia, which in turn can exaggerate the problem in an anxious person. Some people have a cardiac neurosis, often related to identification with a relative or friend. A family history of cardiac disease can engender this particular anxiety. Evidence of anxiety and depression should be sought in patients presenting with palpitations without clinical evidence of cardiovascular disease.

THE CLINICAL APPROACH

Careful attention to basic detail in the history and examination can point the way clearly to the clinical diagnosis.

History

Ask the patient to describe the onset and offset of the palpitations, the duration of each episode and any associated features. Then ask the patient to tap out on the desk the rhythm and rate of the heartbeat experienced during the 'attack'. If the patient is unable to do this, tap out the cadence of the various arrhythmias to find a matching beat.

An irregular tapping 'all over the place' suggests atrial fibrillation, while an isolated thump or jump followed by a definite pause on a background or a regular pattern indicates premature beats (ectopics/extrasystoles) usually of ventricular origin. The thump is not the abnormal beat but the huge stroke volume of the beat following the compensatory pause.

Key questions

- Do the palpitations start suddenly? How long do they last?
- What do you think brings them on?
- Are they related to stress or worry or excitement?
- What symptoms do you notice during an attack?
- Do you have pain in the chest or breathlessness during the attack?
- Do you feel dizzy or faint during the attack?
- What medications do you take?
- How much coffee, tea, Coke do you drink?
- Have you been using nasal decongestants?
- Did you eat Chinese food before the attack?
- Do you smoke cigarettes, and how many?
- Do you take any of the social drugs, such as cocaine or marijuana?

- Have you ever had rheumatic fever?
- Have you lost weight recently or do you sweat a lot?

Chest pain may indicate myocardial ischaemia or aortic stenosis; breathlessness indicates anxiety with hyperventilation, mitral stenosis or cardiac failure; dizziness or syncope suggests severe arrhythmias such as SSS and complete heart block, aortic stenosis and associated cerebrovascular disease.

Physical examination

The ideal time to examine the patient is while the palpitations are being experienced. often this is not possible and the physical examination is normal. Measurement of the heart rate may provide a clue to the problem.

As a working guide, a rate estimated to be about 150 beats per minute suggests PSVT, atrial flutter/fibrillation or ventricular tachycardia (Figure 70.1). A rate less than 150 beats per minute is more likely to be sinus tachycardia, which may be associated with exercise, fever, drugs or thyrotoxicosis.[3]



Figure 70.1 Heart rate guide to causes of various arrhythmias

The nature of the pulse, especially the pulse pressure and rhythm, should be carefully evaluated (Figure 70.2). Look for evidence of fever and infection and features of an anxiety state or depressive illness.



Form
Significance
Normal
Weak, small volume
'shock'
mitral stenosis
constrictive pericarditis
pericardial effusion
Large volume, bounding
hypertension
Collapsing
aortic incompetence
high output states
e.g. anaemia
fever
A-V fistula
Plateau
aortic stenosis
Pulsus alternans
severe myocardial dysfunction
Pulsus bigeminus
premature ectopic complex following every sinus beat
— consider digoxin toxicity

Figure 70.2 Various pulse forms

Have the patient hyperventilate for 3 minutes to determine whether the arrhythmia is induced. Evidence of underlying disease such as anaemia, thyroid disorder, alcohol abuse or cardiac disease including the JVP and pulmonary congestion should be sought. Also look for evidence of a mitral valve prolapse (mid-systolic click; late systolic murmur). Possible signs in the patient presenting with palpitations are shown in Figure 70.3.[4]


bright eyes 'thyroid glitter'
ocular fundal changes
exophthalmos
of hyperthyroidism
Cannon waves in jugular vein
(CHB)
hypertension
collapsing pulse
senile arcus
malar flush
goitre
signs of • aortic stenosis
• aortic incompetence
• mitral incompetence
• mitral stenosis
• others
cardiac enlargement
lung—basal crackles
thickened
atherosclerotic
radial artery
palm signs
• sweaty (hyperthyroidism)
• pallor of anaemia
peripheral vascular disease
ankle oedema

Figure 70.3 Signs to consider in a patient with palpitations

Diagnostic investigations

The number and complexity of investigations should be selected according to the problem and test availability. A checklist would include:

- blood tests (for underlying disease)
  - haemoglobin and film
  - thyroid function tests
  - serum potassium and magnesium
  - serum digoxin ? digitalis toxicity
  - virus antibodies ? myocarditis
- chest X-ray
- cardiac (ischaemia and function)
  - ECG (12 lead)
  - ambulatory 24-hour ECG monitoring
  - echocardiography (to look for valvular heart disease and assess left ventricular function)
  - electrophysiology studies
  - exercise stress test (? underlying CAD)

PALPITATIONS IN CHILDREN

Children may complain of palpitations which may be associated with exercise, fever or anxiety. Various arrhythmias can occur with three requiring special consideration - PSVT, heart block and ventricular arrhythmias.[5]

PSVT is characterised by beats at 200 - 300 per minute, the fastest rates occurring in infants. The cause is often not found but some children have ECG abnormalities compatible with the Wolff-Parkinson-White (WPW) syndrome. The recommended firstline treatment of PSVT is vagal stimulation via the application of ice packs to the upper face (forehead, eyes and nose) of the affected infant. Intravenous adenosine will usually terminate the episode.

A particular concern is these children who have the familial long QT syndrome. They are prone to develop ventricular tachyarrhythmias, which may lead to sudden death. Consider it in children developing syncope on exertion.

PALPITATIONS IN THE ELDERLY

The older the patient the more likely the onset of palpitations due to cardiac disease such as myocardial infarction/ischaemia, hypertension, arrhythmias and drugs, especially digoxin. Occasional atrial and ventricular arrhythmias, especially premature complexes (ectopics), occur in 40% of old people[6] and treatment is rarely required. Atrial fibrillation occurs in 5 - 10% of patients over 65 years of age, 30% of whom have no clinical evidence of cardiovascular disease. A rapid ventricular rate with symptoms is the only indication for digoxin in the elderly but beware of SSS, especially if dizziness or syncope accompanies the fibrillation.

In the elderly, thyrotoxicosis may present as sinus tachycardia or atrial fibrillation with only minimal signs - the so-called ‘masked thyrotoxicosis’ - so it is easy to overlook it. The only clue may be bright eyes (‘thyroid glitter’) due to conjunctival oedema.

ARRHYTHMIAS

Facts and figures

- Cardiac arrhythmias account for about 25% of management decisions in cardiology (Table 70.3).

Table 70.3 Types of arrhythmias

Non-pathological sinus rhythms
• sinus arrhythmia
• sinus bradycardia
• sinus tachycardia
Pathological bradycardias
• sinus node disease (sick sinus syndrome)
• atrioventricular (AV) block
- first degree AV block
- second degree AV block
- third degree (complete) AV block
Pathological tachyarrhythmias
1 *Atrial*
• atrial premature (ectopic) complexes
• PSVT
• atrial flutter
• atrial fibrillation (AF)
2 *Ventricular*

- ventricular premature complexes
- ventricular tachycardia
- ventricular fibrillation
- torsade de pointes (twisting of points)

- Commonest are premature (ectopic) ventricular beats and atrial fibrillation.
- PSVT is next most common - 6 per 1000 of population.
- The commonest mechanism of paroxysmal tachycardias is re-entry (Figure 70.4).



Figure 70.4 Diagrammatic mechanism of re-entry tachycardia

- Electrophysiological studies are the gold standard investigation for tachycardias but are rarely needed for diagnosing most arrhythmias.
- Almost all antiarrhythmic drugs have a proarrhythmic potential (i.e. they may worsen existing arrhythmias or provoke new arrhythmias in some patients) (refer to Table 70.4). Always consider the no-treatment option.

Table 70.4 Electrophysiological classification of common antiarrhythmic drugs (after Vaughan Williams)

Class	Drug	Usual dosage	Common side-effects
1_A	Disopyramide	100 - 200 mg qid	Blurred vision, dry mouth, urinary problems in males (avoid in men > 50)
	Procainamide	1 g qid IV use	Anorexia, nausea, urticaria
	Quinidine	2 - 3 SR tabs (0.25 g) bd	Diarrhoea, headache, tinnitus
1_B	Lignocaine	IV use	Nausea, dizziness, tremor

	Mexiletine	200 mg tid	Nausea, vomiting, tremor, dizziness
1_C	Flecainide	100 mg bd	Nausea, dizziness, rash
II	Beta-blockers	various	Fatigue, insomnia, nightmares, hypotension, bronchospasm. Avoid in asthmatics
III	Amiodarone	SVT: 200 mg daily VT: 400 mg daily	Rash, pulmonary fibrosis, thyroid, hepatic and CNS effects
	Sotalol	80 - 160 mg bd	As for beta-blockers
IV	Verapamil	(SR) 160 - 480 mg daily	Constipation, dizziness, hypotension
	Diltiazem	(CR) 180 - 360 mg daily	Hypotension, headache

Note: Sotalol is a beta-blocker and thus is a class II and III agent. All drugs are taken orally unless IV indicated. Adenosine and digoxin are not classified.

- Avoid digoxin in cases with an accessory pathway.
- If ‘quinidine syncope’ occurs, consider torsade de pointes as the cause.
- Any patient commencing anti-arrhythmic therapy should have a 12-lead ECG 1 - 2 weeks later to check the QT interval. If prolonged, treatment should usually be ceased.
- The two main indications for permanent pacemaking are SSS (only if symptomatic) and complete heart block.

Management strategies

- Treat the underlying cause.
- Give appropriate reassurance.
- Provide clear patient education.
- Explain about the problems of fatigue, stress and emotion.
- Advise moderation in consumption of tea, coffee, caffeine-containing soft drinks and alcohol.
- Advise about cessation of smoking and other drugs.

PREMATURE (ECTOPIC) COMPLEXES

Atrial premature complexes

- These are usually asymptomatic.
- Management is based on reassurance.
- Check lifestyle factors such as excess alcohol, caffeine, stress and smoking; avoid precipitating factors.
- Treatment is rarely required and should be avoided if possible.
- At present there is no ideal antiectopic agent.
- They may be a forerunner of other arrhythmias (e.g. PSVT, atrial fibrillation).

Ventricular premature complexes

- These are also usually asymptomatic (90%).
- They occur in 20% of people with ‘normal’ hearts.
- Symptoms are usually noticed at rest in bed at night.
- Check lifestyle factors as for atrial premature beats.
- Drugs that can cause both types of premature beats include digoxin and sympathomimetics.
- Look for evidence of ischaemic heart disease, mitral valve prolapse (especially women), thyrotoxicosis and left

ventricular failure.

- Ventricular premature complexes may be a forerunner of other arrhythmias (e.g. ventricular tachycardia).
- If symptomatic but otherwise well with a normal chest X-ray and ECG, reassure the patient.
- Drug therapy: Never commence drug therapy without performing an echocardiograph. This will help to guide the choice of agent. Class 1 agents can make the arrhythmia worse or even life-threatening if there is reduced ventricular function. If this is the case, the patient should be referred to a cardiologist.

SUPRAVENTRICULAR TACHYCARDIA[7]

- SVT can be paroxysmal or sustained.
- Rate is 150 - 220/minute.
- There are at least eight different types of SVT with differing risks and responses to treatment.
- PSVT commonly presents with a sudden onset in otherwise healthy young people.
- Passing copious urine after an attack is characteristic of PSVT.



normal sinus rhythm



sinus bradycardia and sinus arrhythmia—rate approx. 55 per minute



sinus tachycardia—rate approx. 100 per minute



complete heart block



atrial flutter



atrial fibrillation



atrial premature complexes



ventricular premature complexes



supraventricular tachycardia



ventricular tachycardia



ventricular fibrillation

Figure 70.5 Tracings of important arrhythmias

- Look for predisposing factors such as an accessory pathway and thyrotoxicosis.
- Approximately 60% are due to atrioventricular (AV) node re-entry and 35% due to accessory pathway tachycardia (e.g. WPW).[8]
- Look for evidence of accessory pathways after reversion because accessory pathways can lead to sudden death (avoid digoxin in WPW).
- Consider SSS in a patient with SVT and dizziness.

Wolff-Parkinson-White syndrome

The structural basis for the arrhythmia of SVT in WPW syndrome is an accessory pathway that bypasses the AV node. A typical ECG shows a short PR interval and slurred upstroke of the QRS complex (delta wave). Patients are prone to sudden attacks of SVT. Up to 30% of patients will develop atrial fibrillation or flutter. Even one episode of PSVT requires consideration for radiofrequency ablation.[9]

Management of PSVT

1. Vagal stimulation can be attempted. Carotid sinus massage is the first treatment of choice. Other methods of vagal stimulation include:
    - Valsalva manoeuvre (easiest for patient)
    - self-induced vomiting
    - ocular pressure (avoid)
    - cold (ice) water to face
    - immersion of the face in water
2. If vagal stimulation fails, give adenosine IV (try 6 mg first over 5 - 10 seconds, then 12 mg in 2 minutes if unsuccessful, then 18 mg in 2 minutes if necessary). Second-line treatment is verapamil IV 1 mg/min up to 10 - 15 mg (provided patient is not taking a beta-blocker).[8]
    Precautions

  - Adenosine causes less hypotension than verapamil but may cause bronchospasm in asthmatics
  - Use only if narrow QRS and BP > 80 mm Hg
  - Carefully monitor blood pressure
  - AVOID verapamil if taking beta-blockers and persistent tachycardia with QRS complexes > 0.14 s (suggests ventricular tachycardia)

3. In the rare event of failure of medical treatment, consider DC cardioversion or overdrive pacing.

Prophylaxis

To prevent recurrences use flecainide (only if no structural heart damage) or sotalol. If these agents fail, consider amiodarone. Do an echocardiograph first to exclude structural heart disease. Radiofrequency catheter ablation, which is usually curative, is indicated for frequent attacks not responding to medical therapy.

Carotid sinus massage[1]

Carotid sinus massage causes vagal stimulation and its effect on SVT is all or nothing. It has no effect on ventricular tachycardia. It slows the sinus rate and breaks the SVT by blocking AV nodal conduction.

Method

- Locate the carotid pulse in front of the sternomastoid muscle just below the angle of the jaw (Figure 70.6).



Figure 70.6 Carotid sinus massage

- Ensure that no bruit is present.
- Rub the carotid with a circular motion for 5 - 10 seconds.
- Rub each carotid in turn if the SVT is not ‘broken’, but never both together.

In general, right carotid pressure tends to slow the sinus rate and left carotid pressure tends to impair AV nodal conduction.

Precautions

In the elderly (risk of embolism or bradycardia).

ATRIAL FIBRILLATION

Facts and figures

- A common problem (9% incidence in the over 70 age group).
- It usually presents with an irregular ventricular rate of about 160 - 180 beats per minute in untreated patients with a normal AV node.
  It tends to fall into one of the 'three Ps' patterns:
    - paroxysmal AF
    - persistent AF
    - permanent (chronic) AF

All types appear to have a similar risk for thromboembolism.

- Remember to look for the underlying cause: myocardial ischaemia (15% of cases), mitral valve disease, thyrotoxicosis, hypertension, cardiomyopathy including chronic alcohol dependence, alcohol binge.
- No cause is found in 12% - isolated[9, 10] atrial fibrillation.
- All patients should have thyroid function tests and an echocardiograph to help find a cause.
- With sustained atrial fibrillation there is a 5% chance per annum of embolic episodes. There is a fivefold risk of CVA overall.
- The risk of CVA is greater in those with previous CVA, valvular heart disease, prosthetic mitral valve and cardiac failure.
- For reversion anticoagulate with warfarin for 4 weeks beforehand and maintain for 4 weeks after.
- Digoxin controls the ventricular rate but does not terminate or prevent attacks.
- Sotalol, flecainide and amiodarone are used for conversion of atrial fibrillation and maintenance of sinus rhythm. Flecainide should never be prescribed in patients with reduced LV function.
- Evidence basis: RCTs show that digoxin was beneficial for lowering the ventricular rate in the short term but no better than placebo in restoring rhythm. Beta-blockers and calciumchannel antagonists benefited rate control but verapamil was much less effective than amiodarone at restoring cardiac rhythm.[11]

Atrial flutter

The ECG of atrial flutter has a regular saw-tooth baseline ventricular rate of 150 with narrow QRS complexes. This is a 2:1 AV block. It is often misdiagnosed as SVT. Rarely, conduction occurs 1:1, giving a ventricular rate of 300/minute.

Treatment for atrial fibrillation/flutter[8, 9]

The treatment needs to control the arrhythmia either as the rate or the rhythm and also from the viewpoint of prophylaxis against thromboembolic complications. A recent study (AFFIRM)12 confirms that patients fare better with just rate control rather than trying to get them back into sinus rhythm if they are asymptomatic in atrial fibrillation.

Rate control

For rapid urgent control of ventricular rate:

- digoxin
  0.5 - 1.0 mg (o) immediately then 0.25 - 0.5 mg (o) every 4 - 6 hours to maximum of 1.5 - 2.0 mg in first 24 hours
  or
- verapamil
  1 mg/min IV up to maximum 15 mg (provided no evidence of heart failure and well-monitored BP)

Routine control:

- digoxin 0.0625 - 0.25 mg (o) daily according to age, plasma creatinine and digoxin level

Maintenance:

- digoxin (as above)
  ±
- verapamil SR 160 - 480 mg (o) daily *or*
  diltiazem CR 180 - 360 mg (o) daily *or*
- atenolol or metoprolol 25 - 100 mg (o) bd

Rhythm control

This should be considered if the patient is symptomatic and the arrhythmia is of recent onset - less than 6 months.

Medical cardioversion

Sotalol or amiodarone or flecainide

If the rate cannot be well controlled despite maximal medical therapy, consider AV node ablation and a permanent pacemaker. Atrial fibrillation with a rapid ventricular response over a long period gradually causes left ventricular dysfunction.

Electrical DC cardioversion

This can be given for first-line treatment or for failed medical conversion.

The use of warfarin in atrial fibrillation

Warfarin is effective in preventing stroke in patients with lone or non-rheumatic atrial fibrillation. The decision to use it or an antiplatelet agent, especially in the younger patient, is difficult and should be made in consultation with a cardiologist. As a general rule all patients should start on warfarin unless < 65 years old or have a major contraindication to its use. If using warfarin, start with a low dose (e.g. 2 - 4 mg) and maintain a relatively low INR of 2 - 3 with regular checks.

ADVANCES IN TREATMENT OF ARRHYTHMIAS

Apart from special rate-responsive pacemakers for bradycardia, there are several new modalities of treatment for complex arrhythmias, including means of blocking the re-entry phenomenon.

Surgery

Guided by electrophysiologic monitoring, surgeons can dissect a small section of the AV ring to ensure that all aberrant connections between the atria and the underlying ventricular muscle are severed.

Catheter electrode ablation

Specific abnormal foci in the conducting pathways can be ablated using direct current electrical surgery or radiofrequency ‘burns’ via a catheter electrode. Radiofrequency ablation, which will probably supplant surgery as a form of treatment, is indicated for recurrent episodes of supraventricular tachycardia, accessory pathways and nodal re-entry tachycardia with success rates of up to 95%. Catheter ablation for atrial fibrillation is more complex and still evolving. Success rates at best are 60 - 80%.

Automatic implantable cardiac defibrillator (AICD)

This expensive implant is the most effective therapy yet devised for the prevention of sudden cardiac death in patients with documented sustained ventricular tachycardia or fibrillation. Operative mortality should be less than 10%, after which survival at 1 year is over 90%. These new defibrillators incorporate an antitachycardia pacemaker. Patients can either be paced out of arrhythmia or, if they develop ventricular fibrillation, they can be defibrillated using higher energy.

Practice tips

- Atrial fibrillation and dizziness (even syncope) are suggestive of SSS (bradycardia - tachycardia syndrome), which is made worse by digoxin.
- Consider thyrotoxicosis as a cause of atrial fibrillation or sinus tachycardia even if clinical manifestations are not apparent.
- Check for a history of palpitations in a patient complaining of dizziness or syncope (and vice versa). Consider an arrhythmia, especially in the elderly.
- PSVT is rarely caused by organic heart disease in young patients.
- Arrhythmia of sudden onset suggests PSVT, atrial flutter/fibrillation or ventricular tachycardia.
- A normal ECG in sinus rhythm does not exclude an accessory pathway.
- Consider conduction disorders such as the WPW syndrome in PSVT. Avoid digoxin in WPW syndrome.
- Common triggers of premature beats and PSVT are smoking, anxiety and caffeine (especially 8 or more cups a day).
- Many antiarrhythmic drugs have proarrhythmic potential:
  - never use digoxin in WPW syndrome and SSS (without pacemaker back-up)
  - never use digoxin or verapamil for atrial fibrillation in WPW syndrome
  - beware of proarrhythmic side-effects of class I and class III antiarrhythmics
  - beware of using verapamil or diltiazem with a beta-blocker
- There is no ideal antiarrhythmic agent for ventricular premature beats.

Table 70.5 presents summary of the treatment of arrhythmias.

Table 70.5 Summary of treatment of arrhythmias[7, 8]

Arrhythmia	First line	Second line	Third line
Sinus tachycardia	Treat cause Reduce caffeine intake	Metoprolol or atenolol or Verapamil (rarely indicated)	
Bradycardias			
Sick sinus syndrome	Permanent pacing if symptomatic		
AV block			
first degree	No treatment		
second degree			
• Mobitz I	No treatment	Pacing if problematic	
• Mobitz II	Consider pacing		
third degree			
• acute (e.g. MI)	Temporary pacing or Isoprenaline IV		
• chronic	Permanent pacing		
Atrial tachyarrhythmias			
PSVT	Valsalva Carotid sinus massage	Adenosine IV or Verapamil IV	DC cardioversion Class III drug

			? Ablation
Atrial fibrillation Rate control Rhythm control	Digoxin	Beta-blocker or diltiazem or verapamil	AV node ablation + permanent pacemaker
Atrial flutter Rate control Rhythm control	Consider if symptomatic and recent onset < 6 months	Cardioversion → *or*	Electrical or chemical
		Maintenance of sinus rhythm →	Sotalol, flecainide, amiodarone
Atrial premature complexes	Treat cause Check lifestyle	Metoprolol or atenolol or Verapamil	
Ventricular tachyarrhythmias			
Ventricular premature beats	Treat cause Check lifestyle	Beta-blocker (especially mitral valve prolapse)	Class I or III drugs (rarely needed)
Ventricular tachycardia			
• non-sustained	Lignocaine IV	amiodarone	Class III drug
• sustained	Lignocaine IV if stable - if not: DC shock		DC cardioversion
Ventricular fibrillation	DC cardioversion	IV adrenaline if fine VF then DC cardioversion	amiodarone (maintenance) Class III (if recurrent)

WHEN TO REFER[13]

Patients should be referred to a cardiologist[12] when:

- a sustained supraventricular tachycardia is suspected
- a sustained ventricular tachycardia is suspected
- an ECG shows sustained delta waves of WPW syndrome, even if asymptomatic
- syncope or dizziness suggests a cardiovascular cause
- a paroxysmal arrhythmia may be the cause of unexplained cardiovascular symptoms
- anticoagulation has to be considered

REFERENCES

1. Boxall J. Annual update course for general practitioners. Course abstracts. Melbourne: Monash University, 2002: 16.
2. Lewit K. *Manipulative Therapy in Rehabilitation of the Motor System*. London: Butterworths, 1985: 338 - 9.
3. Davis A, Bolin T, Ham J. *Symptom Analysis and Physical Diagnosis* (2nd edn). Sydney: Pergamon, 1985.
4. Sandler G, Fry J. *Early Clinical Diagnosis*. Lancaster: MTP Press, 1986: 327 - 59.
5. Robinson MJ, Roberton DM. *Practical paediatrics* (5th edn.) Melbourne: Churchill Livingstone, 2003: 501 - 2.
6. Merriman A. *Handbook of International Geriatric Medicine*. Singapore: PG Publishing, 1989: 99 - 100.
7. O'Connor S, Baker T. *Practical Cardiology*. Sydney: MacLennan & Petty, 1999.
8. Smith A (Chair). *Therapeutic Guidelines: Cardiovascular* (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2003: 121 - 39.
9. Vohra J. Management of Palpitations. Aust Doc, 2001; 19 January: 111.
10. ACC/AHA/ESC. Guidelines for the management of patients with atrial fibrillation. 2001.

< http://www.acc.org >.

11. Barton S ed. *Clinical Evidence*. London: BMJ Publishing Group, 2001: 1 - 6.
12. Wyse DG et al. The Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) Investigators. NEJM, 2002; 347(23): 1825 - 33.
13. Ross DL. Cardiac arrhythmias. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 93 - 6.

CHAPTER 71. SLEEP DISORDERS

Sleep … is the first great natural resource to be exhausted by modern man. The erosion of the nerves, not to be halted by any reclamation project, public or private.

IRWIN SHAW 1949, 'THE CLIMATE OF INSOMNIA', *The New Yorker*

SLEEP DISORDERS

Sleep is one of the five great innate drives in humans. Disorder of this basic function is one of the most common health-related problems presenting to the GP. It may represent the clue to some very important disorders, such as depression, anxiety, adverse drug reactions, drug abuse and obstructive sleep apnoea (OSA). About half of the population report having some sleep-related problem in a year, with 25% of the Australian population reporting trouble getting enough sleep.[1] Normal sleep re-quirement varies considerably.

Disorders of the sleep - wake cycle, which are invariably caused by a disruption of the body's endogenous time clock, can result in insomnia or hypersomnolence (excessive sleepiness) or a combination of both. This is a feature of people experiencing the jet lag of international travel and shift workers.

KEY FACTS AND CHECKPOINTS

- Normal sleep: in a fit young person the ideal is 7.5 - 8 hours; latency < 30 minutes; wakefulness within sleep usually < 5% of time.
- Humans can stay awake without a problem for 16 - 18 hours. Sleepiness is wake-state instability.
- The evaluation of sleep disorders involving the sleep - wake cycle is enhanced by the patient keeping a sleep chart.
- It is important to take a drug history from patients complaining of insomnia or hypersomnolence.
- Drugs that can disturb sleep include alcohol, nicotine, antihistamines, selective serotonin reuptake inhibitors (SSRIs), caffeine, hypnotics, venlafaxine, selected β-blockers, β_2-agonists, theophylline, corticosteroids, sympathomimetic agents.
- Sleep disorders in children, including snoring, should be taken seriously and investigated. They have many consequences, such as learning difficulties, hyperactivity, behavioural disorders, failure to thrive and short stature.
- Be wary of young people and others presenting with insomnia with some urgency, especially requesting temazepam capsules - they may be dependent on benzodiazepines.
- People with OSA usually present with the TATT syndrome - 'tired all the time'. These patients are often unaware of waking or becoming aroused during the night.
- A patient who snores, has witnessed apnoeas and sleepiness is likely to have OSA.
- The majority of cases of excessive somnolence are caused by OSA and narcolepsy.[2]
- Non-pharmacological therapies, which include basic education and practice of sleep hygiene and behavioural therapy, should be used in management wherever possible.
- Referral to the new generation, specialist sleep disorder centres provides enhanced objective evaluation, diagnosis and treatment of the more complex disorders.
- It is illegal for a driver with a commercial driver's licence to continue to drive while suffering from untreated OSA.

SLEEP-RELATED DISORDERS

Disorders of sleep are a common and significant contribution to community illness and death. For example, it is known that untreated moderate to severe OSA has an 11 - 13% 5-year mortality and a 37% 8-year mortality, mainly from cardiovascular and motor vehicle accident related deaths.[3, 4] A classification of sleep disorders is presented in Table 71.1.[5]

Table 71.1 Classification of sleep disorders[5] (modified DSM - IV - TR, with key examples)

Dyssomnias

Primary insomnia

Other disorders initiating or maintaining sleep

- Periodic limb movements (nocturnal myoclonus)
- Restless legs syndrome

Excessive somnolence

- Primary hypersomnia
- Narcolepsy

Breathing-related sleep disorder

- Obstructive sleep apnoea
- Central sleep apnoea
- Central alveolar hypoventilation syndrome

Circadian rhythm sleep disorder

- Jet lag type
- Shift work type
- Delayed sleep phase type

Parasomnias

- Nightmare (dream anxiety) disorder
- Sleep terror disorder
- Sleepwalking disorder

Secondary sleep disorders

- Medical condition disorder
- Mental disorder
- Substance abuse

Many conditions may disturb breathing during the night (Figure 71.1). Nocturnal dyspnoea may result from cardiac causes (mitral stenosis, ischaemic cardiomyopathy, cardiac arrhythmias, fluid overload or retention), which usually present with orthopnoea, pulmonary crepitations and peripheral oedema. Asthma is another common cause of nocturnal dyspnoea, cough (with or without wheeze) occurring classically between 2 am and 5 am. Gastro-oesophageal reflux with or without aspiration may disturb respiration at night, but it usually presents with daytime or postural reflux. All these conditions can usually be differentiated from sleep apnoea clinically or with further investigation.


mental disorder
• major depression
• anxiety
drugs (e.g. hypnotics)
neurological disorders
TMJ dysfunction
thyroid disorders
(e.g. goitre, hypothyroidism)
asthma
sleep apnoea
neuromuscular disorder
(e.g. motor neurone disease)
nasal obstruction
tonsillar-adenoidal hypertrophy
macroglossia
micrognathia
pharyngeal tumours
gastro-oesophagael
cardiac disorders
genitourinary disorders
Others
• obesity
• alcohol excess
nocturnal myoclonus
restless legs

Figure 71.1 Significant causes of sleep disturbance

The sleep apnoea syndromes are a common group of disorders that result in periodic hypoventilation during sleep. They occur in about 2% of the general population in all age groups, and in about 10% of middle-aged men.

INSOMNIA

Insomnia, also referred to as DIMS (disorders of initiating and maintaining sleep), is defined as the inability to function well with the amount and quality of sleep experienced. Insomniac patients may complain of difficulty getting to sleep or staying asleep, of frequent intermittent nocturnal arousals, early morning awakening or combinations of these.

A careful history is required because some patients have unrealistic expectations about the required amount of sleep they need or have misperceptions of how long they have slept.

When taking the history, explore lifestyle factors, especially psychological reasons, painful conditions, drug use and abuse, appetite, energy, sexual issues and physical factors. Check thyroid status, especially hyperthyroidism.

Management[6]

1. Contract with the patient: discuss and agree to the therapeutic objective with the patient (e.g. to reinstate sleep without medication).
2. Sleep - wake history: take a sleep - wake history (preferably with a sleep diary) and evaluate daytime functioning.
3. Exclude and treat any underlying problem, for example:
    - drugs (e.g. caffeine, alcohol, β-blockers, nicotine)
    - anxiety, stress
    - depression
    - restless legs syndrome
    - sleep apnoea
    - parasomnias - nightmares, sleepwalking

   - physical disorders (e.g. CCF, arthritis, asthma)
   - bed-wetting
   - reflux disease
   - thyroid disorders (e.g. thyrotoxicosis)
   - menopausal symptoms
   - snoring partner
   - lower urinary tract symptoms with nocturia

4. Explanation and reassurance, including patient education handout.
5. Sleep hygiene advice.
   - Try to recognise what helps patient to settle best (e.g. warm bath, listening to music).
   - Establish a routine before going to bed.
   - Avoid daytime naps.
   - Regular daytime exercise.
   - Regular time of arising.
   - Avoid strenuous exercise close to bedtime.
   - Avoid alcohol and drinks containing caffeine in evening, especially close to retiring.
   - Avoid a heavy evening meal.
   - Avoid pets in the room.
   - Avoid smokings.
   - Try a drink of warm milk before retiring.
   - Organise a comfortable quiet sleep setting with the right temperature.
   - Remove pets from the bedroom.
   - Avoid lights, including poorly screened windows and highly illuminated clocks in the room.
   - Sex as the last thing at night where appropriate is usually helpful.

6. Non-pharmacological treatment.
   There are various techniques that should be adopted according to patient personality and preference, clinician's expertise and available resources. These include relaxation therapy, meditation and stress management, which are all highly recommended.
   Other measures include cognitive-behaviour therapy, structured problem solving and electromyographic feedback. Hypnosis is worth considering.
7. Pharmacological treatment.[6]
   It is advisable to avoid hypnotic agents as first-line treatment. If any form of continuous agent is necessary it is best to limit it to 2 weeks.
   Options:
   - temazepam 10 - 20 mg tablets (o) nocte
   - zopiclone 3.75 - 7.5 mg (o) nocte, or
   - zolpidem 10 mg (o) nocte

   Note:
   - The cyclopyrrlone derivative zopiclone and the imidazopyridine derivative, zolpidem, are non-benzodiazepine hypnotics with a similar action to the benzodiazepines represented by temazepam. However, warnings have been issued about adverse neurological and psychiatric reactions.[4]
   - Tricyclic antidepressants with sedative effects (e.g. amitriptyline) are often used as hypnotics but should generally be avoided in the absence of depressive disorders.

SLEEP APNOEA[7]

The term ‘sleep apnoea’ is used to describe cyclical brief interruptions of ventilation, each cycle lasting 15 - 90 seconds and resulting in hypoxaemia, hypercapnia and respiratory acidosis, terminating in an arousal from sleep (often not recognised by the patient). The interruption is then followed by the resumption of normal ventilation, a return to sleep, and further interruption of ventilation.

Sleep apnoea is broadly classified into obstructive and central types.

Obstructive sleep apnoea (OSA) is the commonest type and involves an intermittent narrowing or occlusion of the pharyngeal area of the upper airway (Figures 71.2 and 71.3). The effects include snoring and hypopnoea, sometimes apnoea.

Predisposing causes include:

- diminished airway size (e.g. macroglossia obesity, tonsillar-adenoidal hypertrophy)
- upper airway muscle hypotonia (e.g. alcohol hypnotics, neurological disorders)
- nasal obstruction



Figure 71.2 Normal airway when sleeping



Figure 71.3 Sleep apnoea: obstructed airway when sleeping

Central sleep apnoea (CSA) is less common and is due mainly to neurological conditions such as brain stem disorders leading to reduced ventilatory drive, and neuromuscular disorders such as motor neurone disease.

Clinical effects of sleep apnoea syndromes[5]

Important clinical presentations include:

- excessive daytime sleepiness and tiredness
- nocturnal problems (e.g. loud snoring, thrashing, ‘seizures’, choking, pain reactions)

- morning headache
- subtle neuropsychiatric disturbance - learning difficulties, loss of concentration, personality change, depression
- sexual dysfunction

Causes of excessive daytime sleepiness are presented in Table 71.2. In OSA, sleepiness results from repeated arousals during sleep and the effects of hypoxaemia and hypercapnia on the brain. Physical examination may reveal few or no signs.

Table 71.2 Causes of excessive somnolence

Inadequate sleep duration
Sleep apnoea syndromes
Narcolepsy
Endocrine (e.g. hypothyroidism)
Drug induced
Purposeful sleep deprivation
Nocturnal myoclonus
Bereavement
Idiopathic

Management of sleep apnoea[7]

Referral to a comprehensive sleep disorder centre is advisable if this disorder is diagnosed or suspected.

The general principles are as follows:

1. Lifestyle modification.
    - Weight loss (e.g. loss of 7 - 10 kg can significantly reduce severity).
    - Achieve physical fitness with regular exercise.
    - Good sleep hygiene and adequate sleep hours.
    - Reduce or cease sedatives/hypnotics.
    - Avoid alcohol for up to 3 hours before going to sleep.
    - Cease cigarette smoking (increases nasal resistance).
    - Medical management of nasal obstruction (e.g. short-course nasal decongestants).
    - Avoid supine sleep.
2. Continuous positive airway pressure (CPAP).
    - CPAP is currently the most effective treatment for OSA (consider it for CSA).
    - Delivered by nasal (or facial) mask.
    - Provides an air splint to the upper airway and prevents pharyngeal collapse.
    - Sleepiness and neurocognitive function improved.
    - Not tolerated by all patients.
3. Surgery. In children, OSA is usually due to tonsillar and/or adenoid hypertrophy and is relieved by surgery (Figure 71.4). In adults, depending on the cause, the options are:
    - correction of the specific upper airway anatomical problem - -up to 2% of problems
    - correction of nasal obstruction (improves snoring and OSA)
    - palatal surgery - uvulopalatopharyngealplasty - conventional or laser assisted
    - tonsillectomy

  - base of tongue surgery
  - radio-frequency treatment to soft palate and base of tongue (‘somnoplasty’)

4. Oral appliances.
  - The mandibular advancement splint - a custom-made dental device that supports the mandible and tongue during sleep to increase pharyngeal dimensions.

5. Medication.
  - Amitriptyline 25 - 100 mg (o) nocte, in severe cases during REM sleep and intolerance of CPAP.



Figure 71.4 Influence of tonsils and adenoids on airways

NARCOLEPSY

Narcolepsy is a specific, permanent neurological disorder that is characterised by brief spells of irresistible sleep during daytime hours, usually at times when the average person simply feels sleepy. It is uncommon with an incidence of two to five per 100 000.

Features

Usual onset between adolescence and 30 years of age - in teens and twenties (has been reported in children as young as 2 years).

Tetrad of symptoms

- Sleepiness: sudden brief sleep attacks (15 - 20 minutes).
- Cataplexy: a sudden decrease or loss of muscle tone in the lower limbs that may cause the person to slump to the floor, unable to move. These attacks are usually triggered by sudden surprise or emotional upset.
- Sleep paralysis: a frightening feeling of inability to move while drowsy (between sleep and waking).
- Hypnagogic (terrifying) hallucinations on falling asleep or waking up.

Several attacks per day are possible.

Diagnosis

The diagnosis is clinical through the taking of an appropriate history. If doubtful, include:

- EEG monitoring
- sleep laboratory studies (sleep latency test - rapid eye movement is a hallmark)

Treatment[6]

Treatment is mainly symptomatic and initiated by a consultant. Central nervous system psychostimulants are of proven effectiveness in increasing alertness:

- dexamphetamine 5 - 10 mg (o), half an hour before breakfast and lunchtime; up to 40 mg daily may be required in slowly increasing doses, or
- methylphenidate (Ritalin) 10 - 20 mg (o) half an hour before breakfast and lunchtime; up to 60 mg daily may be

required

Drug holidays from these drugs may be necessary.

- Tricyclic antidepressants are used to treat cataplexy, sleep paralysis and hypnagogic hallucinations (e.g. clomipramine 20 - 100 mg (o) daily).
- Modafinil is used successfully in some countries.

SNORING

Definition

Snoring is a sonorous sound with breathing during sleep, caused by vibrations in the upper airways from the nose to the back of the throat. It is caused by partially obstructed breathing during sleep (Figure 71.5).



Figure 71.5 Snoring: vibrations of tongue and soft palate

Features

- Sometimes indicates OSA, especially in perimenopausal women.[7]
- Three times more common in obese persons.
- Generally harmless, but if very severe, unusual or associated with periods of no breathing (> 10 s) assessment is advisable.

Aggravating factors

- Obesity
- Old age
- Sleeping on the back
- Excess alcohol
- Neck problems, especially a ‘thick’, inflexible neck
- Various drugs, especially sedatives and sleeping pills
- Hay fever and other causes of nasal congestion
- Problems in the upper airways, such as nasal polyps, enlarged tonsils or a foreign body (e.g. a piece of plastic or metal)

Management

If an examination rules out a physical problem causing obstruction in the back of the nose and OSA, then the following simple advice can be given to patients.

- Obtain and keep to ideal weight.
- Avoid drugs (including sedatives and sleeping tablets), alcohol in excess and smoking.

- Treat nasal congestion (including hay fever) but avoid the overuse of nasal decongestants.
- For neck problems, keep the neck extended at night by wearing a soft collar.
- Consider a trial of an intranasal device such as the Breathing Wonder, which is a hollow intranasal plastic insert. Pharmacists can advise about the range of such devices.
- Try to sleep on your side. If you tend to roll on to your back at night, a maverick method is to consider sewing ping pong balls or tennis balls on the back of the nightwear. Others wear a bra (with tennis balls) back to front.

PERIODIC LIMB MOVEMENTS (NOCTURNAL MYOCLONUS)

Periodic limb movements (PLMs) and restless legs syndrome are important causes of insomnia and excessive daytime sleepiness. They may coexist in the same patient. Periodic limb movements, which are also referred to as nocturnal myoclonus or ‘leg jerks’, tend to occur usually in the anterior tibialis muscles of the leg but can occur in the upper limbs. The prevalence increases with age. Most people with PLMs are completely asymptomatic. The diagnosis is often made during sleep studies.[8] If troublesome, referral to a sleep clinic or neurologist may be appropriate.[8]

Medication

Medication that may help includes:

- levodopa plus carbidopa (e.g. Sinemet 100/25, 2 tablets before bedtime) or
- clonazepam 1 mg (o) nocte increasing to 3 mg (o) nocte or
- sodium valproate 100 mg (o) nocte

RESTLESS LEGS SYNDROME (RLS)

RLS, also known as Ekbom's syndrome, is a rather common movement disorder of the nervous system where the legs feel as though they want to exercise or move when the body is trying to rest. The major complaint of sufferers is of disruption both to sleep and of relaxing activities, such as watching television or reading a book. Prolonged car or airplane travel can be difficult.

RLS is frequently an undiagnosed disorder because people often don't complain about it to their doctor. A Canadian study reported that 15% of people sampled reported ‘leg restlessness’ at bedtime.

The diagnosis is made from the history - there are no special diagnostic tests.

Its prevalence increases with age so it mainly affects elderly people. Women are more prone to get RLS and it is aggravated by pregnancy. The exact cause of primary RLS is not clear. It is not related to exercise and does not appear to follow strenuous exercise.

Symptoms

There is an urge to move legs upon resting, particularly after retiring to bed. This urge is a response to unpleasant sensations in the legs, especially in the calves. The sensations are commonly and variously described as crawling, creeping, prickly, tingling, itching, con-tractions, burning, pulling or tugging, electric shock-like. However, sometimes patients are unable to describe the sensation or refer to it as simply a compulsion to move the legs.

In some patients the arms are affected in a similar way. The symptoms seem to be aggravated by warmth or heat. Many patients with RLS also experience nocturnal myoclonus.

Secondary (medical) causes include:

- anaemia
- iron deficiency
- uraemia

- hypothyroidism
- pregnancy (usually ceases within weeks of delivery)
- drugs (e.g. antihistamines, antiemetics, selective antidepressants, lithium, selective major tranquillisers and antihypertensives)

Management

Iron studies should be performed and, if low, treat with iron and vitamin C tablets. Advise patients that although RLS can come and go for years it usually responds well to treatment.

Self-help advice

- Perform activities that can reduce symptoms, for example, a modest amount of walking before bedtime, massage or prescribed exercises (see Figure 71.6). Note: Getting out of bed and going for a walk or run does not seem to help RLS.



Figure 71.6 Stretching exercises for restless legs

- Good sleep hygiene, namely regular sleeping hours, gradual relaxation at bedtime, avoidance of non-sleep activities in bed (e.g. reading, eating).
- Diet: follow a very healthy diet. Avoid caffeine drinks, smoking and alcohol.
- Try keeping the legs cooler than the body for sleeping.
- Exercises: a popular treatment is gentle stretching of the legs, particularly of the hamstring and calf muscles for at least 5 minutes before retiring. This can be done by using a wide crepe bandage, scarf or other length of material around the foot to stretch and then relax the legs (as shown in Figure 71.6).

Pharmacological treatment

The following may be effective if simple measures fail:

- paracetamol 1000 mg (o) nocte or
- clonazepam 1 mg (o) 1 hour before retiring or
- paracetamol 1000 mg (o) plus clonazepam 1 mg or diazepam 5 mg

Levodopa, gabapentin, codeine, baclofen and propranolol may be helpful. Carbamazepine, quinine, antipsychotics (avoid), antihistamines (avoid) and antidepressants are generally unhelpful.

BRUXISM (TEETH GRINDING)

Bruxism is the habit of grinding, clenching or tapping teeth, which may occur while awake (especially in children) or more commonly while asleep. The usual symptom is annoying, teeth-grinding noises during sleep that disturb family members. It may result in headaches and TMJ dysfunction in the person during the day. The cause may be a habit or a response to subconsciously correct a faulty bite by making contact between the upper and lower teeth when the jaws are closed. It is aggravated by stress and is more common in heavy alcohol drinkers.

Tips on management

- Educate the patient to recognise, understand and try to overcome the habit.
- Practise keeping the jaws (and teeth) apart.
- Slowly munch an apple before retiring.
- Practice relaxation techniques, including meditation, before retiring.
- Consider other stress management techniques (e.g. counselling, relaxation exercises, yoga and tai chi).
- Place a hot towel against the sides of the face before retiring to achieve relaxation.
- If this fails and bruxism is socially unacceptable, a plastic night-guard mouthpiece can be fashioned by a dentist to wear at night.

PARAINSOMNIAS

Parainsomnias are defined as dysfunctional episodes associated with sleep, sleep stages or partial arousal. They are more common in children.

Nightmares (dream anxiety)

These usually occur later in the sleep period and are accompanied by unconscious body movements, which usually cause the person to awaken.

Associations include traumatic stress disorders, drug withdrawal (e.g. alcohol, barbiturates, drugs such as SSRIs, β-blockers, benzodiazepines). Violent behaviour can occur during these dreams due to a REM behaviour disorder and this requires a sleep study.

Psychological evaluation with cognitive behaviour therapy (CBT) is appropriate. Medication that may help includes phenytoin, clonazepam or diazepam.

Somnambulism (sleepwalking)[2]

This is a complex motor activity in which the person performs some repetitive activity in bed or walks around freely. No treatment is usually required but, if it is repetitive and potentially dangerous, then the sleeping environment should be rendered safe. Psychological assistance is required for recurring episodes. Benzodiazepines such as diazepam may be useful but withdrawal usually leads to rebound problems.

Night terrors

These are part of the same sleep cycle disorder as somnambulism. Characteristics of night terrors are sharp screams, violent thrashing movements and autonomic overactivity, including sweating and tachycardia. The sufferers may or may not awake and usually cannot recall the event. They also require psychological evaluation and therapy. Similar medication as used for nightmares may help (e.g. a 6-week trial of phenytoin, diazepam or clonazepam).

SLEEP DISORDERS IN CHILDREN

Sleep disorders in children are very common in late infancy, toddlerhood and early preschool age groups. The majority do not sleep throughout the night until 6 months of age. Over 50% of toddlers and preschool children resist going to bed.[9] At least 30% of infants and toddlers wake at least once during the night every night. Toddlers begin to have dreams coinciding with language development in the second year of life.[9]

The child who wakes during the night needs reassurance, protection and the parent's presence, but it must be given discreetly. Although psychosocial stresses can trigger sleep problems, serious psychological problems in children with sleep disorders is uncommon.[10]

Management

Aim tjo see both parents together and get them to agree on the approach, including sharing the workload.

Advice to parents:

- Resist taking the child into bed during the night unless they are happy to encourage this.
- Avoid giving attention to the child in the middle of the night - it encourages attention seeking.
- Avoid extra feeding or other pacifiers during the night.
- Return the child to bed promptly and spend only a brief time to give reassurance.
- A rigid series of rituals performed before bedtime helps the child to develop a routine. Settling to sleep may be assisted by soft music, a soft toy and a gentle night light.
- Take the child into the bedroom while still awake.
- Encourage the parents to keep a sleep dary.

Medication has a minimal place in the management of sleep disturbances, although the judicious use of a sedative/hypnotic for a short term may break the sleepless cycle. Such drugs include chloral hydrate 25 - 50 mg/kg per dose and trimeprazine (Vallergan) 1 - 2 mg/kg per dose (not for infants under 6 months).[11]

Para-insomnias (night terrors, sleepwalking and -talking)[2]

These are not true sleep disorders or night-time arousals. They occur in deep won-REM sleep. With night terrors, which usually develop within 2 hours of sleep and last 1 - 2 minutes, the child is usually inconsolable and has no memory of the event. These events cluster in two age ranges, 2 - 4 years and 6 - 9 years, and are self-limiting over a period of months. Usually, no active treatment is needed but for persistent severe problems a 6-week trial of phenytoin, diazepain or imipramine is worthwhile.

SLEEP PROBLEMS IN THE ELDERLY

The elderly constitute the bulk of long-term users of hypnotics and benzodiazepines. Two key issues to consider are sleep in the elderly and confusion in the elderly. Problems associated with long-term benzodiazepine use are dependence, confusion, memory impairment and falls.

A recent study of elderly patients with insomnia showed that:[12]

- 25% had insomnia either coexisting with or related to other sleep disorders, such as sleep apnoea or periodic limb movement disorder
- 10% had insomnia related to medical or psychiatric conditions
- 13% had insomnia associated with an inability to stop taking sedative - hypnotic agents

Principles of management in the elderly

- Exclude underlying causes of sleep disturbance.
- Educate patients and carers about the changing needs with ageing and the rational use of medicines.
- Avoid hypnotics if possible.
- Avoid hypnotics combined with alcohol.
- Beware of risks of long-term use and drug accumulation.
- Consider non-drug measures where possible (e.g. CBT).
- Avoid the ‘why bother’ factor when carers and patients are comfortable with hypnotics.
- In nursing homes, practise ‘bothering’ and use team effort to encourage less prescribing.

Medication[6]

If medication is necessary, prescribe a short-acting benzodiazepine for as limited a time as possible. An alternative non-benzodiazepine hypnotic (e.g. zopiclone or zolpidem) taken just before retiring may be useful in the elderly (beware of adverse effects). Tricyclic anti-depressants with a sedative action are frequently used, especially if there is coexistent depression, but side-effects limit their use. However, commonsense prescribing needs to be used and the best option may be to continue the long-term prescribing of hypnotics in those with chronic medical problems or long-term dependents on a low effective dose.

REFERENCES

1. Wilson CW, Lack L. Sleeping habits of people living in the Adelaide metropolitan area - a telephone survey. Aust Psychol, 1983; 18: 368 - 76.
2. Tiller JWG, Rees VW. Sleep disorders. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 475 - 8.
3. Partinen M, Guilleminault C. Daytime sleepiness and vascular morbidity: a seven year follow-up in obstructive sleep apnoea patients. Chest, 1990; 97: 27 - 32.
4. Prescribing benzodiazepines - ongoing dilemma for the GP. NPS News, 2002; 24: 1 - 2.
5. Pierce R, Naughton M. Sleep-related breathing disorders: recent advances. Aust Fam Physician, 1992; 21: 397 - 405.
6. Dowden J (Chair). Therapeutic Guidelines: Psycho-tropic (Version 5). Melbourne: Therapeutic Guidelines Ltd, 2005: 163 - 71.
7. Dowden J (Chair) Therapeutic guidelines: Respiratory (Version 5). Melbourne: Therapeutic Guidelines Ltd, 2005.
8. Laks L. Sleep disorders. CHECK, 2000; issue 345: 3 - 10.
9. Jarman R. Sleep problems. Aust Paed Rev, 1992; 2(2): 1 - 2.
10. Smart J. Paediatric Handbook (6th edn). Melbourne: Blackwell Science, 2000: 145 - 7.
11. Ramchandni P, Wiggsh et al. A systemic review of treatments for settling problems and night waking young-Children. BMJ, 2000;320:209 - 13
12. Morin CM et al. Behavioral and pharmacological therapies for late-life insomnia: a randomised controlled trial. JAMA, 1999; 281 (11): 991 - 9.

CHAPTER 72. SORE MOUTH AND TONGUE

The key to managing diseases of the oral mucosa is to evaluate lifestyle factors carefully, including causes of immune suppression.

DR JONATHAN TVERSKY 2002

SORE MOUTH AND TONGUE

Evaluating the sore mouth and tongue is basically an understanding ofdisorders of the oral mucosa. Disorders of the oral mucosa are a common problem in general practice, with recurrent aphthous ulceration being the most common oral mucosal disease in humans.

There are three main types of epithelium in the oral mucosa:[1]

1. masticatory epithelium - orthokeratinised stratified squamous attached to underlying periosteum (e.g. palate and gingivae)
2. lining epithelium - parakeratinised stratified squamous (e.g. lip and buccal mucosa, alveolar mucosa, floor of mouth, soft palate andtongue - lateral and undersurface)
3. specialised epithelium - orthokeratinised stratified squamous with taste buds and papillae - filiform, fungiform and circumvallate - on dorsum of tongue.

KEY FACTS AND CHECKPOINTS

- Dental trauma or neglect is an important cause of many oral mucosal disorders, such as ulceration, bleeding gums and hyperplasia.
- Non-healing oral ulcers warrant biopsy to exclude squamous cell carcinoma (SCC).
- If oral mucosal carcinoma is suspected, palpate the lesions to check for induration or a firm discrete edge and check regional nodes.
- Erythroplasia or leukoplakia persisting for 3 weeks after injury, e.g. sharp tooth or denture, should have an incisional biopsy.
- Any oral ulcer or soft-tissue lesion that persists 3 weeks after the apparent cause has been removed should be biopsied.
- Consider Epstein-Barr virus (EBV) infectionwith unusual faucial ulceration and petechial haemorrhages of the soft palate.
- Apthous ulcers are usually 3 - 5 mm indiameter - minor ones have an erythematous margin.
- Intraoral bony exostoses, other than palatal and mandibular tori, are often variations from normal or less commonly part of a syndrome, for example, Gardner's syndrome. No treatment is usually required.[2]
- Shared care of more complex lesions of the mouth and tongue with an oral or dental surgeon is best practice.

ORAL ULCERATION

The histology of oral ulceration is usually non-specific, with fibrinslough covering granulation tissue, and the aetiology is varied. The ulceration is a breach in the epithelial compartment with inflammatory cell infiltrate in the submucosa. The most common form is recurrent aphthous ulceration. Always inquire about a history of skin problems, medication, bowel function and psychological stress.

A list of causes is presented in the diagnostic strategy model (Table72.1). Depending on the clinical picture, investigations may include FBE, swabs, autoantibody screen, syphilis serology, blood sugar, vitamin B_{12} and folate levels, and biopsy.

Table 72.1 Mouth ulcers: diagnostic strategy model

Q. ***Probability diagnosis*** A. Recurrent aphthous ulceration Trauma Acute herpes gingivostomatitis Candidiasis
Q. ***Serious disorders not to be missed*** A Carcinoma: SCC Leukaemia Agranulocytosis HIV Syphilitic - chancre or gumma Tuberculosis
Q. ***Pitfalls (often missed)*** A. Aspirin burn Inflammatory bowel disease (e.g. Crohn's) Herpes zoster virus Glandular fever (EBV) Lichen planus Coxsackie virus: herpangine hand, foot and mouth Immunosuppression therapy Lupus erythematosus *Rarities*: Behçet's syndrome Pemphigoid and pemphigus Erythema multiforme Radiation mucositis
Q. ***Seven masquerades checklist*** A. Depression - Diabetes ✔ Candida Drugs ✔ Anaemia ✔ iron-deficiency Thyroid disease - Spinal dysfunction - UTI -
Q. ***Is the patient trying to tell me something?*** A. Unlikely

Recurrent aphthous ulceration

Aphthous ulcers are round/oval ulcers usually 3-5 mm in diameter withan erythematrus margin and sloughing base.

Aphthous ulcers occur in all ages on para-keratinised mucosa such as the buccal and labial mucosa and the floor of the mouth (not on orthokeratinised mucosa). The frequency of aphthous ulcers in the population varies from 5% to 25%

(average 20%). The cause isunknown although human herpes virus 6 has been implemented, as have nutritional and autoimmune factors.[3] People with recurrent aphthous ulcers have a genetic predisposition.

Precipitating factors

- Trauma (e.g. cheek and tongue biting, toothbrush, dental pressure)
- Drug reaction (e.g. new medication)
- Stress
- Allergy
- Systemic factors (e.g. iron, folate, vitamin B_{12} deficiency, hormonal)

*Note:*Exclude blood dyscrasias, Crohn's disease, Behcets disease, coeliac disease, drug therapy (e.g. phenytoin, cytotoxics, corticosteroids, immunosuppressants).

Rules

- Minor ulcer < 5 mm in diameter: lasts 5 - 10 days and heals without scarring.
- Major ulcer > 5 mm: can persist for up to 6 weeks.[3]
- Major ulcers usually occur on lips, soft palate and fauces and sometimes on the tongue.
- Minor ulcers are usually found on the buccal and labial mucosa and the floor of the mouth.
- Non-healing ulcers: consider SCC (biopsy required).
- Recurrent ulcers: consider Behçet's syndrome. Check serum iron and folate.

DxT: recurrent oral and genital ulcers + uveitis + arthritis ° Behçet's syndrome

Treatment

There are multiple methods but none are specific.

Symptomatic relief

- Apply topical lignocaine (e.g. 2% jelly or 5% ointment) with cotton bud: after 2 minutes apply lignocaine gel or paint (e.g. SM-33 adult paint formula or SM-33 gel children) every 3 hours, or
- Eutectic EMLA cream 5% - apply on a cotton bud or gauze for 5 minutes.

Healing: options

- Tetracycline/nystatin mouthwash[1] - highly recommended despite terrible taste (see formulation under practice tips).
- Triamcinolone 0.1% (Kenalog in Orobase) paste, apply 8 hourly and nocte (preferred method but be careful of herpes simplex ulcers).
- Other topical steroids (e.g. betamethasone 0.5%, hydrocortisone 1%).
- Hydrocortisone lozenges (if available) dissolve in contact with ulcer grid.
- 10% chloramphenicol in propylene glycol - apply with cotton bud for 1 minute (after drying the ulcer) 6 hourly for 3 - 4 days.
- Beclomethasone spray onto ulcer tds.
- Sucralfate 1 g in 20 - 30 mL of warm water - use this as a mouthwash.
- Diclofenac 3% in hyaluronan 2.5%.

All of the above treatments have been shown to be effective in controlled trials.

Major ulceration

Consider:

- injection of steroids into the base of the ulcer, and/or

- oral prednisolone, 5 - 7 days

Referral: patience with a non healing ulcer within 3 weeks of presentation.

Complementary measures

1. Teabag method. Consider applying a wet, squeezed-out, black teabag directly to the ulcer regularly (the tannic acid promotes healing). Must be used when ulcer is worse.
2. Melaleuca (tea-tree) oil. 1% tea-tree oil used as a mouth rinse for 1 minute has been shown to prevent secondary infection.[2, 4]
3. Acupuncture. This is advocated by its protagonists. It has been proven to improve salivary flow.

Traumatic ulceration

- This is commonly caused by sporting injuries, biting of the cheek, lip and hot food.
- Factitious causes include scratching of an ‘itchy’ mouth or over-brushing of the teeth.
- Other relevant causes include dentures, sharp tooth surface, orthodontic bands and sharp objects, such as pencils and hard food.
- Aspirin ‘burns’ are caused bypeople leaving salicylate-based tablets to dissolve against oral mucosa.
- Iatrogenic causes include surgical procedures, such as intubation and endoscopy, and dental treatment, such as retractors and removing drycotton rolls.

Management[1]

- Explanation, including removal of the cause.
- Warm salt-water mouthwashes, and/or local anaesthetic mouthwashes:
  - benzocaine compound (Cepacaine), swirl in mouth 10 - 15 mL for 10 - 15 seconds and expel, every 3 hours prn, or
  - benzydamine hydrochloride (Difflam), swirl in mouth 15 mL for 30 seconds and expel

These ulcers can take up to 10 days to resolve.

Lichenoid drug reaction

Several drugs can induce a lichenoid drug reaction of the oral mucosa, that is, cause shallow mucosal erosions similar to lichen planus. The drugs include gold, the NSAIDs, carbimazole, selected antihypertensives and cytotoxics.

Herpes infection

- It is important to beware of herpes simplex virus lesions.
- Primary herpetic gingivostomatitis is usually obvious but herpes infection has an extraordinary ability to present in many ways. It can spread from the hands to the mouth.
- Application of a topical corticosteroid, such as Kenalog in Orobase, can aggravate and spread the herpetic lesion.
- Treatment: acidovir or similar antiviral if seen early, eg. 48 hours from onset; fluids+ + +; analgesee mouth rinses, eg. Difflam; counter admission for IV acidovir and hydration.
- Lesions of herpes zoster virus affecting the maxillary division of the trigeminal nerve, for example, can involve the buccal mucosa in a unilateral pattern.

RED PATCHES

A reduction in the surface epithelial layer causes erythematous patches. Causes include trauma (e.g. cheek biting), infection (e.g. Candida albicans), geographic tongue, haematologic disorders, the dermatoses and neoplasia.

Neoplasia that can look red includes squamous cell carcinoma, Kaposi's sarcoma and erythroplakia. Erythroplakia is similar in significance to leukoplakia except for the erythematous feature. It is an important condition to recognise since about 70% of cases are either dysplastic or carcinoma.[5]

WHITE PATCHES

White patches occur where the epithelial compartment thickens. Causesinclude inflammation due to trauma or infection, especially *Candida*, dermatoses and neoplasia.

An interesting condition is hyperkeratotic burns on the dependent floor of the mouth, which appear white. Causes include tea-tree oil mouthwash and the sucking of aspirin.

Leukoplakia is any white lesion that cannot be removed by rubbing the mucosal surface (unlike oral candidiasis). About 5% of cases represent either dysplasia or early SCC.[5]Any persistent white patch should be biopsied.

Specific conditions causing red and/or white patches follow.

Oral candidiasis (thrush)

This is usually tender and looks like white or yellowish curd-like patches overlying erythematous mucosa. Unlike lichen planus or leukoplakia, they are usually readily rubbed off and hence only the underlying red patch may be seen.

Patients may also complain of a bad metallic taste or halitosis and dysphagia. They often complain of sensitivity to toothpaste or acidic substances in general.

Consider predisposing factors:

- immunodeficiency and cytotoxic therapy
- medication, especially broad-spectrum antibiotics and corticosteroids, including inhalers
- debility and anaemia (iron, folic acid, vitamin B_6 deficiency)
- diabetes mellitus and HIV infection

The carriage rate of *Candida albicans* in the oral cavity is 60 - 75%. The diagnosis is made clinically but a wet preparation using potassium hydroxide will reveal spores and perhaps mycelia.

Treatment

Attend to underlying cause. Consider multi-vitamin preparations.

Topical therapy

- nystatin suspension, rinse and swallow qid or
- miconazole oral gel (as directed by manufacturer) or
- amphotericin 10 mg or nystatin 100 000 U lozenges dissolved slowly in oral cavity, 6 hourly, for 7 - 14 days

Oral therapy

Use if unresponsive to tropical therapy and the immunocaupromised:

- fluconazole 50 mg (0) daily.

Denture therapy[1]

Dentures need to be decontaminated, especially if acrylic. Use:

- chlorhexidine denture scrub (care with bleaching), or
- dilute Milton's denture scrub (e.g. $^1/_4$ teaspoon White King in a cup of water)

If oral thrush, brush the dentures each night with a thin coat of nystatin cream or oral miconazole.

ANGULAR CHELITIS

Feature is redness, soreness and maceration of the corners of the mouth. Usually, associated with oral candidiases. Counder poor-fitting denfuses. Treat with topical nystatin or miconazole. ‘Golden’ crusting indicates S. *auteus*.

BLEEDING OR PAINFUL GUMS

Erythematous bleeding gums are a common worldwide problem, which is almost always a localised inflammation associated with poor dental hygiene.[6] Systemic problems usually as part of a bleeding diathesis need to be considered.

The causes are summarised in the diagnostic strategy model (Table 72.2).

Table 72.2 Bleeding/painful gums: diagnostic strategy model (modified)

Q. ***Probability diagnosis***
A. Gingivitis/periodontal (gum) disease
Trauma: poor fitting or partial dentures
Factitious: excessive brushing
Drugs: warfarin overdose
Q. ***Serious disorders not to be missed***
A. Oral carcinoma/benign neoplasms, eg. Epilus
Blood dyscrasias (e.g. AML)
Acute herpetic gingivostomatitis
Q. ***Pitfalls (often missed) but uncommon***
A. Acute necrotising ulceration gingivitis (Vincent's infection)
Autoimmune disease (e.g. lichen planus, SLE)
Hereditary haemorrhagic telangiectasia
Malabsorption
Scurvy

Acute necrotising ulcerative gingivitis (Vincent's infection or trench mouth) caused by anaerobic organisms is rarely seen but is more common in undernourished or ill young adults under stress. Treatment is with procaine penicillin IM daily for 3-5 days followed by oral phenoxymethyl penicillin and metronidazole.[7]

Gingivitis

Features

- Red, swollen gingivae adjacent to teeth (Figure 72.1)



Figure 72.1 Gingivitis, showing plaque and marginal gingivitis

- Bleeds with gentle probing
- Halitosis
- Usually no pain
- Dental plaque accumulation with calculus (tartar) secondary to poor oral hygiene

Periodontitis

This is inflammation of the periodontal space. It is a sequel to gingivitis and shows periodontal ligament breakdown with recession or periodontal pocketing and alveolar bone loss. There is possible loosening of teeth and periodontal abscess formation (Figure 72.2). An underlying medical condition must be suspected.



Figure 72.2 Periodontitis with more widespread gingival inflammation that invades the supporting alveolar bone

Treatment of the above conditions[6]

- Chlorhexidine 0.1 - 0.2% aqueous solution as a mouthwash bd for 10 days: this can be followed by a phenolic compound mouthwash long term.
- Systemic antibiotics (not topical) for periodontal abscess formation (e.g. amoxycillin 250 mg (o) tds for 5 days - drug of choice or phenoxymethyl penicillin).
- For severe or unresponsive cases add metronidazole.[7]

Prevention

- Use a fluoride, abrasive type of toothpaste.
- Brush with a medium-to-soft, nylon-tufted, small-headed toothbrush.
- Direct at gingival margin with a small horizontal motion.
- Keep interdental spaces clean with dental flossing in a vertical direction or tooth picking.
- Regular dental review - eliminate plaque.

ORAL DERMATOSES

The dermatoses include lichen planus, pemphigus vulgaris (uncommon), mucous membrane pemphigoid (uncommon) and lupus erythematosus. The clinical appearance of theseconditions in the mouth is quite different to the skin condition because of the environment, especially due to the presence of saliva.

Diagnosis

Histopathological examination (after approp-riate biopsy) with immunofluorescence is recommended, especially because of the similarity of the lesions of lupus erythematosus and lichen planus, which are both considered to be potentially premalignant in the mouth.[8]

Main features

Lichen planus:

- affects 2% of the population, usually over 45+10 years
- can vary from asymptomatic to severely painful
- usually white lace-like patterns on mucosa, cheeks and tongue
- may form superficial ulcers

Lupus erythematosus:

- oral lesions may be first sign of SLE
- usually on lateral aspects of the hard palate
- can resemble lichen planus

Treatment of oral dermatoses

Consider specialist referral.

Oral hygiene and pain control:

- chlorhexidine mouthwash or
- tetracycline/nystatin mouthwash or
- topical analgesics (e.g. lignocaine preparation)

Corticosteroids:

- topical (e.g. Kenalog in Orobase; betamethasone dipropionate 0.05%)
- intralesional (e.g. triamcinolone 10 mg/mL, especially for lichen planus)
- systemic - may be necessary in severe cases

THE PAINFUL TONGUE

Pain in the tongue is a reasonably common symptom in general practice. The cause is usually obvious upon examination but there are some obscure causes. As for many other oral mucosal problems, shared care with a dental or oral medical specialist is important. The causes of a sore or painful tongue are similar to that of the sore throat or mouth. Xerostomia is common in the elderly.

Investigations may include an FBE, serum vitamin B_{12}, folate and ferritin levels, a swab or a biopsy of a suspicious lesion.

A diagnosis strategy with lists of causes is presented in Table 72.3.

Table 72.3 Sore tongue: diagnostic strategy model

Q. *Probability diagnosis*
A. Geographic tongue
Atrophic glossitis
Trauma (bites, teeth, hot food/drink)
Aphthous ulceration
Herpes simplex virus (children)
Fissured tongue
Q. Disorders not to be missed
A. Carcinoma
HIV
Q. *Pitfalls (often missed)*

A. Anaemia: iron, vitamins B_6 and B_{12}, folate deficiency

Glossopharyngeal neuralgia

Lichen planus

Fissured tongue (rarely causes soreness)

Median rhomboid glossitis

Behçet's syndrome

Crohn's disease

Coeliac disease

Q. *Seven masquerades checklist*

A. Depression ✔

Diabetes ✔ Candida

Drugs ✔ mouthwashes, aspirin

Anaemia ✔ various

Thyroid disease -

Spinal dysfunction -

UTI -

Q. *Is the patient trying to tell me something?*

A. Possible with glossodynia.

Tongue tips

- Look for evidence of trauma, especially from a sharp tooth.
- A miserable child with a painful mouth and tongue is likely to have acute primary herpetic gingivostomatitis.
- When taking the history, take note of self-medications, especially sucking aspirin, a history of skin lesions (e.g. lichen planus) and consider underlying diabetes or immunosuppression.
- A long history of soreness with spicy or other foods indicates benign migratory glossitis (geographic tongue) or median rhomboid glossitis(Figure 72.3).



Figure 72.3 Disorders of the tongue: (a)median rhomboid glossitis; (b)geographic tongue; (c)black tongue

- Any non-healing or chronic ulcer requires urgent referral.
- Macroglossia (large tongue): consider acromegaly, myxvedesma, amyloidosis, lymphangioma.
- Strawberry tongue: Consider scarlet fever, Kawasaki's disease.
- Glossodynia (painful tongue) characteristically presents as a burning pain on the tip of the tongue. It can be a real ‘heartsink’ presentation. Consider depressive illness as an underlying cause.

Erythema migrans (geographic tongue)

Also known as benign migratory glossitis, this benign condition shows changing patterns of desquamatous areas and erythema on the dorsum and edges of the tongue. With the smooth red patches and raised whitish grey edges, the pattern resembles a relief map with mountain ridges. The border changes shape within weeks. It is irregular and slightly reddered.

It is considered to be a hypersensitivity reaction but the offending allergen has not been identified. Stress, tobacco, alcohol, marijuana and spicy foods can aggravate the condition in some individuals.

Management

- The condition is self-limiting and there is no specific therapy.
- Explanation and reassurance is important.
- No treatment is recommended if asymptomatic.
- If tender, benzocaine compound (Cepacaine) gargles 10 mL tds.
- If persistent and troublesome, low-dose inhaled glucocorticoid (e.g. beclomethasone 50 mg tds - don't rinse after use).

Black or hairy tongue

This is due to overgrowth of papillae or reduced wear of papillae, eg. debility and lack of fibrous foods.

- Appearance: dark, elongated filiform papillae giving brownish appearance to dorsum (posterior) of tongue.
- Symptoms: bad tastes and malodorous oral cavity.

Causes

- Unknown
- Poor oral hygiene/debility
- Iatrogenic (e.g. antibiotics)

Treatment

Brush tongue to remove stained papillae. Use a topical keratolytic agent such as salicylate, with pineapple being the most practical (95% cases are helped).

Method:

- Cut a thin slice of pineapple into eight segments. Slowly suck a segment on the back of the tongue for 40 seconds and then slowly chew it. Repeat until all segments are completed. Do this bd for 7 - 10 days.[1] Repeat if recurs.

Note: The salicylate in pineapple can aggravate irritable bowel syndrome.

ORAL DYSAESTHESIA

The classic chronic burning sensation of the oral cavity appears to have a neuropathic and/or psychological basis.[1] Symptoms include:

- altered sensitivity - burning pain or 'raw' sensation
- altered taste - sweet, salty or bitter
- altered saliva (subjective) - quality and quantity
- altered tooth sensation (e.g. 'phantom tooth pain')

Consider the underlying cause:

- haematinic deficiency - iron, folate, vitamin B_{12}
- autoimmune disorder (e.g. Sjögren's syndrome)
- endocrine disorder (e.g. diabetes)
- psychological disorder

Treatment

Consider:

- clonazepam 0.5 - 1 mg bd
- or, if resistant, gabapentin (Neurontin)

ORAL CARCINOMA

Cancer of the lip and oral cavity accounts for 2 - 3% of all newly diagnosed cancers in Australia.[9]

SCC is the most common malignancy of the oral cavity, accounting for 90% of cases. It has a 5-year survival rate of 68% without lymph node involvement and 25% with local node metastases.[10] Cancer of the lip is usually treated successfullyby excisional biopsy but intraoral cancer has significant morbidity and mortality.[9]

Other malignancies include mucoepi-dermoid carcinoma, lymphoma, Kaposi's sarcoma and malignant melanoma, which is usually found on the palate.

Predisposing or associated factors for SCC include tobacco and marijuana abuse, alcohol abuse, excessive sunlight and immune suppressive disorders such as HIV, lymphoma and various medications.

SCC is usually found as a chronic indurated ulcer on the ventral and lateral surfaces of the tongue followed by the floor of the mouth and buccal mucosa. It may present as a white patch or, more commonly, as a speckled white and red nodular patch or a red velvety patch.

The red patches of erythroplakia (in particular) and the white patches of leukoplakia may be premalignant or early invasive carcinoma and necessitate further investigation, particularly incisional biopsy.

Treatment for oral carcinoma is surgery ± radiotherapy and chemotherapy.

BENIGN INTRAORAL SWELLINGS AND TUMOURS

Epulis

An epulis is a benign, localised gingival swelling. It is a very ancient term with no pathological significance, meaning a 'tumour situatedon the gum'. There are two distinct types - a fibrous epulis and giant cell epulis. An epulis emerges between two teeth from the periodontal membrane where there is usually dentaldecay or a site of irritation, such as a partial denture. It appears to be more common during pregnancy where the epulis has a more vascular appearance.

Treatment is usually by excision, curettage of the origin and extracting associated teeth. The 'pregnancy' epulis should be left for several weeks after child birth before treatment.

Typical locations of intraoral tumours are shown in Figure 72.4.


blood blisters
(from dentures)
mucocele
torus
palatinus
tonsillar
retention
cyst
carcinoma
of tongue
ranula
epulis
fibrous mucosal
hyperplasia
mucocele

Figure 72.4 Typical location of benign swelling in the oral cavity

Pyogenic granuloma

These may occur on the gums or oral mucosa of the lips and look like pyogenic granulomas of the skin, which also are associated with minor trauma. It is best treated by excision.

Retention cysts (Mucous cysts)

The oral mucosa contains numerous mucous cysts and accessory mucous and serous salivary glands.

Small retention cysts are probably caused by minor trauma to the duct. They may rupture spontaneously. They commonly occur on the mucosa of the lower lip. Treatmentis by incision and enucleation under local anaesthesia. Larger ones require marsupialisation. Others occur on the tonsils where they usually appear as sessile yellow swellings. A special typeof retention cyst is the ranula.

Ranula

A ranula is a large transparent mucocele occurring in the floor of the mouth. A blue colour and small tortuous veins stretched across the surface is typical.

It is usually unilateral and simple but may extend into the tissues of the floor of the mouth and neck (plunging ranula). Patients may give a history of a cyst thatbursts and then returns.

Treatment is usually by marsupialisation.

Fibrous (fibroepithelial) hyperplasia

Hyperplasia of the oral mucosa, a very common condition, is usually seen on the floor of the mouth and is due to chronic irritation from ill-fitting dentures. Removal of the offending irritation is necessary. The hyperplasia may resolve but if it doesn't, surgical removal of the residual mass is necessary.

Haemangioma

These appear as a dark blue/purple sessile or modular swelling anywhere in and around the mouth, especially on the vermillion border of the lips, fesor of the mouthand tongue. They blanch or pressure. No treatment is needed except for pressing cosmetic reasons. Copious bleaching can occur with attempted removal.

Other soft tissue swellings

Swelling that may be encountered includes squamous papillomas (like viral warts), fibroepithelial polyps (inner side cheek), mixed salivary tumours, vascular haemartomas and giant cell granulomas.

The most common benign intraoral salivary neoplasm is the pleomorphicadenoma usually presenting as an asymptomatic swelling of the hard palate or cheeks.[2] Excision is recommended.

Bony exostoses

Bony outgrowths of the maxilla and mandible are reasonably common andthe hard intraoral lump may cause concern. The most common is known astorus palatinus, which is situated in the centre of the hard palate. A similar exostosis is torus mandibularis, which occurs inside the mandible, opposite the premolar teeth and is usually bilateral. These lesious are hamartomes and do not removal except if there is impending dental obstruction.

XEROSTOMIA (DRY MOUTH)

This is a symptom rather than a disease entity. It occurs in about 10% of the population and approximately 70% of patients have a systemic cause.[11]

The most common cause is a side-effect of drug therapy and it is relative rather than absolute. Some patients who have a dry mouth on clinical examination do not complain of it while others who do complain of it may be found to have a normal salivary flow rate (a feature of depressive illness).

Other obvious causes are dehydration, mouth breathing and psychogenic.

Primary xerostomia

Causes

- Salivary gland atrophy due to ageing
- Salivary gland infections
- Autoimmune salivary gland disease (e.g. Sjögren's syndrome)

> DxT: dry eyes + dry mouth + arthritis ° Sjögren's syndrome

Secondary xerostomia

Causes

- Drugs: antidepressants (especially tricyclic agents), diuretics, anticholinergics, tranquillisers, antihistamines, antiemetics, antihypertensives (some), antimigraine (some), antiparkinsonian, lithium and opioids
- Depression and anxiety (e.g. public speaking)
- Thirst/hunger
- Dehydration (e.g. diabetes, diarrhoea, renal failure)
- Anaemias: iron, folate, vitamin B_{12} deficiency

Consequences

Xerostomia interferes with speech, mastication and swallowing and causes difficulty in managing oral hygiene, especially dentures.

Symptoms include a burning sensation, a decrease in taste or a bad taste and fetid breath.

There is an increase in dental decay and perhaps a tendency to Candida albicans infection.

Treatment

This involves education, especially the need for meticulous oral hygiene, including topical fluoride preparations to the teeth and regular dental checks.

The cause must be diagnosed and treated if possible, especially a review of (and replacement of if necessary) drug therapy.

Avoid decongestants and antihistamines.

Strategies to consider

- Sip sugarless fluids frequently and chew sugarless gum (avoid mouthwashes containing sugars and alcohol).
- Use a saliva substitute (e.g. Aquae, SalivaOrthana) or frequent mouthwashes (e.g. lemon and glycerin, 5 - 10 mL in 100 mL water as required - can be used in a plastic squeeze bottle).
- Use sodium fluoride 0.5% mouthwash for 5 minutes each day.
- Topical applications of glycerin or paraffin oil to the lips.

HALITOSIS

Causes

The diagnostic strategy model for chronic halitosis (bad breath) is presented on p. 164. The commonest causes are orodental disorders secondary to poor oral hygiene and inappropriate diet. Bacterial putrefaction of dental and food debris, together with inflammation of the gums, are largely responsible for the oral malodour. Smoking, alcohol and a dry mouth will aggravate the problem. [A 1999 survey showed that 87% of patients with halitosis had an oral cause, 8% an ear, nose and throat cause with 5% having otheror unidentified causes.]

Management

- First exclude dental disease, malignancy (esp. nasopharyngeal carcinoma), pulmonary TB, hairy tongue, nasal and sinus infection.
- Refer for a dental check. Treat gingivitis.
- Consider drugs such as isosorbide dinitrateand various antidepressants as a cause.
- Cease smoking if this is a factor.
- Avoid or limit onions, garlic, peppers, curries, spicy salami and similar meats.
- Avoid or limit strong cheeses.
- Avoid smoking and excessive nips of alcohol.
- Brush teeth regularly during the day - immediately after meals.
- Gently brush the dorsum of the tongue with special, available soft brushes.
- Rinse mouth out with water after meals.
- Avoid fasting for long periods during the day.
- Drink copious amounts of water during the day.
- Gargle with mouthwash regularly (e.g. Listerine; Cepacol Mint mouthwash; 0.2% aqueous chlorhexidine).
- Use dental floss regularly to clean the teeth.
- Chew sugarless gum to help moisten the mouth.

Tip: use an oil/water wash (e.g. equal volumes of cetylpyridinium chloride (Cepacol) and olive oil), gargle a well-shaken mixture and spit out, qid.

Practice tips

- The tetracycline/nystatin mouthwash is an effective treatment for widespread mouth ulceration be it viral, aphthous, autoimmune or cytotoxic chemotherapy. It provides symptomatic relief and facilitates healing.[1]
  Formulation:
  - tetracycline 250 mg capsule - empty the capsule contents into 10 - 15 mL of warm water and shake vigorously
  - nystatin drops - add 1 mL to this mixture and stir

  Use 10 mL of the mouthwash held in the mouth for 2 minutes and expectorate. Do this 4 times a day for 5 days.
- Recurrent herpes simplex ulceration is not common in the oral mucosa and if suspected should be confirmed by laboratory investigation. Thetreatment is different to aphthous ulceration so clinical distinction is important. Herpes simplex virus is aggravated by topical steroids.
- For unusual mouth ulceration consider acute leukaemia, carcinoma, blood dyscrasias, Crohn's disease and drug therapy such as antiepileptics and antihypertensives.

REFERENCES

1. Tversky J. Oral mucosal disease. In: *Skin and Cancer Foundation Dermatology Conference Proceedings.* Melbourne, 2002.
2. Angel CM et al. Non-neoplastic oral swellings. Aust Fam Physician, 1992; 21: 188 - 9.
3. Vickers R. Oral ulcers. Med Observer, 2000; 12 May: 78 - 9.

4. Rogers AH, Gully NJ. Melaleuca (tea tree oil) for mouth ulcers. J Dent Res 1998; 78: 949.
5. Tierney LM et al. Current Medical Diagnoses and Treatment (41st edn). New York: McGraw-Hill & Lange, 2002: 250.
6. Bastiaan RJ. Periodontal disease. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 403 - 4.
7. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (12th edn). Melbourne: Therapeutic Guidelines, 2003: 118 - 130.
8. Reade PC, Rich AM. Oral dermatoses. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 362 - 4.
9. Rich AM, Reade PC. Oral cancer. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 360 - 1.
10. Beers MH, Berkow R. The Merck Manual of Diagnosis and Therapy (17th edn). New Jersey: Merck Research Laboratories, 1999: 758.
11. Reade PC, Rich AM. Xerostomia. In: *MIMS Disease Index*(2nd edn). Sydney: IMS Publishing, 1996: 583.

CHAPTER 73. SORE THROAT

I believe there are hundreds of young adults who have erroneously suffered tonsillectomy because of the tonsillitis of undiagnosed glandular fever.

SCEPTICAL GP (ANONYMOUS)

SORE THROAT

A sore or painful throat is one of the commonest symptoms encountered in general practice. The most usual cause is viral pharyngitis, which is self-limiting and usually only requires symptomatic treatment.

Definitions

- Pharyngitis ° inflammation of pharynx ± tonsils
- Tonsillitis ° inflammation of tonsils only
- Quinsy ° a peritonsillar abscess

KEY FACTS AND CHECKPOINTS

- In the National Morbidity Survey (UK)[1] nine episodes per annum of acute pharyngitis or acute tonsillitis were diagnosed for every 100 patients.
- Sore throats account for about 5% of consultations in general practice per annum.[2]
- In one UK general practice it was the third most common new presenting symptom-5.4% of presenting problems.
- Although throat infections are common from infancy, children under 4 years of age rarely complain of a sore throat.
- Complaints of a sore throat are prevalent in children between 4 and 8 years and in teenagers.
- Sore throats continue to be common up to the age of 45 and then decline significantly.
- The common causes are viral pharyngitis (approximately 60 - 65%) and tonsillitis due to Streptococcus pyogenes (approximately 20%).
- The sore throat may be the presentation of serious and hidden systemic diseases, such as blood dyscrasias, HIV infection and diabetes (due to candidiasis).
- A very important cause is tonsillitis caused by Epstein-Barr mononucleosis (EBM). Treating this cause with penicillin can produce adverse effects.
- As a general rule antibiotics should not be prescribed to treat a sore throat, excluding evidence of group A beta-haemolytic Streptococcus (GABHS) infection.[3]

Presentation

Sore throat may be present as part of a complex of the common upper respiratory infections, such as the common cold and influenza. However, sore throat often presents as a single symptom. The pain is usually continuous and aggravated by swallowing. In those under 4 years old the presentation of acute pharyngitis or tonsillitis may be confusing as the presenting complaints may be vomiting, abdominal pain and fever rather than sore thoat and swallowing difficulty. It is appropriate to consider sore throat as acute or chronic. Most presentations come as acute problems, the causes of which are listed in Table 73.1.

Table 73.1 Causes of acute sore throat

Bacteria
Beta-haemolytic streptococci (GABHS) Diphtheria (rare) Gonococcal pharyngitis *Haemophilus influenzae*

Quinsy *Staphylococcus aureus* (rare) Syphilis (rare) Vincent's angina Viral
Severe-moderate soreness Epstein-Barr mononucleosis Herpangina Herpes simplex pharyngitis *Mild-moderate soreness* Adenovirus Coronavirus Enterovirus Influenza virus Picornavirus Rhinovirus Human immunodeficiency virus Varicella (chicken pox) Other infections
Candida albicans, especially in infants *Mycoplasma pneumoniae* *Chlamydia pneumoniae* **Blood dyscrasias**
Agranulocytosis Leukaemia Irritants Tobacco smoke Antiseptic lozenges (oral use)

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 73.2.

Table 73.2 Sore throat: diagnostic strategy model

Q. Probability diagnosis
A. Viral pharyngitis
Streptococcal tonsillitis Chronic sinusitis with postnasal drip **Q. Serious disorders not to be missed**
A. Cardiovascular

- angina
- myocardial infarction

Neoplasia

- carcinoma of oropharynx, tongue

Blood dyscrasias (e.g. agranulocytosis, acute leukaemia)

Severe infections

- acute epiglottitis (children and adults)
- peritonsillar abscess (quinsy)
- pharyngeal abscess
- diphtheria (very rare)
- HIV/AIDS

Q. Pitfalls (often missed)

A. Foreign body (e.g. fish bone)

Epstein-Barr mononucleosis

Candida

- common in infants
- steroid inhalers

STIs

- gonococcal pharyngitis
- herpes simplex (type II)
- syphilis

Irritants (e.g. cigarette smoke, chemicals)

Reflux oesophagitis → pharynolaryngitis

Chronic mouth breathing

Aphthous ulceration

Thyroiditis

Rarities

Systemic sclerosis

Sarcoidosis

Malignant granuloma

Tuberculosis

Q. Seven masquerades checklist

A. Depression

Diabetes (Candida)

Drugs

Anaemia possible

Thyroid disorder thyroiditis

Spinal dysfunction cervical

UTI -

Q. Is the patient trying to tell me something?

A. Unlikely, but the association with depression is significant.

Probability diagnosis

At least 50% of sore throats, mainly pharyngitis, will be caused by a virus. A viral infection is supported by the presence of coryza prodromata, hoarseness and nasal stuffiness.

Serious disorders not to be missed

It is vital to be aware of Haemophilus influenzae infection in children, especially between 2 and 4 years, when the deadly problem of epiglottitis can develop suddenly. These patients present with a short febrile illness, respiratory difficulty (cough is not a feature) and are unable to swallow.

Apart from acute epiglottitis it is important not to overlook carcinoma of the oropharynx or tongue, or the blood dyscrasias, including acute leukaemia. (Refer to pp. 240 - 1.) The severe infections not to be missed include streptococcal pharyngitis with its complications, including quinsy, diphtheria and HIV infection (including AIDS).

Pitfalls

There are many pitfalls, the classic being to diagnose the exudative tonsillitis of EBM as streptococcal tonsillitis and prescribe one of the penicillins, which may precipitate a severe rash. Primary HIV infection can present with a sore throat along with other symptoms. Adenovirus pharyngitis can also mimic streptococcal pharyngitis, especially in young adults.

Traumatic episodes are important but are often not considered, especially in children. They include:

- a foreign body-may cause a sudden onset of throat pain, then drooling and dysphagia
- vocal abuse-excessive singing or shouting can cause a sore throat and hoarseness
- burns-hot food and drink, acids or alkalis

Various irritants, especially cigarette smoke in the household and smoke inhalation from fires, can cause pharyngeal irritation with sore throat, especially in children.

The mouth and pharynx may become dry and sore from mouth breathing, which is often associated with nasal obstruction (e.g. adenoid hypertrophy, allergic rhinitis).

Seven masquerades checklist

Depression may be associated with a sore throat. Diabetes and aplastic anaemia and drugs are indirectly associated through candidiasis, neutropenia and agranulocytosis respectively. NSAIDs can cause a sore throat. The possibility of thyroiditis presenting as a sore throat should be kept in mind.

Making a diagnosis

The issues of making a reliable diagnosis and prescribing antibiotics are rather contentious and at times difficult. It is difficult to distinguish clinically between bacterial and viral causes. The main issue is to determine whether the sore throat has a treatable cause by interpretation of the clinical and epidemiological data.

The appearance of the pharynx and tonsils is not always discriminating. A generalised red throat may be caused by a streptococcal or a viral infection, as may tonsils that are swollen with follicular exudates. On probability, most sore throats are caused by a virus and generally do not show marked inflammatory changes or purulent-looking exudates (Figure 73.1). Such throats should be treated symptomatically.



Figure 73.1 Viral pharyngitis: the signs may be minimal but mild redness of pharynx and prominent lymphoid patches on the oropharynx are typical

THE CLINICAL APPROACH

History

It is necessary to determine whether the patient has a sore throat, a deep pain in the throat or neck pain. Instruct the patient to point to exactly where the pain is experienced. Enquire about relevant associated symptoms such as metallic taste in mouth, fever, upper respiratory infection, other pain such as ear pain, nasal stuffiness or discharge and cough.

Note whether the patient is an asthmatic and uses a corticosteroid inhaler, or is a smoker, or exposed to environmental irritants. Check the immunisation history, enquiring especially about diphtheria.

The history should give a clue to the remote possibility that the painful throat is a manifestation of angina.

Examination

An inspection should note the general appearance of the patient, looking for 'toxicity', the anaemic pallor of leukaemia, the nasal stuffiness of infectious mononucleosis, the characteristic halitosis of a streptococcal throat.

Palpate the neck for soreness and lymph-adenopathy, inspect the ears and check the sinus areas.

Then inspect the oral cavity and pharynx. Look for ulcers, abnormal masses and exudates. Note whether the uvula and soft palate, tonsils, fauces or pharynx are swollen, red or covered in exudate. The typical appearances of various conditions causing a sore throat are shown in Figures 73.1 to 73.7, and important causes to exclude in Figure 73.7

Guidelines

- Small patches of exudate on the palate or other structure indicate Candida albicans (oral thrush) (Figure 73.2).



Figure 73.2 Oral thrush due to Candida albicans in a diabetic patient. Small patches of yellow white exudate are seen on the palate, dorsum of the tongue, pharynx and mucosa. Photo courtesy Hugh Newton-John

- A large whitish-yellow membrane virtually covering both tonsils indicates EBM (Figure 73.3).



Figure 73.3 Tonsillitis of Epstein-Barr mononucleosis showing swollen red tonsils, with a whitish-yellow membranous exudate, swollen uvula and petechiae on the soft palate. Photo courtesy Hugh Newton-John

- A generalised red, swollen appearance with exudate indicates GABHS infection (Figures 73.4 and 73.5).



Figure 73.4 Acute follicular tonsillitis due to Streptococcus pyogenes: the tonsils are red and swollen with pockets of pus. Photo courtesy Hugh Newton-John



Figure 73.5 Streptococcal tonsillopharyngitis: severe inflammation involves both tonsils and pharynx with marked redness, swelling and exudate. Consider herpes simplex and mononucleosis as alternative diagnoses

Investigations

Investigations can be selected from:

- throat swab
- haemoglobin, blood film and white cell count
- mononucleosis test
- random blood sugar (? diabetes)

To swab or not to swab

Throat swabs are about 90% effective in isolating GABHS from the infected throat. Authorities are divided about management. Some recommend that throat cultures be performed for all sore throats and antibiotics given only when GABHS is found. Others regard throat cultures as being unnecessary and recommend therapy based on clinical judgments. Swabs are seldom helpful because the isolation of GABHS often represents asymptomatic carriage.[4] Still others recommend throat cultures for selected patients only.[5]

Generally, throat cultures are not necessary except to verify the presence of S. pyogenes, especially in closed institutions such as boarding schools, or if diphtheria is suspected in the non-immunised. One study has found that toothbrushes harbour GABHS and should not be shared.6 A positive culture and a fourfold rise in the ASOT titre is necessary for a precise diagnosis.

Epstein-Barr mononucleosis screening

It is important initially if tonsillar exudate is present to consider the possibility of EBM. If suspected, an IgM antibody test should be ordered, rather than the older tests, such as a Paul Bunnell test.

Supportive symptomatic treatment

Supportive measures include:

- adequate soothing fluids, including icy poles
- analgesia:
  adults-2 soluble aspirin
  children-paracetamol elixir (not alcohol base) or ibuprofen
- rest with adequate fluid intake
- soothing gargles (e.g. soluble aspirin used for analgesia)
- advice against overuse of OTC throat lozenges and topical sprays, which can sensitise the throat; limited use (3 days) of decongestants for nasal decongestion is helpful

SORE THROAT IN CHILDREN

An acute sore throat in a child usually means viral or, less commonly, bacterial infection of the tonsillopharynx. A bacterial cause is more common in children aged 3 - 13 years than in children < 3 years. Other causes to consider are:

- gingivostomatitis, especially primary herpes simplex
- epiglottitis
- laryngotracheobronchitis (croup)
- laryngitis
- oral candidiasis (more a bad taste than pain)
- aphthous ulcers
- foreign bodies
- postnasal drip (e.g. allergic rhinitis)
- irritation
  - low environmental humidity
  - smoke (e.g. household smoke)

SORE THROAT IN THE ELDERLY

Sore throat in the elderly may be caused by a viral infection but otherwise needs to be treated with considerable respect. It is important to exclude pharyngeal carcinoma which can present with the classic triad.

DxT: painful swallowing + referred ear pain + hoarseness ° pharyngeal carcinoma

Oropharangeal lesions may occur with herpes zoster but vesicles are usually present on the face.

A metallic taste in the mouth with or without a complaint of a sore throat indicates C. albicans and hence diabetes must be excluded.

STREPTOCOCCAL TONSILLOPHARYNGITIS

This infection may involve the pharynx only and vary from mild to severe, or it may involve both tonsils and pharynx. It is uncommon under 3 years or over 40 years.[7]

Guidelines for streptococcal throat

The four diagnostic features are:

- constitutional symptoms
    - fever ≥ 38°C
    - toxicity
- tender anterior cervical lymphadenopathy
- tonsillar swelling and exudate
- absence of cough

Other symptoms include:

- difficulty in swallowing
- significant pain, including pain on talking
- foul-smelling breath

Examination

Pharynx very inflamed and oedematous. Tonsils swollen with pockets of yellow exudate on surfaces (Figures 73.4 and 73.5). Very tender enlarged tonsillar lymph nodes.

Treatment

Indications for antibiotic therapy:[4]

- severe tonsillitis with above features of GABHS
- existing rheumatic heart disease at any age
- scarlet fever
- peritonsillar cellulitis or abscess (quinsy)
- patients 3 - 25 years with presumptive GABHS from special communities (e.g. remote Indigenous) with a high incidence of acute rheumatic fever

One evidence-based review recommended that if a cold follows its natural course, no antibiotics will be helpful. If there is a sore throat with no cough, but fever > 38°C, tender neck glands and white spots in the throat, antibiotics are indicated.[8] Treatment should be with penicillin or an alternative antibiotic (Table 73.3).[4]

Table 73.3 Treatment for streptococcal throat (proven or suspected)[4]

Children
phenoxymethyl penicillin 50 mg/kg/day (o) in 2 divided doses for 10 days (to max. 1 g/day) or (if sensitive to penicillin) roxithromycin 4 mg/kg up to 150 mg (o) bd for 10 days Adults
phenoxymethyl penicillin 500 mg (o) 12 hourly for 10 days (can initiate treatment with one injection of procaine penicillin) or roxithromycin 300 mg (o) daily for 10 days In poorly compliant patients:
benzathine penicillin 900 mg IM as a single dose in adults In severe cases:

Procaine penicillin 1 - 1.5 mg IM daily for 3 - 5 days plus phenoxymethyl penicillin (as above) for 10 days

Note: Although symptoms and most evidence will disappear within 1 - 2 days of treatment, a full course of 10 days should be given to provide an optimal chance of eradicating S. pyogenes from the nasopharynx and thus minimising the risk of recurrence or complications such as rheumatic fever.[4] Some studies indicate that 7 days may be sufficient.

Antibiotic treatment has a variable effect on the resolution of symptoms. It does not protect against glomerulonephritis but does protect against rheumatic fever.[7] Amoxycillin should be avoided in tonsillitis because of confusion caused should mononucleosis be present. Frequent fluids are advisable.

Recurrent tonsillitis[4, 7]

Treat with prophylactic penicillin for patients with more than five episodes of presumptive bacterial tonsillitis in a year. The decision should be based on the severity of the episode, time lost from work or school, infectivity and response to antibiotics.

QUINSY

Quinsy a peritonsillar abscess characterised by marked swelling of the peritonsillar area with medial displacement of tonsillar tissue (Figure 73.6). It is usually caused by GABHS or anaerobes, occasionally Haemophilus sp. and Staphylococcus aureus. A typical picture of tonsillitis is followed by increasing difficulty in swallowing and trismus.



Figure 73.6 Peritonsillar abscess (quinsy): a tense red bulging mass is noted and the uvula is displaced from the mid-line; a site of incision for drainage is indicated

Treatment

Antibiotics (e.g. procaine penicillin IM or clindamycin) plus drainage under local anaesthetic if it is pointing. Oral penicillin treatment is likely to fail. Subsequent tonsillectomy may, but not always, be necessary.

ACUTE EPIGLOTTITIS

In children this is a life-threatening infection (p. 935). It may be overlooked in adults where, unlike children, the airway is usually not obstructed and the patient presents with a severe sore throat, dysphagia, drooling of saliva and a tender neck. Examination of the throat may appear quite normal. However, it is a severe infection requiring hospitalisation and parenteral antibiotics (e.g. cefotaxime).

VIRAL CAUSES OF SORE THROAT

Epstein-Barr mononucleosis

The angiose form of EBM is a real trap and must be considered in 15 - 25-year old patients (peak incidence) with a painful throat that takes about 7 days to reach its peak. Refer to Chapter 29.

Clinical features

- Sore throat
- Prodromal fever, malaise, lethargy
- Anorexia, myalgia
- Nasal quality to voice
- Skin rash

Examination

- Petechiae on palate (not pathognomonic)
- Enlarged tonsils with or without white exudates (looks, but isn't, purulent)
- Periorbital oedema
- Lymphadenopathy, especially posterior cervical
- Splenomegaly (50%)
- Jaundice ± hepatomegaly (5 - 10%)

The rash

- Primary rash (5%)
- Secondary rash
    - with ampicillin, amoxycillin (90 - 100%)
    - with penicillin (50%)

Note: This rash is not synonymous with penicillin hypersensitivity.

Diagnosis

- Blood film-atypical lymphocytes
- White cell count-absolute lymphocytosis
- Heterophil antibodies or monospot test or
- EBV IgM test (more specific)

Herpangina

An uncommon infection caused by the Coxsackie virus. Presents as small vesicles on soft palate, uvula and anterior fauces. These ulcerate to form small ulcers. The problem is benign and rapidly self-limiting.

Herpes simplex pharyngitis

In adults primary infection is similar to severe streptococcal pharyngitis but ulcers extend beyond the tonsils.

Other viral pharyngitis

Typically, the signs are fewer than with other causes. The typical case has mild redness without exudate and prominent (sometimes pale) lymphoid patches on the posterior pharynx (Figure 73.1). Tonsillar lymph nodes are usually not enlarged or tender. This picture is the commonest encountered in general practice.

DIPHTHERIA

The potentially fatal form of this disease almost always occurs in non-immunised people. The clinical presentation may be modified by previous immunisation or by antibiotic treatment.

Typical clinical features

- Insidious onset
- Mild to moderate fever
- Mild sore throat and dysphagia
- Patient looks pale and ill
- Enlarged tonsils
- Pharynx inflamed and oedematous
- Membrane (any colour but usually grey-green) can spread beyond tonsils to fauces, soft palate, lateral pharyngeal wall and downwards to involve larynx (Figure 73.7)
- enlarged cervical lymph nodes
- soft tissue swelling of neck → 'bull neck' appearance



Figure 73.7 Diphtheria: tonsils and pharynx are red and swollen; a thick grey-green exudate forms on the tonsils as a spreading membrane


trauma
glossopharyngeal neuralgia
carcinoma
• palate
• tonsil
• tongue
aphthous ulcers
tonsils
• tonsillitis
• quinsy
pharyngitis

Figure 73.8 General causes of a sore throat: note the importance of excluding carcinoma

Management

- Throat swabs
- Antitoxin
- Penicillin or erythromycin 500 mg qid for 10 days
- Isolate patient

CANDIDA PHARYNGITIS

Oral candidiasis typically presents as milky-white growths on the palate, buccal and gingival mucosae, pharynx and dorsum of the tongue (Figure 73.2). If scraped away, a bleeding ulcerated surface remains. A bad (metallic) taste is a feature but the patient may complain of a sore throat and tongue and dysphagia.

Causes or predisposing factors to consider:

- HIV infection
- diabetes mellitus
- broad spectrum antibiotics
- corticosteroids, including inhalers
- dentures
- debility

Management

- Determine underlying cause.
- Nystatin suspension, rinse and swallow qid *or* amphotericin 10 mg lozenge dissolved slowly in oral cavity, 6 hourly, for 7 - 14 days.

WHEN TO REFER

- Acute epiglottitis in children (a medical emergency)
- Inaccessible foreign body
- Abscess: peritonsillar or retropharangeal
- Recurrent attacks of tonsillitis and adenoid hypertrophy for an opinion about tonsillectomy and/or adenoidectomy

- Suspicion or evidence of HIV infection or diphtheria
- Patients not responding to treatment
- Patients with more generalised disorders that are not yet diagnosed[9]

GUIDELINES FOR TONSILLECTOMY[9]

- Repeated attacks of acute tonsillitis
- Enlarged tonsils and/or adenoids causing airway obstruction, including OSA
- Chronic tonsillitis
- More than one attack of peritonsillar abscess
- Biopsy excision for suspected new growth

Antibiotic treatment is aimed primarily at streptococcal pharyngitis and this is often based on clinical judgment.

Practice tips

- Consider severe tonsillitis with a covering membrane as EBM.
- If an adult presents with an intensely painful throat with a heavy exudate and seems toxic, consider primary herpes simplex as well as streptococcal throat.
- Reserve swabs of the throat for verification of a streptococcal throat where it is important to do so, for suspected diphtheria and for suspicion of other serious infections such as tuberculosis.
- Be aware of possible complications, such as febrile convulsions in children and abscess formation.
- Do not misdiagnose unusual causes of a sore throat, such as carcinoma (Figure 73.7).
- The triad: hoarseness, pain on swallowing and referred ear pain ° pharyngeal carcinoma.
- The two major considerations in managing the acute sore throat are:
  - Can diphtheria be excluded?
  - Should the patient be treated with an antibiotic?

REFERENCES

1. Office of Population Censuses and Surveys. *Morbidity Statistics from General Practice Studies on Medical and Population Subjects*. No 26. London: HMSO, 1974: 33 - 40.
2. Cormack J, Marinker M, Morrell D. *Practice: A Handbook of Primary Medical Care*. London: Kluwer-Harrap, 1980: 3(25): 1 - 7.
3. Del Mar CB, Glasziou PO, Spinks AB. *Antibiotics for Sore Throat* (Cochrane review). In: The Cochrane Library, Issue 1. Oxford: Update Software, 2002.
4. Spicer J (Chair). *Therapeutic Guidelines: Antibiotics* (13th edn). Melbourne: Therapeutic Guidelines Ltd, 2006: 232 - 4.
5. Yung AP, Stanley PA. The patient with the acute sore throat. Curr Ther, 1985; 26: 54 - 5.
6. Brook I, Gober AE. Persistence of group A β-hemolytic streptococci in toothbrushes and removable orthodontic appliances following treatment of pharyngotonsillitis. Arch Otolaryngol Head Neck Surg, 1998; 124: 993 - 5.
7. Cooper RJ, Hoffman JR et al. Principles of appropriate antibiotic use for acute pharyngitis in adults. Ann Intern Med, 2001; 134: 509 - 17.
8. Rosser W, Shafir MS. *Evidence-Based Family Medicine*. Hamilton: BC Decker Inc, 1998: 108 - 10.
9. Benjamin B. Indications for tonsil removal. AustDoctor, 1999; 3 December: 63 - 5.

CHAPTER 74. TIREDNESS

Chronic fatigue syndrome is not tiredness: there is a difference between feeling tired and 'fatiguey'. Fatigue involves a heaviness in the limbs, a sense of inability to think or move, pain in muscles and joints, nausea etc. Please understand the difference.

CPS PATIENT TO AUTHOR, JANUARY 1995

TIREDNESS

Tiredness or chronic fatigue is not a diagnosis but rather a symptom of illness: it may occur as either a presenting or a supporting symptom. Tiredness is interchangeable with terms such as weariness, loss of energy, listlessness and exhaustion. It is a common and difficult presenting symptom. The symptom of tiredness is likely to be 'hidden' behind the request for a tonic or a physical check-up.[1]

Tiredness can be a symptom of a great variety of serious and uncommon diseases, including malignant disease. The challenge for the family doctor is to diagnose such disorders quickly without extravagant investigation.

KEY FACTS AND CHECKPOINTS

- The commonest cause of tiredness is psychological distress, including anxiety states, depression and somatisation disorder.
- The study by Hickie et al.[2] showed that 25% of a sample of attendees visiting general practices had chronic fatigue. Of these, 70% had psychological distress. The others were more likely to have a current depressive disorder.
- In Jerrett's study[3] no organic cause was found in 62.3% of his patients presenting with lethargy; the constant factors were sleep disturbance and the presence of stress in their lives. Many of them turned out to be suffering from psychological problems or psychiatric illnesses, including depression, anxiety state or bereavement.
- An important cause of daytime tiredness is a sleep disorder such as obstructive sleep apnoea which results in periodic hypoventilation during sleep. It occurs in 2% of the general population in all age groups and in about 10% of middle-aged men.[3] A history of snoring is a pointer to the problem. See Chapter 71.
- Underlying disorders that need to be considered as possible causes of chronic fatigue are endocrine and metabolic disorders, malignancy, chronic infection, autoimmune disorders, primary psychiatric disorders, neuromuscular disorders, anaemia, drugs and cardiovascular disorders.
- Prolonged or chronic tiredness is characterised clinically by disabling tiredness, typically lasting more than 2 weeks, associated with non-restorative sleep, headaches and a range of other musculoskeletal and neuropsychiatric symptoms.[2]
- Sociodemographic correlates are concurrent psychological distress, female sex, lower socioeconomic status and fewer total years of education.[2]
- Chronic fatigue syndrome (CFS) is defined as debilitating fatigue, persisting or relapsing over 6 months, associated with a significant reduction in activity levels of at least 50%, and for which no other cause can be found.

CAUSES OF TIREDNESS

Analysing the symptom and reaching a diagnosis demands considerable skill since tiredness may indicate the first subtle manifestation of a serious physical disease or, more commonly, may represent a patient's inability to deal with the problems of everyday life. Chronic tiredness or fatigue is a feature of the 'high pressure' nature of many people's lifestyles.

Careful consideration must be given to the differentiation of physiological tiredness, as a result of excessive physical activity, from psychological tiredness. Furthermore, before diagnosing tiredness as psychological, pathological as well as physical causes must be excluded.

A summary of causes of chronic tiredness is presented in Table 74.1.

Table 74.1 Causes of chronic tiredness/fatigue

Psychogenic/non-organic
Psychiatric disorders
• anxiety states
• depression
• other primary disorders
• bereavement
• somatisation disorder
Lifestyle factors
• workaholic tendencies and 'burnout'
• lack of exercise/sedentary lifestyle
• mental stress and emotional demands
• exposure to irritants (e.g. carbon monoxide, 'lead' fumes)
• inappropriate diet
• obesity
• sleep deprivation
Organic
Congestive cardiac failure
Anaemia
Malignancy
HIV/AIDS
Subacute to chronic infection (e.g. hepatitis, malaria, Lyme disease)
Endocrine
• thyroid (hyper- and hypo-)
• adrenal (Cushing's syndrome, Addison's disease (p. 224))
• hyperparathyroidism
• diabetes mellitus
Nutritional deficiency
Kidney failure
Liver disorders - chronic liver failure, chronic active hepatitis
Respiratory conditions (e.g. asthma, COPD)
Neuromuscular (e.g. MS, myasthenia gravis, Parkinson's disease)
Metabolic (e.g. hypokalaemia, hypomagnesaemia)
Drug toxicity, addiction or side-effects (see Table 74.3)
Autoimmune disorders
Sleep-related disorders
Postinfectious fatigue syndrome (e.g. influenza, mononucleosis)
Unknown

Fibromyalgia
Chronic fatigue syndrome

Table 74.3 Drugs that can cause tiredness

Alcohol
Analgesics
Antibiotics
Anticonvulsants
Antidepressants
Antiemetics
Antihistamines
Antihypertensives
Anxiolytics
Corticosteroids
Ergot alkaloids
Digoxin
Hormones (e.g. oral contraceptives)
Hypnotics
Nicotine
NSAIDs
Vitamins A and D (early toxic symptoms)

Note: Most drugs have a considerable capacity to cause tiredness.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 74.2.

Table 74.2 Tiredness/chronic fatigue: diagnostic strategy model

Q. Probability diagnosis
A. Stress and anxiety
Depression
Viral/postviral infection
Sleep-related disorders (e.g. sleep apnoea)
Q. Serious disorders not to be missed
A. Malignant disease
Cardiac arrhythmia (e.g. sick sinus syndrome)
Cardiomyopathy
Anaemia
Haemochromatosis
HIV infection

Hepatitis C

Q. Pitfalls (often missed)

A. ‘Masked’ depression

Food intolerance

Coeliac disease

Chronic infection (e.g. Lyme disease)

Incipient CCF

Fibromyalgia

Lack of fitness

Drugs: alcohol, prescribed, withdrawal

Menopause syndrome

Pregnancy

Neurological disorders

- Post head injury
- CVA
- Parkinson's disease

Kidney failure

Metabolic (e.g. hypokalaemia, hypomagnesaemia)

Chemical exposure (e.g. occupational)

Rarities

Hyperparathyroidism

Addison's disease (p. 224)

Cushing's syndrome

Narcolepsy

Multiple sclerosis

Autoimmune disorders

Q. Seven masquerades checklist

A. Depression

Diabetes

Drugs

Anaemia

Thyroid disorder

Spinal dysfunction

UTI

Q. Is the patient trying to tell me something?

A. Highly likely.

Probability diagnosis

The most probable diagnoses to consider are:

- tension, stress and anxiety

- depression
- viral or postviral infection
- sleep-related disorders

Research studies have reported that over 50% (and in some cases as many as 80%) of reported cases of fatigue have been of psychological causation.[2, 4] Overwork is a common cause of fatigue and is often obvious to everyone but the patient. The modern approach to sleep-related disorders has revealed several important factors causing excessive tiredness.

Serious disorders not to be missed

Many serious disorders such as anaemia, malignant disease and subacute or chronic infections (e.g. hepatitis, bacterial endocarditis and tuberculosis) can be 'hidden' or masked in the initial stages or not readily apparent. Neuromuscular diseases such as myasthenia gravis and multiple sclerosis, connective tissue disorders and HIV infection also have to be excluded.

Pitfalls

The symptom of tiredness is fraught with pitfalls. Common ones include depression and other psychoneurotic disorders, and incipient congestive cardiac failure. Drug intake is a very common pitfall, whether it be by self-administration (including alcohol) or iatrogenic.

Tiredness is a feature of pregnancy in many women, so this association is worth keeping in mind, especially in the early stages when a change in menstrual history is not given or a young single woman will attempt to conceal the fact. It is also a presenting symptom of the menopause syndrome, which should not be misdiagnosed. Two classic causes of tiredness are haemochromatosis and coeliac disease.

Seven masquerades checklist

All these important problems are capable of being responsible for tiredness, especially depression, diabetes, drugs, anaemia and urinary infection. Thyroid disorder could certainly be responsible. Spinal pain can indirectly cause tiredness. Drugs that commonly cause tiredness are listed in Table 74.3.

Antihypertensives require special consideration. Drug withdrawal, especially for illicit drugs such as amphetamines, marijuana, cocaine and heroin, has to be considered.

Psychogenic considerations

Tiredness is a symptom that may represent a 'ticket of entry': a plea for help in a stressed, anxious or depressed patient. Any of the primary psychiatric disorders can present as tiredness.

THE CLINICAL APPROACH

In routine history taking, it is mandatory that questions be asked about the following if the information is not volunteered by the patient:

- sleep pattern (it is not uncommon for patients to say they sleep well and yet on questioning it is found they have initial insomnia, or middle insomnia, or both, with or without early morning waking). It is most relevant to talk to any sleeping partners to obtain a history of sleep disturbance.
- weight fluctuations.
- energy - performance - ability to cope.
- sexual activity.
- suicidal ideas.
- self-medication - OTC preparations (e.g. bromides, stimulants, analgesics, alcohol, cigarettes, other drugs). This is particularly important in the drug addiction-prone group: doctors, chemists, nurses, workers in the liquor industry,

truck drivers.

- fears (including phobic symptoms, hypochondriasis).
- precipitating factors (present in over 50% of patients with depressive illness):
  - postpartum
  - postoperative
  - associated with chronic physical illness
  - bereavement
  - pain - chronic pain conditions
  - retirement
  - medication
  - post trauma (e.g. motor vehicle accident)
  - postviral infections, especially hepatitis, mononucleosis, influenza
- work history - determine whether the patient is a workaholic.
- dietary history - determine pattern, including fad diets or skipped meals.
- menstrual history and symptoms related to the menopause syndrome.
- self-question: ‘Is this patient depressed?’.

Physical examination

A routine physical examination is important, followed by a more detailed specific examination relevant to the individual patient. In particular, it is important to ascertain the presence of hepatosplenomegaly and lymphadenopathy. In general the physical examination is unrewarding. In the chronic fatigue syndrome the relevant abnormal findings are muscle tenderness, mild pharyngitis and tender slightly enlarged cervical lymph nodes. When an alternative underlying medical illness is responsible for the tiredness there will usually be evidence for this on physical examination (e.g. positive Babinski reflex in multiple sclerosis, postural hypotension in Addison's disease (p. 224), right ventricular liftwith an atrial septal defect). A mental state assessment should be considered.

Investigations

Investigations should be selected judiciously from the following (tests that most patients should have when the examination is completely normal are marked *):

- haemoglobin, blood count and film *
- ESR */CRP *
- ECG and Holter monitor
- thyroid function tests *
- liver function tests *
- urea/kidney function tests *
- serum electrolytes (including calcium and magnesium) *
- blood sugar *
- plasma cortisol
- serumiron, ferritin, transferrin saturation *
- microscopy and culture of urine *
- tests for autoimmune disorders
  - antinuclear antibodies
  - rheumatoid factor
- HIV screening
- chest X-ray and spirometry
- chronic infection screening (consider): hepatitis A, B, C, D, E, cytomegalovirus, EBM, Ross River virus, Lyme disease, brucellosis, Q fever, tuberculosis, malaria, infective endocarditis, toxoplasmosis
- primary neuromuscular disorders
  - muscle enzyme assay

  - electromyography
- tissue markers for malignancy
- referral to a sleep disorder laboratory for sleep apnoea studies

The diagnosis of CFS can only be made when the minimum investigations (listed *) have been shown to be normal or to demonstrate minor abnormalities in liver function or blood film (atypical lymphocytes).

TIREDNESS IN CHILDREN

Tiredness in children is caused by a range of predictable conditions, such as physiological factors (excessive exercise, lack of sleep, poor diet), infections, allergies including asthma, drugs, depression and various illnesses in general.

Overweight children are likely to fatigue more rapidly than children of normal weight.[5] Any bacterial, viral or other infection may be associated with tiredness, with EBM being very significant in adolescents. Chronic EBV infection causing recurrent episodes of fever, pharyngitis, malaise and adenopathy can occur, especially in teenagers, who present with chronic exhaustion which is frequently mistaken for malignancy.[5]

Tiredness is a presenting feature of depression in adolescents, a serious problem that often goes unrecognised. Tonsillar - adenoidal hypertrophy may be large enough to compromise air exchange, particularly during sleep. Snoring may be a feature plus tiredness and lethargy in the waking state.

TIREDNESS IN THE ELDERLY

Elderly people tend to tire more quickly and recover more slowly and incompletely than younger ones. Sleep in older people is generally shorter in duration and of lesser depth, and they feel less refreshed and sometimes irritable on awakening.

Fatigue may be present as a result of emotional frustration. Whenever the prospect of gratification is small, a person tends to tire quickly and to remain so until something stimulating appears. Since the prospects for gratifying experience wane with the years, easy ‘fatigueability’ or tiredness is common in this age group.

BEREAVEMENT

Although a bereavement reaction is common and a normal human response that occurs at all ages, it is more frequently encountered in the elderly, with the loss of a spouse or a child (young or middle-aged!). Fatigue that occurs during the initial mourning period is striking and might represent a protective mechanism against intense emotional stress. With time, usually around 6 to 12 months, a compensated stage is reached, fatigue gradually abates, and the patient resumes normal activities as the conflicts of grief are gradually resolved. Freud pointed out the complexities of mourning as the bereaved person slowly adjusts to the loss of the loved one. In others, various symptoms persist as an ‘abnormal grief reaction’, including persistence of fatigue. Some factors that may lead to this include:

- unexpected death
- high dependence upon the dead person
- guilt feelings, especially in a love/hate relationship

Studies in general practice have shown that widows see their family doctors for psychiatric symptoms at three times the usual rate in the first 6 months after bereavement. The consultation rate for non-psychiatric symptoms also increases, by almost 50%.

Role of the family doctor

Following bereavement it is important to watch for evidence of depression, drug dependency, especially on alcohol, and suicidal tendencies. In cases of expected death, management should, if possible, start before the bereavement. Supportive care and ongoing counselling are very important.

BURNOUT

Definition

Burnout is a clinical syndrome with:

- emotional exhaustion
- depersonalisation of others
- lack of personal accomplishment

It is similar to stress-related depression but mood lowering is temporary and work-specific.

Patients sometimes claim that they feel ‘burnout’. Burnout can mean many things and include a whole constellation of psychogenic symptoms, such as exhaustion, boredom and cynicism, paranoia, detachment, heightened irritability and impatience, depression and psychosomatic complaints, such as headache and tiredness. Ellard[6] defines burnout as the syndrome that arises when a person who has a strong neurotic need to succeed in a particular task becomes confronted with the impossibility of success in that task. This seems a realistic explanation, but the important factor is to clarify the nature of the problem with care and determine whether the patient has a psychoneurotic disorder, such as hypomania, anxiety state or depression, or a personality disorder or simply unrealistic goals.

CHRONIC FATIGUE SYNDROME

This complex syndrome, which causes profound and persistent tiredness, is also referred to as myalgic encephalomyelitis, chronic neuromuscular viral syndrome,[7] postviral syndrome, chronic EBV syndrome, viral fatigue state, epidemic neuromyasthenia, neurasthenia, Icelandic disease, Royal Free disease and Tapanui disease. CFS is not to be confused with the tiredness and depression that follow a viral infection such as infectious mononucleosis, hepatitis or influenza. These postviral tiredness states are certainly common but resolve within 6 months or so.

Typical features of CFS (Figure 74.1):[8]

- extreme exhaustion (with minimal physical effort)
- headache or a vague ‘fuzzy’ feeling in the head
- aching in the muscles and legs
- poor concentration and memory
- hypersomnia or other sleep disturbance
- waking feeling tired
- emotional lability
- depressive-type illness
- arthralgia (without joint swelling)
- sore throat
- subjective feeling of fever (with a normal temperature)
- tender swollen lymph glands
- usually occurs between 20 and 40 years of age



Figure 74.1 Chronic fatigue syndrome: characteristic symptoms

Epidemiologically it has been related to coxsackie B virus infections.[8] The responsible organism is referred to as a slow virus infection by some authorities.[7]

In approximately two-thirds of patients the illness follows a clearly defined viral illness. However, no single virus has been consistently associated with the development of the syndrome, which is known to develop following a wide range of viral and non-viral infective illness. Immune system dysfunction with chronic overproduction of cytokines (e.g. interferon) is a possible pathogenetic mechanism.

Every family doctor probably has patients with this disorder and the syndrome has been observed in isolated endemics from time to time. Hickie et al.[2] found that only 0.3% of those with prolonged fatigue had been diagnosed with CFS by their family doctor.

There is no doubt that the syndrome is real in these patients. One of the major problems confronting clinicians is that there is no diagnostic test for this illness, so it remains a clinical diagnosis backed up by normal baseline investigations.

Diagnostic criteria for CFS have been published[9] (Table 74.4), which emphasise the positive clinical features of the syndrome and the chronicity of symptoms (greater than 6 months), in addition to the need for careful exclusion of alternative diagnoses by history, physical examination and laboratory investigation.

Table 74.4 Criteria for the diagnosis of chronic fatigue syndrome[9]

Fatigue Clinically evaluated, unexplained, persistent or relapsing fatigue persistent for 6 months or more, that: • is of new or definite onset • is not the result of ongoing exertion • is not substantially alleviated by rest • results in substantial reduction in previous levels of occupational, educational, social or personal activities *and*
Other symptoms Four or more of the following symptoms that are concurrent, persistent for 6 months or more and which did not predate the fatigue: • impaired short-term memory or concentration

- sore throat
- tender cervical or axillary lymph nodes
- muscle pain
- multi-joint pain without arthritis
- headaches of a new type, pattern, or severity
- unrefreshing sleep
- post-exertional malaise lasting more than 24 hour

Physical examination and investigation

Apart from mild pharyngeal infection, cervical lymphadenopathy or localised muscle tenderness, the physical examination is normal.

Investigations should be directed towards excluding possible diagnoses for that patient, such as chronic infection, autoimmune disorders, endocrine and metabolic disorders, primary neuromuscular disorders, malignancy and primary psychiatric disorders. The last mentioned is the most difficult of the differential diagnoses and psychiatric referral will often need to be considered.

Management

Patients who have CFS are really suffering and unhappy people, similar to those with fibromyalgia. They require considerable understanding and support. Symptoms last approximately 2½ years.

Management strategies include:[10]

- CFS recognition - explain that the illness is real but the cause unknown and tests are likely to be normal
- explanation and reassurance that the illness is usually self-limiting with no permanent complications; and that a slow steady improvement can be anticipated with most CSF patients returning to normal health
- provide continued psychological support
- review for diagnostic reappraisal (examine at least every 4 months)
- avoid telling patients they are depressed
- treat symptomatically - pain relief, consider NSAIDs and antidepressants if significant depression
- refer to counselling and support groups
- provide a realistic, regular, graduated exercise program
- reduce relevant stress factors (map a realistic living program)
- psychiatric referral if appropriate
- ask the patient to keep a diary of exercise/stress and symptom severity, in particular
- avoid long-distance travel, which is poorly tolerated

Cognitive therapy appears to help some patients, as do relaxation therapy, meditation, stress management and psychotherapy, where indicated.

The emphasis should be placed on caring, rather than curing, until a scientific solution is found. A systematic review has found that cognitive behaviour therapy administered by skilled therapists and exercise are beneficial. There is insufficient data or evidence to support the use of antidepressants, corticosteroids and dietary supplements. Prolonged rest and immunotherapy was unlikely to be beneficial.[11]

Practice tips

- Always consider underlying psychological distress, especially depressive disorder.
- Do not overlook a sleep disorder.
- Believe the patient's symptoms.
- Be careful of labelling a patient as having CFS.

- Key investigations are:
    - FBE and ESR
    - serum urea and creatinine
    - serum electrolytes, including calcium
    - liver function tests
    - blood sugar
    - thyroid function tests
    - microscopy and culture of urine

FIBROMYALGIA

The fibromyalgia syndrome (see Chapter 38) bears a clinical resemblance to CFS. Musculoskeletal pain is more prominent although tiredness (fatigue) and sleep disturbance are features. According to Schwenk,[12] fibromyalgia affects 5% of the American population with a peak age of 35 years (range 20 - 60) and a female:male ratio of 10:1. The management is similar to CFS but the prognosis less optimistic.

REFERENCES

1. Marinker M, Watter CAH. The patient complaining of tiredness. In: Cormack J, Marinker M, Morrell D eds. *Practice*. London: Kluwer Medical, 1982: Section 3.1.
2. Hickie IB et al. Sociodemographic and psychiatric correlates of fatigue in selected primary care settings. Med J Aust, 1996; 164: 585 - 8.
3. Jerrett WA. Lethargy in general practice. Practitioner, 1981; 225: 731 - 7.
4. French MA. The clinical significance of tiredness. Can Med Assoc J, 1960; 82: 665 - 71.
5. Tennessen WW. *Signs and Symptoms in Paediatrics*. Philadelphia: Lippincott, 1988: 37 - 40.
6. Ellard J. A note on burnout. Mod Med Aust, 1987; January: 32 - 5.
7. Simpson G. Persistent neuromuscular viral syndrome. Aust Fam Physician, 1987; 16: 978 - 81.
8. Blaiklock DA. The chronically tired patient. Patient Management, 1986; 10 (July): 89 - 96.
9. Fukuda K, Straus SE, Hickie I et al. The chronic fatigue syndrome: a comprehensive approach to its definition and study. International Chronic Fatigue Syndrome Study Group. Ann Intern Med, 1994; 121: 953 - 9.
10. Loblay R, Stewart G (Convenors). Chronic fatigue syndrome: clinical practice guidelines 2002. RACGP/MJA 2002; 176: Supplement.
11. Barton S. Chronic fatigue syndrome. In: *Clinical Evidence*. London: BMJ Publishing Group, 2001: 729 - 33.
12. Schwenk TL. Fibromyalgia and chronic fatigue syndrome: solving diagnostic and therapeutic dilemmas. Mod Med (USA), 1992; 60: 50 - 60.

CHAPTER 75. THE UNCONSCIOUS PATIENT

In whatever disease sleep is laborious, it is a deadly symptom; but if sleep does good, it is not deadly.

HIPPOCRATES

THE UNCONSCIOUS PATIENT

The state of arousal is determined by the function of the central reticular formation, which extends from the brain stem to the thalamus. Coma occurs when this centre is damaged by a metabolic abnormality or by an invasive lesion which compresses this centre. Coma is also caused by damage to the cerebral cortex.[1]

The word ‘coma’ is derived from the Greek koma, which is deep sleep. The deeply un-conscious patient is not in deep sleep. Coma is best defined as ‘lack of self-awareness’.[2]

The various levels of consciousness are summarised in Table 75.1; the levels vary from consciousness, which means awareness of oneself and the surroundings in a state of wake-fulness,[3] to coma, which is a state of unrousable unresponsiveness. Rather than using these broad terms in clinical practice it is preferable to describe the actual state of the patient in a sentence.

Table 75.1 The five conscious levels

	State	Clinical features	Simplified classification
↑Degree of consciousness↓	**1** Consciousness	Aware and wakeful	Awake
	2 Clouded consciousness	Reduced awareness and wakefulness ‘Alcohol effect’ Confusion Drowsiness	Confused
	3 Stupor	Unconscious Deep sleep-like state Arousal with vigorous stimuli	Responds to shake and shout
	4 Semicomatose	Unconscious (deeper) Responds only to painful stimuli (sternal rubbing with knuckles) without arousing	Responds to pain
	5 Coma	Deeply unconscious Unrousable and unresponsive	Unresponsive coma

KEY FACTS AND CHECKPOINTS

- Always consider hypoglycaemia or opioid overdose in any unconscious patient, especially of unknown background.
- If a patient is unconscious and cyanosed consider upper airway obstruction until proved otherwise.
- The commonest causes of unconsciousness encountered in general practice are syncope, expecially postural hypotension, concussion and cerebrovascular accidents (CVAs). The main causes are presented in Table 75.2.

Table 75.2 Main causes of loss of consciousness

Episodic causes - blackouts • Epilepsy • Syncope • Drop attacks • Cardiac arrhythmias (e.g. Stokes-Adams attacks) • Vertebrobasilar insufficiency • Psychogenic disorders, including hyperventilation • Breath-holding (children)
Coma (COMA provides a useful mnemonic for four major groups[1] of causes of unconsciousness) **C** ° CO_2 narcosis: respiratory failure **O** ° Overdose of drugs • alcohol • opioids • tranquillisers and antidepressants • carbon monoxide • analgesics • others **M** ° Metabolic • diabetes - hypoglycaemia - ketoacidosis • hypothyroidism • hypopituitarism • hepatic failure • renal failure (uraemia) • others **A** ° Apoplexy Supratentorial • intracerebral haemorrhage • haematoma: subdural or extradural • head injury • cerebral tumour • cerebral abscess Infratentorial (posterior fossa) • pressure from above

- cerebellar tumour
- brain-stem infarct/haemorrhage
- Wernicke's encephalopathy

Meningismus (neck stiffness)

- subarachnoid haemorrhage
- meningitis

Other

- encephalitis
- overwhelming infection

Trauma

- Do not allow the person who accompanies the unconscious patient to leave until all relevant details have been obtained.
- Record the degree of coma as a baseline to determine improvement or deterioration.

URGENT ATTENTION

The initial contact with the unconscious patient is invariably sudden and dramatic and demands immediate action which should take only seconds to minutes. The primary objective is to keep the patient alive until the cause is determined and possible remedial action taken.[3]

Examination	Action
Is the patient breathing? Note chest wall movement	If not, clear airway and ventilate.
Check pulse and pupils.	Perform cardiopulmonary resuscitation if necessary. Consider naloxone.
Is there evidence of trauma?	Consider extradural haematoma.
Is the patient hypoglycaemic? Evidence of diabetes (discs, etc.)	Consider glucometer estimation of blood sugar.
Are vital functions present yet immediate correctable causes eliminated?	Place in coma position.

HISTORY

A history can be obtained from relatives, friends, witnesses, ambulance officers or others. The setting in which the patient is found is important. Evidence of discs or cards identifying an illness such as diabetes or epilepsy should be searched for. Is there a known history of hypertension, heart disease, respiratory disease or psychiatric illness?

Questions to be considered[4]

- Is the patient diabetic?
  Does the patient have insulin injections?
  Has the patient had an infection?
  Has the patient been eating properly?

- Is drug overdose possible?
  Has the patient been depressed?
  Has the patient experienced recent stress or personal ‘mishaps’?
  Has the patient been on any medications?
- Is opioid usage possible in this patient?
  Are the presenting circumstances unusual?
- Is epilepsy possible?
  Was twitching in the limbs observed?
  Did the patient pass urine or faeces?
- Is head injury possible?
  Has the patient been in a recent accident?
  Has the patient complained of headache?
- Has a stroke or subarachnoid haemorrhage occurred?
  Has the patient a history of hypertension?
  Did the patient complain of a severe headache?
  Has the patient complained of weakness of the limbs?

EXAMINATION

General features requiring assessment:

- Breathing pattern:
  Cheyne-Stokes respiration (periodic respiration) ° cerebral dysfunction
  Ataxic respiration: shallow irregular respiration ° brain-stem lesion
  Kussmaul respiration: deep rapid hyperventilation ° metabolic acidosis
- Breath: characteristic odours may be a feature of alcohol, diabetes, uraemia and hepatic coma.
- Level of consciousness: degree of coma (Table 75.1); the Glasgow coma scale (Table 75.3) is frequently used as a guide to the conscious state.

Table 75.3 Glasgow coma scale

	Score
Eye opening (E)	
• Spontaneous opening	4
• To verbal command	3
• To pain	2
• No response	1
Motor response (M)	
• Obeys verbal command	6
Response to painful stimuli	
• Localises pain	5
• Withdraws from pain stimuli	4
• Abnormal flexion	3
• Extensor response	2
• No response	1
Verbal response (V)	

• Orientated and converses	5
• Disoriented and converses	4
• Inappropriate words	3
• Incomprehensible sounds	2
• No response	1

Coma score ° E + M + V

Minimum 3

Maximum 15

If 8 - 10: take care - monitor the airway

- Skin features: look for evidence of injection sites (drug addicts, diabetics) and snake bite marks, colour (cyanosis, purpura, jaundice, rashes, hyperpigmentation) and texture.
- Circulation.
- Temperature: consider infection such as meningitis and hyperpyrexia if raised and hypothermia (e.g. hypothyroidism) if low.
- Hydration: dehydration may signify conditions such as a high fever with infections, uraemia, hyperglycaemic coma.

Examination of the head and neck[3, 4]

The following should be considered:

- facial asymmetry
- the skull and neck - palpation for evidence of trauma and neck rigidity
- eyes, pupils and ocular fundi - look for constricted pupils in opioid overdose
- tongue
- nostrils and ears
- auscultation of the skull

Examination of the limbs

Consider:

- injection marks (drug addicts, diabetics)
- tone of the limbs by lifting and dropping (e.g. flaccid limbs with early hemiplegia)
- reaction of limbs to painful stimuli
- reflexes - tendon reflexes and plantar response

General examination of the body

This should include assessment of the pulses and blood pressure.

Urine examination

Catheterisation of the bladder may be necessary to obtain urine. Check the urine for protein, sugar and ketones.

DIAGNOSING THE HYSTERICAL ‘UNCONSCIOUS’ PATIENT

One of the most puzzling problems in emergency medicine is how to diagnose the unconscious patient caused by a conversion reaction (hysteria). These patients really experience their symptoms (as opposed to the pretending patient) and resist most normal stimuli, including painful stimuli.

Method

- Hold the patient's eye or eyes open with your fingers and note the reaction to light.
- Now hold a mirror over the eye and watch closely for pupillary reaction. The pupil should constrict with accommodation from the patient looking at his or her own image.

INVESTIGATIONS

Appropriate investigations depend on the clinical assessment. The following is a checklist.

- Blood tests

- All patients	blood sugar
	urea and electrolytes
- Selected patients	FBE
	blood gases
	liver function tests
	blood alcohol
	serum cortisol
	thyroid function tests
	serum digoxin

- Urine tests
  - A urine specimen is obtained by catheterisation.
  - Test for glucose and albumin.
  - Keep the specimen for drug screening.
- Stomach contents: aspiration of stomach contents for analysis.
- Radiology: CT scan or MRI are the investigations of choice (if available). If unavailable, X-ray of the skull may be helpful.
- Cerebrospinal fluid: lumbar puncture, necessary with neck stiffness, has risks in the comatose patient. A preliminary CT scan is necessary to search for coning of the cerebellum. If clear, the lumbar puncture should be safe and will help to diagnose subarachnoid haemorrhage and meningitis.
- Electroencephalograph
- ECG; look for ↑QT interval, etc.

BLACKOUTS - EPISODIC LOSS OF CONSCIOUSNESS

Episodic or transient loss of consciousness is a common problem. The important causes of blackout are presented in Table 75.4. The history is important to determine whether the patient is describing a true blackout or episodes of dizziness, weakness or some other sensation.

Table 75.4 Clinical features of blackouts

Cause	Precipitants	Subjective onset	Observation	Recovery
Vasovagal syncope	Posture stress haemorrhage micturition	warning of feeling ‘faint’, ‘distant’, ‘clammy, sweaty’	very pale sweating	gradual feels ‘terrible’ fatigue nausea
Respiratory	Cough weight-lifting trumpet playing	warning (feels faint)	pale	rapid

syncope				
Carotid sinus syncope	Carotid pressure (e.g. tight collar + turning neck) postendarterectomy	warning (feels faint)	pale	rapid
Cardiac syncope	Various	may be palpitations	pale	rapid may be flushing
Migrainous syncope	foods stress sleep deprivation	scotomas	pale	nausea and vomiting throbbing headache
Autonomic syncope	postural change	warning (feels faint)	pale	rapid
Epilepsy	Stress sleep deprivation alcohol withdrawal infection menstruation drug non-compliance	aura with complex partial seizures (CPS)	automatism (e.g. fidgeting, lip smacking) with CPS	slow confused

The clinical features of various types of blackouts are summarised in Table 75.4.

EPILEPSY

Epilepsy is the commonest cause of blackouts. There are various types, the most dramatic being the tonic - clonic seizure in which patients have sudden loss of consciousness without warning.

The typical features (in order) of a tonic - clonic convulsion are:

- aura (sensory or psychological feelings)
- initial rigid tonic phase (up to 60 seconds)
- convulsion (clonic phase) (seconds to minutes)
- mild coma or drowsiness (15 minutes to several hours) - post-ictal confusion

Associated features:

- cyanosis, then heavy ‘snoring’ breathing
- eyes rolling ‘back into head’
- ± tongue biting
- ± incontinence of urine or faeces

It should be noted that sphincter in-continence is not firmly diagnostic of epilepsy. In less severe episodes the patient may fall without observable twitching of the limbs.[5]

In atonic epilepsy, which occurs in those with tonic - clonic epilepsy, the patient falls to the ground and is unconscious for only a brief period.

SYNCOPE

In syncope there is a transient loss of consciousness but with warning symptoms and rapid return of alertness following a brief period of unconsciousness (seconds to 3 minutes).

Vasovagal ‘attack’

Relevant features of vasovagal or common faint (see Table 75.4):

- occurs with standing or, less commonly, sitting
- warning feelings of dizziness, faintness or true vertigo
- nausea, hot and cold skin sensations

- fading hearing or blurred vision
- sliding to ground (rather than heavy full-length fall)
- rapid return of consciousness
- pallor and sweating and bradycardia
- often trigger factors (e.g. emotional upset, pain)

The patient invariably remembers the onset of fainting. Most syncope is of the benign vasomotor type and tends to occur in young people, especially when standing still (e.g. choir boys). It is the main cause of repeated fainting attacks.

The treatment is to avoid precipitating causes (e.g. prolonged standing, especially in the sun) and bend forwards with the head down or lie down with premonitory signs. ‘Smelling salts’ (ammonium carbonate) can be carried and used in these circumstances.

Other forms of syncope

Micturition syncope

This uncommon event may occur after micturition in older men, especially during the night when they leave a warm bed and stand to void. The cause appears to be peripheral vasodilation associated with reduction of venous return from straining.

Cough syncope

Severe coughing can result in obstruction of venous return with subsequent blackout. This is also the mechanism of blackouts with breath-holding attacks.

Carotid sinus syncope

This problem is caused by pressure on a hypersensitive carotid sinus (e.g. in some elderly patients who lose consciousness when their neck is touched).

Effort syncope

Syncope on exertion is due to obstructive cardiac disorders, such as aortic stenosis and hypertrophic obstructive cardiomyopathy.

CHOKING

Sudden collapse can follow choking. Examples include the so-called ‘cafe coronary’ or ‘barbecue coronary’ when the patient, while eating meat, suddenly becomes cyanosed, is speechless and grasps the throat. This is caused by inhaling a large bolus of meat which obstructs the larynx. To avoid death, immediate relief of obstruction is necessary. An emergency treatment is the Heimlich manoeuvre, whereby the patient is grasped from behind around the abdomen and a forceful squeeze applied to try to eject the food. If this fails, the foreign body may have to be manually removed from the throat.

DROP ATTACKS

Drop attacks are episodes of ‘blackouts’ in which the patient suddenly falls to the ground and then immediately gets up again. They involve sudden attacks of weakness in the legs. Although there is some doubt about whether loss of consciousness has occurred, most patients cannot remember the process of falling. Drop attacks occur typically in middle-aged women and are considered to be brain-stem disturbances producing sudden changes in tone in the lower limbs. Other causes of drop attacks include vertebrobasilar insufficiency, Parkinson's disease and epilepsy.[5]

CARDIAC ARRHYTHMIAS

Stokes-Adams attacks and cardiac syncope are manifestations of recurrent episodes of loss of consciousness, especially in the elderly, caused by cardiac arrhythmias. These arrhythmias include complete heart block, sick sinus syndrome and ventricular tachycardia. The blackout is sudden with the patient falling straight to the ground without warning and without convulsive movements. The patient goes pale at first and then flushed.

Twenty-four-hour ambulatory cardiac monitoring may be necessary to confirm the diagnosis.

Patients with aortic stenosis are prone to have exercise-induced blackouts. Consider the prolonged QT interval syndrome in all age groups when the person presents with dizziness or blackouts.

VERTEBROBASILAR INSUFFICIENCY

Loss of consciousness can occur rarely with vertebrobasilar insufficiency (VBI) transient ischaemic attack. Typical preceding symptoms of VBI include dyspnoea, vertigo, vomiting, hemisensory loss, ataxia and transient global amnesia.

HYPOGLYCAEMIA

Hypoglycaemia can be difficult to recognise but must be considered as it can vary from a feeling of malaise and lightheadedness to loss of consciousness, sometimes with a convulsion. There are usually preliminary symptoms of hunger, sweating, shaking or altered behaviour. Hypoglycaemic attacks are usually related to diabetes and can occur with oral hypo-glycaemics as well as insulin.

HEAD INJURIES AND UNCONSCIOUSNESS

Some non-life threatening head injuries are sufficiently serious to cause significant loss of consciousness and retrograde amnesia. The clinical terms used to describe brain injury of concussion, contusion and laceration simply indicate minor to major degrees of a similar injury. Severe individual cases of the above can certainly result in fatal outcomes.

Concussion[6]

Concussion is a transient disturbance of neurological function induced by head injury and resulting in no persistent abnormal neurological signs. There may or may not be brief loss of consciousness. The old definition of loss of consciousness with eventual recovery applies only to more severe forms of con-cussion. The features of the various grades of concussion are shown in Table 75.5.

Table 75.5 Classification of concussion

Grade	Clinical features
Mild (grade 1)	Stunned or dazed Sensorium clears in < 60 seconds No post-traumatic amnesia ± Loss of consciousness
Moderate (grade 2)	Stunned or dazed Sensorium cloudy > 60 seconds

	Headache Amnesia < 60 minutes ± Loss of consciousness
Severe (grade 3)	Sensorium cloudy > 60 seconds Irritable Persistent headache Unsteady gait ± Loss of consciousness

Note: There is no such thing as delayed concussion or progressive deterioration due to concussion.

Post-concussion syndrome

Occasionally a person who has an episode of concussion has persistence of headaches and dizziness for a number of weeks. Poor memory and concentration and sluggish decision making indicate impaired mental capacity. Patients with this problem should be investigated with neuropsychological testing and CT scanning or MRI of the brain.

PSYCHOGENIC FACTORS

Psychogenic factors leading to blackouts represent a diagnostic dilemma, especially if occurring in patients with tonic - clonic epilepsy. If the attacks are witnessed by the practitioner then the possibility of functional origin can be determined.

Hysterical blackouts or fits are not uncommon and have to be differentiated from hyperventilation. It is unusual for hyper-ventilation to cause unconsciousness but it is possible to get clouding of consciousness, especially if the patient is administered oxygen.

Other features that suggest psychogenic rather than organic factors are:

- labile affect
- rapidly changing levels of consciousness
- well-articulated speech
- bizarre thought control

INITIAL MANAGEMENT OF THE UNCONSCIOUS PATIENT

The first principle of management of a person found unconscious is to keep the patient alive by maintaining the airway and the circulation. The basic management essentials are summarised in Table 75.6.

Table 75.6 Basic management essentials

Keep patient alive (maintain airway and circulation)
Get history from witnesses
Examine patient
Give 'coma cocktail' (TONG)
Take blood (for investigations)
CT scan (if diagnosis doubtful)

Before embarking on a secondary survey always consider giving the 'coma cocktail' (also called TONG[2]), which refers to the combination of:

Thiamine	100 mg IM or IV
Oxygenation	
Naloxone	0.1 - 0.2 mg IV
Glucose	i.e. 50 mL, 50% dextrose

The rapid administration of these agents should be considered for any patient[3, 7] with an altered level of consciousness because they may lessen or reverse metabolic insult to the brain.

In the presence of hypoventilation, con-stricted pupils[7] or circumstantial evidence of opioid use, naloxone (the specific opiate antagonist) should be given intravenously. If there is no response the patient should be intubated before further naloxone is given. Use a nasogastric tube to prevent acute gastric dilatation.

Use of flumazenil

Flumazenil is a specific benzodiazepine antagonist and may have an important use in the assessment of the unconscious patient. It can have a dramatic effect on benzodiazepine overdosage. After an initial dose of 0.2 mg IV, 0.3 - 0.5 mg boluses should be given every 1 - 2 minutes with caution until a response is observed.[7]

Opioid (heroin) overdose

A known overdose patient should be treated initially with both IV and IM naloxone:

- naloxone 0.4 mg IV (repeat in 3 minutes if necessary)
- naloxone 0.4 mg IM (to maintain cover)

Practice tips[2]

- The hypotensive patient is bleeding until proved otherwise.
- The presence of a head injury should not prevent rigorous resuscitation of the hypotensive patient.
- Always suspect cervical injury in the presence of patients who are victims of time-critical trauma.
- Tachypnoea is a sign of inadequate oxygenation and not a sign of central nerve damage.
- Always suspect opioid overdosage in the 'unknown' patient brought in with an altered conscious state.
- Consider administration of TONG - the 'coma cocktail'.

REFERENCES

1. Talley N, O'Connor S. *Clinical Examination* (3rd edn). Sydney: MacLennan & Petty, 1996: 414.
2. Wassertheil J. Management of neurological emergencies. Melbourne: Monash University, Update for GP's Course notes, 1996: 1 - 10.
3. Kumar PJ, Clark ML. *Clinical Medicine* (5th edn). London: Bailliere Tindall, 2003:1161 - 2.
4. Davis A, Bolin T, Ham J. *Symptom Analysis and Physical Diagnosis* (2nd edn). Sydney: Pergamon Press, 1990: 276 - 9.
5. Kincaid-Smith P, Larkins R, Whelan G. *Problems in Clinical Medicine*. Sydney: MacLennan & Petty, 1990: 156 - 75.
6. Brukner P, Khan K. *Clinical Sports Medicine* (2nd edn). Sydney: McGraw-Hill, 2001: 189 - 94.
7. McGirr J, McDonagh T. Management of acute poisoning. Curr Ther, 1995; 36(5): 51 - 9.

CHAPTER 76. URINARY DISORDERS

As men draw near the common goal, Can anything be sadder Than he who, master of his soul, Is
servant to his bladder?

ANONYMOUS 1938, SPECULUM

URINARY DISORDERS

Disturbances of micturition are a common problem in general practice with an annual incidence of about 20 per 1000 patients at risk.[1] Such disturbances include dysuria, frequency of micturition, difficulty or inability to initiate micturition, stress incontinence and haematuria. These symptoms are three times as common in women as in men.[1] The combination of dysuria and frequency is the most common of the symptoms with an incidence of about 14 per 1000 patients and a female to male ratio of 5 to 1.[1]

In children and the elderly the patient may complain of urinary incontinence unassociated with stress. However, with the exception of enuresis (Chapter 81), disturbances of micturition are uncommon in children.

DYSURIA AND FREQUENCY

Dysuria, or difficult and/or painful micturition, which is characterised mainly by urethral and suprapubic discomfort, indicates mucosal inflammation of the lower genitourinary tract, (i.e. the urethra, bladder or prostate). The passage of urine across inflamed mucosa causes pain. Frequency can vary from being negligible to extreme. It can be 'habit frequency' or associated with anxiety, which is typically long term and worse with stress and cold weather. In these conditions urinalysis is normal. Sometimes haematuria and systemic symptoms can accompany dysuria and frequency.

A summary of the diagnostic strategy model for dysuria is presented in Table 76.1.

Table 76.1 Dysuria: diagnostic strategy model

Q. Probability diagnosis
A. UTI esp. cystitis (female)
Urethritis
Urethral syndrome (female)
Vaginitis
Q. Serious disorders not to be missed
A. Neoplasia
• bladder
• prostate
• urethra
Severe infections
• gonorrhoea
• NSU
• genital herpes
Reiter's syndrome
Calculi (e.g. bladder)
Q. Pitfalls (often missed)
A. Menopause syndrome

Prostatitis

Foreign bodies in LUT

Acidic urine

Acute fever

Interstitial cystitis

Urethral caruncle/diverticuli

Vaginal prolapse

Obstruction

- benign prostatic hyperplasia
- urethral stricture
- phimosis
- meatal stenosis

Q. Seven masquerades checklist

A. Depression

Diabetes

Drugs

Anaemia -

Thyroid disorder -

Spinal dysfunction -

UTI

Q. Is the patient trying to tell me something?

A. Consider psychosexual problems; anxiety and hypochondriasis.

Key facts and checkpoints [1, 2]

- Strangury = difficult and painful micturition with associated spasm.
- Inflammation usually results in the frequent passage of small amounts of urine and a sense of urgency.
- Urethritis usually causes pain at the onset of micturition.
- Cystitis usually causes pain at the end of micturition.
- Suprapubic discomfort is a feature of bladder infection (cystitis).
- Vesicocolonic fistulas (e.g. prostatic cancer) cause severe dysuria, pneumaturia and foul-smelling urine.
- Dysuria and frequency are most common in women aged 15 to 44 years.
- They are four times more common in sexually active women.
- Vaginitis is an important cause and must be considered.
- Dysuria and discomfort is a common feature of postmenopausal syndrome, due to atrophic urethritis. The urethra and lower bladder are oestrogendependent.
- Unexplained dysuria could be a pointer to *Chlamydia* urethritis.
- Urinary infection and other disorders can be quite asymptomatic.

Is it really a urinary tract infection?

Although UTIs account for the majority of cases of dysuria in women it must be remembered that vaginitis and postmenopausal atrophic vaginitis can cause dysuria (Figure 76.1). Vaginitis is the most common cause of dysuria in the adolescent age group and is a relatively common cause of dysuria in family practice, estimated at around 15%. Postmenopausal oestrogen deficiency is estimated at 5 - 10%.[3] In the latter it is worthwhile prescribing oestrogen, either topically or systemically. Acute bacterial cystitis accounts for about 40% of causes of dysuria.



Figure 76.1 Relative causes of dysuria in women

The dysuria associated with vaginitis may be described as burning 'on the outside' with the discomfort usually felt at the beginning or end of micturition. If vaginitis is suspected, a pelvic examination should be carried out to inspect the genitalia and obtain swabs.[3]

The clinical approach

History

It is important to determine whether dysuria is really genitourinary in origin and not attributable to functional disorders, such as psychosexual problems. Disturbances of micturition are uncommon in the young male and if present suggest venereal infection.

Key questions

- Could you describe the discomfort?
- What colour is your urine?
- Does it have a particular odour?
- Have you noticed a discharge?
- If so, could it be sexually acquired?
- Do you find intercourse painful or uncomfortable (women)?
- Have you any fever, sweats or chills?

Examination

The general inspection and examination should include measurement of the basic parameters of pulse, temperature and blood pressure. The possibility of underlying kidney disease, especially in the presence of an obstructive component, should be kept in mind.

Abdominal palpation is important with a focus on the loins and suprapubic areas. The possibility of sexually transmitted diseases should also be considered and vaginal examination of the female and rectal and genital examination in the male may be appropriate. In the menopausal female a dry atrophic urethral opening, a urethral caruncle or urethral prolapse may give the clue to this important and neglected cause of dysuria.

Investigations

Basic investigations include:

- dipstick testing of urine
- microscopy and culture (midstream specimen of urine, or suprapubic puncture in children), and possibly urethral swabs or first pass urine for sexually transmitted infections

Further investigations depend on initial findings and referral for detailed investigation will be necessary if the primary cause cannot be found.

HAEMATURIA

Haematuria is the presence of blood in the urine and can vary from frank bleeding (macroscopic) to the microscopic detection of red cells. Haematuria can occur in a wide variety of disorders but a careful history and examination can often lead to the source of the bleeding and help with the selection of investigations. It is often a sign of a serious underlying disorder.

Key facts and checkpoints

- Macroscopic haematuria is the presence of blood visible to the naked eye. It is always abnormal except in menstruating women.
- Small amounts of blood (1 mL/1000 mL urine) can produce macroscopic haematuria.
- Microscopic haematuria is the presence of blood in the urine that can only be detected by microscopic or chemical methods.
- Microscopic haematuria includes the presence of red blood cells (RBC) > 8000 per mL of urine (phase contrast microscopy) or > 2000 per mL of urine (light microscopy) representing the occasional RBC on microscopic examination.
- Joggers and athletes engaged in very vigorous exercise can develop transient microscopic haematuria.
- Microscopic (asymptomatic haematuria) can be classified as either:
  - glomerular (from kidney parenchyma) *or*
  - non-glomerular (urinary tract)
- Common sources of macroscopic haematuria are the bladder, urethra, prostate and kidney.[4]
- Macroscopic haematuria occurs in 70% of people with bladder cancer and 40% with kidney cancer.[4]
- Common urological cancers that cause haematuria are the bladder (70%), kidney (17%), kidney pelvis or ureter (7%) and prostate (5%).[5]
- It is important to exclude kidney damage, so patients should have blood pressure, urinary protein and plasma creatinine levels measured as a baseline.
- All patients presenting with macroscopic haematuria or recurrent microscopic haematuria require judicious investigation, which may involve both radiological investigation of the upper urinary system and visualisation of the lower urinary system to detect or exclude pathology.
- The key radiological investigation is the intravenous urogram (pyelogram), unless there is a history of iodine allergy, severe asthma or other contraindications, when ultrasound is the next choice.

The clinical approach

History

Is it really haematuria? In many patients the underlying disorder may be suspected from a detailed enquiry about associated urinary symptoms. The presence of blood can be verified rapidly by microscopy so that red discolouration due to haemolysis or red food dye can be discounted.

The time relationship of bleeding is useful because, as a general rule, haematuria occurring in the first part of the stream suggests a urethral or prostatic lesion, while terminal haematuria suggests bleeding from the bladder. Uniform haematuria has no localising features.

The possibility of sexually acquired urethritis should be kept in mind. It is most unusual for haematuria to cause anaemia unless it is massive. Massive haematuria is a feature of radiation cystitis.

Painful haematuria is suggestive of infection, calculi or kidney infarction, while painless haematuria is commonly associated with infection, trauma, tumours or polycystic kidneys. Loin pain can occur as a manifestation of nephritis and may be a feature of bleeding in carcinoma of the kidney or polycystic kidney.

A drug history is relevant, especially with anticoagulants and cyclophosphamide. A diet history should also be considered.

It is worth noting that large prostatic veins, secondary to prostatic enlargement located at the bladder neck, may rupture when a man strains to urinate. A summary of the diagnostic strategy model for haematuria is presented in Table 76.2.

Table 76.2 Haematuria: diagnostic strategy model

Q. Probability diagnosis
A. Infection
• cystitis/urethrotrigonitis (female)
• urethritis (male)
• prostatitis (male)
Calculi - kidney, ureteric, bladder
Q. Serious disorders not to be missed
A. Cardiovascular
• kidney infarction
• kidney vein thrombosis
• prostatic varices
Neoplasia
• kidney tumour
• urothelial: bladder, kidney pelvis, ureter
• carcinoma prostate
Severe infections
• infective endocarditis
• kidney tuberculosis
• acute glomerulonephritis
• Blackwater fever
IgA nephropathy
Kidney papillary necrosis
Other kidney disease
Q. Pitfalls (often missed)
A. Urethral prolapse/caruncle
Pseudohaematuria (e.g. beetroot, porphyria)
Benign prostatic hyperplasia

Trauma: blunt or penetrating

Foreign bodies

Bleeding disorders

Exercise

Radiation cystitis

Menstrual contamination

Rarities

- hydronephrosis
- Henoch-Schönlein purpura
- bilharzia
- polycystic kidneys
- kidney cysts
- endometriosis (bladder)
- systemic vasculitides

Q. Seven masquerades checklist

Depression -

Diabetes -

Drugs cytotoxics anticoagulants

Anaemia -

Thyroid disorder -

Spinal dysfunction -

UTI

Q. Is this patient trying to tell me something?

A. Consider artefactual haematuria.

Key questions

- Have you had an injury such as a blow to the loin, pelvis or genital area?
- Have you noticed whether the redness is at the start or end of your stream or throughout the stream?
- Have you noticed any bleeding elsewhere, such as bruising of the skin or nose bleed?
- Have you experienced any pain in the loin or abdomen?
- Have you noticed any burning or frequency of your urine?
- Have you had any problems with the flow of your urine?
- Have you been having large amounts of beetroot, red lollies or berries in your diet?
- Could your problem have been sexually acquired?
- Have you been overseas recently?
- What is your general health like?
- Have you been aware of any other symptoms?
- Do you engage in strenuous sports such as jogging?
- Have you had any kidney problems in the past?

Physical examination

The general examination should include looking for signs of a bleeding tendency and anaemia, and recording the parameters of temperature, blood pressure and the pulse (see Figure 76.2). The heart should be assessed to exclude atrial fibrillation or infective endocarditis with emboli to the kidney, and the chest should be examined for a possible pleural effusion associated with perinephric or kidney infections.


General
fever
anaemia
purpura
Cardiovascular
heart murmurs
evidence SBE
atrial fibrillation
pleural effusion (kidney infections)
hypertension
splenomegaly
Urinary tract
kidney mass
bladder
• palpable
• tenderness
urethra
• caruncle
• blood
prostate
• inflammation
• mass

Figure 76.2 Haematuria: diagnostic strategy model

The abdomen should be examined for evidence of a palpable enlarged kidney or spleen. The different clinical findings for an enlarged left kidney and spleen are shown in Table 76.3. Kidney enlargement may be due to kidney tumour, hydronephrosis, or polycystic disease. Splenomegaly suggests the possibility of a bleeding disorder.

Table 76.3 Differences between spleen and left kidney on abdominal examination

	Spleen Left	kidney
Palpable upper border	Impalpable	Palpable
Movement with inspiration	Inferomedial	Inferior
Notch	Yes	No
Ballotable	No	Yes
Percussion	Dull	Resonant (usually)
Friction rub	Possible	Not possible

The suprapubic region should be examined for evidence of bladder tenderness or enlargement. In men the prostate should be examined rectally to detect benign o malignant enlargement or tenderness from prostatitis.

In women a vaginal examination should be performed to search for possible pelvic masses. The urethral meatus should be inspected to exclude a urethral caruncle or urethral prolapse.

Investigations

It is important to identify the cause, especially if a possible sequel is impaired kidney function.

- Urinalysis by dipstick testing (e.g. 'Hemastix' and its derivatives - affected by Vitamin C intake).
- Urine microscopy
  - formed RBCs in true haematuria
  - red cell casts indicate glomerular bleeding
  - deformed (dysmorphic) red cells indicate glomerular bleeding
- Urinary culture: early culture is important because of the common association with infection and consideration of early treatment with antibiotics. If tuberculosis is suspected, three early morning urines should be cultured for tubercle

bacilli.

- Urinary cytology: this test, performed on a urine sample, may be useful to detect malignancies of the bladder and lower tract but is usually negative with carcinoma of the kidney.
- Blood tests: appropriate screening tests include a full blood count, ESR and basic kidney function tests (urea and creatinine). If glomerulonephritis is suspected, antistreptolysin O titres and serum complement levels should be measured.
- Radiological techniques - available tests include:
  - intravenous urography (IVU; intravenous pyelogram) - the key investigation
  - ultrasound (less sensitive at detecting LUT abnormalities)
  - CT scanning
  - kidney angiography
  - retrograde pyelography
- Direct imaging techniques: these include urethroscopy, cystoscopy and ureteroscopy. In all patients, regardless of the IVU findings, cystoscopy is advisable.
- Kidney biopsy: indicated if glomerular disease is suspected, especially in the presence of dysmorphic red cells on microscopic examination.

Pseudohaematuria

Pseudohaematuria is red urine caused by pigments other than red blood cells that simply stain the culture red.

Causes include:

- anthrocyanins in food (e.g. beetroot, berries)
- red-coloured confectionery
- porphyrins
- free haemoglobin (e.g. haemoglobinuria)
- myoglobin (red-black colour)
- drugs (e.g. pyridium, phenolphthalein - alkaline urine)

Exercise haematuria

Exercise or sports haematuria is the passage of a significant number of red cells in the urine during or immediately after heavy exercise. It has been recorded in a wide variety of athletes, including swimmers and rowers. Dipstick testing is usually positive in these athletes. Despite the theory that it is largely caused by the posterior wall of the bladder impacting repetitively on the base of the bladder during running, there are other possible factors and glomerular disease must be excluded in the athlete with regular haematuria, especially if dysmorphic red cells are found on microscopy.

Artefactual haematuria

Macroscopic haematuria is a common presenting ploy of people with Munchausen's syndrome and pethidine addicts simulating kidney colic. If suspected, it is wise to get these people to pass urine in the presence of an appropriate witness before examining the urine.

Glomerulonephritis [6]

Glomerulonephritis means kidney inflammation involving the glomeruli. It can be simply classified into:

- nephritic syndrome: oedema + hypertension + haematuria
- nephrotic syndrome: oedema + hypoalbuminaemia + proteinuria
- asymptomatic kidney disease

The main causes of glomerulonephritis - nephritic syndrome are:

- IgA nephropathy (commonest)
- thin glomerular basement membrane disease

- post-streptococcal glomerulonephritis
- systemic vasculitis
- others

IgA nephropathy

Typically presents as haematuria in a young adult at the time of or within 1 - 2 days of a mucosal infection (usually throat, influenza or URTI) and persists for several days. Other presentations are as incidentally found microscopic haematuria or as previously unsuspected chronic kidney failure.

Due to deposition of IgA antibody complexes in the glomeruli it runs a variable course. There is no specific treatment to date.

Acute post-streptococcal glomerulonephritis

Typically seen in children (> 5 years), especially in Indigenous communities following GABHS throat infection or impetigo. Presents after a gap of 7 - 10 days or so.

Features

- Irritable lethargic sick child
- Haematuria: discoloured urine (‘eCoke’ urine)
- Periorbital oedema (may be legs, scrotum)
- Rapid weight gain (from oedema)
- Scanty urine output
- Hypertension → may be complications

Usual course

- Oliguria 2 days
- Oedema and hypertension 2 - 4 days
- Invariably resolves
- Good longterm prognosis

Diagnosis

- GABHS antigens
- Blood urea, creatinine, C_3 & $_4$ (complement),
- ASOT, DNAase B

Treatment

- Hospital admission
- Bed rest
- Strict fluid balance chart
- Daily weighing
- Penicillin (if GABHS +ve)
- Fluid restriction
- Low protein, high carbohydrate, low salt diet
- Antihypertensives and diuretics (as necessary)

Follow-up: monitor BP and kidney function. Regular urinalysis (microscopic haematuria may last for years)

> **DxT**: discoloured urine + periorbital oedema + oliguria = post-streptococcal glomerulonephritis

PROTEINURIA

Proteinuria is an important and common sign of kidney disease. The protein can originate from the glomeruli, the tubules or the LUT. Healthy people, however, do excrete some protein in the urine, which can vary from day to day and hour to hour; hence the value of collecting it over 24 hours. While proteinuria can be benign, it always requires further investigation. Important causes of proteinuria are presented in Table 76.4.

Table 76.4 Important causes of proteinuria

Transient • Contamination from vaginal secretions • Urinary tract infection • Pre-eclampsia These all require exclusion and follow-up.
Kidney disease • Glomerulonephritis • Nephrotic syndrome • Congenital tubular disease, e.g. - polycystic kidney - kidney dysplasia • Acute tubular damage • Kidney papillary necrosis, e.g. - analgesic nephropathy - diabetic papillary necrosis • Overflow proteinuria, e.g. - multiple myeloma • Systemic diseases affecting the glomeruli - diabetes mellitus - hypertension - SLE - malignancy - drugs (e.g. penicillamine, gold salts) - amyloid - vasculitides
No kidney disease • Orthostatic proteinuria • Exercise • Fever • Postoperative • Heart failure

Key facts and checkpoints[7]

- The amount of protein in the urine is normally less than 100 mg/24 hours.
- Greater than 150 mg protein/24 hours is abnormal in adults.
- Greater than 300 mg/24 hours is abnormal for children and adults.
- Proteinuria > 1 g /24 hours indicates a serious underlying disorder.
- If accompanied by dysmorphic haematuria or red cell casts, this tends to confirm glomerular origin.
- Routine dipstick testing will only detect levels greater than 300 mg/24 hours and thus has limitations.
- In diabetics, microalbuminuria is predictive of nephropathy and an indication for early blood pressure treatment.

If proteinuria is confirmed on repeated dipstick testing it should be measured more accurately by measuring daily albumin excretion with a 24 hour urine or the albumin creatinine ratio (ACR), which is preferred. High values require referral for investigation. The minimum investigations are microurine and assessment of kidney function (eGFR). Nephrotic range proteinuria (> 3 g/24 hours) is due to one or other form of glomerulonephritis in over 90% of patients.[6] Possible contamination from vaginal secretions or from a low UTI needs to be excluded.

Orthostatic proteinuria

Orthostatic proteinuria is the presence of significant proteinuria after the patient has been standing but is absent from specimens obtained following recumbency for several hours, such as an early morning specimen.

It occurs in 5 - 10% of people,5 especially during their adolescent years. In the majority it is of no significance and eventually disappears without the development of significant kidney disease. However, in a small number the proteinuria can foreshadow serious kidney disease.

Diabetic microalbuminuria

The presence of protein in the urine is a sensitive marker of diabetic nephropathy, so regular screening for microalbuminuria in diabetics is regarded as an important predictor of nephropathy and other possible complications of diabetes. Dipstick testing for microalbuminuria is now available but more accurate measurement can be performed with radioimmunoassay techniques. The use of ACE inhibitors at the microalbuminuria stage may slow the development of overt nephropathy. The gold standard is a 24-hour collection.

Consequences of proteinuria

While proteinuria is usually simply a marker of kidney disease, heavy proteinuria in excess of 3 g/ 24 hours may have severe clinical consequences, including oedema, intravascular volume depletion, venous thromboembolism, hyperlipidaemia and malnutrition.

Minimal change glomerulonephritis is the commonest cause of the nephrotic syndrome in childhood and accounts for about 30% of adult nephrotic syndrome.[7] It is steroid responsive.

Nephrotic syndrome [6, 7]

DxT: proteinuria + generalised oedema + waxy pallor = nephrotic syndrome

Features

- Proteinuria > 3 g/day (3 - 4+ on dipstick)
- Generalised oedema
- Hypoalbuminaemia < 25 g/L
- Hypercholesterolaemia > 4.5 mmol/
- Waxy pallor
- Normal BP
- Dyspnoea

Predisposes to sepsis (e.g. peritonitis, pyelonephritis, thromboembolism).

Causes

- 1 in 3 (approx.)
    - systemic kidney disease (e.g. diabetes)
- 2 in 3 (approx.)
    - idiopathic nephrotic syndrome (based on kidney biopsy)
        - minimal change disease (commonest)
        - focal glomerular sclerosis
        - membranous nephropathy
        - membranoproliferate glomerulonephritis

Treatment

- Referral
- Bed rest
- Diet: low fluid, high protein, low salt
- Diuretics
- Prednisolone
- Phenoxymethylpenicillin
- Aspirin

URINARY INCONTINENCE

Definitions

Urinary incontinence The involuntary loss of urine during the day or night.

Nocturnal enuresis (or bed-wetting) Involuntary urine loss during sleep.

Urge incontinence An urgent desire to void followed by involuntary loss of urine.

Overactive bladder (detrusor instability) The commonest cause of urge incontinence; synonymous with an irritable or unstable bladder; characterised by involuntary bladder contractions, resulting in a sudden urge to urinate.

Stress incontinence The involuntary loss of urine on coughing, sneezing, straining or lifting, or any factor that suddenly increases intra-abdominal pressure.

Voiding dysfunction Includes urinary difficulties, detrusor instability and overflow incontinence.

Functional incontinence Loss of urine secondary to factors extrinsic to the urinary tract.

A summary of the types of incontinence and their causes is presented in Table 76.5.

Table 76.5 Types of incontinence and their implied causes

Type of incontinence	Likely cause
Simple stress incontinence (with cough/sneeze)	Sphincter incompetence
Urge incontinence Giggle incontinence Stress and urge incontinence Enuresis	Unstable bladder, with or without sphincter weakness

Complex stress incontinence (with exercise) Quiet dribble incontinence	Sphincter incompetence and unstable bladder or overflow
Continuous leakage	Fistula, ectopic ureter, patulous urethra
Reflex incontinence	Neuropathic bladder

The basic requirements for continence are:

- adequate central and peripheral nervous function
- an intact urinary tract
- a compliant stable bladder
- a competent urethral sphincter
- efficient bladder emptying

Causes of incontinence (a mnemonic)[8]

D delirium

I infection of urinary tract

A atrophic urethritis

P pharmacological (e.g. diuretics)

P psychological (e.g. acute distress)

E endocrine (e.g. hypercalcaemia)

E environmental (e.g. unfamiliar surrounds)

R restricted mobility

S stool impaction

S sphincter damage or weakness

Female urinary incontinence

Urinary leakage affects at least 9% of women aged 15 - 64 years: a conservative estimate. Successful treatment depends on accurate assessment of the LUT storage and emptying functions. The most common contributing factor is weakness of the pelvic floor muscles.

Assessment

The basic assessment of the incontinent patient requires a careful history and examination, exclusion of infection and keeping of a micturition or bladder chart. Use of a severity index questionnaire is very helpful. Drugs that adversely affect urinary function are presented in Table 76.6. Investigations may be required to dispel doubt about the diagnosis or to exclude intravesical or kidney disorders: these include cystometry, uroflowmetry, cysto-urethroscopy, micturating cystourethrogram, IVU and also residual volume (> 100 mL is abnormal).

Table 76.6 Drugs that can cause or aggravate incontinence

Antihypertensive/vasodilator drugs → stress incontinence • ACE inhibitors

- phenoxybenzamine (Dibenyline)
- prazosin
- labetalol

Bladder relaxants → overflow incontinence

- anticholinergic agents
- tricyclic antidepressants

Bladder stimulants → urge incontinence

- cholinergic agents
- caffeine

Sedatives → urge incontinence

- antidepressants
- antihistamines
- antipsychotics
- hypnotics
- tranquillisers

Others → urge incontinence

- alcohol
- loop diurectics (e.g. frusemide), other diuretics
- lithium

Management approach

1. Exclude UTI and drug causes.
2. Is it stress incontinence?
    - key symptoms: involuntary loss with coughing, jumping, etc.
    - demonstrable (e.g. patient coughs when standing with full bladder)

Treatment

- Weak pelvic floor - exercises
- Obesity - weight reduction
- Menopause - HRT/vaginal oestrogen creams
- Chronic cough - physiotherapy

If urodynamic studies of LUT function show genuine stress incontinence (GSI) due to urethral sphincter weakness, consider surgery (e.g. suprapubic urethral suspension - better than vaginal repair)

The severity index questionnaire

- How often do you experience urine leakage? 0 = never, 1 = less than once a month, 2 = one or several times a month, 3 = one or several times a week, 4 = every day and/or night
- How much urine do you lose each time? 1 = drops or little, 2 = more The total score is the score for the first question multiplied by the score for the second question. 0 = dry, 1 - 2 = slight, 3 - 4 = moderate, 6 - 8 = severe

1. 3 Is it urge incontinence?
    - urge symptoms prominent
    - no residual urine

Treatment

- neurological signs → neurologist
- abnormal voiding pattern → bladder retraining (e.g. void more urine less frequently)

2. Is it voiding dysfunction?
    - symptoms of voiding difficulty (e.g. frequency, urgency, nocturia, incomplete emptying)
    - large residual urine

Treatment

- Neurological signs → neurologist
- Gynaecological cause (e.g. pelvic mass) → gynaecologist
- If bladder atony → anticholinergic drugs
- Intravescical treatment:
    - oxybutrin
    - botalinum toxin
    - others, e.g. capsaicin
- May require catheterisation

Anticholinergic drugs[9]

These may be worth a trial for bladder instability or voiding dysfunction:

- propantheline 15 mg (o) bd or tds
- oxybutynin 2.5 - 5 mg (o) bd or tds
- tolterodine 2 mg (o) bd
- imipramine 10 - 75 mg (o) nocte

The future:

- neuromodulation e.g. sacral nerve

Pelvic floor exercises

- The mainstay of treatment of most problems,→especially GSI (e.g. 40 ‘squeezes’ daily).
- 75% improved and 25% cured.
- Best in motivated young women with→bladder GSI.
- At least 3 months trial with supervision→(physiotherapist or continence nurse→adviser).

Basic techniques

- Advise the patient to pull up her pelvic muscles to imagine herself stopping passing urine (or controlling diarrhoea) and hold the ‘squeeze’ for a count of 10. Repeat many times daily. Refer to Patient Education.[10]
- Bladder retraining[11] includes getting the patient to delaying micturition for 10 - 15 minutes whenever she wishes to void.

BLADDER DYSFUNCTION (IN WOMEN DURING NIGHT)

Women with urethral syndrome constantly wake at night with urge to micturate but produce only a small dribble of urine.

- Instruct patient to perform a pelvic tilt exercise by balancing on upper back, lift ing her pelvis with knees flexed and holding position for 30 seconds
- Squeeze pelvic floor inwards (as though holding back urine or faeces)
- Repeat a few times

UTEROVAGINAL PROLAPSE[12]

Uterovaginal prolapse is very common, affecting 50% of parous women. The main complaint is of ‘heaviness’ in the vagina and a sensation of ‘something coming down’. Relevant symptoms that are of considerable distress for the patient and, depending on the type of prolapse, include voiding difficulties, urinary stress incontinence, faecal incontinence, incomplete rectal emptying and recurrent cystitis. Backache is a common associated symptom, usually relieved by lying down.

Classification of prolapse (Figure 76.3)



Figure 76.3 Uterovaginal prolapse

- Cystocele - bladder descends into vagina
- Urethrocele - urethra bulges into vagina
- Rectocele - rectum protrudes into vagina
- Enterocele - loop of small intestine bulges into vagina (usually posterior wall)
- Uterine - uterus and cervix descend toward vaginal introitus:
  - first degree - cervix remains in vagina
  - second degree - cervix protrudes on coughing/straining
  - third degree (procidentia) - uterus lies outside vagina

Examination

This is best performed with women in the left lateral position using a Sims’ speculum or posterior blade of the Grave's speculum. Ask the patient to cough or bear down (several times) - observe anterior, posterior and lateral vaginal walls and descent of cervix.

Management

As a rule asymptomatic prolapse does not need invasive treatment, just basic reassurance and education, including pelvic floor exercises (see p. 834). Consider referral to a physiotherapist. Lifestyle measures include optimal nutrition, weight loss if obese, smoking cessation and exercise. Aggravating comorbidities such as constipation, menopause/atrophic vaginitis and COPD require optimal treatment. Consider topical oestrogens.

Prevention

Promote optimal obstetric management, especially postpartum exercises, lifelong pelvic floor exercises, ideal weight and sensible bladder and bowel function.

Ring pessaries

Pessaries are an option for those who are poor anaesthetic risks, too frail for surgery, don't want surgery, are young and have not completed their family or are awaiting surgery. Each patient needs to be fitted individually with the correct-sized pessary. Topical oestrogens will improve comfort. The pessary needs to be cleaned or changed every 4 - 6 months.

Surgery

Refer to a gynaecological surgeon if the patient who is fit for surgery has symptomatic prolapse that warrants surgery especially with associated incontinence and recurrent UTIs. The principles of reconstructive pelvic surgery are to:

- reposition pelvic structures to normal anatomical relationships
- restore and maintain urinary and/or faecal continence
- maintain coital function
- correct coexisting pelvic pathology

Options include repair procedures (vaginally, sometimes abdominally, per laparoscopy), colpo/ vaginal suspension, hysteroscopy and hysterectomy (vaginal or abdominal).

REFERENCES

1. Cormack J, Marinker M, Morrell D. Practice. *A Handbook of Primary Medical Care*. London: Kluwer-Harrap Handbooks, 1980: 3.51: 1 - 10.
2. Kincaid-Smith P, Larkins R, Whelan G. Problems in Clinical Medicine. Sydney: MacLennan & Petty, 1990: 105 - 8.
3. Sloane PD, Slatt PD, Baker RM. *Essentials of Family Medicine*. Baltimore: Williams & Wilkins, 1988: 169 - 74.
4. Walsh D. Symptom Control. Boston: Blackwell Scientific Publications, 1989: 229 - 33.
5. George C. How to treat haematuria and proteinuria. Aust Dr Weekly, 1991; 15 March: I - VIII.
6. Faull R. How to treat: glomerulonephritis. Aust Doctor, 2002; 8 February: I - VIII.
7. Thomson N. Managing the patient with proteinuria. Curr Ther, 1996; 9: 7 - 28.
8. Jayasuriya P. How to treat: urinary incontinence. Aust Doctor, 2001; 11 May: I - VIII.
9. Haylen B. Advances in incontinence treatment. Aust Doctor, 1999; 10 September: 66 - 70.
10. Murtagh J. Incontinence of urine. In: Patient Education (4th edn). Sydney; McGraw-Hill, 2005: 70.
11. Benness C. Female urinary incontinence. Medical Observer, 2004; 25 June: 31 - 3.
12. Benness C. Uterovaginal prolapse. Medical Observer, 2005; 3 June: 29 - 3s1.

CHAPTER 77. VISUAL FAILURE

> All those, therefore, who have cataract see the light more or less, and by this we distinguish cataract from amaurosis and glaucoma, for persons affected with these complaints do not perceive the light at all.
>
> PAUL OF AEGINA (615-690)

VISUAL FAILURE

The commonest cause of visual dysfunction is a simple refractive error. However, there are many causes of visual failure, including the emergency of sudden blindness, a problem that requires a sound management strategy. Apart from migraine, virtually all cases of sudden loss of vision require urgent treatment.

The 'white' eye or uninflamed eye presents a different clinical problem from the red or inflamed eye.[1] The 'white' eye is painless and usually presents with visual symptoms and it is in the 'white' eye that the majority of blinding conditions occur.

CRITERIA FOR BLINDNESS AND DRIVING

This varies from country to country. The WHO defines blindness as 'best visual acuity less than 3/60', while in Australia eligibility for the blind pension is 'bilateral corrected visual acuity less than 6/60 or significant visual field loss'; e.g. a patient can have 6/6 vision but severely restricted fields caused by chronic open-angle glaucoma. The minimum standard for driving is 6/12 (Snellen's system).

KEY FACTS AND CHECKPOINTS

- The commonest cause of blindness in the world is trachoma.
- In Western countries the commonest causes are senile cataract, glaucoma, age-related macular degeneration, trauma and the retinopathy of diabetes mellitus.[2]
- The commonest causes of sudden visual loss are transient occlusion of the retinal artery (amaurosis fugax) and migraine.[3]
- 'Flashing lights' are caused by traction on the retina and may have a serious connotation: the commonest cause is vitreoretinal traction, which is a classic cause of retinal detachment.
- The presence of floaters or 'blobs' in the visual fields indicates pigment in the vitreous: causes include vitreous haemorrhage and vitreous detachment.
- Posterior vitreous detachment is the commonest cause of the acute onset of floaters, especially with advancing age.
- Retinal detachment has a tendency to occur in short-sighted (myopic) people.
- Suspect a macular abnormality where objects look smaller or straight lines are bent or distorted.

THE CLINICAL APPROACH

History

The history should carefully define the onset, progress, duration, offset and the extent of visual loss. An accurate history is important because a longstanding visual defect may only just have been noticed by the patient, especially if it is unilateral. Two questions need to be answered:

- Is the loss unilateral or bilateral?
- Is the onset acute, or gradual and progressive?

The distinction between central and peripheral visual loss is useful. Central visual loss presents as impairment of visual acuity and implies defective retinal image formation (through refractive error or opacity in the ocular media) or macular or optic nerve dysfunction. Peripheral field loss is more subtle, especially when the onset is gradual, and implies extramacular retinal disease or a defect in the visual pathway.

It is important to differentiate the central field loss of macular degeneration from the hemianopia of a CVA.

A drug history is very important (Table 77.1). Treatment for tuberculosis with ethambutol or treatment with quinine/chloroquine has to be considered as these drugs are oculotoxic. The family history is relevant for diabetes, migraine, Leber's hereditary optic atrophy, amaurotic familial idiocy and retinitis pigmentosa.

Table 77.1 Visual disorders associated with drugs

Disorder	Drug
Corneal opacities	Amiodarone
Precipitating of acute narrow angle glaucoma	Hydroxychloroquine Chlorpromazine Vitamin D Indomethacin Chlorpropamide Mydriatic drops
Refractive changes	Tricyclics Antihistamines Thiazides
Lens opacities	Corticosteroids
Retinopathy	Phenothiazines Hydroxychloroquine
Papilloedema (secondary to benign intracranial hypertension)	Chloroquine Thioridazine (other phenothiazines less commonly) Tamoxifen Oral contraceptives
Optic neuropathy	Corticosteroids Tetracyclines Nalidixic acid Vitamin A Ethanol Tobacco Ethambutol Disulfiram

Reprinted from Central Nervous System: Clinical Algorithms, published by the Br Med J, with permission

Questions directed to specific symptoms

- Presence of floaters → normal ageing (especially ≥ 55 years) with posterior vitreous detachment or may indicate haemorrhages or choroiditis
- Flashing lights → normal ageing with posterior vitreous detachment or indicates traction on retina ? retinal

detachment

- Coloured haloes around lights → glaucoma, cataract
- Zigzag lines → migraine
- Vision worse at night or in dim light → retinitis pigmentosa, hysteria, syphilitic retinitis
- Headache → temporal arteritis, migraine, benign intracranial hypertension
- Central scotomata → macular disease, optic neuritis
- Pain on moving eye → retrobulbar neuritis
- Distortion, micropsia (smaller), macropsia (larger) → macular degeneration
- Visual field loss:
  - central loss - macular disorder
  - total loss - arterial occlusion

It is worth noting that if a patient repeatedly knocks into people and objects on a particular side (including traffic accidents), a bitemporal or homonymous hemianopia should be suspected.

Diseases/disorders to exclude or consider

- Diabetes mellitus
- Giant cell (temporal) arteritis
- Hypopituitarism (pituitary adenoma)
- Cerebrovascular ischaemia/carotid artery stenosis (emboli)
- Multiple sclerosis
- Cardiac disease (e.g. arrhythmias, and SBE - emboli)
- Anaemia (if severe can cause retinal haemorrhage and exudate)
- Marfan syndrome (subluxated lenses)
- Malignancy (the commonest cause of eye malignancy is melanoma of the choroid)

Examination

The same principles of examination should apply as for the red eye. Testing should include:

- visual acuity (Snellen chart) - with pinhole testing
- pupil reactions, to test afferent (sensory) responses to light
- confrontation fields (using a red pin)
- colour vision
- Amsler grid (or graph paper)
- fundus examination with dilated pupil (ophthalmoscope), noting
  - the red reflex
  - appearance of the retina, macula and optic nerve
- tonometry

General examination

General examination should focus on the general features of the patient, the nervous system, endocrine system and cardiovascular system.

Perimetry

Various defects in the visual fields are depicted in Figure 77.1.


Typical defects
in
visual fields
L
R
1
unilateral blindness
2
bitemporal hemianopia
3
R homonymous hemianopia
4
quadrantic field defect
5
R homonymous hemianopia
Causes of sudden blindness
vitreous haemorrhage
retinal detachment
disciform macular degeneration
papilloedema
central retinal artery occlusion
central retinal vein occlusion
optic neuritis
optic nerve
1
optic chiasma
2
3
optic
tract
geniculate body
optic radiation
5
occipital cortex

Figure 77.1 Diagrammatic representation of important causes of sudden painless loss of vision (right side) and typical defects in the visual fields (left side)

Investigations

Depending on the clinical examination the following tests can be selected to confirm the diagnosis:

- blood tests
    - full blood ? anaemia, lead poisoning, leukaemia
    - ESR ? temporal arteritis
    - blood sugar ? diabetes mellitus
- temporal artery biopsy ? temporal arteritis
- CT/MRI scan ? CVA, optic nerve lesions, space occupying lesions
- formal perimetry and Bjerrum screen
- fluorescein angiography ? retinal vascular obstruction, diabetic retinopathy
- visual evoked responses ? demyelinating disorders
- carotid Doppler ultrasound

VISUAL FAILURE IN CHILDREN

There are long lists of causes for visual failure or blindness in children. An approximate order of frequency of causes of blindness in children is cortical blindness, optic atrophy, choroidoretinal degeneration, cataracts and retinopathy of prematurity. Almost half the Figure. 77.1 Diagrammatic representation of important causes of sudden painless loss of vision (right side) and typical defects in the visual fields (left side) causes of blindness are genetically determined, in contrast to the nutritional and infective causes which predominate in third world countries.[4] About 3% of children will fail to develop proper vision in at least one eye.

The eyes of all babies should be examined at birth and at 6 weeks.

Amblyopia

Amblyopia is defined as a reduction in visual acuity due to abnormal visual experience in early childhood. It is the main reason for poor unilateral eyesight until middle age and is usually caused by interference with visual development during the early months and years of life.

The common causes are:

- strabismus

- large refractive defect, especially hypermetropia
- congenital cataract

Principles of management[5]

- Most cases are treatable.
- Early diagnosis and intervention is fundamental to achieving useful vision.
- No child is too young to have the visual system assessed.
- The good eye should be patched in order to utilise the affected eye.
- Remove a remedial cause such as strabismus.
- Correct any refractive error, usually by prescription of glasses.

Some important guidelines in children

Referral

Refer if any of the following are present in infants:

- nystagmus
- a wandering eye
- a lack of fixation or following of movements
- photophobia
- opacities (seen with ophthalmoscope set on + 3, held 30 cm from baby's eye)
- delayed development.

Strabismus

- The two serious squints are the constant and alternating ones, which require early referral. Transient squint and latent squint (occurs under stress e.g. fatigue) usually are not a problem.
- Always refer children with strabismus (squint) when first seen to exclude ocular pathology such as retinoblastoma, congenital cataract and glaucoma, which would require emergency surgery.
- Children with strabismus (even if the ocular examination is normal) need specialist management because the deviating eye will become amblyopic (a lazy eye with reduced vision i.e. ‘blind’ if not functioning by 7 years of age). The younger the child, the easier it is to treat amblyopia; it may be irreversible if first detected later than school age. Surgical correction of a true squint is preferred at 1 - 2 years of age. (see also chapter 82)

Cataracts

Children with suspected cataracts must be referred immediately; the problem is very serious as the development of vision may be permanently impaired (amblyopia).[2] Cataracts are diagnosed by looking at the red reflex and this should be a routine part of the examination of a young child. Common conditions causing cataracts are genetic disorders and rubella but most causes are unknown. Rarer conditions, such as galactosaemia, need to be considered.

Refractive errors

Refractive errors, with the error greater in one eye, can cause amblyopia. Detection of re-fractive errors is an important objective of screening.

Retinoblastoma

Retinoblastoma, although rare, is the com-monest intraocular tumour in childhood. It must be excluded in any child presenting with a white pupil. Such children also have the so-called ‘cat’s eye reflex’. In 30% of patients the condition is bilateral with an autosomal dominant gene being responsible.

VISUAL FAILURE IN THE ELDERLY

Most patients with visual complaints are elderly and their failing vision affects their perception of the environment and their ability to communicate effectively. Typical problems are cataracts, vascular disease, macular degen-eration, chronic simple glaucoma and retinal detachment. Retinal detachment and diabetic retinopathy can occur at any age, although they are more likely with increasing age. Macular degeneration in its various forms is the commonest cause of visual deterioration in the elderly. For the elderly with cataracts the decision to operate depends on the patients' vision and their ability to cope. Most patients with a vision of 6/18 or worse in both eyes usually benefit from cataract extraction, but some can cope with this level of vision and rely on a good, well-placed (above and behind) reading light.[6]

Sudden loss of vision in the elderly is suggestive of temporal arteritis or vascular embolism, so this problem should be checked.

FLOATERS AND FLASHES

When the vitreous gel shrinks as part as the normal ageing process, it fugs on the retina (rods and cones) causing flashing lights. When the gel separates from the retina, floaters (which may appear as dots, spots or cobwebs) are seen. Floaters are more commonly seen with age, but are also more common in people who are myopic or who have had eye surgery such as removel of cataracts. It is important to consider retinal detachment but if floaters remain constant there is little cause for concern. The appearance of a fresh onset of flashes or floaters is if concern.

REFRACTIVE ERRORS

Indistinct or blurred vision is most commonly caused by errors of refraction.

In the normal eye (emmetropia) light rays from infinity are brought to a focus on the retina by the cornea (contributing about two-thirds of the eye's refractive power) and the lens (one-third). Thus, the cornea is very important in refraction and abnormalities such as keratoconus may cause severe refractive problems.[6]

The process of accommodation is required for focusing closer objects. This process, which relies on the action of ciliary muscles and lens elasticity, is usually affected by ageing, so that from the age of 45 close work becomes gradually more difficult (presbyopia).[6]

The important clinical feature is that the use of a simple ‘pinhole’ in a card will usually improve blurred vision or reduced acuity where there is a refractive error only.[1]

Myopia (short-sightedness)

This is usually progressive in the teens. Highly myopic eyes may develop retinal detachment or macular degeneration.

Management

- Glasses with a concave lens
- Contact lenses
- Consider radial keratotomy or excimer laser surgery

Hypermetropia (long-sightedness)

This condition is more susceptible to closed-angle glaucoma. In early childhood it may be associated with convergent strabismus (squint). The spectacle correction alone may straighten the eyes. It is mostly overcome by the accommodative power of the eye, though it may cause reading difficulty. Typically, the long-sighted person needs reading glasses at about 30 years.

Presbyopia

There is a need for near correction with loss of accommodative power of the eye in the 40s.

Astigmatism

This creates the need for a corrective lens that is more curved in one meridian than another because the cornea does not have even curvature. If uncorrected, this may cause headaches of ocular origin. Conical cornea is one cause of astigmatism.

Pinhole test

The pinhole reduces the size of the blur circle on the retina in the uncorrected eye. A pinhole acts as a universal correcting lens. If visual acuity is not normalised by looking through a card with a 1 mm pinhole, then the defective vision is not solely due to a refractive error. The pinhole test may actually help to improve visual acuity with some cataracts. Further investigation is mandatory.

CATARACTS

The term ‘cataract’ describes any lens opacity. The symptoms depend on the degree and the site of opacity. Cataract causes gradual visual loss with normal direct pupillary light reflex.

The prevalence of cataracts increases with age: 65% at age 50 to 59, and all people aged over 80 have opacities.[3] Significant causes of cataracts are presented in Table 77.2 and causes of progressive visual loss in Table 77.3.

Table 77.2 Causes of cataracts

Advancing age
Diabetes mellitus
Steroids (topical or oral)
Radiation: long exposure to UV light
Trauma
Uveitis
Dystrophia myotonia

Table 77.3 Progressive bilateral visual loss

Globe	Chronic glaucoma
	Senile cataracts
Retina	Macular degeneration
	Retinal disease
	• diabetic retinopathy
	• retinitis pigmentosa
	• choroidoretinitis
Optic nerve	Optic neuropathies
	Optic nerve compression (e.g. aneurysm, glioma)
	Toxic damage to optic nerves
Optic chiasma	Chiasmal compression: pituitary adenoma, craniopharyngioma, etc.
Occipital cortex	Tumours
	Degenerative conditions

Note: Unilateral causes (e.g. cataract, refractive errors, uveitis, glaucoma, progressive optic atrophy and tumours) can affect the second eye.

Typical symptoms:

- reading difficulty
- difficulty in recognising faces
- problems with driving, especially at night
- difficulty with television viewing
- reduced ability to see in bright light
- may see haloes around lights

The type of visual distortion seen by patients is illustrated in Figure 77.2.



Figure 77.2 (a) Normal vision; (b) advanced cataract vision

Examination

- Reduced visual acuity (sometimes improved with pinhole)
- Diminished red reflex on ophthalmoscopy
- A change in the appearance of the lens

The red reflex and ophthalmoscopy

The ‘red reflex’ is a reflection of the fundus when the eye is viewed from a distance of about 60 cm (2 feet) with the ophthalmoscope using a zero lens. This reflex is easier to see if the pupil is dilated. Commencing with the plus 15 or 20 lens, reduce the power gradually and, at plus 12, lens opacities will be seen against the red reflex which may be totally obscured by a very dense cataract. The setting up of the ophthalmoscope to examine intraocular structures is illustrated in Figure 77.3.



Figure 77.3 Settings of the ophthalmoscope used to examine intraocular structures

Management

Advise extraction when the patient cannot cope. Contraindications for extraction include intraocular inflammation and severe diabetic retinopathy. There is no effective medical treatment for established cataracts. The removal of the cataractous lens requires optical correction to restore vision and this is usually performed with an intraocular lens implant. Full visual recovery may take 2 - 3 months. Complications are uncommon yet many patients may require YAG laser capsulotomy to clear any opacities that may develop behind the lens implant.

Postoperative advice to the patient

- Avoid bending for a few weeks.
- Avoid strenuous exercise.
- The following drops may be prescribed:
    - steroids (to reduce inflammation)
    - antibiotics (to avoid infection)
    - dilators (to prevent adhesions)

Prevention

Sunglasses, particularly those that wrap around and filter UV light, may offer protection against cataract formation.

GLAUCOMA

Chronic simple glaucoma is the commonest cause of irreversible blindness in middle age.[1] At a very late stage it presents as difficulty in seeing because of loss of the outer fields of vision due to optic atrophy (Figure 77.4). Acute glaucoma, on the other hand, has a relatively rapid onset over a few days.



Figure 77.4 Typical visual field loss for chronic simple glaucoma; a similar pattern occurs with retinitis pigmentosa and hysteria

Clinical features of chronic glaucoma

- Familial tendency
- No early signs or symptoms

- Central vision usually normal
- Progressive restriction of visual field

Investigations

Tonometry

- Upper limit of normal is 22 mmHg

Ophthalmoscopy

- Optic disc cupping > 30% of total disc area

Screening

- Adults 40 years and over: 2 - 5 yearly (at least 2 yearly over 60)
- Start about 30 years, then 2 yearly if family history

Management

- Treatment can prevent visual field loss
- Medication (for life) usually selected from
    - timolol or betaxolol drops bd
      Note: These beta-blockers can cause systemic complications, e.g. asthma
    - latanoprost drops, once daily
    - pilocarpine drops qid
    - dipivefrine drops bd
    - acetazolamide (oral diuretics)
- Surgery or laser therapy for failed medication

RETINITIS PIGMENTOSA

Primary degeneration of the retina is a hereditary condition characterised by a degeneration of rods and cones associated with displacement of melanin-containing cells from the pigment epithelium into the more superficial parts of the retina.

Typical features

- Begins as night blindness in childhood
- Visual fields become concentrically narrowed (periphery to centre)
- Blind by adolescence (sometimes up to middle age)

Ophthalmoscopic examination

- Irregular patches of dark pigment, especially at periphery
- Optic atrophy

TRAUMA

Trauma to the eye may cause only a little discomfort so it is important to keep this in mind.

Intraocular foreign body

A small metal chip may penetrate the eye with minimal pain and the patient may not present with an ocular problem until the history of injury is long forgotten.

If infection does not supervene, presentation may be delayed for months or years until vision deteriorates due to metal degradation. The iris becomes rust-brown. It is important to X-ray the eye if it has been struck by a hammered fragment or if in any doubt at all about the mechanism of the injury.[1]

CHRONIC UVEITIS

Pain and redness may be minimal with this chronic inflammation. If untreated, visual loss often develops from secondary glaucoma and cataract. The pupil is bound to the lens by synechiae and is distorted.

HIV INFECTION

AIDS may have serious ocular complications, including Kaposi's sarcoma of the conjunctivae, retinal haemorrhage and vasculitis.[3] Another problem is ocular cytomegalovirus infection, which presents as areas of opacification with haemorrhage and exudates.

SUDDEN LOSS OF VISION

It is important to remember that the problem is alarming and distressing to the patient; considerable empathy is needed and care must be taken not to diagnose seemingly in-appropriate behaviour as of psychogenic origin.

A comparison of bilateral and unilateral causes of sudden loss of vision is presented in Table 77.4, and the diagnostic strategy model in Table 77.5. A simplified classification is:

- Unilateral: Retinal detachment retinal artery occlusion retinal vein thrombosis temporal arteritis optic neuritis migraine
- bilateral: bilateral optic nerve lesion hysteria

Table 77.4 Causes of sudden loss of vision[7]

		Unilateral	
	Bilateral	**Transient**	**Permanent**
Vascular causes	Occipital cortex ischaemia	Amaurosis fugax	Central retinal artery occlusion
	Pituitary apoplexy	Transient ocular ischaemia	Central retinal artery occlusion
	Homonymous hemianopia - vascular	Retinal emboli	Vitreous haemorrhage
		Malignant hypertension	Ischaemic optic neuropathy
Other causes	Bilateral optic neuritis	Acute angle closure glaucoma	Optic neuritis
	Toxic damage to optic nerve	Uhthoff's phenomenon	Retinal detachment
	• methanol • ethanol	Papilloedema	Optic nerve compression
	• tobacco • lead	Posterior vitreous detachment	Carcinomatous optic neuropathy
	Leber's optic atrophy		
	Quinine poisoning of retina		Intraocular tumour
	Cerebral oedema		

	Occipital lobe trauma
	Craniopharyngioma
	Hysteria

A flow chart for the diagnosis of painless loss of vision is presented in Figure 77.5.


Painless loss of vision
Is vision normal through pinhole?
no
yes
refractive error
Was onset sudden?
no, gradual
yes
Was visual loss transient?
no, prolonged
yes
Is direct pupil response normal?
no
yes
• retinal detachment
• vitreous haemorrhage
• central retinal vein occlusion
• cerebrovascular accident
• posterior vitreous detachment
• temporal arteritis
• central retinal artery occlusion
• optic neuritis
? associated headache
yes
no
• migraine (without headache)
• amaurosis fugax
Is red reflex normal?
yes
no
• chronic simple glaucoma
• diabetic retinopathy
• macular degeneration
• optic atrophy
• cataract
• malignant melanoma

Figure 77.5 Diagnosis of painless loss of vision Reproduced with permission Dr. J Reich and Dr J. Colvin.

Amaurosis fugax

Amaurosis fugax is transient loss of vision (partial or complete) in one eye due to transient occlusion of a retinal artery. It is painless and lasts less than 60 minutes. It is usually caused by an embolus from an atheromatous carotid artery in the neck. The most common emboli are cholesterol emboli, which usually arise from an ulcerated plaque.[7] Other causes include emboli from the heart, temporal arteritis and benign intracranial hypertension. Other symp-toms or signs of cerebral ischaemia, such as transient hemiparesis, may accompany the symptom. The source of the problem should be investigated. The risk of stroke after an episode of amaurosis fugax appears to be about 2% per year.[7]

Transient ocular ischaemia

Unilateral loss of vision provoked by activities such as walking, bending or looking upwards is suggestive of ocular ischaemia.[7] It occurs in the presence of severe extracranial vascular disease and may be triggered by postural hypotension and stealing blood from the retinal circulation.

Retinal detachment

Retinal detachment may be caused by trauma, thin retina (myopic people), previous surgery (e.g. cataract operation), choroidal tumours, vitreous degeneration or diabetic retinopathy.

Clinical features

- Sudden onset of floaters or flashes or black spots
- Blurred vision in one eye becoming worse
- ‘A curtain came down over the eye’, grey cloud or black spot
- Partial or total loss of visual field (total if macula detached)

Ophthalmoscopy may show detached retinal fold as large grey shadow in vitreous cavity.

Management

- Immediate referral for sealing of retinal tears
- Small holes treated with laser or freezing probe
- True detachments usually require surgery

Vitreous haemorrhage

Haemorrhage may occur from spontaneous rupture of vessels, avulsion of vessels during retinal traction or bleeding from abnormal new vessels.[6] Associations include ocular trauma, diabetic retinopathy, tumour and retinal detachment.

Table 77.5 Acute or subacute painless loss of vision: diagnostic strategy model

Q. Probability diagnosis
A. Amaurosis fugax
Migraine
Retinal detachment
Q. Serious disorders not to be missed
A. Cardiovascular
• central retinal artery occlusion
• central retinal vein occlusion
• hypertension (complications)
Neoplasia
• intracranial tumour
• intraocular tumour
- primary melanoma
- retinoblastoma
- metastases
AIDS
Temporal arteritis
Acute glaucoma
Benign intracranial hypertension
Q. Pitfalls (often missed)
A. Acute glaucoma
Papilloedema
Optic neuritis
Intraocular foreign body

Q. Seven masquerades checklist

A. Depression -

Diabetes ✔ diabetic retinopathy

Drugs ✔

Anaemia -

Thyroid disorder ✔ hyperthyroidism

Spinal dysfunction -

UTI -

Q. Is this patient trying to tell me something?

A. Consider ‘hysterical’ blindness, although it is uncommon.

Clinical features

- Sudden onset of floaters or ‘blobs’ in vision
- May be sudden loss of vision
- Visual acuity depends on the extent of the haemorrhage; if small, visual acuity may be normal

Ophthalmoscopy may show reduced light reflex: there may be clots of blood that move with the vitreous (a black swirling cloud).

Management

- Urgent referral to exclude retinal detachment
- Exclude underlying causes such as diabetes
- Ultrasound helps diagnosis
- May resolve spontaneously
- Bed rest encourages resolution
- Surgical vitrectomy for persistent haemorrhage

Central retinal artery occlusion

The cause is usually arterial obstruction by atherosclerosis, thrombi or emboli. There may be a history of TIAs. Exclude temporal arteritis (perform immediate ESR).

Clinical features

- Sudden loss of vision like a ‘curtain descending’ in one eye
- Vision not improved with 1 mm pinhole
- Usually no light perception

Ophthalmoscopy

- Initially normal
- May see retinal emboli
- Classic ‘red cherry spot’ at macula

Management

If seen early, use this procedure within 30 minutes:

- massage globe digitally through closed eyelids (use rhythmic direct digital pressure)
- rebreathe carbon dioxide (paper bag) or inhale special CO_2 mixture (carbogen)
- intravenous acetazolamide (Diamox) 500 mg

- refer urgently (less than 6 hours) - exclude temporal arteritis

Prognosis is poor. Significant recovery is unlikely unless treated immediately (within 30 minutes).

Central retinal vein thrombosis

Thrombosis is associated with several possible factors, such as hypertension, diabetes, anaemia, glaucoma and hyperlipidaemia.

Clinical features

- Sudden loss of central vision in one eye (if macula involved)
- Vision not improved with 1 mm pinhole

Ophthalmoscopy shows swollen disc and multiple retinal haemorrhages, “stormy sunset” appearance.

Management

No immediate treatment is effective. The cause needs to be found first and treated accordingly. Some cases respond to fibrinolysin treatment. Laser photocoagulation may be necessary in later stages to prevent thrombotic glaucoma.

Macular degeneration

There are two types: exudative (acute) and pigmentary (slow onset).

- Caused by neovascular membranes which develop under the retina of the macular area and leak fluid or bleed
- More common with increasing age (usually over 60) and those with myopia (relatively common)
- May be familial

Clinical features

- Sudden fading of central vision
- Distortion of vision
- Straight lines may seem wavy and objects distorted
- Use a grid pattern (Amsler chart): shows distorted lines
- Central vision eventually completely lost
- Peripheral fields normal

Ophthalmoscopy

- White exudates, haemorrhage in retina
- Macula may look normal or raised

Management

Urgent referral for fluorescein angiography and laser photocoagulation where indicated. The recent Age-Related Eye Disease Study has provided confirmatory evidence that the chronic pigmentation type responds to free-radical treatment with the antioxidants vitamins A, C, E, and zinc using betacarotene, 15 mg; vitamin C, 500 mg; vitamin E, 400 IU; and 80 mg zinc oxide.[9]

Temporal arteritis

With temporal arteritis (giant cell arteritis) there is a risk of sudden and often bilateral occlusion of the short ciliary arteries supplying the optic nerves, with or without central retinal artery involvement.[10]

Clinical features

- Usually older person: over 65 years

- Sudden loss of central vision in one eye (central scotoma)
- Can rapidly become bilateral
- Associated temporal headache (not invariable)
- Temporal arteries tender, thickened and nonpulsatile (but often normal)
- Visual acuity severely impaired
- Afferent pupil defect on affected side
- Usually elevated ESR > 40 mm

Ophthalmoscopy shows optic disc swollen at first, then atrophic. The disc may appear quite normal.

Management

- Other eye must be tested
- Immediate corticosteroids (100 mg prednisolone daily for at least 1 week)
- Biopsy temporal artery (if there is a localised tender area)

Migraine

Migraine may present with symptoms of visual loss. Associated headache and nausea may not be present.

Clinical features

- Zigzag lines or lights
- Multicoloured flashing lights
- Unilateral or bilateral field deficit
- Resolution within a few hours

Posterior vitreous detachment

The vitreous body collapses and detaches from the retina. It may lead to retinal detachment.

Clinical features

- Sudden onset of floaters
- Visual acuity usually normal
- Flashing lights indicate traction on the retina

Management

- Refer to an ophthalmologist urgently.
- An associated retinal hole or detachment needs exclusion.

Optic (retrobulbar) neuritis (77.6)

Causes include multiple sclerosis, neurosyphilis and toxins. A significant number of cases eventually develop multiple sclerosis.

Clinical features

- Usually a woman 20 - 40 years
- Loss of vision in one eye over a few days
- Retro-ocular discomfort with eye movements
- Variable visual acuity
- Usually a central field loss (central scotoma)
- Afferent pupil defect on affected side

Ophthalmoscopy

- Optic disc swollen if 'inflammation' anterior in nerve
- Optic atrophy appears later
- Disc pallor is an invariable sequel

Management

- Test visual field of other eye.
- Consider MRI.
- Most patients recover spontaneously but are left with diminished acuity.
- Intravenous steroids hasten recovery and have a protective effect against the development of further demyelinating episodes.

Corneal disorders

Patients with corneal conditions typically suffer from ocular pain or discomfort and reduced vision. The common condition of dry eye may involve the cornea while contact lens disorders, abrasions/ulcers and infections are common sensors problems that threaten eye sight. Inflammation of the cornea - **Keratitis** - is caused by factors such as ultraviolet light eg. 'arc eye', herpes simplex, herpes zoster ophthalmicus and the dangerous 'microbio/keratitis'. Bacterial keratitis is an ophthalmological emergency energy which should be considered in the contact lens nearer presently with pain and reduced vision. Topical corticosteroids should be avoided in the undiagnosed red eye.



PITFALLS

- Mistaking the coloured haloes of glaucoma for migraine.
- Failing to appreciate the presence of retinal detachment in the presence of minimal visual impairment.
- Omitting to consider temporal arteritis as a cause of sudden visual failure in the elderly.
- Using eyedrops to dilate the pupil (for fundal examination) in the presence of glaucoma.

WHEN TO REFER

- Most problems outlined need urgent referral to an ophthalmologist.
- Acute visual disturbance of unknown cause requires urgent referral.
- Any blurred vision - sudden or gradual, painful or painless - especially if 1 mm pinhole fails to alter visual acuity.
- Refer all suspicious optic discs.

Practice tips

- Tonometry is advised routinely for all people over 40; those over 60 should have tests every 2 years.
- Any family history of glaucoma requires tonometry at earliest age.
- Sudden loss of vision in the elderly suggests temporal arteritis (check the ESR and temporal arteries). It requires immediate institution of high-dose steroids to prevent blindness in the other eye. A time-scale

guide showing the rate of visual loss is presented in Table 77.6.

- Temporal arteritis is an important cause of retinal artery occlusion.
- Suspect field defect due to chiasmal compression if people are misjudging when driving.
- Pupillary reactions are normal in cortical blindness.
- Central retinal artery occlusion may be overcome by early rapid lowering of intraocular pressure.
- Retinal detachment and vitreous haemorrhage may require early surgical repair.
- Keep in mind antioxidant therapy (vitamins and minerals) for chronic macular degeneration.
- Consider multiple sclerosis foremost if there is a past history of transient visual failure, especially with eye pain.
- If the patient has been using a hammer, always X-ray if a fragment of metal has hit the eye but nothing can be seen.

Table 77.6 Time-scale guide for rate of visual loss[3, 7]

Sudden: less than 1 hour
• Amaurosis fugax
• Central retinal artery occlusion
• Hemianopias from ischaemia (emboli)
• Migraine
• Vitreous haemorrhage
• Acute angle glaucoma
• Papilloedema
Within 24 hours
• Central retinal vein occlusion
• Hysteria
Less than 7 days
• Retinal detachment
• Optic neuritis
• Acute macular problems
Up to several weeks (variable)
• Choroiditis
• Malignant hypertension
Gradual
• Compression of visual pathways
• Chronic glaucoma
• Cataracts
• Diabetic maculopathy
• Retinitis pigmentosa
• Macular degeneration
• Refractive errors

REFERENCES

1. Colvin J, Reich J. CHECK Programme, Unit 219-220. Melbourne: RACGP, 1990, 1 - 32.
2. Beck ER, Francis JL, Souhami RL. *Tutorials in Differential Diagnosis* (2nd edn). Edinburgh: Churchill Livingstone, 1988: 141 - 4.
3. Enoch B. Painless loss of vision in adults. Update, 1987; 5 June: 22 - 30.
4. Robinson MJ. Roberton, DIH *Practical Paediatrics* (5th edn). Melbourne: Churchill Livingstone, 2003: 756 - 770.
5. Cole GA. Amblyopia and strabismus. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 20 - 4.
6. Elkington AR, Khaw PT. *ABC of Eyes*. London: British Medical Association, 1990: 20 - 38.
7. King J. Loss of vision. Mod Med Aust, 1990; May: 52 - 61.
8. Hodge C, Ng D. Eye emergencies. CHECK Program, Unit 400. Melbourne: RACGP, 2005: 1 - 34.
9. Age-Related Eye Disease Study Research Group. A Randomised, placebo-controlled, clinical trial of high dose supplementation with vitamins C and E, beta-carotene, and zinc for age-related macular degeneration and vision loss. Arch Ophthalmol, 2001; 119: 1417 - 36.
10. Warne R, Prinsley D. A *Manual of Geriatric Care*. Sydney: Williams & Wilkins, 1988: 191 - 5.

CHAPTER 78. WEIGHT GAIN

Persons who are naturally very fat are apt to die earlier than those who are slender.

HIPPOCRATES

WEIGHT GAIN

Obesity is the most common nutrition-related disorder in the Western world; as Tunnessen puts it, ‘Obesity is the most common form of malnutrition in the United States’.[1] Most overweight adults and children who are obese have exogenous obesity, which tends to imply that ‘they ate too much’, but the problem is more complex than relative food input. Physical activity, and environmental and genetic influences must also be taken into account. There is still a persisting tendency of affected families to blame ‘glandular’ problems as a cause of obesity. It is now considered that there is a strong genetic basis to obesity and that attributing it to overeating and lack of exercise is an oversimplistic viewpoint.

KEY FACTS AND FIGURES

- The cause of exogenous obesity is multifactorial, the end result being increased body fatness (greater than 30% of total body weight in females and greater than 25% in males).[2]
- Abdominal obesity gives a higher cardiovascular risk at any weight.
- The onset of obesity can occur at any age.
- Secondary or pathologic causes are rare.
- Less than 1% of obese patients have an identifiable secondary cause of obesity.[3]
- Two conditions causing unexplained weight gain that can be diagnosed by the physical examination are Cushing's syndrome and hypothyroidism.
- After pregnancy, obesity may result from a failure to return to prepartum energy requirements.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 78.1.

Table 78.1 Weight gain: diagnostic strategy model

Q. Probability diagnosis A. Exogenous obesity
Q. Serious disorders not to be missed A. Cardiovascular • cardiac failure Hypothalamic disorders • craniopharyngiomas • optic gliomas Liver failure Nephrotic syndrome
Q. Pitfalls (often missed) A. Pregnancy (early) Endocrine disorders • hypothyroidism

- Cushing's syndrome
- insulinoma
- acromegaly
- hypogonadism
- hyperprolactinaemia
- polycystic ovarian disease

Idiopathic oedema syndrome

Klinefelter's syndrome

Congenital disorders

- Prader-Willi syndrome
- Laurence-Moon-Biedl syndrome

Q. Seven masquerades checklist

A. Depression

Diabetes -

Drugs

Anaemia -

Thyroid disorder   hypothyroid

Spinal dysfunction -

UTI -

Q. Is the patient trying to tell me something?

A. Yes: the reasons for obesity should be explored.

Probability diagnosis

The outstanding cause of weight gain in exogenous obesity is excessive calorie intake coupled with lack of exercise. Overweight people often deny overeating but the true situation can be determined by recording actual food intake and energy expenditure, and by interviewing reliable witnesses.

Serious causes not to be missed

It is important not to misdiagnose hypothalamic disorders, which may result in hyperphagia and obesity. Injury to the hypothalamus may occur following trauma and encephalitis and with a variety of tumours, including craniopharyngiomas, optic gliomas and pituitary neoplasms. Some of these tumours may cause headaches and visual disturbances.

It is important not to overlook major organ failure and kidney disorders as a cause of increased body weight, especially cardiac failure, liver failure and the nephrotic syndrome. The associated increase in body water needs to be distinguished from increased body fat.

Pitfalls

Endocrine disorders

The endocrine disorders that cause obesity include Cushing's syndrome, hypothyroidism, insulin-secreting tumours and hypogonadism. They should not represent difficult diagnostic problems.

An insulin-secreting tumour (insulinoma) is a very rare adenoma of the B cells of the islets of Langerhans. The main features are symptoms of hypoglycaemia and obesity.

Congenital disorders

The rare congenital disorders that cause obesity, such as Prader-Willi and Laurence-Moon-Biedl syndromes, should be easy to recognise in children (see p. 852).

Chromosomal abnormalities

An important abnormality to bear in mind is Klinefelter's syndrome (XXY karyotype), which affects one out of every 400 - 500 males. The boys show excessive growth of long bones and are tall and slim. Without testosterone treatment they become obese as adults.

Some girls with Turner's syndrome (XO karyotype) may be short and overweight.

Some gender pointers

Consider polycystic ovarian disease in women and obstructive sleep apnoea in obese men.

Seven masquerades checklist

The important masquerades include hypothyroidism and drug ingestion. Hypothyroidism is usually not associated with marked obesity. Drugs that can be an important contributing factor include tricyclic antidepressants, corticosteroids, pizotifen, thioridazine, Depo-Provera and the contraceptive pill. Obesity (overeating) may be a feature of depression, especially in the early stages. Prescribed tricyclic antidepressants may compound the problem.

Psychogenic considerations

An underlying emotional crisis may be the reason for the overweight patient to seek medical advice. It is important to explore diplomatically any hidden agenda and help the patient to resolve any conflict.

THE CLINICAL APPROACH

A careful history is very valuable in ascertaining food and beverage intake and perhaps giving patients insight into their calorie intake, since some deny overeating or will underestimate their food intake.[4]

Relevant questions

- Do you feel that you have an excessive appetite?
- Tell me in detail what you ate yesterday.
- Give me an outline of a typical daily meal.
- Tell me about snacks, soft drink and alcohol that you have.
- What exercise do you get?
- Do you have any special problems, such as getting bored, tense and upset or depressed?
- What drugs are you taking?

Physical examination

In the physical examination it is very important to measure body weight and height and calculate the BMI, and assess the degree and distribution of body fat and the overall nutritional status. Record the blood pressure and test the urine for sugar. Keep in mind that a standard blood pressure cuffon a large arm may give falsely elevated values. Remember the rare possibilities of Cushing's syndrome, acromegaly and hypothyroidism. Search for evidence of atherosclerosis and diabetes and for signs of alcohol abuse.

An extensive working up of the CNS is not indicated in obesity without the presence of suspicious symptoms such as visual difficulties.

Investigations

It is essential to perform two measurements:

- weight and height (to calculate BMI)
- waist - hip ratio

Important investigations

- Cholesterol/triglycerides
- Glucose (fasting)
- Liver function tests
- Electrolytes and urea

Investigations to consider

- Thyroid function tests
- Cortisol (if hypertensive)
- Testosterone (suspected sleep apnoea)
- ECG and chest X-ray (older than 40)

Anthropometric measurements

Useful measuring instruments include:

- BMI: ‘healthy’ range is between 20 and 25
- Waist circumference: risk of comorbidities
    - in men > 94 cm
    - in women > 80 cm
- waist - hip circumference ratio (W/H ratio): healthy range < 0.9 - a better predictor of cardiovascular risk than BMI
- single skinfold thickness (> 25 mm suggests increased body fat)
- upper arm circumference
- 4 skinfold thickness (sum of suprailiac, subscapular, triceps and biceps skinfolds) - for calculation of percentage body fat

Abdominal fatness is defined as a W/H ratio of > 0.85 in women and > 0.95 in men.

Body mass index

The easiest and possibly most accurate assessment of obesity is the BMI (refer to Appendix VI):

$$\text{BMI} = \frac{\text{weight (kg)}}{\text{height (M}^2\text{)}}$$

Garrow[5] has produced a simple classification of the BMI associated with the relative degree of risk increase and suggested therapy (Table 78.2).

Table 78.2 Classification of obesity (based on WHO guidelines)[5]

BMI	Grading	Suggested therapy
< 18.5	underweight	diet and counselling
18.5 - 25	healthy weight	
25 - 30	overweight	more exercise diet: less alcohol
30 - 35	Grade I: obesity	combined program: • behaviour modification • diet • exercise

35 - 40	Grade II: obesity	consider medical therapy if > 35
≥ 40	Grade III: morbid obesity	combined program plus medical therapy consider gastric surgery

WEIGHT GAIN IN CHILDREN

Various studies have found that approximately 10% of prepubertal and 15% of adolescent age groups are obese.[1]

Parents often blame obesity in children on their ‘glands’, but endocrine or metabolic causes are rare and can be readily differentiated from exogenous obesity by a simple physical examination and an assessment of linear growth. Children with exogenous obesity tend to have an accelerated linear growth whereas children with secondary causes are usually short.

Congenital or inherited disorders associated with obesity

Prader-Willi syndrome

The characteristic features are bizarre eating habits (e.g. binge eating), obesity, hypotonia, hypogonadism, intellectual disability, small hands and feet and a characteristic facial appearance (narrow bifrontal diameter, ‘almond-shaped’ eyes and a ‘tented’ upperlip). Progressive obesity results from excessive intake in addition to decreased caloric requirements.

Laurence-Moon-Biedl syndrome

The characteristic features are obesity, intellectual disability, polydactyly and syndactyly, retinitis pigmentation and hypogonadism.

Beckwith-Wiedemann syndrome

Characteristics include excessive growth, macrosomia, macroglossia, umbilical hernia and neonatal hypoglycaemia. Children appear obese as they are above the 95th percentile by 18 months of age. Intelligence is usually in the normal range.

Endocrine disorders

Endocrine disorders in children that can rarely cause obesity include hypothyroidism (often blamed as the cause but seldom is), Cushing's syndrome, insulinomas, hypothalamic lesions, Fröhlich's syndrome (adiposogenital dystrophy) and Stein-Leventhal syndrome in girls.

Managing obesity in children

Childhood obesity usually reflects an underlying problem in the family system. It can be a very difficult emotional problem in adolescents, who develop a poor body image. An important strategy is to meet with family members, determine whether they perceive the child's obesity as a problem and whether they are prepared to solve the problem. The family dynamics will have to be assessed and strategies outlined. This may involve referral for expert counselling. It is worth pointing out that children eat between one-third and two-thirds of their meals at school so schools should be approached to promote special programs for children who need weight reduction.

CUSHING'S SYNDROME

Cushing's syndrome is the term used to describe the chemical features of increased free circulating glucocorticoid. The most common cause is iatrogenic with the prescribing of synthetic corticosteroids. The spontaneous primary forms such as Cushing's disease (pituitary dependent hyperadrenalism) are rare. As the disorder progresses the body contour tends to assume the often quoted configuration of a lemon with matchsticks (Figure 78.1).



Figure 78.1 The traditional 'lemon with matchsticks' configuration of Cushing's syndrome

Typical clinical features

- Change in appearance
- Central weight gain (truncal obesity)
- Hair growth and acne in females
- Muscle weakness
- Amenorrhoea/oligomenorrhoea (females)
- Thin skin/spontaneous bruising
- Polymyalgia/polydipsia (diabetes mellitus)
- Insomnia
- Depression

Signs

- Moon face
- 'Buffalo bump'
- Purple striae
- Large trunk and thin limbs

The patient should be referred for diagnostic evaluation, including plasma cortisol and overnight dexamethasone suppression tests.

Untreated Cushing's syndrome has a very poor prognosis, with premature death from myocardial infarction, cardiac failure and infection; hence early diagnosis and referral is essential.

OBESITY

Obesity and overweight are the most common pathological conditions in our society and are caused by an accumulation of adipose tissue (Table 78.3). It is not the extra weight *per se* that causes problems but excess fat. The calculation of the BMI gives a better estimate of adiposity and it is convenient and preferable to use this index when assessing the overweight and obese. However, recent data suggests that the distribution of body fat is as important a risk factor as its total amount. Abdominal fat (upper body segment obesity, or 'apple' obesity, is considered a greater health hazard than fat in the thighs and buttocks (lower body segment obesity, or 'pear' obesity) (Figure 78.2).

Table 78.3 Factors predisposing to primary obesity

Genetic	- familial tendency
Sex	- women more susceptible
Activity	- lack of physical activity
Psychogenic	- emotional deprivation; depression
Social class	- poorer classes
Alcohol	- problem drinking
Smoking	- cessation of smoking
Prescribed drugs	- tricyclic derivatives



Figure 78.2 Comparison of two types of obesity according to distribution of body fat

Obese patients with high waist - hip ratios (> 1.0 in men and > 0.9 in women) have a significantly greater risk of diabetes mellitus, stroke, coronary artery disease and early death than equally obese people with lower waist - hip ratios.[3]

In regard to the BMI reference scale it is worth noting that the risks follow a J-shaped curve (Figure 78.3) and are only slightly increased in the overweight range but increase with obesity so that a BMI of > 40 carries a threefold increase in mortality.



Figure 78.3 Body mass index (BMI) reference scale

The consequences of obesity include:

- cardiovascular
  - increased mortality (stroke, ischaemic heart disease etc.)
  - hypertension
  - varicose veins
- metabolic
  - dyslipidaemia
  - type 2 diabetes
  - insulin resistance
  - hyperuricaemia/gout
  - infertility
  - PCOS
- mechanical
  - osteoarthritis
  - obstructive sleep apnoea
  - spinal dysfunction
  - low back pain
- other
  - gall bladder disease
  - cancer (various)
  - kidney disease (check microalbuminuria)
  - psychological problems

Management

Treatment is based on four major interventions, the choice of which depends on the degree of obesity, the associated health problems and the health risk posed:[2]

1. reduction in energy intake

2. change in diet composition
3. increased physical activity
4. behavioural therapy

Pharmacological agents are not used for first-line therapy although they may have a place in management, especially at grade III level of obesity. Surgery is an option for the treatment of morbid obesity.

There is no single effective method for the treatment of obesity, which is a difficult and frustrating problem. A continuing close therapist/patient contact has a better chance of success than any single treatment regimen.

Goals

1. no further weight gain
2. loss of 5 - 10% initial body weight
3. improve activity

Most successful programs involve a multidisciplinary approach to weight loss, embracing the four major interventions. The first goal should be no further weight gain. Emphasis must be on maintenance of weight loss. Behaviour modification is important and the most valuable strategy is to emphasise planning and record keeping with a continuous weekly diary of menus, exercise and actual behaviour.

Social support is essential for a successful weight loss program. A better result is likely if close family members, especially the chief cook, are involved in the program, preferably striving for the same goals.[3]

A doctor - patient strategy

A close therapeutic supportive relationship with a patient can be effective using the following methods:[6]

1. *Promote realistic goals*. Lose weight at the same rate that it was gained (i.e. slowly) - for example, 5 - 10 kg a year. A graph can be used for this purpose with an ‘exaggerated’ scale on the vertical axis so that small variations appear highly significant and encouraging (Figure 78.4).



Figure 78.4 Weight loss chart to encourage patient. After Kaczmarczyk.[6]

Promote the equation: Energy In = Energy Out + Energy Stored

The only way to reduce the stored energy (fat) is either to reduce energy in (eat less) or increase energy out (exercise).

2. *Dietary advice*. It is important to be realistic and allow patients to eat their normal foods but advise them about quantity and frequency. Give advice on simple substitutions (e.g. fortified skim milk in place of whole milk, high-fibre wholemeal bread instead of white bread, and fruit and vegetables instead of biscuits and cakes as in-between snacks).[6] A strategy that seems to work effectively is to advise patients, especially those who are overweight (grade I obesity), to eat onethird less than they usually do and discipline themselves not to 'pick' and to avoid second helpings.
3. *Counselling* is simple and commonsense. It involves being supportive, interested and encouraging. A list of tips on coping is provided (see below 'a practical plan' for grade II and III obesity) and the patient advised to keep a food, exercise and behaviour diary.
4. *Review*. 'Review is the most vital part of the weight loss programme as it stimulates and revitalises motivation and enables assessment of progress'.[6] It should be frequent initially (e.g. fortnightly), then monthly until the goal weight has been achieved and then 3-monthly. It is important never to be judgmental or critical if progress is unsatisfactory.

A practical plan

The following patient education sheet to be handed to patients represents useful advice to offer the obese patient.[7] The emphasis is on a healthy lifestyle program.

Physical activity

- A brisk walk for at least 20 - 30 minutes each day is the most practical exercise.
- Other activities such as tennis, swimming, golf and cycling are a bonus.

Dietary advice

Provide patients with a glycaemic index guide (Table 10.1 at p. 83).

Breakfast:

- oatmeal (soaked overnight in water); after cooking, add fresh or dried fruit; serve with fat-reduced milk or yoghurt *or*
- muesli (homemade or from a health food store) - medium serve with fat-reduced milk, perhaps add extra fruit (fresh or dried)
- slice of wholemeal toast with a thin scraping of margarine, spread with Vegemite, Marmite or sugar-free marmalade
- fresh orange juice or herbal tea or black tea/coffee

Morning and afternoon tea:

- piece of fruit or vegetable (e.g. carrot or celery)
- freshly squeezed juice or chilled water with fresh lemon

Midday meal:

- salad sandwich with wholemeal or multigrain bread and thin scraping of margarine (for variety use egg, salmon, chicken or cheese fillings)
- drink as for breakfast

Evening meal:

- Summer (cold) - lean meat cuts (grilled, hot or cold), poultry (skin removed) or fish; fresh garden salad; slices of fresh fruit
- Winter (hot) - lean meat cuts (grilled), poultry (skin removed) or fish; plenty of green, red and yellow vegetables and small potato; fruit for sweets

Weight-losing tips

- Have sensible goals; do not ‘crash’ diet but have a 6 - 12 month plan to achieve your ideal weight.
- Go for natural foods; avoid junk foods.
- Avoid alcohol, sugary soft drinks and high calorie fruit juices.
- Strict dieting without exercise fails.
- If you are mildly overweight, eat one-third less than you usually do.
- Do not eat biscuits, cakes, buns, etc. between meals (preferably at no time).
- Use high-fibre foods to munch on.
- A small treat once a week may add variety.
- Avoid seconds and do not eat left overs.
- Eat slowly - spin out your meal.

Pharmacological agents

A variety of these are available or imminent but have limitations and need to be used with caution, if at all. Adverse effects can be problematic. Consider for those with a BMI > 30.

The agents are:[8, 9]

- Local, acting on GIT
  - bulking agents (e.g. methylcellulose)
  - lipase inhibitor - orlistat (Xenical) 120 mg (o) tds ac (used with a low-fat eating program)
- Centrally acting agents
  - amphetamine derivatives (reduce hunger)
    - phentermine 15 - 40 mg (o) daily
    - diethylpropion 25 mg (o) tds or SR 75 mg daily
  - serotonin analogues (enhance satiety)
    - fluoxetine 20 - 40 mg (o) daily
    - sibutramine 10 - 15 mg (o) daily
    - sertraline

A summary of systemic reviews to date indicates that the effectiveness of serotonin analogues is unknown or, with sibutramine and phentermine, of some benefit, which has to be weighed against adverse effects, including the potentially fatal serotonin syndrome. The same applies to orlistat, which together with a low-fat eating program, does produce extra weight loss over placebo.[10]

Meal replacements

Food replacement agents such as Modifast and SlimFast have been promoted but there is insufficient evidence so far to evaluate their effectiveness and the reduced intake of key vitamins.

Surgery

In those with morbid obesity (about 2% of the population) unresponsive to behaviour modification therapy and a course of dexfenfluramine for 3 months or so, gastric banding has a place. One example is Lap-Band, which is inserted laparoscopically and can be adjusted and eventually removed with no significant residual adverse defect left in the stomach. It appears to be effective for 10 years.[9, 11] Gastric stapling and gastric bypass are other techniques to consider.

OEDEMA

Oedema (dropsy) is an excessive accumulation of fluid in tissue spaces. It may be generalised or localised - periorbital, peripheral or an arm (lymphoedema, refer p. 768).

Generalised oedema

The site of generalised oedema is largely determined by gravity. It is due to an abnormal excess of sodium in the body, which leads to accumulation of water. The causes can be generally divided into two groups - oedema associated with a decreased plasma volume and oedema associated with an increased plasma volume (see Table 78.4).

Table 78.4 Causes of generalised oedema

Decreased plasma volume
• Hypoalbuminaemia (e.g. nephrotic syndrome, chronic liver disease, malnutrition)
Increased plasma volume
• Congestive cardiac failure
• Chronic kidney failure
• Drugs (e.g. corticosteroids, NSAIDs, certain antihypertensives, oestrogens, lithium, others)
Idiopathic oedema

Diagnosis

Clinical examination, including urinalysis, is usually sufficient to establish the cause of the oedema. In other cases, investigation of kidney or liver function may be required.

Treatment of generalised oedema

- Treat the cause where known
- Salt (sodium) restriction
- Diuretics
    - a loop diuretic (e.g. frusemide)
    - a potassium-sparing diuretic (e.g. spironolactone)

Idiopathic oedema

Idiopathic oedema, also known as cyclical or periodic oedema, is a common problem and the diagnosis is made on a characteristic history:

- exclusive to women
- may be cyclical or persistent
- usually unrelated to menstrual cycle
- excessive diurnal weight gain (worse on prolonged standing)
- abdominal bloating
- may affect hands and face as well as feet
- often made worse by diuretics
- may be associated with headache, depression, tension

Treatment of this condition is difficult. Most diuretics can aggravate the problem. Supportive stockings and a nutritious diet (with restricted sodium intake) is recommended as first-line treatment. A trial of spironolactone is often recommended.

Swelling (puffiness) of the face and eyelids

The causes are similar to those for generalised oedema. Important specific causes to consider are:

- kidney disease (e.g. nephrotic syndrome, acute nephritis)
- hypothyroidism

- Cushing's syndrome and corticosteroid treatment
- mediastinal obstruction
- angio-oedema
- skin sensitivity (e.g. drugs, cosmetics, hair dryers)

Swelling of the legs

Refer to p. 768.

‘CELLULITE’

Many patients seek advice about ‘cellulite’ in the buttocks and thighs in particular. Explain that the best way to overcome it is to maintain an ideal weight. If overweight, lose it slowly and exercise to improve the muscle tone in the buttocks and thighs.

WHEN TO REFER

- Patients with grade II or III obesity (BMI > 30) who are resistant to simple weight control measures.[2]
- Patients with associated medical problems such as angina or severe osteoarthritis who require rapid weight reduction.
- Possibility of endocrine cause of obesity.
- Suspicion of congenital or inherited disorder in children.

Practice tips

- Avoid a critical or judgmental attitude to the overweight patient.[12]
- Seek diplomatic independent information from a spouse or parent about food and beverage intake.
- Obtain a chronological history of the patient's weight from infancy onwards and attempt to correlate any significant changes to stressful life events.
- Central or visceral obesity carries a large risk factor for medical complications. People are advised to keep the waist circumference to less than 100 cm.

REFERENCES

1. Tunnessen WW Jr. *Signs and Symptoms in Paediatrics* (2nd edn). Philadelphia: JB Lippincott, 1988: 33 - 41.
2. Marks S, Walqvist M. Obesity. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 354 - 6.
3. Schroeder SA et al. *Current Medical Diagnosis and Treatment*. East Norwalk: Appleton & Lange, 1990: 874 - 5.
4. Sloane PD, Slatt PD, Baker RM. *Essentials of Family Medicine*. Baltimore: Williams & Wilkins, 1988: 200 - 8.
5. WHO Technical Report Series Number 894. Geneva: WHO, 2000.
6. Kaczmarczyk W. The obese patient. In: How I manage my difficult problems. Aust Fam Physician, 1991; 20: 417 - 21.
7. Murtagh JE. Obesity: how to lose weight wisely. In: *Patient Education* (4th edn). Sydney: McGraw-Hill, 2004: 104.
8. Caterson ID. Obesity: effective management in general practice. Mod Med Aust, 1996; 39(6): 70 - 80.
9. Shenfeld G (Chair). *Therapeutic Guidelines: Endocrinology* (Version 3). Melbourne: Therapeutic Guidelines Ltd, 2003: 41 - 50.
10. Arterburn D, Noel P. Obesity. In: Barton S ed. *Clinical Evidence* (issue 5). London: BMJ Publishing Group, 2001: 412 - 19.
11. O'Brien P. Advances in gastrointestinal surgery. Update Course handbook. Melbourne: Monash University, 1996: 29.
12. Kincaid-Smith P, Larkins R, Whelan G. *Problems in Clinical Medicine*. Sydney: Maclennan & Petty, 1990: 105 - 8.

CHAPTER 79. WEIGHT LOSS

> Among young women dieters in modern society about 1 in 20 will become preoccupied with their appearance and progress to the eating disorders of anorexia nervosa and bulimia.
>
> Professor Doris Young 1988

WEIGHT LOSS

In family practice complaints of loss of weight are more frequent than complaints about being too thin. Of great significance is the problem of recent loss of weight. A very analytical history is required to determine the patient's perception of weight loss. The equivalent problem in children is failure to gain weight or thrive.

Weight loss is an important symptom because it usually implies a serious underlying disorder, either organic or functional. It may or may not be associated with anorexia and thus diminished food intake.

KEY FACTS AND CHECKPOINTS

- Any loss of more than 5% of normal body weight is significant.
- The most common cause in adults of recent weight loss is stress and anxiety.[1]
- Serious organic diseases to consider are:
  - malignant disease
  - diabetes mellitus
  - chronic infections (e.g. tuberculosis)
  - thyrotoxicosis
- The most important variable to consider in evaluating weight loss is appetite. Eating and weight go hand in glove.
- Two conditions commonly associated with weight loss are anaemia and fever; they must be excluded.
- Early detection of eating disorders improves outcome.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 79.1.

Table 79.1 Weight loss: diagnostic strategy model (other than deliberate dieting or malnutrition)

Q. Probability diagnosis
A. Stress and anxiety
Non-coping elderly
Q. Serious disorders not to be missed
A. Chronic heart failure
Malignant disease, including
• stomach
• pancreas
• lung
• myeloma
• caecum
• lymphoma
Chronic infection

- HIV infections (AIDS, ARC)
- tuberculosis
- hidden abscess
- infective endocarditis
- brucellosis
- others

Q. Pitfalls (often missed)

A. Drug dependence, esp. alcohol

Malabsorption states

- ? intestinal parasites/infestations
- coeliac disease

Other GIT problems

Chronic kidney failure

Connective tissue disorders (e.g. SLE)

Dementia

Rarities

Addison's disease (p. 224)

Hypopituitarism

Q. Seven masquerades checklist

A. Depression

Diabetes

Drugs

Anaemia

Thyroid disorder   hyperthyroidism

Spinal dysfunction -

UTI

Q. Is the patient trying to tell me something?

A. A possibility. Consider stress, anxiety and depression. Anorexia nervosa and bulimia are special considerations.

Probability diagnosis

Excluding planned dietary restriction, psychological factors are the most common cause, particularly recent stress and anxiety.[1] Elderly people with adverse psychological factors, neglect and possibly drug effects can present with wasting.

Serious disorders not to be missed

Many of the problems causing weight loss are very serious, especially malignant disease.

Malignant disease

Weight loss may be a manifestation of any malignancy. With carcinoma of the stomach, pancreas and caecum, malignant lymphomas and myeloma, weight loss may be the only symptom. Occult malignancy must be regarded as the most common cause of weight loss in the absence of major symptoms and signs. The mechanisms may be multiple, with anorexia and increased metabolism being important factors.

Chronic infections

These are now less common but tuberculosis must be considered, especially in people from less developed countries. Some cases of infective endocarditis may progress only very slowly with general debility, weight loss and fever as major features.[2]

Other infections to consider are brucellosis, and protozoal and systemic fungal infection. Infection with HIV virus must be considered, especially in high-risk groups.

Pitfalls

Drug dependency, including alcohol and narcotic drugs, must be considered, especially when the problem may result in inappropriate nutrition. Apart from malignant disease there is a whole variety of gastrointestinal disorders that require consideration - these include malabsorption states, gastric ulceration, and intestinal infestations that should be considered, especially in people returning from a significant stay in tropical and under-developed countries.

Addison's disease (p. 224) can be very difficult to diagnose. Symptoms include excessive fatigue, anorexia, nausea and postural dizziness. Hyperpigmentation is a late sign.

Seven masquerades checklist

Depression and the endocrine disorders, diabetes mellitus and hyperthyroidism, are important causes.

Diabetes

The diabetic who presents with weight loss will be young and insulin-dependent. The initial presentation may be ketoacidosis. The triad of symptoms is thirst + polyuria + weight loss.

Hyperthyroidism

This is usually associated with weight loss although in some, such as an elderly male, it may not be obvious. An important clue will be weight loss in the presence of an excellent appetite and this helps to distinguish it from a psychoneurotic disturbance.

Depression

Weight loss is a common feature of depression and is usually proportional to the severity of the disease. In the early stages of depression weight gain may be present but when the classic loss of the four basic drives (appetite, energy, sleep and sex) becomes manifest, weight loss is a feature.

Drugs

Any prescribed drugs causing anorexia can cause weight loss. Important drugs include digoxin, narcotics, cytotoxics, NSAIDs, some antihypertensives and theophylline.

Psychogenic considerations

Weight loss is a feature of anxiety as well as depression. Some patients with psychotic disturbances, including schizophrenia and mania, may present with weight loss.

Anorexia nervosa is quite common and is almost entirely confined to females between the ages of 12 and 20 years. The main differential diagnosis is hypopituitarism, although anorexia nervosa can cause endocrine disturbances through the hypothalamic pituitary axis.

THE CLINICAL APPROACH

History

It is important to document the weight loss carefully and evaluate the patient's recordings. The same set of scales should be used. It is also important to determine the food intake. However, in the absence of an independent witness such as a spouse or parent, this can be difficult. Food intake may be diminished with psychogenic disorders and carcinoma but increased or steady with endocrine disorders, such as diabetes and hyperthyroidism, and with steatorrhoea. Figure 79.1 shows the possible causes of weight loss.


Malignancy
lymphoma
melanoma
lung
stomach
kidney
pancreas
colon
caecum
ovary
others
stress and anxiety
depression
eating disorders
drugs
hyperthyroidism
congestive cardiac failure
tuberculosis
diabetes (type 1)
GIT disorders
• malabsorption
• post surgery

Figure 79.1 Weight loss: causes to consider

General questions

- Exactly how much weight have you lost and over how long?
- Have you changed your diet in any way?
- Has your appetite changed? Do you feel like eating?
- Have your clothes become looser?
- What is your general health like?
- How do you feel in yourself?
- Do you feel uptight (tense), worried or anxious?
- Do you get very irritable or tremulous?
- Do you feel depressed?
- Do you ever force yourself to vomit?
- Are you thirsty?
- Do you pass a lot of urine?
- Do you have excessive sweating?
- Do you experience a lot of night sweats?
- What are your motions like?
- Are they difficult to flush down the toilet?
- Do you have a cough or bring up sputum?
- Do you get short of breath?
- Do you have any abdominal pain?
- Are your periods normal (for females)?
- What drugs are you taking?
- How many cigarettes do you smoke?

Physical examination

A careful general examination is essential with special attention to:

- vital parameters
- the thyroid and signs of hyperthyroidism
- the abdomen (check liver, any masses and tenderness)
- rectal examination (test stool for occult blood)
- reflexes

Investigations

Basic investigations include:

- haemoglobin, red cell indices and film
- white cell count
- ESR
- thyroid function test
- random blood sugar
- chest X-ray
- urine analysis

Others to consider:

- upper GIT (endoscopy or barium meal)
- ultrasound of abdomen (or CT if suspected abnormality not found)
- colonoscopy
- LFTs

WEIGHT LOSS IN CHILDREN

Weight loss in children can be considered as:

1. failure to thrive (FTT): the child up to 2 years below 3rd percentile (refer to Chapter 81, pp. 877 - 8)
2. weight loss in a child after normal development

Loss of weight in the older child

Acute or chronic infections are the most common causes of weight loss in children beyond infancy.[3] In acute infections the weight loss is transient, and once the infection clears the child generally regains the lost weight. In chronic infections signs may be more difficult to detect; for example, urinary tract infection, pulmonary infection, osteomyelitis, chronic hepatitis. In common with the younger child who fails to thrive, the older child may be suffering from malabsorption syndrome, chronic infection of the urinary tract or a rare chromosomal or metabolic disorder.[4] Tuberculosis, diabetes and malignant disease may present as weight loss and it is necessary to exclude organic disease before considering the more common emotional disorders.

EATING DISORDERS IN THE ADOLESCENT

Concerns about body image and dieting are very common among young women in modern society. Among these dieters 5 - 10% become abnormally preoccupied with dieting and slimness and progress to the eating disorders of anorexia nervosa and bulimia. Sufferers often have extremely low self-esteem and feel ineffective. They tend to be perfectionists with obsessive - compulsive traits. A history of childhood sexual abuse may be relevant. Media images alone do not cause eating disorders but have a role - genetic vulnerability, temperament, psychological and environmental factors also mediate these illnesses: ‘Genes load the gun - the environment pulls the trigger’.[5]

The DSM IV criteria for diagnosing these disorders, which have serious physical and psychological consequences, are presented in Table 79.2. The diferential diagnosis of anorexia nervosa includes most of the problems listed in

Table 79.1 at p. 859.

Table 79.2 DSM IV criteria for diagnosing anorexia nervosa and bulimia

Anorexia nervosa

A Refusal to maintain normal body weight Loss to < 85% of expected body weight

B Intense fear of becoming fat, despite current under weight status

C Disturbance of body image (body size or shape)

D Amenorrhoea in postmenarcheal females

Bulimia nervosa

A Recurrent episodes of binge eating, that is:

1. eating in a discrete period of time an abnormal quantity of foodcompared with the average person
2. a sense of lack of control during the binge

B Recurrent inappropriate compensatory behaviour to prevent weight gain (e.g. self-induced vomiting; misuse of laxatives, enemas etc.)

C A and B both occur, on average, at least twice a week for 3 months

D Self-evaluation is unduly influenced by body shape and weight

E Does not occur exclusively during periods of anorexia nervosa

Anorexia nervosa (Figure 79.2)



Figure 79.2 18 year old adolescent with severe anorexia nervosa (BMI 8.7). This patient survived after care by her general practitioner. Photo and history courtesy Dr. M M O'Brien

Anorexia nervosa is a syndrome characterised by the obsessive pursuit of thinness through dieting with extreme weight loss and disturbance of body image.[6] The mortality rate may be as high as 18%. It has the highest mortality and suicide rate of any psychiatric disorder.[5]

Typical features

- Adolescent and young adult females
- Up to 1% incidence among schoolgirls aged 16 [7]
- Bimodal age of onset: 13 to 14 and 17 to 18 years[6]
- Unknown cause
- Severe emaciation
- Amenorrhoea
- Loss of body fat
- Dry and scaly skin
- Increased lanugo body hair
- BMI < 17.5

Bulimia

Bulimia is episodic secretive binge eating followed by self-induced vomiting, fasting or the use of laxatives or diuretics. This binge - purge syndrome is also referred to as bulimarexia. It is more difficult to detect than anorexia nervosa but has a higher incidence. There are two types - the purging type and the nonpurging type where fasting or excessive exercise are the compensatory behaviours. The purging type is the most life-threatening behaviour because of the danger of hypokaelaemia.

Typical features

- Young females (F:M ratio 10:1)
- Begins at later age, usually 17 - 25 years
- Associated psychoneurotic disorders
- Fluctuations in body weight
- Periods irregular - amenorrhoea rare
- Physical complications of frequent vomiting (e.g. dental decay, effects of hypokalaemia)
- Recurrent laxative, stimulant or enema abuse

Laboratory evaluation should include electrolytes, iron studies, TFTs, kidney function tests, coeliac disease screening and LFTs, ESR.

MANAGEMENT OF EATING DISORDERS

Early detection and intervention are essential to reduce the risk of chronicity. Treatment can be conducted on an outpatient basis but if there are marked trends, such as severe weight loss, a family crisis, severe depression and a suicide risk, the patient requires hospital admission. The burden on caregivers of people with eating disorders is high. There are often problematic family interrelationships which require exploration.

Important goals are:

- establish a good and caring relationship with the patient
- resolve underlying psychological difficulties
- restore weight to a level between ideal and the patient's concept of optimal weight
- provide a balanced diet of at least 3000 calories per day (anorexia nervosa)

Structured behavioural therapy, intensive psychotherapy and family therapy may be tried but supportive care by physicians and allied health staffappears to be the most important feature of therapy.[8] Antidepressants, especially of the SSRI group, may be helpful for selected patients with a comorbid depressive illness. Fluoxetine is the preferred agent for bulimia. It is important to provide ongoing support for both patient and family.

WEIGHT LOSS IN THE ELDERLY

General weight loss is a relatively common physiological feature of many elderly people. However, abnormal weight loss is commonly encountered in the socially disadvantaged elderly, especially those who live alone and lack drive and interest in adequate food preparation. Other factors include relative poverty and poor dentition, including ill-fitting and painful false teeth. An important cause that should always be considered is malignant disease.

Congestive cardiac failure, especially secondary to ischaemic heart disease, is a common cause of weight loss. This is due to visceral congestion.

GASTROINTESTINAL CAUSES OF WEIGHT LOSS

The following conditions may lead to weight loss:

- coeliac disease
- poor oral hygiene
- chronic vomiting or diarrhoea (e.g. pyloric stenosis)
- gastric ulcer
- carcinoma of the stomach, oesophagus, large bowel
- problem alcohol drinking
- partial or total gastrectomy
- other GIT surgery
- inflammatory bowel disease (e.g. Crohn's disease, ulcerative colitis)
- steatorrhoea
- lymphoma of the gut
- parasitic infestation
- cirrhosis of the liver

The mechanisms of weight loss include anorexia, malabsorption, obstruction with vomiting and inflammation.

Practice tips

- Ask patients what they really believe is the cause of their weight loss.
- An anxiety state and hyperthyroidism can be difficult to differentiate. Consider the latter and perform thyroid function tests.
- Laboratory tests are rarely needed to establish the diagnosis of an eating disorder. Hormonal levels return to normal following weight gain.
- A high index of suspicion by the family doctor is required to diagnose eating disorders. Think of it in a mid-teen female; weight loss through dieting; wide fluctuation in weight; amenorrhoea and hyperactivity.

WHEN TO REFER

- Any unexplained weight loss, especially if an endocrine cause or malignancy is suspected
- Weight loss related to a serious psychological illness
- A serious eating disorder

REFERENCES

1. Cormack J, Marinker M, Morrell D. Practice: *A Handbook of Primary Health Care*. London: Kluwer-Harrap Handbooks, 1980; 3.42: 1 - 2.
2. Beck ER, Francis JL, Souhami RL. *Tutorials in Differential Diagnosis* (2nd edn). Edinburgh: Churchill Livingstone, 1988: 117 - 20.
3. Tunnessen WW. *Symptoms and Signs in Paediatrics* (2nd edn). Philadelphia: Lippincott, 1988: 25 - 8.

4. Robinson MJ, Roberton DM. *Practical Paediatrics* (5th edn). Melbourne: Churchill Livingstone, 2003: 140 - 4.
5. The Bronte Foundation. < http://www.thebrontefoundation. com.au>
6. Young D. Eating disorders. In: Jones R et al. Oxford Textbook of Primary Medical Care. Oxford: Oxford University Press, 2004: 972 - 5.
7. Crisp AH, Palmer RL, Kalucy RS. How common is anorexia nervosa? A prevalence study. Br J Psychiatry, 1976; 128: 549 - 54.
8. Tierney LM et al. *Current Medical Diagnosis and Treatment*. New York: The McGraw-Hill Companies, 2006: 1270 - 1.

CHAPTER 80. AN APPROACH TO THE CHILD

In every child who is born, under no matter the circumstances, and of no matter what parents, the potentiality of the human race is born again; and in him, too, once more, and of each of us, our terrific responsibility towards human life; towards the utmost idea of goodness, of the horror of error, and of God.

JAMES AGEE

AN APPROACH TO THE CHILD

The diagnostic approach to the child is based on the ability to achieve good lines of communication with both the child and the parent. In the diagnostic approach, the relative importance of the majority of the information required to reach a diagnosis comes from the history with a smaller amount from the physical examination.

The establishment of rapport can be achieved by showing a genuine interest in the child with strategies such as:

- asking them what they like to be called
- passing a compliment about the child such as a clothing item or a toy or book they are carrying
- taking time to converse with them
- asking them if they would like to be a doctor when they grow up
- asking about their teacher or friends
- ask them about their pets
- have special stickers to place on them e.g. backs of hands, T-shirts

This process will help set the scene for easier history taking and a sound physical examination.

Children's general behaviour patterns and personality can be classified into broad identifiable categories according to age group. Although there is considerable variation and generalisations can be inappropriate, the following stereotypes are helpful guidelines for parents.

- terrible 2s mischievous, explorative, dangerous activity, conflicts
- trusting 3s friendly, amenable to reason, loving
- frustrating 4s cheeky, inquisitive, hard to reason, no social graces
- fascinating 5s more co-ordinated and independent
- sociable 6s enjoys tasks for temporary interests, loves to be wooed
- problematic 7s tendency to wrongdoing, stubborn, searching for independence
- steady 8s
- noisy and adventurous 9s

HISTORY

Obtaining information in the following sequence is recommended:

- presenting problem (focus on this first)
    - allow the parents to elaborate without interruption
    - be a listener and believe the story
- state of health prior to the present complaint
- past history
    - general features
    - pregnancy and neonatal features
    - feeding and diet
    - immunisation

  - toilet training
- family history
  - inherited disorders
  - other
- systems review
  - general features (e.g. fever, energy)
  - feeding and elimination
  - hearing
  - vision
- developmental history
  - check list of milestones (Table 80.1)

Table 80.1 Developmental milestones (m = months) (after Jarman and Oberklaid[2])

Gross motor	***Expressive language***
Chin up (1 m)	Coos (3 m)
Lifts head (4 m)	Babbles (6 m)
RollsÑprone to supine (4 m)	Da-DaÑinappropriate (8 m)
RollsÑsupine to prone (5 m)	Da/MaÑappropriate (10 m)
Sits unsupported (8 m)	First word (11 m)
Pulls to stand (9 m)	Two to six words (15 m)
Cruises (10 m)	Two-word phrases (21 m)
Walks alone (13 m)	Speech all understandable (27 m)
Walks up stairs (20 m)	Names one colour (30 m)
Kicks ball forward (24 m)	Uses plurals (36 m)
Walks up stairsÑalternate feet (30 m)	Names four colours (42 m)
Rides tricycle (36 m)	Gives first and last names (44 m)
Two-wheeler bike (36 m)	Names two opposites (50 m)
Hops on one foot (60 m)	Strings sentences together (60 m)
Fine motor	***Receptive language***
Unfisting (3 m)	Gesture games (9 m)
Reach and grasp (5 m)	Understands ‘no’ (9 m)
Transfer (6 m)	Follows one-step command (12 m)
ThumbÐfinger grasp (9 m)	Points to animal pictures (19 m)
Tower of 2 cubes (16 m)	Points to 6 body parts (20 m)
Handedness (24 m)	Follows two-step command (24 m)
Scribbles (24 m)	
Tower of 4 cubes (26 m)	***Cognitive***
Tower of 8 cubes (40 m)	Shows anticipatory excitement (3 m)
Social/self-help	Plays with rattle (4 m)
Social smile (6 weeks)	Plays peek-a-boo (8 m)

Recognises mother (3 m)	Finds hidden object (9 m)
Stranger anxiety (9 m)	Pulls string to obtain toy (14 m)
Finger feeds (10 m)	Activates mechanical toy (20 m)
Uses spoon (15 m)	Pretend play (24 m)
Uses fork (21 m)	Seeks out other for play (36 m)
Assists with dressing (12 m)	
Pulls off socks (15 m)	
Unbuttons (30 m)	
Buttons (48 m)	
Ties shoelaces (60 m)	
Dresses without supervision (60 m)	

- social history
- psychological history
    - behavioural problems
    - reaction to other people and situations

PARENT-CHILD INTERACTION

It is advisable to observe carefully the parentÐchild interaction at all times, including in the waiting room. The parent's manner in talking to and handling the child will provide useful clues about possible problems related to the parent's ability to nurture the child adequately.

PHYSICAL EXAMINATION

It is convenient to consider the physical examination for two main groups:[2]

- the infant and child up to the age of 3 years
- the child from 3 years onwards

An important aspect of assessment is to note the development of the child by comparing its growth with standard developmental charts (Appendices I to IV). Developmental milestones are summarised in Table 80.1 [1] and the incidence of developmental problems under 5 years in Table 80.2.[3] The examination includes attention to any unusual appearance, which is the beginning of the process of diagnosis of the dysmorphic child.

Table 80.2 Incidence of developmental problems under 5 years (after Hutchins)[3]

More common, less severe	Less common, more severe
10Ð20% behaviour problems	3.0% intellectual handicap (IQ < 70)
10% specific learning deficits	1% intellectual handicap (IQ < 50)
10% conductive hearing loss	0.3% cerebral palsy
10% eye problems (e.g. squint)	0.2% neural tube defects
5% isolated speech problems	0.17% severe deafness
3% attention deficit disorder	0.06% blind
1% specific language disorder (e.g. comprehension)	0.1% autistic spectrum features

0.05% classic autism

VISION AND HEARING

Vision is present at birth and matures gradually to adult vision at about 12 months. Hearing is present at birth.

ACHIEVING CO-OPERATION OF INFANTS

A good aphorism is ‘Never examine the child until you have made the mother laugh’.

Children, especially if sick and irritable, can be very difficult to examine and may be most unco-operative, particularly if distressed by past experience. However, they can be readily distracted, a characteristic that the family doctor can use effectively to achieve some degree of cooperation for examination, especially for the ears, throat and chest.

Try to examine them on their parent's lap.

Children respond very positively to playing games such as a flashing light, tickling or peek-a-boo, and to any type of noise, particularly animal noises, and good humour from a friendly patient practitioner. Some doctors have strategies such as small animal images on stethoscopes to distract attention.

Distracting children[4]

In the consulting room, a small duck with a rattle inside it can be used for palpating the abdomen of young children. This seems more acceptable to them, as it becomes a game and you obtain the same information as if you had palpated with your hand. When examining the ears of young children sitting on their mother's lap, difficulty is encountered when the child follows the auroscope light and moves its head. A small rabbit or other animal on the desk which, at the press of a button under the desk, will play a drum, distracts the child to the right and enables you to get a good look into the left ear.

Similarly, a clockwork revolving musical toy over the examination couch will distract the child for examination of the ear. It is also a distraction for the general examination of children on the couch, and can become a most useful instrument.

Spatula sketches for children[5]

Many young patients have quickly forgotten any inspection of their throats while observing the preparation of a ‘present’ in the form of a drawing on the wooden spatula used for the examination.

After the examination they are informed of their special present, and you can then proceed to draw on the unused end of the spatula. The drawings take about 15 seconds. Figure 80.1 illustrates three sketches from one repertoire: a penguin (with optional bow tie), a caterpillar, and a racing car.



Figure 80.1 Spatula sketches for children

RECOGNITION OF SERIOUS ILLNESS IN INFANCY

It is vital to diagnose serious life-threatening disease in children, especially in early infancy. Certain symptoms and signs that provide a reliable indicator to such a problem are:[6]

- drowsiness
- irritability
- decreased activity (lies quietly)
- child moves eyes (rather than head) to follow you
- pallor
- vomiting (persistent)
- whimpers and lies quietly (as opposed to crying lustily)
- reduced feeding (< 50% normal intakeÑover 24 hour period)
- less than four wet nappies in 24 hours
- increased respiratory rate, may be chest wall retraction
- noisy breathing
- cold extremities
- sunken eyes
- sign: capillary refill time over 2 seconds

Serious illnesses to consider include:

- Haemophilus influenza type B infection
  - acute epiglottitis
  - meningitis
  - pneumonia
  - septicaemia
  - septic arthritis/osteomyelitis

- meningococcal infection
    - septicaemia
    - meningitis
- other forms of meningitis
- acute myocarditis
- asthma/bronchitis
- intussusception

THE CHILD AS A BAROMETER OF THE FAMILY

A disturbed child is a very common indicator of family disharmony. There is a saying that ‘love is to a child what sunlight is to a flower’.

Children need from their parents:[7]

- affection (acceptance for what they are, not for what they might have been)
- security (freedom from fighting parents, child abuse and sibling problems)
- consistent discipline
- stable figures to act as role models (parents are their heroes)
- freedom to develop a personality without emotional entanglements
- play (a need to be active and creative)
- honesty

Parents with their own problems and conflicts will have difficulty meeting the needs of their children. Family tensions, financial problems, marital disharmony, separation and divorce have a major effect on children. Parent frustration may eventuate in child abuse. Depressive illness in a parent can have a profound effect on the child.

The child's reaction to the family disharmony may manifest in three ways (with significant overlap):[7]

- behavioural problems
- psychosomatic symptoms
- school difficulties

The importance of the family doctor

The family doctor is in an important position to detect disharmony in the family through presenting problems that give subtle clues; for example, several uncharacteristic visits from parent and child, inappropriate non-verbal behaviour, such as a trembling hand or voice, or somatic symptoms incompatible with physical findings.

It is important to consider the environment of the disturbed child. Search for a possible source of disturbances at home, such as parental quarrelling, economic hardship, drug abuse including alcohol, physical or sexual abuse and maternal depression. If detected and addressed, the family dysfunction may be resolved satisfactorily.

GUIDELINES FOR FEEDING INFANTS

Starting rules for parents

- It is best to breastfeed for the first 12 months.
- Cow's milk-based formulas should be used if baby is not breastfed until baby is at least 12 months.
- Cow's milk should ideally be left until 12 months.
- In the first3-4 months ‘baby knows best’.
- Formula choice for healthy term infants can be based on cost.

- The rule is 150 ml/kg per does.
- The only reliable measure of adequate nutrition is weight gain.
- Your baby only needs breast milk or formula for the first 5-6 months.
- It is good to introduce soft solid foods form 5-6 months but introduce them slowly.
- Babies don't need teeth to chew soft foods.

Starting solid foods

Solids should be gradually introduced at about 5 - 6 months, one at a time. Food should never be forced but introduced slowly.

Solids should be offered after a feed or between feeds of milk. Breast milk or formula remains the most important food.

Examples of solid foods for beginners are:

- baby rice cereal mixed with their usual milk or cooled boiled water (best first option)
- cooked pumpkin, potato or carrot
- fruits such as banana, cooked apple or pear

The texture should be pureed (no lumps).

Introduce a new food only after 3-4 day, early in the day, and check for any allergic reaction. Start with 1-2 teaspoons of solids and build up to 3 meals a day at your baby's own pace.

Lumpy food can be introduced at 6 - 9 months, as by this time babies learn to chew.

By 9-10 months more solids should be eaten each mealtime and the milk gradually decreased. Cooked boiled water should be introduced as it is better than fruit juices.

From 12 months onwards cow's milk can be introduced and more solid foods, especially meats, vegetables and fruits.

Note: Babies on a cow's milk diet who eat little are prone to iron deficiency anaemia.

GUIDELINES FOR TOILET TRAINING

As a rule children will learn to use the toilet when they are ready.

The age that most are fully trained are:

- daytime - between 2-1/2 and 4 years
- night-time - by 8 years of age

Once they start training it usually takes 4 months before they are dry, but some can take several months.

General rules for parent

- Be relaxed about toilet training.
- Avoid rushing toilet training.
- Do not force your child to go to the toilet.
- Nagging does not work well.
- Punishing well not work.

Best times to sit child on the toilet

- First things in the morning

- After meals
- When you sense their need to go
- Before going out
- Upon returning home

Key points for parents

- Use a potty or toilet with a seat ring and a step.
- Explain the process in simple terms.
- They will learn to use the toilet when they are ready.
- Sit both boys and girls on the toilet to pass urine.
- Do not force them if they refuse to use it.
- Make a fuss of success - praise and reward them.
- Help the child relax on the toilet.
- Stop using nappies (except when sleeping).
- If the training upsets them, wait for a month and try again.

PERSONAL HEALTH RECORD

Many family practices issue a personal health record (PHR) to the child's parents as a means of improving health care delivery, including the enhancement of preventive care. The PHR, also referred to as the 'parent-held health record' or 'health passport', is usually distributed by hospitals when the baby is born and research has shown it to be well received by both health practitioners and parents.[8]

The PHR is a small, loose-leaf booklet with a sturdy plastic cover which can be easily carried around by the parent. The contents can vary from one producer to another but generally it contains:

- records of birth details and newborn examination
- percentile charts for weight gain
- visual check
- hearing check
- developmental check
- immunisation schedules and recordings
- progress notes
- advice on accident prevention (Table 80.3)

Table 80.3 Accidents don't have to happen

Six to eighteen months
Have cupboards made child-resistant for the storage of medicinals and household chemicals. Pesticides and petroleum products should be locked away in the shed. Don't store in ordinary food and drink containers.
Fires and radiators should be adequately guarded.
Cords on electrical food and drink heaters need to be shortened or hooked up out of a toddler's reach. Do not use tablecloths. Put hot food and drinks into the centre of the table. Take care with buckets of hot water.
Fit dummy plugs in unused power points.
From 9 kg (20 lb) body weight, baby's car rides should be in an ASA-approved child seat.
Supervise your toddler at all times in or near water. The swimming pool needs to be adequately fenced.
Keep matches in the child-resistant cupboard in the kitchen. Put scissors, needles and pins well out of reach.
Have the play yard safely fenced from the street.

Parents: Walk right round the car before reversing down the drive, or place your child in the car first.

Do not allow your small child to be unsupervised in the bathroom.

Never give the child nuts to eat because it cannot chew them properly. Peanuts present a particular hazard because of their shape and hardness. They can cause the child to choke.

- other health educational material

The PHR provides a very practical method of promoting communication between various health professionals involved in the child's care and also between the family and their doctor. It promotes the concept of ‘self-care’ by encouraging a sense of responsibility by parents for the child's health and is also a medium for enhancing preventive care, especially with immunisation, hearing tests and development.[8]

Immunisation shedule

Refer to chapter 9, page 78.

DIAGNOSTIC TRIADS FOR CHILDREN

The following are a selection of childhood disorders. The equals sign represents a pointer to the diagnosis.

AcuteĐsubacute onset

- arthralgia (lower limbs) + rash (buttocks, legs) ± abdominal pain = Henoch-Schnlein purpura
- respiratory symptoms + vomiting + mental changes ± seizure/coma = Reye's syndrome
- pallor + drowsiness + fever = meningitis
- pallor + abdominal pain (severe and intermittent) + inactivity = intussusception
- malaise + fever + polyarthritis (flitting) ± rash ± chorea = rheumatic fever
- (< 12 months): drowsiness + cough + wheezing = bronchiolitis
- (< 3 months, (usually male): weakness + weight loss + vomiting (severe, intermittent) = pyloric stenosis
- (neonate): vomiting (after first feeds) + drooling + abdominal distension = oesophageal or duodenal atresia
- malaise + pallor + bone pain = acute lymphatic leukaemia
- malaise + pallor + oral problems (gingival hypertrophy, bleeding, ulceration) = acute myeloid leukaemia
- abdominal pain + pallor + a/n/v = acute appendicitis
- abdominal pain + malar flush + fever ± URTI = mesenteric adenitis
- (during gastroenteritis epidemic): malaise (extreme) + diarrhoea + severe abdominal pain = haemolytic uraemic syndrome
- drowsiness + tachypnoea + chest wall recession = pneumonia
- drowsiness + fever + purpuric rash = meningococcal infection
- lethargy (extreme) + drooling saliva + snoring stridor = acute epiglottitis
- URTI + brassy cough + inspiratory stridor = croup
- coughing + wheezing + chest wall recession = asthma or aspirated foreign body
- fever + conjunctivitis + skin changes (cracked red lips, maculopapular rash, erythema of palms/soles, desquamation fingertips) = Kawasaki's syndrome

Developmental delay

- large ears + long narrow face + large genitals = fragile X syndrome
- small extremities (hands, feet, genitals) + narrow forehead + eating disorder = Prader-Willi syndrome
- ‘elfin’ face + low set ears + cardiac murmur = Williams syndrome
- (female): short + webbed neck + pigmented naevi ? cardiac disorder = Turner's syndrome

- short + webbed neck + facial disproportion (broad forehead, ptosis, low set ears etc.) ? cardiac disorder = Noonan's syndrome

Chronic

- fever (FUO) + abdominal mass + haematuria (uncommon) = Wilm's tumour
- malaise + abdominal pain (vague) + abnormal behaviour = lead poisoning
- (< 2 years): lethargy + irritability + pallor = iron deficiency anaemia or thalassaemia major
- fever + malaise (extreme) + a/n/v ? anaemia = neuroblastoma
- headache + a/n/v + ataxia = medulloblastoma

Older children

- (male): ? snorting, blinking etc. + oral noises (e.g. grunts, hisses) ? loud expletives = Tourette's syndrome
- irritability + disorders of appetite + abdominal pain (vague) = substance abuse (nicotine, heroin etc.)
- mid to low back pain/discomfort + inability to touch toes + kyphosis = Scheuermann's syndrome
- knee pain (after activity) + tender knee ‘lump’ + pain on kneeling = Osgood-Schlatter disorder
- (adolescent): limp + knee pain + hip pain = slipped capital femoral epiphysis



Figure 80.2 The normal child can sit without support form 6 - 9 months (avrage 8 months)

REFERENCES

1. Jarman FC, Oberklaid F. The detection of developmental problems in children. Aust Fam Physician, 1992; 21: 1079 - 88.
2. Robinson MJ, Roberton DM. Practical Paediatrics (5th edn). Melbourne: Churchill Livingstone, 2003;22 - 23.
3. Hutchins P. The young child with developmental problems. Aust Dr Weekly (How to treat series), 1990; 20 July: I - viii.
4. Trollor J. Distracting children. Aust Fam Physician, 1987; 16: 1372.
5. Malcher G. Spatula sketches for children. Aust Fam Physician, 1990; 19: 1441.
6. Hewson P. Recognition of serious illness in early infancy. Aust Paed Review, 1992; (6): 1.
7. Connell HM. The child as a barometer of the family. Aust Fam Physician, 1980; 9: 759 - 63.
8. Jeffs D, Harris M. The personal health recordÑmaking it work better for general practitioners. Aust Fam Physician, 1993; 22: 1417 - 27.

CHAPTER 81. SPECIFIC PROBLEMS OF CHILDREN

Children are not simply micro-adults, but have their own specific problems.

BELA SCHICK (1877 - 1967)

SPECIFIC PROBLEMS OF CHILDREN

The family doctor usually treats children for common minor complaints such as skin disorders and respiratory infections, and for preventive measures such as immunisation. However, in many instances, parents consult their doctor for advice on normal or abnormal behavioural disorders so doctors need to be well versed in normal behaviour in order to provide appropriate advice and reassurance. Many of these everyday problems are discussed in this chapter. Childhood management of problems such as constipation, anaemia, diarrhoea and cough are included in the relevant chapters on problem solving.

CRYING AND FUSSING IN INFANTS

Crying and fussing is a common problem in the first 3 months but is now considered to be a normal physiological aspect of a maturing central nervous system.[1]

Crying is excessive if it lasts for long periods when the baby should be sleeping or playing. It usually occurs at 6 - 9 pm. In most instances an organic cause, such as infection, milk allergy or reflux cannot be demonstrated.

Parents should be made aware of a checklist of common causes:

- hunger (underfeeding is the main feeding problem causing crying)
- wet or soiled nappy
- loneliness: crying usually ceases when the baby is picked up
- infant colic: a possibility at 2 - 16 weeks
- individual temperament
- teething (more likely to cause discomfort after 12 months)
- reflux oesophagitis

Management

- Perform careful physical examination, including assessment of child's temperament.
- Give parental reassurance and education.
- Reassure parents that extra attention will not affect the baby but overstimulation should be avoided.
- Provide soothing alternatives (e.g. use of dummy - pacifier, extra cuddling and carrying, gentle massage).

Practice tip - the role of 5 Ss

1. Swaddling - firm clothing, not too loose
2. Lie baby on Side or Stomach
3. Shush, i.e. ‘sshhusshhing’ as loud as the child
4. Swing - sway them from side to side
5. Suckling - nipple, teat or dummy

Diet and medications do not have a significant place in management.

Infantile colic

Refer to pp. 318 - 9.

Typical features

- Baby 2 - 16 weeks old, especially 10 weeks
- Prolonged crying - at least 3 hours
- Crying during late afternoon and early evening
- Child flexing legs and clenching fists because of the ‘stomach ache’

Management

- Reassurance and explanation to the parents
- Pacifying methods

Medication

Avoid medications if possible, but consider:

- simethicone preparations (e.g. Infacol wind drops) *or*
- dicyclomine syrup

Teething

Baby teeth (milk or deciduous teeth)

- Babies usually cut their teeth from age 6 months until 2 - 3 years.
- The first teeth to appear (which seldom cause discomfort) are the lower incisors (during first year).
- The first and second molars (ages 1 - 3) tend to cause problems.
- Usually the first set (20 teeth) is complete soon after the second birthday (Figure 81.1).



Figure 81.1 The lower set of primary teeth with average times of eruption

Symptoms

- The gum is slightly swollen and red.
- This may cause little or no discomfort but may be quite painful.
- The child is more clinging, fretful and dribbling than usual.
- Chews on something such as fingers.
- Irritability and crying (on and off for no more than a few days).
- Difficulty with sleeping.

Note: Teething does not cause fever.

Treatment

Reassure the parents that the problem will soon settle.

Soothing methods

- Gentle massaging of the gum with the parent's forefinger wrapped in a soft cloth or gauze pad is comforting. A gel such as Oro-Sed can be massaged into the gums every 3 hours if extremely troublesome. *or*

- Allow the baby to chew on a clean, cold, lightly moistened facewasher (a piece of apple can be placed inside the facewasher). *or*
- Give the baby a teething ring (kept cold in the refrigerator) or a teething biscuit.

Medication

Medicine is usually not necessary for teething. Paracetamol mixture should be used for any discomfort. For more severe problems, especially affecting sleep, an antihistamine can be given at night or a combined mixture of antihistamine and analgesic.

Pitted dark teeth

Some children who are breastfed for long periods (e.g. 3 years) may develop unsightly pitting of the front surface of their teeth. This will not go away but parents should be reassured that the adult teeth will be normal when they appear.

THUMB SUCKING

Thumb sucking involves placing the thumb or finger on the roof of the mouth behind the teeth (hard palate) and sucking with the mouth closed. It is basically a habit and should be regarded as normal. It is one of the first pleasurable acts that the infant can manage. It occurs in children up to the age of 12 years but is most common under the age of 4 years. It usually settles by age 6 or 7. If it persists beyond this age it can cause problems with the permanent teeth, which begin to appear at about age 7. One effect is that the pressure on the front teeth may cause protrusion of these teeth (i.e. buck teeth).

Prevention

Provide alternative comfort measures, such as a dummy (pacifier) if the habit is developing. If it persists, avoid making an issue and drawing attention to it.

Treatment (advice to parents)

- No special diet or medication is necessary.
- For a child over 6 years, carefully observe trigger factors and find ways of avoiding them. Provide extra attention and organise pleasant distractions.
- Help the child explore other solutions.
- Give praise and rewards for efforts to stop.

Referral for specialised treatment may be necessary for a persistent problem.

SNUFFLING INFANT

Snuffles occur in about one in three normal babies in the few weeks after birth and is not a problem un less it affects feeding.

Snuffling in older infants is usually caused by rhinitis due to an intercurrent viral infection. The presence of yellow or green mucus should not usually be cause for concern.

Treatment

Reassure the parents.

- Paracetamol mixture or drops for significant discomfort.
- Get the parents to perform nasal toilet with a salt solution (1 teaspoon of salt dissolved in some boiled water); using a cotton bud, gently clear out nasal secretions every 2 waking hours.

- Once the nose is clean, saline nose drops or spray (e.g. Narium nasal mist) can be instilled.
- Stronger decongestant preparations are not advised unless the obstruction is causing a significant feeding problem, when they can be used for up to 4 - 5 days.

BLOCKED NASOLACRIMAL DUCT

About 20% of infants develop watery eyes, but most resolve by 12 months. Excessive eye watering in infants is the key sign that there is inherited narrowing of the nasolacrimal ducts (Figure 81.2a). It usually becomes obvious in infants between 3 and 12 weeks and affects one or both eyes. Mucus and muco-pus may appear in the tears. The discharge is worse on waking. In some infants, watering and discharge soon after birth indicate that the ducts have failed to open. Infection may intervene (refer to p. 560) and conjunctivitis can be problematic.


(a)
watery red eye
blockage can occur at any of these points in the tear collecting ducts
nasolacrimal duct
(b)
direction of finger strocking

Figure 81.2 (a) Blocked nasolacrimal duct **(b)** Nasolacrimal duct blockage: method of massage

Outcome

In most cases the problem improves spontaneously. Self-correction usually occurs from 6 months of age onwards or even earlier.

Treatment

The mother or baby's carer should massage the drainage ducts several times daily. This is done by firmly placing the tip of the little finger over the inside corner of the eye and stroking firmly downward to the tip of the nose (Figure 81.2b).

Minor infection can be treated with warm cottonwool soaks. For more severe blockage or when eye watering has not settled by 12 months, the ducts should be probed and dilated under light anaesthesia, followed by irrigation with saline.

Very rarely, an artificial duct will need to be fashioned surgically.

GROWTH AND DEVELOPMENT PROBLEMS

Recommendations for routine growth monitoring

- Check weight regularly.
- Check height/length, e.g. 18 - 24 m, 5 years.
- Check head circumference:
  - neonate
  - 6 - 8 weeks

Failure to thrive (FTT) Refer Appendices I - IV

The long list of possible causes includes malfunction of any of the organ systems of the body as well as nutritional, environmental, social and psychological factors.

FTT is best determined by the important health and nutrition procedure of anthropometry (i.e. sequentially plotting the weight, length and head circumference on growth charts, see growth charts in Appendix). The infant with FTT has a decreased growth rate or is losing weight, and may be below the 3rd percentile. The percentile charts mean little without considering the context of the baby's growth (e.g. premature babies, children of small parents).

On an average, babies put on 150 - 200 g a week.[2] A doubling of birth weight is achieved by age 4 - 5 months. A classification of FTT is presented in Table 81.1, divided into organic and non-organic causes. The distinction is not always easy. Psychosocial problems may coexist with organic problems. Feeding problems are common to both.

Table 81.1 Failure to thrive: general causes (after Robinson)[4]

Non-organic causes
1 Inadequate parenting
2 Poor nutrition
Organic causes
1 Failure of intake
• underfeeding (e.g. nipple disorders)
• congenital abnormalities (e.g. cleft palate)
• dyspnoea (e.g. congenital heart disease)
• neurological lesions (e.g. cerebral birth injuries)
• behavioural factors
2 Abnormal losses
• vomiting (e.g. pyloric stenosis, galactosaemia)
• stools (e.g. steatorrhoea)
• urine (e.g. kidney disease)
3 Failure of utilisation
• chronic infection (e.g. cystic fibrosis)
• metabolic disorders (e.g. phenylketonuria)
• endocrine disorders (e.g. hypothyroidism)
• constitutional (e.g. Down syndrome)
4 Sleep apnoea

Non-organic failure to thrive

Non-organic FTT can be caused by emotional deprivation or by poor nutrition from inadequate intake. Emotional deprivation might be anticipated by a knowledge of the mother, her family background, marital relationships, attitude to the pregnancy, delivery and early bonding experience. In her book *The Abused Child*, Martin lists factors influencing such bonding.[3]

Factors in the parent

- Expectation of the child
- Desire for the child
- Capacity to give
- Ego-strength to adapt to stress
- Ability to accept imperfection
- Realistic fantasies of the child

Factors in the child

- Absence of defects
- The ability to match the parent's expectations
- Good health
- Loving behaviour, including smiling, cuddling and thriving

Disturbance of any of these factors may lead to relationship difficulties. The management of FTT due to psychological factors may be complex.[5] At the simplest level the recognition by the mother that she is having difficulty in relating to her baby is essential, and the reassurance that not all babies are as lovable and easy to manage as portrayed may help. A home visit to evaluate the home environment will provide invaluable information. These mothers require considerable caring support and encouragement.

Organic failure to thrive

Any chronic disease will cause FTT. Serious organic diseases include kidney failure, hypothyroidism, cystic fibrosis, other causes of malabsorption such as coeliac disease, and various inborn errors of metabolism such as galactosaemia (Table 81.1).

Poor developmental progress may indicate intellectual disability. Babies born to mothers who are HIV carriers present with FTT in the first 5 months, with or without other signs of disease, such as infections.[2] Another possible cause of growth failure in a baby who has a good intake may be sleep apnoea and this requires investigation.

Examination of the baby

Examine for developmental problems, including cerebral palsy, cleft palate, respiratory disorders and abdominal abnormalities.

Investigations

Simple screening tests should be performed if either the history or physical examination suggests organic disease. Tests include routine blood counts, urinalysis and urine culture, Guthrie test for PKU, IVP, thyroid function tests and chromosomal and hormone analysis. It is important to exclude urinary tract infection.

Main causes of FTT (account for up to 90%):[6]

- normal variants, and
- nutritional deprivation

Most important considerations

- Manner of feeding
- Home visit
- Environmental factors
- Parental problems

- Admission to hospital

Rare possibilities

- HIV infection
- Sleep apnoea
- Hypopituitarism (growth hormone ↓)
- Chromosomal abnormalities

Management

As the diagnosis is made on the history, extensive investigations are inappropriate. It is useful to watch the child feed and assess their degree of hunger. Consultation with the child's child health/infant welfare nurse where applicable is usually very helpful. Straight-forward problems will respond to careful dietary advice. Enlist the expertise of an early childhood intervention support service. More difficult problems may require admission to hospital for supervised feeding and parental education.

SHORT STATURE

Short stature is generally considered to be a psychosocial and physical handicap if the definitive height is in:[6]

- males < 162.6 cm (5′4″)
- females < 152.4 cm (5′0″)

The three major growth factors are genetic, nutritional and hormonal. The hormones that are essential for a normal growth process are growth hormone and insulin-like growth factor 1 (the key), thyroxine, cortisol and sex steroids.[6]

In general, assessment is important to differentiate between normal physiological variants of growth, namely familial short stature, and constitutional growth delay and pathological causes.

Ten essential questions to consider:[6,7]

1. Is the child actually short?
2. Is the child short compared with other children (i.e. below the 3rd percentile)?
3. Is the child unexpectedly short from a genetic viewpoint?
4. Is the child's growth slowing (i.e. falling behind the height percentile)?
5. If the growth is slow, what is the reason?
6. How does the child feel about the short stature?
7. How does the height percentile match against a growth velocity (GV) chart?
8. Has puberty commenced?
9. Is there any specific investigation warranted?
10. Is there any specific therapy warranted?

Causes to consider:

1. Constitutional delay in maturation - a common and normal variant in which the growth spurt is later than average. Bone age is delayed.
2. Familial short stature - this follows the family trend of a genetically small family. Skeletal proportions and growth velocity are normal. Determine parental height.

Note: The lower segment should be > ½ the height beyond the age of 8 years.[7] Bone age is equivalent to chronological age.

Rough rule for expected adult height based on parental height

- Boys - mean of parent's heights plus 7 cm

- Girls - mean minus 7 cm

1. Organic causes - of the many causes some are rare but serious conditions, such as coeliac disease, Crohn's disease and chronic kidney failure may present with slow growth as the only abnormal sign. Such disorders should be suspected in children who have unexpected abnormal GV.

Physical examination

- General inspection includes dysmorphic features and nutritional status. Measure all anthropometry (height, weight, GV, upper/lower segment ratio) and compare with percentile charts
- Measure skeletal proportions
- Assess pubertal status

Investigations

If the GV is < 25th percentile for bone age, consider:

- thyroid function tests
- FBE and ESR (↑with Crohn's)
- coeliac disease tests (e.g. antiendomysial antibodies)
- chromosomes in all girls (only) - regardless of appearance. Karyotype to exclude Turner's 45 XO
- growth hormone studies - 1. exercise test and 2. a glucagon stimulation test
- kidney function
- bone age X-ray (left wrist and hand) - to compare bone age with height age: if equivalent, this negates the need for growth hormone studies

Management[7]

Referral to a consultant is appropriate. The growth promoting agent is recombinant growth hormone given by subcutaneous injection, which is very expensive.

Criteria for treatment:

- height below the first percentile
- GV < 25th percentile for bone age
- bone age under 13.5 years for girls or under 15.5 years for boys

Beneficial results in:

- growth hormone deficiency
- growth retardation secondary to kidney insufficiency
- Turner's syndrome (can improve 8 - 10 cm)

Dosage: somatropin 14 - 22 U/m^2 per week divided into 6 - 7 doses/week.

TALL STATURE

The estimated mature height is, for:

- females - 182.9 cm (6′)
- males - 193.1 cm (6′4″)

It is not a common presenting childhood problem in general practice.

Causes:

- familial
- precocious puberty

- growth hormone excess (pituitary gigantism)
- hyperthyroidism
- syndromic: Marfan, Klinefelter's, XYY males, homocystinuria

Management

As tall stature is now more socially acceptable, reassurance, counselling and education may alleviate the family's concerns. If treatment is considered appropriate, high-dose oestrogen is used in very tall girls (accelerates epiphyseal maturation and reduces final height) while high-dose testosterone is used for boys. The management should be undertaken by experienced therapists. The ideal time to commence hormone therapy is just after the appearance of the first pubertal changes.[7]

OBESITY IN CHILDHOOD

Obesity, which is defined as > 120% of expected weight per height, is an increasingly prevalent disorder, with up to 20% of Australian children being considered as overweight.[7] Obesity in childhood increases the risk of adult obesity. Nutritional obesity is associated with growth acceleration and advancement of bone age, while endocrine obesity has the opposite effect.[7] Refer to Chapter 78 (pp. 850 - 1).

DELAYED PUBERTY

This is the absence of pubertal development in:

- girls > 14 years
- boys > 15 years

Significant causes include:

- constitutional delay (usually familial and the commonest cause) - associated with delayed growth and bone age
- chronic illness (e.g. severe asthma, cystic fibrosis, kidney failure)
- poor nutrition and exercise
- anorexia nervosa

Other less common causes include chromosomal abnormalities (e.g. Turner's syndrome) and gonadal failure.

Investigations

Delayed puberty, either simple familial or constitutional, usually does not warrant investigation. If in doubt a bone age X-ray is useful. Otherwise the following can be considered:

- FBE and ESR
- kidney function
- thyroid function tests
- chromosomal analysis (usually in girls)
- serum FSH, LH, prolactin, testosterone

Management

- Treat according to findings (e.g. chronic asthma), otherwise referral to a paediatric endocrinologist.
- Treatments include testosterone 100 - 500 mg IM every 2 - 4 weeks in boys or oestradiol in girls.

PRECOCIOUS PUBERTY[6,7]

This is the appearance of true puberty in:

- girls < 8 years
- boys < 9.5 years

It is about four times more common in girls than boys and girls are less likely to have underlying pathology (e.g. hypothalamic hamartoma) than boys.

Clinical presentations include early secondary sexual characteristics, accelerated GV, mood disorders and inappropriate sexual behaviour.

If considered appropriate investigations include serum FSH and LH, gonadal steroid (testosterone or oestradiol), bone age and MRI of cranium if FSH or LH increased.

Management

- If the condition is evolving slowly no treatment is needed.
- If concerned refer to a paediatric endocrinologist.

PREMATURE THELARCHE[6]

This is the isolated breast development in girls under 8 years old. It occurs in girls under 2 years and spontaneous regression can be expected. It may present at birth. Observation with reassurance for this benign condition is appropriate.

PREMATURE ADRENARCHE[6]

This is the isolated appearance of pubic hair (usually in a girl) under 8 years old. There are no other features of virilisation or oestrogenisation. It is usually a normal variant (no specific treatment available) but may signify atypical congenital adrenal hyperplasia. Referral is indicated if any concerns.

PUBERTAL GYNAECOMASTIA

This should be considered as a normal variant of puberty. It is common in normal adolescent males, with a prevalence of about 40 - 50% and is a transient phenomenon.

There is a palpable disc of breast tissue, which in boys may feel quite firm and needs to be distinguished from adipose tissue. There is no hormonal or medical treatment and surgical removal should rarely be necessary.

ASYMMETRICAL BREAST DEVELOPMENT

This may occur in both sexes - in males being a variant of pubertal gynaecomastia and in females part of a normal development process - and reassurance that the breast sizes will normalise in time is all that is required. If the discrepancy persists and is causing psychological problems, further strategies, such as prostheses or reconstructive surgery, may be advisable.

GROWING PAINS (BENIGN NOCTURNAL LIMB PAIN)

'Growing pains' is a term often used inappropriately for diffuse aches and pains in the legs of children.

Features[8]

- Typical age 3 - 7 years, may start at 2 years

- Positive family history
- Pain wakes child - usually distressed
- Poorly localised in leg - knee, shin, calf (Figure 81.3)



Figure 81.3 Growing pains: typical sites of pain

- May affect the arms (uncommon)
- Lasts 20 - 30 minutes, regardless of treatments
- May recur during the night
- Normal examination
- No associated symptoms
- No pain or disability next morning (runs ok)
- Very active days may lead to a ‘bad’ night

Management

- Problem resolves spontaneously in time
- Reassurance
- Consider analgesic and heat packs (usually unsuccessful)
- Massage is a reasonable option - appears to help
- Check ESR if in doubt

DEVELOPMENTAL DISABILITY AND DELAY

The family doctor is in an ideal position to recognise and initiate evaluation of the child with a developmental disability, whether it is a physical disability or delay, an intellectual disability or a learning disability.

Many developmental problems will be obvious but others, such as fragile X syndrome, are subtle. Several disabilities may evade a diagnosis. Success depends on the ability of the family doctor, early referral and the level of sophistication of genetic services.

Transient developmental delay may be associated with factors such as prematurity, family stress, physical illness and learning opportunities, while persistent delay can be caused by intellectual disability, cerebral palsy, autism, and hearing and visual impairment.

There are many rare dysmorphic syndromes which are becoming more recognised and referral to genetic disorder units will help in appropriate evaluation.

Evaluation

An appropriate history includes a careful look at developmental milestones and family history. The physical examination includes testing of hearing and vision.

Investigations to consider:

- screen for congenital infection (e.g. rubella, toxoplasmosis, CMV)
- chromosome (karyotyping) studies
- urinary metabolic screen (e.g. PKU)
- creatinine phosphokinase in males
- thyroid function studies
- DNA specific tests (e.g. fragile X, Prader-Willi, Williams syndrome)
- CT scan or MRI

Intellectual disability

This is regarded as a component of developmental disability and refers to significant substandard intellectual functioning (2 SDs < mean IQ) with deficits in adaptive behaviour and is manifested during the development period. Presentation includes learning difficulties, language delay and behavioural problems.

The two most common causes are trisomy 21 and fragile X syndrome.

Fragile X syndrome, Prader-Willi syndrome and Williams syndrome are presented in Chapter 19.

Cerebral palsy

Definition

A persistent motor disorder of movement and posture resulting from prenatal developmental abnormalities or perinatal or postnatal CNS damage (to the immature brain). Features include spasticity, ataxia and involuntary movements.

Facts

- Cerebral palsy is not a diagnosis but a diverse group of disorders.
- In most cases the cause is unknown.
- Fewer than 1 in 10 cases result from hypoxia.
- Prevalence is about 2 per 1000 live births.

Classification of syndrome

- The type of motor disorder (e.g. spasticity - 70% of cases, athetosis, ataxic, mixed)
- The distribution (e.g. hemiplegia, paraplegia, diplegia)
- The severity of the motor disorder

Associated disorders

- Seizures (30% of cases)
- Visual problems (e.g. strabismus)
- Hearing defects
- Intellectual disability - may be normal intelligence
- Perceptual problems
- Hyperactivity
- Short attention span

Diagnosis

This is rarely made during infancy. Formal diagnosis usually made by 2 years following referral.

Management

- Accurate diagnosis
- Genetic counselling
- Education materials
- Assessment of child's capabilities
- Referral to several agencies for assessment (e.g. hearing, vision, dietitian, speech pathologist, various other allied health professionals)
- Monitor problematic areas (e.g. constipation)
- Refer to a multidisciplinary team such as a cerebral palsy clinic (major hospital); includes orthopaedic assessment with special attention to legs (e.g. hips, knees, hamstrings)

Specific learning disabilities

A specific learning disability (SLD) is an unexpected and unexplained condition, occurring in a child of average or above intelligence, with a significant delay in one or more areas of learning. These areas include reading, spelling, writing, arithmetic, language (comprehension and expression), attention and organisation, co-ordination and social and emotional development. An SLD can vary from very mild to quite severe. It may, in turn, cause a general learning disability. The primary cause is unknown.

Diagnosis

If undetected by parents, any undisclosed SLD will soon be detected in the classroom. Sometimes the disability is not detected until later (8 years or more) when more demanding schoolwork is required. Speech delays, reading difficulties and calculation problems are among the first signs. It is important to check hearing and vision. These children may also present with a behaviour disorder as they are often subject to ridicule by other children and tend to develop a poor self-image and low self-esteem.

Dyslexia

The word ‘dyslexia’ is derived from the Greek term meaning ‘difficulty with words’. The condition was originally called ‘word blindness’, referring to an SLD with reading. It affects about 4% of the population. Dyslexic children have a normal IQ and no physical problems, but their reading skills are below average. Other SLDs may also be present, particularly in spelling, writing and clear speaking.

The two main features are reading and spelling difficulties because dyslexic children confuse certain letters whose shape is similar, perhaps a mirror image of each other (e.g. confusing *b* with *d* and *p* with *q*). This means that affected children cannot properly use and interpret the knowledge they have acquired.

Characteristics include:

- a reluctance to read aloud
- a monotonous voice when reading

- following the text with the finger when reading
- difficulty repeating long words

These features, of course, are seen in all or most learners but, if they persist in a bright child, dyslexia should be considered. The most important factor in management is to recognise the problem and the earlier the better.

Refer children with:

- poor educational achievement, especially with learning to write, read and spell
- difficulty answering questions on paper
- a lack of understanding of time and tense
- poor concentration in reading and writing

Management of specific learning disabilities

It is important to build the child's self-esteem by explaining the problem carefully, removing any sense of self-blame and encouraging efforts towards progress. Parents can play an important role in building up their child's self-esteem and in helping learning. Parents are the most important teachers.

Children with SLDs are usually referred to an experienced professional or to a clinic (e.g. a dyslexia clinic) for assessment. Management may involve a clinical psychologist, an audiologist, an optometrist or a speech pathologist. A specific method of correcting the problem and promoting learning will be devised. It is also worthwhile seeking the help of a support organisation.

CHILDHOOD CARDIAC MURMURS

Innocent murmurs

Many children and infants will be found to have systolic murmurs on routine examination, especially in the presence of a fever. The majority, which will be innocent or physiological, are found in asymptomatic children and due to normal turbulence of flow within the heart and great vessels.

History

Indicators:

- innocent murmurs - asymptomatic
- significant serious murmurs
  - age less than 12 months
  - other congenital abnormalities
  - symptoms of cardiac dysfunction (e.g. cyanosis, dyspnoea)

Physical examination

Innocent murmur characteristics:

- mid-systolic
- has a musical quality
- best heard between left sternal edge and apex
- soft, grade 2/6
- quieter sitting up
- associated normal heart sounds
- usually ejection in quality
- variable with change in posture or exercise or respiration

Note: Exclude a wide or fixed splitting of second heart sound (VSD, ASD).

Diagnostic tests

- Chest X-ray
- ECG
- Echocardiography

Note: A venous hum is a continuous roaring noise usually heard in the neck above the clavicles, especially the right side. It is loudest when the child is sitting up and disappears when supine. It is of no pathological significance.

When to refer

- Murmurs heard in infancy, especially the first 6 months
- Relevant symptoms (e.g. cardiac symptoms, feeding difficulties, poor growth)
- Doubt about diagnosis

ENURESIS

Enuresis can be defined as daytime wetting (diurnal enuresis) after age 4 years or night-time wetting (nocturnal enuresis) after 6 years.[9] These are primary enuresis and appear to be due to delayed maturation of achieving bladder continence. Secondary enuresis is wetting after normal continence of at least 3 months.

DSM-IV diagnostic criteria for enuresis

1. Repeated voiding of urine into bed or clothes
2. Clinically significant: frequency twice weekly for at least 3 consecutive months or significant distress in social or other important areas of functions
3. Chronological age at least 5 years
4. Behaviour not due to physiological effect of a substance (e.g. diuretic) or a general medical condition

Nocturnal enuresis

Nocturnal enuresis refers to the involuntary passage of urine during sleep in the absence of any identified physical abnormality in children (or adults) at a time when control of urine could reasonably be expected (usually the age of 5).

What is normal?

Bed-wetting at night is common in children up to the age of 5. About 50% of 3 year olds wet the bed, as do 20% of 4 year olds and 15% of 5 year olds. It is considered a problem if regular bedwetting occurs in children of 6 years and older, although many boys do not become dry until about 8 years. About 2% of 14 year olds are affected.[10] Of children with diurnal enuresis, 60% also have nocturnal enuresis while only 10% with nocturnal enuresis also have diurnal enuresis.

Aetiology

There is usually no obvious cause and most of the children affected are normal in every respect but seem to have a delay in development of bladder control. Others may have a small bladder capacity or a sensitive bladder. It tends to be more common in boys and has a genetic predisposition. The cause of secondary enuresis can be psychological; it commonly occurs during a period of stress or anxiety, such as separation from a parent or the arrival of a new baby.

Underlying disorders to be excluded:

- urinary tract infection
- diabetes mellitus

- diabetes insipidus
- neurogenic bladder
- urinary tract abnormality

After the age of 6, investigations including an intravenous urogram or ultrasound are necessary to exclude urinary tract abnormalities.

Advice for parents on managing the child

If no cause is found, reassure the child that there is nothing wrong and that it is a common problem that will eventually go away (spontaneous resolution rate is 15% per year). There are some important ways of helping the child to adjust to the problem.

- Do not scold or punish the child.
- Praise the child often, when appropriate. Consider using a star chart whereby the child places a sticky star on it for each dry night.
- Do not stop the child drinking after the evening meal.
- Do not wake the child at night to visit the toilet.
- Use a night-light to help the child who wakes.

Some parents use a nappy to keep the bed dry, but special absorbent pads beneath the bottom sheet are more appropriate. Make sure the child has a shower or bath before going to kindergarten or school.

Treatment

Many methods have been tried, but the bedwetting alarm system is generally regarded to be the most effective. If the child has emotional problems, counselling or hypnotherapy may be desirable. Enuresis clinics currently favour two trials of the alarm system and, if persistent, desmopressin acetate nasal spray.

The bed alarm. There are various types of alarms: some use pads in the pyjama pants and under the bottom sheet, but recently developed alarms use a small bakelite chip, which is attached to the child's briefs by a safety pin. A lead connects to the buzzer outside the bed, which makes a loud noise when urine is passed. The child wakes, switches off the buzzer and visits the toilet. This method is based on a conditioned response to the release of urine. It is important to make sure that both the mat and alarm are properly functional. Persist with this method for a least 3 months. Alarms can be obtained through enuresis clinics, pharmacies and community health clinics; works especially well in older children.

Desmopressin acetate. This is the treatment of choice after failed trial of bed alarm. The dose is one spray to the lower part of each nostril per night. Avoid water loading before bed. It is very useful for children in school camps.

The persistent problem. For the 1 - 2% of patients whose bed-wetting persists beyond adolescence, a formal urodynamic assessment is advisable. Many of these patients also have daytime symptoms.

Stepwise management trial

1. Conditioning therapy
    - pad or bell alarm *or*
    - body worn alarm (e.g. Malem night-trainer) If a trial fails, have a 3-month break and try again with close supervision
2. Desmopressin acetate nasal spray to lower part of nostril each night (avoid water load before bed)
3. Desmopressin + alarm
4. Planned waking

Diurnal enuresis

Suggested management program[9]

- Urinary containment exercises: visit toilet upon urge but sit and hold urine stream for 1 minute. Then stop and start urine flow on three occasions before emptying bladder.
- Structural toilet program: the child sits on the toilet and urinates at scheduled intervals during the day irrespective of urge. Start at 1 hour intervals, increasing to 2 - 3 hours as control is obtained.
- Medication: useful short-term drugs include the anticholinergics - oxybutynin, imipramine.

Secondary enuresis

Secondary enuresis can develop at any age and should always be fully investigated. It is often caused by urinary infection, especially in the elderly, and may be associated with some neurological disorders and chronic retention of urine associated with prostatic enlargement. Treatment is directed at the cause, which may be a psychologically traumatic event.

ENCOPRESIS

Encopresis is the involuntary passage of formed or semi-formed stools into underwear, occurring repeatedly for at least one month in children over 4 years. The cause is almost always physiological, which is often related to poor diet.

Features

- Incidence 1 - 2 per 100 children
- More prevalent in boys 3:1
- Inadequate toileting
- Poor diet
- Faecal retention (in most)
- Rectal dilatation and insensitive urge to stool
- Unawareness of passage of stools
- Enuresis common

The key feature is significant chronic faecal retention leading to rectal dilatation and insensitivity to normal defecation reflex.

Assessment

- History
- Examination
- Abdominal X-ray (serves as baseline) - may show a loaded colon

Management

A structured toileting program is the basis of management and the initial task is to empty the bowel of faeces.

The majority are cured with the following:

- ongoing interest and support (critical)
- education and counselling
- a good normal diet, adequate fluids and exercises
- structured toileting program (e.g. regular sitting on toilet for at least 10 minutes, 3 times per day after each meal)
- laxative medication, e.g. 3-day cycle: repeated 3 - 4 times after initial cleanout[11]
    - Day 1. Microlax enema

  - Day 2. Durolax rectal suppository
  - Day 3. Oral bowel stimulant (e.g. Senokot, 1 - 2 doses)

Then lubricant or softener (e.g. paraffin oil preparation) *or*

- rapid bowel emptying with Go Lytely or twice daily laxatives, then paraffin oil
- encourage keeping a star chart diary of sitting on toilet and successful results
- regular follow-up with encouragement (maintain program for at least 6 months)
- consider encopresis clinic if problematic

Once the colon and rectum are emptied and of normal size, the frequency of accidents and soiling usually decreases gradually.

CONSTIPATION

Some infants are subject to constipation - hard stool ± pain and bleeding. Exclude an anal fissure. Consider adding strained fruit juice to feeds for infants and at time of weaning. If still problematic give 2.5 mL lactulose daily (refer p. 437).

COMMON SKIN PROBLEMS[12,13]

Many of the common problems (e.g. acne, psoriasis, atopic dermatitis) are covered in more detail in Chapter 113. The following are disorders of the neonatal period and early infancy.

Toxic erythema of newborn

This is a self-limiting benign condition with onset usually 24 - 48 hours after birth (up to 14 days). Erythematous macules mainly on face and trunk. Resolves spontaneously in a few days.

Transient neonatal pustular dermatosis

This is a blistering eruption with pustules presenting at birth or in the first few hours of life. Occurs mainly on the trunk and buttocks. No treatment is required.

Naevus flammeus (‘stork bite’/salmon patch)

Dilated capillaries form on the face and eyelids (about 50% of babies) and nape of neck (almost 100%). Present from birth, they fade over 6 - 12 months but neck patches may persist into adult life. No treatment is required.

Haemangioma[6]

The classic superficial haemangioma is the so-called cavernous haemangioma or ‘strawberry naevus’. It usually develops on the head and neck, starts as a pinpoint red lesion at birth or soon after and grows up to 6 - 12 months, then with slow resolution up to 7 years. Ulceration is a complication and dressings with Duoderm or Intrasite gel can be used. Lasers can promote healing but treatment is usually unnecessary. Consider corticosteroid treatment or interferon. Refer lesions on eyelids.

Capilliary malformation (‘port wine stain’)[6]

These are also present from birth. Assessment for underlying vascular malformations are advisable if the lesion is in the area supplied by the ophthalmic or maxillary divisions of the trigeminal nerve. Consider the Sturge-Weber syndrome - associated with intellectual disability and epilepsy. The stains can be considered for pulsed dye laser therapy in the first two years or when the colour changes to bluishred, usually in early adult life. Cosmetic camouflage is useful.

Lymphatic malformation (lymphangioma)[7]

These present as a cystic tumour on the neck, face and oral cavity usually and then tend to enlarge. Previously known as cystic hygroma, they resemble clusters of vesicles with red ‘dots’. If treatment is needed excision has to be wide and deep to tie off the lymphatic channels.

Mongolian blue spot

This condition presents as bluish discolouration of the skin over the lower back and sacrum in darkskinned babies. These are of no clinical significance but may be mistaken for bruising or non-accidental injury. Usually disappears over 1 year.

Frey's syndrome

The child develops a red superficial rash or discolouration on the face (upper cheeks) upon eating or drinking. It is presumed to be related to auriculotemporal nerve damage due to forceps delivery. It usually improves with age.

Sebaceous hyperplasia

Hyperplastic sebaceous glands appear as tiny yellow - white papules on the nose, especially at the tip. Disappear in several weeks.

Milia

Blocked sebaceous glands, especially on the face, are present in 50% of neonates. The firm white papules are about 1 - 2 mm in diameter and differ from the yellowish papules of sebaceous hyperplasia. Also disappear after several weeks.

Miliaria

This is related to overheating and appears as two types:

- ‘crystallina’ - beads of sweat trapped under the epidermis, mainly on the forehead
- ‘rubia’ or ‘heat rash’ - mainly on forehead, scalp, face and trunk

It is a benign condition that disappears after a few weeks.

If problematic:

- keep skin dry and cool (e.g. fan, airconditioner)
- dress in loose-fitting cotton clothing
- reduce activity
- avoid frequent bathing and overuse of soap
- treatment: salicyclic acid 2%, menthol 1%, chlorhexidine 0.5% in alcohol
- prevention: Ego Prickly Heat Powder

Sucking blisters

These are common on the upper lip. Reassure that these will settle.

Umbilical discharge

The discharge may be pus, mucus, urine or faeces. Usually infected (fungal or bacterial) dermatitis, often with offensive discharge.

Precautions: consider umbilical fistula, carcinoma, umbilical calculus.

Management (for pus or seropus):

- swab for micro and culture
- toilet - remove all debris and clean
- keep dry and clean - daily dressings
- consider Kenacomb or similar ointment

Urinary discharge suggests a persistent communication with the bladder in the form of a patent urachus. Refer for surgical correction.

Ectopic bowel mucosa has a glistening red appearance and discharges mucus.

Bleeding umbilical cord

Small amounts of bleeding may occur as the cord is separating and requires no treatment unless it is more profuse (consider infection or a bleeding disorder).

Umbilical granuloma

This can cause a seropurulent discharge. Gently apply a caustic pencil daily for about 5 days.

Breast hyperplasia

A breast ‘bud’ is common in most term babies and may enlarge with breastfeeding (Figure 81.4). Milk may discharge from some (‘witches' milk’) but reassurance is all that is required.



Figure 81.4 Breast hyperplasia in a 15-week-old child

CHILDHOOD SKIN PROBLEMS (TREATMENT REGIMENS)

Atopic dermatitis (eczema) (Figure 81.5)



Figure 81.5 Atopic dermatitis (eczema) in a 3-year-old child with widespread distribution and severe pruritus

Refer to Chapter 113 on page 1158 - 61.

Mild atopic dermatitis

- Soap substitutes, such as aqueous cream
- Emollients - choose from:
  - aqueous cream
  - paraffin preparation (e.g. Dermeze)
  - sorbolene with 10% glycerol
  - bath oils (e.g. Alpha-Keri)
- 1% hydrocortisone (if not responding to above)

Moderate atopic dermatitis

- As for mild
- Topical corticosteroids (twice daily):
  - vital for active areas
  - moderate strength (e.g. fluorinated) to trunk and limbs
  - weaker strength (e.g. 1% hydrocortisone) to face and flexures
- Oral antihistamines at night for itch

Severe dermatitis

- As for mild and moderate eczema
- Potent topical corticosteroids to worse areas (consider occlusive dressings)
- Consider hospitalisation
- Systemic corticosteroids (rarely used)

Chronic dermatitis (on limbs)

- Zinc and tar combinations

- Corticosteroids (short course)

Pityriasis alba

- These are white patches on the face of children and adolescents.
- Can occur on the neck and upper limbs, occasionally on trunk.
- Full repigmentation occurs eventually.

Treatment

- Reassurance
- Simple emollients
- Restrict use of soap and washing
- May prescribe hydrocortisone ointment (rarely necessary)

Seborrhoeic dermatitis (Figure 81.6)



Figure 81.6 Seborrhoeic dermatitis in a 10-weekold child showing a red scaly rash affecting the scalp, forehead, face, axillae and napkin area. Both cradle cap and nappy rash are present

This is quite different to adult seborrhoeic dermatitis and appears in the first 2 to 3 months.

Refer to Chapter 111.

Medication: children

Scalp

- 1 or 2% sulphur and 1 or 2% salicylic acid or 2% salicylic acid in aqueous cream or Egozite cradle cap lotion.
- Apply overnight to scalp, shampoo off next day with a mild shampoo.
- Use 3 times a week.

Face, flexures and trunk

- 2% salicylic acid ± 2% sulphur in aqueous or sorbolene cream

- hydrocortisone 1% (for irritation on face and flexures) tds until clearance
- betamethasone 0.02 - 0.05% (if severe irritation on trunk)

Napkin area

Mix equal parts 1% hydrocortisone with nystatin or ketoconazole 2% cream

Napkin rash

Refer to Chapter 111.

Irritant dermatitis (commonest cause)

- Keep the area dry.
- Change wet or soiled napkins often - disposable ones are good.
- Wash gently and pat dry (do not rub).
- Avoid excessive bathing and soap.
- Avoid powders and plastic pants.
- Use emollients to keep skin lubricated (e.g. zinc oxide and castor oil cream).

Treatment (for specific conditions)[14]

Atopic dermatitis	1% hydrocortisone
Seborrhoeic dermatitis	1% hydrocortisone and ketoconazole ointment
Psoriasis	methylprednisolone aceponate 0.1% in a fatty ointment base daily until resolution
Candida albicans	topical nystatin at each nappy change
Widespread nappy rash	1% hydrocortisone and nystatin ointment or clotrimazole cream (qid after changes)

Impetigo

- Remove crusts with gentle washing (antibacterial soap and water).
- If mild and limited: antiseptic cleaning with chlorhexidine or povidone - iodine; then mupirocin (Bactroban) tds for 10 days (avoid around mouth).
- If extensive: oral cephalexin or flucloxacillin or erythromycin for 10 days.

Refer to Chapter 83.

Head lice

Permethrin scalp preparation or pyrethrins/piperonyl butoxide (Lyban) foam or shampoo:

- massage well into wet hair
- leave at least 20 minutes but preferably overnight
- wash off thoroughly
- comb with a fine-toothed comb with conditioner next day
- repeat after 7 - 10 days
- treat all household contacts

Refer to Chapter 83.

Scabies

Permethrin 5% cream (preferable):

- apply to whole body from jawline down
- leave overnight, then wash off

- single application *or* benzyl benzoate 25% emulsion (dilute with water if under 10 years)

Use either for all ages except children under 2 months.

Refer to Chapter 112.

Tinea capitis

Griseofulvin 10 mg/kg/day (max. 250 mg) 4 - 6 weeks course or until non-fluorescent

Take hair plucking and scale for culture.

Refer to Chapter 113.

Papular urticaria (hives)

- Prevent bites by using insecticide sprays and repellents, and treating pets.
- Lukewarm baths with Pinetarsol or similar soothing bath oil.
- Topical Liquor Picis Carb 2% in calamine lotion or 0.5% hydrocortisone - apply every 4 hours for itching.
- Antihistamine (e.g. cyproheptadine, promethazine).

Refer to Chapter 110.

Henoch-Schönlein purpura

- Characteristic rash over buttocks and back of legs.
- Prognosis is generally excellent.
- Consider analgesia (paracetamol), bed rest and crutches if symptoms a problem.
- No specific therapy; follow-up required.

Refer to Chapter 32.

Molluscum contagiosum

There are many methods but aim to provoke an immune response by applying benzyl peroxide 2.5% gel/ointment and then hypoallergenic sticking paper (Micropore) cover on a daily basis.

Vaginal skin tags

A small tag of vaginal skin commonly appears between the labia in newborn females. It is of no concern as it disappears with enlargement of the labia.

Vulvovaginitis

This is common in girls between 2 and 8. Refer to pp. 1029 - 30.

Warts

Spontaneous resolution may occur so avoid invasive treatment, including painful procedures. Do not freeze warts in children under 10 years old.[13]

If problematic and causing embarrassment, use a simple method:

Common warts:	pare every 2 - 3 days; apply a keratolytic agent with salicylic and lactic acid (e.g. Dermatech Wart Treatment) daily
Plane warts:	treat as for common warts but beware of facial lesions
Plantar warts:	pare, then apply one of the preparations given on p. 758

Refer to Chapter 67.

HAIR PROBLEMS IN CHILDREN

Refer to Chapter 118.

LEAD POISONING[15]

It is important to keep in mind that young children are susceptible to lead poisoning. They are more likely than adults to be exposed to lead because of their exploratory behaviour and they absorb more of any ingested dose. The most common source seen in general practice is home renovation, involving paint removal in houses built before the 1960s or 1970s.

The desirable level of blood lead is < 0.48 μmol/L

Children with mild to moderate lead exposure (< 2.17 μmol/L) are usually asymptomatic and tend to be symptomatic with > 2.64 μmol/L. When symptoms appear, they are usually non-specific and may include lethargy, intermittent abdominal pain, irritability, headache, abnormal behaviour and encephalopathy. Toxicity may be a cause of unexplained iron-deficiency anaemia.

An important feature is that lead poisoning may present insidiously in children with developmental delay, learning difficulties, hyperactivity or other behaviour problems. However, it is a relatively uncommon cause. It is interesting that the US Centers for Disease Control have recommended that all children between 6 months and 6 years of age should be screened with a blood lead measurement.[16]

The following children are at risk of elevated blood lead levels:

- those aged 9 - 48 months living in, or visiting, older dilapidated houses with peeling paint or such houses undergoing renovation
- those with pica
- those living near lead-contaminated areas such as lead smelters, battery-breaking yards or heavy traffic areas

High blood lead levels should be considered in the presence of unexplained iron-deficiency anaemia.

The treatment of toxicity is chelation therapy. It may involve administration of sodium calcium edatate (calcium disodium versenate) or dimercaprol in hospital. The new oral preparation, succimer, is likely to become the drug of choice for less severe degrees of poisoning.

REFERENCES

1. Oberklaid F. Crying and fussing in infancy. Aust Paediatric Rev, 1995; 5(4): 1 - 2.
2. Caswell A, Hutchins P. How to treat failure to thrive. Aust Dr Weekly, 1990; 13 April: I - VIII.
3. Martin HP. *The Abused Child.* Cambridge, MA: Ballinger, 1976.
4. Robinson MJ, Roberton DM. *Practical Paediatrics* (5th edn). Melbourne: Churchill Livingstone, 2003: 82 - 3.
5. Cormack J, Marinker M, Morrell D. *Practice: A Handbook of Primary Health Care.* London: Kluwer-Harrap Handbooks, 1980; 3.7A: 1 - 8.
6. Oates K, Currow K, Hu W. *Child Health.* Sydney: Maclennan & Petty, 2001: 198 - 205.
7. Zacharin M, Warne G, Cameron F. Endocrine conditions. In: Paxton G, Munro J. *Paediatric Handbook* (7th edn). Melbourne: Blackwell Science, 2003: 336 - 40.
8. Ibid.: 589.

9. Ibid.: 563 - 6.
10. Walsh D. *Symptom Control.* Boston: Blackwell Science, 1989: 229 - 33.
11. Jarman F. Encopresis. Aust Paediatr Rev, 1993: 3(2): 1 - 2.
12. Rogers M. Benign skin conditions of the neonatal period and early infancy. Aust Paediatr Rev, 1994: 4(1): 1 - 3.
13. Varigos G, Phillips R. Dermatologic conditions. In: Efron D. *Paediatric Handbook* (5th edn). Melbourne: Blackwell Science, 1996: 113.
14. Marley J (Chair). *Therapeutic Guidelines: Dermatology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2004: 287 - 305.
15. Campbell B. Lead poisoning. Aust Fam Physician, 1993; 22: 1139 - 45.
16. Centers for Disease Control. Preventing lead poisoning in young children. Atlanta: Department of Health & Human Services, October 1991.

CHAPTER 82. SURGICAL PROBLEMS IN CHILDREN

A surgeon should have three diverse properties in his person. That is to saie, a harte as the harte of a lyon. His eyes like the eyes of a hawke, and the handes of a woman.

JOHN HALLE (1529 - 1568)

SURGICAL PROBLEMS IN CHILDREN

An imperative task for the general practitioner is not only to diagnose surgical conditions in infancy and children as early as possible, but to be aware of the degree of urgency and the optimal times for intervention. In many instances the emphasis should be placed on a non-surgical solution using natural resolution with time and simple ‘tricks of the trade’.

HEAD DEFORMITY

The neonate's head may become distorted because of the position in utero or after the passage through the birth canal. The head shape can recover to a normal shape within about 8 weeks following birth. If the abnormal shape persists consider deformational plagiocephaly or craniostenosis.

Plagiocephaly

This is asymmetry of the skull with a normal head circumference. The shape, which can be likened to a tilted parallelogram (Figure 82.1), is the most common cause of an abnormal head shape. On the side with the flat frontal area the ear and the parietal eminence sit more posteriorly. It is either congenital or acquired and often results from the infant sleeping in one position. There is no impairment of cerebral development or intellect. If the sutures are ridged or the sleeping position causation is ruled out, a skull X-ray should be performed. Management involves initially changing the side to which the child usually faces for sleeping, then regularly changing sides and encouraging time in the prone position while awake. If not responsive a cranial remodelling helmet can be tried - best from 4 to 8 months.[1]



Figure 82.1 Plagiocephaly, congenital or acquired. The long axis is deflected from the saggital plane. In this case the right ear is more posterior

Craniosynostosis

This is premature fusion of one or more sutures of the cranial vault and base which act as lines of growth. The abnormality of head shape depends on the sutures involved. The diagnosis is confirmed by radiography. Prompt referral to a paediatric craniofacial surgeon is necessary as planning for possible complex surgery, best at 5 to 10 months, is required.

Hydrocephalus

This condition, which is due to an imbalance between the production and absorption of CSF usually caused by obstruction to circulation, requires early referral for diversion of ventricular fluid. Prognosis is generally good with early intervention and regular supervision.

Macrocephaly and microcephaly

Macrocephaly and microcephaly are defined as a head circumference greater than the 97th percentile and less than the 3rd percentile respectively. Infants whose head circumference measurements cross these percentile lines require expert assessment and investigation. It is appropriate to undertake regular head circumference measurements in the early childhood years.

EARS, NOSE, FACE AND ORAL CAVITY

Prominent/bat/shell ears

The ears are almost adult size and firmness by 5 to 6 years of age but the ear cartilage is not strong enough to cope with surgery less than 3 years. For this reason and because it is best to correct the problem when the child is in a position to support a decision to operate, the optimal time for surgical correction is after 5 to 6 years. It may be possible to correct an ear deformity by moulding the ear with tape or splinting within the first 6 months of life.[1]

Facial deformity

It is best to refer any facial deformity as soon as it is detected

External angular dermoid

This dermoid cyst, which has a readily identifiable constant position, lies in the outer aspect of the eyebrow. It is noticed in infancy as it progressively enlarges. Excision is advisable.

Cleft lip and cleft palate

Congenital clefts of the lip and palate occur in approximately 1:600 of all births. ‘It is very important that the simplest and least obvious form should be recognised in time for adequate repair. This is the sub mucus cleft, frequently not recognised in infancy because the palate appears to be intact.[2] ’ The sub mucus cleft can be diagnosed on close inspection as the uvula is bifid and there is a deep groove in the midline of the palate covered only by mucous membrane. The ideal age for repair of the cleft lip is less than 3 months of age. Secondary surgery can then be performed at various ages. The repair of the palate, which requires preliminary diagnostic ultrasound, is best performed before the child begins to speak. The optimal time is 6 to 12 months of age.

Nasal disorders

Rhinoplasty is best deferred to late adolescence. If performed early there is a higher incidence of secondary surgery.

Choanal atresia may be unilateral, leading to delayed diagnosis, or bilateral where there is no instinctive reaction to breathe through the mouth, leading to asphyxia. Since the obstruction is usually by a very thin membrane one side can be perforated with a urethral sound as an emergency procedure.

Septoplasty can be considered if the problem is symptomatic.

Tongue tie

Unless there are related breastfeeding problems formal correction is not undertaken until the diagnosis is confirmed. The child has an inability to protrude the tongue beyond the lips. A useful guideline is a strong family history of speech problems corrected by tongue tie surgery. Surgery is not usually recommended before 2 years of age.

Pre-auricular sinus

This common condition can get recurrently infected with pus discharge from a small opening immediately anterior to the ear at the level of the meatus in front of the upper crus of the helix. It also causes cosmetic problems. It is not a branchial sinus. Refer when diagnosed for surgical excision although it can be left alone if it is causing no problems.

Branchial sinus/cyst/fistula

This is a rare condition and is located inferior to the external auditory meatus or anterior to the sternomastoid muscle. The opening may discharge mucopus. A skin tag or cartilage remnant may be present. Refer when diagnosed for excision.

STRABISMUS (SQUINT)

A squint is rarely obvious in the first weeks of life, but tends to show up when the baby learns to use the eyes, from about 2 weeks to 3 or 4 months of age. However, it may appear late, even as an adult. Vision, which is present at birth, continues to develop until 7 - 8 years of age.

Main types of squint (Figure 82.2)


Convergent squint (affected eye on right of page)
Divergent squint (affected eye on right of page)
Pseudo squint (due to shape of eyelids)

Figure 82.2 Types of squint

- *Constant or true squint* is one that is permanent - always present.
- *Latent squint* is one that only appears under stressful conditions such as fatigue.
- *Transient squint* is one that is noticeable for short periods and then the eye appears normal.

- *Alternating squint* is one that changes between the eyes so the child can use either eye to fix vision.
- *Pseudo squint* is not a true squint but only appears to be one because of the shape of the eyelids, i.e. broad epicanthic folds.

A useful way to differentiate a true squint from a pseudosquint is to observe the position of the light in the eyes (corneal reflections) when a torch is shone into them from about 40 cm away. This light reflex will be in exactly the same position in both eyes in the pseudosquint but in different spots with the true squint.

- If one eye is 'lazy' (that is, not being used), it is standard practice to wear a patch (maybe on glasses) over the good eye for long periods in order to use the inactive eye and have both eyes eventually capable of vision.
- The two serious squints are the constant and alternating ones, which require early referral. Transient squint and latent squint usually are not a problem.
- Always refer children with strabismus (squint) when first seen to exclude ocular pathology such as retinoblastoma, congenital cataract and glaucoma, which would require emergency surgery.
- Children with strabismus (even if the ocular examination is normal) need specialist management because the deviating eye will become amblyopic (a lazy eye with reduced vision, i.e. 'blind', if not functioning by 7 years of age). The younger the child, the easier it is to treat amblyopia; it may be irreversible if first detected later than school age. Surgical correction of a true squint is preferred at 1 - 2 years of age.

BLOCKED NASOLACRIMAL DUCT

Refer Chapter 81, page 874.

NECK LUMPS

Sternomastoid tumour/fibrosis

Features in infants:

- Hard painless lump (2 - 3 cm long) within sternomastoid muscle
- Tight and shortened sternomastoid muscle
- Usually not observed at birth
- Appears at 20 - 30 days of age
- Associated torticollis - head turned away from tumour
- Restricted head rotation to side of tumour

Most tumours resolve spontaneously within 1 year. The mother should be reassured and the child referred to a physiotherapist early. Mother should frequently gently massage the lump and stretch the neck towards the tumour. If surgery for a persistent fibrotic shortened muscle is required it is best before 12 months.

Older children can present with torticollis and a tight, short fibrous sternomastoid muscle. It is associated with rotation of the head to the affected side, hemihypoplasia of the face and a wasted ipsilateral trapezius muscle. It requires surgical repair.

Thyroglossal cyst

This is the most common childhood midline neck swelling. It moves with swallowing and tongue protrusion. It is prone to infection, including abscess formation. The cyst and its tract are best excised before it becomes infected.

Lymphatic malformation/lymphangioma/cystic hygroma

These usually present as soft cystic tumours of the neck, face or oral cavity. They resemble clusters of vesicles and are often poorly localised. Some have visible red dots due to haemangiomatous inclusions. If located in the floor of the mouth or peripharyngeal area they endanger the airway and can precipitate an emergency requiring surgery. Surgery is advisable in the early years.

Cervical lymphadenopathy

Refer to p. 650.

SKIN TUMOURS

Haemangioma (strawberry naevus)

These start soon after birth as a red pinpoint lesion and grow rapidly for the first 6 months, then involute and become pale. Full resolution may take several years. Reassure parents and demonstrate how to stop any bleeding by applying pressure. Possible treatment options include oral or intralesional steroids, vascular laser, interferon and surgery. Surgical intervention is usually not necessary. Exceptions are locations in critical areas such as periorbital, nose, lips and face. Refer lesions on the eyelid early since visual obstruction can lead to amblyopia. Stridor accompanying a haemangioma on the face is suggestive of laryngeal haemorrhage so refer urgently.

Capillary malformation (port wine stain)

These are present from birth and surgical intervention is inadvisable. They may be treated by pulsed dye laser, which is best initiated as early as possible as the response is best in the first 2 years.[3] (Refer to p. 883.)

Venous malformations

These are aggregations of abnormal subcutaneous veins that may infiltrate deeper tissues. In the past these lesions were treated surgically, but now the emphasis is on specialised sclerosant agents injected under fluoroscopic guidance and specialised laser techniques. Referral to Vascular Malformation Clinics in larger centres is worthy of enquiry.

Lymphatic malformations

These appear sometimes as skin lesions because of the red discolouration on the surface of the tumour. Management is as described above.

Congenital naevi

These have to be treated on an individual basis. If giant naevi they can be dermabraded at ideally less that 6 weeks.

Benign juvenile melanoma

These pigmented lesions, which typically appear on the face, are usually surgically excised because of their rapid growth and family concerns.

CHEST AND BREAST DISORDERS

Breast asymmetry

If necessary surgery should be performed in late adolescence after breast development is complete. It can take the form of a unilateral implant, different-sized bilateral implants or unilateral breast reduction.

Macromastia

Reduction surgery should also be delayed until breast growth is complete, at late adolescence.

Gynaecomastia

If this develops in the pubertal stage gynaecomastia may resolve spontaneously within 1 or 2 years. If necessary simple mastectomy can be performed.

Subareolar hyperplasia in boys

This presents as a firm discoid subareolar lesion similar to premature breast hyperplasia of girls. It typically occurs at about 12 - 14 years. There is no indication for surgical treatment. Give reassurance that the problem will dissipate.

Chest wall skeletal deformity

Surgical correction is best performed in adolescence.

Poland's syndrome

This syndrome is an absent sternal head of pectoralis major with associated chest wall deformity plus a hypoplastic or absent breast and nipple - areolar complex. Surgery correction can be undertaken from 10 to 20 years.

CONGENITAL HEART DISORDERS

GPs have an important role in the diagnosis of congenital heart disorders as many of the affected infants develop cyanosis, murmurs or congestive heart failure. Early diagnosis and intervention helps prevent serious problems such as bacterial endocarditis and paradoxical emboli.

Ventricular septal defect (VSD)

VSD is the commonest congenital heart lesion (1:500 births).

The defect connects the two ventricles with a L→R shunt (Figure 82.3).



Figure 82.3 Ventricular septal defect: defect in muscle wall

Symptoms and signs depend on size of hole. All have a palpable thrill at the left sternal edge and a pansystolic down right sternal edge.

Small VSD (‘maladie de Roger’): harsh murmur, usually asymptomatic and close spontaneously.

Larger VSD: symptoms appear in infancy.

- Breathlessness on feeding and crying (i.e. early CHF)
- Recurrent chest infections
- Failure to thrive
- Heart failure from about 3 months with large defects

Refer early, especially if failure - early surgery performed by 6 months but can be performed at any age from the newborn period. A patch can close the defect through open-heart surgery. Some may be closed by sealing with an occlusive device through a percutaneous cardiac catheter. A cardiologist will make the appropriate decision. As a general rule about 50% of all VSDs will close spontaneously.

Atrial septal defect (ASD)

In ASD the defect connects the two atria with two distinct types - ostium secundum with holes higher in the septum (most common) and ostium primum with holes lower in the septum (more serious) (Figure 82.4). Signs are a mid-systolic murmur in the pulmonary area, a split 2nd sound and a loud P2. An echocardiogram is diagnostic.



Figure 82.4 Atrial septal defect: primum defect

Symptoms are uncommon in infancy and childhood with o. secundum but heart failure with pulmonary hypertension develops early with o. primum.

Refer these patients early. Prophylactic antibiotics are needed for patients with o. primum. Follow other cases with regular echocardiograms and growth/development monitoring. Closure is advisable where there is evidence of a troublesome shunt. Options are repair by direct surgical suture or an insertion of a patch or a device closure using a self-expanding ‘double umbrella device’ manipulated into the defect via cardiac catheterisation.

All patients require prophylactic antibiotics before procedures.

Patent ductus arteriosus

The ductus fails to close after birth. A loud continuous machinery murmur is heard. Symptoms related to shunt size.

Child presents with a murmur with possible respiratory infections, failure to thrive and heart failure. Refer for possible surgical closure by ligation. Alternatives include device closure with placement of an occlusive device or by embolisation coils.

Coarctation of the aorta

Usually presents in infancy with heart failure. Refer for early surgery to remove the narrowed portion of the aorta.

HERNIAS AND GENITAL DISORDERS

Inguinal hernias

These usually present in the first 3 to 4 months with an incidence of 1 in 50 males and 1 in 500 females (Figure 82.5).



Figure 82.5 Sites of common hernias

They should be referred urgently as early surgery is advisable to avoid the high risk of strangulation. It is useful to follow the ‘6 - 2’ rule (see p. 1083).

Hydroceles

Scrotal hydroceles are painless cystic swellings around the testis. The opening of the processus vaginalis is narrow and often closes spontaneously. Two types can be identified - slack, often bilateral, which disappear within 12 months, and tense, often unilateral, which often persist after the first year. Ninety per cent resolve by 18 months of age and for those that persist referral is recommended with a view to surgical intervention if present longer than 2 years.

Undescended testes

Testes can still descend up to 3 months after birth. Refer by 6 months with a view to correction between 9 and 12 months but definitely before 2 years (see p. 1088).

Hypospadias

Refer to disorders of the penis (see p. 1094). Look for other abnormalities. Refer as soon as possible if the child is not producing a good urinary stream. Non-urgent cases should be evaluated by 6 months with a view to surgery at around 12 months; these patients should not be circumcised.

The foreskin and circumcision

For more detail refer to penile problems (pp. 1093). If not circumcised in the neonatal period it is best performed under general anaesthetic after 6 months of age following consultation, counselling and with the consent of both parents.

Phimosis

Real phimosis is uncommon and almost all cases of tight foreskin with narrowing of the preputial orifice resolve naturally. Treatment with corticosteroid creams should be considered (see p. 1091). Probably the only indication for circumcision is persistent difficulty passing urine.

Paraphimosis

Management of this painful condition is outlined on page 1091.

Umbilical hernia

Surgery is not usually required for umbilical herniae as most close naturally by 4 years of age. Refer for possible repair if still present at 4 years of age. A good guideline is that if the hernial orifice is greater than 1 cm at 12 months then surgical intervention is a possibility. It is usual to operate at 4 to 5 years.

Para-umbilical hernia

This is due to a defect in the linea alba adjacent to the umbilicus proper. Most lie just above the umbilicus. The defect is feels like an elliptical slit with firm edges. Spontaneous closure rarely occurs and they are more likely to incarcerate. Refer for operation at any age preferably after 6 - 12 months.

Epigastric hernia

An epigastric hernia (not to be confused with divarification of the rectus muscles) lies between the umbilicus and the xiphisternum. It is unlikely to close naturally, is likely to incarcerate and causes pain by strangulation of herniated fat. Indications for repair are pain (reproducible on palpation of the hernia) and cosmetic.

Anal fissure

Anal fissures are often seen in infants and toddlers with uncomfortable defecation and minimal bight bleeding. The anal mucosa is split in the midline either anterior or posterior. It is caused by the passage of hard stool. The fissure usually heals within a few days.

Fused labia (labial agglutination)

Labial fusion is caused by adhesions considered to be acquired from perineal inflammation (see p. 1030). They are certainly not present at birth. Most authorities recommend no treatment if the child can void readily and allow natural healing to occur.

NEONATAL LEG AND FOOT DEFORMITIES

Developmental dysplasia of hip

- Detected by clinical examination (Ortolani and Barlow tests) and ultrasound examination (see p.700).
- Infants are usually treated successfully by abduction splinting e.g. Pavlik harness.
- Open reduction may be required, especially in older babies and toddlers.

Bow legs (genu varum)

- Most are physiological (which are symmetrical) and improve with age.
- Consider rickets in children at risk.
- Toddlers are usually bow-legged until 3 years of age.
- Invariably resolve spontaneously by 3 years except in severe cases.
- Monitor intercondylar separation (ICS): distance between medial femoral condyles.
- Refer when ICS > 6 cm at 4 years, not improving or asymmetric (Figure 82.6).



Figure 82.6 Postural variance of lower limbs. Reprinted from D. Efron. *Paediatric Handbook*. Melbourne: Blackwell Science, 1996, with permission.

Knock knees (genu valgum)

- Most are physiological and children are usually knock-kneed from 2 - 8 years (maximal 3 - 4 years).
- Running is awkward, but improves with time.
- Reassure parents about spontaneous improvement.
- Monitor intermalleolar separation (IMS): distance between medial malleoli.
- Refer if IMS > 8 cm (Figure 82.6).

In-toeing (pigeon toes)

In-toeing does not cause pain or affect mobility.

Causes of in-toeing (Figure 82.7) are metatarsus varus, internal tibial torsion and medial femoral torsion. Children with femoral torsion tend to sit in a characteristic ‘W’ sitting position (Figure 82.8). These features are compared in Table 82.1.[4]



Figure 82.7 Causes of in-toeing. Reprinted from D. Efron. *Paediatric Handbook*. Melbourne: Blackwell Science, 1996, with permission



Figure 82.8 The classic ‘W’ position of femoral torsion (inset hips).

Table 82.1 In-toeing in childhood[1]

	Metatarsus varus	Internal tibial torsion	Medial femoral torsion
Synonyms	Metatarsus adductus		Inset hips
Age at presentation	Birth	Toddler	Child
Site of problem	Foot	Tibia	Femur
Examination	Sole of foot bean-shaped	Thigh - foot angle is inwards	Arc of hip rotation favours internal rotation
Management	Observe or cast	Observe and measure	Observe, rarely surgery
Resolution (usually by)	3 years	3 - 4 years	8 - 9 years
When to refer if not resolved	3 months after presentation	6 months after presentation	8 years after presentation

Reprinted from D. Efron. *Paediatric Handbook*. Melbourne: Blackwell Science, 1996 with permission

Out-toeing

Infants

- Have restricted internal rotation of hip due to an external rotation contracture
- Exhibit a ‘Charlie Chaplin’ posture between 3 and 12 months - up to 2 years
- Child weight-bears and walks normally
- No treatment required as spontaneous resolution occurs

Surgery may be necessary in older children.

Club foot (congenital talipes equinovarus)

Most abnormal-looking feet in infants are not a true club foot deformity; the majority have postural problems referred to as 'postural talipes' such as talipes calcaneovalgus, metatarsus varus and postural talipes equinovarus. Such conditions are usually quite mobile and mild, and all resolve spontaneously without treatment. True club foot deformity is usually stiff and severe, and requires orthopaedic correction.[4]

Inset hips (medial femoral torsion)

In children with inset hips the femur tends to rotate inwards especially when the child is about 5 - 6 years old and is normal up to 12 years. The children tend to sit in a 'W' position (Figure 82.8). Fortunately, most children outgrow this condition before the age of 12.

Flat feet (pes plano valgus)

The majority are physiological and are usually hereditary. All newborns have flat feet but 80% develop a medial arch by their sixth birthday and most by 11 years.[4] The presence of the arch can be demonstrated to parents by the tiptoe test. The arch can be seen better when the feet are hanging in the air and even better still when the child is standing on tiptoes (Figure 82.9). Flat feet are present in about 10% of teenagers. No treatment is required unless painful and stiff. Good roomy footwear is important. Studies in California have shown no benefit from wearing orthoses or other form of arch supports. Arches develop naturally.



Figure 82.9 The tiptoe test for flat feet

Curly toes

Usually the third toe curls inward under the second toe so that the second toe lies above the level of the first and third toes. The toes can usually be straightened if necessary so ignore the problem until 2 years. Refer if necessary to have a severe deformity corrected by flexor tenotomy.

Table 82.2 Optimal times for surgery/intervention in children's surgical disorders

Disorder	Surgery/intervention
Squint (fixed or alternating)	12 - 24 months: absolutely before 7 years
Tongue tie	2 - 6 years

Ear deformity	After 6 years
Cleft lip	Less than 3 months
Cleft palate	6 - 12 months
Inguinoscrotal lumps	
• Undescended testes	Best assessed before 6 months
	Surgery best at 6 - 12 months
• Inguinal hernia	ASAP, especially infants and irreducible hernias
	Reducible hernias: the '6 - 2' rule birth - 6 weeks: surgery within 2 days 6 weeks-6 months: surgery within 2 weeks over 6 months: surgery within 2 months
• Femoral hernia	ASAP
• Torsion of testicle	Surgery within 4 hours (absolutely 6 hours)
• Hydrocele	Leave to 12 months, then review (often resolve)
• Varicocele	Leave and review
Other hernias	
• Umbilical hernia	Leave to age 4
	Surgery after 4 if persistent (tend to strangulate)
	Never tape down
• Para-umbilical hernia	Any age - best after 6 months
• Epigastric hernia	Any age - best after 6 months
Leg and foot development problems	
• Developmental dysplasia hip	Most treated successfully by abductor bracing with Pavlic harness
• Bowed legs (genu varum)	Normal up to 3 years
	Usually improve with age: refer if ICS > 6 cm
• Knock knees	Normal 3 - 8 years, then refer if IMS > 8 cm
• Flat feet	No treatment unless stiff and painful
• Internal tibial torsion	Refer 6 months after presentation if unresolved
• Medial tibial torsion	Leave to 8 years, then refer if unresolved
• Metatarsus varus	Refer 3 months after presentation if unresolved

APPENDICITIS IN CHILDREN

Refer to p. 320.

Surgical causes of vomiting in children

Refer to p. 640.

NEONATAL SURGICAL EMERGENCIES

It is worthwhile knowing those non-traumatic conditions that demand immediate attention. Danger signs include excessive drooling of frothy secretions, bile-stained vomiting and delayed passage of meconium.

Neonatal emergencies[5, 6]

- Oesophageal atresia: rattling respiratory distress + excessive drooling and secretions + choking with feeding (passage of 10F catheter stops about 10 cm)
  Action: nil orally, oropharyngeal suction, IV fluids
- Diaphragmatic hernia: severe respiratory distress + barrel-shaped chest + scaphoid abdomen (X-rays of chest/abdomen show loops of bowel in chest)
  Action: Give O_2, nasogastric tube (avoid bag and mask)
- Bilious (green vomiting) = bowel obstruction or malrotation (abdominal X-ray)
  Action: nasogastric drainage and refer
- Neonatal intestinal obstruction: bilious vomiting+ distension + delayed stools
- Imperforate anus and rectum
- Bile duct atresia: neonatal jaundice (conjugated bilirubin) = white stools
- Congenital lobar emphysema: respiratory distress + cyanosis + signs of emphysema
- Congenital cystic disease of the lungs: respiratory distress soon after birth
- Congenital heart disease (severe forms)
- Exomphalos (intestinal contents in a sac)
- Gastroschisis (exposed bowel contents through anterior wall defect
- Pierre-Robin syndrome: micrognathia + cleft palate + respiratory obstruction from tongue
- Tension pneumothorax
- Myelomeningocele and meningocele

Note the importance of plain X-rays as urgent first-line investigations.

Other important childhood emergencies

- Pyloric stenosis: projectile vomiting weeks 3 - 6, epigastric ‘ptumour’
- Torsion of testis: severe groin/low abdominal pain + vomiting
- Intussusception: pallor + abdominal pain (severe and intermittent) + inactivity
- Acute appendicitis: abdominal pain + a/n/v + guarding
- Peritonitis
- Intestinal obstruction: colicky pain + vomiting + distension
- Cerebral abscess
- Meckel's diverticulum - diverticulitis or haemorrhage
- Various neoplasias

REFERENCES

1. MacGill KA. Paediatric plastic surgery. Aust Doctor, 2005; 9 September: 32.
2. Hutson JM, Woodward A, Beasley SW. *Jones Clinical Paediatric Surgery*. Oxford: Blackwell Publishing, 2003; 18 - 72.
3. Marley J (Chair). *Therapeutic Guidelines: Dermatology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2004: 309.
4. Kerr G, Barnett P. Orthopaedic conditions. In: Smart J (ed). *Paediatric Handbook* (6th edn). Melbourne: Blackwell Science, 2000; 454 - 9.
5. Jones PG. *Clinical Paediatric Surgery*. Sydney: Ure Smith, 1970: 16-19.
6. Smart J, Munro J. *Paediatric Handbook* (7th edn). Melbourne: Blackwell Science 2003: 602 - 4.

CHAPTER 83. COMMON CHILDHOOD INFECTIOUS DISEASES (INCLUDING SKIN ERUPTIONS)

> The physical signs of measles are nearly the SAME as those of smallpox, but nausea and inflammation is more severe. The rash of measles usually appears at once, but the rash of smallpox spot after spot…
>
> AVICENNA (980 - 1037)

COMMON CHILDHOOD INFECTIOUS DISEASES (INCLUDING SKIN ERUPTIONS)

Children are subject to a variety of infectious diseases, mainly causing acute skin eruptions. Fortunately, many of these diseases, such as scarlet fever, measles and rubella, are being seen less frequently by the family doctor.

REYE'S SYNDROME AND ASPIRIN

The concern about the ingestion of aspirin for febrile illness in children is the suspected causal relationship between it and Reye's syndrome, particularly in children with varicella and influenza infections. However, there is some controversy about the connection. Orlowski and colleagues at the Children's Hospital in Sydney found no association between aspirin use and Reye's syndrome from 1973 - 1985.[1] It is possible that the connection is coincidental or at least confounded with other factors.

Despite these doubts, aspirin should not be recommended for the treatment of fever in young children in view of our knowledge of the beneficial effects of fever on the immune response and the availability of a safe alternative antipyretic such as paracetamol.

Aspirin should probably not be used under 16 years (previous recommendation under 12) except for Kawasaki disease and juvenile arthritis.

Reye's syndrome

Clinical features:

- a rare complication of influenza, chickenpox and other viral diseases (e.g. Coxsackie virus)
- rapid development of:
  - encephalopathy (seizures and coma)
  - hepatic failure (seizures and coma)
  - hypoglycaemia (seizures and coma)
- 30% fatality rate and significant morbidity
- treatment is supportive and directed at cerebral oedema

VARICELLA (CHICKENPOX)

Varicella, a common and highly infectious disease, affects people mainly during childhood, especially between 2 and 8 years, but no age is exempt. The characteristic crops of small vesicles have a central distribution (face, scalp and trunk). It is caused by the varicella zoster virus, one of the human herpes viruses, which remains latent after infection. Clinical reactivation later in life results in herpes zoster.

Epidemiology

Varicella has a worldwide distribution, causing endemic (occasionally epidemic) disease, with little clear evidence of seasonal incidence in temperate climates. About 75% of people in urban communities have had the infection by 15 years of age and at least 90% by young adulthood.

It is one of the most easily transmitted viruses, probably by airborne spread, usually via a person with chickenpox (occasionally with herpes zoster). Varicella is contagious only while the patient has symptoms and vesicles remain; drying of the vesicles indicates that infectivity has stopped. The scabs are not infectious.

The incubation period is 10 - 21 days (usually 15 - 16). Laboratory diagnosis is by serology[2] or immunofluorescence of vesicular fluid.

Clinical features

The clinical features of varicella are shown in Table 83.1 and the complications in Table 83.2. Children are not normally very sick but tend to be lethargic and have a mild fever. Adults have an influenza-like illness. The typical distribution is shown in Figure 83.2.

Table 83.1 Clinical features of varicella

Onset • Children: no prodrome • Adults: prodrome (myalgia, fever, headaches) for 2 - 3 days
Rash • Centripetal distribution, including oral mucosa (Figure 83.1) • Scalp lesions can become infected • 'Cropping' phenomenon: vesicles, papules, crusting lesions present together • Pruritic **Degrees of severity** • Number of vesicles can vary from fewer than ten to thousands • Mild cases can be missed • More severe in adults, especially the immunocompromised • Viral pneumonia rare in children, uncommon in adults • Death rare except in the immunocompromised and neonates with congenital varicella



Figure 83.1 Varicella (chickenpox) in a 12-year-old girl showing a maculopapular vesicular rash with a centripetal distribution

Table 83.2 Complications of varicella

Common • Bacterial infection of cutaneous lesions (usually staphylococcal or streptococcal); can take form of cellulitis or bullous impetigo • Can leave pitted scars
Uncommon • Viral pneumonia • Thrombocytopenia • Acute cerebellitis (ataxia, normal mental state)
Rare • Meningoencephalitis • Purpura fulminans



Figure 83.2 Chickenpox: typical distribution

Treatment

Treatment is symptomatic and usually no specific therapy is required. Many people worry about scarring but the lesions invariably heal, leaving normal skin, unless they become infected.

Advice to parents

- The patient should rest until feeling well.
- Give paracetamol for the fever (avoid aspirin).
- Daub calamine or a similar soothing lotion to relieve itching, although the itch is usually not severe.
- Avoid scratching; clean and cut the fingernails of children short. Provide cotton mittens if necessary.
- Keep the diet simple. Drink ample fluids, including orange juice and lemonade.
- Daily bathing is advisable, with the addition of mild antiseptic or sodium bicarbonate if pruritic (half a cup to the bath water). Pat dry with a clean, soft towel; do not rub.

Medication

Antihistamines can be prescribed for itching. Aciclovir or similar agents can be life-saving in the immunocompromised host. Antibiotics (e.g. flucloxacillin/dicloxacillin) are reserved for bacterial skin infection.

Use of antiviral agents for varicella

The antiviral agents, aciclovir and others, have an important place in the management of severe chickenpox. Although generally used in adolescents and adults with a severe eruption or the likelihood of a severe eruption, there is no set age for the use of an antiviral agent. It should be commenced during the first 3 days of the eruption (preferably day 1) and can be introduced in a contact (often the second case in a family) experiencing a severe prodromal syndrome.

In general, it is not used in the very young, or in those with a mild/short prodrome, and in those who are not ill, do not have many spots and are not compromised.

Exclusion from school

Exclusion is recommended until full recovery, usually for 7 days. A few remaining scabs are not an indication to continue exclusion. Except for immunocompromised children, contacts should not be excluded from school.

Exclusion and incubation times are given in Table 83.3.

Table 83.3 Basic childhood infectious diseases: incubation periods, minimum exclusion periods from school, preschool and child care centres (times in days)[*]

	Incubation period (days)	Patient exclusion (least time from onset rash or symptoms) (days)	Contact exclusion (days)
Measles	10 - 14	5	14 in non-immunised
Mononucleosis	?30 - 50	Nil	Nil
Mumps	14 - 21	9	Nil
Pertussis	7 - 14	5 (after starting antibiotics)	14 in non-immunised
Parvovirus (erythema infectiosum)	4 - 21	Nil	Nil
Rosesla infantum	7-17	Nil	Nil
Rubella	14 - 21	5	Nil
Scarlet fever	1-7	24 hours (after starting antibiotics)	Nil
Varicella and zoster	10 - 21	7	Only those immune deficient
Hepatitis			
A	15 - 45	7 or recovery	Nil
B	40 - 180	Nil	Nil
C	14 - 180	Nil	Nil
Infective diarrhoea	varies	24 hours after cessation diarrhoea	Nil

[*] Based on NHMRC recommendations.

Prevention

Prevention in contacts who are immuno-compromised, or premature infants in contact with varicella, is possible with zoster immune globulin. An attenuated live virus vaccine (0.5 mL sci)is available and suitable for healthy children from the age of 12 months up to and including 13 years. It is given routinely at 18 months with the immunisation schedule.

MEASLES

Measles (rubeola) is a highly contagious disease caused by an RNA paramyxovirus. It presents as an acute febrile exanthematous illness with characteristic lesions on the buccal mucosa called Koplik's spots (tiny white spots like grains of salt).

The disease is endemic throughout the world and complications are usually respiratory in nature. If an acute exanthematous illness is not accompanied by a dry cough and red eyes, it is unlikely to be measles. Laboratory diagnosis is by serology, nasopharyngeal aspirate immuno-fluorescence and culture.[2]

Epidemiology

Measles is transmitted by patient-to-patient contact through oropharyngeal and nasopharyngeal droplets expelled during coughing and sneezing.

The incubation period is 10 - 14 days and the patient is infectious for about 5 days, but especially just before the appearance of the rash. Morbidity and mortality are high in countries with substandard living conditions and poor nutrition.

Immunity appears to be lifelong after infection. Measles, like smallpox, could be eradicated with public health measures.

Clinical features

The clinical presentation can be considered in three stages.

1. *Prodromal stage*. This usually lasts 3 - 4 days. It is marked by fever, malaise, anorexia, diarrhoea, and 'the three Cs': cough, coryza and conjunctivitis (Figure 83.4). Sometimes a non-specific rash appears a day before the Koplik's spots (opposite the molars).



Figure 83.4 Measles: typical symptoms. Note the 3C's: cough, coryza, conjunctivitis

2. *Exanthema (rash) stage*. Identified by a typically blotchy, bright red maculopapular eruption; this stage lasts 4 - 5 days. The rash begins behind the ears and on the first day spreads to the face,(Figure 83.3) the next day to the trunk and later to the limbs. It may become confluent and blanches under pressure. The patient's fever usually subsides within 5 days of the onset of the rash.



Figure 83.3 Measles showing the typical blotchy, bright red maculopapular rash on the face in a miserable child with 'the three Cs' (cough, coryza and conjunctivitis). *Photo courtesy Hugh Newton-John*

3. *Convalescent stage*. The rash fades, leaving a temporary brownish 'staining'. The patient's cough may persist for days, but usually good health and appetite return quickly.

Complications

Respiratory

The patient could develop secondary bacterial otitis media or sinusitis. If pneumonia develops it is more likely to be bacterial superinfection than viral. Laryngotracheobronchitis (croup) is a common complication of measles.

Central nervous system

Encephalitis has an incidence of 1 in 1500 and although the mortality rate is low there is significant CNS morbidity. Febrile convulsions are another common complication.

Late complications

Two rare complications are bronchiectasis and subacute sclerosing panencephalitis.

Treatment

There is no specific treatment although some symptoms can be relieved (e.g. a linctus for the cough, paracetamol for fever). The patient should rest quietly, avoid bright lights and stay in bed until the fever subsides.

The management of complications is determined by their nature and severity. Children should be kept away from school until they have recovered or for at least 5 days from the onset of the rash.

Prevention

Vaccination programs have been most successful. Young adults are probably at highest risk of measles these days and consideration should be given to immunising those 18 - 30 years of age. Live attenuated measles virus vaccinations combined with mumps and rubella (MMR) are recommended at the age of 12 months and then 4 - 5 years. Consider normal immunoglobulin for infants less than 12 months and the immunocompromised when MMR is contraindicated, given as soon as possible after exposure.

RUBELLA

Rubella (German measles) is a viral exanthema caused by a togavirus. Because of immunisation programs it is seen less frequently now in family practice. It is a minor illness in children and adults, but devastating when transmitted in utero. Congenital rubella is still the most important cause of blindness and deafness in the neonate. It is completely preventable.

Epidemiology

Rubella has been reported from virtually every country and is endemic in heavily populated communities. Epidemics occur every 6 - 9 years in non-immunised populations, the disease being spread by droplets from the nose and throat. It is not as communicable as varicella and measles. Intrauterine infection occurs via the placenta.

Approximately one-third of infections are asymptomatic (subclinical). Infection usually confers lifelong immunity. Infection is proved either by virus culture or by specific serology. Incubation period is 14 - 21 days.

Clinical features

The clinical features of rubella are presented in Table 83.4 and Figure 83.5 and the comp-lications in Table 83.5.

Table 83.4 Clinical features of rubella

• There is no prodrome. • A generalised, maculopapular rash, sometimes pruritic, may be the only evidence of infection. • Other symptoms are usually mild and short-lived. • There is often a reddened pharynx but sore throats are unusual. An exudate may be seen as well as palatal exanthem. • Fever is usually absent or low-grade. • Other features: headache, myalgia, conjunctivitis and polyarthritis (small joints). • Lymphadenopathy may be noted; usually postauricular, suboccipital and postcervical. • The patient is infectious for up to 10 days from onset of rash (this aspect is often not appreciated as the patient is asymptomatic by that time).
The rash • A discrete pale pink maculopapular rash (not confluent as in measles). • Starts on the face and neck - spreads to the trunk and extremities. • Variable severity: may be absent in subclinical infection. • Exaggerated on skin exposed to sun. • Brief duration - usually fades on the third day. • No staining or desquamation.


postauricular lymphadenopathy
rash first appears on neck and face
scattered rash on body

Figure 83.5 Rubella: typical symptoms

Table 83.5 Complications of rubella

- Encephalitis (1 in 5000)
- Polyarthritis, especially in adult women (this complication abates spontaneously)
- Thrombocytopenia with bleeding (1 in 3000)
- Congenital rubella

Treatment

Treatment is symptomatic, especially as rubella is a mild disease. Patients should rest quietly until they feel well and take paracetamol for fever and aching joints. Prevention is by vaccine, recommended at 12 months and 4 - 5 years.

School exclusion

The child is usually excluded until fully recovered or for at least 5 days from the onset of the rash.

Congenital rubella

Infection of the mother in the first trimester can lead to abortion or stillbirth, or to foetal malformation, including congenital heart disease, deafness and blindness (cataract or glaucoma). It can also produce lesions (such as microcephaly), intellectual disability, retarded growth, thrombocytopenic purpura (with a 30% mortality rate), jaundice/hepatospleno-megaly and bone abnormalities.

Rubella in pregnancy

Ideally all women of child-bearing age should know their rubella immune status by having serology performed. A history of immunisation is not good enough evidence of immunity. However, in Victoria, Australia, almost 95% of women aged 15 - 40 are immune.[3]

If the immune status is not known, then serology testing should be ordered at the first antenatal visit. Rubella vaccine, while not shown to be embryopathic, should not be given during pregnancy. If maternal rubella antibodies are in adequate titre, there is no risk to the foetus from rubella infection.

VIRAL EXANTHEMA (FOURTH SYNDROME)

This mild childhood infection may be caused by a number of viruses, especially the enteroviruses, and produces a rubella-like rash which may be misdiagnosed as rubella. The rash, which is usually non-pruritic and mainly confined to the trunk, does not desquamate and often fades within 48 hours. (Figure 83.6)The child may appear quite well or can have mild constitutional symptoms, including diarrhoea.



Figure 83.6 Viral exanthema (fourth syndrome). This mild rubella-like maculopapular rash, which may be caused by a number of viruses, is usually non-pruritic

ERYTHEMA INFECTIOSUM (FIFTH SYNDROME)

Erythema infectiosum, also known as ‘slapped cheek’ syndrome or fifth disease, is a childhood exanthem caused by parvovirus B19. It occurs typically in young school-aged children. Incubation period is 4 - 21 days. The bright macular rash erupts on the face first (Figure 83.7) then, after a day or so, a maculopapular rash appears on the limbs.[4] The rash lasts for only a few days but may recur for several weeks. By the time the rash appears the individual usually is no longer infective.



Figure 83.7 Erythema infectiosum (fifth syndrome) showing the typical slapped cheek appearance. *Photo courtesy Frank Mansfield*

Clinical features

- Mild fever (30%) and malaise
- Possible lymphadenopathy (especially cervical)

The rash (Figure 83.8):


possible lymphadenopathy
erythematous 'slapped' cheek appearance in well child
erythematous rash, mainly on forearms and thighs

Figure 83.8 Erythema infectiosum: typical distribution of rash

- bright red flushed cheeks with circumoral pallor for 2 - 3 days
- maculopapular rash on limbs (especially) and trunk (sparse)
- reticular appearance on fading
- may be pruritic

Typically, the cheeks become reddened again for the next few weeks on exposure to sunlight or wind or after a hot bath.[2] Erythema in-fectiosum is a mild illness but, if the parvovirus infection occurs during pregnancy, foetal complications including death in utero can occur.[3] It is advisable to avoid close contact with pregnant women. Adults can be infected and the side-effects, especially arthritis, can be quite severe. Diagnosis is by serology.

Treatment

Treatment is symptomatic

- Ample fluid intake
- Paracetamol for fever
- If itchy daub a soothing anti- itch lotion such as piretarsol or calamine lotion.
- Wear a broad-brimmed hat when outside

Adults may need stronger analgesics or NSAIDs for joint pain.

ROSEOLA INFANTUM (EXANTHEMA SUBITUM OR SIXTH SYNDROME)

Roseola infantum is a viral infection (human herpes virus 6) of infancy, affecting children at the age of 6 - 18 months; it is rare after this time. Constitutional symptoms are generally mild.

Clinical features

- High fever (up to 40°C)
- Runny nose
- Temperature falls after 3 days (or so) then
- Red macular or maculopapular rash appears

The rash:

- largely confined to trunk (Figure 83.9)



Figure 83.9 Roseola infantum: typical distribution of rash

- usually spares face and limbs
- appears as fever subsides
- disappears within 2 days
- no desquamation
- mild cervical lymphadenopathy

The infection runs a benign course, although a febrile convulsion can occur. Diagnosis is by serology and treatment is symptomatic. Encourage high fluid intake.

HAND, FOOT AND MOUTH (HFM) DISEASE

This is a mild vesicular eruption caused by a coxsackie A virus (usually A $_{16}$). HFM disease affects both children and adults but typically children under the age of 10. Known as 'creche disease' it often occurs among groups of children in child care centres.

Clinical features

- Incubation period 3 - 5 days
- Initial fever, headache and malaise
- Sore mouth and throat
- The rash appears after 1 or 2 days.
- Starts as a red macule, then progresses to vesicles.
- Vesicles lead to shallow ulcers on buccal mucosa, gums and tongue
- Greyish vesicle with surrounding erythema
- On hands, palms and soles (usually lateral borders)
- May appear on limbs especially buttocks and genilah.
- Lesions resolve in 3 - 5 days
- Healing without scarring
- Spread by direct contact or aerosol droplets

- Virus excreted in faeces and saliva for several week
- Children are infectious until the blisters hare disappeared.

Management

- Reassurance and explanation
- Symptomatic treatment
- Careful hygiene
- Exclusion not usually recommended.

SCARLET FEVER

Scarlet fever results when a Group A *Streptococcus pyogenes* organism produces erythrogenic toxin. The prodromal symptoms prior to the acute exanthem comprise about 2 days of malaise, sore throat, fever (may be rigors) and vomiting. A throat swab should be taken.

Features of the rash

- Appears on second day of illness
- First appears on neck
- Rapidly generalised
- Punctuate and red
- Blanches on pressure
- Prominent on neck, in axillae, groin, skinfolds (Figure 83.10)


circumoral pallor
strawberry tongue
background erythema with tiny fine superimposed puncta (scarlet spots)
Pastia's lines

Figure 83.10 Scarlet fever: typical presentation of rash

- Absent or sparse on face, palms and soles
- Circumoral pallor
- Feels like fine sandpaper
- Lasts about 5 days

- Fine desquamation

Treatment

Phenoxymethylpenicillin (dose according to age) with rapid resolution of symptoms. Children can return to school 24 hours after taking antibiotics.

KAWASAKI'S DISEASE (MUCOCUTANEOUS LYMPH NODE SYNDROME)

This is an uncommon acute multisystemic disorder in children, characterised by an acute onset of fever > 39 °C of 5 days or more and accompanied by the following features:

- bilateral non exudative conjunctivitis
- maculopapular polymorphous rash
- ± cervical lymphadenopathy > 1.5 cm
- dryness, redness and cracking of the lips
- erythema of the oral cavity
- erythema of palms and soles with induration and oedema
- desquamation of fingertips (a characteristic), also of feet and groin in convalescence.
- tender mass in right hypochondrium

Kawasaki's disease can be elusive as there are variations with incomplete manifestations. There is no specific test but the ESR and CRP are usually elevated. Think of it in a miserable child with a high fever > 5 days with poor response to paracetamol.

The disease is generally benign and self-limiting but it is important to make an early diagnosis because early treatment may prevent complications. The major complication is vasculitis, which causes coronary aneurysms in 17 - 31% of cases, with an overall case fatality rate of 0.5 - 2.8% [5] due to the aneurysm that usually develops between the second week and the second month of the illness. Early treatment with immunoglobulin and aspirin has been shown to be effective in reducing the prevalence of coronary artery abnormalities. Echocardiography is indicated to detect these aneurysms and determine prognosis. Avoid corticosteroids in these patients.

MUMPS (EPIDEMIC PAROTITIS)

Mumps is an acute infectious disease caused by a paramyxovirus, with an affinity for the salivary glands and meninges. Although it most often affects children (90% present before adolescence), no one is exempt.[3]

Mumps has a worldwide prevalence. Most adults have antibodies to it, whether or not they have had the clinical infection. Because the antibody crosses the placenta in pregnancy, the infant will be immune for the first 6 - 9 months of life. One episode of the illness is sufficient to confer permanent immunity.

The patient is most infective during the prodrome, less so by the time the parotid glands are enlarged. Spread of infection is by aerosol droplets from the saliva and nasopharynx and can be rapid in school classrooms and throughout a household.

Mumps in a woman in early pregnancy occasionally causes abortion or foetal abnormalities.

General course and symptoms

The incubation period is 2 - 3 weeks.

The patient might be free of symptoms but a high fever, headache and malaise, for 5 - 7 days (occasionally 2 - 3 weeks), is usual. Involvement of the salivary glands is common. Dry mouth and discomfort on eating or opening the mouth occur.

Major manifestations

Unilateral or bilateral inflammation of the parotid gland is usual: one parotid gland swells first and in 70% of cases the opposite side swells after 1 - 2 days. The submandibular and sublingual glands are less commonly involved. About 6% of patients will have presternal oedema resembling cellulitis of the neck.

Complications

The complications are summarised in Table 83.6.

Table 83.6 Complications of mumps

Common
• Orchitis
• Aseptic meningitis (benign)
• Abdominal pain (transient)
Rare
• Oophoritis
• Encephalitis
• Arthritis (one or several joints)
• Deafness (usually transient)
• Pancreatitis

Orchitis, usually unilateral, occurs in 25% of postpubertal males, developing 3 - 4 days after parotitis. Subsequent sterility is rare, even if both testes are affected.

Aseptic meningitis is common but benign. Many patients suffer transient abdominal pain and vomiting: severe pancreatitis is a rare complication.

Clinical diagnosis

Enlargement of the cervical lymph glands can be mistaken for parotitis but the correct diagnosis is indicated by the anatomy of this area. Lymph nodes are posteroinferior to the ear lobe; the parotid gland is anterior and, when enlarged, obscures the angle of the mandible.

Bacterial (suppurative) parotitis is associated with toxaemia and results in a high leucocyte count. The skin over the parotid gland is tense and shiny and Stensen's duct might discharge pus.

Rare disorders such as Sjögren's syndrome can be misdiagnosed as mumps.

Virological diagnosis

The diagnosis of mumps is usually clinical; virological confirmation is rarely required but the virus can be isolated from the nasopharynx or saliva during the acute illness (and from cerebrospinal fluid in mumps meningitis).

A serological test for antibodies is available.

Management

Treatment is symptomatic. Paracetamol may be prescribed for fever, meningitis and orchitis. Ample fluid intake and a bland diet is advisable. Bed rest should be taken only according to the symptoms: it does not seem to have an in-fluence on the development of complications.[3]

Children should not return to school until the symptoms subside but contacts need not be excluded.

The patient with orchitis should use supportive underwear. Steroids may be prescribed to relieve severe pain but will have no other effect; nor will they reduce the risk of testicular atrophy.

Prevention

Isolation is generally ineffective. The best protection is immunisation of all children given at 12 months and 4 - 5 years.

EPSTEIN-BARR MONONUCLEOSIS

Although glandular fever is more common in adolescents and young adults, it can occur in young children but is often asymptomatic or atypical. The differential diagnosis includes cytomegalovirus infection and acute lymphatic leukaemia. Diagnosis is confirmed by specific antibody tests. Refer to Chapter 29.

PERTUSSIS

Pertussis (whooping cough) is a respiratory infection caused by *Bordetella pertussis* and occurs worldwide. The incidence of this infectious disease has diminished because of immunisation programs and improvements in standards of living, but the infection is still seen frequently, often modified by partial immunity.

Pertussis is predominantly an illness of infants under 2 years of age (up to 50% of all cases). Approximately 70% of unimmunised children will eventually develop pertussis, the majority by their fifth birthday.[3] However, no age is exempt. The source of infection is older children or young adults who have relatively mild disease.

The illness is characterised by three stages - catarrhal, paroxysmal and convalescent - with the person being most infectious during the catarrhal stage.

Suspect pertussis in an illness lasting 2 weeks or more with one of:

- paroxysms of coughing *or*
- inspiratory 'whoop' without other apparent causes *or*
- post-tussive vomiting

CLINICAL FEATURES

- Incubation period 7 - 14 days
- Catarrhal stage (7 - 14 days)
  - anorexia
  - rhinorrhoea
  - conjunctivitis/lacrimation
  - dry cough
- Paroxysmal stage (about 4 weeks)
  - paroxysms of severe coughing with inspiratory 'whoop'
  - vomiting (at end of coughing bout)
  - coughing mainly at night
  - lymphocytosis (almost absolute)
- Convalescent stage
  - coughing (less severe)

Note: Physical findings are minimal or absent.

DIAGNOSIS

The diagnosis is basically a clinical one - virtually no other acute infectious illness in children causes a cough that lasts 4 - 8 weeks.[3] Confirmed by culture and/or PCR of nasopharyngeal aspirate (within 1 week from onset of cough) or IgA serology (late in disease), although this can be misleading. High-grade lymphocytosis (12 - 25 × 10^9/L) on an FBE is strongly suggestive of pertussis. New methods include immunofluorescence and ELISA techniques.[6]

Differential diagnosis

Viral pneumonia, acute bronchitis, influenza. Chlamydia respiratory infection can cause a 'pseudopertussis' type of illness in infants.

Complications

These include asphyxia, hypoxia, convulsions and cerebral haemorrhage. Also pulmonary complications (e.g. atelectasis, pneumonia, pneumothorax).

Treatment

Clarithromycin (child 7.5 mg/kg up to) 500 mg 12 hourly for 7 days (or) erythromycin estolate for 7 days may help reduce the period of communicability (but not the symptoms) if given early (cough less than 3 weeks). There is no evidence that antibiotics produce an improvement in the patient.[7] Cough mixtures are ineffective.[3] Good ventilation is important: avoid dust and smoke, and also emotional excitement and overfeeding during the paroxysmal phase.

Almost all infants under 6 months and some who are older require admission to hospital[2] because the danger of apnoea and choking. School exclusion is necessary until at least 5 days of antibiotic use.

Prevention

Active immunisation with pertussis vaccine.

Prophylaxis

A 7 day course of erythromycin (cotrimoxazole if contraindicated) is recommended for household and other close contacts, regardless of immunisation status, commenced within 3 weeks of onset of cough in the patient.[6]

HERPES SIMPLEX

HSV infection is common and widespread. Primary HSV infection is basically a disease of childhood, presenting as severe acute gingivostomatitis. However, the infection may be subclinical in children; based on antibody studies, approximately 90% of the population acquire HSV infection before the age of 4 or 5 years.[5]

The primary infection

Typical clinical features:

- children 1 - 3 years
- fever and refusal to feed
- ulcers on gums, tongue and palate
- prone to dehydration
- may be lesions on face and conjunctivae
- resolution over 7 - 14 days

These children are generally very miserable and ill, and some may require hospitalisation for intravenous therapy to correct fluid and electrolyte loss. Push oral fluids. Monitor the urine output. Treatment is usually symptomatic (e.g. oral lignocaine gel). Careful nursing and prevention of secondary infection is important. The latter includes gentle mouth toilets. Children with severe infections, those who are immuno-suppressed and those with eczema herpeticum can have aciclovir IV or orally.

Serious complications:

- encephalitis can develop in otherwise healthy children
- eczema herpeticum - children with eczema can get widespread herpetic lesions
- disseminated HSV infection in neonates (avoid contact until recovered)

HERPES ZOSTER

Herpes zoster (shingles) is caused by reactivation of varicella zoster virus (acquired from the primary infection of chickenpox) in the dorsal root ganglion. It occurs at all ages and can occur in children, including infants, who have been exposed to varicella in utero.[3]

Recurrences are uncommon except in immunocompromised patients. The diagnosis is a clinical one but can pose difficulties, especially as it is not so common in childhood and may present with only a few vesicles.

IMPETIGO

Impetigo (school sores) is a contagious superficial bacterial skin infection caused by *Streptococcus pyogenes or Staphylococcus aureus* or a combination of these two virulent organisms.

There are two common forms:

1. vesiculopustular with honey-coloured crusts (either *Strep.* or *Staph.*)
2. bullous type, usually *S. aureus*

Ecthyma is a deeper form of impetigo, usually on the legs and other covered areas.

Treatment

If mild with small lesions and a limited area:

- Topical antiseptic cleansing with gentle removal of crusts, using antibacterial soap, salme chlorhexidine or povidone-iodine. Then mupirocin (Bactroban), a small amount tds for 10 days. Topical antibiotics other than mupirocin 2%, (Bactroban) are not recommended.[7]
- Other measures to minimise recurrence or transmission of bacteria:[8]
  - daily bath with Oilatum Plus bath oil for 2 - 4 weeks
  - hot water wash for clothes, towels and linen for 2 - 4 weeks
  - regular hand washing

If extensive and causing systemic symptoms:[7]

- Flucloxacillin/dicloxacillin 6.25 mg/kg up to adult dose (250 mg) (o) 6 hourly for 10 days *or*
- Cephalexin 6.25 mg/kg up to adult dose (250 mg) (o) 6 hourly for 10 days (first choice)

Boils (furunculosis) and carbuncles - same treatment as impetigo.

If *Streptococcus pyogenes* is confirmed use benzathine penicillin IM *or* pheoxy methyl penicillin (orally)[9]

The child should be excluded from child-care settings until sores have healed fully.

HEAD LICE

Head lice is an infestation caused by the louse *Pediculus humanus capitis*. The female louse lays eggs (or 'nits') which are glued to the hairs; they hatch within 6 days, mature into adults in about 10 days and live for about a month. Head lice spread from person to person by direct contact, such as sitting and working very close to one another. They can also spread by the sharing of combs, brushes and headwear, especially within the family. Children are the ones usually affected, but people of all ages and from all walks of life can be infested. It is more common in overcrowded living conditions. Resistance to the usual agents is becoming a problem.



Figure 83.11 Louse (enlarged)

Clinical features

- Asymptomatic or itching of scalp
- White spots of nits can be mistaken for dandruff
- Unlike dandruff, the nits cannot be brushed off
- Diagnosis by finding lice (or 'nits')
- 'Wet combing' improves detection rate

Treatment (topical)[10]

- Pyrethrins/piperonyl foam or shampoo (e.g. Lyban foam) - leave minimum 20 minutes *or*
- Permethrin 1% leave minimum 20minutes *or*
- Maldison 0.5% (leave 8 hours)

Method

Massage well into wet hair, then wash off thoroughly. Repeat treatment at 7 - 10 days.

Treat household child contacts at the same time.

Remove nits after first treatment with hair conditioner and fine tooth comb.

Note: The hair does not have to be cut short. All members of the family must be treated if lice or nits are found on inspection. Wash clothing, towels and sheets after treatment using a normal machine wash or hot soapy water. The hot cycle of a dryer is also effective for killing lice on bedding. School exclusion should not be necessary after proper treatment. For eyelash involvement, apply petrolatum bd for 8 days and then pluck off remaining nits.

Follow up with combing hair wetted by conditioner weekly.

Resistant head lice

Repeat treatment using another insecticide.

A US RCT showed that 1% permethrin plus a 10 day course of cotrimoxazole was the best treatment for resistant head lice.[12]

Handy tips for removing nits

- Combing with a hair conditioner is useful to remove the nits *or*
- Apply a 1:1 mixture of water and vinegar, leave 15 minutes, then comb with a fine-toothed comb.

Prevention

- Apply conditioner to dry hair once a week and them comb it out with a fine comb.

WORMS (HELMINTHS)[13]

Worms that inhabit the human intestine can be classified into:

- nematodes (roundworms)
- cestodes (tapeworms)
- trematodes (flukes)

The roundworms, which include pinworm (*Enterobius vermicularis*), whipworm (*Trichuris trichiura*), human roundworm (*Ascaris lumbricoides*), human threadworm (*Strong-yloides stercoralis*) and hookworm are the most prevalent worldwide and are usually asymp-tomatic in infected people in first world countries.

Pinworm

Also known as ‘threadworm’, it is a ubiquitous parasite infesting mainly children of all social classes. They are tiny white worms about 1 cm long which multiply profusely and are spread readily between individuals by close contact. Virtually all children have been infected by the time they reach high school but at any one time approximately 50% of the 5 - 10 year old age group will harbour pinworms.[13]

Symptoms (usually asymptomatic)

- Pruritus ani (in about 30% of cases)
- Diarrhoea (occasionally)

Diagnosis

- Inspect anus in child about 1 hour after going to sleep (Figure 83.12)
- Collect eggs with adhesive tape on perianal skin early morning - send to laboratory



Figure 83.12 Pinworms: female worms appearing soon after patient asleep to lay eggs

Management (troublesome cases)

- Scrupulous hygiene by family
- Hands should be washed thoroughly after toileting and before handling food
- Clip fingernails short (eggs lodge under nails)
- Patient should wear pyjamas (not nightgowns) and shower each morning
- Bed linen, nightwear and underwear changed and washed in very hot water daily for several days
- Vacuum room of affected person daily
- Have a veterinarian check any pets, especially dogs

Medication

- Any one of pyrantel, albendazole or mebendazole - as single dose orally
  - Pyrantel 10mg/kg up to 750 mg *or*
  - Mebendazine 100 mg (child < 10 kg: 50 mg) *or*
  - Albendazole 400mg (child < 10 kg :200 mg)
- Repeat in 2 - 3 weeks - both patient and household contacts

Human roundworm

Adult worms are about 20 - 40 cm long and are usually acquired from contaminated food and water overseas. They are mainly light infections that rarely cause problems but come to notice when they emerge from the anus (cause anxiety in the family!) or invade the lung (cause respiratory symptoms) or seen in radiological contrast examinations. Diagnosis is by finding eggs in the faeces. The worm is very sensitive to any of the three agents used for pinworm. A first-line option is pyrantel 20 mg/kg up to 750 mg orally, as a single dose - to be repeated after 7 days if a heavy infestation.

Whipworm

These used to be common in Indigenous communities, possibly causing failure to thrive, anaemia abdominal pain and diarrhoea with heavy chronic infestation. The worms are about 1 - 2 cm long. Diagnosis is by faecal microscopy. Treatment is with single large doses of mebendazole or albendazole.

Hookworm

Found in humid tropical regions but now uncommon in Northern Australia. About 1 - 1.5 cm long they can cause iron/protein deficiency anaemia in chronic infestation in addition to respiratory and abdominal symptoms. As with other helminths, diagnosis is by finding eggs on microscopy of faeces. Treatment is with a single dose of mebendazole or pyrantel.

Human threadworm (Strongyloides)[13]

These are tiny parasites (2 mm or so) and have a worldwide distribution. Infestation can cause symptoms such as recurrent abdominal pain and diarrhoea with blood eosinophilia. Strongyloides can live and reproduce in the body for many years. The problem is aggravated by corticosteroid therapy. High-risk people include migrants and refugees from tropical developing countries, returned soldiers, former prisoners of war from South-East Asia and workers or residents in northern Aboriginal communities.

DxT: abdominal pain (low grade) + recurrent diarrhoea + blood eosinophilia = strongyloides

Diagnosis

- Detecting faecal larvae or duodenal biopsy

Treatment

- Ivermectin 200 mg/kg (o) as single dose or albendazole 200 mg bd for 3 days

REFERENCES

1. Jarman R. A word about aspirin in children. Aust Pediatric Rev, 1991; 1(6): 2.
2. Efron D. *Paediatric Handbook* (5th edn). Melbourne: Blackwell Science, 1996: 69.
3. Robinson MJ. *Practical Paediatrics* (2nd edn). Melbourne: Churchill Livingstone, 1990: 217 - 27.
4. Mansfield F. Erythema infectiosum. Slapped face disease. Aust Fam Physician, 1988; 17: 737 - 8.
5. Wilson JD et al. Harrison's Principles of Internal Medicine (12th edn). New York: McGraw-Hill, 1991: 1462 - 3.
6. Golledge C. A case of persistent cough. Aust Fam Physician, 1997; 26: 1219.
7. Spicer WJ (Chair) *Antibiotic Guidelines* (version 12). Melbourne: Therapeutic Guidelines, 2006: 249 - 51.
8. Hogan P. Impetigo. Aust Fam Physician, 1998; 27(8): 735 - 6.
9. Marley J (Chair). *Dermatology Guidelines* (Version 2). Melbourne: Therapeutic Guidelines, 2004: 203 - 4.
10. Ibid.: 227 - 8.
11. *Rudolph's Paediatrics* (20th edn.) Stamford, CA: Appleton & Lange; 1996: 12 - 13.
12. Hipolito RB et al. Head lice infestation: single drug versus combination therapy with one percent permethrin and trimethoprim/sulfamethoxazole. Pediatrics, 2001; 107: e30.
13. Prociv P. Gastrointestinal worm infections. Aust Fam Physician, 2001; 30(8): 755 - 61.

CHAPTER 84. BEHAVIOUR DISORDERS IN CHILDREN

> The habit of constantly gratifying every wayward wish and temper under the plea of illness, and the constant indulgence which it meets with in this form from a mother's overkindness, exert a most injurious influence on the child's character, and it grows up a juvenile hypochondriac.
>
> CHARLES WEST (1816 - 98), FOUNDER OF THE GREAT ORMOND STREET HOSPITAL FOR SICK CHILDREN

BEHAVIOUR DISORDERS IN CHILDREN

The prevalence of significant psychiatric disorders in children in Western society is at least 12% in the age range 1 - 14 years, with an increase of 3 - 4% after puberty.[1] Most of these disorders are behavioural but there is a significant incidence of emotional disorders, such as anxiety and depression, which tend to be misdiagnosed, as there is a perception that children do not suffer from these psychiatric disorders in the same way as adults.

The author has observed that most of the personality characteristics and behavioural problems of infancy tend to remain throughout childhood and adolescence and form the personality and behavioural disposition of the adult, although many associated problems do not tend to persist into adult life.

Parry,[2] who describes the five phases of childhood development (Table 84.1), emphasises the importance of the first phase of infancy (where the infant is learning to trust the environment) as crucial to overall normal development.

Table 84.1 The five phases of childhood development

1 Infancy	sense of trust
2 Early childhood	sense of autonomy (independence)
3 Preschool	sense of initiative
4 School age	sense of industry
5 Adolescence	sense of identity

The second phase, in which the child is developing independent skills in the second and third years of life, is also an important phase and needs to be based on a secure and smooth first phase. It is in this toddler stage that many of the behavioural disorders will be discussed.

TEMPER TANTRUMS

The tantrum is a feature of the 'terrible twos' toddler whose protestation to frustration is a dramatic reaction of kicking, shouting, screaming, breath-holding, throwing, or banging of the head. They usually start at 15 - 18 months and often persist until 3 - 4 years.[3] Tantrums are more likely to occur if the child is tired or bored. This behaviour may be perpetuated if the tantrums are inadvertently rewarded by the parents to seek peace and avoid conflict.

A careful history is required to gain insight into the family stresses; it also allows parents to ventilate their feelings. Ask the parents exactly what the child does during a tantrum - what they do during and after and what causes the tantrums.

Management

Reassure parents that the tantrums are relatively commonplace and not harmful. Explain the reasons for the tantrums and include the concept that 'temper tantrums need an audience'.

Advice[2]

- Ignore what is ignorable: parents should pretend to ignore the behaviour and leave the child alone without comment, including moving to a different area (but not locking the child in its room).

- Stay calm and say nothing.
- Don't give in.
- Avoid what is avoidable: try to avoid the cause or causes of the tantrums (e.g. visiting the supermarket).
- Distract what is distractable: redirect the child's interest to some other object or activity.
- Praise appropriate behaviour.

When ignored, the problem will probably get worse for a few days before it starts to improve. Medication has no place in the management of temper tantrums.

BREATH-HOLDING ATTACKS

The age group for the attack is 6 months to 6 years (peak 2 - 3 years). There are two distinct types - one occurring with a tantrum and the other a simple faint in response to pain or fright. They can also manifest as ‘blue attacks’ due to breath-holding with a closed glottis or ‘white attacks’ known as reflex anoxic seizures, often in response to pain. The precipitating event can be of a minor emotional or physical nature. In the tantrum situation children will emit a loud cry, then hold their breath. They become pale, then cyanosed (this distinguishes it from a seizure disorder). If prolonged it may result in jerky movements, unconsciousness or a fit. The episode lasts 10 - 60 seconds.

Management

Place the child in the coma position. Reassure parents that the attacks are self-limiting and are not associated with epilepsy or intellectual disability. Advise parents to maintain discipline and to resist spoiling the child. Try to avoid incidents known to frustrate the child or to precipitate a tantrum by distraction methods. Explain to parents that gently flicking cold water in the child's face may abort the attack.

HEAD BANGING

This behaviour is common, occurring in 5 - 15% of normal infants and toddlers.3 It also occurs in developmental disability and severe emotional deprivation. It is quite different to a child hitting the head with the hands.

Features

- Occurs in children under 4, especially 3 years old
- Usually prior to going to sleep
- Head banging occurs 60 - 80 times/minute
- Lasts several minutes to 60 minutes or more per episode
- Associated repetitive movements (e.g. body rocking, thumb sucking)
- Child usually not distressed and rarely self-injurious

Management

- Reassure parents that it's a normal self-limiting behaviour and usually settles by 3 - 5 years.
- Avoid reinforcing behaviour by excessive attention or punishment.
- Advise distraction or actively ignoring the behaviour.
- Place the bed or cot in the middle of the room away from a wall (reduces disruption from noise).
- Restrict bed time (if appropriate).

OPPOSITIONAL BEHAVIOUR[2]

Resistant and oppositional behaviour is common and perhaps normal from time to time provided it is not associated with antisocial behaviour. It is a feature of 2 - 4 year olds as well as school-age children and adolescents. It is important to interview the child and family to determine whether it is normal or abnormal. Generally, supportive counselling and behaviour modification works effectively. This includes looking for and praising or rewarding good behaviour. It is important to praise (or chastise) behaviour rather than the child. 'Time out' is the preferred disciplinary measure for children over 18 months (maximum 1 minute per year of age) while withdrawing privileges is appropriate for children over 6 years.

CONDUCT DISORDERS

Conduct disorders affect 3 - 5% of children and represent the largest group of childhood psychiatric disorders.

Clinical features[3]

- Antisocial behaviour that is repetitive and persistent
- Lack of guilt or remorse for offensive behaviour
- Generally poor interpersonal relationships
- Manipulative
- Tendency to aggressive, destructive, 'criminal' behaviour
- Learning problems (about 50%)
- Hyperactivity (one-third)

Family and environmental factors[3]

- Disrupted childhood care
- Socially disadvantaged
- Lack of a warm, caring family
- Family violence: emotional, physical or sexual abuse
- Antisocial peer group exposure

Management

- Early intervention and family assistance to help provide a warm, caring family environment
- Family therapy to reduce interfamily conflict
- Appropriate educational programs to facilitate self-esteem and achievement
- Provision of opportunities for interesting, socially positive activities (e.g. sports, recreation, jobs, other skills)
- Behaviour modification programs
- With physically aggressive behaviour, refer for psychotherapy if repeated, severe, causes injury or is associated with other antisocial behaviour

STEALING[3]

Isolated theft, which is common, is not necessarily an indication of serious psychopathology but may reflect normal risk-taking behaviour, a reaction to stress, low self-esteem, searching for peer group acceptance or a 'cry for help and attention'.[3]

Management

- Insist on retribution - return of goods or payment and personal apologies to the 'victim'.
- 'Punish' with withdrawal of appropriate privileges.
- Refer for psychotherapy if persistent.

SLEEP DISORDERS/PARAINSOMNIAS

Refer to Chapter 71.

POOR EATING

Some parents may complain that their toddler ‘eats nothing’. Apart from taking a careful history about what constitutes ‘nothing’, it is useful to describe the typical diet for the age group and then match the child's weight on the normal growth chart. The important aspect of management is to point out what is necessary from a nutritional viewpoint as opposed to what is considered normal for the particular culture.

ATTENTION DEFICIT HYPERACTIVITY DISORDER (ADHD)

This disorder, which is characterised by developmentally inappropriate degrees of inattentiveness, overactivity and impulsiveness, has an estimated prevalence of about 2 - 5%. It is far more common in boys than girls (6:1) and is usually present from infancy.[1] About 60% will carry some degree of the disorder into adulthood. A neurological basis for ADHD has been demonstrated. Accurate diagnosis of ADHD is very important but can be problematic. It is usually inappropriate to make a diagnosis under the age of 4 years.

Diagnostic criteria

1. Either 1 or 2 (refer to diagnostic criteria for ADHD in DSM IV):
    a. inattention
    b. hyperactivity and impulsiveness
2. Onset no later than 7 years of age
3. Symptoms must be present in two or more situations (e.g. at school and at home)
4. Disturbance causes clinically significant distress or impairment in social, academic or occupational functioning
5. Not part of a pervasive developmental disability, psychotic disorder, mood disorder, anxiety disorder, associative disorder or a personality disorder

Other clinical features (may be present)

- Irritability and moodiness
- Poor co-ordination (clumsiness)
- Disorganisation
- Social clumsiness
- Learning difficulties
- Low self-esteem

Differential diagnoses

- Auditory processing disorder
- Mild intellectual disability
- Learning difficulty
- Family psychosocial stresses

Diagnosis

- No foolproof diagnostic tests available
- Psychometric tests available
- Questionnaires (such as Connor's rating scale) for parents and teachers

Assessment should include child and family interviews, neurological examination, assessment of vision and hearing levels, serum lead levels (in high-risk groups) and the testing of formal cognitive achievement. Twenty-five per cent of children with ADHD have coexistent learning disabilities.[3] It is essential that an accurate diagnosis be made before treatment is commenced. Shared care is an important principle and referral to a consultant is recommended.[4]

Management

- Protect child's self-esteem.
- Counsel and support family.
- Involve teachers.
- Refer to appropriate consultant (e.g. child psychiatrist).
- Refer to parent support group.

Diet. Exclusion diet probably ineffective but encourage good diet (consider dietitian's help).

Pharmacological. Based on psychostimulants for >4 years (2 doses: after breakfast and noon):[3,4]

- methylphenidate (Ritalin) 0.3 - 1 mg/kg (o) daily in 2 divided doses (usually 5 mg once or twice daily at first) *or*
- dexamphetamine 0.15 - 0.5 mg/kg (o) daily (usually 2.5 mg once or twice daily at first)

Other available agents:

- methylphenidate extended release tabs
- atomoxetine

Medication should be commenced at the low end of the dose range and gradually increased until satisfactory response or adverse effects (e.g. movement disorders, nervousness, psychosis)

- antidepressants, second line
- clonidine, especially for sleep disturbance and aggression

SIBLING RIVALRY

Sibling rivalry is a real concern as a toddler acts out apparent jealousy towards a new baby. The baby needs help from the inappropriate prodding, pinching and smothering attempts. The jealous toddler needs attention from the mother and a fair share of the comforts, cuddling and love that the toddler has been used to having.

It is important that the toddler is encouraged to feel that it is his or her baby too and to have opportunities to experience warmth and smiles from the baby, so that a sense of belonging is engendered.

STUTTERING AND STAMMERING[3]

This interruption of the orderly flow of speech may be accompanied by blinking and various other tics. It tends to be common in the school years but at least 80% of sufferers become fluent by adulthood.[5] Some children who stutter may avoid speaking.

Features of stuttering

- More common in boys
- Has a familial pattern
- Usually begins under 6 years of age (starts between 2 - 5 years)
- No evidence of neurotic or neurological disorder
- Causes anxiety and social withdrawal

Management

Although most stutterers improve spontaneously, speech therapy from a caring empathic speech pathologist is very helpful in those with stuttering persisting beyond 12 months or so. It is recommended before 5 years of age and preferably from 2½ years. Excellent results are obtained in over 90% of cases.

HABIT COUGH[6]

This is a common problem usually affecting schoolage-children and occurs in the absence of underlying disease. It only occurs when the child is awake - not during sleep.

Habit cough is usually loud, harsh, honking or barking. It may follow an URTI and last for months.

It is a diagnosis by exclusion including PFTs and CXR.

Underlying triggers include inter-family problems and bullying or other perceived stress or anxiety. Management includes explanation, reassurance and CBT.

Referral for resistant cases is appropriate.

Other functional respiratory problems

- Hyperventilation
- Sighing dyspnoea
- Vocal cord dysfunction (see p. 615)

TICS (HABIT SPASM)

Tics are ‘sudden, rapid and involuntary movements of circumscribed muscle groups which serve no apparent purpose’.[5] Most are minor, transient facial tics, nose twitching, or vocal tics such as grunts, throat clearing and staccato semi-coughs. Most of these tics resolve spontaneously (usually in less than a year) and reassurance can be given.

TOURETTE'S DISORDER[4]

Also known as Gilles de la Tourette's syndrome or multiple tic disorder, Tourette's disorder usually first appears in children between the ages of 4 and 15 years (before 18 years) and has a prevalence of 1 in 10 000. Diagnosis is based on recurrent tics over a period > 1 year in which there is never a tic-free period for more than 3 months.

Clinical features

- More common in boys
- Bizarre multiple motor tics
- One or more vocal tics
- Echolia (repetition of others' words)
- Coprolalia (compulsive utterances of obscene words)
- Palilalia (repeating one's own words)
- Familial: dominant gene with variable expression

Treatment

The basis of treatment is psychoeducation, including reassurance that the person is not responsible for their tics.

Promote their self-esteem.

Consider TD support groups.

Medical treatment (if necessary) is haloperidol, clonidine or pimozide.[4]

AUTISM SPECTRUM DISORDERS

Autism spectrum disorders (pervasive development disorders, PDD) are lifelong neurodevelopmental disorders with onset before 36 months of age. A characteristic feature is impairment of social interaction, verbal and non-verbal communication skills and stereotyped behaviour and activities.[7]

The spectrum can be grouped (after DSM IV) as:

- autistic disorder
- Asperger's disorder
- atypical autism or PDD not otherwise specified (PDDNOS)
- Rett syndrome
- childhood disintegrative disorder

However, there is a wide range of presentations within the spectrum, with overlapping between the first three mentioned.

Autistic disorder

Described first by Kanner in 1943, it is a PDD commencing early in childhood; it affects at least four children in 10 000, boys four times as commonly as girls. It is not due to faulty parenting or birth trauma, but is a biological disorder of the CNS which may have multiple organic aetiologies.

Many autistic children appear physically healthy and well developed although there is an association with a range of other disorders, such as Tourette's disorder, tuberous sclerosis, epilepsy (up to 30% onset, usually in adolescence) and rubella encephalopathy. Most have intellectual disability but about 30% function in the normal range.

The children show many disturbed behaviours. The main features are presented in Table 84.2.

Table 84.2 A guide to the diagnosis of autistic disorder (after Tonge)[8]

1 Onset during infancy and early childhood.

2 An impairment of social interactions shown by at least two of the following:

- lack of awareness of the feelings of others
- absent or abnormal comfort seeking in response to distress
- lack of imitation
- absent or abnormal social play
- impaired ability to socialise, which may include gaze avoidance

3 Impairment in communication as shown by at least one of the following:

- lack of babbling, gesture, mime or spoken language
- absent or abnormal non-verbal communication
- abnormalities in the form or content of speech
- poor ability to initiate or sustain conversation
- abnormal speech production

4 Restricted or repetitive range of activities, interests and imaginative development, shown in at least one of the following:

- stereotyped body movements

- persistent and unusual preoccupations and rituals with objects or activities
- severe distress over changes in routine or environment
- an absence of imaginative and symbolic play

5 Behavioural problems:

- tantrums
- hyperactivity
- destructiveness
- risk-taking activity

The earliest signs of autistic spectrum disorder in infancy include:[7]

- excessive crying
- no response to cuddling if crying
- failure to mould the body in anticipation of being picked up
- stiffening the body or resisting when being held
- no babbling by 1 year
- resistance to a change in routine
- appearing to be deaf
- failing to respond or overacting to sensory stimuli
- persistent failure to imitate, such as waving goodbye
- a need for minimal sleep
- no single words by 16 months

The diagnosis remains difficult before the age of 2 years.

Latter features:

- fascination with certain toys/objects
- poor interaction with other children
- not pointing to objects, e.g. grabs parent's hands to show things

Asperger's disorder[4, 7]

Also known as high-functioning autism, Asperger's disorder or syndrome is a developmental disability. As in autistic disorder, features include:

- impairment in social interaction
- impairment in communication skills
- repetitive and restricted interests but (usually) no significant language delay

It is usually diagnosed at age 6 or older. Typical characteristics are:

- marked male preponderance
- normal or borderline intellectual ability
- normal (may be precocious) speech development
- emotional blunting
- have fixed and rigid routines (e.g. bed time routine, coffee time and place in morning) and can become very distressed if not met
- rigid/intolerant of change
- anxiety
- awkward motor skills/clumsiness
- mechanical, almost robotic patterns of speech
- lack of empathy or feeling

- obsessive focus on narrow interest e.g. reciting train schedules, weather patterns, dinosaurs

People with Asperger's usually seek friendships but lack the skills to make and maintain them.

Examples of behaviour

They have difficulty:

- in greeting someone appropriately and taking turns in conversation
- reading body language, such as noticing the signs that someone is bored, happy or sad
- understanding metaphor and common expressions, e.g. they look bemused to their feet when told ‘pull up your socks’

As a rule they can learn social rules and behaviour and so minimise or reduce their disability, but their fundamental difficulties tend to persist throughout life.

People with Asperger's disorder usually seek friendships but lack the skills to maintain them.

(Acknowledgement: Dr Jane Tracey)

Atypical autism (PDDNOS)

This diagnosis applies to the presence of core autistic behaviours but the criteria for autistic disorders are not fully met. However, management follows the same principles.

Assessment

If a child has delayed and deviant development and autistic spectrum disorder is suspected, a comprehensive multidisciplinary assessment is necessary. Referral to professionals, e.g. Child Development Unit, is essential as accurate diagnosis is important.

Treatment[7, 9]

Many treatments have been tried and behavioural treatment methods have proved to be the most helpful. Medications are unhelpful for autism per se although medications such as tranquillisers, antidepressants and anticonvulsants are helpful for associated disorders.

The best results are achieved by early diagnosis, followed by a firm and consistent home management and early intervention program. Remedial education and speech therapy have an important place in management.

CASE HISTORIES AND ‘DRAW A DREAM’

A useful strategy for communicating with disturbed children and getting to the source of a behaviour problem is to ask them to ‘draw a dream’.[10] Professor Tonge believes that the dream is the royal road to the child's mental processes and the family doctor is ideally placed to use this technique. The following case studies concerning insomnia and nightmares illustrate the importance of these symptoms as reflecting a deep emotional problem in the child.

CASE STUDY 1[11]

Steven, aged 6, was a bright, happy little boy until he developed an extraordinary and puzzling episode of insomnia which was solved eventually by his teacher.

He presented to our group practice with his bemused mother who claimed that, suddenly, he would not and could not sleep. His parents would be startled at night by the eerie vision of Steven standing silent and motionless beside their bed. When not in his bed at night he would be found hiding under it or in his wardrobe.

His behaviour was normal otherwise, but his teacher reported that his schoolwork had deteriorated and that he was constantly falling asleep at his desk. On direct questioning Steven was shy and evasive, claiming nothing was worrying him. We considered it was a temporary phase of abnormal behaviour and advised conservative measures such as hot beverages, baths and exercises before retiring, but this strategy failed. He was referred to a consultant who also failed to find the cause of the insomnia and advised long midnight jogs.

Eventually Steven's teacher had the bright idea of asking all the children to draw the thing that scared or worried them the most, stipulating that it would be a ‘make believe’ picture.

Looking at the drawing depicting two robbers stealing his moneybox as he slept (Figure 84.1), she tactfully confronted Steven, who admitted that his playmate had told him robbers would come one night, steal his moneybox and ‘bash’ him.



Figure 84.1 Steven's drawing

The final chapter of this story saw a happy Steven perched on a bank counter watching his money being counted, deposited in a huge safe and exchanged for a bank book. Steven has slept normally ever since.

CASE STUDY 2[10]

George, the second child of four children, seemed a normal healthy 3 year old when his mother presented him for assessment.

For about 3 months George had been having nightmares, episodes that fractured the entire household. His mother, Mary, was absolutely frustrated by his nocturnal behaviour and said she was ‘at her wit’s end'. As she excitedly rattled off details of the family dilemma, I noted that she was intense and rather domineering but obviously a very conscientious and dutiful wife and mother.

She explained that George would wake her at night calling out to her because of a monster in his room or outside his window. She had no idea about any causes for this problem and explained that ‘our household is very normal - no problems really’. She said George's behaviour was otherwise normal and he was a healthy boy.

Identifying the monster

I then asked George about his problem but could elicit only very scant information. Recalling the immense value of the ‘draw a dream’ strategy I asked him to draw the monster. George quickly drew the monster as shown in Figure 84.2. I asked him about the monster and finally confronted him with the question: ‘Do you know who or what the monster is?’.



Figure 84.2 George's drawing

‘Mum,’ replied George, very matter-of-factly.

A shocked Mary looked unbelievingly at George and, for once, seemed stuck for words. Realising the delicacy of the situation, I asked George to tell me what it was about his mother that worried him. He offered the very revealing information: ‘I don’t think that she loves me. She's always yelling at me.'

Obviously the monster was George's insecurity because George declared how much he did love his mother and was ‘scared’ of losing her love. With appropriate counselling the outcome was good.

A lesson learned often is that it is important to ‘look close to home’ for any significant behaviour disorder or other psychological problem. It is important to explore the relationship that is most meaningful to the affected person (e.g. mother - daughter, father - son, student - teacher).

The ‘draw a dream’ strategy revealed vital information in this case.

CHILDHOOD BULLYING

Research indicates that childhood bullying is common and up to 50% told no one about it. It was associated with school truancy and depression.

Indicative signs

- School phobia: sham sickness and other excuses
- Being tense, tearful and miserable after school
- Reluctance to talk about happenings at school
- Poor appetite
- Functional symptoms, e.g. habit cough
- Repeated abdominal pains/headache
- Unexplained bruises, injuries, torn clothing, damaged books
- Lack of a close friend; not bringing peers home
- Crying during sleep

- Restless sleep with bad dreams
- Appearing unhappy or depressed
- Unexpected irritability and moods; temper outbursts

It is important to encourage them to talk to their parents and/or family doctor and receive support. Cognitive behaviour therapy works well.

WHEN TO REFER[1]

- When child abuse is known or suspected
- When an underlying medical problem is present
- For assessment of associated psychological, family and related factors
- For failed management, including simple behavioural and family support interventions

REFERENCES

1. Tonge BJ. Behavioural, emotional and psychosomatic disorders in children and adolescents. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 52 - 5.
2. Parry TS. Behavioural problems in toddlers. Aust Fam Physician, 1986; 15: 1038 - 40.
3. Efron D, Davey M, Reilly S. In: Paxton G, Munro J. *Paediatric Handbook* (7th edn). Melbourne: Blackwell Science, 2003: 151 - 67.
4. Dowden J (Chair). *Therapeutic Guidelines: Psychotropic* (Version 5). Melbourne: Therapeutic Guidelines Ltd, 2003: 208 - 16.
5. Robinson MJ. *Practical Paediatrics* (2nd edn). Melbourne: Churchill Livingstone, 1990: 543 - 9.
6. Powell C, Brazier A. Psychological approaches to the management of respiratory symptoms in children and adolescents. Paediatric Respir Rev, 2004; 5: 214 - 24.
7. Curran J, Tonge B. Autism spectrum disorders. In: Lennox N, Diggens J. *Management Guidelines: People with Developmental and Intellectual Disabilities*. Melbourne: Therapeutic Guidelines Ltd, 1999: 197 - 204.
8. Tonge BJ. Autism. Aust Fam Physician, 1989; 18: 247 - 50.
9. Curtis J. Autism. Patient Education, 1993; 22: 1239.
10. Tonge BJ. ‘I'm upset, you're upset and so are my Mum and Dad'. Aust Fam Physician, 1983, 12: 497 - 9.
11. Murtagh J. *Cautionary Tales*. Sydney: McGraw-Hill, 1992: 165 - 74.

CHAPTER 85. CHILD ABUSE

It is customary, but I think it is a mistake, to speak of happy childhood. Children, however, are often overanxious and acutely sensitive. Man ought to be man and master of his fate; but children are at the mercy of those around them.

SIR JOHN LUBBOCK, BARON AVEBURY 1834 - 1913

CHILD ABUSE

The description of the 'battered child syndrome' in 1962 provoked an awareness of a problem facing children which continues to increase in prominence.The possibility of both physical and sexual abuse has to be kept in mind by the family doctor. It maysurface in families known to us as respectable and where a good trustful relationship exists between parents and doctor. Another aspect of child abuse is neglect.

The various types of abuse are classified as:

- physical
- neglect
- emotional
- sexual
- potential

Physical abuse occurs most often in the first 2 years of life, neglect in the first 5 years and sexual abuse from 5 years of age[1] (Figure 85.1). In a Community Services of Victoria study[2] the distribution ofchildabuse was physical15%, emotional 48%, sexual 9% and neglect 28%.



Figure 85.1 Typical relative age patterns for child abuse After Bentovin[1]

In another study,[2] thefindings were:

- Girls are more likely to be abused than boys.
- Girls are more often assaulted by someone they know.
- Most of the adults who sexually abuse are men (> 90%).
- About 75% of offenders are known to the child.
- Abuse is the misuse of a power situation (e.g. a close relative, coupled with the child's immaturity).

DEFINITIONS

Child abuse can be defined by the nature of the abusive act or by theresult of the abuse. A parent, guardian or other carer can harm a child by a deliberate act or by failure to provide adequate care.

Physical abuse (non-accidental injury)

Physical abuse is defined as ‘a child with a characteristic pattern of injuries, the explanation of which is not consistent with the pattern, or where there is definite information through acknowledgment or reasonable suspicion that the injury was inflicted or knowingly not prevented by any person having custody, charge or care of the child’.[1]

Neglect

Neglect is defined as ‘the privation of food, drink, medical care, stimulation or affection’.

Emotional abuse

Emotional abuse is the ‘systematic destruction of the child’s sense of self-esteem and com-petence, where competence is defined as the ability to act in social contexts'.

Sexual abuse

Sexual abuse in children is defined[3] as ‘the involvement of dependent, developmentally immature children and adolescents in sexual activities thatthey do not fully comprehend, to which they are unable to give informed consent, and which violate the social taboos’.

Munchausen syndrome by proxy

This is the term used when a parent or guardian creates an illness ina child so that the perpetrator can develop or maintain a re-lationship with medical staff or transfer their responsibilities. The masquerade may be simple or very sophisticaited. Children may be indirectly abesed by the lengthy or invacive investigatun.

Incest

Incest is legally defined as ‘intercourse between biological family members’.

Female genital mutilation

This comprises all procedures involving partial or total removal of the female external genitalia or other injury to the femal genital organs, whether for cultural or other non-therapeutic reasons (WHO definition). It is also referred to as female circumcision.

ABUSE: WHO AND WHY?[2]

The real cause seems to be a combination of several interrelated factors: personal, familial, social/cultural and societal stress. Abused children exist at all levels of society, although the majority of abused children who come to the attention of authorities are from families where there is high mobility, lack of education, loneliness, poverty, unemployment, in-adequate housing and social isolation. Sexual abuse, occurring alone, does not follow these patterns and can occur under any socio-economic circumstances.

Both men and women physically abuse their children. While women are the parents most responsible in cases of neglect and emotional abuse (probably because of a dominant role in child care, social and economic disadvantage and being the only one responsible for the care of children in a single parent arrangement), men are more likely to abuse their children sexually.

The child can be abused at any age (even adolescents can be victims of abuse and neglect). It is important to keep this in mind - *it does happen*.

UNDERDIAGNOSING AND UNDER-REPORTING[2]

Although the medical profession remains the foremost focus of child abuse reporting (they are the most likely to encounter injured children and are the most qualified to diagnose abuse), they still contribute only a small percentage of the total reporting to central registries. This could be because there is underdiagnosis of the problem but it could also be because of under-reporting.

Reasons given as to why GPs don't report more cases of child abuse include:

- concern about drain on time and finances
- lack of positive feedback about other cases
- lack of undergraduate education on the topic
- risk of alienation and stigmatisation to the family
- the feeling by some GPs that they can work on the problem with the family without outside intervention
- lack of trust or confidence in local officials and agencies
- uncertainty about *what* to do
- personal and legal risks (i.e. fear of court, libel suits, irate parents)
- reluctance until absolutely certain of diagnosis

It will always be difficult to take the first step but it is important and it can help, no matter how small that first step is.

INTERVIEWING PARENTS OR GUARDIANS

A skilled, sensitive, diplomatic interview is fundamental to management. Guidelines include:

- a relaxed non-judgmental approach
- sensitivity to all people involved
- appropriate questions - open-ended, not leading
- use verbatim quotes from the child where possible and wait silently for a reaction.

PHYSICAL ABUSE

Physical abuse should be suspected, especially in a child aged under 3, if certain physical or behavioural indicators in either the child or the parents are present.Bruising, especially fingertip bruising, is the most common sign of the physically abused child.

Physical indicators:[2]

- unexplained injury
- different explanations offered
- injury unlikely to have occurred in manner stated
- unreasonable delays between injury and presentation
- finger-shaped bruises thumb grup marks
- multiple bruises/welts of different ages, especially on face, buttocks (Figure 85.2), genitalia, earlobes



Figure 85.2 Physical abuse: imprint of a boot on the buttock of a child

- bruises in babies who are not yet pulling themselre up to walk
- fractures (especially if child < 2 years old)
- metaphyseal fractures of the proximal humerus and proximal or distal tibia are almost pathognomonic
- other common fractures
    - rib, clavicle, vertebral body
- burns, scalds, dislocations, poisoning, cuts, bites
- cigarette butt type burns
- shaking injuries (e.g. retinal damage, torn frenulum)
- subdural haematoma
- internal injuries
- episodes of unconsciousness

Remember Munchausen syndrome by proxy.

Behavioural indicators[2]

- Wariness of adult contacts
- Inappropriate clothing (e.g. long sleeves on a hot day)
- Apprehension when other children cry or shout
- Behavioural extremes
- Fear of parents
- Afraid to go home
- Child reports injury by parents or gives inappropriate explanation of injury
- Excessive compliance
- Extreme wariness
- Attaches too readily to strangers

Investigations

The following should be considered:[4]

- FBE, including platelets, if bruising
- coagulation studies (PT, APTT) if bruising

- imaging for fractures: specific X-rays, bone scan (especially < 3 years old), skull X-ray, skeletal survey (skull, thorax including clavicle, abdomen, pelvis and extremities)
- photographs: taken on presentation

Management

The family doctor should diplomatically confront the parent or parents and always act in the best interests of the child. Offer to help the family. An approach would be to say, ‘I am very concerned about your child’s injuries as they don't add up - these injuries are not usually caused by what I'm told has been the cause. I will therefore seek assistance - it is my legal obligation. My duty is to help you and, especially, your child'.

Acquiring essential help

- Psychosocial assessment of child and family: involves social worker and multidisciplinary assessment
- Admission to hospital: for moderate and severe injuries
- Case conference (where appropriate)
- Mandatory reporting: notify child protection authorities

The stages of management are:[1]

- recognition or disclosure of abuse
- the family separation phase
- working towards rehabilitation
- finding a new family for the child when rehabilitation fails

EMOTIONAL ABUSE

Physical indicators:

- There are few physical indicators, but emotional abuse can cause delay in physical, emotional and mental development.

Behavioural indicators:

- extremely low self-esteem
- compliant, passive, withdrawn, tearful and/or apathetic behaviour
- aggressive or demanding behaviour
- anxiety
- serious difficulties with peers and/or adult relations
- delayed or distorted speech
- regressive behaviour (e.g. soiling)

NEGLECT

Physical indicators:

- consistent hunger
- failure to thrive, or malnutrition
- poor hygiene
- inappropriate clothing
- consistent lack of supervision
- unattended physical problems or medical needs
- abandonment
- dangerous health or dietary practices

Behavioural indicators:

- stealing food
- extending stays at school
- consistent fatigue, listlessness or falling asleep in class
- alcohol or drug abuse
- child states there is no caregiver
- aggressive or inappropriate behaviour
- isolation from peer group

SEXUAL ABUSE

Incest and sexual abuse of children within the family occur more frequently than is acknowledged. One difficulty in recognising sexually abused children is to determine what is appropriate physical contact between adult and child, and what is abusive sexual behaviour.

Sexual abuse presents in three main ways:[5]

- allegations by the child or an adult
- injuries to the genitalia or anus
- suspicious presentations, especially:
  - genital infection(Figure 85.3)



Figure 85.3 Genital human papillomavirus infection - a sign of sexual abuse in a child

  - recurrent urinary infection
  - unexplained behavioural changes/psychological disorders

Clinical indicators that may suggest child sexual abuse are presentedin Table 85.1.[6, 7]

Table 85.1 Clinical indicators that may suggest child sexual abuse

• complaint of abuse (rarely invented)
• vaginal discharge
• other STI

- urinary tract infection
- unexplained genital trauma
- unexplained perianal trauma
- overt sexual play
- pregnancy in an adolescent
- deterioration in school work
- family disruption
- indiscriminate attachment
- abnormal sexual behaviour
- poor self-esteem
- psychological disorders
    - behaviour disturbances
    - regression in behaviour
    - sleep disturbances
    - abnormal fears/reactions to specific places or persons
    - psychosomatic symptoms
    - anxiety
    - lack of trust
    - overcompliance
    - aggressive behaviour
- depression
    - self-destructive behaviour
    - substance abuse
    - suicidal tendencies

Examination (abnormal findings uncommon)

- genital trauma
- perforated hymen/lax vagina
- perianal trauma
- vaginal discharge
- look for semen and STIs

Sexual abuse can take many forms, including:

- genital fondling
- digital penetration
- penetration with various objects
- simulated sexual intercourse (anal in boys)
- Infercrural intercousse
- full sexual penetration
- Masturbation of an adult

- Pornography
- Exposure to indectent acts
- prostitution

Clinical approach

Ideally, the child should be assessed by experienced medical officers at the regional sexual assault service, so the temptation for the inexperienced GP to have a quick look should be resisted. For the practitioner having to assess the problem, a complete medical and social history, including a behavioural history, should be obtained prior to examination.

The child's history must be obtained carefully, honestly, patiently and objectively, without leading the child. Use language appropriate to the child and employ aids such as drawings and a model (such as a ‘gingerbread man’) to help the child illustrate what has happened. The history is more important than the physical findings as there are no abnormal physical findings in 40% of confessed cases.[8]

Physical examination[9]

It is recommended that the physical examination of any child suspected of being sexually abused is performed by a paediatrician or forensic physician experienced inthe area of sexual abuse.

It is important to spend time explaining the examination process to the child and the accompanying parent. In prepubescent girls and boys, the examination is limited to visual inspection of the external areas using a good light source. Magnification may be used with a colposcope or magnifying glass, and photographs may be taken for documentation or medicolegal purposes.

Speculum examination is limited to post-pubertal girls or used if there is concern about internal injuries (the latter may necessitate general anaesthesia). Rectal examination is usually limited to visualisation.

Three recommended positions are:

- Supine with legs apart (frog-leg position with soles of feet apposed)
- Prone knee-chest position (the best position)
- Lateral decubitus

Always record the findings and note the examination position.

It is useful to remember that examination of urine in female childrenmay show sperm so, if the child is uncharacteristically passing urine at night, get her mother to collect a specimen.

The crisis situation

It is important to realise that the child will be in crisis.[6] Children are trapped into the secrecy of sexual abuse, often by a trusted adult, by powerful threats of the consequences of disclosure. They are giventhe great re-sponsibility of keeping the secret and holding the family together or disclosing the secret and disrupting the family. A crisis occurs when these threats become reality.

Management

It is important to act responsibly in the best interests of the child.When we encounter real or suspected child abuse, immediate action is necessary. The child needs an advocate to act on its behalf and our intervention actions may have to override our relationship with the family.

Some golden rules are:

- Never attempt to solve the problem alone.
- Do not attempt confrontation and counselling in isolation (unless under exceptional circumstances).
- Seek advice from experts (only a telephone call away).

- Avoid telling the alleged perpetrator what the child has said.
- Refer to a child sexual assault centre or Protective Services Unit where an experienced team can take the serious responsibility for the problem.

Supporting the child

- Acknowledge the child's fear and perhaps guilt.
- Assure the child it is not his or her fault.
- Tell the child you will help.
- Obtain the child's trust.
- Tell the child it has happened to other children and you have helped them.

Confronting the parents

If certain about the diagnosis the doctor should inform the child's mother and encourage her to notify the protective authority. If abuse is suspected, concerns should be raised that the child may have been sexually abused and that you wonder who the perpetrator could be. Ask who has access to the child (e.g. babysitter, member of a creche or kindergarten, teacher, other males, relatives).

PREVENTION OF CHILD ABUSE

Prevention of abuse, particularly self-perpetuating abuse, can be helped by creating awareness through media attention, programs in schools and the community in general, and increased knowledge and surveillance by all professionals involved with children. Clear guidelines on reporting and the accessibility of child abuse clinics are important for the strategies to be effective. Teaching children how to protect themselves offers the greatest potential for prevention.[5]

COUNSELLING THE SECONDARY VICTIMS

Non-offending parents, who are the secondary victims of the abused child, will require help and guidance from their family doctor on how to manage the crisis at home. Parents should be advised to reassure the child of support and safety and to maintain usual routines. The child should be allowed to set the pace, without zealous overattention and pressure from the parents. Siblings should be informed that something has happened but that the childis safe. Ensure that the child will inform if the perpetrator attempts further abuse. Parents need substantial support, including alleviation of any guilt.

An unhappy consequence of the crisis is the problem of broken relationships, which may involve the separation of the child from the family. At least one hitherto unsuspecting parent will be devastated if a parent is responsible for the abuse. The sexually abused child needs to be living with a protective parent with the abusive parent living separately.

Support for doctors

The attending doctor also requires support, and sharing the problem with colleagues, mentors and family is recommended. Some helpful guidelines are:

- Carefully record all examination findings (take copious notes).
- Always keep to the facts and be objective.
- Do not become emotionally involved.
- Work with (not for) the authorities.
- Avoid making inappropriate judgments to the authorities; e.g. do not state ‘incest was committed’, but rather say ‘there is evidence (or no evidence) to support penetration of … ’.
- If called to court, be well prepared; rehearse presentation; be authoritative and keep calm without allowing

yourself to be upset by personal affronts.

The main difficulty in diagnosing child abuse is denial that it could be possible.

ADULT SURVIVORS OF CHILD ABUSE

Putnam and others have reported a chronic post-traumatic stress disorder in many victims, with an increased incidence of somatic complaints, such as asthma, headache, skin disorders, in addition to anxiety and depression. They are also prone to poor self-image, substance abuse and relationship failure.[10] The possibility of child abuse shouldbe tactfully raised in people presenting with these problems.

BASIC RULES[4]

- Suspect child abuse.
- Recognise child abuse.
- Consult the child protection authorities.

Mandatory reporting

In most states of Australia and in many areas throughout the world itis man-datory to notify the relevant statutory authorities about suspected child abuse. All family doctors should become familiar with the appropriate local legislation.

Practice tips and guidelines

- A child's statement alleging abuse should be accepted as true until proved otherwise.
- Children rarely lie about sexual abuse.
- False allegations, however, are a sign of family disharmony and an indication that the child may need help.
- Do not insist that the child ‘has got it wrong’, even if you find the actions by the alleged perpetrator unbelievable.
- Do not procrastinate - move swiftly to solve the problem.
- The genitalia are normal in the majority ofsexually abused children.
- Be supportive to the child by listening, believing, being kind and caring.

WHEN TO REFER

Unless there are exceptional circumstances, referral to an appropriate child abuse centre where an expert team is available is rec-ommended. If doubtful, relativelyurgent referral to a paediatrician is an alternative.

REFERENCES

1. Bentovin A. Child abuse. Med Int, 1987: 1851 - 7.
2. Lewis D. Child abuse. In: Department of Community Medicine. Final year student handbook. Melbourne: Monash University, 1993: 164 - 8.
3. Schechter MD, Roberge L. Sexual exploitation. In: Helfer RE, Kempe CH eds. Child Abuse and Neglect. The Family and the Community. Cambridge, MA: Ballinger, 1976: 127 - 42.
4. Paxton G, Monaj Paediatric Handbook.7th edn.Melbourne: Blackwell publiser 2003:216-8
5. Valman HD. ABC of One to Seven. London: British Medical Association, 1988: 112 - 14.
6. McMichael A. Counselling the victims of child sexual assault. Aust Fam Physician, 1990; 19: 481 - 9.

7. Steven I, Castell-McGregor S, Francis J, Winefield H. Child sexual abuse. Aust Fam Physician, 1988; 17: 427 - 33.
8. Irons TG. Child Sexual Abuse. California: Audio Digest, 1993; 41: 2.
9. Murnane M. Child sexual abuse. Aust Fam Physician, 1990; 19: 603 - 6.
10. Kramer K. Dealing with the trauma of child sex abuse. Aust Doctor, 2001; 7 September: 49 - 50.

CHAPTER 86. EMERGENCIES IN CHILDREN

We can say with some assurance that, although children may be the victims of fate, they will not be the victims of our neglect.

JOHN F. KENNEDY (1917 - 63)

EMERGENCIES IN CHILDREN

Important serious emergencies in children include:

- trauma, especially head injuries and intra-abdominal injuries
- swallowed foreign bodies (FB)
- respiratory problems
    - bronchial asthma
    - epiglottitis
    - croup
    - inhaled FB
    - acute bronchiolitis
- severe gastroenteritis
- septicaemia (e.g. meningococcal septicaemia)
- myocarditis
- immersion
- poisoning
- bites and stings
- seizures
- febrile convulsions
- sudden infant death syndrome (SIDS) and apparent life-threatening episode (ALTE)
- child abuse
    - emotional
    - physical
    - sexual
- psychogenic disturbances
- anxiety/hyperventilation
- suicide/parasuicide

SURVEY BY AGE GROUP

The author's study analysed emergencies into three groups:[1] preschool (0 - 5 years), primary school (6 - 12), adolescence (13 - 17).

The commonest emergency calls in the 0 - 5 years group were poisoning, accidents and violence, dyspnoea, fever/rigors, convulsions, abdominal pain, earache, vomiting.

In the 6 - 12 years age group - accidents and violence, dyspnoea, abdominal pain, vomiting, acute allergy, bites and stings, earache.

In the 13 - 17 years age group - accidents and violence, abdominal pain, psychogenic disorders, acute allergy, bites and stings, epistaxis.

THE SIGNS AND SYMPTOMS OF A SERIOUS ILLNESS

> Babies who are febrile, drowsy and pale are at very high risk and require hospital admission.

The busy general practitioner will see many sick children in a day's work, especially in the winter months with the epidemic of URTIs. It is vital to be able to recognise the very sick child who requires special attention, including admission to hospital. It is unlikely that the commonplace robust, lustily crying, hot, red-faced child is seriously ill but the pale, quiet, whimpering child spells danger. These rules are particularly helpful in the assessment of babies under six months of age.[2, 3] The presence of a fever in itself is not necessarily an indication of serious illness but rather that the baby has an infection.[2]

> The features of a very sick infant include:
>
> - inactive, lying quietly, uninterested
> - increased respiratory rate
> - increased work of breathing
> - noisy breathing
>   - chest wall or sternal retraction
>   - wheezes, grunting, stridor
> - tachycardia
> - sunken eyes
> - cold, pale skin
> - cold extremities
> - drowsiness

A Melbourne study of the sensitivity of clinical signs in detecting serious illness in infants identified five key signs or markers.

Marker	Risk to baby
Drowsiness	58%
Pallor	48%
Chest wall retraction	41%
Temperature > 38.9°c or < 36.4°c	42%
Lump > 2 cm	42%

If sepsis suspected investigate with:

- blood culture
- FBE/ESR/CRP
- lumbar puncture
- urine culture

Serious illnesses to consider include:

- *Haemophilus influenza* type B (Hib) infection
  - acute epiglottitis
  - meningitis (now uncommon since Hib immunisation)
- acute bacterial meningitis
- septicaemia
  - meningococcaemia
  - toxic shock syndrome
  - other bacterial sepsis

- acute viral encephalitis
- acute myocarditis
- asthma/bronchitis/bronchiolitis
- pneumonia
- intussusception/bowel obstruction/appendicitis
- severe gastroenteritis

Predictive combinations

- Pallor + drowsiness + fever = meningitis
- Drowsiness + chest wall recession = pneumonia or severe bronchiolitis
- Pallor + inactivity = intussusception

Two main groups of signs are good indicators of serious illness.[2]

Group 1: common features with reasonable risk

'A, B, C, fluids in and out'

- A = poor arousal, alertness and activity
- B = breathing difficulty
- C = poor circulation (persistent pallor, cold legs to knees)
- 'Fluids in' = feeding less than half normal in 24 hours
- 'Fluids out' = less than 4 wet nappies in 24 hours

Note: The more signs present, the greater the risk.

Group 2: uncommon features with high risk requiring urgent referral

- Respiratory: apnoea, central cyanosis, respiratory grunt
- GIT: persistent bile-stained vomiting, mass > 2 cm other than hydrocele or umbilical hernia, significant faecal blood
- CNS: convulsions
- Skin: petechial rash

Indications for investigations are presented in Table 86.1.

Table 86.1 Indications for investigations in the sick child[2]

Urine microscopy, culture and sensitivity	All with fever
Full blood examination	All < 4 weeks Risk factors present Doctor uncertain
Blood microscopy, culture and sensitivity	All < 3 months Risk factors present Doctor uncertain
Faecal microscopy, culture and sensitivity	All with diarrhoea
Chest X-ray	All with significant respiratory tract symptoms and signs
C-reactive protein	Those on antibiotics Doctor uncertain
CSF examination (lumbar puncture contraindicated in the unresponsive febrile patient)	Suspected meningitis (infant drowsy, pale and febrile)

	Convulsion in febrile child and • source of fever unknown • preceding drowsiness and pallor • infant < 6 months child > 5 years • prolonged convulsion (> 10 minutes) • postictal phase longer than usual (> 30 minutes)

Note: Fever measured rectally with thermometer bulb 3 cm past anal verge.

COLLAPSE IN CHILDREN

Collapse in children is a very dramatic emergency and often represents a life-threatening event. It is important to remember that the child's brain requires two vital factors: oxygen and glucose

There is only a 2 minute reserve once cerebral blood flow stops. Bacterial meningitis should be considered as a cause.

Important causes of collapse are presented in Table 86.2. Keep in mind child abuse as a cause of collapse.

Table 86.2 Collapse in children: causes to consider

Anaphylaxis	Penicillin injection Stings
Asphyxia	Near drowning Strangulation
Airways obstruction	Asthma Epiglottitis Croup Inhaled foreign body
CNS disorders	Convulsions Meningitis Encephalitis Head injury
Severe infection	Gastroenteritis → dehydration Septicaemia Myocarditis
Hypovolaemia	Dehydration (e.g. heat) Blood loss (e.g. ruptured spleen)
Cardiac failure	Arrhythmias Cardiomyopathy
Metabolic	Acidosis (e.g. diabetic coma) Hypoglycaemia
Poisoning	Drug ingestion

	Envenomation
SIDS	Near miss
Functional	Breath-holding attacks Conversion reaction Vasovagal

Note: Consider child abuse.

Initial basic management[5]

1. Lay child on side.
2. Suck out mouth and nasopharynx.
3. Resqre breathe
4. Intubate or ventilate (if necessary).
5. Give oxygen 8 - 10 L/min by mask.
6. Pass a nasogastric tube:
   0 - 3 years 12 FG
   4 - 10 years 14 FG.
7. Attend to circulation: IV access
   ? give blood, Haemaccel or N saline.
8. Take blood for appropriate investigations.
9. Consider ‘blind’ administration of IV glucose.
10. A pulse oximeter is ideal

Cardiopulmonary resuscitation[4]

Sudden primary cardiac arrest is rare in children. Asystole or severe bradycardia is the usual rhythm at the time of arrest.

The following basic life support plan should be followed:

- Check breathing and pulse.
- Inspect oropharynx and clear any debris.
- Basic life support outline the hospital setting is 30:2 compression veutilation rates, including of inifial rescul brekes.
- Tilt head backwards, lift chin and thrust jaw forwards (the sniffing position).
- Ventilate lungs at about 20 inflations/min with bag-valve-mask or mouth to mask or mouth to mouth. An Air-viva using 8 - 10 L/min of oxygen is ideal if available.
- Intubate via mouth and secure, if necessary (must preoxygenate).
- If intubation not possible, use a needle cricothyroidotomy as an emergency.
- Start external cardiac compression if pulseless or < 60 beats/min (see Table 130.2 at p. 1351). Use two fingers for infants < 1 year and one hand for children 1 - 8 years.

Guidelines

Differences in children's airways for intubation:

- epiglottis longer and stiffer, more horizontal
- larynx more anterior → difficult to intubate ‘blind’
- cricoid ring is narrowest position → cuffed tube not required
- shorter trachea → increased risk intubating right main bronchus
- narrow airway → increased airway resistance

Endotracheal size (internal diameter in mm)

Rule:

ETT (mm) = (age in years ÷ 4) + 4 *or* the size of the child's little finger or nares

ETT length (cm) oral = (age in years ÷ 2) + 12 nasal - add 3 cm

For endotracheal size refer to Table 86.3 and for a basic schedule for CPR refer to Table 130.2 at p. 1351.

Table 86.3 Childhood intubation: endotracheal tube size and insertion distance[5]

Age	Internal diameter (mm)	Length to lip (cm)
Newborn	3.0	8.5
1 - 6 months	3.5	10
6 - 12 months	4.0	11
2 years	4.5	12
4	5.0	14
6	5.5	15
8	6.0	16
10	6.5	17
12	7.0	18
14	7.5	19
Adult	8.0	20

Paediatric Advanced Life Support

- Compression ventilation ratio of 15:2 for infants and chidren should be used in an advanced life support situation (i.e. in a hospital setting)
- Give a single shock instead of stacked shocks (single shock strategy) for ventricular fibrillation /pulseless ventricular tachycardia.
- Where the arrest is winessed by a health care professional and a manual defibrillator is available, then up to three shocks may be given (stacked shock strategy) at the first defibrillation attempt.
- Monophasic or biphasic defibrillation-first shock-2 kg, subsequent shocks-4 kg.

POISONING

Poisoning in children is a special problem in toddlers (accidental) and in adolescents (deliberate). Children of 1 - 2 years old are most prone to accidental poisoning. The most common cause of death in comatose patients is respiratory failure.

The common dangerous poisons in the past were kerosene and aspirin. Excluding household chemicals, camphor/moth balls, pesticides, insecticides and opioids, the dangerous drugs are:

- antidepressants, especially tricyclics
- antihistamines
- antihypertensives
- anxiolytics (e.g. benzodiazepines/barbiturates)
- chloral hydrate
- dishwashing power

- digoxin
- disc (button) batteries
- iron tablets
- Lomotil (diphenoxylate)
- Opiates / ‘designer’ drugs
- paracetamol/acetaminophen
- potassium tablets
- quinine/quinidine
- salicylates (e.g. aspirin)
- tricycle antidepressants

In a UK study[6] the main cause of deaths from poisoning were (in order) tricyclics, salicylates, opioids including Lomotil, barbiturates, digoxin, orphenadrine, quinine, potassium and iron.

Principles of treatment[7, 8]

The use of activated charcoal has become the key to treatment and is the ‘universal antidote’.

- Identify the poison
- Support vital functions - ABCD:
  Airway - relieve obstruction
  Breathing - ventilate with oxygen
  Circulation - treat hypotension/arrhythmias
  Dextrose - avoid severe hypoglycaemia
- Dilute the poison - give a cupful of milk or water to drink
- Remove the poison (for witnessed ingestion)
  - induce emesis (rarely employed):
    1. rub back of child's throat with spoon or spatula
    2. give syrup of ipecac (see Table 86.4) (still has limited use in the home in country areas); avoid common salt

Table 86.4 Guidelines for syrup of ipecacuanha

Dosage:
< 1 year: do not give
1 year: 15 mL
2 years: 20 mL
3 years: 25 mL
4 years: 25 mL
> 4 years: 30 mL
Follow with 100 - 200 mL water or juice
Optimal time: within 30 minutes
Mean time to emesis - 20 minutes (range 15 - 60 minutes)
If ineffective - repeat with $^{1/2}$ dose after 20 minutes
Contraindications:
> 1 hour since ingestion
rapidly acting convulsants
acid

alkali strychnine petroleum products • kerosene • petrol patient cannot sit up and hold a vomit bowl impaired consciousness • actual • anticipated

Note: The modern trend is away from emesis which is useful only for very recent ingestion (< 30 minutes) or for iron/slow potassium poisoning where concretions form in the stomach.

- gastric lavage: within 1 hour (refer to Table 86.5 for guidelines) but also limited place

Table 86.5 Guidelines for gastric lavage

Do within 60 minutes of ingestion Ideally indicated for serious poisoning when a child is already intubated
Contraindications: • stuporose or comatose • absent gag reflex (unless endotracheal tube in situ) • acid • alkali
Method: • child on left side • head of bed tilted down • insert orogastric tube (lubricated) < 2 years 12 - 14 size FG 2 - 4 14 - 18 size FG 5 - 12 18 - 22 size FG > 12 22 - 30 • check in stomach by aspiration • instil 50 - 100 mL lukewarm tap water or saline via large syringe or funnel • brief pause, then drain into bucket • repeat often until washings clear • use about 2 - 3 L • restrict the total volume to 40 mL/kg be careful of water intoxication

- gastric aspiration

- Delay absorption
  - activated charcoal (the preferred method)
    - 1 g/kg orally or via gastric tube (best) (refer to Tables 86.6 and 86.7)

Table 86.6 Drugs not absorbed by active charcoal

Acids	Cyanide
Alcohols (e.g. ethanol)	Iodines
Alkalis	Iron
Boric acid	Lithium
Bromides	Other heavy metals

Table 86.7 Drugs successfully treated by repeated doses of activated charcoal

Carbamazepine
Chlorpropamide
Cyclosporin
Dextropropoxyphene
Digoxin
Methotrexate
Phenobarbital
Phenytoin
Salicylate
Theophylline
Tricyclic antidepressants

    - Multiple dose charcoal, 5 - 10 g every 4 hours or 0.25 g/kg per hour for 12 hours, is effective
  - evaporated milk
  - aspirin and petroleum products
- Administer antidote early (see Table 86.8)

Table 86.8 Important antidotes for poisons

Poison	Antidote
Amphetamines	esmolol labetolol (hypertention)
Benzodiazepines	flumazenil
Carbon monoxide	oxygen 100% hyperbaric oxygen
Cyanide	dicobalt edetate sodium nitrite sodium thiosulphate
Digoxin	digoxin-specific antibodies magnesium sulphale

Heavy metals (e.g. Pb, As, Hg, Fe)	Dimercaprol
Iron	desferrioxamine
Isoniazid	Pyridoxine
Methanol, ethylene glycol	ethanol (ethyl alcohol)
Narcotics/opioids	naloxone (Narcan)
Organophosphates	atropine pralidoxamine (2-PAM)
Paracetamol (acetaminophen)	acetylcysteine (IV) (effective within 12 hours) consider up to 36 hours
Phenothiazines	benztropine
Potassium	Sodium bicarbonate salbutamol aerosal

- Treat any complications
    - respiratory failure
        - hypoventilation
        - apnoea
    - pulmonary aspiration of gastric contents
    - arrhythmias
    - hypotension
    - seizures
    - delayed effects (e.g. paracetamol - hepatotoxicity; tricyclics - arrhythmia)

Whole bowel irrigation

This is performed with a solution of polyethylene glycol and electrolytes (Golytely) via a nasogastric tube. It is usually limited to iron and lead, and slow-release drug preparations that don't bind to charcoal.

Investigations

- Drug levels (e.g. paracetamol, aspirin, iron)
- Blood gas analysis
- X-ray
    - chest
    - abdomen (e.g. radiopaque iron tablets)
    - skull
- ECG

Psychosocial care

The reasons for the poisoning need to be carefully evaluated and proper support and advice given.

SWALLOWED FOREIGN OBJECTS

A golden rule

The natural passage of most objects entering the stomach can be expected. Once the pylourus is traversed the FB asually continuese.

This includes:

- coins
- buttons
- sharp objects
- open safety pins
- glass (e.g. ends of thermometers)
- drawing pins

Special cases are:

- very large coins: watch carefully
- hair clips (usually cannot pass duodenum if under 7 years)

Management

- Manage conservatively.
- X-ray all children (mouth to anus espeacally chest and abdomen) on presentation the oesophagus is a concern).
- Investigate unusual gagging, coughing and retching with X-rays of the head, neck, thorax and abdomen (check nasopharynx and respiratory tract).
- Watch for passage of the FB in stool (usually 3 days). Defaecate into a conatainer.
- If not passed, order X-ray in one week.
- If a blunt FB has been stationary for 1 month without symptoms, remove at laparotomy.

Button and disc battery ingestion

If not in stomach these lespeceally lithium batteries create an emergency if is the oesophagus because electrical current generated destroys mucous membranes and perforates within 6 hours (must be removed). There also affles to the ear canal and nares.

FEBRILE CONVULSIONS

Diagnosis based on presence of fever, short duration and no clinical evidence of CNS pathology.

Features:

- The commonest cause is an URTI (e.g. the common cold or similar viral syndrome).
- About 5 per 100 incidence in children.
- Rare under 6 months and over 5 years.
- Commonest age range 9 - 20 months.
- Recurrent in up to 50% of children.
- Consider meningitis and lumbar puncture after first convulsion if less than 2 years or cause of fever not obvious.
- Epilepsy develops in about 2 - 3% of such children.

Management of the convulsion (if prolonged > 15 minutes)

- Undress the child to singlet and underpants to keep cool.
- Maintain the airway and prevent injury.
- Place patient chest down with head turned to one side.
- Oxygen 8 L/min by mask.
- Give diazepam by one of two routes:
- IV 0.2 mg/kg, undiluted or diluted (10 mg in 20 ml N saline) or rectally 0.5 mg/kg (dilute with saline or in preprepared syringe) up to 10 mg or with suppository or rectal gel.

Note: Although the IV route is preferred, the rectal route is ideal in a home or office situation; for example consider a 2 year old child (weight 12 kg) with a persistent febrile convulsion. The dose of diazepam injectable is 0.5 mg/kg so 6 mg (1.2 mL) of diazepam is diluted with isotonic saline (up to 10 mL of solution) and the nozzle of the syringe pressed gently but firmly into the anus and injected slowly. *Observe carefully for respiratory depression*.

- Rectal paracetamol 15 mg/kg statim.

Explain risk of later epilepsy is small < 3%.

Note: Intranasal midazolam given as 1 mL of drops from the vial should be considered.

MENINGITIS OR ENCEPHALITIS

Diagnosing meningitis and encephalitis requires a high level of clinical awareness and watchfulness for the infective problem that appears more serious than normal.

Vial meningitis

There ia basically a childhood infector. The most common cases are human herpes virus 6 (the cause of roseola infection) and enteroviruses.

Most case are sell limiting, but the dimical presentation can mimic baefenal mening. Lawbar puncture is important for diagnosis. Treatineat is symptomatic. The immuno compromised tequere specal manegment.

Bacterial meningitis

Bacterial meningitis is basically a childhood infection. Neonates and children aged 6 - 12 months are at the greatest risk. Meningococcal disease can take the form of either meningitis or septicaemia (meningococcaemia) or both. Most cases begin as septicaemia, usually via the nasopharynx (Figure 86.1).



Figure 86.1 Very sick infant with Pseudomonas pyocyaneus septicaemia with necrotic ulcers. *Photo courtesy Max Robinson*

Clinical presentations (typical)

Infancy:

- fever, pallor, vomiting, ± altered conscious state
- lethargic
- increasing irritability, with drowsiness
- refusing to feed, indifferent to mother
- neck stiffness (not always present)
- Kernig's sign unreliable
- may be bulging fontanelle

Children over 3 years:

- meningeal irritation more obvious (e.g. headache, fever, vomiting, neck stiffness)
- later: delirium, altered conscious state

Note: Antibiotics may mask symptoms. Suspect meningitis if fever > 3 days in reasonably well child on antibiotics.[2]

Fulminating:

- dramatic sudden-onset shock, purpura ± coma
- usually due to meningococcal septicaemia, also H. influenzae type B, *Streptococcus pneumoniae*

Treatment

First: oxygen + IV access.

If bacterial meningitis suspected, consider IV ceftriaxone or cefotaxime ± IV benzylpenicillin and admit to hospital for lumbar puncture and ongoing management. Consider dexamethasone also.

MENINGOCOCCAEMIA[10, 11]

Note: Treatment is urgent once suspected (e.g. petechial or purpuric rash on trunk and limbs).

- Take blood culture if time and facilities permit (often impractical) then
- Benzylpenicillin 60 mg/kg IV (max. 2 g), 4 hourly for 5 days or
- Ceftriaxone 50 mg/kg IV (max. 4 g) statim then 4 hourly for 5 days

A simple plan for children: benzylpenicillin[10]

- infants < 1 year: 300 mg IV or IM
- 1 - 9 years: 600 - 600 mg
- ≥ 10 years: 1200 mg

- Admit to hospital
- Continue antibiotics for 7 - 10 days

Treat contacts who:

- live in the household and < 24 months
- have kissed patient in the previous 10 days
- have attended the same daycare centre

For prophylaxis: rifampicin dose:

adult - 600 mg bd for 3 days

child < 1 month - 5 mg/kg

child > 1 month - 10 mg/kg

give bd for 2 days or (if unsuitable)

ceftriaxone 125 - 250 mg IM or ciprofloxacin 500 mg (o) as single dose Give meningococcus type C vaccine.

Practice point

In a very sick child with fever give IV antibiotics while awaiting culture.

ACUTE EPIGLOTTITIS

Acute epiglottitis due to *H. influenzae* is a life-threatening emergency in a child. A toxic febrile illness, with sudden onset of expiratory stridor, should alert one to this potentially fatal condition. A high index of suspicion of epiglottitis is always warranted in such presentations.

Differential diagnosis

The main alternative diagnosis is viral laryngotracheobronchitis (croup). There are, however, significant clinical differences.

Epiglottitis is characterised by fever, a soft voice, lack of a harsh cough, a preference to sit quietly (rather than lie down) and especially by a soft stridor with a sonorous expiratory component (Figure 86.2).


sick and toxic child
sitting quietly upright
leans forward
pale
lethargic
febrile
eyes follow
people in room
mouth open
drooling of saliva
snoring stridor
T
↑↑
P
↑↑
BP
↓
R
↑↑

Figure 86.2 Typical features of acute epiglottitis

Croup is distinguished by a harsh inspiratory stridor, a hoarse voice and brassy cough.

Other differential diagnoses include tonsillitis, infectious mononucleosis and bacterial tracheitis. The clinical features of croup and epiglottitis are compared in Table 86.9.

Table 86.9 Comparison of clinical features of croup and acute epiglottitis

Croup	Epiglottitis

Epidemiology		
Periodicity	Winter months, late autumn	Any time of the year
Age	6 months to 6 years occasionally older	6 months to 6 years
Incidence	Common	Infrequent
Clinical		
Onset	Prodrome of URTI or coryza 2 days	Rapid 2 - 6 hours
Fever	Variable, rarely above 39°C	Usually above 39°C
Toxicity	Often not anorexic	Very lethargic
	Drinking fluids	looks ill, pale, drooling
	looks like URTI	
Stridor	Loud inspiratory, increased if upset	Soft stridor, often barely audible expiratory ‘flutter’
	Harsh Brassy cough	
Pathology		
Causative organism	Viral - mostly Parainfluenza 1	Bacterial, mostly *H. influenzae* some beta-haemolytic *Streptococcus*
Site	larynx, trachea, bronchi	Epiglottis
Laboratory		Toxic FBC, increased WCC
		+ve blood culture
		+ve epiglottis culture
Treatment (refer to detailed notes)		
	Mild cases:	• airway support with nasotracheal intubation (e.g. 48 hours)
	• at home with moist air	
	moderate:	• blood culture
	• admit to hospital	
	• cool humified air	
	• oral steroids	Antibiotic: 3rd generation cephalosporin (e.g. cefotaxime or ceftriaxone)
	• observe	
	Severe:	
	• nurse in intensive care	
	• nebulised adrenaline 1:1000 solution (max. 5 mL)	Prevention
		• Hib vaccine
	• dexamethasone IV 0.2 mg/kg	
	• oral steroids	

Diagnostic tip

The child with epiglottitis usually sits still and the eyes follow you around the room because limited head movement protects the compromised airway.

Physical examination

DO NOT EXAMINE THE THROAT.

A swollen cherry-red epiglottis recognised on examination of the nasopharynx confirms the diagnosis. However, the initial diagnosis should be made on the clinical history and appearance of the child.

Direct examination using a spatula and torch should not be performed in the office but only where there are appropriate facilities for suction, endotracheal intubation and tracheostomy, as this procedure can precipitate laryngeal obstruction.

Almost all children with epiglottitis require nasotracheal intubation

Management

Transporting the patient to hospital. After ringing the hospital to warn about the emergency the practitioner should escort the child to hospital in an ambulance (with the child sitting propped up on mother's knee) prepared to perform cricothyroidotomy using a large bore cannula in the unlikely event of sudden obstruction.

The primary objective, during trans-portation, is to keep the child calm. This is enhanced by having the mother nurse the child during transfer.

If the child's condition deteriorates, administer 100% oxygen by mask. Most obstructed patients can be bagged and masked, which is sufficient to maintain oxygenation. If obstruction occurs, use the cannula to provide an airway.

Method of emergency cricothyroidotomy (last resort)

- Lay the child across your knees with neck fully extended.
- Insert a number 14 needle or angiocath through the cricothyroid membrane.

Always try to intubate once before resorting to cricothyroidotomy.

Hospital treatment

- Intubation: in theatre suck away profuse secretions and perform nasotracheal intubation
- Antibiotics[10]: cefotaxime 75 mg/kg/day IV (max. 3 g/day in 3 divided doses) *or* ceft riaxone 25 mg/kg to max. 1 g/day IV daily

Note: Continue therapy for 5 days. Early transfer to oral therapy (e.g. amoxycillin/clavulanate) is desirable.

CROUP: TREATMENT[12]

The grading system for croup is outlined in Table 86.10

Table 86.10 Grading System for croup

Croup scoup:
Grade 1 - Stridor at rest without retractions and no distress
Grade 2 - Stridor at rest with sternal and chest wall retraction
Grade 3 - marked respiratory distress indicated byirritability, pallor, cyanosis, tachycardia and exhaustion, i.e. impending airway obstraction

Grade1 croup

Mild croup (barking cough, no stridor, or stridorat rest without chest retraction, hoarse voice):

- manage at home, keep child relaxed
- consider oral steroids (e.g. 0.15 - 0.3 mg/kg dexamethasone)

A recent randomised controked trial showed no additional benefit from most supersatured air therapy when coupled with oral wrticosteroids (level II evedence)[13]

Grade 2 croup

Moderate croup (inspiratory stridor when upsetor at rest with chest wall retraction):

- admit to hospital (e.g. emergency department)
- cool humidified air
- oral steroids dexamethasone 0.15 - 0.3 mg/kg *or* prednisolone (tablets or oral solution) 1 mg/kg (2 - 3 doses) *and/or* (for children 2 or more years) budesonide 100 μg × 20 puffs or 2 mg nebulised
- nebulesed adrenaline - if poor response to steroids.
- observe for at least 4 hours

Grade 3 croup

Severe croup (inspiratory stridor at rest, use of accessory muscles, patient restless and agitated). Adrenaline is first-line therapy:

- nurse in intensive care
- nebulised adrenaline 1:1000 solution 0.5 mL/kg/dose (to max. 5 mL) (beware of possible rebound effect after 2 - 3 hours-must be observed)

Note: Can use 5 ampoules of 1:1000 solution in a nebuliser run with oxygen 8 L/min. Repeat the dose if no response at 10 - 15 minutes, do not dilute solution.

- dexamethasone 0.2 mg/kg IV *or* 0.6 mg/kg IM followed by oral steroids
- have facilities for artificial airway
- may need endotracheal intubation for 48 hours

There is no place for cough medicines or antibiotics. Steaming methods are discouraged.

BRONCHIOLITIS

- An acute viral illness usually due to RSV
- The commonest acute LRTI in infants
- Usual age 2 weeks to 9 months (up to 12 months)
- Prodromal symptoms for 48 hours (e.g. coryza, irritating cough, then 3 - 5 days of more severe symptoms)
- Wheezy breathing - often distressed
- Tachypnoea
- Hyperinflated chest.barre-shaped usually subcostal recession

Auscultation

- Widespread fine inspiratory crackles (not with asthma)
- Frequent expiratory wheezes

X-ray. Hyperinflation of lungs with depression of diaphragm

Management

Admission to hospital is usual, especially with increasing respiratory distress reflected by difficulty in feeding (particularly less than half normal over 24 hours). Dehydration ia a serious problem especially with exhaustel infant.

- Minimal handling/good nursing care
- Observation: colour, pulse, respiration, oxygen saturation (pulse oximetry)
- Oxygen: by nasal prongs to maintain P_aO_2 above 90% (preferably ≥ 93%)
- Fluids IV preferable or by nasogastric tube if unable to feed orally
- Antibiotics not indicated unless secondary bacterial infection

Note: Increased tendency to asthma later.

VERY SEVERE ASTHMA

Very severe asthma in children should be referred to an intensive care unit, stepwere action includes:

- inhaled salbutamol (6 puffs) from spacer, or
- continuous nebulised 0.5% salbutamol via mask[4]
- oxygen flow 6 L/min through the nebuliser
- IV infusion of salbutamol 5 μg/kg/min hydrocortisone 4 mg/kg IV statim then 6 hourly

Common mistakes

- Using assisted mechanical ventilation inappropriately (main indications are physical exhaustion and cardiopulmonary arrest - it can be dangerous in asthma)
- Not giving high flow oxygen
- Giving excessive fluid
- Giving submaximal bronchodilator therapy

BREATH-HOLDING ATTACKS

This is a dramatic emergency. There are two types: one is related to a tantrum (description follows) and the other is a simple faint.

Clinical features

- Age group - usually 6 months to 6 years (peak 2 - 3 years)
- Precipitating event (minor emotional or physical)
- Children emit a long loud cry, then hold their breath
- They become pale and then blue
- If severe, may result in unconsciousness or even a brief tonic - clonic fit
- Lasts 10 - 60 seconds

Management

- Reassure the parents that attacks are self-limiting, not harmful and not associated with epilepsy or intellectual disability.
- Advise parents to maintain discipline and to resist spoiling the child.
- Try to avoid incidents known to frustrate the child or to precipitate a tantrum.

Note: Important childhood emergency drugs with dosages are presented in Table 86.11.

Table 86.11 Important childhood emergency drugs (after Pitt)[14]

Drug	Route	Dose	Notes

Adrenaline 1:10 000	IV	0.1 - 0.2 mL/kg/dose	Anaphylaxis, asystole (repeat every 5 minutes until response)
Adrenaline 1:1000	Nebuliser	0.5 mL/kg/dose (max. 5 mL)	LTB* (patient must be admitted)
Aminophylline	IV slowly	5 mg/kg loading	Moderate to severe asthma
Atropine	IV	0.02 mg/kg	Bradycardia producing shock
Dextrose 50%	IV	1 mL/kg	Hypoglycaemia
Diazepam	IV PR	0.2 mg/kg 0.5 mg/kg	Seizures
Glucagon	IV or IM	0.1 mg/kg (max. 1 mg)	Hypoglycaemia
Hydrocortisone	IV	4 - 8 mg/kg	Anaphylaxis, asthma
Morphine	IV or IM	0.1 - 0.2 mg/kg	Sedation, pain relief
Paraldehyde	PR	0.3 mL/kg (dilute 1:2 in peanut oil)	Seizures
Paracetamol	O	15 - 20 mg/kg loading	Fever
Salbutamol	Nebuliser Spacer	0.3 mL/kg 6 puffs	Asthma
Salbutamol	IV	5 μg/kg	
Sodium bicarbonate 8.4%	IV	2 mL/kg	Titrate against blood gases
Soluble insulin	IV infusion	0.1 U/kg/h	Only if glucose > 14 mmol/L

*LTB = laryngotracheal bronchitis (croup)

Note: Volume resuscitation: IV fluid bolus 20 mg/kg statim of crystalloid (e.g. N saline).

ASPIRATED FOREIGN BODY

Parents or guardians may not be able to give a history of inhalation. One in eight episodes is not witnessed.[9]

Symptoms

- Choking or coughing episodes while eating nuts or similar food or while sucking a small object (e.g. plastic toy)
- Persistent coughing and wheezing (‘all that wheezes is not asthma’)
- Sudden onset of first wheezing episode in a toddler with no past history of allergy, especially after a choking bout

Signs

- Reduced or absent breath sounds over whole or part of a lung
- Wheeze

Investigations

Chest X-ray (full inspiration and full expiration) to exclude an area of collapse or obstructive hyperinflation.

Note: Normal X-rays do not absolutely exclude an FB.

Management

First aid:

- most cough out the FB, so encourage coughing
- a finger sweep helps, as do back slap, raferal chest compression and the Heimlich manoeuvre (take care with viscera)

If complete obstruction - attempt removal of the FB with forceps. If unsuccessful, perform a tracheostomy or cricothyroidotomy.

Note: Once an FB has passed through the larynx it is very rare for there to be an immediate threat to life, so referral is usually quite safe.

Note: Do not instrument the airways if the child is coping.

Bronchoscopy

Bronchoscopy is necessary in almost every child where there is a strong suggestion of an inhaled FB. It is difficult and requires an expert with appropriate facilities.

Prevention

- No child < 15 months should have ppopcorn, hard lotties, raw carrot apples
- Children < 4 year-No peanuls
- Children < 3 year - no toys with small parts

ANAPHYLAXIS

The management of airway obstruction and hypotension can be summarised as:

- oxygen 6 - 8 L/min by mask
- adrenaline 0.1 mL/kg of 1:10 000 solution IV slowly or adrenaline 0.01 mL/kg of 1:1000 IM (Repeat adrenaline obery 5 - 10 minutes as necessary - it ia the key antidote of give a continuous infusion.)
- nebulised salbutamol for bronchospasm
- colloid or crystalloid solution IV

If necessary: corticosteroid 8 - 10 mg/kg IV and antihistamine.

If persistent upper airways obstruction - try nebulised 1% adrenaline (max. 4 mL); intubation may be necessary. Admit to hospital and observe for at least 12 hours.

STATUS EPILEPTICUS

Ensure adequate oxygenation: attend to airway (e.g. Guedel tube): give oxygen. Check blood glucose.

Antiepileptic options include:[5]

- diazepam 0.2 mg/kg IV or 0.5 mg/kg per rectum
- clonazepam 0.25 mg (< 1 year), 0.5 mg (1 - 5 years), 1 mg (> 5 years)
- phenobarbitone 20 mg/kg slowly, repeated every 15 - 30 minutes if required
- phenytoin 15 mg/kg slowly over 20 - 30 minutes
- thiopentone: titrate the dose (usually 2 - 5 mg/kg)
- intranasal midazolam

If refractory (up to 60 minutes) use full anaesthetic.

Consider hyponatraemia as the cause of convulsions with meningitis.Performan ECG (look for prolonged QT interval).

DROWNING

The differences between salt-water and freshwater drowning are usually not clinically significant. If global hypoxic cerebral ischaemia and pulmonary aspiration, treat as follows:[5]

- adequate oxygenation and ventilation
- decompress stomach with nasogastric tube
- support circulation with IV infusion of colloid solution and dopamine 5 - 20 µg/kg per minute
- mannitol 0.25 - 0.5 g/kg IV if cerebral oedema
- correct electrolyte disturbances (e.g. hypokalaemia)
- give prophylactic penicillin

INTRAOSSEOUS INFUSION[3]

In an emergency situation where intravenous access in a collapsed person (especially children) is difficult, parenteral fluid can be infused into the bone marrow (an intravascular space). Intra-osseous infusion is preferred to an intravenous cutdown in children under 5 years. It is useful to practise the technique on a chicken bone.

Site of infusion

- Adults and children over 5: distal end of tibia
- Children under 5: proximal end of tibia
- The distal femur: 2 - 3 cm above condyles in midline is an alternative

Avoid growth plates, midshaft and the sternum.

Method for proximal tibia

Note: Strict asepsis is essential (skin preparation and sterile gloves).

- Inject local anaesthetic (if necessary).
- Choose 16 gauge intraosseous needle (Dieckmann modification e.g. Cook critical needle 15.5 gauge) or a 16 - 18 gauge lumbar puncture needle (less expensive).
- Hold it at right angles to the anteromedial surface of the proximal tibia about 2 cm below the tibial tuberosity (Figure 86.3). Point the needle slightly downwards, away from the joint space (< 90° to long axis).


tibial tubercule
insert midway between level of tibial tubercle and medial border of tibia, and 2 cm distal to the tibial tubercle

Figure 86.3 Intraosseous infusion

- Carefully twist the needle to penetrate the bone cortex; it enters bone marrow with a sensation of giving way.
- Remove the trocar, aspirate a small amount of marrow to ensure its position.
- Hold the needle in place with a small plaster of paris splint.
- Fluid, including blood, can be infused with a normal IV infusion - rapidly or slowly.
- The infusion rate can be markedly increased by using a pressure bag at 300 mmHg pressure.

SERIOUS GASTROENTEROLOGICAL CONDITIONS

GIT conditions that cause vomiting require careful evaluation because of potentially fatal outcomes.

Red flags

- Bile-stained vomitus mdicatesurgentrefenal to consider possible intestinal matrotation and mid-qut volualus
- Failure to passmeconium beyond 24 hours: may represent congemtal intesting atresent and stenosis, meconium ileus or Hirsch spring's disease

Gastroenteritis

This condition should not be treated lightly. Assessment of general signs such as arousal, presence of pallor, degree of weight loss, fluids in and fluids out is important (pp. 475 - 7). If concerned or in doubt, arrange hospitalisation. The social situation needs to be taken into account.

Intussusception

It is important to recognise this condition as about 50% of infants with intussusception are not diagnosed on initial presentation.[2] Characteristic features include sudden-onset pallor which persists, episodic crying and vomiting. Rectal bleeding and an abdominal mass (see p. 319 - 20) are present in only 40% of cases.

Pyloric stenosis

Pyloric stenosis, which appears from 2 weeks to 3 months of age, should be suspected with projectile vomiting, acute weight loss and alkalosis. It must not be confused with projectile vomiting from overfeeding. If in doubt, expert ultrasound examination of the pylorus will assist diagnosis(refer p. 644).

SUDDEN INFANT DEATH SYNDROME

Facts

- SIDS is the major cause of death between 1 and 12 months of age (peak incidence 4 months).
- The incidence is around 1 per 500 live births but improving.
- The causes are unknown but risk factors have been identified. Over whelming infection is a possible.
- No investigations have identified susceptible infants.
- Although SIDS can recur in a family, the risk is small.
- Association with low socio economic state.

Risk factors

- Prone sleeping position
- Smothered airways (debatable)
- Artificial feeding (possible)

- Passive smoking (before or after birth)
- Hyperthermia or excess warmth
- Extreme prematurity < 32 weeks
- Parental narcotic/cocaine abuse
- Intercurrent viral infections

Preventive advice

After baby is born.

- Place baby to sleep on its back (preferable) with no pillow or side (unless special reason for placing it on its stomach, e.g. gastro-oesophageal reflux).
- Ensure the head is uncovered.
- Breastfeeding.
- Ensure the baby is not exposed to cigarette smoking.(before and after bath)
- Ensure the baby does not get overheated (sweating around the head and neck indicates the baby is too hot).
- Bed coverings no more than adults require.
- Nothing else in cot (e.g. soft toys).

Note : monitoring offer no protector.

Reactions of bereaved parents

- May be hostility to GP, especially if recent examination
- May ‘hear’ the baby cry
- Distressing dreams
- Guilt/self-blame, especially mother
- Psychiatric morbidity

Management of SIDS

- Allow parents to see or hold baby.
- Give explanations, including reasons for coroner's involvement.
- Provide bereavement counselling.
- Early contact with counsellors and continuing support.
- Contact the SIDS support group.
- Revisit the home.
- Provide hypnotics (limited).
- Offer advice on lactation suppression.
- Remember: siblings can also experience grief reactions.
- The police and coroner must be notified - the law requires an autopsy.

Apparent life-threatening episode

ALTE, or ‘near-miss SIDS’, is defined as a ‘frightening’ encounter of apnoea, colour change or choking. At least 10% will have another episode. Management includes admission to hospital for investigation and monitoring.

Guidelines for home apnoea monitoring

- ALTE
- Subsequent siblings of SIDS victims
- Twins of SIDS victims

- Extremely premature infants

Obstructive sleep apnoea syndrome

A childhood disorder of breathing during sleep characterised by noisy, disturbed breathing and periods of apnoea. Leads to daytime sleepiness, disturbed behaviour and cognitive dysfunction. Requires referral.

REFERENCES

1. Murtagh J. The anatomy of a rural practice. Aust Fam Physician, 1981; 10: 564 - 7.
2. Hewson P, Oberklaid F. Recognition of serious illness in infants. Mod Med Aust, 1994; 37(7): 89 - 96.
3. Tibballs J. Endotracheal and intraosseous drug administration for paediatric CPR. Aust Fam Physician, 1992; 21: 1477 - 80.
4. Hewson p, Recognition of serious illness in babes. J Paediatrics child health 2000; 36:221 - 5.
5. Paxtion G, et al. Paediatric Handbook (7th edn). Melbourne: Blackwell Science, 2003: 1 - 34.
6. Fraser NC. Accidental poisoning deaths in British children 1958 - 77. Br Med J, 1980; 280: 1595.
7. Stevenson N. Drug overdose. Aust Fam Physician, 1989; 18: 1235 - 8.
8. Yuen A. Accidental poisoning in children. Patient Management, 1991; November: 39 - 45.
9. Paxton G, Munno J. *Paediatric Handbook* (7th edn). Melbourne: Blackwell Science, 2003: 26.
10. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 12). Therapeutic Guidelines Ltd, Melbourne: 2000: 55 - 247.
11. Patel MS et al. New guidelines for management and prevention of meningococcal disease in Australia. Med J Aust, 1997; 166: 598 - 601.
12. Fitzqerald DA, Kilhan Itlc. Croup : assessment and based management Med J Aust 2003; 179:372 - 7.
13. Neto 6M, kental 0 et al. A randomised controlled tral of with in the acute tratment of modernate croup. Acad Emerg Med, 2002; 9: 873 - 9.
14. Pitt R. Common paediatric emergencies. Aust Fam Physician, 1989; 18: 1228 - 34.

CHAPTER 87. ADOLESCENT HEALTH

> The feeling of apartness from others comes to most with puberty, but it is not always developed to such a degree as to make the difference between the individual and his fellows noticeable to the individual.
>
> W. SOMERSET MAUGHAM (1874 - 1965), OF HUMAN BONDAGE

ADOLESCENT HEALTH

Adolescence is the transitional period of development between relatively dependent childhood and relatively independent adulthood.[1] The time of onset and duration varies from one person to another but it is generally considered to occur between the ages of 11 and 19 years. It is a difficult period of considerable physical and mental change in which the young person is trying to cope with an inner conflict of striving for independence while still relying on adult support. There are inevitable clashes with parents, especially during the turbulent years of 13 to 16.

Adolescent patients require special understanding and caring from their doctor. They are often hesitant in approaching adult caregivers but they have a great capacity to appreciate a caring empathic approach. In this setting the family doctor has an excellent opportunity to anticipate their problems, educate them and improve their health.[2]

In recent times reference is made to the young person in the context of health policies and young people are defined as those between the ages of 10 and 24 years.[1]

ADOLESCENT DEVELOPMENT PERIODS[3]

Early adolescence (puberty: 11 - 14 years) is dominated by adjustment to the physical and psychosexual changes and by the beginnings of psychological independence from parents. Girls generally advance through this stage more rapidly than boys.

Middle adolescence (the search for independence: 14 - 17 years) is a time where boys have caught up physically and psychologically with girls, so that peer group sexual attractions and relationships are common preoccupations at this stage. It is a phase of peer group alliances, clothes, music, jargon and food and drink.[3] The average age for first sexual intercourse of both sexes is 16 years. It is a stage where intellectual knowledge and cognitive processes become quite sophisticated. Experimentation and risktaking behaviour is a feature.

Late adolescence (maturity: 17 - 19 years) is the stage of reaching maturity and leads to more self-confidence with relationships and successful rapport with parents. Thought is more abstract and reality-based.

MAJOR AREAS OF HEALTH PROBLEMS[1]

- Psychological health problems, including depression, suicide and attempted suicide
- Injuries, including sporting injuries, motor vehicle accidents and interpersonal violence
- Risk-taking behaviour, including drug abuse
- Sexual adjustment, including unsafe sexual practices and teenage pregnancy
- Eating disorders, including obesity, fast food, bulimia nervosa and anorexia nervosa (refer to Chapters 78 and 79)
- Chronic illness and disability, including survivors of inherited disorders
- Asthma, which is the leading reason for Victorian Public Hospital admissions for both sexes in the 10 - 14 year age group[1]
- Overexposure to sunlight
- Acne, which can be very distressing for adolescents. One UK study found a link between it and suicide[4]

MYTHS ABOUT THE ADOLESCENT PATIENT

The following are myths that some practitioners feel apply to the adolescent:

- different from adults in needs
- ‘superficial’ thinkers
- represent a ‘quick’ consultation
- shun personal questions
- resent invasion of space

It is important to treat them as normal human beings.

AGE AND INFORMED CONSENT

As a rule parents and physicians should not exclude children and adolescents from decision making without persuasive reasons.

On sensitive issues such as contraceptive advice, the legal concept of ‘Gillick judgement’ acts as a reasonable guideline[5]. This suggests that clinicians can identify children aged under 16 with the maturity, competence and legal capacity to consent to medical examination and treatment.

Guidelines according to age

≥ 18 - adulthood

≥ 16 - consenting age

14 - 16 - ideally involve parents, but decisions according to ‘Gillick competence’

< 14 - as above

HALLMARKS OF THE ADOLESCENT

The main hallmarks of the adolescent[2] are:

- self-consciousness
- self-awareness
- self-centredness
- lack of confidence

These basic features lead to anxieties about the body, and so many adolescents focus on their skin, body shape, weight and hair. Concerns about acne, curly hair, round shoulders and obesity are very common.

There are usually special concerns about boy - girl and same sex relationships and maybe guilt or frustration about sexual matters. Many adolescents therefore feel a lack of self-worth or have a poor body image. They are very private people, and this must be respected. While there are concerns about their identity, parental conflict, school, their peers and the world around them, there is also an innate separation anxiety.

NEEDS OF THE ADOLESCENT

Adolescents have basic needs that will allow them the optimal environmental conditions for their development:

- ‘room’ to move
- privacy and confidentiality
- security (e.g. stable home)

- acceptance by peers
- someone to ‘lean on’ (e.g. youth leader)
- special ‘heroes’
- establishment of an adult sexual role
- respect
- at least one close trustworthy friend

REBELLIOUSNESS

It is quite normal for normal parents and normal teenagers to clash and get into arguments. Adolescents are usually suspicious of and rebellious against convention and authority (parents, teachers, politicians, police and so on). This attitude tends to fade after leaving school (at around 18 years of age).

Common signs are:

- criticising and questioning parents
- putting down family members or even friends
- unusual, maybe outrageous, fashions and hairstyles
- experimenting with drugs such as nicotine and alcohol
- bravado and posturing
- unusual, often stormy, love affairs Signs of out-of-control behaviour are:[5]
- refusal to attend school
- vandalism and theft
- drug abuse
- sexual promiscuity
- eating disorders: anorexia, bulimia, severe obesity
- depression

Note: Beware of suicide if there are signs of depression.

THE CLINICAL APPROACH

Managing behavioural disorders or out-of-control behaviour demands tact and sensitivity on the part of the family doctor. It is important to interview the adolescent separately. The usual comprehensive medical history including psychosocial features is vital, particularly the family interrelationships (Table 87.1). Specifically it is appropriate to enquire about the adolescent's family relationships (parents and siblings), relationships with peers, drug taking, medical problems in the family, and parental abuse (sexual, physical, emotional or neglect).

Table 87.1 Basic clinical information (after Young[6])

History
• General history
• Drug history
• Psychological
- Personality
- introverted
- withdrawn
- anxious
- Stress

- school
- peers
- home
- Depression
- Parent - adolescent relationship
- Overprotectiveness/distant
- Separation anxiety
- Physical or sexual abuse
- Family interrelationships
- Marital conflict
- Medical problems
- Alcohol abuse

Physical examination

Investigations Keep to bare minimum

Consider the mnemonic HEADS in the history:[2]

- **H** - home
- **E** - education, employment, economic situation
- **A** - activities, affect, ambition, anxieties
- **D** - drugs, depression
- **S** - sex, stress, suicide, self-esteem

During this process it is necessary to be aware of the fundamental development tasks of adolescence, namely:[6]

- establishing identity and self-image
- emancipation from the family and self-reliance
- establishing an appropriate adult sexual role
- developing a personal moral code
- making career and vocational choices
- ego identity and self-esteem

It is necessary to conduct a physical examination and order very basic investigations if only to exclude organic disease and provide the proper basis for effective counselling. The physical examination should be conducted with sensitivity.

COUNSELLING

Counselling the adolescent involves several important principles and strategies, including:

- seeing the patient alone
- seeing parents and patient together from time to time
- confidentiality and trust
- sensitivity
- engendering the feeling that you are their doctor
- encouraging free talking and then listening carefully
- time and patience
- non-judgmental behaviour

- reassurance
- explanation
- acting as their advocate and friend
- showing genuine respect for their concerns and viewpoint

Intervention strategies on behalf of the adolescent are outlined in Table 87.2.

Table 87.2 Intervention strategies on behalf of the adolescent (after Young[6])

School
• Academic assessment (student services) • Pupil welfare co-ordinator
Family
• Simple counselling (e.g. letting go) • Family therapy
Adolescent
• direct communication about stress • be the adolescent's advocate, not the parents' • psychiatric or psychologist referral

Areas of counselling and anticipatory guidance that are most relevant are:

- emotional problems/depression
- significant loss (e.g. breakdown of ‘first love’)
- sexuality
- contraception
- guilt about masturbation or other concerns

Advice to parents

Wise parenting can be difficult because one cannot afford to be either overprotective or too distant. A successful relationship depends on good communication, which means continuing to show concern and care but being flexible and giving the adolescent ‘space’ and time.

Important management tips are:

- Treat adolescents with respect.
- Be non-judgmental.
- Stick to reasonable ground rules of behaviour (e.g. regarding alcohol, driving, language).

- Do not cling to them or show too much concern.
- Listen rather than argue.
- Listen to what they are *not* saying.
- Be flexible and consistent.
- Be available to help when requested.
- Give advice about diet and skin care.
- Talk about sex and give good advice, but only when the right opportunity arises.

Healthy distraction

Most authorities say that the best thing to keep adolescents healthy and adjusted is to be active and interested. Regular participation in sporting activities and other hobbies such as bushwalking, skiing and so on with parents or groups is an excellent way to help them cope with this important stage of their lives.

DEPRESSION, PARASUICIDE AND SUICIDE

When dealing with adolescents it is important always to be on the lookout for depression and the possibility of suicide, which is the second most common cause of death in this group. Males successfully complete suicide four times more often than females, while females attempt suicide 8 - 20 times more often than males.

The features of depression are presented in Chapter 20 but it is worth looking for the following indicators of depression:

- persistent sadness
- sleep disturbances
- eating disorders
- apathy towards friends, school and family
- sense of worthlessness
- deterioration of school performance
- crying and emotional lability
- psychosomatic symptoms
- persistent boredom, low energy
- acting out/ ↑risk taking behaviour
- preoccupation with death and dying
- suicide attempts (parasuicide)

It is important not to be afraid to enquire about thoughts of suicide as it gives teenagers a chance to unburden themselves; it is not provoking them to contemplate suicide. Parasuicide is a term coined to differentiate suicide attempts from suicide itself. Identification of risk factors as presented in Table 87.3 [9] certainly requires positive intervention in the depressed teenager. Such teenagers are especially vulnerable to a precipitating event, which can be unemployment, significant loss such as death, divorce, separation, relationship break-up, anniversary of a loss or some special celebration, additional stress or conflict and poor health.

Table 87.3 Risk factors in suicide attempts (after Birleson[9])

• Previous threats or attempts at suicide [*]
• Limited problem-solving and coping strategies
• Unsupportive families with or without marital conflict
• History of separations, psychiatric disorder, alcohol or drug abuse in family
• Family history or culture of suicide attempts
• Family disorganisation and actual neglect or abuse
• Male [*]
• Major relationship disburbances with social isolation and/or aggression
• Indicators of psychiatric disorders, especially:
- major depression [*]
- school refusal
- self-injurious behaviour
- psychosis
- alcohol/drug abuse
- personality disorder

- Availability of guns, psychotropic drugs, ropes, etc.[*]

[*] These items are the most lethal risk factors.

Depression and suicidal thoughts can respond very well to basic counselling but psychiatric referral is advisable.

The basic tasks facing the GP in managing suicidal behaviour in adolescents are summarised in Table 87.4.

Table 87.4 The 4 R tasks of managing suicidal behaviour in adolescents

Task	Strategies
Recognising the signs	• be aware and alert • ask good questions
Raising the issue	• establish rapport and listen • be direct in questioning • ask directly about suicidal thoughts • do not swear secrecy
Risk assessment(see Table 20.2 at p. 190, sad person's index)	• suicidal ideations • suicidal plan • previous attempts • precipitating events • change in daily routine • mood changes • substance abuse • supports • access to weapons/drugs
Responding	• genuine concern and support • proposed management plan • access supports • anti-depressants generally of little value • share the care • appropriate referral • assertive follow-up

MAJOR DEPRESSION

Major depression affects an estimated 8% of Australian teenagers with 40 - 80% experiencing suicidal ideation and up to 35% making a suicide attempt.[7] It is a national concern and needs to be taken very seriously in children. The incidence of depression increases markedly after puberty, especially in females. Problems adjusting to sexual orientation is a factor that should be considered in assessment. Another factor is alcohol and other substance abuse. The classic indications for depression are summarised opposite. In the history enquire about substance use/abuse, school performance, self-esteem, diet, family relationships, peer relationships, possible bullying, abuse (physical, sexual or emotional), sexuality, significant loss and energy.

Special care has to be taken with treatment, especially regarding the adverse effects of anti-depressants, especially the SSRIs. The non-pharmacological interventions that are important in all grades of depression (mild to severe) are:

- general support and education
- family therapy
- interpersonal psychotherapy
- cognitive behaviour therapy

Anti-depressant medication should be considered for moderate to severe depression (not for mild depression). No such medication is approved in Australia for the treatment of major depression in people aged under 18 while in the UK and USA fluoxetine alone is approved[7]. Fluoxetine is the treatment of choice because several randomised clinical trials support its efficacy, which is enhanced by combined psychological treatments. Tricyclic antidepressants and the SSRI, paroxetine, are currently not recommended.

Recommended treatment for moderate or severe depression[7, 8]

Combined psychosocial - CBT and family therapy + medication: fluoxetine 10 mg/day initially, increasing to 20 mg/day (if tolerated)

Continue the medication for 6 - 12 months after recovery.

Practice tips in handling adolescents

- Adolescents need support and understanding.
- They need just one good personal relationship, that one good friend they can relate to and rely on.
- They also need a good relationship with their family doctor.
- Two good questions to ask the problematic adolescent, including the obsessive compulsive:
  - Do you ever have silly thoughts?
  - Do you do silly things?
- Don't be judgmental.
- Confidentiality is of great importance to adolescents.

REFERENCES

1. Bowes G, Patton G. Adolescent health. In: Efron D. *Paediatric Handbook* (5Thedn). Melbourne: Blackwell Science, 1996: 309 - 17.
2. Bennett DL. Understanding the adolescent patient. Aust Fam Physician, 1988; 17: 345 - 6.
3. Clarke S. How to treat the adolescent. Aust Dr Weekly, 1991; 7 June: i - viii.
4. Cotterill JA, Cunliffe WJ. Suicide in dermatological patients. Br J Dermatol, 1997; 137: 246 - 50.
5. Wheeler R. Gillick or Fraser? A plea for consistency over competence in children. Editorial. BMJ, 2006; 332: 807.
6. Young D. The troubled adolescent. Aust Fam Physician, 1991; 20: 395 - 7.
7. Rey JM, Chan RT. How to treat: Depression in teenagers. Australian Doctor, 2005; 2 December: 25 - 32.
8. Dowden J. (Chair). *Therapeutic Guidelines: Psychotropic* (Version 5). Melbourne: Therapeutic Guidelines Ltd, 2003: 186 - 8.
9. Birleson P. Depression and suicide in adolescence. Aust Fam Physician, 1988; 17: 331 - 3.

CHAPTER 100. INFECTIONS IN PREGNANCY

> After autopsies I concluded that the newborn died of childbed fever, or in other words, they died from the same disease as the maternity patients … from hands contaminated with cadaverous particles brought into contact with the genitals of delivering maternity patients. If those particles are destroyed chemically, so that in examinations patients are touched by fingers but not by these particles, the disease must be reduced. To destroy cadaverous matter adhering to hands I used chlorine liquida … starting in May 1847 …
>
> When I look back upon the past, I can only dispel the sadness which falls upon me by gazing into the happy future when the infection will be banished.
>
> IGNAZ SEMMELWEIS (1818 - 1865), AUTOBIOGRAPHICAL INTRODUCTION semmelweis discovered the infectious nature of puerperal fever and how physicians transmitted it, but was not believed at the time. he died in a mental institution.

URINARY TRACT INFECTION

Urinary tract infection includes pyelonephritis, cystitis and asymptomatic cases.

Acute pyelonephritis

This infection, usually due to *Escherichia coli*, is one of the most common infective complications of pregnancy. Symptoms include fever, chills, vomiting and loin pain. Bladder symptoms such as frequency and dysuria are commonly absent. The patient should be hospitalised and usually requires intravenous antibiotic therapy and possibly rehydration.

Treatment of acute pyelonephritis

- Amoxycillin 1 g IV 6 hourly for 48 hours, then 500 mg (o) 8 hourly (if bacteria sensitive) for 14 days[1, 2]
- Alternatives: cephalosporins (e.g. cephalothin IV and cephalexin 500 mg (o))

Acute cystitis

Patients with acute cystitis typically have dysuria and frequency. Treat for 10 - 14 days.

Treatment

- Cephalexin 250 mg (o) 6 hourly[2] *or*
- Amoxycillin/potassium clavulanate (500/125 mg) (o) 12 hourly *or*
- Nitrofurantoin 50 mg (o) 6 hourly, if a betalactam antibiotic is contraindicated

  Note: Nitrofurantoin is contraindicated in the third trimester of pregnancy as it may lead to haemolytic diseases in the newborn. Cotrimoxazole and sulphonamides should be avoided. Amoxycillin is recommended only if susceptibility of the organism is proven.
- A high fluid intake should be maintained during treatment

Asymptomatic bacteriuria[1]

- 5 - 10% of pregnant asymptomatic women have positive cultures during pregnancy.
- Ideally all women should be screened for bacteriuria at their first visit.
- Less than 1% will subsequently develop bacteraemia.
- Approximately 5% of such women subsequently develop pyelonephritis during pregnancy with an increased risk of preterm labour, mid-trimester abortion and pregnancy-induced hypertension.

Treatment

Treatment is recommended according to culture sensitivities. It is preferable to delay it until the first trimester has passed.[2]

Puerperal infection

Puerperal infection is defined as a wound infection of the genital tract arising as a complication of childbirth. It especially involves the placental site in the uterus and laceration or incisions of the birth canal. Chorioamnionitis is infection of the placenta and membranes usually from normal vaginal flora e.g. Group B streptococcus, *E Coli*. It is worth recalling that Lancefield group A streptococcus infection was the outstanding cause of septic maternal death before the introduction of penicillin. The group B streptococcus has only recently been identified as a human pathogen.[3] It is carried by 15 - 20% of pregnant women in the vagina, usually without causing problems.

Maternal puerperal group B streptococcal infection usually has the following features:

- high fever > 38°C on any 2 days from days 1 to 14
- tachycardia
- endometritis - offensive or purulent discharge
- maybe abdominal pain, collapse or delirium if severe

Swab the genital tract and culture and order FBE (neutrophilia + leucocytosis).

If the woman is febrile but not clinically ill, treat with amoxicillin + clavulanate 875/125 (o) bd for 5 - 7 days. If septic treat with IV amoxicillin + gentamicin + metronidazole.

About 1 - 4% of neonates per 1000 live births develop group B strep sepsis with a high mortality rate, up to 50%.[3] The onset can be early or late in the neonate with a severe clinical presentation of rapid deterioration with septic shock. High-risk situations are premature rupture of the membranes, early labour and maternal intrapartum fever and also vaginal delivery. Routine group B streptococcus screening is impractical; therefore prophylactic treatment for these situations is IV penicillin or amoxicillin during labour.

Vaginal candidiasis

Candida (thrush) is common in pregnancy since pregnancy is a predisposing factor to the growth of the fungus. Treatment follows conventional lines with topical creams and vaginal tablets as described in pp. 1020 - 1. Clotrimazole is a first-line treatment. Oral medication with fluconazole or itraconazole should be avoided.

TRANSMISSIBLE VIRAL INFECTIONS

Rubella

Facts:

- Fewer than 5% of women are not immune (IgG negative).
- Rubella IgM indicates recent infection and a real risk if pregnant.
- Reinfection can occur even after vaccination so test all pregnant contacts.
- Congenital rubella features on pp. 903 - 4.
- Effects on foetus:[4]
  - < 8 weeks gestation - up to 85% infected with all clinically affected
  - < 12 weeks gestation - 50 - 80% infected with 65 - 85% clinically affected
  - 13 - 16 weeks - 30% infected with 1:3 sensorineural deafness
  - 16 - 19 weeks - 10% infected - clinical effects rare, possible deafness
  - > 19 weeks - no apparent risk[3]

Diagnosis: fourfold rise in IgG titres or rubella specific IgM antibody

Vaccination:

- routine vaccination advisable - gives 95% protection[5]
- do not vaccinate during or within 3 months pregnancy

- if inadvertent vaccination in early pregnancy - negligible risk to foetus[4]
- any IgG negative woman should be offered immunisation in the puerperium

Varicella/chickenpox

Facts:

- Greatest risk if infection is in first trimester and very late pregnancy.
- Does not apply to herpes zoster (shingles).
- Establish diagnosis by IgG antibody test.
- Foetal varicella syndrome is rare - includes limb abnormalities, microcephaly, optic atrophy, mental retardation, IUGR.
- Appears to occur in 3% of pregnancies where mother contracts varicella.[6]

Contacts: if no history clinical varicella do IgG antibody test in mother. If seronegative give varicella zoster immunoglobulin (VZ-Ig) immediately (within 3 - 4 days of contact)

Maternal infection early pregnancy: greatest risk < 20 weeks gestation. Give course antiviral e.g. acyclovir, valaciclovir; consider ultrasound.

Maternal infection late pregnancy[7]: greatest risk is 5 days before delivery up to 4 weeks after delivery - 30% foetal mortality if infected. Consider VZ-Ig for baby if < 7 days before delivery and up to 4 weeks after. Isolate mother from baby until not contagious.

Parvovirus B19

Facts:

- This virus causes slapped face syndrome (pp. 904 - 5) and can be mistaken for rubella.
- Non-immune at risk.
- Risk of transplacental infection exists throughout pregnancy.
- Screen for immunity with HPV B19 IgG antibodies (reassure if positive).
- Screen for infection with acute and convalescent sera for IgM antibodies.
- Miscarriage rate is 4% < 20 weeks.[8]
- Foetal parvovirus syndrome is anaemia - hydrops fetalis with cardiac failure - possibly death.

If infection occurs during pregnancy, refer for foetal monitoring by ultrasound - if hydrops, consider early blood transfusion.

Cytomegalovirus

Facts:

- CMV is the commonest cause of intrauterine infection.
- Commonest viral cause of birth defects.
- Primary infection is usually asymptomatic.
- Foetus most vulnerable in early pregnancy.
- Specific IgM antibody implies acute infection.
- The risk from maternal infection is about 40%.
- Incidence of congenital CMV is about 0.3%.
- Effects variable and unpredictable - can be severe or mild (refer pp. 258 - 9).
- Up to 30% of CMV-affected infants have mental retardation.
- Up to 50% are restricted to hearing loss.
- There is no therapy or preventive strategy to alter the disease.

If foetal infection is likely or suspected consider referral and amniocentesis.

Hepatitis B

Facts:[8]

- Women should be screened for hepatitis B during pregnancy.
- +ve HBsAg indicates acute infection.
- +ve anti-HBs indicates recovery and immunity.
- +ve HBeAg indicates high infectivity but low transmission in utero.
- Vertical transmission usually during labour is the concern.
- Higher risk (90%) if mother HBeAg seropositive.
- Infected infants - 90% risk of becoming chronic carriers with liver disease.

Prevention by immunisation:

At delivery or ASAP give newborn babies of carrier mothers both hepatitis B vaccine and immunoglobulin (HBIg). This gives efficacy of about 90 - 95%. Follow up with booster doses of vaccine at 2, 4 and 6 (or 12) months.

Hepatitis C

HCV screening recommended at first antenatal visit for those from significant risk group. If positive transmission rate to foetus is 5% and much higher if maternal infection during pregnancy.[3] Evidence unclear where HCV transmission occurs during breastfeeding. Screen infants at risk at 12 months and treat positive cases accordingly under specialist care.

SEXUALLY TRANSMITTED INFECTIONS

Genital herpes

Both primary genital herpes (in particular) and recurrent herpes poses a risk to the neonate. The risk from primary infection is greatest if it occurs after 28 weeks gestation. Risk factors for intrapartum infection include primary infection, multiple lesions, premature rupture of the membranes, premature labour. The main problem is vertical transmission during labour.

Management[9]

- Perform cervical swab for HSV infection.
- Consider prophylactic antiviral, e.g. acyclovir, for mother from 38 weeks until time of delivery - to try and prevent recurrent herpes in late pregnancy.
- Arrange caesarean section if:
  - active lesions present (cervix/vulva) at time of delivery or within preceding 4 days
  - membranes ruptured > 4 hours
- If vaginal delivery give acyclovir to the neonate (check with neonatal paediatrician).

Genital warts

Although estimates indicate that up to 70% of the world's adult population carry one or more strains of human papilloma virus, the risk of transmission of the virus from the maternal genital tract to the foetus is very low.[3] Caesarean section is rarely indicated for obstruction from or the presence of gross numbers of cervical warts.

Syphilis

Facts:[3]

- Incidence 2/1000.
- Usually transmitted in second trimester.
- May cause foetal death; congenital infection with handicap.
- Tests - VDRL, TPHA, FTA-Abs.

Treatment:

- Acquired early syphilis including latent < 12 months: benzathine penicillin 1.8 g IM as single dose.
- Late latent syphilis (incubation period > 12 months): benzathine penicillin 1.8 g IM once each week for 3 doses.
- Treat the partner.

HIV

The foetal infection rate from a HIV positive mother is about 15 - 25%.[10]

If screening detects a HIV positive mother both she and her newborn infant require antiretroviral therapy. Consult with a specialist experienced in treating HIV infection.

Breastfeeding is inadvisable because it doubles the risk of vertical transmission.

The risk of transmission can be reduced to < 5%:[10, 11]

- by treatment with zidovudine prescribed for the mother antenatally and during labour and to the neonate for the first 6 weeks postpartum
- by elective caesarean, and
- by avoiding breastfeeding

Chlamydia/gonorrhoea

Both gonorrhoea and *Chlamydia* urethritis can transmit infection to the foetus, causing neonatal conjunctivitis which develops in the first 2 weeks of life. *Chlamydia* can also cause neonatal pulmonary infection such as pneumonia usually appearing at 2 or 3 months of age. PCR testing of maternal urine as appropriate and eye swabs from the neonate are advisable. Treat according to antibiotic guidelines.

INFECTIONS TRANSMITTED FROM CONTAMINATED FOOD

Toxoplasmosis

Facts:

- Incidence in pregnancy in Australia less than in Europe.
- Caused by *Toxoplasma gondii* - a parasite found in cat's (and ? possum's) faeces and raw meat.
- Acquired by close contact with infected cats or eating uncooked or undercooked meat.
- About 2/1000 maternal infection rate with about 30% passed to foetus.
- Most maternal cases are asymptomatic.
- Greater risk to foetus before 20 weeks gestation.
- Congenital toxoplasmosis rare, but may result in various CNS and eye abnormalities. Refer to p. 259.
- May cause IUGR, miscarriage, stillbirth.
- Overall up to 90% of infected babies escape long-term damage.[12]
- Diagnosis by serological tests.

Treatment in pregnant patients is with spiramycin.

Listeria

Facts:

- Infection caused by a Gram-positive bacterium.
- High risk foetal mortality (30 - 50%) if contracted in pregnancy.
- May cause recurrent abortion, stillbirth, premature labour.
- Maternal symptoms - fever, headache, myalgia, rigors, vomiting, diarrhoea, abdominal pain. Suspect if persistent high fever/flu-like illness. Refer for expert advice.

- Non-pregnant women are usually asymptomatic.
- Transmission to foetus mainly in later pregnancy with a high risk.
- Infection is usually by infected food, e.g. soft cheese, pate, milk esp. unpasteurised, raw seafoods, smoked seafoods, coleslaw.
- Investigations - mainly blood culture, also amniotic fluid, cervical swab culture.
- Newborn may acquire pneumonia, liver abscesses, meningitis, sepsis.
- Maternal listeriosis is treated with amoxicillin.

Practice tips

- Well-proven transplacental vertically transmissible pathogens include cytomegalovirus, rubella, syphilis, toxoplasmosis and varicella.
- The best serological evidence of recent infection is IgG seroconversion so the first specimen should be collected ASAP after the onset of symptoms.
- Many relevant viral infections such as CMV, toxoplasmosis, parvovirus, and rubella can be asymptomatic.
- No treatment or vaccine is available for specific infections such as CMV and parvovirus.
- Patients with a history of contact or clinical infection with congenitopathic infections require good counselling, education and support - their anxiety levels can be extreme.

Preventive advice for toxoplasmosis and listeriosis

- Protection is afforded by good personal and food hygiene.
- Avoid eating the above products.
- Only eat well-cooked meat/poultry - thoroughly cook all food of animal origin.
- Carefully wash soil from raw vegetables and fruit.
- Reheat leftover foods and ready to eat food until steaming hot.
- Always thoroughly clean utensils and food surfaces after preparing uncooked food.
- Pregnant women should get another person to clean cat litter boxes daily and wear disposable gloves for handling soil likely to be contaminated with cat's faeces.
- Carefully wash hands after gardening or handling raw meats.

REFERENCES

1. Humphrey M. Common problems in an otherwise normal pregnancy. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996; 117 - 20.
2. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 12). Melbourne: Therapeutic Guidelines Ltd, 2003: 196 - 8.
3. Humphrey M. *The Obstetrics Manual* (revised edition). Sydney. McGraw-Hill, 1999: 74 - 81.
4. Goh J, Flynn M. *Examination Obstetrics & Gynaecology* (2nd edition). Marrickville: Elsevier 2004: 140 - 7.
5. Jones G et al. Congenital rubella in Great Britain. Health Trends, 1990; 22: 73 - 6.
6. Gilbert GL. Chickenpox during pregnancy. BMJ 1993; 306;1079-80.
7. Sterner G et al. Varicella-zoster infections in late pregnancy. Scand J Infect Dis, 1990; 71: Suppl 30.
8. Gilbert G. Infections in pregnant women. Med J Aust, 2002; 176: 229 - 36.
9. Braig S et al. Acyclovir prophylaxis in late pregnancy prevents recurrences of genital herpes and viral shedding. Eur J Obs Gyn Reprod Biol, 2001; 96: 55 - 8.
10. Doherty R. Preventing transmission of HIV from mothers to babies in Australia, Med J Aust, 2001; 174: 433.
11. Connor EM et al. Reduction of maternal-infant transmission of HIV type 1 with zidovudine treatment. New Eng J Med, 1994; 331: 1173 - 80.
12. Khot A, Polmear A. *Practical General Practice* (4th edn). London: Butterworth Heinemann, 2003: 262.

CHAPTER 101. HIGH-RISK PREGNANCY

Grace the wyffe of William Baxter, beinge aboute three weeks before her tyme, was brought in bed the first day of December of two children, their bellies were growne and joyned together, from their breastes to their navells, and their faces were together.

JOHN RICHARDSON, GIVING DETAILS OF SIAMESE TWINS, BORN in 1655

DEFINITION[1]

A high-risk pregnancy is one in which the fetus is at increased risk of stillbirth, neonatal morbidity or death, and/or the expectant mother is at increased risk for morbidity or mortality.

High-risk pregnancies may be predicted before conception in some women, especially those with serious medical problems and a poor obstetric history. Others at high risk can be identified at the first antenatal visit and others develop during the course of pregnancy. The first antenatal visit is the most important visit and demands time and care to make an accurate assessment.

The primary goal in obstetric care is to avoid maternal and perinatal morbidity and mortality (in particular) wherever possible through appropriate risk management including proactive strategies.

MATERNAL MORTALITY

The WHO definition of maternal mortality is the death of a woman during pregnancy, childbirth or in the 42 days of the puerperium, irrespective of duration or site of the pregnancy, from any cause related to or aggravated by the pregnancy or its management. This excludes deaths from assisted reproductive technology where pregnancy has not resulted, but includes incidental causes of deaths. In Australia the classification comprises direct maternal deaths, indirect obstetric deaths (where diseases such as diabetes, cardiovascular and kidney are complicated by pregnancy) and incidental deaths such as road accidents, suicide and malignancy.[2]

The maternal mortality ratio is the number of deaths per 100 000 confinements. In first world countries it is approximately 10. The latest triennium statistics for Australia was 8 deaths/100 000 confinements (c.f. Africa approx. 900).[2] The main causes of direct maternal deaths in Australia are (in order):

- haemorrhage
- pulmonary embolism
- amniotic fluid embolism
- eclampsia/pre-eclampsia
- ectopic pregnancy
- septic shock

Contributing factors to those who died from haemorrhage are delays in diagnosis, treatment to arrest the haemorrhage and giving blood transfusion. Intracerebral haemorrhage was the main cause of death in pre-eclampsia.

Some Australian obstetric statistics for 2003:

- average age of all mothers was 29.5 years
- spontaneous vaginal births - 60.3%
- caesarean section rate - 28.5%
- instrumental delivery rate - 10.7%
- multiple pregnancies - 1.7%

PERINATAL MORTALITY

The perinatal mortality is the total number of deaths of children within 28 days of birth (early neonatal deaths) plus fetal deaths at a minimum gestation period of 20 weeks or a minimum fetal weight of 400 g expressed per 1 000 births.[3]

In Australia the average current perinatal mortality rate is approximately 7 to 8/1 000. The average for the 10 year period 1994 - 2003 in NSW[4] was 8.2 with a steady decrease over the period. The major factors associated with perinatal mortality are very premature birth, congenital abnormalities and birth asphyxia. A review of perinatal deaths occurring in 2003 in Australia found that 30.9% of perinatal deaths (or 45.7% of stillbirths) were unexplained antepartum deaths. Common causes included spontaneous preterm labour (< 37 weeks gestation) - 16%, congenital abnormality - 16%, antepartum haemorrhage - 7.4%. For neonatal deaths, extreme prematurity was the most common cause of death (44.8%), followed by congenital abnormality (19.3%) and neurological disease (12.5%).[4]

ACCURATE DETERMINATION OF EDD

It is vital to determine the expected due date of delivery (EDD) based on the exact time of the last normal menstrual period (sometimes misleading), the fundal height, the time of first fetal movements and, if in doubt, ultrasound assessment (see Figure 99.1).The earlier ultrasound is performed after 6 - 7 weeks of gestation, the more accurate the determination.

HIGH-RISK OBSTETRIC PATIENTS

Guidelines for high-risk pregnancy are presented in Table 101.1. Recognition of these high-risk pregnancies is important for the family doctor involved in shared care. Common categories that require special surveillance are:

- elderly primigravida (aged > 35 years)
- grand multigravida (fifth or greater pregnancy, especially if > 35 years)
- those with a poor obstetric history
- previous caesarean section
- severely disadvantaged social problem (e.g. sole teenage parent with drug problem)
- hypertension ± chronic kidney disease
- obesity
- short stature
- diabetes mellitus
- prolonged infertility or essential drug or hormone treatment
- heavy smoking or alcohol intake

The onset during pregnancy of the following also requires surveillance:

- little or no weight gain during the first half of the pregnancy
- complications such as pre-eclampsia, multiple pregnancies or those with antepartum haemorrhage
- abnormal presentation
- abnormal fetal growth

Table 101.1 Guidelines for specialist obstetric consultation

	Major risk factors Obstetric consultation mandatory	**Other risk factors Consultation should be considered**
Past problematic obstetric history	• Previous caesarean section	Multiple spontaneous or elective abortions
	• Incompetent cervical os	
	• 2nd trimester spontaneous abortion	• Premature delivery
	Rhesus/other blood group incompatibility	Previous stillbirth

	• Thromboembolic disease • Premature labour	• Neonatal death • Grand multigravida
Problems related to current pregnancy	• Pre-eclampsia • Rhesus/other blood group incompatibility • Significant vaginal bleeding • Placenta praevia • Postmaturity (especially if > 41 weeks • Multiple pregnancy • Polyhydramnios • Need for amniocentesis: - genetic concerns - abnormal AFP - other	• Recurrent urinary infections • Abnormal uterine growth • Inadequate maternal weight gain • Hypertension
	General factors	• Age > 35 years; < 18 years • Obesity > 110 kg • Prepregnancy weight < 45 kg • Psychosocial problems • Short stature < 152 cm • Smoking > 10/day
Maternal disorders	• Diabetes mellitus • SLE • Sickle cell anaemia or other haemoglobinopathy • Thrombophilia	• Anaemia: Hb < 100 g/L • Cardiovascular disease • Chronic kidney disease • Alcohol or drug abuse • Genital herpes
Perinatal problems	• Premature labour • Postmaturity (> 41 weeks) • Disproportion • Malpresentation • Placental insufficiency	• Non-vertex presentation (at term) • Fetal arrhythmia • Membranes ruptured > 18 h
Inadequate antenatal care		• Late presentation (after 20 weeks) • No antenatal care • Failed or poor attendance

Diabetes is always a concern and may manifest in pregnancy. Glucose screening testing is now performed routinely in many hospitals at 26 - 28 weeks gestation.

HYPERTENSIVE DISORDERS IN PREGNANCY[5, 6]

Hypertensive disorders complicate about 10% of all pregnancies. Pregnancy may induce hypertension in previously normotensive women or may aggravate pre-existing hypertension. Pre-eclampsia, which in fact complicates 2 - 8% of pregnancies, can occur at any time in the second half of pregnancy.

Types

Hypertension without proteinuria

Pre-eclampsia - hypertension + proteinuria

Pre-eclampsia - hypertension + oedema + proteinuria

Eclampsia - hypertension + convulsions

A classification of hypertensive disorders in pregnancy:

- *Pregnancy-induced hypertension*
  - SBP > 140 mmHg and DBP > 90 mmHg, occurring for first time after 20th week of pregnancy and regressing postpartum *or*
  - Rise in SBP > 25 mmHg or DBP > 15 mmHg from readings before pregnancy or in first trimester
- *Mild pre-eclampsia* BP up to 170/110 mmHg in absence of associated features (see following)
- *Severe pre-eclampsia* BP > 170/110 mmHg and/or associated features, such as kidney impairment, thrombocytopenia, abnormal liver transaminase levels, persistent headache, epigastric tenderness or fetal compromise
- *Essential (coincidental) hypertension* Chronic underlying hypertension occurring before the onset of pregnancy or persisting postpartum
- *Pregnancy-aggravated hypertension* Underlying hypertension worsened by pregnancy

Risk factors

The following are risk factors for pregnancy-induced hypertension:

- nulliparity/primigravida
- family history of hypertension/pre-eclampsia
- chronic essential hypertension
- diabetes complicating pregnancy
- obesity
- donor sperm or oocyte pregnancy
- multiple pregnancy
- hydatidiform mole
- hydrops fetalis
- hydramnios
- kidney disease
- autoimmune disease, e.g. SLE

Clinical features of superimposed pre-eclampsia include hypertension, excessive weight gain, generalised oedema and proteinuria (urinary protein > 0.3 g/24 hours). Late symptoms include headache related to severe hypertension), epigastric pain and visual disturbances.

Risks of severe pre-eclampsia/hypertension[6]

Maternal risks (poor control)

- Kidney failure

- Cerebrovascular accident
- Cardiac failure
- Coagulation failure

Risks to baby

- Hypoxia
- Placental separation
- Premature delivery

Management

The optimal treatment is delivery, and induction of labour needs to be timed appropriately - based on parameters such as the BP level and the development of proteinuria. The BP level must be kept below 160/100 mmHg because at this level intrauterine fetal death is likely to occur and there is a risk of maternal stroke.

Antihypertensive drugs[5]

Contraindicated drugs are ACE inhibitors and diuretics. There is no place for the use of diuretics alone unless cardiac failure is present.

Commonly used medications:

- beta-blockers, e.g. labetalol, oxprenolol and atenolol (used under close supervision and after 20 weeks gestation)
- methyldopa: good for sustained BP control
- nifedipine

Labetalol, hydralazine and diazoxide are useful for rapid control of BP in hypertensive crises, e.g. hydralazine 5 mg IV bolus every 20 - 30 minutes or continuous infusion.

Guidelines for urgent referral/admission to hospital

Maternal factors

- Progressing pre-eclampsia including development of proteinuria
- Inability to control BP
- Deteriorating liver, blood (platelets), kidney function
- Neurological symptoms and signs, e.g. headache, drowsy and confused, twitching, rolling eyes, vomiting, visual disturbances, hyperreflexia, clonus, i.e. imminent eclampsia

Fetal factors

- Abnormal cardiotocograph (CTG) indicating fetal distress
- Intrauterine growth retardation

Refer to eclampsia ward under specialist care.

Treatment of severe pre-eclampsia: prevention of convulsions

- Control BP : use IV hydralazine or diazoxide. don't suppress to < 140/80
- Magnesium sulphate 4 g IV (given over 10. 15 minutes) followed by an infusion 1 g/hour for 24 hours (if normal kidney function)[7, 8]
- Corticosteroid therapy IM for fetal lung maturity if gestation ≤ 34 weeks.

Monitor fetus and maternal BP, urine output, urine protein, coagulation profile

Note: The best treatment is termination of pregnancy with early delivery - by caesarean section or vaginal delivery if favourable circumstances. There is a risk of fitting at delivery and after delivery (usually < 24 hours).

Treatment of convulsion[8]

- Secure airway and ventilate if necessary - give intranasal oxygen.
- Magnesium sulphate as above; if further fit use bolus of 2 g $MgSO_4$.[7]
- Strict fluid balance required.
- Monitor and treat any coagulopathy, pulmonary oedema.
- Avoid ergometrine in the third stage.

Note: Be aware of the risk of more fits over the next 24 - 48 hours.

HELLP syndrome

Haemolysis Elevated Liver enzymes Low Platelets. A severe form of pre-eclampsia. Treat as for severe pre-eclampsia.

MEDICAL CONDITIONS IN PREGNANCY

Anaemias

During the course of a normal pregnancy the haemoglobin level should remain above 110 g/L. Anaemia is defined as a haemoglobin < 110 g/L. Levels below this, particularly less than 100 g/L, require investigation. Total iron demands during pregnancy are 725 mg, especially during the third trimester.

Risks for anaemia include poor diet, multiple pregnancy and starting with anaemia.

Important types of anaemia in pregnancy:

- iron deficiency (approximately 50%)
- megaloblastic anaemia (usually due to folic acid deficiency)
- thalassaemia (most commonly β-thalassaemia)

Management

- If anaemia is found, measure serum ferritin level, MCV, red cell folate and serum vitamin B_{12} as indicated
- Treatment is according to cause:
  - iron deficiency: ferrous sulphate 0.9 g (o) daily iron infusion may be required
  - megaloblastic anaemia: folic acid 5 mg (o) bd

Note: A blood transfusion is rarely required.

Diabetes mellitus

Diabetes mellitus, especially type 1, in pregnancy requires meticulous management skills to minimise the morbidity and perinatal wastage associated with diabetes. Optimal management demands a team approach from the compliant patient, GP, obstetrician, diabetic physician, diabetic nurse - educator and dietician.

Effects on the fetus

Large for dates (macrosomia), ↑ fetal abnormalities (neural tube, cardiac, kidney, vertebral, etc. defects), hypoxia and intrauterine death (FDIU), miscarriage, malpresentation, IUGR, preterm delivery.

Postnatal

Early hypoglycaemia, respiratory distress syndrome, jaundice.[9]

Effects on mother

Increased risk pre-eclampsia, diabetic ketoacidosis, polyhydramios, intercurrent infection, psychological effects, first trimester miscarriage, obstructed labour (↑ shoulder dystocia), placental abruption, caesarean section.[9]

Management

Pre-pregnancy

Education and counselling, review of results, dietary compliance, medication especially insulin, advice as on pp. 1037 - 8, including folate.

Aim for diabetic control: fasting blood sugar 4 - 7 mmol/L: HbA1c < 7%

During pregnancy

- Early referral to a diabetic physician and obstetrician is best management.
- Assess basic parameters.blood sugar, HbA1c, BP, kidney function inc. 24-hour urinary protein, urine, electrolytes.
- Reassess insulin requirements as pregnancy advances (↑ × 2-3).
- Screen for fetal morphology and growth: ultrasound at 18 weeks then 4 weekly and as required; cardiotocography as required. usually weekly until delivery.
- Aim to deliver at term at latest:
    - vaginally if optimal control
    - caesarean section if large fetus (> 90th weight percentile or 4 000 g) or evidence fetal distress or breech presentation
- In labour monitor fetus with CTG and keep maternal blood glucose levels normal with intermittent insulin injections.
- Cease insulin infusion and ↓ insulin to prepregnancy regimen immediately after delivery; organise contraception; avoid oral hypoglycaemics during breastfeeding.

Gestational diabetes

Gestational diabetes is the onset or initial recognition of abnormal glucose tolerance during pregnancy. If suspected a diagnostic oral glucose (75 g) tolerance test is indicated.

Diagnosis: fasting blood glucose > 5.5 mmol/L *or* 2-hour level > 8.0 mmol/L

The same risks to mother and fetus such as macrosomia, FDIU, congenital malformations, hyaline membrane disease and neonatal hypoglycaemia as for pre-existing diabetes are applicable.[10]

Note: Glycosuria in pregnancy is unhelpful for screening because it is common in pregnancy and lacks specificity.

Principles of management

These are the same as for pre-existing diabetes (above). If insulin is required antenatally, cease insulin postpartum.

Aim to deliver vaginally at term at the latest, unless earlier delivery indicated by obstetric complications.

Post delivery

Follow up GTT at 6 weeks and then every 5 years. Gestational diabetes is likely in subsequent pregnancies and there is a 30% risk of developing diabetes in later life - even < 10 years.

Thyroid disorders

Hypothyroidism in pregnancy is uncommon and usually mild. It is associated with infertility. It is associated with a higher rate of fetal loss, miscarriage, fetal abnormalities and IUGR. TFTs should be checked at first presentation if past history is relevant and at 36 weeks. Optimal dosage of thyroxine replacement should be monitored by thyroid function tests. Normal maintenance dosage is usually increased to meet increased needs in pregnancy.

Hyperthyroidism in pregnancy is usually caused by Graves’ disease, but the outcomes are generally good with most remaining euthyroid throughout pregnancy.

If untreated there is an increased risk of maternal and fetal complications. Propylthiouracil is usually the preferred agent. Patients should be referred for specialist management.

Cardiac disorders

Key points:

- Maternal cardiac disease is not common in pregnancy.
- The highest risk of maternal mortality is where pulmonary blood flow cannot be increased, e.g. pulmonary hypertension, Eisenmenger's syndrome.
- Cardiac output rises in pregnancy.
- Syncope and dyspnoea may be a pointer to a cardiac disorder.
- Murmurs are common - 90% are physiological. Refer if in doubt.
- Patients with an existing cardiac condition require combined specialist cardiologist/ obstetrician care.
- Antibiotic prophylaxis is important for those with structural cardiac problems, e.g. valvular problems, most congenital malformations.
- Aim for spontaneous vaginal delivery at term if possible.
- As a rule avoid lithotomy, ergometrine, sympathomimetic drugs.
- Watch fluid balance, especially after delivery.
- Administer oxygen during labour and early postpartum.

Kidney disease

Key points:[11]

- Outcome in chronic kidney disease depends on severity of prepregnancy kidney function and blood pressure. Good control of hypertension is essential.
- Pre-eclampsia is more common, requiring cautious monitoring.
- Mild kidney failure usually has very good outcomes.
- ↑ maternal and fetal complications in moderate failure (s. creatinine 0.125. 0.25 mmol/L) and severe kidney failure (> 0.25 mmol/L).
- Pregnancy uncommon in dialysis patients. If so, there can be many problems requiring increased use of dialysis.
- Refer these patients and admit if hypertension becomes poorly controlled.
- Aim for delivery at 38.40 weeks at the latest. Delivery is usually required before this time.
- In kidney transplant patients the risk of early miscarriage is ↑. Surviving pregnancies have > 90% success. ↑ risk hypertension and preeclampsia but low chance of kidney rejection. Pelvic site of transplant is not a problem.

Systemic lupus erythematosus

Key points:

- Pregnancy does not seem to cause exacerbations of SLE.
- SLE can adversely affect pregnancy according to disease severity.
- Increased incidence of spontaneous abortions and stillbirths.related to lupus anticoagulant and anticardiolipin antibodies.
- ↑ risk pre-eclampsia, prematurity, IUGR, perinatal mortality.
- Neonatal lupus syndrome includes blooddisorders and cardiac abnor malities in the neonate.
- Increased maternal morbidity.kidney complications, pre-eclampsia.

Management

- Careful preconception counselling - plan pregnancy with remission phase.
- Refer for review of drugs.
- Many investigations required under specialist supervision including lupus antibodies, AAPT, FBE, kidney function, ultrasound scans.

- Corticosteroids, e.g. prednisone are mainstay of therapy.
- Low dose aspirin (100 mg daily) if anticardiolipin antibodies present.
- Low molecular weight heparin may be used as alternative to aspirin and in presence of prolonged APTT.
- Timing of delivery according to assessment of progress and status of fetus.

Thrombocytopenia in pregnancy

The two most common causes of significant thrombocytopenia (CTP) in an otherwise normal pregnancy blood film are gestational thrombocytopenia and immune thrombocytopenia. Other causes that need to be considered are SLE, anti-phospholipid syndrome (APS), drug induced thrombocytopenia and HIV infection.

Gestational thrombocytopenia

In up to 4% of pregnancies, mild CTP in the range 75 - 150/nL develops, usually in the third trimester. Because of the increasing hazard of epidural anaesthesia in platelet counts under 75/nL, a 2-week course of prednisolone is often prescribed at 37 - 38 weeks gestation, aiming for a platelet count in excess of 100/nL at the time of delivery. It is prone to occur with subsequent pregnancies.

Immune TCP

Although less common than gestational ITP, it is clinically more significant since it is typically severe and arises earlier in pregnancy. ANA factor and other antibody studies are necessary to exclude SLE and APS. Platelet-specific antibodies are found in at least 50% of ITP patients. Life-threatening cases (< 10 nL) require hospitalisation and corticosteroid therapy or IV immunoglobulin (if unresponsive).

Fetal ITP

Fetal ITP results from the transplacental passage of maternal IgG anti-platelet antibody into the fetal circulation. The risk of severe fetal ITP is increased in severe ITP. The platelet count should be followed carefully postpartum. If severely reduced, IV immunoglobulin and even platelet transfusion may be required (even in utero if necessary).

Jaundice specific to pregnancy

Specific types of jaundice related to pregnancy are rare. Viral hepatitis accounts for 40% of all cases of jaundice during pregnancy. Severe pre-eclampsia, eclampsia and hyperemesis gravidarum may cause hepatic damage with jaundice. Two interesting specific conditions are cholestatic jaundice of pregnancy and acute fatty liver of pregnancy - the latter condition is now very rare.

Cholestasis of pregnancy

This condition is due to an oestrogen sensitivity. The symptoms, which are mild, include low grade jaundice and pruritus during the latter half of pregnancy. It can be associated with fetal distress, fetal death and preterm delivery. Refer for monitoring of fetus with a view to deliver if problematic. Perform LFTs and consider vitamin K supplements. Drug treatment is generally unrewarding.

The condition clears up rapidly after delivery, but it often recurs in future pregnancies and if the patient is prescribed oral contraceptives, which are contraindicated.

Acute fatty liver of pregnancy

This is a serious condition of unknown aetiology and may follow the administration of hepatotoxic agents, especially in the more debilitated patient. Acute fatty liver presents in the last trimester with symptoms of fulminant hepatitis - jaundice, vomiting, abdominal pain, headache and possibly coma. It has a high mortality (about 50%) and necessitates urgent termination of pregnancy, which may be life-saving for mother and baby.

Epilepsy in pregnancy

Although the outcome is successful (normal pregnancy with healthy baby) for more than 90% of epileptic women, there is a slightly increased risk of prematurity, low birth rate, perinatal mortality, defects and intervention. About 25% of women have an increased number of seizures, due mainly to a fall in antiepileptic drug levels with a small increased frequency during labour and the puerperium. On the other hand 25% have a reduction in frequency. It is important to take oral folic acid supplementation (5 mg daily) prepregnancy and up to 12 weeks' gestation, and to share the care with the patient's neurologist who can advise on the most appropriate antiepileptic and dosage for the patient. All antiepileptic drugs are potentially teratogenic, with different drugs being related to different defects: phenytoin has been related to cleft lip and palate and congenital heart disease, while sodium valproate (in particular) and carbamazepine have been associated with spina bifida.

All antiepileptic drugs are excreted in breast milk but in such reduced concentrations as not to preclude breastfeeding. For subsequent contraception a higher dose oestrogen pill is recommended because the agent usually increases liver enzyme activity.

Multiple sclerosis

Patients with multiple sclerosis usually manage very well in pregnancy, which appears to have a stabilising effect on the disorder. Preventive advice against urinary tract infections is advisable and spinal anaesthesia should be avoided.

ANTEPARTUM HAEMORRHAGE

If haemorrhage occurs less than 24 weeks treat as for threatened miscarriage. If it occurs after 26 weeks admit to hospital for management. Remember to give anti-D gamma globulin if the mother is Rhnegative. Do not perform a vaginal examination. The main causes are placental, namely placenta praevia (unavoidable APH) and placental abruption (accidental). Placental abruption in particular has a high risk of causing fetal death in utero with coagulopathy complications.

Placenta praevia

The placenta has a low attachment onto the lower uterine segment and may cover the cervix. Incidence is about 1%. Rather than classify placenta praevia according to grades I - IV it is best to divide it into 'minor' and 'major' grades. Presentation usually includes painless bleeding at 28 - 30 weeks gestation. There is a high presenting part on palpation.

Principles of management

- Resuscitation of the mother as required - insert IV line and cross-match blood.
- Assess fetus with ongoing serial growth scans.
- Confirm diagnosis by ultrasound.
- Aim to prolong pregnancy, especially if minor grade, to reduce prematurity.
- If major grade admit to hospital for observation with a view to rest and watchful expectancy, and subsequent delivery by caesarean section (the safest approach).
- Give anti-D prophylaxis if mother is Rh D-negative.
- Seek an opinion whether vaginal birth is a safe option to consider with a marginal placenta praevia. If bleeding restarts or continues, arrange termination by caesarean section.

Caesarean section is always required for major placenta praevia. Vaginal delivery may be possible in a specialist unit after careful assessment recommends a trial of labour in a patient with a minor degree of praevia where the fetal head lies below the lower edge of the placenta.

Placental abruption[10]

Placental abruption (incidence 1%) is retroplacental bleeding from a normally situated placenta resulting in detachment of a segment of decidua from the uterine wall. The patient presents with mid-abdominal pain, bleeding PV, and a tense and tender uterus (large for dates) and signs of hypovolaemic shock. The cases can vary from mild to severe. It is the commonest obstetric cause of coagulopathy.

Principles of management

- Admit to hospital for full assessment of mother and baby.
- Resuscitate and restore circulatory blood volume if necessary. Give blood until patient recovers and ‘shocked’ uterus recovers (four times the measured loss is generally needed). Coagulation products may be necessary.
- Perform tests - FBE, coagulation profile, Kleihauer test to define any feto-maternal haemorrhage, blood for cross-matching, kidney function, electrolytes.
- Cardiotocography (CTG) for fetal assessment, especially accurate heart activity. May need to maintain continuous CTG. Fetal death is the rule with severe placental abruption.
- Ultrasound examination of uterus and contents, to exclude placenta praevia and presence/size of any retroplacental clot.
- Rest in bed and pain relief if required.
- Give course of corticosteroids to mature the baby's lungs as urgent delivery may be necessary.
- Introduce Foley's catheter to monitor urine output.

The objective is to aim for vaginal delivery, especially if the baby is dead. Induce when the condition is stable to prevent further retroplacental bleeding. This is normally done by rupturing the membranes and using a Syntocinon drip. Use ergometrine in the third stage unless there is evidence of pre-eclampsia. Caesarean section has been recommended where the baby's life is immediately threatened, but this is hazardous in the presence of coagulopathy.

Vasa praevia

Vasa praevia is a rare cause of APH due to rupture of fetal blood vessels. It coincides with rupture of the membranes and it is a vital diagnosis to make as it can lead rapidly to fetal exsanguination. Diagnosis is by a characteristic ominous pattern on CTG and the Apt's test. Emergency delivery is indicated.

PRIMARY POSTPARTUM HAEMORRHAGE

Primary postpartum haemorrhage is loss of > 500 mL of blood within 24 hours of delivery. A severe PPH is defined as > 1 000 mL blood loss.

Causes

- Uterine atony
- Retained placenta/placental fragments
- Coagulation disorder
- Soft tissue laceration of genital tract, e.g. episiotomy, cervical tear
- Ruptured or inverted uterus

Principles of management

- Rub up the uterine fundus.
- Resuscitate.
- Stop the bleeding.
- Make the diagnosis - inspect placenta/speculum examination of genital tract.
- Catheterise (a full bladder can aggravate the problem).
- An urgent examination and exploration under general anaesthetic may be necessary.

Treatment

- IV access - commence crystalloid IV fluids
- Tests: FBE, cross-match blood, coagulation profile

- High-flow oxygen by mask
- IV oxytocin 10 IU followed by 40 IU in IV infusion of Hartman's solution
- If continuing heavy bleeding ergometrine 0.25 - 0.5 mg IM or IV
- If retained placenta - deliver with cord traction or manual removal
- If continuing heavy bleeding - bimanual compression for 3 minutes
- Treat any coagulation disorder, e.g. fresh plasma/platelet transfusion

If a persistent atonic uterus is not controlled by oxytocics, 1 - 2.5 mg doses of intramyometrial prostaglandin F2-α can be injected through the abdominal wall. Life-saving measures can include uterine artery ligation, internal iliac artery ligation (usually bilateral) or hysterectomy.

BLOOD GROUP ISOIMMUNISATION

Blood group or red cell isoimmunisation is primarily related to Rhesus D (RhD) isoimmunisation leading to haemolytic disease of the newborn from the effect of the development of anti-D antibodies.[10] These antibodies develop from fetomaternal haemorrhage/ transfusion in RhD negative women carrying a RhD positive fetus.

Effects of haemolytic disease on the fetus includes hydrops (oedema), FDIU.

Effects on the neonate include anaemia, heart failure, jaundice and hepatosplenomegaly.

Screening Rh negative mothers: at presentation, 28 weeks, and 34 - 36 weeks.

Do not perform screening if anti-D has been given.

Rhesus immunoprophylaxis

Indications for giving anti-D Ig to the RhD negative mother free of immune anti-D:

- after spontaneous miscarriage at any stage of pregnancy
- after threatened miscarriage
- after delivery of an RhD positive baby
- following termination of pregnancy or ectopic pregnancy
- following any sensitising event during pregnancy (that may provoke a transplacental haemorrhage), e.g. amniocentesis or CVS, APH, external cephalic version, significant closed abdominal trauma
- prophylactically at 28 and 34 weeks in an apparently normal pregnancy

Kleihauer test

This is a test on maternal blood after a sensitising event to detect the degree of fetomaternal transfusion and whether additional anti-D Ig is required.

ABO incompatibility

This occurs when the mother is group O and the baby A or B and can occur in the first pregnancy without a tendency to become increasingly severe in subsequent pregnancies. A small number of babies have mild jaundice while severe haemolytic consequences are rare.

THROMBOEMBOLISM IN PREGNANCY

Pregnancy is associated with an increased risk of thromboembolism with an incidence of about 1% of deep venous thrombosis (DVT). Untreated DVT carries about 15% risk of pulmonary embolism. DVT or pulmonary embolism should be suspected in a woman in the antenatal or postpartum period who complains of pain or swelling in the leg, mild unexplained fever, dyspnoea or chest pain.

Risk factors include past history of DVT, prolonged bed rest, operative delivery, multiparity, postpartum surgical procedure, anaemia, inherited thrombophilia disorders or antiphospholipid antibodies. If a DVT is suspected, low molecular weight heparin is recommended until investigation and specialist advice are obtained.

HYDATIDIFORM MOLE[8]

Hydatidiform mole is an overgrowth of gestational trophoblastic tissue. The moles may be complete (no fetal tissue) or partial (some fetal tissue). Incidence 1:1400 pregnancies. There is a risk that some persistent gestational trophoblastic may become invasive and penetrate the uterus and metastasize to the lungs. 1:20 progress to develop choriocarcinoma.

Presentation

- Bleeding in early pregnancy ± passage of grapelike debris.
- May be exaggerated symptoms of pregnancy, e.g. hyperemesis.
- Uterus large for dates.

Management

- Investigations: FBC, blood group and crossmatch, hCG level (very high), ultrasound pelvis (typical ‘snow-storm’ appearance), chest X-ray.
- Suction curette with oxytocin drip.
- Consider hysterectomy if patient has completed family planning.
- Register in the trophoblastic registry.

Follow-up

- Chest X-ray: ? metastatic disease.
- Weekly serum (or urine) hCG until zero (usually takes 8 - 12 weeks), then monthly for 12 months.
- Avoid pregnancy for 12 months after hCG levels normal.
- The oral contraceptive pill is appropriate.
- Later a putrid vaginal discharge indicates malignancy.
- Refer for possible cytotoxic therapy, e.g. methotrexate and folinic acid.

MULTIPLE PREGNANCY

This is associated with increased risks to mother and children.

Facts

- Spontaneous incidence[8, 10]: twins 1:80; triplets 1: 80^2 (1:6400); quads 1: 804
- Increased incidence with clomiphene induced ovulation and IVF
- Twins: monozygotic (identical) 30%, dizygotic 70%
- Conjoint twins are a special problem
- Predisposing factors: family history, previous twins, infertility treatment, race esp. African blacks, ↑ age
- Diagnosis: hyperemesis, ultrasound, polyhydramnios, large for dates, palpation two fetal heads/three poles ± many limbs, two different hearts heard
- Presentation twins: first twin cephalic 70%, both cephalic 40%, cephalic + breech 30%

Complications

- Maternal: increased risk anaemia: symptoms of pregnancy, e.g. morning sickness, varicose veins; pre-

eclampsia × 3, antepartum and postpartum haemorrhage; malpresentation; cord prolapse; caesarean section
- Fetal/neonatal: increased risk abnormalities, preterm delivery (premature labour, premature rupture membranes); intrauterine growth restriction of one fetus; twin - twin transfusion; perinatal mortality × 5; prematurity; malformations × 2 - 4; (also those of mother)

Principles of management

- Increased supplements (iron/folic acid), nutrition requirements and rest
- Increased frequency of antenatal visits and associated care, e.g. weekly visits from 28 weeks
- Education and counselling about increased incidence of above, e.g. preterm birth
- Early referral to self-help groups
- Ultrasound examination at 28 weeks, then serial scans to check fetal growth of both babies
- Intensive fetal monitoring
- If threatened premature delivery admit to hospital for rest; consideration of tocolytics to gain time and antenatal corticosteroids for fetal lung maturation
- Aim to deliver at 38 weeks if possible either by vaginal delivery (if favourable conditions such as normal growth and double cephalic presentations) or caesarean section, e.g. malpresentation first twin, conjoint twins
- Careful attention to third stage.↑ risk of haemorrhage

PRETERM LABOUR

Preterm or premature labour is confirmed labour after 20 weeks and before 37 weeks gestation. The incidence is 5 - 10% (average 7%) of all deliveries and is associated with 85% of neonatal deaths. There is often a past history. Causes of spontaneous preterm labour:

- unknown (approx. 40%)
- multiple pregnancy
- cervical incompetence
- polyhydramnios
- uterine abnormal maternal medical conditions, e.g. diabetes, drug abuse, infection

The patient may present with regular contractions or premature rupture of the membranes.

Management

- Admit to an obstetric unit.
- Consider tocolysis (inhibition of uterine contractions) with atosiban, nifedipine or β-sympathomimetic agents (salbutamol, ritodrine, fenoterol).
- Give corticosteroid, e.g. betamethasone, hydrocortisone, for ↑ fetal lung maturity (maximum benefit 28.32 weeks).

PREMATURE RUPTURE OF MEMBRANES

Premature rupture of the membranes (PROM) is rupture of the membranes with amniorrhoea before labour commences.

Preterm PROM (PPROM) is rupture of the membranes < 37 weeks gestation.

Key points

- 50% of PPROM progress to labour within 24 hours (80% within 7 days).
- Smoking is a risk factor for PROM.
- Differential diagnosis of PROM includes profuse vaginal discharge, incontinence of urine.20% false alarms for

aminorrhoea (aminohexis).

- A vaginal examination should not be performed.
- Signs of infection (chorioamnionitis): maternal fever, tachycardia, ↑ WCC, ↑ CRP, signs of fetal distress.
- Don't give antibiotics if cervical smears and culture normal.

Management guidelines

- Perform routine examination including vital signs.
- Speculum examination - liquor pools in posterior fornix.
- Do not perform a vaginal examination.
- Admit to hospital.
- Take vaginal - cervical smears for culture.
- Perform WCC or CRP every 2 - 3 days.
- Continuous observations.
- If no evidence of infection continue the pregnancy; deliver if signs of infection.
- CTG every second day or daily.
- Give corticosteroid therapy if delivery prior to 34 weeks likely.
- Induce if CTG abnormal or infection present.
- Induce if not in labour by 36 weeks.

PROLONGED PREGNANCY

Prolonged pregnancy is pregnancy lasting longer than 42 weeks. The due date is based on pregnancy lasting 40 weeks when 65% will proceed to spontaneous labour within the next week.

Normal delivery is between 37 and 42 weeks.

At 41 weeks check CTG and amniotic fluid index (AFI) - allow to proceed if normal.

Induce labour at 42 weeks as perinatal mortality rate is ↑ × 2 from 42 - 43 weeks and more so after 43 weeks. Induction is probably best achieved using prostaglandin E2 vaginal gel, or oral or vaginal misoprostol, followed by ARM if labour does not follow the use of prostaglandin alone.

INTRAUTERINE GROWTH DISORDERS

Fundus greater than dates

Consider the following causes:

- polyhydramnios
- multiple pregnancy
- macrosomic baby > 90th percentile - diabetes or history of large babies
- uterine abnormality, e.g. fibroids
- wrong dates

Polyhydramnios

Features

- Liquor volume: usually > 2000 mL
- Multiple risks, e.g. PROM, prem labour, cord
- prolapse, APH, malpresentation

Causes

- Fetal abnormalities: CNS, upper GIT atresia, ectopic vesicae
- Hydrops fetalis
- Diabetes
- Multiple pregnancy
- Chorioangioma of placenta
- Fetal infection - cytomegalovirus, toxoplasmosis
- Unknown causes

Refer for tests - diabetes, ultrasound.

Refer for specialist advice.

Fundus less than dates

Consider:

- oligohydramnios - liquor volume usually < 500 mL
- small baby
- intrauterine growth restriction
- wrong dates

Oligohydramnios is associated with conditions such as fetal abnormality, prolonged pregnancy, kidney disease, pre-eclampsia, congenital infections (CMV, toxoplasmosis), PROM and placental insufficiency.

Investigation should include ultrasound, kidney function tests, lupus antibodies and regular CTG. Refer for specialist management.

Intrauterine growth restriction (IUGR)[9]

IUGR is defined as an estimated birth weight < 10th percentile. Apart from the disorders outlined in the preceding topics, causes to also consider for a small baby include previous history of a small baby, racial factors, prematurity, various infections and maternal factors such as smoking, drugs, alcohol, anaemia and nutrition.

Sometimes it is difficult to detect IUGR on routine clinical assessments, but most cases are detected when the examiner finds that the symphysis - fundal height is below expectations. Other telltale signs include oligohydramnios and reduced fetal movements. Appropriate management is referral for an ultrasound and specialist advice.

MECONIUM-STAINED LIQUOR

Key points

- Fresh meconium is dark green and sticky.
- Meconium staining is common post-term (13% all births).
- It is a sign of fetal distress but meconium alone is an inadequate marker of perinatal asphyxia.
- It is more common for the baby to be normal.
- Cord prolapse is an important cause.
- Aspiration of meconium causes pneumonitis in the neonate.

Management

- Assess fetal condition with continuous cardiotocography.
- Perform a pelvic examination - assess progress and cord prolapse.

- If all is well, allow labour to proceed to a vaginal delivery.
- If minor abnormality on CTG, measure fetal scalp pH or lactate.

If abnormal - caesarean section or manipulative vaginal delivery.

Note: As the head is born aspirate the oropharynx (visualise the vocal cords) or nose (same applies to CS). The presence of a paediatrician is ideal.

OTHER HIGH-RISK CONDITIONS

Malpresentations

The important malpresentations are breech (4% of all babies) and transverse or oblique lie. A primary concern is the high risk of cord presentation and prolapse. The usual standard practice is to perform CS as the best option.

Breech presentation

The general rule is to deliver by caesarean section (CS), especially since studies have shown that the risk of vaginal delivery was much higher than that associated with CS. However, in selected patients and circumstances, vaginal delivery is safe, especially with spontaneous labour that progresses appropriately with a normal-sized baby and normal pelvic dimensions. However, if in doubt - caesar!

Transverse or oblique lie (1:300)

These presentations are more common in multigravida. Perform an ultrasound examination to exclude placenta praevia. The lie may convert to a longitudinal one. Should it persist beyond 37 weeks admit to hospital and if it persists or labour commences CS is the best option.

Cord prolapse and presentation[11]

When the diagnosis is made on pelvic examination and the cord is still pulsating, push the presenting part away and also push the cord up as far as possible and keep it there by continuous digital pressure. This is a particular problem if the mother is in labour. Place her into the knee - chest position and organise urgent delivery, usually by CS while maintaining the position of the hand. If the woman is fully dilated (second stage) and circumstances are favourable, a vaginal delivery with forceps or vacuum extraction, avoiding cord pressure, can be performed.

Malpositions

Important malpositions include occipito-posterior (the commonest), occipito-transverse, face and brow. The general rules to follow are that with good analgesia (e.g. epidural), no evidence of cephalo-pelvic disproportion and no fetal or maternal distress, allow labour to proceed and await events. Aim for a vaginal delivery assisted by instruments if necessary, especially with the occipito-posterior and occipitotransverse positions. A notable exception is the face presentation where the head rotates back into the mento-posterior position making vaginal delivery impossible and CS is necessary. Otherwise CS is also the best option in prolonged labour with maternal exhaustion, obstructed labour, disproportion, fetal distress and selected primigravida.

Shoulder dystocia

Impacted shoulders causing sudden arrest of delivery after delivery of the baby's head is a terrifying complication of childbirth requiring expert assistance. Various manoeuvres can be attempted to achieve the widest possible diameter of the pelvic outlet and thus facilitate delivery of the shoulders. (Consider McRobert's manoeuvre11 - suprapubic pressure, shoulder rotation performed vaginally, delivery of the posterior shoulder.)

Inverted uterus

This potentially lethal complication occurs when the fundus of the uterus turns inside out, either partially or completely. If it is complete the inside of the uterine fundus may appear outside the introitus. The placenta is usually still attached to the uterus and often the problem is caused by over-traction of the cord without waiting for firm uterine contractions. The easiest way to manage this complication is to return the uterus to its normal position with the cord still attached immediately after the event. If unsuccessful resuscitate and attempt hydrostatic replacement of the inverted uterus.[10] Otherwise surgical intervention will be necessary.

CONSIDERATION FOR INDUCTION

Consideration for induction:

- post-term (41 weeks or over) - a metaanalysis of 11 RCTs found that there were benefits to both mother and fetus in being induced between 10 and 14 days after the due date[11]
- maternal hypertension
- maternal distress
- diseases of pregnancy (e.g. pre-eclampsia)
- intrauterine growth restriction
- intrauterine fetal death
- diabetes mellitus
- isoimmunisation
- unstable lie

TRAUMA: MOTOR VEHICLE ACCIDENTS

Abdominal trauma in pregnancy is usually associated with seat-belt restraints during motor vehicle accidents. However, these injuries are far less severe than those that occur when people are not wearing seat belts. Women should be encouraged to wear seat belts and should not be given certificates stating that seat belts should not be worn in pregnancy.

The incidence of placental abruption following accidents is related to the severity of the accident and the extent of the external injuries. Injured patients should be admitted to a unit where cardiotocography can be performed regularly for 48 hours and perinatal intensive care can be provided if needed. Consider anti-D gammaglobulin injection.

DRUGS IN PREGNANCY

Drugs have to be used with great care during pregnancy. An Australian categorisation of drug risk is presented in summary in Table 101.2. It is worth noting that β_2-agonists used to treat asthma have a category A rating.

Table 101.2 Examples of medicines in pregnancy: an Australian categorisation of risk (Australian Drug Evaluation Committee)

	Category*
Iron and haemopoietic agents	
• Folic acid	A
• Iron preparations (all types)	A
Antihistamines and antiemetics	

• Phenothiazines (e.g. prochlorperazine)	C
• Meclozine, cyclizine	A
• Other antihistamines	A or B2
Alimentary system agents	
• Antacids	A
• H_2-receptor antagonists	B1
• Proton pump inhibitors	B3
Cardiovascular	
• ACE inhibitor	D
• Methyldopa	A
• Calcium-channel blockers	C
• Beta-blockers	C
• Digoxin	A
• Diuretics (except spironalactone, B3)	C
• Glyceryl trinitrate	B2
Analgesics	
• Aspirin	C
• Paracetamol/acetaminophen	A
• Codeine	A
• Opioid analgesics	C
Hypnotics, sedatives, antipsychotic agents	
• Barbiturates	C
• Benzodiazepines	C
• Chloral hydrate	A
• Phenothiazines and butyrophenones	C
Antidepressants	
• SSRIs	C
• Tricyclics (e.g. amitriptyline)	C
• Tetracyclics (e.g. mianserin)	B2
Anticonvulsants (all groups)	D
NSAIDs	C
Antimicrobials	
• Penicillins	A
• Cephalexin, cephalothin	A
• Aminoglycosides	D
• Nitrofurantoin	A
• Tetracyclines	D
• Guanine analogues, e.g. acyclovir	B3
• Azithromycin, roxithromycin	B1

• Ciprofloxacin	B3
• Erythromycin	A
• Nystatin	A
• Norfloxacin	B3
Corticosteroids	
• Systemic	C
• Inhalation	B3
Quinine	D

[*] A - No harmful fetal effect recorded.

B - No harmful effects to date but limited experience (see ADEC guidelines for subgroups B1, B2, B3).

C - Have caused or suspected of causing harmful effects on fetus or neonates without causing malformations (reversible).

D - Have caused, are suspected to cause or may be expected to cause an increased incidence of fetal malformation or irreversible damage. Also may have adverse pharmacological effects.

X - Have such a high risk of causing permanent damage to the fetus that they should not be used in pregnancy or where there is a possibility of pregnancy.

WHEN TO REFER

If there is a possibility of cervical incompetence; refer for a specialist opinion before 14 weeks.

Referral to specialist centre[12]

The key to an optimal outcome is early identification of the high-risk pregnancy and early referral to a specialist team to supervise the management of the remainder of the pregnancy. This has been shown to improve neonatal morbidity and mortality significantly. It is important that family physicians, obstetricians, perinatologists and neonatalogists work as a harmonious team.

REFERENCES

1. Shires DB, Hennen BK, Rice DI. *Family medicine*. New York: McGraw-Hill, 1987: 136 - 51.
2. Slaytor EK, Sullivan EA, King JF. *Maternal deaths in Australia* 1997 - 1999. AIHW Cat. No. PER 24. Sydney: AIHW National Perinatal Statistics Unit, 2004.
3. Laws PJ, Sullivan EA. Australian mothers and babies 2003. AIHW Cat. No. PER 29. Sydney: AIHW National Perinatal Statistics Unit, 2005. (Perinatal Statistics Series No. 16).
4. Centre for Epidemiology and Research. New South Wales Mothers and Babies 2003. NSW Public Health Bulletin, 2004: 15(S-5).
5. Michael CA. Hypertensive disease in pregnancy. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 260 - 3.
6. Brown MA (ed). Pregnancy and hypertension. Bailliere's Best Practice and Research. Clinical Obstetrics & Gynaecology. Vol. 13. Cambridge: Bailliere Tindall, 1999.
7. Altman D, Carrol, G. Do women with pre-eclampsia and their babies benefit from magnesium sulphate? The Magpie Trial: a randomised placebo-controlled trial. Lancet 2002; 359(9321): 1877 - 90.
8. The Eclampsia Trial Collaborative Group. Which anticonvulsant for women with eclampsia? Evidence from the Collaborative Eclampsia Trial. Lancet, 1995; 345: 1455.

9. Goh J, Flynn M. *Examination Obstetrics and Gynaecology* (2nd edn). Sydney: Maclennan+Petty, 2005.
10. Martin FIR. The diagnosis of gestational diabetes. MJA, 1991; 15: 112.
11. Humphrey MD. *The Obstetrics Manual* (revised edn). Sydney: McGraw-Hill, 1999.
12. Peat B. Antenatal care: common issues facing GPs in shared care. Med Today, 2001; June: 81 - 8, 260 - 3.

CHAPTER 102. POSTNATAL CARE

As concerning the bringing up, nourishment and giving of suckle to the child, it shal be beste that the mother give her child sucke her selfe, for the mothers milk is more convenient and agreeable to the infant than any other woman's or other milke.

THOMAS RAYNALDE 1540, *THE BYRTHE OF MANKYNDE*

POSTNATAL CARE

Education for the puerperium and caring for the baby should begin during pregnancy so that a new mother is familiar with the basic principles of motherhood, especially infant feeding.[1] The puerperium is defined as the period of approximately 42 days from the completion of the third stage to the return of the normal physiological state.

Postnatal care really begins with the birth of the baby. Once the airways are cleared the baby should be given to the mother as soon as possible and not taken from her except for essential management.

The mother should remain in the labour ward (if delivered in hospital) for at least an hour after giving birth and until she has passed urine. She should be inspected frequently to exclude the possibility of a silent postpartum haemorrhage and vital signs checked before transferring to a lying-in ward. It is worth remembering that one-third of eclamptic convulsions occur postpartum.

GUIDELINES FOR THE LYING-IN WARD[1]

- Every mother needs rest but should have full toilet and shower facilities.
- The baby should be in a bassinet beside the mother and may be taken into bed any time the mother likes.
- Room in: the baby should not go to the nursery unless it is sick or the mother requests it.
- There should be no visiting restrictions on close relatives but restrictions should be put on other visitors for the first 2 - 3 days.
- Demand feed to appetite.
- No test weighing.
- No complementary feeding unless mother is empty and baby screaming.
- A golden rule is that breastfeeding and the supply of mother's milk is a classic case of ‘supply and demand’.
- The doctor should listen carefully to what the mother is saying (and not saying) during visits.
- Check that anti-D vaccine was given (if necessary).

POSTNATAL CONSULTATIONS

The two-week consultation

Mother:

- Assess the coping ability of the new mother
- Look for signs/symptoms of postpartum depression
- Provide encouragement and advice
- Check breastfeeding

Baby:

- Routine examination
- Perform a Phenistix test on the baby's napkin (in case the Guthrie test has been missed in hospital)

The six-week consultation

This is basically a repeat of the previous consultation and a checklist is presented in Table 102.1.

Table 102.1 Checklist for postnatal check at 4 - 6 weeks

Mother
Pap smear (if not performed at first visit)
Check rubella status
Check hepatitis B status
Review antenatal screening test for follow up action
Check for adequate contraception
Check if intercourse has been resumed and give advice (if appropriate)
Check bowel and urine control
Encourage abdominal and pelvic floor exercises
Check for back problems
Check weight, blood pressure and urine
Check breasts
Check abdomen (uterus should be impalpath)
Caesarean round
Check perineum
Check psychological health, including coping ability
Check for postpartum thyroiditis
Discuss adequate diet, rest and personal care
Perform pelvic examination
Check pelvic floor strength
Further follow-up if necessary
Give Personal Health Record folder to mother
Baby
Routine examination
Check growth and feeding
Educate mother regarding immunisation schedule

CONTRACEPTION

Breastfeeding, when it is truly on demand, is an extremely good contraceptive, but in reality some supplement is necessary for about 3 months in the average lactating woman.

Oral contraception

Delay to after 21 days because ↑ risk of throurbosis (COC) and bleeding (POP)

- The mini pill (progestogen only)
    - norethisterone 350 µg/day or
    - levonorgestrel 30 µg/day taken every night
- Transfer to COC when breastfeeding completed (oestrogens can suppress lactation).

IUCD: If used, it should be fitted at or after 6 weeks.[2]

AFTER-PAINS[1]

After-pains, which are more common and most intense after the second and subsequent pregnancies, are characterised by intermittent lower abdominal pains, like period pains, which are often worse during and after feeding in the first 2 weeks. They are caused by oxytocin released from the posterior pituitary, which also causes the let-down reflex of nursing. Suspect endometritis if offensive lochia, fever and poor involution of uterus.

Treatment, after examination, is reassurance and analgesics in the form of paracetamol every 4 hours for 3 days or as long as necessary.

BREASTFEEDING

Insufficient milk supply

This is sometimes a problem in mothers who are under a lot of stress and find it hard to relax. A let-down reflex is necessary to get the milk supply going, and sometimes this reflex is slow. If there is insufficient supply, the baby tends to demand frequent feeds, may continually suck its hand and will be slow in gaining weight.[1]

There are three important factors in establishing breastfeeding:

1. positioning the baby on the breast
2. the ‘let-down’
3. supply and demand

The breasts produce milk on the principle of supply and demand. This means that the more the breasts are emptied, the more milk is produced.

Advice to mother

- Try to practise relaxation techniques.
- Put the baby to your breast as often as it demands, using the ‘chest to chest, chin on breast’ method.
- Express after feeds, because the emptier the breasts, the more milk will be produced.
- Make sure you get adequate rest, but if you feel tired go to your doctor for a check-up.

Engorged breasts

Engorgement occurs when the milk supply comes on so quickly that the breasts become swollen, hard and tender. There is an increased supply of blood and other fluids in the breast as well as milk. The breasts and nipples may be so swollen that the baby is unable to latch on and suckle.

Advice to mother

- Feed your baby on demand from day 1 until the baby has had enough.
- Finish the first breast completely; maybe use one side per feed rather than some from each breast. Offer the second breast if the baby appears hungry.
- Soften the breasts before feeds or express with a warm washer or shower, which will help to get the milk flowing.
- Avoid giving the baby other fluids.
- Express a little milk before putting the baby to your breast (a must if the baby has trouble latching on) and express a little after feeding from the other side if it is too uncomfortable.
- Massage any breast lumps gently towards the nipple while feeding.
- Apply cold packs after feeding and cool washed cabbage leaves (left in the refrigerator) between feeds. Change the leaves every 2 hours.

- Wake your baby for a feed if your breasts are uncomfortable or if the baby is sleeping longer than 4 hours.
- Use a good, comfortable bra.
- Remove your bra completely before feeding.
- Take paracetamol regularly for severe discomfort.

Regular feeding and following demand feeding is the best treatment for engorged breasts.

Suppression of lactation[2]

Women may seek suppression of lactation for a variety of reasons such as weaning the baby, not wishing to breastfeed initially, or after stillbirth.

Mechanical suppression

The simplest way of suppressing lactation once it is established is to transfer the baby gradually to a bottle or a cup over a 3-week period. The decreased demand reduces milk supply, with minimal discomfort. If abrupt cessation is required, it is necessary to avoid nipple stimulation, refrain from expressing milk and use a well-fitting bra. Use cold packs and analgesics as necessary. Engorgement will gradually settle over a 2 - 3 week period.

Hormonal suppression

Hormonal suppression can be used for severe engorgement but only as a last resort. It is more effective if given at the time of delivery but may produce side-effects. Avoid oestrogens.

- Cabergotine 1mg (0) statim or
- Bromocriptine 2.5 mg (o) bd for 10 - 14 days[2] (nausea, dizziness and hypo tension) are problems with both drugs.

Drugs and lactation

Drugs that can affect lactation or a breastfed infant are listed in Table 102.2.

Table 102.2 Drugs taken by nursing mother that can affect breastfed infant or lactation

Contraindicated drugs
Antibiotics:
• aminoglycosides
• chloramphenicol
• nitrofurantoin
• metronidazole
• tetracycline
• sulphonamides
Antihistamines
Antineoplastics/cytotoxics
Benzodiazepines
Bromocriptine
Combined oral contraceptive/oestrogens
Ergotamine
Gold salts
H_2-receptor antagonists (e.g. cimetidine, ranitidine)
Illicit drugs (e.g. cocaine, cannabis, LSD)

Lithium Quinidine Laxatives (e.g. senna)
Alcohol (no harmful effects unless taken in excess) Nicotine (increased respiratory distress in infants exposed)

Nipple problems with breastfeeding

Sore nipples

Sore nipples is a common problem, thought to be caused by the baby not taking the nipple into its mouth properly, often because of breast engorgement. The problem is preventable with careful attention to the feeding position of the baby.

Management advice to mother

It is important to be as relaxed and comfortable as possible (with your back well supported) and for your baby to suck gently.

- Try to use the ‘chest to chest, chin on breast’ feeding position.
- Vary the feeding positions (make sure each position is correct).
- Start feeding from the less painful side first if one nipple is very sore.
- Express some milk first to soften and ‘lubricate’ the nipple. (Avoid drying agents such as methylated spirits, soap and tincture of benzoin, and moisturising creams and ointments, which may contain unwanted chemicals and germs.)
- Gently break the suction with your finger before removing the baby from the breast. (Never pull the baby off the nipple.)
- Apply covered ice to the nipple to relieve pain.
- Keep the nipples dry by exposing the breasts to the air and/or using a hair dryer on a low setting.
- If wearing a bra, try Cannon breast shields inside the bra. Do not wear a bra at night.

Note: Raynauds phenomenon can affect the nipple and cause painful breast feeding. It is often mistaken for candida albicans infection.

Cracked nipples

Cracked nipples are usually caused by the baby clamping on the end of the nipple rather than applying the jaw behind the whole nipple. Not drying the nipples thoroughly after each feed and wearing soggy breast pads are other contributing factors. Untreated sore nipples may progress to painful cracks.

Symptoms

At first, the crack may be so small that it cannot be seen. The crack is either on the skin of the nipple or where it joins the areola. A sharp pain in the nipple with suckling probably means the crack has developed. Feeding is usually very painful, and bleeding can occur.

Management advice to mother[3]

Cracked nipples nearly always heal when you get the baby to latch onto the breast fully and properly. They usually take only 1 - 2 days to heal.

- Follow the same rules as for sore nipples.
- Do not feed from the affected breast - rest the nipple for 1 - 2 feeds.
- Express the milk from that breast by hand.
- Feed that expressed milk to the baby.

- Start feeding gradually with short feeds.
- A sympathetic expert such as an experienced nursing mother will be a great help if you are having trouble coping.
- Take paracetamol just before feeding to relieve pain.

Inverted nipples

An inverted nipple is one that inverts or moves into the breast instead of pointing outwards when the baby tries to suck from it. When the areola is squeezed, the nipple retracts inwards.

Treatment[3]

During pregnancy, rolling and stretching the nipple by hand can be helpful. The partner can assist with gentle oral and manual stimulation.

A simple treatment, which should start at the beginning of the seventh month of pregnancy, is the Hoffman technique:

1. Draw an imaginary cross on the breast with the vertical and horizontal lines crossing at the nipple.
2. Place the thumbs or the forefingers opposite each other at the edge of the areola on the imaginary horizontal line. Press in firmly and then pull the thumbs (or fingers) back and forth to stretch the areola.
3. In the vertical position, pull the thumbs or fingers upwards and downwards.

This procedure should be repeated about five times each morning. The nipple will become erect and is then easier to grasp, so that it can be slowly and gently drawn out.

MASTITIS

Mastitis which has a high incidence (up to 20%) is basically cellulitis of the interlobular connective tissue of the breast. Usually restricted to lactating women it is caused mainly by a cracked nipple or poor milk drainage. The infecting organism is usually Staphylococcus aureus or more rarely Escherichia coli or Candida albicans. It is a serious problem and requires early treatment. Breastfeeding from the affected side can continue as the infection is confined to interstitial breast tissue and doesn't usually affect the milk supply.

Note: Mastitis must be treated vigorously - it is a serious condition.

Refer to Chapter 90.

Clinical features

- A lump and then soreness (at first)
- A red wedge-shaped possibly tender area
- Fever, tiredness, muscle aches and pains

Note: Candida infection usually causes severe breast pain - a feeling like a hot knife or shooting pains, especially during and after feeding. It may occur after a course of antibiotics.

Management

Prevention (in lactation)

- Maintain free breast drainage
- Attend to breast engorgement and cracked nipples

Treatment[4]

- Antibiotics: resolution without progression to an abscess will usually be prevented by antibiotics: dicloxacillin 500 mg (o) qid for 7-10 days or flucloxacillin 500 mg (o) qid for 7-10 days or cephalexin 500 mg (o) qid for 7-10 days
  If severe cellulitis: flucloxacillin/dicloxacillin 2g IV 6 hourly

- Therapeutic ultrasound (2 W/cm^2 for 6 minutes) daily for 2 - 3 days
- Aspirin or paracetamol for pain
- Treat candidiasis with fluconazole 200 - 400 mg (o) daily for 2 - 4 weeks

Instructions to patients

- Keep the affected breast well drained.
- Continue breastfeeding: do it frequently and start with the sore side.
- Heat the sore breast before feeding (e.g. hot shower or hot face washer).
- Cool the breast after feeding: use a cold face washer from the freezer.
- Massage any breast lumps gently towards the nipple while feeding.
- Empty the breast well: hand express if necessary.
- Get sufficient rest.
- Keep to a nutritious diet and drink ample fluids.

BREAST ABSCESS

If tenderness and redness persist beyond 48 hours and an area of tense induration develops, then a breast abscess has formed. It requires surgical drainage under general anaesthesia, antibiotics, rest and complete emptying of the breast. Temporary weaning of breastfeeding from the affected side is necessary because of the surgical disruption.

For a description of surgical drainage refer to pages 970 - 1.

SECONDARY POSTPARTUM HAEMORRHAGE[5]

Primary PPH is presented on page 1057 - 58.

Secondary Postpartum haemorrhage is any bright bleeding from the birth canal 24 hours or more after delivery. It may vary from very slight to torrential and may occur at any time up to 6 weeks postpartum. It tends to peak at 5-10 days.

Causes

- Retained products of conception (POC)
- Infection, especially at placental site
- Laceration of any part of the birth cana
- Coagulation disorder

No cause is found in 1/3 of cases i.e. idiopathic subinvolution

Treatment

Rule: An empty and contracted uterus will not bleed.

- Investigation: ultrasound (? Retained POC); cervical shiear and culure; FBE
- IV oxytocin 10 IU followed by infusion of 40 IU in Hartman's solution
- Ergometrine 0.25 - 0.5 mg IM or IV (if continuing heavy bleeding)
- Exploration under general anaesthetic if blood loss > 250 mL
    - gentle blunt curettage required in the postpartum uterus
- Antibiotics (e.g. amoxycillin/clavulanate 500 mg (0) 8 hourly) while awaiting culture and blood (if Hb < 100 g/L) as indicated

Note: Referral is necessary after the ergometrine injection. Occasionally a life-saving hysterectomy or ligation of the internal iliac arteries may be necessary.

Lochia discharge

The discharge of lochia should be followed. Normal:

1. Bloody loss = lochia rubra 2 - 12 days
2. Serous loss = lochia serosa: up to 20 days
3. White loss = lochia alba
4. offensive lochia = endometritis

Lochia loss persists for 4 to 8 weeks. Abnormal lochia rubra indicates a retained placenta. If there is a problem examine with a speculure and take cervical/vaginal swab.

PUERPERAL FEVER

Puerperal fever is defined as raised temperature of = 38°C from day 1 to day 10. If fever - think of the three B's - birth canal, breast, bladder. The cause is genital infection in about 75% of patients. Other causes include urinary tract infection, mastitis and an intercurrent infection. Investigations include a vaginal swab for smear, culture and sensitivities (include anaerobic culture) and a midstream specimen of urine for microscopy and culture.

Treatment

- Amoxycillin/potassium clavulanate plus metronidazole (while awaiting sensitivities)

Beware of severe puerperal sepsis such as Gram-negative septicaemia or Clostridium welchii septicaemia.

POSTNATAL DEPRESSIVE DISORDERS

It is quite common for women to feel emotional and flat after childbirth; this is apparently due to hormonal changes and to the anticlimax after the long-awaited event. There are three separate important problems:

1. postnatal blues
2. postnatal adjustment disorder
3. postnatal (or postpartum) depression

POSTNATAL BLUES

'The blues' is a very common problem (occurs in 80%) that arises in the first 2 weeks (usually days 3 - 10) after childbirth.

Clinical features

- Feeling flat or depressed
- Mood swings
- Irritability
- Feeling emotional (e.g. crying easily)
- Feeling inadequate
- Tiredness
- Insomnia
- Lacking confidence (e.g. in bathing and feeding the baby)
- Aches and pains (e.g. headache)

Fortunately ‘the blues’ is a passing phase and lasts only 4 - 14 days. Management is based on support, reassurance and basic counselling. Contact friends and relatives to help.

Advice to mother[2]

All you really need is encouragement and support from your partner, family and friends, so tell them how you feel.

- Avoid getting tired and rest as much as possible.
- Talk over your problems with a good listener (perhaps another mother with a baby).
- Accept help from others in the house.
- Allow your partner to take turns getting up to attend to the baby.

If ‘the blues’ lasts longer than 4 days, it is very important to contact your doctor.

POSTNATAL ADJUSTMENT DISORDER

- Occurs in first 6 months
- Similar symptoms to ‘the blues’
- Anxiety with handling baby
- Psychosomatic complaints
- Fearful of criticism

Treatment

- Support and reassurance
- Cognitive therapy
- Parentcraft support
- Settles with time

POSTNATAL DEPRESSION

Some women develop a very severe depression after childbirth. Always consider it in the frequent attender. Symptoms are present for at least 2 consecutive weeks with onset in the first few days postpartum.

- Occurs in 10 - 30% women
- In first 6 - 12 months (usually first 6 months): peaks about 12th week.
- Anxiety and agitation common
- Marked mood swings
- Poor memory and concentration
- Typical depressive features

Treatment

- Support, reassurance, counselling
- Group psychotherapy
- Couple therapy
- Postnatal depression support group
- Hospitalisation may be necessary (especially if suicidal or infanticidal ideations)
- Medication - amitriptyline (nortriptyline), SSRIs (setraline, paroxitine agents of choice)

Note: Beware of puerperal psychosis with onset usually within first 2 weeks.

POSTPARTUM PSYCHOSIS

Symptoms that appear within the first month include unusual bebaviour, agitation, delusions, hallucinations, mania and suicidal ideations.

It is rare, occurring in about 1:500 births.

Past history may be a pointer. Suspect if severe depression not responding to treatment. Organise impotent psychiatric care.

SLEEP DEPRIVATION

Give advice and counselling. Use the ‘sleep when baby sleeps’ rule. Avoid sedatives.

TIREDNESS

Tiredness is very common in the first few months after delivery. It may be a presenting symptom of anaemia, depression, hypothyroidism, anxiety or depression.

Perform a FBE and TFTs.

POSTPARTUM HYPOTHYROIDISM

Postpartum hypothyroidism (postpartum thyroiditis) may be misdiagnosed as postpartum depression and should always be considered in the tired, apparently depressed woman in the first 6 months after delivery.

HAIR LOSS

Increased hair shedding as telogen effluvium is common about 4-6 months after delivery. Large clumps of hair with white buffs come out easily with combing or shampooing. Reassure that it reverts to normal in 3-6 months.

BACK PAIN AND COCCYGODYNIA

Postnatal backache occurs in about 50% of women and may persist for several weeks. Management includes simple analgesics, massage, exercises, hot or cold packs, education regarding lifting of body and referral to a physiotherapist for persistent pain.

SEXUAL DIFFICULTIES

Decreased libido is a common problem and often related to sleep deprivation. Only 50% of couples achieve intercourse by the 6 weeks check.[2] Decreased libido can also be due to one of the postnatal depressive disorders or to tension in adjusting to the new relationships.

Dyspareunia is common and should be treated symptomatically and with education. Simple lubrication or vaginal oestrogens can help until perineal healing is achieved.

Early intercourse is risky with deaths from air embolism reported in the first 2 weeks. Intercourse is not recommended in the first 6 weeks.

ELIMINATION DISORDERS[2]

Always enquire how the patient is coping with her bowels and urination. Simple advice such as stool softening and pelvic floor exercises will help. However, serious problems such as faecal incontinence secondary to a fistula from a third-degree tear or urinary retention due to neuropraxia of the pelvic floor can develop and need urgent attention.

REFERENCES

1. Smibert J. Practical postnatal care. Aust Fam Physician, 1989; 18: 508 - 11.
2. McKenna M. Postnatal problems. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 423 - 5.
3. Murtagh J. Patient Education (4th edn). Sydney: McGraw-Hill, 2004: 4-10.
4. Spicer J (Chair). Therapeutic Guidelines: Antibiotic (Version 12). Melbourne: Therapeutic Guidelines Ltd, 2003: 226.
5. Smibert J. Common puerperal complications. Aust Fam Physician, 1989; 18: 824 - 7.

CHAPTER 88. CERVICAL CANCER AND PAP SMEARS

If all women have regular Pap smears, one every two years, we can prevent 90% of all cervical cancers.

DR GABRIELE MEDLEY, TIME, 24 APRIL 1995

CARCINOMA OF THE CERVIX

Carcinoma of the cervix is the most common malignancy in women worldwide; it is the sixth most common in Australia[1] and seventh in the United States.[2] The incidence of invasive cervical cancer rises steadily from age 20 to 50 and then remains relatively steady.

The most common form of cervical cancer is squamous cell carcinoma (SCC) 85-90, with adenocarcinoma representing 10-15%.[2]

A striking epidemiological feature about cervical cancer is that it is a disorder related to sexual activity. It is almost non-existent in virgins but has an increased incidence in women with multiple partners and those who began sexual activity at an early age. Thus, epidemiological studies indicate that cervical cancer is a sexually transmitted disorder (Table 88.1).

Table 88.1 Cervical cancer and risk factors

Age	Increased after 55
Sexuality	Increased • with multiple and or promiscuous sex partners • early age for first intercourse • early age first pregnancy
Viruses	Increased after herpes II or wart virus infection (probable)
Occupation	Increased in prostitutes (decreased in nuns)
Parity	Increased multiparity
Socioeconomic status	Increased with low socioeconomic status

Facts and figures

- Invasive cervical cancer is almost unknown in women under the age of 20, and very rare before age 25.
- There are two small peaks of incidence, in the late 30s and late 60s.[1]
- The lifetime probability of an Australian woman developing cancer is 1 in 90.[3]
- On average, cervical cancer takes at least a decade to develop from a focus of a cervical squamous intraepithelial lesion[4]
- SCC of the cervix occurs almost exclusively in women who have had coitus.
- The earlier the age of first intercourse the greater the chance of developing cervical cancer.
- Invasive cervical cancer is a disease for which definite curable premalignant lesions can be identified using a Papanicolaou's (Pap) smear as a screening test.
- The incidence of cervical cancer has been decreased significantly through the screening procedures of the Pap smear, colposcopy and colposcopically directed cervical biopsy.
- Poor Pap smear technique is a common cause of a false negative result.
- The GP needs to achieve the best possible cervical cell sample and forward it to the best possible cytology laboratory.
- Despite the availability of liquid-based smears, a well taken conventional Pap smear is still a very good screening test.
- New methods of laboratory examination of the smear include Papnet, which involves computer scanning of the smear, and ThinPrep, whereby a liquid-based sample is prepared.

Basic pathology

The focus of attention is the transformation zone (Figure 88.1) where columnar cells lining the endocervical canal undergo metaplasia to squamous cellsóin the region of the squamocolumnar junction. It is important clinically to realise that this transformation zone can extend with progressive metaplasia of columnar epithelium and so the squamo-columnar junction may recede into the endocervical canal. This is a feature in postmenopausal women (Figure 88.2). As squamous cell carcinoma almost always arises in the transformation zone, it is vital that cells are taken from it when performing a Pap smear.



Figure 88.1 The transformation zone: it is vital that cells are taken from this zone with Pap smears


Appearance of cervix
reproductive age
menopausal
endocervical canal
transformation zone (squamocolumnar junction)
external os
vagina
Instruments for PAP smear
cervex sampler broom or spatula (blunt end) + endocervical brush
spatula (pointed end) or cervex sampler broom + endocervical brush

Figure 88.2 Changing position of the transformation zone with age, and a selection of sampling instruments according to its position (the Cervex sampler can be used as an alternative to the spatula)

Cervical intraepithelial neoplasia

Cellular changes can occur in the transformation zone for a variety of reasons, including invasion with human papilloma virus (HPV). One such important change is cervical dysplasia, previously known as cervical intraepithelial neoplasia (CIN) and squamous intraepithelial lesion in the now adopted modified Bethesda system.[5, 6] These dysplasias have the potential to become invasive cervical cancer.

Natural history of cervical dysplasia

Dysplasia may return to normal, persist or eventually progress to invasive cervical cancer. The reported progression times to cervical cancer range from 1 to 30 years. On average it takes at least 10 years, so it is considered that 2-yearly Pap smears are a reasonable safety margin. However, women with histologically confirmed moderate to sever dysplasia require a colposcopic assessment.

Figure 88.3 and Table 88.2 illustrates the disease spectrum of cervical neoplasia.



Figure 88.3 Illustration of the various grades of squamous intraepithelial lesions and CIN (comparison of nomenclature)

Table 88.2 Squamous cell abnormalities and different nomenclatures [5,6]

Description	CIN grade	Australian modified Bethesda system
1 Normal	Normal	Within normal limits
2 Atypia : reactive or neoplastic	Atypia	ASCUS
3 HPV	HPV	LSIL
4 Mild dysplasia	CIN 1	LSIL
5 Moderate dysplasia	CIN 2	HSIL
6 Severe dysplasia	CIN 3	HSIL
7 Carcinoma in situ	CIS	HSIL
8 Invasive carcinoma	Invasive carcinoma	Invasive carcinoma

ASCUS = Atypical squamous cell of undetermined significance

LSIL = Low-grade squamous intraepithelial lesion

HSIL = High-grade squamous intraepithelial lesion

CIS = Carcinoma in situ

CIN = Cervical intraepithelial neoplasia

Clinical presentation

Many patients with cervical cancer are asymptomatic and when early symptoms do arise they are often dismissed as of little consequence.

Symptoms, if present, may be:

- vaginal bleeding, especially postcoital bleeding
- vaginal discharge
- symptoms of advanced disease (e.g. vaginal urine or flatus, weakness)

Screening recommendations

Routine Pap smears:

- Perform every 2 years for women18-70 years of age with no clinical evidence of cervical pathology and who have ever had sex.
- Perform from beginning of sexual activity up to 70 years.
- Begin Pap smears at 18 - 20 years or 1 - 2 years after first sexual intercourse (whichever is later).
- Cease at 70 years in those who have had two normal Pap smears within the last 5 years.
- Perform a Pap smear on women over 70 years if they request it or if they have never had a smear.
- Ideally, practices should have a reminder or a recall system.

Women who have never engaged in coitus do not need Pap smears. Six-monthly or 12-monthly screening on young asymptomatic women provides only minimal benefit compared with 2-year intervals.

Hysterectomy

Smears are needed if the cervix was not completely removed. However, vaginal vault smears are needed if there is a history of gynaecological dysplasia or malignancy, exposure to diethyl stilboestrol *in utero* and in immune suppressed women.

TAKING A PAP SMEAR[1,7]

The importance of a good specimen

The optimal Pap smear contains:

- sufficient mature and metaplastic squamous cells to indicate adequate sampling from the whole of the transformation zone
- sufficient endocervical cells to indicate that the upper limit of the transformation zone was sampled; and to provide a sample for screening of adenocarcinoma and its precursors

Optimal timing of specimens

- The best time is any time after the cessation of the period.
- Avoid smear-taking during menstruation.
- Avoid in the presence of obvious vaginal infection.
- Avoid within 48 hours of use of vaginal creams or pessaries or douching.
- Avoid within 24 hours of intercourse.
- Avoid lubrication or cleaning of cervix with preliminary pelvic examination.

Communicating with the pathologist

Good communication with the pathologist is essential. It is important to provide basic details about the reason for the Pap smear and the clinical history on the pathology form sent to the laboratory. Include patient age, LMP, hormone in take previous treatment and clinical findings. Use the opportunity also for breast examination and checking of breast self-examination (BSE) technique.

The method

1. *Education and explanation*
   Take time to explain the reason for taking the Pap smear, especially if it is the first. Emphasise that it is mainly a preventive measure to detect and treat early cell changes that could develop into cancer. Anatomical models, sample instructions or charts are useful in describing the procedure. Explain that it does not hurt and doesn't take long, that it may be uncomfortable but slow deep breathing will help relaxation and make it easier. It is preferable to talk to the patient during the examination with appropriate explanation. It is advisable for a male doctor to have a chaperone present.
2. *Equipment*
   Prepare the following equipment:
   - adequate light source
   - speculum (preferably bivalve) warmed under lukewarm water
   - glass slide labelled in pencil with the woman's name and date of birth
   - spray fixative
   - plastic gloves for both hands
   - smear-taking instruments; choose from:
     - Ayer's spatula, wooden or plastic
     - Cervex sampler broom
     - endocervical brush
3. *Positioning*
   The supine position is usually best (Figure 88.4). The left lateral position can be used if smears are difficult to obtain (e.g. older women with lax anterior vaginal walls, older women with poor hip mobility and the very embarrassed patient). The Sims exaggerated left lateral position (Figure 88.5) provides better exposure of the vulva but requires more manipulation of the patient. Better visualisation of the cervix is obtained if the patient elevates her buttocks with her hands (best as fists).



Figure 88.4 The supine or dorsal position is the best position for the speculum examination and subsequent bimanual palpation (Patient should be appropriately clothed and/or draped)



Figure 88.5 The Sims exaggerated left lateral position

4. Inserting the speculum[6]
   Avoid using lubricating jelly on the speculum blades. Warming the speculum with water should provide adequate lubrication. Gently spread the labia with a gloved hand and introduce the speculum with the blades vertical or at 458 from the vertical. Gently advance the blades with slow firm pressure towards the rectum as far as possible. Rotate the blades during the process until they are horizontal and exerting gentle pressure against the posterior wall of the vagina. Remember that the cervix is situated in the upper sixth of the anterior vaginal wall (not in the apex of the vagina).
5. *Visualising the cervix*
   Good lighting and exposure of the cervix is essential. Note any significant features or abnormalities of the cervix. Reassure the woman if the cervix looks normal with a comment such as 'Your tissues look very healthy'.
   A cervical ectropion is normal in most premenopausal women and was formerly incorrectly called an erosion.
6. *Taking the smear*
   Choose the sampling instrument that best suits the shape of the cervix and os. Place Ayer's spatula firmly on the os and rotate it through 3608, ensuring that the whole transformation zone is sampled (Figure 88.6a).



Figure 88.6 Method of smear taking and preparing the slide

If the squamous columnar junction is not visible (lying within the endocervical canal), use both spatula (first) and the cytobrush (Figure 88.6b). The cytobrush (tends to cause bleeding) should be advanced until only the lower bristles are still visible, then rotated twice through 3608. The cytobrush should be avoided in pregnant women.

After removing the speculum, perform a bimanual pelvic examination.

7. *Preparing the slide*

   Transfer the cervical cell sample on to a glass slide with an even spreading motion (Figure 88.6c, d). Fix immediately within 30 seconds to prevent air drying, which distorts cellular features with an aerosol or pump-action alcohol spray (Figure 88.6e).

   The *HPV and chlamydia sample*

   If appropriate after the smear, place the brush and spatula in the tube with the transport medium. Swirl it around vigorously to release material. The specimen tube can be forwarded with the slide to the laboratory with a request to test for HPV and chlamydia.

   Follow-up

   Discuss mutually suitable arrangements to ensure that the woman obtains the result of the smear whether it is positive or negative. Inform her when her next smear is likely to be due (special cards are available) and arrange to send a reminder note.

   The explanation of the results, especially if there is an abnormality present (a variety of abnormal smear), should be crystal clear to the patient.

Abnormal cervical cytology

Confirmation of the Pap smear result is by colposcopy and/or by a biopsy and appropriate referral should be arranged without delay.

Inaccurate results can be caused by:[8]

- using dirty glass slides

- using lubricants or doing pelvic examinations before taking the smear
- insufficient material
- endocervical cells not being taken in the smear (i.e. taken from the wrong site)
- a thick film with an inadequate spread of material
- air-drying before fixing
- smear not being fixed for long enough or the solution of alcohol being too weak
- the slide not being dry before being placed in the cardboard container (this encourages fungal overgrowth).

Abnormal Pap smears

Follow the guidelines in Table 88.3 and Figure 88.7 for the abnormal smear result.

Table 88.3 Guidelines for abnormal Pap smears[7]

Pap smear report	**Investigation and management**
No endocervical cells	Repeat in 2 years.
Negative smearóinflammatory cells	Repeat smear in 2 years.
Unsatisfactory smear	Repeat smear in 6-12. weeks (allows regeneration of cells).
Low-grade epithelial lesion	
Possible LSIL and Defined LSIL	Repeat Pap smear at 12 months. If the woman is 30^{+} years, and has no negative 2-3 years, refer for colposcopy or repeat the test in 6 month
High-grade epithelial Lesion	
Possible HSIL Definite HSLI	Refer for colposcopy
Glandular abnormalities including adenocarcinoma *in situ*.	Refer to a expert gynaecologist.
Invasive squamous cell carcinoma or adenocarcinoma	Refer to appropriate specialist gynaecologist or unit
Inconclusiveóraising possibility of high-grade disease	Refer for colposcopy and possible biopsy


Low grade (LSIL)
(definite or possible)
If woman is 30+ years, and has no negative cytology in previous 2–3 years, immediate colposcopy or repeated test in 6 months.
Repeat Pap test at 12 months
Negative
Low grade (LSIL)
(definite or possible)
High grade (HSIL)
Repeat Pap test at 12 months
Colposcopy
Negative
Return to 2 yearly screening

Figure 88.7 Algorithm for management of low-grade squamous cell abnormalities (based on NH&MRC guidelines)

Post-treatment assessment of HSIL

A woman treated for HSIL should have a colposcopy and cervical cytology at 4-6 months after treatment. Cervical cytology and HPV typing should be done at 12 months and then annually until the woman has tested negative by both tests on two consecutive occasions. Return to usual 2 yearly screening when all four tests are negative.

PREVENTION OF CERVICAL CANCER

‘In other words, chastity and fidelity are recommended for those who can, and condoms for those who cannot’.[9] This statement is a succinct recommendation for prevention and includes:

- Ideally people should have intercourse with only one partner.
- The male should use a condom on each occasion if either sexual partner is unsure of the other's previous behaviour.
- Those at risk should be counselled accordingly.

Other preventive measures include:

- Women should have Pap smears at least 2 yearly.
- Identification of high-risk forms of persistent HPV will aid surveillance. If absent, no treatment is needed as smears become normal.
- Use of beta-carotene has a protective effect against cervical cancer, so ‘both sexes would be well advised to ensure regular intake of green leaf and orange vegetables in their diet’.[9]
- Advise against smoking.

VACCINATION

A new quadrivalent human papilloma virus (types 6,11,16,18) recombinant vaccine is indicated for the prevention of cancer and pre-cancers due to vaccine HPV type in females aged 9 - 26.

MEDICOLEGAL ISSUES[7]

Cervical cancer screening is a potential minefield of litigation, which has increased greatly especially over missed cancers following a false negative Pap smear (a particular dilemma for cytology laboratories).

Common claims made against GPs include:

- failure to offer cervical screening
- failure to adequately investigate abnormal vaginal bleeding (especially post coital bleeding)
- poor communication including inappropriate use of phone contact.
- failure to arrange adequate specialist referral for women with abnormal cytological results or a clinically suspicious cervical lesion.

Advice and reassurance should be given in a diplomatic way that does not produce guilt feelings. This includes reassurance that not all cervical cancer is sexually transmitted, that women with only one partner may develop cervical cancer and that sexual contact with a male partner who has had the wart virus does not always result in cancer of the cervix.[9]

REFERENCES

1. Screening for the prevention of cervical cancer. Canberra: Department of Health, Housing and Community Services, 1991: 1 - 26.
2. Rakel RE. *Essentials of Family Practice*. Philadelphia: Saunders, 1993: 130 - 1.
3. Giles G, Armstrong GK, Smith LR eds. *Cancer in Australia*. Melbourne: National Cancer Statistics Clearing House. Scientific Publications No. 1, Australasian Association of Cancer Registries and Australian Institute of Health, 1987.
4. Day NE. Screening for cancer of the cervix. J Epidemiol Community Health, 1989; 43: 103 - 6.
5. Kurman RJ, Solomon D. *The Bethesda System for reporting Cervical/vaginal Cytologic Diagnoses*. New York: Springer-verlag,1994
6. NH&MRC. *Screening to prevent cervical cancer. Guidelines for the Management of Asymptomatic women with Screen Detected Abnormalities. The Australian Modified Bethesda System*. Canberra: NH&MRC, 2005.
7. Reid R, Hyne S. Taking better Pap smears Medicine Today, 2004; 5(1): 59-65
8. Craig S. The smear test. Aust Fam Physician, 1985; 14: 1092 - 4.
9. Tattersall M. *Preventing Cancer*. Sydney: Australian Professorial Publications, 1988: 182 - 97.

CHAPTER 89. FAMILY PLANNING

> The Membranous Envelope (condom) is prepared from the bladder of a fish caught in the Rhine. Its extreme thinness does not in the least interfere with the pleasure of the act … [its use] is of the greatest utility because, while it is a sure preventive of conception, it also prevents either party from contracting disease.
>
> EDWARD BLISS FOOTE 1864, *Medical Common Sense*

FAMILY PLANNING

Effective family planning requires a good understanding of the function of the menstrual cycle, whether it is for the purpose of conception or contraception.

The main consultation is the presentation of a young woman for contraceptive advice. It is a very critical visit and provides an excellent opportunity to develop a good rapport with the patient and provide education and counselling about important health concerns, such as health promotion, menses regulation, sexual activity, planned parenthood, fertility and infertility, pregnancy prevention, STI prevention, immunisation and cervical smears.[1]

In counselling and treating patients, especially teenagers, confidentiality is of paramount importance. The issues and contraceptive methods can be confusing so careful education using charts and other aids is recommended to enhance the therapeutic relationship and facilitate better compliance.[1]

It is worth discussing the patient's attitude to pregnancy, including the fear of pregnancy and the possible reaction to contraceptive failure.

FERTILITY CONTROL

The choice of contraceptive methodology will be determined not only by individual needs, personal preference and resources but also by its safety and incidence of side-effects.

It is worth emphasising that the estimated risk of death associated with child-bearing (1 in 10 000 in developed countries) is higher than the risk of death associated with all methods of contraception, with two exceptions: women over 35 years of age who smoke and take the combined oestrogen - progestogen oral contraceptive, and those over 40 years of age taking this type of preparation.[2] In developed countries of the Western world the most widely used methods, in order of preference, are combined oral contraceptives (COC), condoms, diaphragms, intrauterine devices, spermicidal agents and rhythm.[2]

A comparison of the efficacy of the various contraceptive methods is presented in Table 89.1.

Table 89.1 Effectiveness of contraceptive methods[3, 4]

Method	Effectiveness (pregnancies per 100 years of use) Lowest expected (reliable consistent user)	Pearl index
Natural rhythm methods	20 - 30	
• Billings ovulation (cervical mucus) method	3	2 - 3
Withdrawal (coitus interruptus)	20 - 25	18
Spermicides		
• vaginal sponge	10	-
• diaphragm (with spermicide)	15	6
• condoms	10 - 15	3♂ 5♀

Intrauterine devices	3 - 5	0.1 - 1
Vaginal ring	1 - 3	0.65
Oral contraceptives		
• combined	1 - 3	0.1
• progestogen only	3	0.5
Depomedroxyprogesterone acetate	0.1	0.3
Implant	0.06	0.09
Female sterilisation	0.02	0.4
Male sterilisation	0.15	0.1

Pearl Index = (total accidental pregnancies × 1200)/total months of exposure

More than half the pregnancies in the United States are unintended and occur because of nonuse of contraception, failure of a specific method or discontinuation of contraception.[1, 3]

For women at risk of acquiring STIs the choice of contraception has to consider methods that protect against both pregnancy and STIs.

STEROIDAL CONTRACEPTION

Methods of steroidal contraception include:[5]

- combined oral contraceptive pill
- progestogen-only pill (POP)
- injectables
- postcoital contraception
- implants (Implanon)
- levonorgestrel-releasing IUCD (Mirena)
- progestogen-releasing vaginal rings
- oestrogen - progestogen-releasing vaginal rings

Combined oral contraception

COCs usually contain a low-dose oestrogen and a moderate dose of progestogen. The main mode of action of COC is inhibition of hypothalamic and pituitary function leading to anovulation.[2]

Which oestrogen to use[5]

Mestranol and ethinyloestradiol (EO) are about equipotent. Mestranol undergoes metabolic conversion to EO in the liver before it exerts its contraceptive effect. EO is therefore the oestrogen of choice.

Which progestogen to use[5]

All progestogens are nor-testosterone derivatives and exhibit a variety of non-progestogenic actions.

The norethisterone (NET) group includes norethisterone acetate, ethynodiol acetate and lynestrenol. The last three progestogens are converted to NET before exerting any contraceptive activity.

Levonorgestrel (LNG) is 10 times more potent than NET. It has less effect on the coagulation system than NET and is therefore the preferred progestogen.

Gestogens are the ‘third generation’ progestogens and include desogestrel, gestodene, norgestimate and cyproterone acetate. These agents, which are less androgenic than NET and LNG, have been implicated with an increased risk of thromboembolism but the data are of doubtful validity. The latest progestogen is drospireone, which is an analogue of the diuretic spironolactone.

Starting the pill: which COC to use[5]

The aim is to provide good cycle control and effective contraception with the least side-effects using a pill of the lowest dose. The past menstrual history and contraceptive use of the patient should be documented and taken into account in selecting the appropriate COC. Various COC preparations available in Australia are listed in Table 89.2.[6, 7]

Table 89.2 Combined oral contraceptive pill formulations

Oestrogen	Dose (µg)	Progestogen	Dose (µg)	Trade name
Monophasic				
Ethinyloestradiol	20	Levonorgestrel	100	Microgynon 20, Loette, Microlevlen
Ethinyloestradiol	30	Levonorgestrel	150	Nordette, Levlen ED Microgynon 30, Monofeme
Ethinyloestradiol	30	Gestodene	75	Femoden ED, Minulet
Ethinyloestradiol	30	Desogestrel	150	Marvelon
Ethinyloestradiol	35	Cyproterone acetate	2000	Brenda-35 ED, Diane-35 ED, Estelle ED
Ethinyloestradiol	35	Norethisterone	500	Brevinor, Norimin
Ethinyloestradiol	35	Norethisterone	1000	Brevinor-1, Norimin-1
Ethinyloestradiol	35	Drospirenone	3000	Yasmin
Ethinyloestradiol	50	Levonorgestrel	125	Nordette 50, Microgynon 50
Ethinyloestradiol	50	Levonorgestrel	250	Nordiol
Mestranol	50	Norethisterone	1000	Norinyl-1
Triphasic				
Ethinyloestradiol	30, 40	Levonorgestrel	50, 75, 125	Triphasil, Triquilar Trifeme 28, Logynon ED
Ethinyloestradiol	35, 40	Norethisterone	500, 1000	Synphasic, Improvil 28 day

A suitable first choice is a monophasic pill containing 30 µg ethinyloestradiol (EO) with levonorgestrel or norethisterone (e.g. Nordette, Microgynon 30, Monofeme, Levlen ED). Cover starts immediately if started on day 1 of cycle (includes switching from POP).

The high-dose monophasic (50 µg oestrogen) should be reserved for the following situations:

- breakthrough bleeding on low-dose COCs
- control of menorrhagia
- concomitant use of enzyme-inducing drugs
- low-dose pill failure

Education and counselling is very important for the woman starting the pill. Suitable patient education should be given. The pill can be used safely up to 50 years of age. Cover starts immediately if COC commenced on day 1 of the cycle.

Specific patient groups[7, 8]

Adolescents. The COC can be prescribed once menstruation has commenced, with appropriate counselling about safe sex and responsibilities. The monophasic low-dose combined preparation should be selected.

Epilepsy. Use a COC with a high dose of oestrogen (e.g. 50 µg).

Women with hirsutism. Use a less androgenic preparation (e.g. Diane-35).

Women over 35 years. Use a low-dose monophasic COC provided the woman is a non-smoker. If continued until about 50 years, the hot flushes of the perimenopause are controlled. It is usual to cease the pill at around 50 - 51, wait several weeks and then measure the serum FSH and oestradiol levels. If the oestradiol levels are low and FSH high, the woman can be regarded as menopausal and can start HRT if desired.

Menstrual disorders: menorrhagia/dysmenorrhoea. Start with a standard low-dose monophasic COC but a higher-dose oestrogen (50 μg) pill may be necessary.

Acne. For women with acne (not on COC), commence with a less androgenic progestogen (e.g. Diane-35 ED, Marvelon).

The high-dose monophasic (50 μg EO) should be reserved for the following situations:

- breakthrough bleeding on low-dose COCs
- control of menorrhagia
- concomitant use of enzyme-inducing drugs
- low-dose pill failure

Contraindications to COC usage are shown in Table 89.3.

Table 89.3 Contraindications for use of the COC[6]

Absolute
• Pregnancy (known or suspected)
• First 2 weeks postpartum
• History of thomboembolic disease, including known thrombophilia
• Cerebrovascular disease
• Focal migraine
• Coronary artery disease
• Oestrogen-dependent tumours
• Recent impaired liver function
• Polycythaemia
Relative
• Heavy smoking
• > 35 years and smoking or other risks of CAD
• Undiagnosed abnormal vaginal bleeding
• Breastfeeding
• 4 weeks before surgery
• 2 weeks after surgery
• Gall bladder or liver disease
• Hypertension
• Diabetes mellitus
• Long-term immobilisation
• Complicated valvular heart disease
• Hyperlipidaemia
• Chloasma
• Severe depression

Efficacy of COCs

Under ideal circumstances the pregnancy rate in women taking COCs is 1 - 3 per 100 women years of use, but in practice varies from 2 - 6 per 100 women years.[2]

Non-contraceptive advantages of COCs

A number of significant beneficial effects arising from the use of COCs have now been documented:

- reduction in most menstrual cycle disorders
- reduction in the incidence of functional ovarian cysts
- 50% reduction in the incidence of PID
- reduced incidence of ovarian and endometrial carcinoma
- benign breast disease reduced
- fewer sebaceous disorders
- reduced incidence of thyroid disorders

Serious side-effects of COCs

The most serious side-effects to be considered are the effects of COCs on the circulatory system and the incidence of cancer.

Cardiovascular effects[5]

The following circulatory disorders have been linked with pill usage:

• Venous	deep vein thrombosis pulmonary embolism rarely: mesenteric, hepatic and kidney thrombosis
• Arterial	myocardial infarction thrombotic stroke haemorrhagic stroke rarely: retinal and mesenteric thrombosis

The risk of circulatory disease has not been related to duration of use and there is no increased risk in perpetual users.

The oestrogen content of the pill is considered to be the aetiological factor and the problem is increased in women taking high-oestrogen-content COCs, but now that the oestrogen content of each pill has been reduced to as low as 20 μg EO, these risks of morbidity and mortality have been reduced.

The progestogen effect on lipid metabolism is not considered significant in the aetiology of circulatory disease. Circulatory diseases have now been recognised as occurring predominantly in certain high-risk groups - the ‘at - risk female’, particularly the smoker over 35 years of age.

Other risk groups include those with hyperlipidaemia, diabetes, hypertension, and a family history of cardiovascular disease or immobilisation.

Provided low-dose COCs are prescribed in lowrisk females it would appear safe to use the COC pill up to 50 years of age.

COCs and cancer

There appears to be no overall increase in the incidence of cancer in women using COCs. Possible effect (not absolutely proven):

- cervix (take regular smears at yearly intervals)
- breast

Protective effect:

- endometrial
- epithelial ovarian

No effect:

- melanoma
- chorioncarcinoma
- prolactinomas

Common side-effects

The relatively minor side-effects listed in Table 89.4 may discourage women from persisting with oral contraception in the absence of appropriate explanation and reassurance. Management of these side-effects is listed in the same table. It is useful in practice to have this list available as a ready reference for manipulating the COC if necessary.

Table 89.4 Management of common side-effects of COC[7, 9]

Symptom change	Change	Examples of pill change
Acne	Increase oestrogen, reduce or change progestogen	Triphasil/Triquilar to Diane ED/Marvelon
Amenorrhoea	Increase oestrogen or decrease progestogen	Nordette/Microgynon 30 to Nordette 50/Microgynon 50
Breakthrough bleeding		
• early to mid cycle	Increase oestrogen	Triphasil/Triquilar to Biphasil/ Sequilar
• late cycle	Increase progestogen or change type	Triphasil to Nordette Nordette to Norinyl-1
Breast problems		
• fullness/tenderness	Decrease oestrogen	Biphasil/Sequilar to Triphasil/Triquilar
• mastalgia	Decrease progestogen	Nordette/Microgynon 30 to Triphasil/Triquilar
Chloasma	Stop oestrogen Try progestogen-only pill Avoid direct sun (use blockout)	
Depression	Decrease or change progestogen	Nordette/Microgynon 30 to Triphasil/Triquilar or Brevinor
Dysmenorrhoea/menorrhagia	Increase progestogen	Triphasil/Triquilar to Nordette/Microgynon
Libido loss	Increase oestrogen Change from anti-androgenic progestogen to an alternative	Microgynon 30 etc. to Femoden/Minulet
Headache		
• focal migraine	Discontinue pill	
• in pill-free week	Add 10 - 30 μg ethinyloestradiol daily during pill-free week	
Nausea/vomiting	Decrease or change oestrogen or stop oestrogen	Use Microgynon 20, etc. or progestogenonly pill
Weight gain		
• constant	Decrease or change progestogen	Triphasil/Triquilar to Brevinor or Marvelon
• cyclic	Decrease oestrogen	Biphasil/Sequilar to Triphasil/Triquilar or progestrogen-only pill

Important advice for the patient

- Periods tend to become shorter, regular and lighter.

- No break from the pill is necessary.
- Drugs that interact with the pill include vitamin C, antibiotics, griseofulvin, rifampicin and anticonvulsants (except sodium valproate). Warfarin and oral hypoglycaemics requirements may change for those starting the pill.
- Diarrhoea and vomiting may reduce the effectiveness of the pill.
- Yearly return visits are recommended to update the history and examination and repeat the Pap smear.

Missed pills

The essential advice is 'just keep going' i.e. take a pill as soon as possible and then resume usual pill - taking schedule.

Also

If the missed pills are in week three, she should omit the pillfree interval.

Also

Condoms or abstinence should be used for 7 days if the following numbers of pills are missed:

'Two for twenty' (i.e. if two or more 20 μg pills are missed)

'Three for thirty' (i.e. if three or more 30 - 35 μg pills are missed)

The seven-day rule for the missed or late pill (more than 12 hours late)

- Take the forgotten pill as soon as possible, even if it means taking two pills in one day. Take the next pill at the usual time and finish the course.
- If you forget to take it for more than 12 hours after the usual time there is an increased risk of pregnancy so use another contraceptive method (such as condoms) for 7 days.
- If these 7 days run beyond the last hormone pill in your packet, then miss out on the inactive pills (or 7-day gap) and proceed directly to the first hormone pill in your next packet.
- You may miss a period. (At least seven hormone tablets should be taken.)

Other useful rules for missed pills

If 1 or 2 × 30-35 μg EO pills or 1 × 20 μg EO pill

- take the most recent missed pill ASAP
- continue taking remaining pills as usual

No additional contraception or emergency contraception needed.

If > 3 × 30-35 μg EO pills *or* > 2 × 20 μg EO pills

- take the most recent missed pill ASAP
- continue taking remaining pills
- use condoms or abstinence until pill is taken for 7 consecutive days

Practice tip

Delaying a period

Prescribe norethisterone 5 mg bd or tds for 3 days prior to expected period.

Period resumes 2 - 3 days after stopping tablets.

If taking COC:

- continue taking the hormone tablets (skip the inactive pills) until end of next pack.

PROGESTOGEN-ONLY CONTRACEPTIVE PILL

The POP is perhaps an underutilised method of contraception, although it is not as efficacious as the COC.

The two common formulations are:

- levonorgestrel 30 µg/day and
- norethisterone 350 µg/day

Providing the mini-pill is taken regularly at the same time each day, the pregnancy rate is 3 per 100 women years.[2] The failure rate decreases with age. There are no serious side-effects but compliance is a problem because of cycle irregularity, especially with irregular bleeding. The mini-pill often reduces the cycle length to less than 25 days or alters the regularity of the bleeding phase.

Indications for the POP include age 45 years or more, smokers aged 45 years or more, contraindications to or intolerance of oestrogens, diabetes mellitus, migraine, chloasma, lactation and well-controlled hypertension.

Contraindications include pregnancy, undiagnosed genital tract bleeding, past history of or increased risk of ectopic pregnancy and concomitant use of enzyme-inducing drugs (absolute).

INJECTABLE CONTRACEPTIVES

Depo-Provera

Medroxyprogesterone acetate (Depo-Provera) is the only injectable intramuscular contraceptive available in Australia. It is very effective for up to 14 weeks.

Dose: 150 mg by deep IM injection in first five days of the menstrual cycle. The same dose is given every 12 weeks to maintain contraception.

Failure rate: 1 per 1000 women years.[2]

Side-effects include a disrupted menstrual cycle (amenorrhoea rate 70% or irregular or prolonged uterine bleeding), excessive weight gain, breast tenderness, depression and a delay in return of fertility (average 6 months).[5] There is no effect on cardiovascular disease or the incidence of cancer.

There are no absolute contraindications.

Etonogestrel implant (Implanon)

This is a subdermal contraceptive implant that is a 3 year system consisting of a single rod containing the progestogen, etonorgestrol. It inhibits ovulation and has an anti-mucus effect. Irregular bleeding is the most common side-effect. It requires a minor surgical procedure to insert it and also to remove it.

EMERGENCY CONTRACEPTION

- Postinor-2 [Plan B (USA)]: one 750 µg levonorgestrel tablet followed by another tablet 12 hours later. Limited to first 72 hours.
- Yuzpe method: use high oestrogen containing COC, e.g. 50 µg EO + 250 µg LNG (Nordiol) - two pills initially, then repeated 12 hours later. Failure rate: 2.6%.2
- Danazol 200 mg tablets (e.g. two initially and repeated 12 hours later).

PILL FAILURE

Causes of oral contraceptive failure include errors in administration, decreased absorption, missed pills, drug interactions and high doses of vitamin C. It is possible that the use of triphasics may be a factor. Management options include using a higher-dose pill, improved education and compliance and an alternative method.

INTRAUTERINE CONTRACEPTIVE DEVICES

IUCDs are usually small devices made of an inert material to which may be added a bioactive substance such as copper (e.g. Multiload-Cu375), or a progestogen (e.g. Mirena).[2] The mechanism of IUCDs is not well understood, but copper devices affect sperm motility and transport.

Efficacy: IUCDs give 96 - 99% protection against pregnancy.[2]

Contraindications for IUCD use:[5]

- absolute
  - known or suspected pregnancy
  - active PID
  - undiagnosed abnormal genital tract bleeding
  - previous ectopic pregnancy
  - severe uterine cavity distortion
- relative
  - menorrhagia
  - dysmenorrhoea
  - lesser uterine cavity distortion
  - very large or very small uterus (> 9.0 or < 5.5 cm)
  - anaemia
  - defective immune system
  - impaired clotting mechanism
  - valvular heart disease
  - acutely anteverted or retroverted uterus
  - increased risk of PID (multiple sex partners)

Problems associated with IUCD usage[2]

Pregnancy/ectopic pregnancy

If pregnancy occurs there is a 40 - 50% increased risk of abortion and intrauterine sepsis during the second trimester. There is an increased risk of ectopic pregnancy (up to 10 times compared with COC usage) so, if pregnancy occurs, ultrasound examination should be performed to determine the location.

Early removal of the IUCD is essential.

Pelvic inflammatory disease

There is evidence of an increased risk of PID in the first 30 days post-insertion. Prophylactic doxycycline reduces this risk.[5] As this risk is related to sexual activity and the number of partners, those at risk of STIs should avoid using IUCDs.

Extrusion, perforation of uterus and translocation

Spontaneous extrusion is greatest during the first month after insertion and the woman is not always aware of this. Perforation of the uterus occurs once in every 1000 insertions and review at 6 weeks post-insertion is essential. If translocation is proved by X-ray and pelvic ultrasound, removal is mandatory.

Bleeding

Intermenstrual bleeding may follow insertion of an IUCD for 2 - 3 months and then disappear. If menstrual loss is excessive, the device should be removed. However, the Mirena system works to reduce menstrual bleeding.

Pain

Lower abdominal cramp-like pains of uterine origin and backache may occur soon after insertion and persist intermittently for several weeks. Rarely is the pain severe enough to warrant removal of the IUCD.

Checking the IUCD

Women should be taught how to examine themselves vaginally to check if the device remains in situ by palpating the strings or threads which protrude from the cervical canal. They should have a medical check 2 - 3 months after the device has been fitted and again after 12 months.

VAGINAL RING

The first available contraceptive vaginal ring is NuvaRing®, a flexible polymear ring with 15 μg ethinyloestradiol and 120 μg etonogestrel. It is inserted into the vagina once a month (in the first 5 days after a period) and removed after 21 days with a break of 7 days.

BARRIER METHODS

Barrier methods include condoms, vaginal diaphragms, cervical caps and vaginal vault caps. If used correctly, some, particularly condoms, are very effective contraceptives, with pregnancy rates of 5 or less per 100 women years.[2]

Condoms are also very effective in preventing the spread of STIs, including HIV infection.

*Diaphragms*have to be individually fitted. After being liberally coated on both sides with a spermicidal cream they are inserted at any convenient time before intercourse and removed after 6 hours have elapsed since the last act of intercourse.

SPERMICIDES

These are useful adjuncts to barrier methods of contraception. When used alone they have a pregnancy rate of less than 10 per 100 women years. They are available as creams, jellies, foams or pessaries containing nonoxynol 9 or octoxinol.

NATURAL METHODS

These methods require high motivation and regular menstrual cycles.

Basal body temperature method

Coitus should only occur after there has been a rise in basal body temperature of 0.2°C for 3 days (72 hours) above the basal body temperature measurement during the preceding 6 days, until the onset of the next menstrual period.

Billings ovulation method [4]

This method is based on careful observation of the nature of the mucus so that ovulation can be recognised. Fertile mucus is wet, clear, stringy, increased in amount and feels lubricative. The peak mucus day is the last day with this oestrogenised mucus before the abrupt change to thick tacky mucus associated with the secretion of progesterone. The infertile phase begins on the fourth day after the peak mucus day. Abstinence from intercourse is practised from the first awareness of increased, clearer wet mucus until 4 days after maximum mucus secretion. If taught correctly and followed as directed, the method is most effective, with a failure rate of only 1 - 2 (average 3) per 100 women years.[4] There is a failure rate of at least 15 if the rules are not followed properly.

Coitus interruptus

Male withdrawal before ejaculation is still a widely used method of contraception and despite theoretical objections will probably continue to have a definite place in contraceptive practice.

STERILISATION

Vasectomy

With vasectomy it is important to confirm the absence of spermatozoa in the ejaculate 2 - 3 months after the operation, before ceasing other contraceptive methods. It takes about 12 - 15 ejaculations to clear all the sperm from the tubes proximal to the surgical division. Vasectomy reversal is successful in up to 80% of patients.[2]There is a 1 in 500 - 1000 chance of recanalisation.

Tubal ligation

Female sterilisation is usually performed by minilaparotomy or laparoscopy, at which time clips (Filshie or Hulka) or rings (Falope) are applied to each Fallopian tube. These are potentially reversible methods of contraception with a 50 - 70% success rate of reversal.[2] There is a subsequent pregnancy rate of 3 - 4 per 1000 women sterilised.

The Essure procedure

This procedure for permanent female birth control involves the placement of a flexible titanium microinsert into each fallopian tube with a hysteroscope. The insert expands and over time (usually 3 months) reactive tissue growth occludes the tubes.

REFERENCES

1. Sloane PD, Slatt ML, Baker RM. Essentials of Family Medicine. Baltimore: Williams & Wilkins, 1988: 175 - 89.
2. Walters W. Fertility control. In: MIMS Disease Index. Sydney: IMS Publishing, 1991 - 92: 185 - 90.
3. Stovall TG. Clinical Manual of Gynaecology. New York: McGraw-Hill, 1992: 263 - 6.
4. Billings E, Westmore A. The Billings Method. Melbourne: Anne O'Donovan, 1992: 11 - 49.
5. O'Connor V, Kovacs G. Obstetrics, Gynaecology and Women's Health. Cambridge: Cambridge University Press, 2003: 395 - 413.
6. Benn M. Prescribing oral contraceptives and the medical record. Aust Fam Physician, 1991; 20: 1469 - 70.
7. Weisberg E. Choosing an oral contraceptive. Mod Med Aust, 1997; 40(1): 18 - 26.
8. Shenfield G (Chair). Therapeutic Guidelines: Endocrinology (Version 3). Melbourne: Therapeutic Guidelines Ltd, 2004: 127 - 44.
9. Miller C. The combined oral contraceptive: a practical guide. Aust Fam Physician, 1990; 19: 897 - 905.

CHAPTER 90. BREAST PAIN (MASTALGIA)

Many women suffer breast pain so severe that it affects their lifestyles, marriages and sexual relationships, and even prevents them from hugging their children.

DR JOHN DAWSON (1990)

BREAST PAIN (MASTALGIA)

Breast pain, or mastalgia, is a common problem accounting for at least 50% of breast problems presenting in general practice.[1] Many women suffer breast pain so severe that it affects their lifestyles, marriages and sexual relationships, and even prevents them from hugging their children. If no obvious physical cause is found, the problem is all too often dismissed, without appropriate empathy and reassurance, as a normal physiological effect.

A careful, sympathetic clinical approach, however, followed by reassurance after examination, will be sufficient treatment for most patients.

SYMPTOMS

Mastalgia usually presents as a heaviness or discomfort in the breast or as a pricking or stabbing sensation. The pain may radiate down the inner arm when the patient is carrying heavy objects or when the arm is in constant use, as in scrubbing floors.

KEY FACTS AND CHECKPOINTS

- The typical age span for mastalgia is 30-50 years.
- The peak incidence is 35-45 years.
- There are four common clinical presentations:
    1. diffuse, bilateral cyclical mastalgia
    2. diffuse, bilateral non-cyclical mastalgia
    3. unilateral diffuse non-cyclical mastalgia
    4. localised breast pain
- The specific type of mastalgia should be identified.
- The commonest type is cyclical mastalgia.
- Premenstrual mastalgia (part of type 1) is common.
- An underlying malignancy should be excluded.
- Less than 10% of breast cancers present with localised pain.
- Only about 1 in 200 women with mastalgia are found to have breast cancer.
- The problems, especially types 2 and 3, are difficult to alleviate.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 90.1.

Table 90.1 Mastalgia: diagnostic strategy model

Q. Probability diagnosis
A. Pregnancy Cyclical mastalgia

- benign mammary dysplasia

Q. Serious disorders not to be missed

A. Neoplasia

Infection

- mastitis
- abscess

Myocardial ischaemia

Q. Pitfalls (often missed)

A. Pregnancy

Chest wall pain e.g. costochondritis

Costochondritis

Pectoralis muscle spasm

Referred pain, esp. thoracic spine

Mechanical

- bra problems
- weight change
- trauma

Rarities

Hyperprolactinaemia

Nerve entrapment

Mammary duct ectasia

Sclerosing adenosis

Ankylosing spondylitis

Q. Seven masquerades checklist

A. Depression ✔

Diabetes -

Drugs ✔

Anaemia -

Thyroid disorder -

Spinal dysfunction ✔

UTI -

Q. Is the patient trying to tell me something?

A. Yes. Fear of malignancy. Consider psychogenic causes.

Probability diagnosis

In the non-pregnant patient, generalised pain which may be cyclical or non-cyclical is commonest. Typical patterns are illustrated in Figure 90.1.



Figure 90.1 Pain patterns for cyclical and non-cyclical mastalgia

Cyclical mastalgia is the commonest diffuse breast pain (see p. 969). It occurs in the latter half of the menstrual cycle, especially in the premenstrual days, and subsides with the onset of menstruation. It obviously has a hormonal basis, which may be an abnormality in prolactin secretion. The main underlying disorder is benign mammary dysplasia, also referred to as fibroadenosis, chronic mastitis, cystic hyperplasia or fibrocystic breast disease.

Non-cyclical mastalgia is also quite common and the cause is poorly understood. It may be associated with duct ectasia and periductal mastitis (see p. 970).

Serious disorders not to be missed

The three important serious disorders not to be missed with any painful chest condition - neoplasia, infection and myocardial ischaemia - are applicable for breast pain.

Neoplasia

We must avoid the trap of considering that breast pain is not compatible with malignancy. Mastalgia can be a presenting symptom (although uncommon) of breast cancer. 'Mastitis carcinomatosa', which is a rare florid form of breast cancer found in young women, often during lactation, is red and hot but not invariably painful or tender.[2] Pain may also be a symptom in juvenile fibroadenoma, a soft rapidly growing tumour in adolescents, and in the fibroadenoma of adult women.

Infection

Mastitis is common among nursing mothers. It should be regarded as a serious and urgent problem because a breast abscess can develop quickly. Apart from bacterial infection, infection with *Candida albicans* may occur following the use of antibiotics. *Candida* infection usually causes severe breast pain, producing a feeling like 'hot cords', especially during and after feeding.

Myocardial ischaemia

A constricting pain under the left breast should be regarded as myocardial ischaemia until proved otherwise.

Pitfalls

These include various causes of apparent mastalgia, such as several musculoskeletal chest wall conditions and referred pain from organs such as the heart, oesophagus, lungs and gall bladder and, in particular, from the upper thoracic spine.

Musculoskeletal conditions include costochondritis, pectoralis muscle strains or spasm, and entrapment of the lateral cutaneous branch of the third intercostal nerve. Ankylosing spondylitis can affect the chest wall under the breasts. Mastalgia may be the first symptom of pregnancy. Pregnancy should be excluded before commencing drug treatment.

Seven masquerades checklist

Of these, depression, drugs and spinal dysfunction are probable causes. Drugs that can cause breast discomfort include oral contraceptives, HRT and methylxanthine derivatives such as theophylline. Drugs that cause tender gynaecomastia (more applicable to men) include digoxin, cimetidine, spironolactone and marijuana.

Dysfunction of the upper thoracic spine and even the lower cervical spine can refer pain under a breast. If suspected, these areas of the spine should be examined.

Psychogenic considerations

The symptoms may be exaggerated as a result of an underlying psychogenic disorder, but with a symptom such as breast pain most women fear malignancy and need reassurance.

THE CLINICAL APPROACH

History

It is important to relate the pain to the menstrual cycle and determine whether the patient is pregnant or not.

Key questions

- Could you be pregnant?
- Is your period on time or overdue?
- Is the pain in both breasts or only one?
- Do you have pain before your periods or all the time during your menstrual cycle?
- Do you have pain in your back or where your ribs join your chest bone?

Physical examination

The breasts should be systematically palpated to check for soreness or lumps. The underlying chest wall and thoracic spine should also be examined.

Investigations

The following specialised tests could be considered.

Mammography should be considered in older women. It is unreliable in young women. With few exceptions it should not be used under 40 years.

Ultrasound can be complementary to mammography for it is useful to assess a localised mass or tender area. It is inappropriate to evaluate a diffuse area. It is not so useful for the postmenopausal breast, which is fatty and looks similar to cancer on ultrasound.

Excision biopsy can be useful for an area of localised pain, especially in the presence of a possible mass.

MASTALGIA IN CHILDREN

Breast pain is uncommon in children, including puberty, but it may be a presenting problem in the late teens. Pubertal boys may complain of breast lumps under the nipple (adolescent gynaecomastia) but these are rarely tender and do not require specific treatment.

MASTALGIA IN THE ELDERLY

Breast pain is rare after the menopause but is increasing with increased use of HRT, where it tends to present as the diffuse bilateral type. If the problem is related to the introduction of HRT, the oestrogen dose should be reduced or an alternative preparation used.

CYCLICAL MASTALGIA

The features of cyclical mastalgia are:

- the typical age is 35 years
- discomfort and sometimes pain are present
- usually bilateral but one breast can dominate
- mainly premenstrual
- breasts diffusely nodular or lumpy
- variable relationship to the pill

It is rare after the menopause.

Management

After excluding a diagnosis of carcinoma and aspirating palpable cysts, various treatments are possible and can be given according to severity.[3]

Mild

- Reassurance
- Regular review and breast self-examination
- Proper brassiere support
- Proper low-fat diet, excluding caffeine
- Aim at ideal weight
- Adjust oral contraception or HRT (if applicable)
- Analgesia (e.g. paracetamol)

Moderate

As for mild, plus options (use one or a combination):

- mefenamic acid 500 mg, three times daily
- vitamin B_1 (thiamine) 100 mg daily, and
- vitamin B_6 (pyridoxine) 100 mg daily
- consider ceasing OCP

If no response

As for mild, plus options (one of the following):

- norethisterone 5 mg daily (for second half of cycle)
- danazol 200 mg daily

Some of these treatments, particularly vitamin therapy, have not been scientifically tested but some empirical evidence is favourable. The value of diuretics is not proven, and testosterone or tamoxifen treatment is generally not favoured.

Evening primrose oil contains an essential fatty acid claimed to be lacking in the diet, and replacement allows for the production of prostaglandin E, which counters the effect of oestrogen and prolactin on the breast. However, it is no more effective than placebo.

Bromocriptine and danazol have significant sideeffects but clinical trials have proved their efficacy for this condition.[4, 5]

Systemic reviews from RTCs provide limited evidence to alleviate mastalgia but the suggestions indicate that tamoxifen and a low-fat, high-carbohydrate diet is beneficial. Danazol provides benefit but has a high incidence of side-effects. Bromocriptine (also high adverse effect profile) and HRT are unlikely to be beneficial. The following have unknown effectiveness - evening primrose oil, pyridoxine, vitamin E and diuretics.[6]

A summary of a treatment strategy for cyclical mastalgia is presented in Table 90.2.

Table 90.2 Management plan for cyclical mastalgia

Progressive stepwise therapy	
Step 1	Reassurance
	Proper brassiere support
	Diet - exclude/reduce caffeine, low fat
	Exercises (e.g. aerobics for upper trunk)
	Analgesics (on days of pain)
Step 2	Add (as a trial)
	Vitamin B_1 100 mg daily
	Vitamin B_6 100 mg daily
Step 3	Add
	Danazol 200 mg daily

NON-CYCLICAL MASTALGIA

The features of non-cyclical mastalgia are:

- the typical age is the early forties
- bilateral and diffuse
- pain present throughout the cycle
- no obvious physical or pathological basis

Typical pain patterns are presented in Figure 90.1.

Management

Non-cyclical mastalgia is very difficult to treat, being less responsive than cyclical mastalgia. It is worth a therapeutic trial.

First-line treatment

- Exclude caffeine from diet
- Weight reduction if needed
- Vitamin B_1 100 mg daily
- Vitamin B_6 100 mg daily

Second-line treatment

- Norethisterone 5 mg daily

LOCAL LESIONS

Surgical excision may be required for local lesions. If there is no discrete lesion but a tender trigger point (including costochondritis), the injection of local anaesthetic and corticosteroid may relieve the problem.

COSTOCHONDRITIS (TIETZE'S SYNDROME)

This is a common cause of referral to a breast pain clinic. The cause is often obscure, but the costochondral junction may become strained in patients with a persistent cough. The pain can appear to be in the breast with intermittent radiation round the chest wall and is initiated or aggravated by deep breathing and coughing.

Features:

- the pain is acute, intermittent or chronic
- the breast is normal to palpation
- palpable swelling about 4 cm from sternal edge due to enlargement of costochondral cartilage
- X-rays are normal
- self-limiting, but may take several months to subside

Treatment. Infiltration with local anaesthetic and corticosteroid with care.

MASTITIS

Mastitis is basically cellulitis of the interlobular connective tissue of the breast. Mostly restricted to lactating women, it is associated with a cracked nipple or poor milk drainage. The infecting organism is usually *Staphylococcus aureus*, or more rarely, *Escherichia coli* or *C. albicans*. It is a serious problem and requires early treatment. Breastfeeding from the affected side can continue as the infection is confined to interstitial breast tissue and doesn't usually aff ect the milk supply.

Clinical features:

- a lump and then soreness (atfirst)
- a red tender area possibly
- fever, tiredness, muscle aches and pains

Note: Candida infection usually causes severe breast pain - a feeling like a hot knife or hot shooting pains, especially during and after feeding. It may occur after a course of antibiotics.

Management

Prevention (in lactation):

- maintain free breast drainage
- attend to breast engorgement and cracked nipples

Treatment

If systemic symptoms develop:

- Antibiotics: resolution without progression to an abscess will usually be prevented by antibiotics[7] di/flucloxacillin 500 mg (o) 6 hourly for 7-10 days *or* cephalexin 500 mg (o) 6 hourly for 7-10 days If severe cellulitis di/flucloxacillin 2 g (IV) 6 hourly
- Therapeutic ultrasound (2 W/cm2 for 6 minutes) daily for 2-3 days
- Aspirin or paracetamol for pain
- For *Candida albicans* infection fluconazole 200-400 mg (o) daily for 2-4 weeks second line - nystatin 500 000 U (o) tds

Instructions to patients

- Keep the affected breast well drained.
- Continue breastfeeding: do frequently and start with the sore side.

- Heat the sore breast before feeding (e.g. with hot shower or hot face washer).
- Cool the breast after feeding: use a cold face washer from the freezer.
- Massage any breast lumps gently towards the nipple while feeding.
- Empty the breast well: hand express if necessary.
- Get sufficient rest.
- Keep to a nutritious diet and drink ample fluids.

BREAST ABSCESS

If tenderness and redness persist beyond 48 hours and an area of tense induration develops, then a breast abscess has formed. It requires surgical drainage under general anaesthesia or aspiration with a large bore needle under local anaesthetic every second day until resolution, antibiotics, rest and complete emptying of the breast. Temporary weaning of breastfeeding from the affected side is necessary because of the surgical disruption.

Surgical drainage

Method

1. Make an incision over the point of maximal tenderness, preferably in a dependent area of the breast. A curvilinear transverse incision which does not continue into breast tissue, following Langer's lines, is optimal (Figure 90.2a).
2. Use artery forceps to separate breast tissue to reach the pus.
3. Take a swab for culture.
4. Introduce a gloved finger to break down the septa that separate the cavity into loculations (Figure 90.2b).
5. Insert a corrugated drainage tube into the cavity. Fix it to the skinedge with a single suture (Figure 90.2c).

Remove the tube two days after the operation. Change the dressings daily until the wound has healed. Continue antibiotics until resolution of the inflammation.



Figure 90.2 Surgical drainage of breast abscess: **(a)** transverse incision; **(b)** exploring abscess cavity; **(c)** drainage

tube in situ

WHEN TO REFER

- Undiagnosed localised breast pain or lump.

Practice tips

- The basis of management for benign mastalgia is firm reassurance.
- Although breast cancer rarely causes mastalgia, it should be excluded.
- Think of *C. albicans* if mastitis is very severe with hot shooting pains, especially after antibiotic treatment.
- Look for underlying disorders of the chest wall if examination of the breasts is normal.
- Consider caffeine intake as a cause of benign diff use mastalgia.
- Mastitis should be treated vigorously - it is a serious condition.
- Fibroadenomas and breast cysts are capable of causing localised pain and tenderness.

REFERENCES

1. McAvoy PA. Breast problems. In: Cormack J, Marinker M, Morrell D eds. *Practice.* London: Kluwer Medical, 1982: Section 3.54.
2. Ryan P. *A Very Short Textbook of Surgery* (2nd edn). Canberra: Dennis & Ryan, 1990: 10.
3. Barraclough B. The fibrocystic breast - clinical assessment, diagnosis and treatment. Mod Med Aust, 1990; 33(4): 16 - 25.
4. Mansel RE et al. Controlled trial of the antigonadotrophin danazol in painful nodular benign breast disease. Lancet, 1982; 1: 928.
5. Hinton CP et al. A double blind controlled trial of danazol and bromocriptine in the management of severe cyclical breast pain. Br J Clin Pract, 1986; 40: 326.
6. Barton S ed. *Clinical Evidence*. London: BMJ Publishing Group, 2001: 1247 - 52.
7. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 282.

CHAPTER 91. LUMPS IN THE BREAST

Neither the cause of breast cancer, one of the most feared and emotion-engendering diseases,
nor the means of preventing it are absolutely known.

Lecturer on breast cancer (Anon)

LUMPS IN THE BREAST

Breast lumps are common and their discovery by a woman provokes considerable anxiety and emotion (which is often masked during presentation) because to many a ‘breast lump’ means cancer. Many of the lumps are actually areas of thickening of normal breast tissue. Many other lumps are due to mammary dysplasia with either fibrosis or cyst formation or a combination of the two producing a dominant (discrete) lump.[1] However, a good working rule is to consider any lump in the breast as carcinoma until proved otherwise. See Table 91.1 for causes of breast lumps in a specific outpatient study.

Table 91.1 Causes of breast lumps (a surgical outpatient study)

Common	%
• Mammary dysplasia	32
• Fibroadenoma	23
• Carcinoma	22
• Cysts	10
• Breast abscess/periareolar inflammation	2
Less common	
• Mammary duct ectasia	
• Duct papilloma	
• Lactation cysts (galactocele)	
• Paget's syndrome of the nipple	
• Fat necrosis/fibrosis	
• Sarcoma	
• Lipoma	

Statistics courtesy MA Henderson, PBR Kitchen, PR Hayes, University of Melbourne Department of Surgery, Breast Clinic, St Vincent's Hospital, Melbourne

The genetic predisposition to breast cancer continues to be delineated with the strong predisposition from mutations in the genes *BRCA1* and *BRCA2*. Refer to Chapter 19.

KEY FACTS AND CHECKPOINTS

- The commonest lumps are those associated with mammary dysplasia (32%).[2] See Table 91.1.
- Mammary dysplasia is also a common cause of cysts, especially in the premenopause phase.
- Over 75% of isolated breast lumps prove to be benign but clinical identification of a malignant tumour can only definitely be made following aspiration biopsy or histological examination of the tumour.[2]
- The investigation of a new breast lump requires a very careful history and the triple test.
- Breast cancer is the most common cancer in females, affecting 1 in 12 - 15 women.[2]
- About 25% of all new cancers in women are breast neoplasms.

- A ‘dominant’ breast lump in an older woman should be regarded as malignant.

The triple test

1. Clinical examination
2. Imaging - mammography ± ultrasound
3. Fine needle aspiration ± core biopsy

THE CLINICAL APPROACH

This is based on following a careful history and examination.

History

The history should include a family history of breast disease and the patient's past history, including trauma, previous breast pain, and details about pregnancies (complications of lactation such as mastitis, nipple problems and milk retention).

Key questions[1]

- Have you had any previous problems with your breasts?
- Have you noticed any breast pain or discomfort?
- Do you have any problems such as increased swelling or tenderness before your periods?
- Have you noticed lumpiness in your breasts before?
- Has the lumpy area been red or hot?
- Have you noticed any discharge from your nipple or nipples?
- Has there been any change in your nipples?
- Did your mother or sisters or any close relatives have any breast problems?

Breast symptoms

- Lump (76%)
- Tenderness or pain (10%)
- Nipple changes (8%)
- Nipple discharge (2%)
- Breast asymmetry or skin dimpling (4%)
- Periareolar inflammation

Important ‘tell-tale’ symptoms are illustrated in Figure 91.1.


lump
skin dimpling
peau d'orange
nipple inversion
bloodstained nipple discharge
eczema of nipple (Paget's sign)

Figure 91.1 Important 'tell-tale' symptoms of carcinoma of the breast

Nipple discharge[3]

This may be intermittent from one or both nipples. It can be induced by quadrant compression.

- bloodstained
  - intraduct papilloma (commonest)
  - intraduct carcinoma
  - mammary dysplasia
- green-grey
  - mammary dysplasia
  - mammary duct ectasia
- yellow
  - mammary dysplasia
  - intraduct carcinoma (serous)
  - breast abscess (pus)
- milky white (galactorrhoea)
  - lactation cysts
  - lactation
  - hyperprolactinaemia
  - drugs (e.g. chlorpromazine)

Periareolar inflammation

This presents as pain around the areola with reddening of the skin, tenderness and swelling. Causes may be inverted nipple or mammary duct ectasia.

Paget's syndrome of the nipple

This rare but interesting sign and condition usually occurs in middle-aged and elderly women. It starts as an eczematous-looking, dry scabbing red rash of the nipple and then proceeds to ulceration of the nipple and areola (Table 91.2). It is always due to an underlying malignancy.

Table 91.2 Differences between Paget's syndrome and eczema of the nipple[2]

Paget's syndrome	Eczema

Unilateral	Bilateral
Older patients	Reproductive years/lactation
Possible nipple discharge	No discharge
Not pruritic	Pruritic
No pustules	Pustules
Deformity of nipple	Normal nipple
Possible palpable lump	No lump

EXAMINATION OF THE BREASTS

Objectives

- Identify a dominant lump (one that differs from the remainder of the breast tissue).
- Identify a lump that may be malignant.
- Screen the breasts for early development of carcinoma.

Time of examination: ideally, 4 days after the end of the period.

Method[1]

1. Inspection: sitting - patient seated upright on side of couch in good light, arms by sides, facing the doctor, undressed to waist.
    a. Note
        - Asymmetry of breasts or a visible lump
        - Localised discolouration of the skin
        - Nipples
            - for retraction or ulceration
            - for variations in the level (e.g. elevation on one side)
            - for discharge (e.g. blood-stained, clear, yellow)
        - Skin attachment or tethering →sdimpling of skin (accentuate this sign by asking patient to raise her arms above her head)
        - Appearance of small nodules of growth
        - Visible veins (if unilateral they suggest a cancer)[4]
        - Peau d'orange due to dermal oedema

    b. Raise arms above the head (renders variations in nipple level and skin tethering more obvious). Hands are pressed on the hips to contract pectoralis major to note if there is a deep attachment of the lump.
2. Examination of lymph glands in sitting position: patient with hands on hips. Examine axillary and supraclavicular glands from behind and front.
   Note: The draining lymphatic nodes are in the axillae, supraclavicular fossae and internal mammary chain.
3. Palpation
    a. Patient still seated: palpate breast with flat of hand and then palpate the bulk of the breast between both hands.
    b. In supine position:
        - patient lies supine on couch with arms above head
        - turn body (slight rotation) towards midline so breasts ‘sit’ as flat as possible on chest wall

    Method

    - Use the pulps of the fingers rather than the tips with the hand laid flat on the breast.

- Move the hand in slow circular movements.
- Examine up and down the breast in vertical strips beginning from the axillary tail (Figure 91.2).



Figure 91.2 Systematic examination of the breast

- Systematically cover the six areas of the breast (Figure 91.3)


Right breast
axillary tail
6
upper outer
2
upper inner
1
5
lower outer
4
lower inner
3

Figure 91.3 The six areas of the breast

  - the four quadrants
  - the axillary tail
  - the region deep to the nipple and areola

4. If a suspicious lump is present, inspect liver, lungs and spine.
5. Inspect the bra. Note possible pressure on breast tissue from underwiring of the bra, usually on the upper outer quadrant.

If a solitary lump is present, assess it for:

- position (breast quadrant and proximity to nipple)
- size and shape
- consistency (firm, hard, cystic, soft)
- tenderness
- mobility and fixation
- attachment to skin or underlying muscle

Note:

- 40 - 50% of carcinomas occur in the upper outer quadrant.[3]
- A hard mass is suspicious of malignancy but cancer can be soft because of fat entrapment.
- The inframammary ridge, which is usually found in the heavier breast, is oft en nodular and firm to hard.
- Lumpiness (if present) is usually most marked in the upper outer quadrant.
- A useful diagram to record the findings is shown in Figure 91.4.
- Lumps that are usually benign and require no immediate action are: tiny (< 4 mm) nodules in subcutaneous tissue (usually in the areolar margin); elongated ridges, usually bilateral and in the lower aspects of the breasts; and rounded soft nodules (usually < 6 mm) around the areolar margin.[5]



Figure 91.4 Diagrammatic scheme for recording the features of breast lumps and any lymphadenopathy (axilla and supraclavicular triangles)

INVESTIGATIONS

X-ray mammography

Mammography can be used as a screening procedure and as a diagnostic procedure. It is currently the only effective screening tool for breast cancer.[6] Positive signs of malignancy include an irregular infiltrating mass with focal spotty microcalcification.

Screening:

- established benefit for women over 50 years
- possible benefit for women in their 40s

- follow-up in those with breast cancer, as 6% develop in the opposite breast
- localisation of the lesion for fine needle aspiration

Breast ultrasound

This is mainly used to elucidate an area of breast density and is the best method of defining benign breast disease, especially with cystic changes. It is generally most useful in women less than 35 years old (as compared with X-ray mammography).

Useful for:

- pregnant and lactating breast
- differentiating between fluid-filled cysts and solid mass
- palpable masses at periphery of breast tissue (not screened by mammography)
- for more accurate localisation of lump during fine needle aspiration

Note: CT and MRI have limited use. An age-related schemata for likely diagnosis and appropriate investigations is presented in Table 91.3.

Table 91.3 Age-related schemata for likely diagnoses and appropriate investigations (after Hirst)[5]

1 Very young women - 12 to 25 years

• inflamed cysts or ducts, usually close to areola

• fibroadenomata, often giant

• hormonal thickening, not uncommon

• malignancy rare

Investigations - mammography contraindicated
- ultrasound helpful

2 Young women - 26 to 35 years

• classic fibroadenomata

• mammary dysplasia with or without discharge

• cysts uncommon

• malignancy uncommon

Investigations - mammography: breasts often very dense
- ultrasound often diagnostic

3 Women - 36 to 50 years (premenopausal)

• cysts

• mammary dysplasia, discharges, duct papillomas

• malignancy common

• fibroadenomata occur but cannot assume

• inflammatory processes not uncommon

Investigations - mammography useful
- ultrasound useful

4 Women - over 50 years (postmenopausal)

• any new discrete mass - malignant until proven otherwise

• any new thickening - regard with suspicion

• inflammatory lesions - probably duct ectasia (follow to resolution)

• cysts unlikely

Investigations - mammography usually diagnostic
- ultrasound may be useful

5 Women - over 50 years, on hormones

• any new mass - regard with suspicion

• cysts may occur - usually asymptomatic

• hormonal change not uncommon

Investigations - mammography usually diagnostic but breast may become more dense
- ultrasound may be useful

Reprinted with permission

Needle aspiration and biopsy techniques

- Cyst aspiration
- Fine needle aspiration biopsy: this is a very useful diagnostic test in solid lumps, and has an accuracy of 90 - 95% (better than mammography)[3]
- Large needle (core needle) biopsy
- Incision biopsy

Tumour markers

Oestrogen receptors are uncommon in normal breasts but are found in two-thirds of breast cancers, although the incidence varies with age. They are good prognostic indicators. Progesterone receptors can also be estimated.

Fine needle aspiration of breast lump

This simple technique is very useful, especially if the lump is a cyst, and will have no adverse effects if the lump is not malignant. If it is, the needle biopsy will help with the preoperative cytological diagnosis.

Method of aspiration and needle biopsy:

1. Use an aqueous skin preparation without local anaesthesia.
2. Use a 23-gauge needle and 5 mL sterile syringe.
3. Identify the mass accurately and fix it by placing three fingers of the non-dominant hand firmly on the three sides of the mass (Figure 91.5).
4. Introduce the needle directly into the area of the swelling. Once in subcutaneous tissue, apply gentle suction as the needle is being advanced (Figure 91.6). If a cyst is involved it can be felt to ‘give’ suddenly.
5. If fluid is obtained (usually yellowish-green), aspirate as much as possible.
6. If no fluid is obtained, try to get a core of cells from several areas of the lump in the bore of the needle.
7. Make several passes through the lump at different angles, without exit from the skin and maintain suction.
8. Release suction before exit from the skin to keep cells in the needle (not in the syringe).
9. After withdrawal, remove syringe from needle, fill with 2 mL of air, reattach needle and produce a fine spray on one or two prepared slides.
10. Fix to one slide (in Cytofix) and allow one to air dry, and forward to a reputable pathology laboratory to be examined by a skilled cytologist.


third finger
fourth finger
direction of syringe
second finger

Figure 91.5 Aspiration of breast lump: fixation of cyst


third finger
second finger
thumb

Figure 91.6 Aspiration of breast lump: position of the hand - second (index) finger and thumb steady the syringe while the third (middle) finger slides out the plunger to create suction

Follow-up: the plan for aspiration is outlined in Figure 91.7.



Figure 91.7 Scheme for management of a breast lump by fine needle aspiration

Indications for biopsy or excision of lump

- The cyst fluid is bloodstained.
- The lump does not disappear completely with aspiration.
- The swelling recurs within 1 month.

Summary: investigation of a breast lump

If the patient presenting with a breast lump is younger than 35, perform an ultrasound;[7] if older than 35 perform a mammogram and an ultrasound. If the lump is cystic - aspirate; if solid - perform a fine needle biopsy and then manage according to outcomes. If it is suspicious, an excisional biopsy is the preferred option.

CARCINOMA OF THE BREAST

Breast carcinoma is uncommon under the age of 30 but it then steadily increases to a maximum at the age of about 60 years.[3] About one-third are premenopausal and two-thirds postmenopausal. About 1 in 11 - 15 women (1 in 11 in Australia) develop breast cancer. Ninety per cent of breast cancers are invasive duct carcinomas, the remainder being lobular carcinoma, papillary carcinomas, medullary carcinomas and colloid or mucoid carcinomas.[2]

Risk factors include increasing age (> 40 years), Caucasian race, pre-existing benign breast lumps, alcohol, HRT > 5 years, personal history of breast cancer, family history in a first-degree relative (raises risk about threefold), nulliparity, late menopause (after 53), obesity, childless until after 30 years of age, early menarche,[6] ionising radiation.

Familial breast cancer

Up to 5% of cases are familial, with most being autosomal dominant. Refer to Chapter 19.

Clinical features

- The majority of patients with breast cancer present with a lump (Figure 91.8).
- The lump is usually painless (16% associated with pain).

- Usually the lump is hard and irregular.
- Nipple changes, discharge, retraction or distortion.
- Rarely cancer can present with Paget's syndrome (nipple eczema).
- Rarely it can present with bony secondaries (e.g. back pain, dyspnoea, weight loss, headache).

Note: There are basically two presentations of the disease:

- the vast majority present with a local breast lump[2] (see Figure 91.9)
- some present with metastatic disease



Figure 91.8 Advanced adenocarcenoma of breast in patient showing denial for a problem of 2 years duration. *Photo courtesy Dr. Victor Vanco*



Figure 91.9 Relative frequencies of breast cancer at various anatomical segments

Of those who present with local disease, approximately 50% will develop metastatic disease.

Management

Immediate referral to an expert surgeon on suspicion or proof of breast cancer is essential. The treatment has to be individualised according to the nature of the lump, age of the patient and staging. Accurate staging requires knowledge of whether the draining lymph nodes are involved with the tumour, as this is the single most powerful predictor of subsequent metastases and death. Staging for systemic disease also requires full blood examination and liver function tests (including alkaline phosphatase). A bone scan may be used as a valuable baseline. Size and histological grading of tumour plus nodal status and receptor status are the most important prognostic factors.

Optimal management of locally advanced breast cancer is a combined approach that uses chemotherapy, radiotherapy, surgery and/or endocrine therapy if applicable (level IV evidence).

Most relapses[8] after surgery occur in the first 3 years.

Adjuvant therapy for breast cancer

The consultant will choose the most appropriate surgical and adjuvant treatments for the individual patient.

The National Breast Cancer Consensus report emphasised that ‘continuing care should be coordinated through the patient's GP as the impact of treatment may last longer than therapy and support must continue’. The report made the following recommendations:[9]

- Tumour excision followed by whole breast irradiation was the most preferred local therapy for most women with stage I or II cancer.
- Total mastectomy and breast-conservation surgery had an equivalent effect on survival.
- Total mastectomy is preferred for a large tumour, multifocal disease, previous irradiation and extensive tumour on mammography.
- Current recommendations for radiotherapy after mastectomy are:[10]
    - tumours > 5 cm in diameter
    - axillary node involvement of > 3 nodes
    - the presence of positive or close tumour margins
- Adjuvant hormonal therapy by the antioestrogen agent tamoxifen 20 mg (o) daily which is a specific modulating agent, is widely used and is most suitable in postmenopausal women.

Adjunct agents available for treatment include:[11]

- anti-oestrogens: tamoxifen, toremifene
- aromatase inhibitors: anastrozole, letrazole, exemestane
- monoclonal antibodies: trastuzumab (Herceptin)
- progesterones, e.g. medroxyprogesterone acetate

Guidelines for adjuvant treatments are presented in Table 91.4.

Table 91.4 Adjuvant treatment favoured by trial meta-analyses[9]

Menopausal status	Node status	Oestrogen receptor	Other factors	Treatment
Premenopausal	Positive	Positive		Chemotherapy[†] + tamoxifen
	Positive	Negative		Chemotherapy
	Negative	Positive	Poor prognosis[*]	Chemotherapy and/or tamoxifen
	Negative	Negative	Poor prognosis	Chemotherapy
Postmenopausal Positive Positive	Positive	Negative	> 60 years	Tamoxifen

		Positive	< 60 years	Chemotherapy + tamoxifen
		Negative	< 60 years	Chemotherapy
Postmenopausal	Negative	Positive		Tamoxifen
		Negative	< 60, poor prognosis	Chemotherapy
		Negative	> 60	?

[*] Poor prognostic features are defined as tumours > 20 mm, or tumours 11 - 20 mm with additional poor prognostic features such as oestrogen and progesterone receptor negativity, or high histological grade. For patients with good prognostic features (e.g. those with tumours < 10 mm diameter) there has been no demonstrated benefit of adjuvant therapy.

[†] Chemotherapy is usually a combination of CMF (cyclophosphamide, methotrexate and fluorouracil) for 6 cycles. Adapted from NH & MRC Recommendations for the selection of adjuvant systemic therapy after surgical treatment for breast cancer. Clinical Practice Guidelines 1995. Canberra: NH & MRC 1995.

MAMMARY DYSPLASIA

Synonyms: fibroadenosis, chronic mastitis, fibrocystic disease, cystic hyperplasia.

Features

- Most common in women between 30 and 50 years
- Hormone-related (between menarche and menopause)
- Pain and tenderness and swelling
- Premenstrual discomfort or pain and increased swelling
- Fluctuation in the size of the mass
- Usually settles after the period
- Unilateral or bilateral
- Nodularity ± a discrete mass
- Ache may extend down inner aspect of upper arm
- Nipple discharge may occur (various colours, mainly green-grey)
- Most cysts are premenopausal (final 5 years before menopause)

Examination. Look for lumpiness in one or both breasts, usually upper outer quadrant.

Management

- Consider mammography if diffuse lumpiness is present in patient > 40 years.
- Perform needle biopsy if a discrete lump is present and aspirate palpable cysts.
- Reassure patient that there is no cancer.
- Give medication to alleviate mastalgia (see treatment for cyclical mastalgia in Chapter 90).
- Use analgesics as necessary.
- Surgically remove undiagnosed mass lesions.

BREAST CYST

- Common in women aged 40 - 50 years (perimenopausal)
- Rare under 30 years
- Associated with mammary dysplasia

- Tends to regress after the menopause
- Pain and tenderness variable
- Has a 1 in 1000 incidence of cancer
- Usually lined by duct epithelium

Examination. Look for a discrete mass, firm, relatively mobile, that is rarely fluctuant.

Diagnosis

- Mammography
- Ultrasound (X-ray of choice)
- Cytology of aspirate

LACTATION CYSTS (GALACTOCELES)

- These milk-containing cysts arise during pregnancy and present postpartum with similar signs to perimenopausal cysts.
- They vary from 1 - 5 cm in diameter.
- Treat by aspiration: fluid may be clear or milky.

FIBROADENOMA

Clinical features

- A discrete asymptomatic lump
- Usually in twenties (range: second to sixth decade, commonly 15 - 35 years)
- Firm, smooth and mobile (the ‘breast mouse’)
- Usually rounded
- Usually in upper outer quadrant
- They double in size about every 12 months7

Management

Ultrasound and fine needle aspiration with cytology is recommended plus mammography in older women. The lump may be left in those in the late teens but as a rule it is removed to be certain of the diagnosis in all patients.

PHYLLODES TUMOUR[12]

These are giant fibroadenoma-like tumours which are usually benign but 25% are malignant and metastatise. They are completely excised with a rim of normal breast tissue.

FAT NECROSIS

Fat necrosis is usually the end result of a large bruise or trauma which may be subtle, such as protracted breastfeeding. The mass that results is often accompanied by skin or nipple retraction and thus closely resembles carcinoma. If untreated it usually disappears but the diagnosis can only be made on excision biopsy.

DUCT PAPILLOMAS

These are benign hyperplastic lesions within large mammary ducts and are not premalignant (nor usually palpable). They present with nipple bleeding or a bloodstained discharge and must be differentiated from infiltrating carcinoma. Mammography and ductography are usually of limited value. The involved duct and affected breast segment should be excised.[12]

MAMMARY DUCT ECTASIA

Synonyms: plasma cell mastitis, periductal mastitis.

In this benign condition a whole breast quadrant may be indurated and tender. The larger breast ducts are dilated. The lump is usually located near the margin of the areola and is a firm or hard, tender, poorly defined swelling. There may be a toothpaste-like nipple discharge. It is a troublesome condition with a tendency to repeated episodes of periareolar inflammation with recurrent abscesses and fistula formation. Many cases settle but often surgical intervention is necessary to make the diagnosis. The condition is most common in the decade around the menopause.

THE PROBLEM OF MAMMARY PROSTHESES[5]

Clinical examination is still necessary and fortunately the residual mammary tissue is usually spread over the prosthesis in a thin easily palpable layer. The areas of clinical difficulty lie at the margin of the prosthesis, especially in the upper outer quadrant where most of the breast tissue is displaced. It should be noted that mammography may be of limited value in the presence of prostheses, especially if a fibrous capsule exists around the prosthesis. Ultrasound examination may be helpful.

BREAST LUMPS IN CHILDREN

There are several benign conditions that can cause a breast lump in children, although the commonest presentation is a diffuse breast enlargement.

Neonatal enlargement[13]

Newborn babies of either sex can present with breast hyperplasia and secretion of breast milk. This is due to transplacental passage of lactogenic hormones. The swelling usually lasts 7 - 10 days if left alone. Any attempts to manipulate the breasts to facilitate emptying will prolong the problem.

Premature hyperplasia[13]

The usual presentation is the development of one breast in girls commonly 7 - 9 years of age but sometimes younger. The feature is a firm discoid lump 1 - 2 cm in diameter, situated deep to the nipple. The same change may follow in the other breast within 3 - 12 months. Reassurance and explanation is the management and biopsy must be avoided at all costs.

Subareolar hyperplasia in boys

A discoid subareolar lesion similar to the premature hyperplasia of girls can occur in boys in one or both breasts at about 12 - 14 years. No specific treatment is necessary, simply explanation and reassurance.

Gynaecomastia

This is not to be confused with pseudogynaecomastia due to fat in obese preadolescents. However, gynaecomastia in thin boys does occur and requires referral for assessment if it cannot be attributed to drugs such as oestrogen. If no cause is found, simple mastectomy may be performed.

BREAST LUMPS IN MEN

Virtually no breast tissue is palpable in normal men and malignancy is rare.

Gynaecomastia

This is a ‘true’ enlargement of the male breasts, not to be confused with false enlargement of obese men. Gynaecomastia occurs in up to 50% of adolescent boys.[4]

If present in adult men, look for evidence of hypogonadal states such as Klinefelter's syndrome and secondary testicular failure (e.g. orchitis, orchidectomy, traumatic atrophy). Other causes include drugs (e.g. oestrogen, digoxin, calcium antagonists, marijuana, spironolactone, amiodarone, tricyclic antidepressents, cimetidine), liver failure, testicular feminisation syndrome, and oestrogen-secreting tumours, such as adrenal carcinoma and Leydig cell tumour.

COUNSELLING OF PATIENTS

‘Treat the whole woman, not merely her breasts.’ [5] Extreme anxiety is generated by the discovery of a breast lump and it is important that women are encouraged to visit their doctor early, especially as they can learn that there is a 90% chance of their lump being benign. It is possible that denial may be a factor or there is a hidden agenda to the consultation. The decision to perform a lumpectomy or a mastectomy should take the patient's feelings into consideration - many do fear that a breast remnant may be a focus for cancer. Long-standing doctor - patient relationships are the ideal basis for coping with the difficulties.

SCREENING

Screening mammography should be encouraged for women between 50 and 70 years, and performed at least every 2 years. Technically it is a better diagnostic tool in older women because of the less dense and glandular breast tissue. It has a specificity of around 90%. A management program for women at high risk of breast cancer is presented in Table 91.5.

Table 91.5 Management program for women at high risk of breast cancer (after Barraclough)[6]

- Monthly breast self-examination
- At least an annual consultation with general practitioner - if aged 40 or older
- Aspiration of cysts
- Mammography, ultrasound and/or fine needle biopsy to diagnose any localised mass
- Ultrasound alone for further assessment of young, dense breasts
- Regular screening mammography after 50 years of age - every 2 years

Breast self-examination is a controversial issue and has no proven benefit in reducing morbidity and mortality. The false positive rate is high, especially in those under 40 years. However, regular BSE is recommended for all women 35 years and over.

WHEN TO REFER

- Patients with a solitary breast mass
- Following cyst aspiration:
    - blood in aspirate

  - palpable residual lump
  - recurrence of the cyst
- Patients given antineoplastic drugs, whether for adjuvant therapy or for advanced disease, require skilled supervision

Lumps that require investigation and referral are presented in Table 91.6.

Table 91.6 Lumps that require investigation and referral (after Hirst)[5]

• A stony hard lump or area, regardless of size, history or position
• A new palpable 'anything' in a postmenopausal woman
• A persisting painless asymmetrical thickening
• An enlarging mass - cyclic or not
• A 'slow-to-resolve' or recurrent inflammation
• A bloodstained or serous nipple discharge
• Skin dimpling, of even a minor degree, or retraction of the nipple
• A new thickening or mass in the vicinity of a scar

Practice tips

- Any doubtful breast lump should be removed.
- Fibroadenomas commonly occur in women in their late teens and twenties, benign breast cysts between 35 years and the menopause, and carcinoma is the most common cause of a lump in women over 50 years.[3]
- Never assume a palpable mass is a fibroadenoma in any woman over 30 years of age.[5]
- Gentle palpation is required. Squeezing breast tissue between finger and thumb tends to produce 'pseudolumps'.
- Any eczematous rash appearing on the nipple or areola indicates underlying breast cancer.
- Mammary duct ectasia and fat necrosis can be clinically indistinguishable from carcinoma of the breast.
- 9 out of 10 women who get breast cancer do not have a strong family history.
- The oral contraceptive pill has been generally shown *not* to alter the risk of breast cancer.
- Never assume that a lump is due to trauma unless you have seen the bruising and can observe the lump decrease in size.[5]
- Never assume a lesion is a cyst - prove it with ultrasound or successful aspiration.[5]
- Never ignore skin dimpling even if no underlying mass is palpable.[5]
- Never ignore a woman's insistence that an area of her breast is different or has changed.[5]
- Mammography can detect breast cancers which are too small to feel.
- Mammography is not a diagnostic tool.
- Recommended mammography screening for women 50 - 69 years and those aged 40 - 49 who request it.

REFERENCES

1. Davis A, Bolin T, Ham J. *Symptom Analysis and Physical Diagnosis* (2nd edn). Sydney: Pergamon Press, 1990: 118.
2. Green M. Breast cancer. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 83 - 5.
3. Hunt P, Marshall V. *Clinical Problems in General Surgery.* Sydney: Butterworths, 1991: 63 - 71.
4. Talley N, O'Connor S. *Clinical Examination* (3rd edn). Sydney: MacLennan & Petty, 1996: 113 - 35.
5. Hirst C. Managing the breast lump. Solving the dilemma - reassurance versus investigation. Aust Fam

Physician, 1989; 18: 121 - 6.

6. Barraclough B. The fibrocystic breast - clinical assessment, diagnosis and treatment. Mod Med Aust, 1990; April: 16 - 25.
7. Crea P. Benign breast diseases: a management guide for GPs. Mod Med Aust, 1995; 38(8): 74 - 88.
8. NH&MRC. *Management of Advanced Breast Cancer: Clinical Practice Guidelines*. NH&MRC Clinical Practice Guidelines 2001. Canberra: NH&MRC, 2001.
9. Coates A. Breast Cancer Consensus report, Med J Aust, 1994; 161: 510 - 13.
10. Wetzig NR. How to Treat: Breast Cancer. Aust Doctor, 2001; 19 October: I - VIII.
11. Bochner F (Chair). *Australian Medicines Handbook.* Adelaide: Australian Medicines Handbook Pty Ltd, 2006: 559 - 63.
12. Burkitt H, Quick C, Gatt D. *Essential Surgery* (2nd edn). Edinburgh: Churchill Livingstone, 1996: 542.
13. Hutson JM, Beasley SW, Woodward AA. *Jones Clinical Paediatric Surgery*. Melbourne: Blackwell Scientific Publications, 1992: 266 - 7.

CHAPTER 92. ABNORMAL UTERINE BLEEDING

It is advisable that menstruation begin before the individual ceases to be a virgin.

SORAMUS OF EPHESUS (2ND CENTURY), TEXT ON DISEASES OF WOMEN

ABNORMAL UTERINE BLEEDING

Abnormal uterine bleeding is a common problem encountered in general practice. Heavy menstrual bleeding is the commonest cause of iron-deficiency anaemia in the Western world. A classification of abnormal uterine bleeding is presented in Table 92.1.

Table 92.1 Classification of abnormal uterine bleeding

Abnormal rhythm
• Irregularity of cycle
• Intermenstrual bleeding (metrorrhagia)
• Postcoital bleeding
• Postmenopausal bleeding
Abnormal amount
• Increased amount = menorrhagia
• Decreased amount = hypomenorrhoea
Combination (rhythm and amount)
• Irregular and heavy periods = metromenorrhagia
• Irregular and light periods = oligomenorrhoea

KEY FACTS AND CHECKPOINTS

- Up to 20% of women in the reproductive age group complain of increased menstrual loss.[1]
- At least 4% of consultations in general practice deal with abnormal uterine bleeding.
- Up to 50% of patients who present with perceived menorrhagia (or excessive blood loss) have a normal blood loss when investigated.[2] Their perception is unreliable.
- The possibility of pregnancy and its complications, such as ectopic pregnancy, abortion (threatened, complete or incomplete), hydatidiform mole or choriocarcinoma should be kept in mind.
- The mean blood loss in a menstrual cycle is 30 - 40 mL.
- A menstrual record is a useful way to calculate blood loss.
- Blood loss is normally less than 80 mL.
- Menorrhagia is a menstrual loss of more than 80 mL per menstruation.
- Menorrhagia disposes women to iron deficiency anaemia.
- Two common organic causes of Menorrhagia are fibroids and adenomyosis (presence of endometrium in the utrine myometrium).[3]
- Various drugs can alter menstrual bleeding (e.g. anticoagulants, cannabis, steroids).

DEFINING WHAT IS NORMAL AND WHAT IS ABNORMAL

This feature is based on a meticulous history, an understanding of the physiology and physiopathology of the menstrual cycle and a clear understanding of what is normal. Most girls reach menarche by the age of 13 (range 10 - 16 years).[1]

Dysfunctional bleeding is common in the first 2 - 3 years after menarche due to many anovulatory cycles resulting in irregular periods, heavy menses and probably dysmenorrhoea.

Once ovulation and regular menstruation are established the cycle usually follows a predictable pattern and any deviation can be considered as abnormal uterine bleeding (see Table 92.2). It is abnormal if the cycle is less than 21 days, the duration of loss is more than 8 days, or the volume of loss is such that menstrual pads of adequate absorbency cannot cope with the flow or clots.[4]

Table 92.2 Normal menstruation in the reproductive age group (after Fung)[1]

	Mean	Range
Length of cycle	26 - 28 days	21 - 35 days
Menstrual flow	3 - 4 days	2 - 7 days
Normal blood loss	30 - 40 mL	20 - 80 mL

Relationship of bleeding to age

Dysfunctional uterine bleeding (DUB) is more common at the extremes of the reproductive era (Figure 92.1).[4] The incidence of malignant disease as a cause of bleeding increases with age, being greatest after the age of 45, while endometrial cancer is predicted to be less than 1 in 100 000 in women under the age of 35.[1]



Figure 92.1 The relationship between age and various causes of abnormal uterine bleeding. DUB is more common in the extremes of the reproductive era, while the incidence of cancer as a cause of bleeding is greatest in the perimenopausal and postmenopausal phases After Mackay et al.[4]

MENORRHAGIA

Menorrhagia, which is excessive blood loss (> 80 mL per period),[5] is essentially caused by hormonal dysfunction (e.g. anovulation), local pathology (e.g. fibroids) or medical disorder (e.g. blood dyscrasia). Heavy bleeding, possibly with clots, is the major symptom of menorrhagia. Dysmenorrhoea may accompany the bleeding and, if it does, endometriosis or PID should be suspected. With care a 60 - 80% accuracy can be achieved in clinical assessment.[5] A summary of the diagnostic strategy model is presented in Table 92.3.

Table 92.3 Menorrhagia: diagnostic strategy model

Q Probability diagnosis
A Dysfunctional uterine bleeding - ovulatory

Fibroids

Complications of hormone therapy

Adenomyosis

Q. Serious disorders not to be missed

A. Disorders of pregnancy

- ectopic pregnancy
- abortion or miscarriage

Neoplasia

- cervical cancer
- endometrial cancer
- leukaemia
- benign tumours
  - polyps, etc.

Severe infections

- PID

Q. Pitfalls (often missed)

A. Genital tract trauma

IUCD

Adenomyosis/endometriosis

Pelvic congestion syndrome

SLE

Rarities

- endocrine disorders (e.g. thyroid disorder)
- bleeding disorder
- liver disease

Q. Seven masquerades checklist

A. Depression association

Diabetes

Drugs

Anaemia association

Thyroid disorder   hypothyroidism

Spinal dysfunction -

UTI -

Q. Is the patient trying to tell me something?

A. Consider exaggerated perception. Note association with anxiety and depression.

By far the most common single ‘cause’ of menorrhagia is ovulatory dysfunctional uterine bleeding. The most common organic causes are fibromyomatas (fibroids), endometriosis, adenomyosis (‘endometriosis’ of the myo-metrium), endometrial polyps and PID.[4]

Clinical approach for menorrhagia

History

Bearing in mind that abnormal uterine bleeding is a subjective complaint, a detailed history is the key initial step in management. The patient's perception of abnormal bleeding may be quite misleading and education about normality is all that is necessary in her management. A meticulous history should include details of the number of tampons or pads used and their degree of saturation. A menstrual calendar (over 3+ months) can be a very useful guide. A history of smoking and other psychosocial factors should be checked. For unknown reasons, cigarette smokers are five times more likely to have abnormal menstrual function.[3]

Questions need to be directed to rule out:[1]

- pregnancy or pregnancy complications (e.g. ectopic pregnancy)
- trauma of the genital tract
- medical disorders (e.g. bleeding disorder)
- endocrine disorders
- cancer of the genital tract
- complications of the pill

Physical examination[1]

A general physical examination should aim at ruling out anaemia, evidence of a bleeding disorder and any other stigmata of relevant medical or endocrine disease.

Specific examinations include:

- speculum examination: ? ulcers (cervical cancer) or polyps
- Pap smear
- bimanual pelvic examination: ? uterine or adnexal tenderness, size and regularity of uterus

It is prudent to avoid vaginal examination in selected patients, such as a young virgin girl, as the procedure is unhelpful and unnecessarily traumatic.

Investigations

Investigations, especially vaginal ultrasound scans, should be selected very carefully and only when really indicated. Abnormal pelvic examination findings, persistent symptoms, older patients and other suspicions of disease indicate further investigation to confirm symptoms of menorrhagia and exclude pelvic or systemic pathology.

Consider foremost:

- full blood count (to exclude anaemia and thrombocytopenia)
- iron studies: serum ferritin
- hysteroscopy and endometrial sampling (use directed endometrial biopsy with an instrument such as a Pipelle or Gynoscann, or curettage under general anaesthetic)

Special investigations (only if indicated):

- pregnancy testing
- laparoscopy where endometriosis, PID or other pelvic pathology is suspected
- serum biochemical screen
- coagulation screen
- thyroid function tests
- tests for SLE: antinuclear antibodies
- ultrasound

Note: Hysteroscopy and D&C remain the gold standard for abnormal uterine bleeding. In some women, a transvainal ultrasound, Pipelle endometrial sauyling or hysteroscopy and D&C will be indicated.[6]

DYSFUNCTIONAL UTERINE BLEEDING

DUB, which is a diagnosis of exclusion, is defined as 'excessive bleeding, whether heavy, prolonged or frequent, of uterine origin, which is not associated with recognisable pelvic disease, complications of pregnancy or systemic disease'.[4]

Features

- It is a working clinical diagnosis based on the initial detailed history, normal physical examination and normal initial investigation.
- Very common: 10 - 20% incidence of women at some stage.
- Peak incidence of ovulatory DUB in late thirties and forties (35 - 45 years).
- Anovulatory DUB has two peaks: 12 - 16 years and 45 - 55 years.
- The majority complain of menorrhagia.
- Up to 40% with the initial diagnosis of DUB will have other pathology (e.g. fibroids, endometrial polyps) if detailed pelvic endoscopic investigations are undertaken.

Symptoms

- Heavy bleeding: saturated pads, frequent changing, 'accidents', 'flooding', 'clots'
- Prolonged bleeding
    - menstruation > 8 days *or*
    - heavy bleeding > 4 days
- Frequent bleeding - periods occur more than once every 21 days
- Pelvic pain and tenderness are not usually prominent features

Management principles[5]

- Establish diagnosis by confirming symptoms and exclude other pathology.
- If no evidence of iron deficiency or anaemia, and significant pathology has been excluded, prospective assessment of the menstrual pattern is indicated using a menstrual calendar.
- Conservative management is usually employed if the uterus is of normal size and there is no evidence of anaemia.
- Drug therapy is indicated if symptoms are persistently troublesome and surgery is contraindicated or not desired by the patient and D&C doesn't alleviate.
- Provide reassurance about the absence of pathology, especially cancer, and give counselling to maximise compliance with treatment.
- Consider surgical management if fertility is no longer important and symptoms cannot be controlled by at least 3 - 4 months of hormone therapy.
- General rule:
  < 35 years - medical treatment
  > 35 years - hysteroscopy and direct endometrial sample (diagnostic - sometimes therapeutic)

Drug therapy[5]

Treatment regimens are presented in Tables 92.4 and 92.5. First-line treatment is with fibrinolytic inhibitors or antiprostaglandin agents, given as soon as possible and throughout the menses. These agents are simple to use, generally very safe and can be used over long periods of time. About 60 - 80% of patients with ovulatory menorrhagia will respond if compliance is good.[5] Such agents include tranexamic acid, mefenamic acid, naproxen, ibuprofen and indomethacin. The agent of first choice is usually mefenamic acid, which reduces blood loss by 20 - 25% as well as helping dysmenorrhoea. Ideally, it should be started at least 4 days before the menses. Evidence-based reviews confirm the benefit of NSAIDs and tranexamic acid for menorrhagia over the other agents.[7]

Table 92.4 Regimens used in management of menorrhagia

NSAIDs (prostaglandin inhibitors) [*]
Mefenamic acid 500 mg tds (4 days before menses due to end of menses) *or* Naproxen 500 mg statim then 250 mg tds *or* Ibuprofen 800 mg statim then 400 mg 6 - 8 hourly
Combined oestrogen - progesterone OC
This is an important first-line therapy. e.g. 50 μg oestrogen + 1 mg norethisterone (Norinyl)
Progestogens (especially for anovulatory patients)
Norethisterone 5 - 15 mg/day for 14 days (days 15 - 28) *or* Medroxyprogesterone acetate 20 - 30 mg/day Try progestogens from days 5 - 25 (ovulatory patients) or if no response to days 15 - 28 therapy.
Danazol [*]
Approved for short-term treatment (6 months or less) of severe menorrhagia - dosage 100 - 200 mg daily. Stops menstruation.
Antifibrinolytic agents [*]
Tranexamic acid 1 g (o) qid, days 1 - 4
GnRH agonists
Administer by nasal spray (Synarel) or monthly implant (Zoladex) to induce medical 'menopause' 3 - 6 months.
Progestogen-releasing IUDs (levonorgestrel)
e.g. Mirena → amenorrhoea in 20 - 50% after 1 year

[*] Effectiveness supported by EBM.[7] A trade-off between benefits and harms applies to danazol.

Table 92.5 Typical treatment options for acute and chronic heavy bleeding[5, 8]

Acute heavy bleeding
• Curettage/hysteroscopy • IV oestrogen (Premarin 25 mg) *or* oral high-dose progestogens (e.g. norethisterone 5 - 10 mg 2 hourly until bleeding stops then 5 mg bd or tds for 14 days[3])
Chronic bleeding
• For anovulatory women - cyclical oral progestogens for 14 days - tranexamic acid • For ovulatory women - cyclical prostaglandin inhibitor (e.g. mefenamic acid) *Or* oral contraceptive

- antifibrinolytic agent (e.g. tranexamic acid 1 g (o) qid, days 1 - 4)

- progesterone-releasing IUDs (e.g. Mirena)

Hormonal agents include progestogens, combined oestrogen - progestogen oral con-traceptives and danazol. Oestrogens can be used but generally are not recommended except in the occasional patient with very heavy bleeding, when intravenous conjugated oestrogens 25 mg can be used (repeated in 2 hours if no response) and always followed by a 14-day course of oral progestogen. The COC constitutes important firstline therapy in both ovulatory and anovulatory patients, but at least 20% of patients do not respond. It is preferable to use a pill with a higher oestrogen dose, which works better (50 µg rather than 30 µg or 35 µg of oestrogen), and one that contains nor-ethisterone[1] (e.g. Norinyl - 1).

Progestogens can be given via several routes. Oral use is usually of no benefit in ovulatory DUB. In the adolescent with anovulatory DUB, cyclical oral progestogens may be required for 6 months until spontaneous regular ovulation eventuates.[9] Intramuscular medroxypro-gesterone acetate (Depo-Provera) will induce amenorrhoea in 50% of users in 1 year.

The most effective agent for both ovulatory and anovulatory DUB is tranexamic acid, which inhibits endometrial plasminogen activation. The dose is 1 g (up to 1.5 g if necessary) orally qid for the first 4 days of the menstrual cycle commencing at the onset of visible bleeding.[8, 10]

The intrauterine progesterone implant system (Mirena) releasing 20 µg of levonorgestrel/day has shown considerable effectiveness.[11] It is regarded as the most efficacevess of the hormone treatments with a mean blood loss of 94% of women with menorrhagia.[6]

Practice tip

Acute flooding

Horethisterone 5 - 10 mg 2 hourly till bleeding stops, then 5 mg bd or tds (or 10 mg daily) for 14 days)

Surgical options

Surgical treatment for menorrhagia is more appropriate if the uterus is enlarged, especially if greater than the size of a 12 week gestation (grapefruit size) or if the patient is anaemic.[1] It is indicated if menorrhagia interferes with lifestyle despite medical (drug) treatment. The surgical options are:

- endometrial ablation or electrodiathermy excision - to produce amenorrhoea
- hysterectomy (up to 25% of Australian women will have this before age 50); it requires a very carefully planned approach

CYCLE IRREGULARITY[12]

For practical purposes patients with irregular menstrual cycles can be divided into those under 35 and those over 35 years.

Patients under 35 years:

- the cause is usually hormonal, rarely organic, but keep malignancy in mind
- management options[1]
  - explanation and reassurance (if slight irregularity)
  - COC pill for better cycle control - any pill can be used
  - progestogen-only pill (especially anovulatory cycles) norethisterone (Primolut N) 5 - 15 mg/day from day 5 - 25 of cycle

Patients over 35 years should be referred for investigation for organic pathology, usually by endometrial sampling and/or hysteroscopy. If normal, the above regimens can be instituted.

INTERMENSTRUAL BLEEDING AND POSTCOITAL BLEEDING

These bleeding problems are due to factors such as cervical ectropion (often termed cervical ‘erosion’), cervical polyps, the presence of an IUCD and the oral contraceptive pill. Cervical cancer and intrauterine cancer must be ruled out, hence the importance of a Pap smear in all age groups and endometrial sampling, especially in the over 35 age group. Mismanagement of these presentations is a legal ‘Minefield’. A Pap smear should be taken, using the speculum carefully so as not to provoke bleeding, if one has not been taken within the previous 3 months, and sent to a laboratory that uses appropriate quality control procedures with notation of the bleeding on the smear request form. Remember that it is only a screening test. Refer women with these bleeding problems with an abnormal smear or even without any unusual features. Those with a friable ectropion that is causing persistent symptoms should also be referred.[13] Thus, intermenstrual bleeding (IMB) should always be investigated. Order a pregnancy test if appropriate.

Cervical ectropion, which is commonly found in women on the pill and post partum, can be left untreated unless intolerable discharge or moderate postcoital (PCB)[1] bleeding is present. An IUCD should be removed if causing significant symptoms and the causative pill should be changed to one with a higher oestrogen dose (e.g. from 30 μg oestrogen to 50 μg oestrogen).

UTERINE FIBROIDS (LEIOMYOMA)

Fibroids are benvygn tumours of smooth muscle of the myometrium. They are classified according to their location: Subserosal, intramusal, Subeulometreal or Intrauterine. They are oesteogen dependent and shrink with ouset of menopause.

Features

- Present in 30% of women > 35
- Only 1 in 800 develop malignancy
- Usually asymptomatic

Symptoms (especially if large)

- Menorrhagia
- Dysmenorrhoea
- Pelvic discomfort ± pain (pressure)
- Bladder dysfunction
- Pain with forsion of peduncnlated fibroid
- Pain with ‘red degeneration’ - only in pregnancy (pain, fever, local tenderness)

Other features

- Infertility (acts like IUCD if submucosal)
- Calcification

Examination

- Bulky uterns

Investigation

- Pelvic ultrasound
- FBE ? anaemia
- Uterine biopey (malignancy suspected)

Management

- Consider COC
- GnRH analognes - especially > 42 can shrink fibroids (maximum 6 months)
- Surgical option:
  - myomectomy (remove fibroids only, esp. child-bearing years)
  - hysteroscopie resection
  - hysterectomy

CARCINCOMA OF THE CERVIX

This should be the diagnosis until proved otherwise for post-coital intermenstrual or postmenopausal bleeding.

Features

- Peak incidence in sixth decade
- 80% due to squamoies cell carcinoma
- Risk factors (refer p. 951)

Symptoms

- Post-coital bleeding
- Inemenstrual bleeding
- Vaginal discharge - may be offensive

Mainly diagnosed on routine screening

Examination

- Ulceration or mass on cervix
- Bleeds readily on contact - may be friable

Management

- Urgent gynaecological referral

ENDOMETRIAL CARCER

This is the diagnosis until proved otherwise in any woman presenting with post - menopausal bleeding.

Features

- Peak incidence 50 - 70 years
- Risk factors:
  - age
  - obesity
  - nulliparity
  - late menopause
  - diapetes mellitus
  - polycystic ovary
  - diabetes mellitus
  - drugs, eg. Unopposed oestrogen tamoxifen
  - family history - breast, ovarion, colon cancer

Symptoms

80% percent with abnormal bleeding, especially post-menopausal bleeding.

Examination

- Uterus usually feelinormal, but may be bulky.

Investigations

- Smear (after pregnancy exclude) - detects some cases
- Transvaginal ultrasound

Management

- Urgent gynaecological seperral

AMENORRHOEA AND OLIGOMENORRHOEA

Amenorrhoea is classified as primary or secondary.

Primary amenorrhoea is the failure of the menses to start by 16 years of age.[3] Secondary amenorrhoea is the absence of menses for over 6 months in a woman who has had established menstruation.

The main approach in the patient with primary amenorrhoea is to differentiate it from delayed puberty. It is important to keep in mind the possibility of an imperforate hymen. A good rule is to note the presence of secondary sex characteristics.[4] If absent it implies that the ovaries are non-functional. Causes of primary amenorrhoea include genital malformations, ovarian disease, pituitary tumours, hypo-thalamic disorder and Turner's syndrome. Refer early as appropriate.

In secondary amenorrhoea, consider a physiological cause such as pregnancy, or the menopause, failure of some part of the hypothalamic - pituitary - ovarian - uterine axis (e.g. PCOS), or a metabolic disturbance. Important causes to consider are emotional, psychiatric and constitutional causes such as anorexia nervosa, hyperprolactinaemia, strenuous exercise, weight loss below 75% of ideal, and drugs/hormone therapy (e.g. oral contraceptives).

Oligomenorrhoea is the term for infrequent and usually irregular periods, where the cycles are between 6 weeks and 6 months.

POSTMENOPAUSAL BLEEDING

Postmenopausal bleeding is vaginal bleeding of any amount occurring 6 months or more after the menopause.[4] It suggests cervical or uterine body cancer (up to 25%).[4] Other causes include polyps, atrophic vaginitis, endometrial hyperplasia and urethral caruncle. Care has to be taken with women on HRT who have irregular bleeding - they require investigation.

Early referral is usually indicated with a view to a diagnostic procedure. If bleeding recurs despite curettage, hysterectomy should be performed since early cancer of the uterus may be missed.

WHEN TO REFER

- Women with persistent IMB and/or PCB unthout any unusual feaxenes
- Women with IMB/PCB and an abnormal smeal[13]
- Women with a friable ectropion

- To exclude intrauterine pathology
- The patient does not respond to initial therapy
- There is evidence of underlying disease (e.g. endometriosis, SLE)
- Surgery is indicated (minor or major)

Practice tips

- Remember that mental dysfunction can obscure the organic causes of menorlagia.
- Non-menstrual bleeding suggests cancer until proved otherwise: it may be postcoital (cervical cancer); intermenstrual (common with progestogen-only pill); postmenopausal (endometrial cancer).
- Think of a foreign body, especially an IUCD: if it is an IUCD, remove it.
- Hysteroscopy is more effective than the traditional curettage. Studies have shown that usually less than 50% of the uterine cavity is sampled by curettage.[14]

REFERENCES

1. Fung P. Abnormal uterine bleeding. Mod Med Aust, 1992; May: 58 - 66.
2. Fraser IS, Pearce C, Shearman RP et al. Efficacy of mefenamic acid in patients with a complaint of menorrhagia. Obstet Gynaecol, 1981; 58: 543 - 51.
3. Wood C. How to treat: Menorrhagia. Australian Doctor, 1999; 12 march: I - VIII
4. Mackay EV, Beischer NA, Pepperell RJ, Wood C. *Illustrated Textbook of Gynaecology* (2nd edn). Sydney: WB Saunders, 1992: 77 - 107.
5. Fraser IS. Dysfunctional uterine bleeding. In: *MIMS Disease Index*. Sydney: IMS Publishing, 1991 - 92: 165 - 7.
6. Quinlivan J, Petersen RW. Menorrhagia. Medreal Obsenver, 16 April 2004: 31 - 34.
7. Barton S ed. *Clinical Evidence*. London: BMJ Publishing Group, 2001: 1311 - 16.
8. Bonnar J, Sheppard BL. Treatment of menorrhagia during menstruation: randomised controlled trial of ethamsylate, mefenamic acid and tranexamic acid. Br Med J, 1996; 313: 579 - 82.
9. Hickey M. Management options in menorrhagia. Curr Ther, 1996; July: 17 - 21.
10. Gleeson NC, Buggy F et al. The effect of tranexamic acid on measured menstrual blood loss and endometrial fibrinolytic enzymes in dysfunctional uterine bleeding. Acta Obstet Gynaecol Scand, 1994; 73: 274 - 7.
11. Anderson JK, Rybo G. The levonorgestrel-releasing intrauterine contraceptive device in the treatment of menorrhagia. Br J Obstet Gynaecol, 1990; 97: 690 - 4.
12. Hewson A. Menstrual disorders. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 307 - 10.
13. RACOG and RACGP. Joint Guidelines for Referral for Investigation of Intermenstrual and Postcoital Bleeding. Melbourne: RACOG & RACGP 1996.
14. Warton B. *Gynaecology*. Melbourne: CHECK programme, RACGP Unit 240, 1992: 2 - 20.

CHAPTER 93. LOWER ABDOMINAL AND PELVIC PAIN IN WOMEN

Man endures pain as an undeserved punishment, woman accepts it as a natural heritage.

ANONYMOUS

LOWER ABDOMINAL AND PELVIC PAIN IN WOMEN

Pain in the lower abdomen and pelvis is one of the most frequent symptoms experienced by women. The diagnostic approach requires a wide variety of consultative skills, especially when the pain is chronic. The examination of acute abdominal pain has been simplified by the advent of sensitive serum pregnancy tests, ultrasound investigation and the increasing use of laparoscopy. However, an accurate history and examination for all types of pain will generally pinpoint the diagnosis. The ever-present problem of Pid, the leading cause of infertility in women, demands an early diagnosis and appropriate management.

KEY FACTS AND CHECKPOINTS

- A distinction has to be made between acute, chronic and recurrent pain.
- Ectopic pregnancy remains a potentially lethal condition and its diagnosis still requires a high index of suspicion.
- Sudden sharp pain in the pelvis which becomes more generalised indicates rupture of an ectopic pregnancy or an ovarian cyst.
- Recurrent sharp self-limiting pain indicates a ruptured Graafian follicle (mittelschmerz).
- Recurrent pain related to menstruation is typical of dysmenorrhoea or endometriosis.
- A UK study[1] of chronic lower abdominal pain in women showed the causes were adhesions (36%), no diagnosis (19%), endometriosis (14%), constipation (13%), ovarian cysts (11%) and Pid (7%). An Australian study found that endometriosis accounted for 30% and adhesions 20%.[2]
- The principal aff erent pathways of the pelvic viscera arise from T10 - 12, L1 and S2 - 4. Th us disorders of the bladder, rectum, lower uterus, cervix and upper vagina can refer pain to the low back, buttocks and posterior thigh.[3]

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 93.1.

Table 93.1 Lower abdominal and pelvic pain in women: diagnostic strategy model

Q. Probability diagnosis
A. Primary dysmenorrhoea Mittelschmerz Pelvic/abdominal adhesions Endometriosis
Q. Serious disorders not to be missed
A. Ectopic pregnancy Neoplasia • ovary • uterus • other pelvic structures Severe infections

- Pid

Acute appendicitis

Q. Pitfalls (often missed)

A. Endometriosis/adenomyosis

Torsion of ovary or pedunculated fibroid

Constipation

Pelvic congestion syndrome

Referred pain (to pelvis)

- appendicitis
- cholecystitis
- diverticulitis
- urinary tract infection

Q. Seven masquerades checklist

A. Depression ✔

Diabetes -

Drugs ✔

Anaemia -

Thyroid disorder -

Spinal dysfunction ✔

UTI ✔

Q. Is the patient trying to tell me something?

A. Can be very relevant. Consider various problems and sexual dysfunction.

Probability diagnosis

The commonest causes are primary dysmenorrhoea, the pain of a ruptured Graafian follicle (mittelschmerz), endometriosis and adhesions. In many instances of pain no diagnosis is made as no pathological cause can be found.

Serious disorders not to be missed

The potentially lethal problem of a ruptured ectopic pregnancy must not be missed, hence the axiom ‘be ectopic minded’. Pid can be overlooked, especially if chronic, and requires early diagnosis and aggressive treatment. Neoplasia must be considered, especially malignancy of pelvic structures, including the ‘silent’ carcinoma of the ovary.

Pitfalls

Several disorders are very difficult to diagnose and these include haemorrhage into the ovary or a cyst, torsion of the ovary or pedunculated fibroid. Endometriosis may be missed so it is important to be familiar with its symptoms. Chronic constipation may be a trap. Another relatively common problem is the so-called ‘pelvic congestion syndrome’, which tends to occur in somewhat neurotic patients and also tends to be a diagnosis of exclusion.

Seven masquerades checklist

Two important conditions to consider are urinary tract infection and spinal dysfunction. Just as disorders of the pelvic organs, such as endometriosis and Pid, can refer pain to the low back and buttocks, so can disorders of the lumbosacral spine cause referred pain to the lower abdomen and groin.

Psychogenic considerations

These can be extremely relevant. Problems in the patient's social, marital or sexual relationships should be evaluated, especially in the assessment of chronic pain. Many patients with undiagnosed chronic pain exhibit psychoneurotic traits and this renders management very complex. Some appear to have the 'pelvic congestion syndrome' and need to be handled with sensitivity and tact, especially if the help of a psychiatrist or psychologist is sought.

THE CLINICAL APPROACH

History

The pain should be linked with the menstrual history, coitus and the possibility of an early pregnancy. For recurrent and chronic pain, it is advisable to instruct the patient to keep a diary over two menstrual cycles. The severity of the pain can be assessed as follows:[2]

- does not interfere with daily activity
- results in days off work
- results in confinement to bed

In this way the pain can be classified objectively as mild, moderate or severe.

Risk factors in the past history should be assessed, for example:

- IUCD (salpingitis, ectopic pregnancy)
- infertility (endometriosis, salpingitis)
- tubal surgery (ectopic)

The typical pain patterns in relation to menstruation are shown in Figure 93.1.


endometriosis
secondary dysmenorrhoea (other than endometriosis)
primary dysmenorrhoea
distension of ovarian capsule pain
mittelschmerz
0
menses
5
ovulation
28
menses
5

Figure 93.1 Typical pain patterns for the menstrual cycle related gynaecological pain

Physical examination

One objective is to correlate any palpable tenderness with the patient's statement of the severity of the pain. Use the traditional abdominal and pelvic examination to identify the site of tenderness and rebound tenderness, and any abdominal or pelvic masses. The pelvis should be examined by speculum (preferably bivalve type) and bimanual palpation.

Proper assessment can be difficult if the patient cannot relax or overreacts, if there is abdominal scarring or obesity, or if extreme tenderness is present. It is therefore important, especially in the younger and apprehensive patient, to conduct a gentle caring vaginal examination with appropriate explanation and reassurance. Explanation of the procedure during vaginal examination, preferably using eye contact with the patient, can help her relax and be more confident in the procedure.

Investigations

Investigations may be selected from:

- haemoglobin level
- white blood cell count (limited value)
- haematocrit
- ESR/CRP
- microbiology (limited value)
    - urine for microscopy and culture ± *Chlamydia* PCR
    - endocervical, urethral, cervical and vaginal swabs
- serum μ-HCG assay
- urinary HCG tests (can be negative in the presence of an ectopic pregnancy)

Diagnostic imaging:

- vaginal ultrasound - to define a gestation sac
- pelvic ultrasound - to differentiate a cystic from a solid pelvic mass

Indicated for:

- pelvic pain
- a palpable pelvic mass
- a palpable lower abdominal mass
- ascites

Laparoscopy is indicated if the history and examination are suggestive of ectopic pregnancy and ultrasound fails to confirm an intrauterine pregnancy.

ACUTE PAIN

The causes of acute pain are summarised in Table 93.2. The patient is usually young (20 - 30 years old), sexually active and distressed by the pain, and should be considered foremost to have a bleeding ectopic pregnancy. Important differential diagnoses include acute Pid, rupture or torsion of an ovarian cyst and acute appendicitis. Cases of acute ruptured ectopics are obviously easier to diagnose in the presence of circulatory collapse.

Table 93.2 Causes of acute lower abdominal and pelvic pain in women (after Soo Keat Khoo)[3]

Genital
• Acute salpingitis
• Pelvic peritonitis
• Bleeding
• Rupture or torsion of ovarian cyst
• Threatened or incomplete abortion
• Rupture or aborting tubal ectopic pregnancy
• Rupture or bleeding endometrioma
Non-genital
• Acute appendicitis
• Bowel obstruction
• Urinary tract infection (cystitis)
• Ureteric colic (calculus)

Functional
• Primary dysmenorrhoea
• Retrograde menstruation

CHRONIC PAIN

Features

- Incidence 15% in 18 - 50 year olds
- Endometriosis causes 33%, adhesions 24%
- It is the reason for up to 40% of gynaecological laparoscopies
- Reason for 15% hysterectomies

The pain can be cyclical (e.g. endometriosis, mittelschmerz) or continuous.

The common causes of chronic pain are listed in Table 93.3. Chronic pain is more difficult to diagnose and it is often difficult to differentiate between problems such as endometriosis, Pid, an ovarian neoplasm and the irritable bowel syndrome. A comparison of the clinical features of endometriosis and Pid is presented in Table 93.4. Furthermore it is difficult to distinguish clinically between endometriosis of the uterus (adenomyosis) and pelvic congestion syndrome. Both conditions are associated with dysmenorrhoea and a tender normal-sized uterus.

Table 93.3 Causes of chronic lower abdominal and pelvic pain in women[3]

Genital
• Endometriosis/adenomyosis
• Pelvic inflammatory disease (chronic; adhesions)
• Ovarian neoplasm
• Fibromyomata (rarely)
Non-genital
• Diverticulitis
• Bowel adhesions
Functional
• Pelvic congestion syndrome
• Secondary dysmenorrhoea - IUCD, polyp
• Irritable bowel, chronic bowel spasm

Table 93.4 Comparison of clinical features of Pid and pelvic endometriosis

Feature	Chronic Pid	Endometriosis
History	Acute pelvic infection (e.g. ruptured appendix)	Dysmenorrhoea
		Infertility
	IUCD usage	Dyspareunia
		Pelvic pain
Pelvic pain	+ to + + (mild to moderate)	+ + to + + + (moderate to severe)
	Premenstrual	Premenstrual and menstrual

	Lower abdominal location	Acute pain if rupture of endometrioma
Backache	+ mild	+ + moderate Low sacral pain with menstruation
Secondary dysmenorrhoea	Moderate to severe From onset of acute Pid Decreases with menstruation	Moderate to severe Gradual onset Increases in severity throughout menstruation
Menstruation	Irregular and heavy	Heavy
Dyspareunia	Moderate	Often severe
Infertility	+ + +	+ +
Urinary symptoms	-	Frequency, dysuria and haematuria if bladder wall involved
Bowel symptoms	-	Painful defecation if rectal wall involved
Vaginal symptoms	May be chronic purulent discharge or leucorrhoea	-

PELVIC CONGESTION SYNDROME

this is considered to be due to ovarian dysfunction (similar to PCOS) with various congestion.

Features

- Patient usually para 3 or 4
- Typical age 35 - 40 years
- Unilateral pain, increased with standing and walking
- Relief with lying down
- Deep dysparunia
- Post-coital aching

The patient usually has a multitude of emotional problems.[3] They often undergo hysterectomy, sometimes without relief of symptoms.

ECTOPIC PREGNANCY

Ectopic pregnancy occurs approximately once in every 100 clinically recognised pregnancies. If ruptured it can be a rapid, fatal condition so we have to be 'ectopic minded'. It is the commonest cause of intraperitoneal haemorrhage. There is usually a history of a missed period but a normal menstrual history may be obtained in some instances.

Typical clinical features of a ruptured ectopic pregnancy

- Average patient in mid-twenties
- First pregnancy in one-third of patients
- Patient at risk
    - previous ectopic pregnancy
    - previous Pid
    - previous abdominal or pelvic surgery, especially sterilisation reversal
    - IUCD use
    - in-vitro fertilisation/GIFT

- Pre-rupture symptoms (many cases)
  - abnormal pregnancy
  - cramping pains in one or other iliac fossa
  - vaginal bleeding
- Rupture
  - excruciating pain (Figure 93.2)
  - circulatory collapse

Note: In 10 - 15% there is no abnormal bleeding.

- Pain may radiate to rectum (lavatory sign), vagina or leg
- Signs of pregnancy, e.g. enlarged breasts and uterus, usually not present


cramping pain
in RIF or LIF
if unruptured
excruciating
pain if ruptured
• lower abdominal pain
• amenorrhoea
• abnormal vaginal bleeding
—'prune juice'

Figure 93.2 Clinical features of ectopic pregnancy

DxT: amenorrhoea (65 - 80%) + lower abdominal pain (95+ %) + abnormal vaginal bleeding (65 - 85%) = ectopic pregnancy

Examination

- Deep tenderness in iliac fossa
- Vaginal examination
  - tenderness on bimanual pelvic examination (pain on cervical provocation)
  - soft cervix
- Bleeding (prune juice appearance)
- Temperature and pulse usually normal early

Diagnosis

It is possible to diagnose ectopic pregnancy at a very early stage of pregnancy.

- Urine pregnancy test (positive in < 50%)
- Serum μ-HCG assay (invariably positive if a significant amount of viable trophoblastic tissue present) - may need serial quantitative tests to distinguish an ectopic from a normal interuterine pregnancy
- Transvaginal ultrasound can diagnose at 5 - 6 weeks (empty uterus, tubal sac, fluid in cul-de-sac)

- Laparoscopy (the definitive diagnostic procedure)

Management

Treatment may be conservative (based on ultrasound and μ-HCG assays); medical, by injecting methotrexate into the ectopic sac; laparoscopic removal; or laparotomy for severe cases. Rupture with blood loss demands urgent surgery.

Post management

- Successful pregnancy 60 - 65%
- Subsequent risk of ectopic pregnancy 10 - 15%

RUPTURED OVARIAN (GRAAFIAN) FOLLICLE (MITTELSCHMERZ)

When the Graafian follicle ruptures a small amount of blood mixed with follicular fluid is usually released into the pouch of Douglas. this may cause peritonism (mittelschmerz), which is different from the unilateral pain experienced just before ovulation due to distension of the ovarian capsule.

Typical clinical features

- Onset of pain in mid-cycle
- Deep pain in one or other iliac fossa (RIF > LIF)
- Often described as a ‘horse kick pain’
- Pain tends to move centrally (Figure 93.3)
- Heavy feeling in pelvis
- Relieved by sitting or supporting lower abdomen
- Pain lasts from a few minutes to hours (average 5 hours)
- Patient otherwise well

Note: Sometimes it can mimic acute appendicitis.


pain tends to move centrally
deep pain in RIF or LIF
• deep ache in lower abdomen
• lateral then shifts centrally
• heavy feeling in pelvis

Figure 93.3 Typical clinical features of a ruptured Graafian follicle (mittelschmerz)

Management

- Explanation and reassurance
- Simple analgesics: aspirin or paracetamol
- 'Hot water bottle' comfort if pain severe

OVARIAN TUMOURS

Benign ovarian tumours, particularly ovarian cysts, may be asymptomatic but will cause pain if complicated. They are common in women under 50 years of age. Ovarian cysts are best defined by vaginal ultrasound, which can identify whether haemorrhage has occurred inside or outside the cyst.

Symptoms

- Pain (usually torsion or haemorrhage)
- Pressure symptoms
- Menstrual irregularity

Ruptured ovarian cyst

The cysts tend to rupture just prior to ovulation or following coitus.

Clinical features

- Patient usually 15 - 25 years
- Sudden onset of pain in one or other iliac fossa
- May be nausea and vomiting
- No systemic signs
- Pain usually settles within a few hours

Signs

- Tenderness and guarding in iliac fossa
- PR: tenderness in rectovaginal pouch

Investigation

- Ultrasound ±colour Doppler (for enhancement)

Management

- Appropriate explanation and reassurance
- Conservative
  - simple cyst < 4 cm
  - internal haemorrhage
  - minimal pain
- Needle vaginal drainage by ultrasonography for a simple larger cyst
- Laparoscopic surgery
  - complex cysts
  - large cysts
  - external bleeding

Acute torsion of ovarian cyst

Torsions are mainly from dermoid cysts and, when right-sided, may be difficult to distinguish from acute pelvic appendicitis.

Clinical features

- Severe cramping lower abdominal pain (Figure 93.4)
- Diffuse pain
- Pain may radiate to the flank, back or thigh
- Repeated vomiting
- Exquisite pelvic tenderness
- Patient looks ill


severe cramping lower abdominal pain
may radiate to flank or thigh
• diffuse severe pain
• repeated vomiting
• patient looks ill
• exquisite pelvic tenderness

Figure 93.4 Typical clinical features of acute torsion of an ovarian cyst

Signs

- Smooth, rounded, mobile mass palpable in abdomen
- May be tenderness and guarding over the mass, especially if leakage

Diagnosis

- Ultrasound ±colour Doppler

Treatment

- Laparotomy and surgical correction

MALIGNANT OVARIAN TUMOURS

Ovarian cancer has an incidence of 10 cases per 100 000 women per year and accounts for 5% of all cancers in women and 20% of all genital cancers. It is responsible for more genital cancer deaths because the tumour is often well advanced at the time of clinical presentation.[4] Earlier discovery may sometimes be made on routine examination or because of investigation of non-specific pelvic symptoms.

Ovarian cancer tends to remain asymptomatic for a long period. No age group is spared but it becomes progressively more common after 45 years (peak incidence 60 - 65 years).

The familial causes and relationship to breast cancer are being delineated. Refer to Chapter 19.

Risk factors

- Age
- Family history (first degree relatives)
- Nulliparity

Protective factors

- COC pill
- Pregnancy

Clinical features

- Ache or discomfort in lower abdomen or pelvis
- Abdominal bloating
- Gastrointestinal dysfunction (e.g. epigastric discomfort, constipation, wind)
- Sensation of pelvic heaviness
- Genitourinary symptoms (e.g. ↑ frequency, prolapse)
- ± Menstrual dysfunction
- Postmenopausal bleeding
- Dyspareunia and/or dysmenorrhoea (10 - 20%)
- A combined vaginal-rectal bimanual examination assists diagnosis. Look for mass, ascites, pleural effusion

Note: Any ovary that is easily palpable is usually abnormal (normal ovary rarely > 4 cm).

Diagnosis

- Ultrasound ±colour Doppler
- Tumour markers such as CA 125, HCG (choriocarcinoma) and alpha-fetoprotein are becoming more important in diagnosis and management

Refer urgently to gynaecologist.

DYSMENORRHOEA

Dysmenorrhoea (painful periods) may commence with the onset of the menses (menarche) when it is called primary dysmenorrhoea, or later in life when the term secondary dysmenorrhoea is applied.

PRIMARY (FUNCTIONAL) DYSMENORRHOEA

this is menstrual pain associated with ovular cycles without any pathologic findings. The pain usually commences within 1 - 2 years after the menarche and become more severe with time up to about 20 years. It affects about 50% of menstruating women and up to 95% of adolescents.

CLINICAL FEATURES

- Low midline abdominal pain
- Pain radiates to back or thighs (Figure 93.5)
- Varies from a dull dragging to a severe cramping pain

- Maximum pain at beginning of the period
- May commence up to 12 hours before the menses appear
- Usually lasts 24 hours but may persist for 2 - 3 days
- May be associated with nausea and vomiting, headache, syncope or flushing
- No abnormal findings on examination



Figure 93.5 Typical pain of dysmenorrhoea

MANAGEMENT

- Full explanation and appropriate reassurance
- Promote a healthy lifestyle
  - regular exercise
  - avoid smoking and excessive alcohol
  - Recommend relaxation techniques such as yoga
  - Avoid exposure to extreme cold
  - Place a hot water bottle over the painful area and curl the knees onto the chest

MEDICATION

Options include (trying in order):

- simple analgesics (e.g. aspirin or paracetamol)
- prostaglandin inhibitors (e.g. mefenamic acid 500 mg tds) or NSAids (e.g. naproxen, ibuprofen) at first suggestion of pain (if simple analgesics ineffective)
- Vitamin B1 (thiamine) 100 mg daily
- COC (low-oestrogen triphasic pills preferable)
- progestogen-medicated IUCD

A Cochrane review found that the most beneficial medication was the NSAids, and also vitamin B1 and magnesium proved effective. There was no evidence so far that the vitamin B6, vitamin E or herbal remedies were effective. Spinal manipulation was unlikely to be beneficial.[5]

SECONDARY DYSMENORRHOEA

Secondary dysmenorrhoea is menstrual pain for which an organic cause can be found. It usually begins after the menarche after years of pain free menses; the patient is usually over 30 years of age. The pain begins as a dull pelvic ache 3 - 4 days before the menses and becomes more severe during menstruation.

Commonest causes

- Endometriosis (a major cause)
- Pid (a major cause)
- IUCD
- Submucous myoma
- Intrauterine polyp
- Pelvic adhesions

Investigations

Investigations include laparoscopy, ultrasound and (less commonly) assessment of the uterine cavity by dilation and curettage, hysteroscopy or hysterosalpingography.

Management involves treating the cause.

PELVIC ADHESIONS

Pelvic adhesions may be the cause of pelvic pain, infertility, dysmenorrhoea and intestinal pain. They can be diagnosed and removed laparoscopically when the adhesions are well visualised and there are no intestinal loops firmly stuck together.

ENDOMETRIOSIS

Endometriosis is the condition where ectopically located endometrial tissue (usually in dependent parts of the pelvis and in the ovaries) responds to female sex hormone stimulation by proliferation, haemorrhage, adhesions and ultimately dense scar tissue changes.

Patients experience varying degrees of symptoms and loss of gynaecological function according to the site and severity of the endometriosis deposits. Pregnancy is beneficial but recurrence can follow.

Clinical features

- 10% incidence[6]
- Puberty to menopause, peak 25 - 35 years
- Secondary dysmenorrhoea
- Infertility
- Dyspareunia
- Non-specific pelvic pain
- Menorrhagia
- Acute pain with rupture of endometrioma
- Premenstrual spotting

Possible signs

- Fixed uterine retroversion
- Tenderness and nodularity in the pouch of Douglas/retrovaginal septum
- Uterine enlargement and tenderness

Differential diagnosis

- Pid - see Table 93.4
- Ovarian cysts or tumours
- Uterine myomas

Diagnosis

- can be made only by direct visual inspection at laparoscopy or laparotomy
- ultrasound scanning helpful

Treatment

- Careful explanation
- Basic analgesics
- Treatment can be surgical or medical

  Medical: To induce amenorrhoea (only twothirds respond to drugs)
  - danazol (Danocrine) - current treatment of choice
  - COC: once daily continuously for about 6 months
  - progestogens (e.g. medroxyprogesterone acetate - Depo-Provera)
  - GnRH analogues (e.g. goserelin, nafarelin)

Surgical: Surgical measures depend on the patient's age, symptoms and family planning. Laser surgery and microsurgery can be performed via either laparoscopy or laparotomy.

PELVIC INFLAMMATORY DISEASE

There are great medical problems in the serious consequences of Pid, namely tubal obstruction, infertility and ectopic pregnancy. Pid may be either acute, which causes sudden severe symptoms, or chronic, which can gradually produce milder symptoms or follow an acute episode. Acute Pid is a major public health problem and is the most important complication of STIs among young women. The majority are young (less than 25 years) sexually active women who are also nulliparous.

Some patients may experience no symptoms but others may have symptoms that vary from mild to very severe. The clinical diagnosis can be difficult as signs and symptoms can be nonspecific and correlate poorly with the extent of the inflammation.

Clinical features

Acute Pid

- Fever &x2A7E;38°C
- Moderate to severe lower abdominal pain

Chronic Pid

- Ache in the lower back
- Mild lower abdominal pain

Both acute and chronic

- Dyspareunia
- Menstrual problems (e.g. painful, heavy or irregular periods)
- Intermenstrual bleeding
- Abnormal, perhaps offensive, purulent vaginal discharge
- Painful or frequent urination

The diagnostic criteria for acute Pid are presented in Table 93.5.[7]

Table 93.5 Diagnostic criteria for acute Pid[3]

All three of the following should be present:

1. Lower abdominal tenderness (with or without rebound)
2. Cervical motion tenderness
3. Adnexal tenderness (may be unilateral) plus

One of the following should be present:

1. Temperature &x2A7E;38°C
2. White blood cell count &x2A7E;10 500/mm2
3. Purulent fluid obtained via culdocentesis
4. Inflammatory mass present on bimanual pelvic examination and/or sonography
5. ESR &x2A7E;15 mm/h or CRP > 1.0 mg/dL
6. Evidence for presence of *N. gonorrhoeae* and/or *C. trachomatis* in the endocervix. Gram stain reveals Gram-negative intracellular diplococci. Monoclonal antibody stain for *C. trachomatis*
7. Presence of > 5 white blood cells per oil immersion field on Gram stain of endocervical discharge

Examination

- In acute Pid there may be lower abdominal tenderness ±rigidity.
- Pelvic examination: in acute Pid there is unusual vaginal warmth, cervical motion tenderness and adnexal tenderness. Inspection usually reveals a red inflamed cervix and a purulent discharge.

Causative agents

These can be subdivided into three broad groups:

1. *Exogenous organisms*: those which are community acquired and initiated by sexual activity. They include the classic STIs, *Chlamydia trachomatis* and *Neisseria gonorrhoeae*. this usually leads to salpingitis.
2. *Endogenous infections*: these are normal commensals of the lower genital tract, especially *Escherichia coli* and *Bacteroides fragilis*. They become pathogenic under conditions that interrupt the normal cervical barrier, such as recent genital tract manipulation or trauma (e.g. abortion, presence of an IUCD, recent pregnancy or a dilatation and curettage). The commonest portals of entry are cervical lacerations and the placental site. These organisms cause an ascending infection and can spread direct or via lymphatics to the broad ligament, causing pelvic cellulitis (Figure 93.6).
3. *Actinomycosis:* due to prolonged IUCD use. Look for *Actinomyces israeli* on culture.


Mono-aetiology
Polymicrobial
pyosalpinx
pelvic abscess
tubo-ovarian abscesses
tubal damage
broad ligament
salpingitis
cervicitis
salpingitis
+ genital manipulation
Sexually transmitted pathogens
C. trachomatis
N. gonorrhoeae
Mycoplasma hominis
Vaginal flora
• Streptococci
• Bacteroides fragilis
• Enterobacteriaceae

Figure 93.6 The pathogenesis of Pid

Investigations

A definitive diagnosis is difficult since routine specimen collection has limitations in assessing the organisms. Definitive diagnosis is by laparoscopy but this is not practical in all cases of suspected Pid.

- Cervical swab for Gram stain and culture, PCR (*N. gonorrhoeae)*
- Cervical swab and special techniques e.g. PCR for *C. trachomatis*

Treatment of Pid

Note: Any IUCD or retained products of contraception should be removed at or before the start of treatment. Sex partners of women with Pid should be treated with agents effective against *C. trachomatis* and *N. gonorrhoeae.*

Sexually acquired infection[8]

Mild to moderate infection (treated as an outpatient)

azithromycin 1 g (o), as 1 dose plus (for gonorrhoea)

ceftriaxone 250 mg IM or IV, as 1 dose plus (in all patients)

doxycycline 100 mg (o) 12 hourly for 14 days plus either

metronidazole 400 mg (o) 12 hourly for 14 days *or*

tinidazole 500 mg (o) daily for 14 days

Severe infection (treated in hospital)

- cefotaxime 1 g IV 8 hourly or ceftriaxone 1 g IV once daily plus
- doxycycline 100 mg (o) or IV 12 hourly plus
- metronidazole 500 mg IV 12 hourly until there is substantial clinical improvement, when the oral regimen above can be used for the remainder of the 14 days. If the patient is pregnant or breastfeeding, doxycyline should be replaced by roxithromycin 300 mg (o) daily for 14 days

Infection non-sexually acquired (related to genital manipulation)

Mild to moderate infection

amoxycillin + clavulanate (875 mg/125 mg) (o) 12 hourly for 14 days plus doxycycline 100 mg (o) 12 hourly for 14 days

Severe infection (including septicaemia)

Use the same regimen as for severe sexually acquired infection.

Actinomycosis

Amoxycillin 500 mg tds + metronidazole 400 mg bd for 14 days. Ensure IUCD is removed.

Practice tips

- Think of endometriosis and ovarian cysts in any woman with lower abdominal pain.
- In any woman whose normal activities are disturbed by dysmenorrhoea unrelieved by NSAids, endometriosis should be suspected.
- If an ectopic pregnancy is suspected and there are no facilities for resuscitation, digital vaginal examination should be deferred for it may provoke rupture.[2]
- Acute abdominal and pelvic pain in the presence of a negative μ-HCG is most often due to an ovarian cyst.
- A positive μ-HCG plus an empty uterus and an adnexal mass are the classic diagnostic features of ectopic pregnancy.

WHEN TO REFER

- All cases of 'unexplained infertility'
- All teenagers with dysmenorrhoea sufficient to interfere with normal school, work or recreational activities, and not responding to prostaglandin inhibitors
- Patients with dysmenorrhoea reaching a crescendo mid menses
- Patients with dysmenorrhoea and unexplained bowel or bladder symptoms
- Patients with positional dyspareunia
- Patients with cyclic pain or bleeding in unusual sites

Note: Pelvic disease that can be treated by advanced laparoscopy surgery includes ectopic pregnancy, ovarian cysts, endometriosis and endometriomas, fibromyomata, pelvic adhesions and hydrosalpinx.

An approach to the management of acute abdominal and pelvic pain in premenopausal women is given in Figure 93.7 (on p. 1002).


Age
Pubertal
Reproductive age group
Ovarian cyst ? appendicitis ? functional
Pregnancy test (β-HCG)
Positive
Negative
Ultrasound
Intrauterine pregnancy
No intrauterine pregnancy
Pelvic mass
Pelvic tenderness
Observe
Refer for laparoscopy
Refer for laparoscopy ± laparotomy
Observe for 24 hours
No pathology
Appendicitis
Placental abruption
Inflammatory bowel disease
Constipation
Tubal ectopic pregnancy
Refer for laparoscopy
Pelvic inflammatory disease

Figure 93.7 An approach to management of acute abdominal and pelvic pain in premenopausal women. After Forbes.
[2]

REFERENCES

1. Foy A, Brown R. Chronic lower abdominal pain in gynaecological practice. Update, 1987; 27 March: 19 - 25.
2. Forbes KL. Lower abdominal and pelvic pain in the female: a gynaecological approach. Mod Med Aust, 1991; September: 24 - 31.
3. Soo Keat Khoo. Lower abdominal pain in women. Patient management supplement, 1990; August: 13 - 23.
4. Mackay EV et al. *Illustrated Textbook of Gynaecology* (2nd edn). Sydney: WB Saunders, Bailliere Tindall, 1992: 514 - 24.
5. Barton S ed. *Clinical Evidence*. London: BMJ Publishing Group, 2001: 1255 - 63.
6. O'Connor DT. Endometriosis. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 170 - 2.
7. Hager WD, Eschenbach DA et al. Criteria for diagnosis and grading of salpingitis. Obstet Gynecol, 1983; 61: 113 - 14.
8. Spicer J (Chair). *Th erapeutic Guidelines: Antibiotc* (Version 13). Melbourne: Th erapeutic Guidelines Ltd, 2006: 101 - 4.

CHAPTER 94. PREMENSTRUAL SYNDROME

I'm tired of all this nonsense about beauty being only skin deep. That's deep enough.

What do you want - an adorable pancreas?

JEAN KERR 1961

PREMENSTRUAL SYNDROME

Premenstrual syndrome (PMS) is defined as a group of physical, psychological and behavioural changes which begin 2 - 14 days before menstruation and are relieved immediately the menstrual flow begins.[1]

These symptoms occur in the luteal phase of the menstrual cycle yet the pathogenesis of PMS is stilluncertain. Among the proposed causes are pyridoxine deficiency, excess prostaglandin production and increased aldosterone production in the luteal phase.[2] However, PMS is most probably a disorder of ovarianfunction with a relativeexcess of oestrogen the main determinant.[3]

KEY FEATURES

- PMS increases in incidence after 30 years, witha peak incidence in the 30 - 40 year age group.
- PMS also occurs in the 45 - 50 year age group when it may alternate with menopausal symptoms, causing clinical confusion.[4]
- The symptoms of PMS decrease in severity just before and during menstruation.
- The symptoms cannot be explained by the presence of various psychological or psychiatric disorders.
- The severe form of PMS is now classified in the *Diagnostic and Statistical Manual of Mental Disorders*(4th edn) as Premenstrual Dysphoric Disorder (PMDD).

Incidence

Up to 90% of women may experience premenstrualsymptoms, which can vary from minor to severe. Interestingly, up to 15% of women can feel better premenstrually.[5] Statistics from countries such as Sweden, the UnitedStates and the United Kingdom indicate that up to 40% of women are significantly affected.[6] About 5 - 10% of women experience such severesymptoms that PMS disrupts their quality of life.

Aetiology

Various aetiological factors have been identified as contributing to PMS.[2]

Predisposing factors:

- mental illness
- alcoholism
- sexual abuse
- family history
- stress

Precipitating factors:

- hysterectomy
- tubal ligation
- cessation of the OCP

Sustaining factors:

- diet containing caffeine, alcohol, sugar

- smoking
- stress
- sedentary lifestyle

Symptoms

Various symptoms from among the 150 reportedare summarised in Figure94.1.


Psychological symptoms
• insomnia
• moodiness
• irritability
• anxiety
• tension
• depression
• confusion
• food cravings
Physical symptoms
nausea
headache
dizziness
hot flushes
acne
breast swelling and tenderness
bloated abdomen
constipation
swelling of ankles

Figure 94.1 Symptoms of premenstrual tension

The most common symptoms are depression 71%, irritability 56%, tiredness 35%, headache 33%, bloatedness 31%, breast tenderness 21%, tension 19% and aggression/violence 13%.[7] Other important symptoms include weight gain, lowered performance, decreased libido and feeling out of control.

Classification of PMS

It is convenient to classify PMS in terms of severity of symptoms.[4]

1. *Mild:*symptoms signal onset of menstruation. No medical advice sought or needed.
2. *Moderate:*symptoms annoying but insufficient to interfere with function at home or work. Medical advice sought in about one-third.
3. *Severe:*symptoms are such that functions at work or home are disrupted. Medical advice is usually sought. This disruptive form is labelled PMDD (see Table 94.1).

Table 94.1 Summary of PMDD criteria[*]

ASymptoms must occur during the week before menses and remitafew days after onset of menses. Five of the following symptoms must be present, with at leastone being 1, 2, 3 or 4.

1. Depressed mood or dysphoria
2. Anxiety or tension
3. Affective lability
4. Irritability
5. Decreased interest in usual activities
6. Concentration difficulties
7. Marked lack of energy
8. Marked change in appetite, overeating, or food cravings
9. Hypersomnia or insomnia
10. Feeling overwhelmed
11. Other physical symptoms (e.g. breast tenderness, bloating)

B Symptoms must interfere with work, school, usual activities or relationships

C Symptoms must not merely be an exacerbation of another disorder

D Criteria for A, B and C must be confirmed by prospective daily ratings for at least two cycles

[*] Adapted from *DSM-IV-TR*.

Differential diagnosis[4]

- Menopause syndrome
- Mastalgia
- Other causes of fluid retention - kidney or adrenal
- Thyroid disorder (hyper- or hypo-activity)
- Polycystic ovary syndrome (PMS may be a feature of oestrogen excess)
- Psychiatric disorders: depression, mania

Diagnosis

- Thorough history - including diet, exercise habits, psychosocial background, emotional influences and family history.
- Menstrual calendar - for 3 months, showing timing of the three main symptoms.[4]
- Physical examination to exclude gynaecological, endocrine or other systemic disease; and also include:
  - breast examination
  - vaginal examination and Pap smear
- Investigations (if considered appropriate, perform one or more serological tests):
  - thyroid function tests
  - serum progesterone and oestradiol in midluteal phase of three representative cycles
  - electrolytes and creatinine
  - prolactin - if galactorrhoea or oligomenorrhoea present

Management

The basic aim of management is to reassure and treat the woman in such a way that she makes changes in her lifestyle to cope with the hormonal dysfunction rather than rely on medication. Themanagement strategies include the following.

Explanation, reassurance and insight

Cognitive based therapy is very helpful for the patient to understandthe nature of her symptoms and to receive appropriate support and rapport. Advise her to be open about her problem and inform her family and close friends about her symptoms.

Keeping a diary[4]

Advise the patient to keep a daily diary of all her symptoms and when they occur over a 2 - 3 month period. This information should help her to plan around her symptoms: for example, avoid too many social events and demanding business appointments at the time when PMS symptoms are worst.

Dietary advice[4]

Advise the patient to eat regularly and sensibly; eat small rather than large meals and aim for ideal weight (if necessary).

Increase amount of complex carbohydrates (whole grains, vegetables and fruit), leafy green vegetables and legumes.

Decrease or avoid: refined sugar, salt, alcohol, caffeine (tea, coffee, chocolate), tobacco, red meat and excessive fluid intake during premenstrual phase. Decrease total protein to 1 g/kg/day; decrease fats.

Exercise

Recommend a program of regular exercise such as swimming, aerobics, jogging or tennis. Such exercise has been proven to decrease depression, anxiety and fluid retention premenstrually.[8]

Relaxation

Advise patients to plan activities that they find relaxing and enjoyable at the appropriate time. Consider stress reduction therapy, including appropriate counselling.

Appropriate dress

Advise sensible dressing to cope with breast tenderness and a bloated abdomen, such as a firm- fitting brassiere and loose-fitting clothes around the abdomen.

Medication

Pharmaceutical agents that have been used with success in some patients and little or no relief in others include diuretics (e.g. spironolactone), vitamins and minerals (e.g. pyridoxine and evening primrose oil), antiprostaglandin preparations (e.g. mefenamic acid, indomethacin), bromocriptine, danazol (suppresses ovulation), GnRH agonists and hormone preparations such as the OCP, progestogens and oestradiol implants. A combination of agents may have to be used.

The evidence base for symptom relief so far is as follows:[9, 10]

- nil or negative - evening primrose oil,[11] gingko biloba, progesterone/progestogens, OCP, bromocriptine
- weak - magnesium, calcium, vitamin E, vitex angus castus (chaste tree)
- moderate - vitamin B_6,[12] St John's wort, spironolactone
- strong (for PMDD) - SSRI agents and clomipramine, GnRH agonists, danazol

If contraception is required it is appropriate to use a monophasic OCP but there is conflicting interest about the efficacy of the OCP.

Mild to moderate symptoms

- Pyridoxine (vitamin B_6) 100 mg daily

Moderate to severe symptoms

- Fluoxetine 20 mg daily for 10 - 14 days before menstruation, or 10 - 20 mg daily continuously[13] *or*
  non-SSRI: clomipramine 25 mg (o) nocte for 2 cycles, increasing if necessary

Individualised therapy[14]

- PMS + fluid retention - use spironolactone 100 mg daily 3 days before expected onset of symptoms to day 1 of menstruation
- PMS + severe mastalgia - consider danazol 200 mg (o) daily from onset of symptoms to onset of menses
- PMS + dysmenorrhoea - mefenamic acid 500 mg tds from onset of symptoms to onset of menses

No matter what medication is taken, up to 70% of women will report animprovement in the early months of treatment, suggesting that a strong placebo factor is involved in management.[15]

Practice tip

Moderate to severe PMDD

- Fluoxetine 20 mg (o) daily for 14 days before menstruation

If the recommended approach of support, education, reassurance and stress management is still not effective.

WHEN TO REFER[4]

- Refer to a gynaecologist if underlying disease is suspected or proven (e.g. polycystic ovary disease, endometriosis)
- Consider referral if prescribing of danazolis contemplated
- Refer to an endocrinologist if an endocrinedisorder such as adrenal, pituitary or thyroid is suspected or proven
- Consider referral if depression or psychosis worsens or is not cyclical

Practice tips

- Keeping a daily diary of symptoms is very helpful for both patient and clinician.
- Aim for lifestyle changes and commonsense non-pharmacological management.
- Allow at least three cycles of treatment to provide a reasonable time for a particular medication.
- Drugs such as danazol are second-line drugs with significant side-effects and should be used with caution.
- High doses of pyridoxine, such as 500 mg a day, are associated with peripheral neuropathy so the dosage should be kept at around 100 mg/day.
- Be careful of overdiagnosing PMS and overlooking disorders such as depression, which may be exacerbated in the premenstrual phase.

REFERENCES

1. Shreeve CM. *The Premenstrual Syndrome.*Th orson Publishers Ltd, 1983: 23 - 4.
2. Smith MA, Yong Kin EQ. Managing the premenstrual syndrome. Clin Pharmacokinet, 1986; 5: 788 - 97.
3. Brush MG. Pharmacological rationale for the management of PMS. J of Psychosom Obstet and Gynaecol, 1983; 2 - 1: 35 - 9.
4. Smith M. Premenstrual syndrome. In: *MIMS Disease Index.* Sydney: IMS Publishing, 1991 - 92: 439 - 41.
5. Farrell E. How to treat: Menstrual disorders. Aust Dr Weekly, 1990; 25 May: IV - VI.
6. Labrum AH. Hypothalamic, pineal and pituitary factors in the premenstrual syndrome. J Reprod Med, 1983; 28(7): 438 - 45.
7. Dalton K. *The Premenstrual Syndrome and Progesterone Therapy.*London: Heinemann, 1984: 3.
8. Tierney LM et al. *Current Medical Diagnosis and Treatment* (41st edn). New York: The McGraw-Hill Companies, 2002: 747 - 8.
9. Primary care reports. American Health Consultants, 1997; 3(16): 146 - 9.
10. Wyatt K et al. Premenstrual syndrome. Clinical Evidence, 2000; 4: 1121.
11. Budeiri DJ et al. Is evening primrose oil of value in the treatment of premenstrual syndrome? Controlled Clin Trials, 1996; 17: 60 - 8.
12. Wyatt K et al. Efficacy of viamin B6 in the treatment of premenstrual syndrome: systemic review. Br Med J,1999; 318:

1375.

13. Stone AB, Perlstein TB, Brown WA. Fluoxetine in the treatment of premenstrual syndrome. Psychopharmacol Bull, 1990; 26: 331 - 5.
14. Roughan P. Premenstrual syndrome. Curr Ther, 1995; 36(11): 53 - 9.
15. Abraham S. The premenstrual syndrome. Mod Med Aust, 1992; September: 80 - 6.

CHAPTER 95. THE MENOPAUSE AND OSTEOPOROSIS

Every woman should use what Mother Nature gave her before Father Time takes it away.

LAURENCE J. PETER 1977

DEFINITIONS

The menopause is the cessation of the menses for longer than 12 months. In most Western women it occurs between the ages of 45 and 55 years, with an average age of 50 - 51 years.

The WHO has defined the menopause as signifying the permanent cessation of menstruation, resulting from the loss of ovarian follicular activity.[1] However, the term is used in a broader sense to include the perimenopausal phase when ovarian function waxes and wanes and the periods become irregular. This may last 2 - 5 years and sometimes longer and involves the premenopausal and menopausal phases.

The postmenopause is the period following the menopause but cannot be defined until after 12 months of spontaneous amenorrhoea, except in women who have had an oophorectomy.

Surgical menopause is known as bilateral oophorectomy.

Summary

The climacteric can be subdivided into four phases:

Phase 1 Premenopausal: up to 5 years before the last menstrual period.

Phase 2 Perimenopausal: the presence of early menopausal symptoms with vaginal bleeding (usually irregular)

Phase 3 Menopausal: the last menstrual period

Phase 4 Postmenopausal: approximately 5 years after the menopause

Osteoporosis

Osteoporosis, which literally means porous bone, is reduced bone mass per unit volume. Osteoporosis is usually addressed in the context of the menopause because it is found mainly in postmenopausal middleaged and elderly women and can be largely prevented by correcting oestrogen deficiency.

PHYSIOLOGY OF THE MENOPAUSE

Inspection of Figure 95.1 provides an overview of how menopausal symptoms are related to ovarian follicular activity and hormonal activity.



Figure 95.1 Schematic representation of some clinical, biological and endocrinological features of the perimenopausal and postmenopausal phases. After Burger.

The number of ovarian primary follicles declines rapidly as the menopause approaches, with few if any being identifiable following the cessation of menstruation. In the postmenopause phase FSH rises to levels 10 - 15 times that of the follicular phase of the cycle while LH levels rise about threefold. The ovary secretes minimal oestrogen but continues to secrete significant amounts of androgens.

An uncomfortable effect of oestrogen withdrawal, often not appreciated by medical practitioners, involves urogenital problems where the epithelium of the vagina, vulva, urethra and the base of the bladder becomes thin and dry, leading possibly to dysuria and frequency, itching, dyspareunia and atrophic bleeding. Hormone replacement therapy (HRT) can ameliorate these urogenital dysfunctions.

CLINICAL FEATURES

Because small amounts of oestrogen are still being produced in the adrenal glands, symptoms other than cessation of periods may be mild or absent.

Symptoms

Vasomotor:[2]

- hot flushes (80%)
- night sweats (70%)
- palpitations (30%)
- lightheadedness/dizziness
- migraine

Psychogenic:

- irritability
- depression
- anxiety/tension
- tearfulness
- loss of concentration
- poor short-term memory
- unloved feelings
- sleep disturbances

- mood changes
- loss of self-confidence

Urogenital (60%):

- atrophic vaginitis
- vaginal dryness (45%)
- dyspareunia
- decline in libido
- bladder dysfunction (e.g. dysuria)
- stress incontinence/prolapse

Musculoskeletal:

- non-specific muscular aches
- non-specific joint aches and pains

Skin and other tissue changes:

- dry skin
- formication (17%)
- new facial hair
- breast glandular tissue atrophy

Other:

- unusual tiredness
- headache

Clinical approach

A thorough evaluation of the patient is important, including a good history.

History

Enquire about any symptoms related to oestrogen deficiency and about other related symptoms, with an emphasis on the menstrual history and hot flushes. Enquire about mental state symptoms, such as anger, irritability, depression, moodiness, loss of self-esteem and other such problems. Ask about sexual history, contraception, micturition and social history, including relationships.

Information on family history of osteoporosis, cancer and cardiovascular disease should be sought.

Physical examination

The general examination should include measurement of blood pressure, weight and height, breast palpation, abdominal palpation, vaginal examination and Pap smear. Note the texture of the vaginal epithelium.

Investigations[2]

Apart from a Pap smear, the following tests should be considered:

- urinalysis
- full blood count, lipids including HDL
- liver function tests
- mammography (all women, preferably before or after 3 months on HRT)
- diagnostic hysteroscopy and endometrial biopsy only if undiagnosed vaginal bleeding
- bone density study (if risk factors)

If diagnosis is in doubt (e.g. perimenopause, younger patient < 45 years, hysterectomy):

- serum FSH diagnostic

- serum oestradiol diagnostic

Differential diagnosis of menopause syndrome

- Depression
- Anaemia
- Thyroid dysfunction
- Hyperparathyroidism
- Gynaecological disorders
    - dysfunctional uterine bleeding

Management

Education and lifestyle

Patients should receive adequate understanding, support and explanation with the emphasis being that the menopause is a natural fact of life. For many, menopausal symptoms will be relatively shortlived; often within 2 years. Emphasise the importance of leading a healthy lifestyle:

- correct diet
- avoid obesity
- adequate relaxation
- adequate exercise
- reduced smoking
- reduced caffeine intake
- reduced alcohol intake
- do pelvic floor exercises regularly

Sexual activity

Advise that it is normal and appropriate to continue sexual relations, using a vaginal lubricant for a dry vagina. Contraception is advisable for 12 months after the last period. The OCP can be used up to 50 - 51 years if there are no risk factors.

HORMONE REPLACEMENT THERAPY

The modern treatment of HRT at the menopause not only reduces climacteric symptoms and enhances the quality of life in the short term but also prevents osteoporosis and fractures. On the other hand, there is evidence of increased risk of endometrial hyperplasia, breast cancer and thromboembolism, especially with long-term use. The future of HRT as a health measure remains uncertain. Further evaluation is awaited.

The Womens Health Initiative (WHI) study[3]

The WHI trial conducted in the United States investigated the use of long-term HRT in postmenopausal women with an intact uterus using long-term combined oral oestrogen and progesterone. The trial did not include people using other forms of HRT, such as patches, gels or implants. The findings were that there was a slightly increased risk of breast cancer (1.26-fold), coronary heart disease (1.29-fold), stroke (1.41-fold) and pulmonary embolism (2.13-fold) with prolonged use greater than 5 years.[4] The study also found a reduction in risk of bowel cancer and fracture in these women.

Points to emphasise:

- there is no firm evidence of an increased risk of breast cancer with use of HRT < 5 years
- HRT cannot be recommended in asymptomatic menopausal women to prevent osteoporosis
- women choosing to cease HRT should reduce their dose gradually over 2 - 3 months

- women without a uterus taking oestrogen are in a different category and further clarification is required

Indications for HRT

- Distressing symptoms
- Significant osteoporosis

HRT has to be tailored to the individual patient and depends on several factors, including the presence of a uterus, individual preferences and tolerance.[5]

The hormones to consider are:

- oestrogen
- progestogen, and
- testosterone

If perimenopausal, the OCP or sequential HRT (non-contraceptive) can be used. If menopausal - use HRT.

Oestrogen

Oestrogen comes in various preparations: oral, patches, implants, injections and topical vaginal preparations (Table 95.1). Injectables are not very effective so the common modes of administration are oral, implants or skin patches. Transdermal patches are the most favoured mode worldwide.

Table 95.1 Oestrogens used in the menopause[5]

Generic name	Proprietary name/s (examples)	Daily dose range	Usual daily protective dose
Oral			
Conjugated equine oestrogen	Premarin	0.3 - 2.5 mg	0.625 mg
Oestradiol valerate	Progynova	1.0 - 4.0 mg	2.0 mg
Oestriol	Ovestin	1.0 - 4.0 mg	2.0 mg
Piperazine oestrone sulphate	Ogen	0.625 - 5 mg	1.25 mg
Implants			
Oestradiol	Oestradiol implants	20 - 100 mg	50 mg
Skin patch			
Oestradiol	Various	25 - 100 μg every 3½ to 7 days	50 μg
Topical gel			
Oestradiol 0.1%	Sandrena	0.5 - 1.5 mg	1 mg
Nasal spray			
Oestradiol	Aerodiol	150 - 600 μg	300 μg
Vaginal preparations			
Creams			
Oestriol 1 mg/g	Ovestin	0.5 g	0.5 g
Pessaries			
Oestradiol	Vagifem	25 μg	1 pessary (25 μg)
Oestriol	Ovestin ovula	0.5 mg	0.5 mg

Note: Vaginal therapy is usually given continuously for 2 weeks, then twice weekly.

Vaginal creams or tablets are usually restricted to women who have mild menopausal symptoms and a dry vagina or urethra, or who cannot tolerate parenteral medication. Most women find the use of vaginal pessaries and creams messy and the older preparations are heavily absorbed, but the new oestradiol tablet, Vagifem, is a very effective topical therapy. The oral oestrogens in common use are Premarin, Ogen and Progynova. Implants of 50 - 100 mg (usually 50 mg) of oestradiol are given 3 - 12 monthly; patches (usually 50 μg) are applied weekly or every three and a half days. Continuous daily oestrogen use is recommended; there is no reason to stop therapy for 1 week.

Progestogen

Progestogen is given to women with a uterus and may be given continuously or cyclically. If it is not given, many women will develop hyperplasia of the uterus and there is a 5 - 10 times increased risk of endocervical cancer with unopposed oestrogen therapy.[2] If given cyclically (postmenopausal) it is given for the first to the twelfth day of the calendar month, generally as Provera or Primolut N (Table 95.2). A withdrawal bleed will occur, which many elderly women find unacceptable. Thus, continuous therapy may be more appropriate. Avoid continuous use in perimenopausal women because of heavy irregular bleeding.

Table 95.2 Progestogens used in the menopause[5]

Generic name	Daily dose range	Usual daily protective dose
Dydrogesterone	10 - 20 mg	10 mg
Medroxyprogesterone acetate	2.5 - 20 mg	10 mg
Norethisterone	1.25 - 5 mg	2.5 mg

Progestogen alone can be given for menopausal symptoms in a woman with an oestrogen-dependent tumour.

Progestogens should be given in the smallest possible dose, to prevent endometrial hyperplasia.

Testosterone

Testosterone is usually reserved for women whose libido does not improve with HRT but this indication is controversial. It is given as an implant of 50 mg and will last 3 - 12 months. An oestrogen implant of 50 mg should be given concurrently.

Tibolone

This is a selective tissue oestrogenic activity regular with combined oestrogenic, progestogenic and androgenic properties that can be used as an alternative in postmenopausal women. Adverse effects with breakthrough bleeding and virulisation are a concern.[6]

Dose: tibolone 2.5 mg (o) daily

Contraindications to HRT

Important contraindications to HRT are listed in Table 95.3. The main absolute contraindications are active oestrogen-dependent neoplasms, such as endometrial and breast carcinoma, acute thrombophlebitis and undiagnosed abnormal vaginal bleeding. HRT is protective for bowel cancer but not ovarian cancer.

Table 95.3 Contraindications (absolute or relative) to HRT

Oestrogen-dependent tumour
• endometrial cancer
• breast cancer

Recurrent thromboembolism
History of coronary artery disease
Uncontrolled hypertension
Undiagnosed vaginal bleeding
Active liver disease
Active SLE
Otosclerosis
Acute intermittent porphyria

HRT regimens

Some commonly used regimens are presented in Figure 95.2. Regimens A and B are in common use. The transdermal system appears to be the most favoured although some women find it unsuitable. A useful regimen, especially for irregular bleeding in the perimenopausal phase, is the combined sequential pill which can be continued for several years if necessary. Sufficient oestrogen should be given to control symptoms and prevent osteoporosis.


Days
1
15
1
15
A. Postmenopausal
oestrogen daily
progestogen 12 days
1–2 days gap
3–4 days bleed
B. Postmenopausal (continuous)
oestrogen daily
progestogen daily
C. Perimenopausal
oestrogen 25 days
progestogen 12 days (from day 13)
1–2 days gap
3–4 days bleed

Figure 95.2 Possible HRT regimens for women with a uterus. After Farrell.[7]

There is a proliferation of combination packs with permutations of various oestrogens and progestogens (pills and patches), usually sequential, some combined. These preparations are most suitable for perimenopausal women. The continuous combined preparation is best suited to women at least 1 year postmenopausal. The newer progesterone, cyproterone acetate, is regarded as useful in the presence of hirsutism.

Informed consent

HRT should be prescribed only after the woman has been informed of the regimens available, their relative benefits and risks and side-effects.[5] It must be emphasised that HRT, especially the combined sequential formulation, is not a contraceptive.

Side-effects of therapy[8]

In the first 2 - 3 months the woman may experience oestrogenic side-effects, but these usually resolve or stabilise.

Starting with a lower dose may minimise these side-effects.

Premenstrual syndrome (in 15%)

Action: decrease progestogen dose *or* change to alternative progestogen

Nausea and breast disorders

Cause: initial sensitivity to oestrogen

Action: reduce oestrogen to starting dose *or* intravaginal oestrogen

Bleeding problems

Heavy bleeding

Action: decrease oestrogen

Breakthrough bleeding

Action: increase progestogen

Irregular bleeding

Action: investigate + endometrial sampling

Intolerance of bleeding

Action: use continuous regimen

No bleeding

Action: reassure that this is not a problem

Leg cramps

Action: decrease oestrogen

FOLLOW-UP AFTER COMMENCING HRT[7]

- Three months (ideal time for mammography if not performed beforehand)
- then 6 monthly

Allow 6 months to stabilise therapy.

DURATION OF TREATMENT

The duration of treatment depends on several factors, including the severity of symptoms, the response to therapy and the long-term aims, such as osteoporosis prevention (at least 10 years) (Figure 95.3). However, long-term therapy should be an informed decision made by the patient in consultation with the doctor. A useful working rule is to aim for treatment for a maximum of 2 years and then review. Current evidence is that combined oestrogen and progestogen therapy will not increase the risk of breast cancer for up to 5 years of use, and oestrogen-only therapy will not increase risk for at least 7.2 years.[8]



Figure 95.3 Graphic illustration of the effects of HRT on the fracture threshold of ageing women

NON-HRT OPTIONS FOR OSTEOPOROSIS PREVENTION

These include raloxifene (a selective oestrogen- receptor modulator), tibolone and the biphosphonates.

NON-HORMONAL THERAPY REGIMENS

Several non-hormonal regimens have been employed to manage menopausal symptoms. These include medical therapies, such as clonidine, tranquillisers and antidepressants, and also natural therapies, such as evening primrose oil, soy products and other phytoestrogens (plants containing oestrogen-like compounds). Although there is considerable anecdotal evidence about the efficacy of phytoestrogens, evidence from long-term double-blind trials is lacking. However, one study by Murkies et al.[9] showed the positive effect of these foods on hot flushes.

There are three broad classes of phytoestrogens - isoflavones, lignans and coumestans. Isoflavones, which are found almost exclusively in legumes, include soy products, lentils and beans. Lignans are found in varying degrees in a wide variety of plants but particularly in cereals, fruit and vegetables. Coumestans occur mainly in alfalfa and soy bean sprouts.

Dr Murkies advises 'a wholegrain diet with porridge or cereal for breakfast, perhaps a tablespoon of linseed on top, wholemeal bread for lunch with sprouts and fruit, especially apple, and an assortment of vegetables and legumes as part of the evening meal'.

EVIDENCE BASE FOR PREVENTION OF MENOPAUSAL SYMPTOMS AND OSTEOPOROSIS

A systematic review from Northern America [10] concluded 'all middle aged women should be counselled about the benefits and risks of HRT. Smoking cessation, low-fat and high-fibre nutrition, 1500 mg calcium daily, 400 - 800 IU vitamin D daily and moderately strenuous weight-bearing exercises every day are recommended'.

Clinical Evidence reports that there is evidence that oestrogen relieves symptoms of urogenital atrophy and vasomotor symptoms, improves short-term quality of life but that long-term use is associated with an increased risk of thromboembolic disease and endometrial and breast cancers. Further RCTs have found that tibolone relieves vasomotor symptoms and improves sexual symptoms and clonidine reduces hot flushes.[11]

Natural remedies

Studies by Professor Eden's group[12] conducted trials for menopause symptoms with several natural products, including red clover (Promensil) and black cohosh extract (Remifemen). Poor evidence was found for red clover, wild yam, evening primrose oil, dongquai and topical progesterone. The only herbal treatment that his team has recommended is black cohosh extract. Furthermore, herbal medicines cannot be recommended for the prevention or treatment of osteoporosis.[12]

An Australian review investigating the effect of natural remedies on bone mineral density commented that there is good evidence that exercise increases bone mineral density in postmenopausal women with osteoporosis, little good quality empirical evidence to support the use of natural progesterone cream and insufficient evidence to support the use of boron, cod liver oil or chelated calcium supplements (as opposed to calcium carbonate).[13]

Phytoestrogens

Limited evidence from restricted RCTs suggests that soy flour, which contains phytoestrogens, may relieve vasomotor menopausal symptoms.[11]

New herbal products

There are a variety of products, such as Menocalm and Menoease, containing herbal agents with a satisfactory evidence base in PMS, including St John's Wort, chaste tree extract (vitex angus castus) and black cohosh. These products await appropriate reviews.

OSTEOPOROSIS

Osteoporosis refers to the increased bone fragility that accompanies ageing and many illnesses. Following the menopause, women begin to lose calcium from their bone at a much faster rate than men, presumably as a direct response to low levels of oestrogen. Within 5 - 10 years of the menopause, women can be seen to suffer from osteoporosis and by the age of 65 the rate of fractures in women has increased to 3 - 5 times that of men (Figure 95.4).



Figure 95.4 Osteoporosis is reduced bone mass per unit volume

Facts and figures

- Osteoporosis is commonest in postmenopausal women.
- Up to 15% of women will develop fractures in their lifetime and 30% of all women reaching 90 years of age will suffer a hip fracture.[14]
- Osteoporosis leads to reduced bone strength and susceptibility to fracture, even with minor trauma.
- Osteoporosis only causes pain when complicated by fracture.
- First presentation is usually a fracture (Colles', femoral neck and vertebra) or height shrinkage.
- Vertebral collapse is the hallmark of osteoporosis.

- The disorder is of bone mass, not calcium metabolism.
- For osteoporosis in a vertebra including a pathological fracture, multiple myeloma may need exclusion.
- The first step in prevention is regular exercise and an adequate dietary intake of calcium (1500 mg per day).

Classification of osteoporosis

Primary

Type 1: postmenopausal (vertebral or distal forearm fractures within first 20 years of menopause)

Type 2: senile osteoporosis (fracture of proximal femur usually 15 - 30 years after menopause)

Osteopenia is 1 - 2.5 SD below the young adult standard mean.

Osteoporosis is > 2.5 SD below this mean.

Secondary

Secondary to various endocrine disorders, malabsorption and malignancies. Various causes and risk factors are presented in Table 95.4.

Table 95.4 Osteoporosis: risk factors and/or causes

Female
Thin build; low BMI < 19
Race: Asian, Caucasian
Family history (e.g. maternal hip fracture < 75 yrs)
Premenopausal oestrogen deficiency (e.g. amenorrhoea)
Early menopause < 45 years (natural or surgical)
Cigarette smoking can be modified
High caffeine intake > 4 cups per day can be modified
High alcohol intake > 2 standard drinks per day can be modified
Low calcium intake can be modified
Physical inactivity can be modified
Chronic corticosteroid use can be modified
Eating disorders (e.g. anorexia nervosa)
Anticonvulsants
Endocrine disorders
• Cushing's syndrome
• diabetes mellitus
• hyperparathyroidism
• thyrotoxicosis
• hypogonadism
• acromegaly
Coeliac disease
Chronic kidney failure
Prolonged immobilisation

Investigations

- Plain radiography is of limited value. Osteoporosis is not detectable until 40 - 50% of bone is lost.
- 25 hydroxy vitamin D (most useful test): normal range 40 - 160 pmol/L.
- Plasma calcium, phosphate and alkaline phosphatase are all usually normal.
- Consider tests for multiple myeloma in an osteoporotic area.
- Densitometry can predict an increased risk of osteoporosis and fracture, the best current modality being dual energy X-ray absorptimetry (DEXA scan) in a facility with high-standard quality control.[6] The spine and femoral neck are targeted: the femoral neck is the most useful index.

Treatment for osteoporosis

The goal of treatment is to prevent osteoporosis or reduce further loss. No treatment has been shown to replace lost bone effectively. Anabolic agents such as nandrolone deconate may reduce further loss but the side-effects are problematic.

Medications of value in decreasing further loss[6]

The following medications may be valuable in preventing further bone loss, possibly reversing the osteoporosis process and preventing further fractures.

- HRT (short-term subject to contraindications) *or*
- calcitriol (a vitamin D metabolite)
    - use for failed trial HRT
    - very useful for corticosteroid induced osteoporosis (avoid calcium supplements during therapy) *or*
- biphosphonates (decrease bone absorption)
    - alendronate 10 mg (o) daily or 70 mg (o) once weekly (take care with potential side-effect of oesophagitis)
    - etidronate
    - risedronate
- raloxifene (a selective oestrogen-receptor modulator) (SERMs) 60 mg (o) daily
- strontium ranelate
    - 2 g (o), as a powder in water, daily

The choice depends on the clinical status, such as the age of the patient and the extent of disease, the patient's tolerance of drugs and further clinical trials of these drugs. The one preferable solution is to give prophylaxis for individuals identified as high risk and the only widely accepted proven therapy is oestrogen therapy but it has limitations.

Recommendations for prevention[6]

- HRT within 2 years of the onset of the menopause
- Adequate dietary intake of calcium:
  800 mg per day - premenopause
  1500 mg per day - postmenopausal

Calcium-rich foods include low-fat calciumenriched milk (500 mL contains 1000 mg), other low-fat dairy products (e.g. yoghurt or cheese), fish (including tinned fish such as salmon with the bone), citrus fruits, sesame and sunflower seeds, almonds, brazil nuts and hazel nuts.

Practice tips

- Careful pretreatment assessment is important.
- Encourage conservative self-help management with an emphasis on lifestyle if symptoms are mild.
- Explain benefits and risks and get informed consent.
- Individualise HRT therapy.

- Regular follow-up is essential.
- Allow about 6 months to stabilise with HRT.
- The prime treatment for an oestrogendeficiency disorder is oestrogen.
- Use oestrogen-only therapy for women without a uterus.
- If a uterus is present give combined oestrogen - progestogen therapy (cyclical or continuous).
- Avoid giving progestogen in the presence of continuing ovarian activity.
- Always start with a low dose of oestrogen.
- Women who have experienced side-effects such as migraine with the COC may have the same problem with HRT.
- Problematic loss of libido can be treated with testosterone in the short term (e.g. as a single parenteral dose or as a short course of oral tablets).
- Oestrogen deficiency results in a loss of elasticity and dryness of the vagina, which can be partially helped with HRT.
- HRT does not always restore the sex drive but does help make sexual intercourse easier and more pleasant.

Calcium supplements will be necessary in postmenopausal women (except if taking calcitriol). Calcium citrate is better absorbed than carbonate.

- Exercise: moderate exercise against gravity - walking (brisk walking for 30 minutes four times a week), jogging or tennis - may make a small contribution to retarding bone loss.
- Lifestyle factors: stop smoking and limit alcohol and caffeine intake.
- Vitamin D and sunlight: there is evidence we need significant exposure to sunlight to produce a gentle ‘pinking’ of the face arms and hands to produce natural vitamin D, e.g. 5 - 15 minutes a day in warm - hot climates, 25 - 50 minutes a day in winter in temperate climates.[5]
- Adequate nutrition: keep BMI > 19.
- Attention to falls prevention, including avoiding sedative medication.
- The evidence for ‘hip protectors’ is not convincing.

WHEN TO REFER

- A problem arises in establishing the correct regimens for HRT
- Complications not corrected by routine measures develop with HRT
- Osteoporosis appears to be secondary to an underlying illness
- Advice is required about the management of a patient with pathological osteoporotic fractures or loss of height

REFERENCES

1. World Health Organization. Technical Report Series 670. WHO Scientific Group, Research on the Menopause. Geneva: 1981.
2. Wren B. Menopause. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 303 - 6.
3. Writing Group for the Women's Health Initiative Investigators. Risks and benefits of oestrogen plus progestin in healthy postmenopausal women: Principal results from the women's health initiative randomised controlled trial. JAMA, 2002; 288(3): 321 - 33.
4. RACGP. HRT advice. Aust Fam Physician, 2002; 31(8): 733 - 4.
5. NH&MRC. Hormone replacement therapy: exploring the options for women. Canberra: NH&MRC, 2005.
6. Moulds RFW (Chair). *Therapeutic guidelines: Endocrinology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2001: 91 - 100.
7. Farrell E. Treatment options and menopause regimens. Aust Fam Physician, 1992; 21: 240 - 6.
8. Baber R. The menopause: update. Medical Observer, 2006; 28 July: 31 - 3.

9. Murkies AL et al. Dietary flour supplementation decreases postmenopausal hot flushes: effect of soy and wheat. Maturitas, 1995; 21: 189 - 95.
10. Prosser WW, Shafir MS. *Evidence-Based Family Medicine*. Hamilton: BC Decker, 1998: 168.
11. Rymer J, Morris E. Menopausal symptoms. In: Barton S ed. *Clinical Evidence*. London: BMJ Publishing Group, 2001: 1304 - 8.
12. Eden J. Managing menopause - HRT or herbal? Mod Med Aust, 1999; August: 32 - 5.
13. Del Mar CB et al. Natural remedies for osteoporosis in postmenopausal women. Med J Aust, 2002; 176: 182 - 3.
14. Seeman E, Young N. Osteoporosis. In: *MIMS Disease Index* (2nd edn). Sydney, IMS Publishing, 1996: 368 - 71.

CHAPTER 96. VAGINAL DISCHARGE

In all cases of abnormal vaginal discharge consider the possibility of the sexually transmitted infections, gonorrhoea and non-specific urethritis.

VAGINAL DISCHARGE

Vaginal discharge is one of the commonest complaints seen by family physicians yet it is one of the most difficult to solve, especially if it is recurrent or persistent. It is present if the woman's underclothes are consistently stained or a pad is required. It is important to make a proper diagnosis, to differentiate between abnormal (physiological) and pathological discharge and to be aware of the considerable variation in secretion of vaginal fluid.

The differential diagnoses should include consideration of normal discharge, vaginitis, either infective or chemical, STIs, and urinary tract infection.

KEY FACTS AND CHECKPOINTS

- A survey of a large family planning clinic found that 17% of women complained of vaginal discharge.[1]
- Vaginal discharge may present at any age but is very common in the reproductive years.
- Vaginal discharge is a common presentation of those STIs responsible for PID.
- The first step in diagnosis is to determine if the discharge is cervical or vaginal in origin.
- One of the simplest methods of making a proper diagnosis is a wet film examination. It saves expensive laboratory investigations.

A DIAGNOSTIC APPROACH

A summary of the safety diagnostic model is presented in Table 96.1.

Table 96.1 Vaginal discharge: diagnostic strategy model

Q. Probability diagnosis
A. Normal physiological discharge Vaginitis • bacterial vaginosis 40 - 50% • candidiasis 20 - 30% • *Trichomonas* 10 - 20%
Q. Serious disorders not to be missed
A. Neoplasia • carcinoma • fistulas STIs/PID (i.e. cervicitis) • gonorrhoea • *Chlamydia* Sexual abuse, esp. children Tampon toxic shock syndrome (staphylococcal infection) Streptococcal vaginosis (in pregnancy)
Q. Pitfalls (often missed)

A. Chemical vaginitis (e.g. perfumes) Retained foreign objects (e.g. tampons, IUCD) Endometriosis (brownish discharge) Ectopic pregnancy (‘prune juice’ discharge) Poor toilet hygiene Genital herpes (possible) Atrophic vaginitis Threadworms
Q. Seven masquerades checklist A. Depression - Diabetes Drugs Anaemia - Thyroid disorder - Spinal dysfunction - UTI (association)
Q. Is the patient trying to tell me something? A. Needs careful consideration; possible sexual dysfunction.

Probability diagnoses

The two most common causes of vaginal discharge are physiological discharge and infective vaginitis.

Physiological discharge

Normal physiological discharge is usually milky-white or clear mucoid and originates from a combination of the following sources:

- cervical mucus (secretions from cervical glands)
- vaginal secretion (transudate through vaginal mucosa)
- vaginal squamous epithelial cells (desquamation)
- cervical columnar epithelial cells
- resident commensal bacteria

The predominant bacterial flora are lactobacilli, which produce lactic acid from glucose derived from the epithelial cells. The lactic acid keeps the vaginal pH acidic (< 4.7). Other commensal bacteria include staphylococci, diptheroids and streptococci.

With physiological discharge there is usually no odour or pruritus.

In addition, the egg-white discharge accompanying ovulation may be noted. The discharge may be aggravated by the use of the pill. The normal discharge usually shows on underclothing by the end of the day. Clear or white, it oxidises to a yellow or brown on contact with air. It is increased by sexual stimulation.

Management:

- reassurance and explanation
- wear cotton underwear (not synthetic)
- bath instead of showering
- avoid douching and feminine deodorants

- use tampons instead of pads

Infective vaginitis

The commonest cause of infective vaginitis is bacterial vaginosis (formerly bacterial vaginitis, *Gardnerella vaginalis* or *Haemophilus vaginalis*) which accounts for 40 - 50% of cases of vaginitis.[2] *Candida albicans* is the causative agent in 20 - 30% of cases while *Trichomonas vaginalis* causes about 20% of cases in Australia. The comparable features are outlined in Table 96.2. Human papilloma virus infection of vaginal epithelium may cause excess discharge.

Table 96.2 Characteristics of discharge for important causes of abnormal vaginal discharge (after Weisberg)[2]

Infective organism	Colour	Consistency	Odour	pH (normal 4 - 4.7)	Associated symptoms
Candida albicans	White	Thick (cream cheese)	None	4	Itch, soreness, redness
Trichomonas vaginalis	Yellow-green	Bubbly, profuse (mucopurulent)	Malodorous, fishy	5 - 6	Soreness
Bacterial vaginosis	Grey	Watery, profuse, bubbly	Malodorous, fishy	5 - 6	Irritation (sometimes)
Cervicitis	Yellow-green (coming from cervix)	Thick (mucopurulent)	Variable - usually malodorous	4 - 4.7	Signs of PID

Serious disorders not to be missed

The 'not to be missed' group includes carcinoma of the vagina, cervix or uterus and STIs, including PIDs caused by *Chlamydia trachomatis* and *Neisseria gonorrhoeae*. Vaginal discharge is the most common presenting symptom of both of these serious STIs. Occasionally, infections of the endometrium and endosalpinx will produce a discharge that gravitates to the vagina.[1] Benign and malignant neoplasia anywhere in the genital tract may produce a discharge. Usually it is watery and pink or blood-stained.

Inspection should include vigilance for fistulas that may be associated with malignancy, inflammation or postirradiation.

Pitfalls

It is common to overlook the problem caused by hygienic preparations. Apart from the vaginal tampon, which may be retained (knowingly or otherwise), there is a variety of preparations that can induce a sensitivity reaction. These include deodorant soaps and sprays and contraceptive agents, especially spermicidal creams. Ironically, the various preparations used to treat the vaginitis may cause a chemical reaction.

Endometriosis of the cervix or vaginal vault may cause a bloody or brownish discharge.

Seven masquerades checklist

Of this group, diabetes mellitus leading to recurrent 'thrush', drugs causing a local sensitivity, and urinary tract infection have to be considered (Table 96.1).

Psychogenic considerations

This question needs to be answered, especially if the discharge is normal. The problem could be related to sexual dysfunction or it may reflect a problematic relationship, and the issue may need to be explored diplomatically. Vaginal discharge is an embarrassing problem for the patient and any discussion needs to be handled thoroughly and sensitively. A relevant sexual history may satisfactorily solve the problem.

THE CLINICAL APPROACH

History

The history is important and should include:

- nature of discharge: colour, odour, quantity, relation to menstrual cycle, associated symptoms
- exact nature and location of irritation
- sexual history: arousal, previous STIs, number of partners and any presence of irritation or discharge in them
- use of chemicals, such as soaps, deodorants, pessaries and douches
- pregnancy possibility
- drug therapy
- associated medical conditions (e.g. diabetes)

Physical examination

Optimal facilities for the physical examination include an appropriate couch and good light, bivalve Sims' specula, sterile swabs (preferably with transport media), normal saline, 10% potassium hydroxide (KOH), slides and cover slips and microscope. Inspection in good light includes viewing the vulva, introitus, urethra, vagina and cervix. Look for the discharge and specific problems such as polyps, warts, prolapses or fistulas. To differentiate between vaginal and cervical discharge, wipe the cervix clear with a cotton ball and observe the cervix. A mucopurulent discharge appearing from the endocervix may be the clue to an STI such as *Chlamydia* and gonorrhoea. Perform a pH test and a wet film.

Pitfalls to keep in mind include:

- The patient may have had a bath or a ‘good wash’ beforehand and may need to return when the discharge is obvious.
- A retained tampon may be missed in the posterior fornix, so the speculum should slide directly along the posterior wall of the vagina.
- *Candida* infection may not show the characteristic curds, ‘the strawberry vagina’ of *Trichomonas* is uncommon and bubbles may not be seen.

Acetic acid 2% is useful in removing the discharge and mucus to enable a clearer view of the cervix and vaginal walls.

Investigations

- pH test with paper of range 4 to 6
- Amine or ‘whiff’ test: add a drop of 10% KOH to vaginal secretions smeared on glass slide
- Wet film microscopy of a drop of vaginal secretions

A culture is necessary if no diagnosis is made after this routine.

A full STI work-up

- First-pass urine and ThinPrep samples - for *Chlamydia* and gonorrhoea PCR
- swabs from the cervix for *Chlamydia, N. gonorrhoeae*
  - swab mucus from cervix first
  - swab endocervix
  - place in transport media
- Pap smear
- Viral culture (herpes simplex)
  - scrape base of ulcer or, ideally, deroof a vesicle
  - immediately immerse in culture medium
  - transport rapidly to laboratory
- Group B *Streptococcus*
  - swabs from endocervix, urethra, rectum

Preparation of a wet film

To make a wet film preparation[2] (Figure 96.1), place one drop of normal saline (preferably warm) on one end of an ordinary slide and one drop of 10% KOH on the other half of the slide. A sample of the discharge needs to be taken with a swab stick, either directly from the posterior fornix of the vagina or from discharge that has collected on the posterior blade of the speculum during the vaginal examination. A high vaginal swab is required for *C. albicans*. A small amount of the discharge is mixed with both the normal saline drop and the KOH drop. A cover slip is placed over each preparation. The slide is examined under low power to get an overall impression, and under high power to determine the presence of lactobacilli, polymorphs, trichomonads, spores, clue cells and hyphae. A summary of various findings on wet film examination is presented in Table 96.3.



Figure 96.1 Wet film method

Table 96.3 Wet film examination (after Weisberg)[2]

	Lactobacilli	Polymorphs	Epithelial cells	Clue cells	Other
Normal	+	None or occasional	+	-	
Candidiasis	+	None or occasional	+	-	Spores/hyphae
Trichomoniasis	Absent or scant	Numerous	+	-	Trichomonads
Bacterial vaginosis	Absent or scant	Numerous	+	2 - 50%	

Lactobacilli are long, thin Gram-positive rods; clue cells are vaginal epithelial cells that have bacteria attached so that the cytoplasm appears granular and often the entire border is obscured. They are a feature of bacterial vaginosis. Trichomonads are about the same size as polymorphs and to distinguish between the two one needs to see the movement of the trichomonad and the beating of its flagella under high power of the microscope. Warming the slide will often precipitate movement.

Refer to Figure 96.2.



Figure 96.2 Relative sizes of various cells or organisms as seen in a wet smear

Other investigations

Gram-stain smear and culture should be contemplated only if a diagnosis cannot be made on wet film.

VAGINAL DISCHARGE IN CHILDREN

Most newborn girls have some mucoid white vaginal discharge. This is normal and usually disappears by 3 months of age. From 3 months of age to puberty, vaginal discharge is usually minimal.[3] Staining on a child's underclothes may be due to excess physiological discharge, especially in the year before the menarche.[1] Vulvovaginitis is the most common gynaecological disorder of childhood, the most common cause being a non-specific bacterial infection (see p. 1030).

VAGINAL DISCHARGE IN THE ELDERLY

Vaginal discharge can occur in the elderly from a variety of causes, including infective vaginitis, atrophic vaginitis, foreign bodies, poor hygiene and neoplasia. It is important to exclude malignancy of the uterus, cervix and vagina in the older patient.

Atrophic vaginitis

In the absence of oestrogen stimulation the vaginal and vulval tissues begin to shrink and become thin and dry. This renders the vagina more susceptible to bacterial attack because of the loss of vaginal acidity. Rarely, a severe attack can occur with a very haemorrhagic vagina and heavy discharge:

- yellowish non-offensive discharge
- tenderness and dyspareunia
- spotting or bleeding with coitus
- the vagina may be reddened with superficial haemorrhagic areas

Treatment

- Oral HRT
- Local oestrogen cream or tablet (e.g. Vagifem) The tablet is preferred as it is less messy *or*
- Zinc and castor oil soothing cream

VAGINAL CANDIDIASIS

Infection with the fungus *C. albicans* is a common and important problem with a tendency to recurrence. However, with the widepread use of over-the-counter antifungals, resistant non-albicans species, such as *Candida glabrata* (in particular), *Candida parapsilosis* and *Candida tropicalis* are becoming more common.[4]

Clinical features

- Intense vaginal and vulval pruritus
- Vulval soreness
- Vulvovaginal erythema (brick red)
- Vaginal excoriation and oedema
- White curd-like discharge (Figure 96.3)



Figure 96.3 Vaginal candidiasis showing typical adherent thick milky white vaginal discharge

- Discomfort with coitus
- Dysuria

Factors predisposing to vaginal candidiasis[1]

Endogenous

- Diabetes mellitus
- AIDS syndrome
- Pregnancy
- Debilitating diseases

Exogenous

- Oral contraceptives
- Antibiotics
- Immunosuppressants
- Carbohydrate-rich diet
- Orogenital/anogenital intercourse

- IUCD
- Tight-fitting jeans
- Nylon underwear
- Humidity/wet bathing suit

Treatment[6]

For the first attack of candidiasis it is appropriate to select one of the large range of vaginal imidazole therapies (clotrimazole, econazole, isoconazole, miconazole) for 1 - 7 days (Table 96.4). There appears to be no significant difference between imidazoles.[5] Nystatin is best reserved for recurrent cases or if there is local reaction to the imidazoles. Some therapists prefer creams to tablets and pessaries because cream can be applied to any tender vulval area, but a tablet and cream can be used simultaneously, especially for a heavy infection.

Table 96.4 Treatment of vaginal candidiasis

		Vaginal therapy	
Length of treatment	**Generic name**	**Tablets**	**Cream (5 g)**
Imidazoles			
1 day (statim dose)	clotrimazole	500 mg × 1	
	isoconazole	300 mg × 2	
3 days	clotrimazole	100 mg × 2	
	clotrimazole		2%
	econazole	150 mg	
	econazole		1.5%
	miconazole	200 mg	
6 days	clotrimazole	100 mg	
	clotrimazole		
7 days	miconazole	100 mg	2%
Nystatin			
7 days	nystatin	100 000 U	100 000 U
Oral therapy for recurrent recalcitrant infections		**Oral tablets**	
1 day	fluconazole	150 mg	
14 days	ketoconazole	200 mg daily	
14 days	nystatin	500 000 U tds	
14 days	fluconazole	50 mg	
14 days	itraconazole	100 mg	

Gentian violet (0.5% aqueous solution) is useful for rapid relief, if available.

Recommended initial regimen:[6]

clotrimazole 500 mg vaginal tablet as a single dose ± clotrimazole 2% cream applied to vulvovaginal and perineal areas 8 - 12 hrly (for symptomatic relief)

An alternative regimen, especially for recurrent infections:

nystatin pessaries twice daily for 7 days *and/or* nystatin vaginal cream (100 000 U per 5 g) twice daily for 7 days

Recalcitrant cases (proven by microscopy and if not pregnant)[6]

fluconazole 150 mg (o) as a single dose *or* itraconazole 100 mg (o) once daily for 14 days

Note: A male sexual partner does not usually require treatment.[4] If symptomatic (usually balanitis in an uncircumcised male), treat with clotrimazole 1% + hydrocortisone 1% topically, 12 hourly until 2 weeks after symptoms resolve.

Candida (Torulopsis) glabrata[6]

A significant number of cases of recurrent vulvovaginal candidiasis are due to non-albicans species of *Candida. Candida glabrata* is the commonest nonalbicans species, which exhibit reduced susceptibility to azoles. In resistant infections use boric acid 600 mg (in a gelatin capsule) intravaginally for 10 to 14 days.

Do not use in pregnancy.

Advice to the patient

- Bathe the genital area gently two or three times a day for symptomatic relief. In preparing for the antifungal preparation, use 1 - 3% acetic acid or sodium bicarbonate solution (1 tablespoon to 1 litre of water). Thoroughly cleanse the vagina, including recesses between rugae and the fornices and also the folds around the vulva.
- Dry the genital area thoroughly after showering or bathing.
- Wear loose-fitting cotton underwear.
- Avoid wearing pantyhose, tight jeans or tight underwear or using tampons.
- Avoid having intercourse or oral sex during the infected period.
- Do not use vaginal douches, powders or deodorants or take bubble baths.

TRICHOMONAS VAGINALIS

This flagellated protozoan, which is thought to originate in the bowel, infects the vagina, Skene's ducts and lower urinary tract in women and the lower genitourinary tract in men. It is transmitted through sexual intercourse and is relatively common in the female after the onset of sexual activity.

Clinical features

- Profuse thin discharge (grey to yellow-green in colour) (Figure 96.4)



Figure 96.4 Trichomonas vaginitis showing profuse thin greyish discharge with erythema of the vaginal walls

- Small bubbles may be seen in 20 - 30%
- Pruritus
- Malodorous discharge
- Dyspareunia
- Diffuse erythema of cervix and vaginal walls
- Characteristic punctate appearance on cervix

Treatment[6]

- Oral metronidazole 2 g as a single dose (preferable) *or* 400 mg bd for 5 days (if relapse) *or*
- tinidazole 2 g as a single dose (Table 96.5)
- Use clotrimazole 100 mg vaginal tablet daily for 6 days during pregnancy
- Attention to hygiene
- The sexual partner must be treated simultaneously
- The male partner should wear a condom during intercourse
- For resistant infections a 3 - 7 day course of either metronidazole or tinidazole may be necessary

Table 96.5 Oral treatment for bacterial vaginosis and *Trichomonas* infections

Length of treatment	Generic name	Oral dosage
1 day (statim dose)	tinidazole	500 mg × 4
	metronidazole	400 mg × 5
7 days (for recurrent)	tinidazole	500 mg daily
	metronidazole	400 mg bd

BACTERIAL VAGINOSIS

Bacterial vaginosis is a clinical entity of mixed aetiology characterised by the replacement of the normal vaginal microflora (chiefly *Lactobacillus*) with a mixed flora consisting of *Gardnerella vaginalis*, other anaerobes such as *Mobiluncus* species, and *Mycoplasma hominis*. This accounts for the alkalinity of the vaginal pH.

Clinical features

- A grey, watery, profuse discharge
- Malodorous
- No obvious vulvitis or vaginitis
- Liberates an amine-like, fishy odour on admixture of 10% KOH (the amine whifftest)
- Clue cells
- ± Dyspareunia and dysuria
- ± Pruritus

Treatment[6]

- Metronidazole 400 mg (o) bd for 5 days or 2 g stat

Clindamycin 300 mg (o) bd for 7 days or 2% clindamycin cream can be used for resistant infections or during pregnancy. Normal vaginal pH can be restored using a variety of topical douches such as povidone iodine solution (1 tablespoon per litre of water), vinegar (3 - 4 tablespoons per litre of water), topical Acigel or a milky solution of yoghurt to restore *Lactobacillus* levels.

There is no evidence that treatment of the male sexual partner reduces the recurrence rate or provides any significant benefit.[7] The STI treatment guidelines of the US Centers for Disease Control state explicitly that such treatment is of no proven benefit.[8]

GROUP B STREPTOCOCCUS VAGINOSIS

Group B *Streptococcus* (*Streptococcus agalactiae*) is a commensal in up to 40% of healthy humans. It is a problem if detected in the pregnant woman because of serious infection in the neonate. In certain at-risk circumstances, such as premature rupture of the membranes or a previous infected neonate, give:

- benzylpenicillin 1.2 g IV stat, then 600 mg IV 4 hourly until delivery

In the non-pregnant woman give amoxycillin 500 mg (o) tds for 7 days if there is significant pyogenic infection.

RETAINED VAGINAL TAMPON

A retained tampon, which may be impacted and cannot be removed by the patient, is usually associated with an extremely offensive vaginal discharge. Its removal can cause considerable embarrassment to both patient and doctor.

Method of removal

Using good vision the tampon is seized with a pair of sponge-holding forceps and quickly immersed under water without releasing the forceps. A bowl of water (an old plastic icecream container is suitable) is kept as close to the introitus as possible. This results in minimal malodour. The tampon and water are immediately flushed down the toilet if the toilet system can accommodate tampons. An alternative method is to grasp the tampon with a gloved hand and quickly peel the glove over the tampon for disposal.

TAMPON TOXIC SHOCK SYNDROME

Staphylococcal infection

This rare dramatic condition is caused by the production of staphylococcal exotoxin associated with tampon use for menstrual protection. The syndrome usually begins within 5 days of the onset of the period.

The clinical features include sudden onset fever, vomiting and diarrhoea, muscle aches and pains, skin erythema, hypotension progressing to confusion, stupor and sometimes death.

Management

Active treatment depends on the severity of the illness. Cultures should be taken from the vagina, cervix, perineum and nasopharynx. The patient should be referred to a major centre if 'shock' develops. Otherwise the vagina must be emptied, ensuring there is not a forgotten tampon, cleaned with a povidone iodine solution tds for 2 days, and flucloxacillin or vancomycin antibiotics administered.

- These women should not use tampons in the future.

Prevention

- Good general hygiene with care in handling and inserting the tampons.
- Change the tampon 3 - 4 times a day.
- Use an external pad at night during sleep.

WHEN TO REFER

- Evidence of sexual abuse in children to an experienced sexual assault centre
- Recurrent, recalcitrant infections
- Presence of carcinoma or fistula
- Staphylococcal toxic shock syndrome

Practice tips

- Failure of treatment may be due to diagnostic error, therapeutic error, sexual reinfection, chemical sensitivity to vaginal tablets, drug resistance or depressed host immunity.
- Patients with an infective cause appreciate the use of patient education material, especially that including preventive measures.
- Advise patients subject to vaginal infection about simple hygiene measures to keep the area dry and cool: avoid nylon underwear, pantyhose, tight jeans, wet swimsuits, perfumed soaps and vaginal deodorisers.
- A serious sequela is subsequent dyspareunia and vaginismus. The patient should be advised to have sufficient lubrication, such as KY jelly, to avoid distressing psychosexual problems.

REFERENCES

1. Mackay EV et al. *Illustrated Textbook of Gynaecology* (2nd edn). Sydney: WB Saunders, Bailliere Tindall, 1992: 296 - 325.
2. Weisberg E. Wet film examination. Aust Fam Physician, 1991; 20: 291 - 4.
3. Tunnessen WW Jr. *Signs and Symptoms in Paediatrics* (2nd edn). Philadelphia: Lippincott, 1988: 458 - 60.
4. Sobel JD. Vulvovaginal candidiasis. In: Holmes KK et al. eds. *Sexually Transmitted Diseases* (3rd edn). New York: McGraw-Hill, 1999: 629 - 39.
5. O'Neill S, Howard S. Recurrent vulvovaginal candidiasis. Aust Fam Physician, 1989; 18: 102.
6. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 110 - 6.
7. Vejtorp M, Bollreup AC, Vejtory L et al. Bacterial vaginosis: a double-blind randomised trial of the effect of treatment of the sexual partner. Br J Obstet Gynaecol, 1988; 95: 920 - 6.

8. US Department of Health and Human Services, Public Health Service, Centers for Disease Control. *Sexually Transmitted Diseases: Treatment Guidelines.* MMWR, 1989; 38: S - 8.

CHAPTER 97. VULVAR DISORDERS

Genital skin is very sensitive. This sensitive organ needs protection from chemical and physical damage. The genital area is also affected by the way you feel and symptoms can appear worse at times of stress.

EXTRACT FROM PATIENT INFORMATION SHEET 'THE DO'S AND DON'TS OF GENITAL HYGIENE ', DERMATOLOGY/VULVAL DISEASES CLINIC, MERCY HOSPITAL FOR WOMEN, MELBOURNE

VULVAR DISORDERS

The dermatoses are the predominant cause of vulvar problems and this chapter focuses mainly on the important female genital skin conditions.

The vulva is that part of the female external genitalia lying posterior to the mons pubis, comprising the labia majora, labia minora, clitoris, vestibule of the vagina, vaginal opening and bulbs of the vestibule.[1]

The vaginal vestibule is an almond-shaped opening between the lines of attachment of the labia minora. The clitoris marks the superior angle and the fourchette the inferior boundary. It is approximately 4 - 5 cm long and 2 cm in width.[1] The four main structures that open into the vestibule are the urethra, vagina and the two secretory ducts of Bartholin's glands. The surface is composed of delicate stratified squamous epithelium.

The genital area is affected by dermatoses found elsewhere on the skin but management is rendered more complex by the sensitivity and thinness of the skin, a tendency to superinfection, in addition to the psychological problems, including the often-resultant dyspareunia.

The vulval area, which is innervated by nerves arising from L1 - 2 and S2 - 4 nerve roots, is sensitive to noxious stimuli but the vagina is not sensitive to pain.[2] Topical creams, soaps, perfumes and other toilet products irritate the vulva easily - it is an area prone to contact dermatitis.

Clinical manifestations of vulvar disorders include itching, pain, irritation, white mucosal patches, lichenification, erosions and intertrigo[3] (Table 97.1).

Table 97.1 Vulvar discomfort/irritation: diagnostic strategy model

Q. Probability diagnosis
A. Atopic dermatitis
Chronic vulvovaginal candidiasis
Irritant contact dermatitis (e.g. douches, bubble baths)
Allergic contact dermatitis (e.g. perfumes, topical antimicrobials)
Fissuring from the above dermatoses
Trauma - 'dry' coitus
Q. Serious disorders not to be missed
A. Neoplasia
Squamous cell carcinoma
- Melanoma
- Lymphomas etc →pruritus
Infection
- Streptococcal vulvovaginitis
- Herpes simplex virus
Vulval vestibular syndrome

Q. Pitfalls (often missed)
A. Lichen sclerosis Urinary incontinence →ammonical vulvitis Faecal soiling Tinea cruris Trichomonal vaginitis Atrophic vaginitis Aphthous ulcers Dysaesthetic vulvodynia Psoriasis Lichen planus Infestations: - Threadworms - Pubic lice - Scabies
Q. Seven masquerades list
A. Depression Diabetes Drugs Anaemia - Thyroid disorder - Spinal dysfunction ?dysaesthesia UTI
Q. Is the patient trying to tell me something?
A. Common: psychosexual problems.

KEY FACTS AND CHECKPOINTS

- If a dermatosis is suspected, check the skin on the body.
- Vestibular hypersensitivity (vulvar vestibular syndrome) is a distressing, reasonably common condition that gives superficial dyspareunia. Diagnosis is by an abnormal response to light touch, even by a cotton bud.
- The vestibule can exhibit pearly papules (the equivalent of pearly penile papules) that look like tiny regular warts - they are normal.
- Urinary incontinence or faecal incontinence may be the basis of the problem in the vulvar area. It can be misdiagnosed as urinary tract infection.
- Avoid using topical mixed steroids and antimicrobial applications (e.g. Kenacomb) to treat vulvar dermatoses.
- Approximately 20% of women carry *Candida albicans* as genital flora but less than 5% suffer from repeated or intractable clinical candidiasis.
- Not all itching and burning of the vulva and vagina is *Candida* infection. Swabs should be taken for diagnosis before committing to treatment empirically.
- The cause of vulvar irritation may be multifactorial (e.g. atopic dermatitis or *Candida* with irritant or allergic contact dermatitis from applications).

- Be alert for malignant melanoma and be aware that an area of benign pigmentation with welldemarcated edges and bluish discolouration called melanocytosis vulvae can develop.

DERMATITIS

As expected, the various common forms of dermatitis collectively represent the prime cause of a pruritic irritating vulvar skin disorder. They classically present with itching, burning and soreness initiated by scratching. The manifestations can vary from symptoms without a rash to a rash without the above symptoms.

The causes of vulvar dermatitis are:

- atopic dermatitis
- irritant contact dermatitis
- allergic contact dermatitis
- seborrhoeic dermatitis
- corticosteroid-induced dermatitis
- psoriasis

Principles of management[4]

- Take an appropriate history, including atopy, skin diseases.
- Check allergens and irritants (e.g. panty liners, soap, bubble bath, perfumed toilet paper, douches, perfumes, condoms, tea-tree oil).
- Check for heat and friction (e.g. synthetic or tight underwear, tight denim jeans, sporting costumes/tights, sweating, vigorous activity - bicycle riding).
- Check gynae-urological history (e.g. oestrogen status, faecal or urinary incontinence, vaginal discharge, 'thrush').
- Check psychosexual history (e.g. dyspareunia, partnership issues, depression).
- Carefully inspect the vulva plus the rest of the skin, scalp and nails. Look for lichenification (Table 97.2).

Table 97.2 Causes of common vulvar signs

White patches	• Lichen simplex (lichenification) • Lichen sclerosus • Lichen planus • Leucoplakia • Carcinoma
Erosions and ulcers	• Herpes simplex virus • Lichen sclerosus • Lichen planus • Carcinoma • Various uncommon dermatoses • Excoriated scabies
Intertrigo	• Atopic dermatitis • *Candida albicans* • Seborrhoeic dermatitis • Tinea • Erythrasma

- Psoriasis

- Appropriate investigations: vaginal swab, PAP smear, perhaps patch testing and vulval biopsy for a rare, premalignant or suspected malignant condition.

Treatment[3, 4]

- Provide supportive education and counselling.
- Correction of underlying factors (e.g. tight clothes, incontinence, anal discharge, overused topical medications and cosmetics).
- Treat any secondary infection.
- Use aqueous cream moisturiser as cleanser.
- Start with potent class III topical steroid and follow with 1% hydrocortisone.

PSORIASIS[5]

Psoriasis can affect the genital or perianal area (especially the natal cleft) and appears as a glazed, beefy red plaque without the classic scale seen elsewhere. There may be minimal or no sign of psoriasis on the skin of the body.

The main symptom is itching. It is usual to take swabs to rule out infection.

Treatment[5]

- Avoid irritants and use a soap substitute.
- First apply a potent topical steroid (e.g. methylprednisolone aceponate) - continue until resolution of rash.
- Second (when controlled) apply LPC 2% in aqueous cream bd, slowly increasing the strength up to 8%. If not tolerated, use ichthammol 2% in aqueous cream.

Note: Maintenance with topical steroids - hydrocortisone 1% or resume potent agent for a flare up.

LICHEN PLANUS[6]

Genital lichen planus is relatively uncommon but may affect both the vulva and vagina and can occur in association with oral lesions. Vulvar lesions appear as whitish reticulated papules and plaques. A delicate white lacy pattern at the periphery of the lesion is a distinctive feature. Erosions may occur.

Symptoms include pruritus, vaginal discharge, dyspareunia (if erosions) and postcoital bleeding. Differential diagnoses of vulvar lichen planus include other causes of desquamative and erosive lesions, such as lichen sclerosis, pemphigus vulgaris, bullous and cicatricial pemphigoid and erythema multiforme.

A biopsy is required to make the diagnosis. Treatment is difficult. Topical steroids provide symptomatic relief and there are a variety of treatment trials, including the encouraging topical cyclosporin.

LICHEN SCLEROSUS[4, 5]

Also known as lichen sclerosis et trophicus, this uncommon chronic inflammatory dermatosis of unknown aetiology (perhaps an autoimmune disorder) presents as well-defined white, finely wrinkled plaques that almost exclusively affects the anogenital skin, although it can occur anywhere on the body. It spares the vagina. It can run a chronic and complicated course with development of squamous cell carcinoma (SCC) in about 4% a concern. The differential diagnosis is atrophic vaginitis.

DxT: genital pruritus + soreness + white wrinkled plaques = lichen sclerosus

Features

- Bimodal peak: prepubertal girls, perimenopause.
- Mean age of onset in adult women is 50 years.
- Pruritus is main symptom.
- Soreness, burning, dyspareunia.

Examination

- Variable distribution
- White wrinkled plaques
- Purpuric and ulcerated areas
- May show figure of 8 pale perianal and perivaginal area

Complications if untreated

- Vulval atrophy and labial (even clitoral hood) fusion
- Lifetime risk of SCC 2 - 6%

Management[5]

- Confirm diagnosis by biopsy (tend to avoid in children).
- Based on potent topical corticosteroids (e.g. betamethasone dipropionate 0.05% ointment or cream applied bd for 4 weeks, then daily for 8 weeks, with applicator - show patients where to apply, using a mirror).
- Reduce to a potent topical steroid once daily for next 3 months, then reduce to hydrocortisone 1% ointment or cream applied daily for long term.
- Lifelong surveillance with 6-monthly check-up.
- A similar topical program is used in children.

INFECTIONS

Chronic vulvovaginal candidiasis

This is different to acute candidiasis and remains difficult to treat because there may be a localised hypersensitivity to *Candida*.

Features

- Chronic vulval itch - scratch cycle.
- Burning, swelling - premenstrual exacerbation.
- Dyspareunia.
- Discharge not usually present.
- Aggravated by courses of systemic antibiotics.

Management[5]

- Swab - low vaginal - with each suspected episode, especially if discharge.
- Aim for symptom remission with continuous antifungal treatment:
    - topical vaginal antifungals (imidazoles or nystatin), or
    - daily oral antifungals (monitor liver function tests) until symptoms clear - ketaconazole 200 mg/day or fluconazole 50 mg/day, or itraconazole 100 mg/day (then weekly for 6 months)
- Relieve itching with hydrocortisone 1% (do not use stronger preparations).

- Use nystatin pessaries in pregnancy.

Streptococcal vulvovaginitis

Swabs from the vulva or vagina can find Streptococcal species or *Staphylococcus aureus*, which are treated accordingly. *Streptococcus* infection may cause a vaginal discharge. It usually presents as an acute beefy red, sore vulva and vagina or a low-grade vulvitis.

Treatment[5]

- Phenoxymethylpenicillin 250 mg (o) 6 hourly for 10 days, or
- Oral antibiotic as indicated on sensitivity testing (e.g. amoxycillin or roxithromycin)
- Topical mupirocin may help prevent recurrences

Tinea

Tinea causes an annular spreading rash with an active border that spreads from the labia to the thigh (see tinea cruris, p. 1154). A problem is the development of tinea incognito from the application of topical steroids. This lacks central clearing but the active margin can be seen. Skin scrapings are necessary for diagnosis. Treatment is with a topical imidazole (avoid nystatin) or oral agents if resistant or extensive.

PRURITUS VULVAE

The causes of an itchy vulva to consider are:

- candidiasis (rash, cottage cheese discharge)
  - broad-spectrum antibiotics
  - diabetes mellitus
  - contraceptive pill
- poor hygiene and excessive sweating
- tight clothing
- sensitivity to soaps, cosmetics and contraceptive agents
- overzealous washing
- local skin conditions
  - psoriasis
  - dermatitis/eczema (uncommon cause)
- post-anal conditions (e.g. haemorrhoids)
- infestations
  - threadworms (children)
  - scabies
  - pediculosis pubis
- infections
  - *Trichomonas*
  - urinary tract infection
  - genital herpes
- menopause: due to oestrogen deficiency
- topical antihistamines
- vulval carcinoma
- psychological disorder (e.g. psychosexual problem, STI phobia)

Treatment is according to the causation.

Management

This depends on the primary cause (e.g. candidiasis, incontinence), which should be treated effectively.

General measures (advice to patients)[3]

- Attend to hygiene and excessive sweating.
- Avoid overzealous washing.
- Take showers of no more than 5 minutes duration.
- Avoid water too hot (lukewarm preferable).
- Avoid toilet soap - use a soap alternative (e.g. aqueous cream, cetaphillotion) and wash it off with water only.
- Use soap alternatives (e.g. Dove, Neutrogena) for rest of body.
- Pat the skin dry after the shower (avoid harsh drying).
- Keep the genital area dry and wash thoroughly at least once a day.
- Do not wear tight pantyhose, tight jeans or tight underwear, or use tampons.
- Do not use vaginal douches, powders or deodorants.
- After the toilet, wipe gently with a soft, noncoloured, non-perfumed toilet paper or baby wipe (e.g. Dove).
- Apply a good moisturiser (e.g. Hydraderm or 5% peanut oil in aqueous cream).

Treatment

- For pruritus, apply cool moisturising cream (kept in refrigerator) when there is an urge to scratch.
- Apply prescribed steroid ointment to the rash.

VULVOVAGINITIS IN PREPUBERTAL GIRLS

Vulvovaginitis is the most common gynaecological disorder of childhood. It can affect women of any age but is particularly common in girls, especially between the ages of 2 and 8 years. It is a type of dermatitis of the vulva and the vagina.

Mild vulvovaginitis

Symptoms of this very common problem include:

- discomfort and soreness
- vulval itching
- redness (Figure 97.1)



Figure 97.1 Typical area of inflammation of vulvovaginitis in girls

- discharge - usually a slight yellow discharge on the underwear
- dysuria

It is important not to confuse it with a urinary infection when the child is clearly uncomfortable due to stinging on urination.

The cause is usually due to a low-grade inflammation in an area with a possible underlying dermatological disorder such as atopic dermatitis, psoriasis and lichen sclerosis, leading to sensitivity to various irritants such as soaps and urine.

Affected girls are often ‘atopic’.

The causal factors include:

- thin vaginal mucosa (the normal prepubescent state)
- dampness from synthetic fibre underwear, tight clothing, wet bathers, obesity
- lack of hygiene
- frequent self-handling, especially with irritation
- irritants (soap residue, bubble baths, antiseptics, chlorinated water)
- ‘sandbox’ vaginitis: girls sitting and playing in sand or dirt may develop irritation from particulate matter trapped in the vagina

Management

- Explanation and reassurance to parents
- Avoidance of the above causal factors, especially wet bathers, synthetic underwear, bubble baths, perfumed soaps and getting overweight
- Attention to good supervised toileting practice
- Attention to bathing and drying
- Regular warm baths (rather than showers)

It is worth soaking the child in a warm shallow bath containing half a cup of white vinegar.

Alternatively, bicarbonate of soda (10 g/10 L water) can be used.

Soothing creams such as soft paraffin creams and nappy rash creams such as zinc and castor oil cream should be applied three times daily as a short-term measure. If a powder is required use zinc oxide (e.g. Curash).

Moderate/persistent vulvovaginitis

The symptoms may be more intense with increased itching, burning and discharge.

Important causes to consider[7]

- ‘Sandbox’ vaginitis
- Skin disorders, especially atopic dermatitis and lichen sclerosus (look for skin problems elsewhere on body)
- Foreign body: consider if a bloody malodorous vaginal discharge
- Candidiasis - uncommon but consider if antibiotic therapy or possibility of diabetes
- Sexual abuse (uncommon but must not be missed)
- Pinworm infestation (*Enterobius*)
- Sexually transmissible organisms - usually postpubertal

Examination

A careful general examination should be performed only if considered appropriate. In infants the best examination method is to place the child on her mother's lap with the legs held well abducted. Lateral traction applied to the labia allows the hymen orifice to be examined. Look for vulval or vaginal infection. Aspirate vaginal secretion with a medicine dropper for appropriate cultures. A Pap smear is advisable for a persistent problem since a sarcoma is a possibility.

An older child can be placed in one of two suitable positions:

1. supine, legs apart in a frog-leg position, with bottom of feet touching (generally preferred)
2. prone, knee/chest position. This allows a better view of the hymenal orifice but many children do not like this position

A rectal examination may be performed to try to feel for suspected foreign bodies in the vagina.

Taking a swab[7]

If the discharge is profuse and offensive, take an introital swab (do not take a vaginal swab). Infective vulvovaginitis in girls is almost always due to a Group A beta-haemolytic streptococcus.

Treat with an appropriate antibiotic.

Treatment of dermatitis

Most cases of vulval dermatitis will respond to short courses 1% hydrocortisone ointment or cream, provided aggravating factors are removed.

Labial adhesions (labial agglutination)

Labial fusion is caused by adhesions considered to be acquired from vulvovaginitis after which sometimes the medial edges of the labia minoria become adherent. The adhesions are certainly not present at birth. Labial fusion is regarded as a normal variant and usually resolves spontaneously in late childhood. Provided the child is able to void easily, no treatment other than reassurance is needed.

For significant adhesions, others prefer the following approach:

- < 18 months - using EMLA cream. Separate with a blunt instrument. This can be distressing for the child and is followed by a high risk of recurrence.
- > 18 months - separate adhesions under general anaesthetic followed by application of Vaseline and/or oestrogen

cream.

However, these measures are not generally recommended.

VULVODYNIA

Vulvodynia describes the symptom of pain (burning, rawness or stinging) and discomfort, where no obvious cause can be found. Itch is not a feature. Causes include vestibular hypersensitivity (vulvar vestibular syndrome), dysaesthetic vulvodynia and various infections (e.g. herpes simplex virus). Virtually every condition of the vulva can be painful at times, even dermatitis can become painful if scratching or splitting leads to open areas and ulceration.

Vestibular hypersensitivity[2, 4]

Definition

Vestibular hypersensitivity is severe vulvar or vestibular pain on touch or entry into the vagina

Also referred to as vulvar vestibular syndrome (VVS) or vestibulitis, it is a very important disorder for the GP to be aware of in the woman with a typical history of introital dyspareunia. It is a difficult problem to treat. The characteristic feature is severe pain with vestibular touch, including attempted vaginal entry. The vestibule is very sensitive, featuring an inappropriate response to light touch. In many instances the cause of the primary condition is not apparent and a history of possible sexual abuse or other psychological provoking factors should be diplomatically elicited. Some patients can develop the problem after years of pain-free sex.[6] It is the most common cause of dyspareunia in the premenopausal female.

Secondary causes include inflammatory triggers such as irritant contact dermatitis and infection. This establishes a conditioned response.

Spontaneous resolution has been reported in up to 50% of cases. Prognosis appears to be reasonably good but depends to some extent on the premorbid personality of the patient.[3]

DxT: young♀ + nulliparous ♀ + dyspareunia = VVS

Features

- Delayed diagnosis (average 2 - 3 years).[4]
- Sexually active women in 20s and 30s.
- Pain provoked by intercourse, tampon insertion, tight underwear.
- Superficial 'entry' dyspareunia.
- Sexual dysfunction.
- Tender vestibule on light pressure.
- Erythema (usually minute red spots) around Bartholin's duct openings (consider *Candida*).

Diagnosis

Inappropriate tenderness to light touch with a cotton bud.

Management[4]

- Investigate underlying cause and treat it.
- Patient education, counselling and support.
- Physiotherapy - rehabilitation of pelvic floor musculature by increasing awareness and increasing elasticity of the tissues of the vaginal opening.
- Reassure that the condition is self-limiting.

- Encourage use of oil-based lubricants.
- Use bland emollients or 2% lignocaine gel prior to intercourse.
- If *Candida* present, treat with fluconazole 150 mg (o) weekly for 6 weeks.

Options

- Biofeedback technique
- Tricyclic antidepressants (start low, e.g. amitryptiline 10 - 20 mg nocte)
- Gabapentin
- Intralesional therapy:
  - triamcinolone
  - interferon
- Vestibulectomy (last resort):
  - excise tender vestibular tissue

DYSAESTHETIC VULVODYNIA[2, 8]

The typical patient with this neuropathic pain problem is a middle-aged to elderly woman who presents with a constant burning pain of the labia. It typically builds up during the course of the day. Examination is often unrewarding. The underlying cause may be pudendal neuralgia (may be secondary to pudendal nerve block), referred spinal pain or simply unknown.

Herpes simplex infection needs to be excluded.

Treatment options include antidepressants and gabapentin.

BARTHOLIN'S CYST

A Bartholin's gland swelling follows obstruction of the duct and presents as a painless vulval swelling at the posterior end of the labia majora, close to the fourchette. A simple non-infected cyst can be left alone and may resolve spontaneously. If it becomes infected an abscess may result, causing a painful, tender, red vulval lump. It may resolve with antibiotics or discharge spontaneously. If the cyst persists and becomes large a surgical marsupialisation procedure, which allows permanent drainage, can be performed (Figures 97.2a, b).



Figure 97.2 (a) Bartholin's cyst: starting the marsupialisation procedure **(b)**Postoperative appearance

WEBSITES REFERENCES

1. International Society for the Study of Vulvovaginal Disease < http://www.issvd.org >
2. Vulval Pain Society http://www.vulvalpainsociety.org

1. Thomas CL ed. *Taber's Cyclopedic Medical Dictionary* (14th edn). Philadelphia: FA Davis, 1997: 2083, 2100.
2. Fischer G. Vulvodynia. Aust Doctor, 2002; 31 May: i - viii.
3. Ang C, Sinclair R. Vulvar dermatoses: part I. Aust Doctor, 1998; 19 June: i - viii.
4. Welsh BM. Vulvar dermatoses. In: Proceedings of Dermatology Conference for General Practitioners. Melbourne: Combined Alfred Hospital/Skin and Cancer Foundation, 2002.
5. Mashford ML (Chair). *Therapeutic Guidelines: Dermatology*. Melbourne: Therapeutic Guidelines Ltd, 1999: 97 - 103.
6. Ang C, Sinclair R. Vulvar dermatoses: part II. Aust Doctor, 1998; 26 June: i - viii.
7. Fisher G. Vulval disease in childhood. Aust Prescr, 2005; 28: 88 - 90.
8. Welsh BM. Management of common vulval conditions. MJA, 2003; 178: 391 - 5.

CHAPTER 98. VIOLENCE AGAINST WOMEN

The root of Solomons seale when applied, taketh away in one or two nights, any bruise, blacke or blue spots gotton by falls or womens wilfulnesse, in stumbling upon their hasty husbands fists, or such like.

JOHN GERARD 1545 - 1612

VIOLENCE AGAINST WOMEN

Domestic violence basically means the physical, sexual or emotional abuse of one partner by the other, almost invariably abuse of a female by a male. However, the abuse can be of an elderly parent by the children or from some other member of the household to another member. It usually results from abuse and/or imbalance of power in close relationships. One person in the relationship consistently dominates or threatens with power and the abused victim gradually gives over more power.

A major problem in dealing with domestic violence is that it is hidden and the victims are reluctant to divulge the cause of their injuries when visiting medical practitioners.

KEY FACTS AND CHECKPOINTS[1]

- Between one-quarter and one-third of relationships experience violence at some time.
- In 90 - 95% of cases the victims are women.
- 10% of women have been violently assaulted in the last year.
- 22% of homicides in Queensland in 1982 - 87 were spouse murders.
- In violent families with children, 90% of children witness the violence and 50% of children are victims of violence.
- 4% of relationships will experience chronic domestic violence (in 20% this occurs before marriage).
- Less than 20% of those who abuse their spouse abuse some other person.[2]
- Alcohol is a factor in 50% of domestic violence incidents (i.e. not the sole cause; it does make violence easier, and is used as an excuse). Other factors include work stress/pressure, financial stress and illness. However, there are no excuses for domestic violence.
- Pregnancy is a high-risk time for victims of domestic battering.
- 50% of people know someone affected by domestic violence, but one-third refuse to speak about it or get involved in any way because they regard it as a ‘private matter’.
- One in five people think that domestic violence is acceptable in certain circumstances.

We usually think of domestic violence in terms of physical violence but it can take many forms.[3] These include:

- physical abuse
- psychological abuse
- economic abuse
- social abuse (e.g. isolation)
- sexual abuse

POSSIBLE PRESENTATIONS[4]

- Physical injuries: usually bruising caused by punching, kicking or biting; also fractures, burns, genital trauma
- Physical symptoms (e.g. back pain, headache, depression, sexual dysfunction, anxiety)
- Psychological problems (in both the woman and her children, e.g. depression, anxiety disorders)
- Pregnancy and childbirth (e.g. unwanted pregnancy, miscarriages, antepartum haemorrhage, poor prenatal care)

The physical injuries are rarely overlooked but the other symptoms are frequently overlooked.

A study by Stark et al.[5] defines a three-stage sequence to the battering syndrome:

- *Stage 1*: woman presents with injuries in the central anterior regions of the body (face, head and torso).
- *Stage 2*: multiple visits to clinics, often with vague complaints.
- *Stage 3*: development of psychological sequelae (alcohol, drug addiction, suicide attempts, depression).

Diagnosis

It is important to have a high index of suspicion and recognise and manage the problem to prevent further violence. If you suspect domestic violence - ASK! Talk to the woman alone:

- How are things at home?
- How are things with your spouse/children?
- Did anything unusual happen to bring about these injuries?
- Has there been any violence?
- You seem to be having a hard time.

It is vital to believe the woman's story. Women are most likely to seek help from their family doctor in preference to any other agency.[6] The doctor has to take the initiative because patients rarely complain about the violence.[3] They may present up to 30 times before they take action to end the violence.[3]

Barriers to communication about abuse[7]

- Concerns about confidentiality
- Perceptions about doctor
  - do not ask directly
  - have no time
  - are not interested
- Embarrassment
- Fear of involving police/courts
- Fear of shaming family
- Fear of partner hurting or killing them

Assessment

- Delineate the problem: pattern of violence; effect on the woman and her children; resources available to women; social/cultural environment.
- Examine and investigate presenting symptoms.
- Check for coexisting injuries (common target areas are breast, chest, abdomen and buttocks). Inspect the ears, teeth and jaw.
- Check the patient's general health status.
- Look for signs of alcohol or drug abuse.
- Keep accurate records and consider taking photographs.
- X-rays are helpful and may show old fractures.

VICTIMS

The victims come from all socioeconomic and cultural groups. As a rule they enter the relationship as normal, independent, competent women but gradually lose their coping ability and self-esteem and may become compliant victims.[1] This has been demonstrated by Hazel-wood and workers in their investigations of sexual sadists.[2] Unfortunately, many victims believe that somehow they deserve their punishment.

Many would like to leave home but the move is not so simple. Some do love their husbands and live in hope that the marriage will eventually work. They may feel that they cannot cope with living alone nor with the guilt and perceived failure of moving out.

PERPETRATORS

Perpetrators come from all walks of life and from all social and ethnic groups. They generally have inner drives to be strong, protective and powerful but can only achieve this at home through an inappropriate show of strength. However, they are basically insecure with poor self-esteem, poor communication skills, learned violence from family origins and an inability to express appropriate emotions, which tend to manifest as anger and violence.[1]

Although they usually control their violence outside the home, there is evidence that some perpetrators are guilty of violent behaviour in the community.

CYCLE OF VIOLENCE

A predictable pattern that is referred to as the 'cycle of violence' has been identified in many marriages. It is controlled by the perpetrator while the victim feels confused and helpless. The cycle repeats itself with a tendency for the violence to increase in severity (Figure 98.1).


Honeymoon phase
• denial
• togetherness
• clinging
• intimacy
Tension build-up phase
EXPLOSION
Violence/ abuse
• self-righteous anger
Remorse phase
• guilt
• denial
• justification
Wooing phase
• denial
• gifts and promises
• retribution

Figure 98.1 The cycle of domestic violence

Management

The key to successful management is initial recognition of the problem and establishment of empathic caring and support for the victim and family. Do not try to fit the victims into the disease model. It must be emphasised that the perpetrators (as in most criminal activity) do not readily change their behavioural pattern and thus there is minimal prospect of the violence decreasing unless there is a dramatic reason to change. As with an alcohol problem, the person has to admit that he has a problem before effective counselling can begin. A management strategy is presented in Table 98.1. The safety of women and children is always the prime working rule.

Table 98.1 Management strategy for domestic violence

Treat the physical injury and suspect domestic violence

↓

Establish the diagnosis

↓

Initiate crisis intervention

• organise admission to a refuge

• ensure informed consent for all actions

• consider notifying police

↓

Establish an empathic trusting relationship

↓

Build the victim's coping skills and self-esteem

↓

Make effective use of community resources

• support services

• women's support group

• domestic violence resource centre

• social services/police

• social workers

Useful strategies

Do believe her.

- Talk openly and explicitly about it.
- Express concern for her safety.
- Give information (i.e. about the course of action available to her, contacts for legal advice).
- Respect her right to make her own decisions.

Harmful strategies

Don't

- deny domestic violence
- minimise the importance of domestic violence
- blame the victim
- treat with tranquillisers
- refer to a psychiatrist
- refer to marriage guidance if the husband isn't interested, but refer to specialist counsellors
- set explicit criteria/rules (takes away her power yet again)

It is uncommon to get the co-operation of the perpetrator in the management process. If they do seek help they require counselling by a skilled and experienced practitioner, as treatment will be prolonged and complex.

As a general rule the most effective intervention in arresting the violence is to arrest the violent person.[3]

SEXUAL ASSAULT

Medical practitioners dealing with the difficult and distressing problem of alleged sexual assault should be trained in the subject and familiar with the laws applicable to sexual assault in their own state. Rape involves considerable violence and physical injury in 5 - 10% of cases,[8] in which the victims fear for their lives. Apart from the inevitable psychological consequences, the possibility of pregnancy or acquired STI should be considered. The inexperienced practitioner should refer the patient to the nearest available resource but continue a caring involvement as the patient's GP. Survivors of sexual assault should be allowed to accept or decline various treatment options offered by the practitioner.

Management of the victim

What you should do for the patients first is to offer and provide privacy, safety, confidentiality and emotional support. Believe them, listen to them and be non-judgmental.

Four important things to say initially to any victim:

- You are safe now.
- We are sorry this happened to you.
- It was not your fault.
- It's good that you are seeing me.

Initial advice to the victim

If victim reporting to police:

1. Notify the police at once.
2. Take along a witness to the alleged assault (if there was a witness).
3. Do not wash or tidy yourself or change your clothing.
4. Do not take any alcohol or drugs.
5. Don't drink or wash out your mouth if there was oral assault.
6. Take a change of warm clothing.

If not reporting to police or unsure, contact any of the following:

1. a friend or other responsible person
2. Lifeline or Lifelink or a similar service
3. a doctor
4. a counselling service

Obtaining information

1. Obtain consent from the patient to record and release information.
2. Take a careful history and copious relevant notes.
3. Keep a record, have a protocol.
4. Obtain a kit for examination.
5. Have someone present during the examination (especially in the case of male doctors examining women).
6. Air-dry swabs (media destroy spermatozoa).
7. Hand specimens to police immediately.

Examination

If possible the patient should be clothed when seen. Have the patient undress while standing on a white sheet to collect debris, and note any injuries as each item is removed. Each part of the body should be examined, under good illumination, and all injuries measured and recorded carefully on a diagram.

Injuries should be photographed professionally. Examine the body and genital area with a Wood's light to identify semen, which fluoresces. Perform a careful speculum examination. Palpate the scalp for hidden trauma. Collect appropriate swabs.

Making reports

Remember that as a doctor you are impartial. Never make inappropriate judgments to authorities (e.g. ‘This patient was raped’ or ‘incest was committed’). Rather say: ‘There is evidence (or no evidence) to support penetration of the vagina/anus’ or ‘There is evidence of trauma to ________________ .’

Postexamination

After the medical examination a discussion of medical problems should take place with the patient. This should be done in private and kept totally confidential. A management plan for physical injuries and emotional problems is discussed.

Consider the possibility of STI and possible referral. Consider also the possibility of pregnancy and the need for postcoital hormone tablets. Organise follow-up counselling and STI screening.

Management issues

- Take swabs and/or first-void specimen for testing gonococcus and chlamydia (PCR).
- Take blood for HIV, syphilis.
- Collect specimens - swab, aspirate of any fluid and keep for DNA analysis.
- Give prophylactic antibiotics (Table 98.2) - depends on type of assault and assailant.
- Emergency contraception.
- Review in 3 weeks - check tests.
- Screen for syphilis and HIV in about 3 months.
- Refer to Rape Crisis centre.

Table 98.2 STI prophylaxis (consider)[9]

- *Chlamydia* - 1 g azithromycin (o)
- Gonorrhoea - ceftriaxone 250 mg IM
- Hepatitis B - Hepatitis B vaccine (dose 1)
  - immune globulin IM (if high risk)
- Syphilis - benzathine penicillin 1.8 g IM
- *Trichomonas* - metronidazole 1 g (o)

Drug-assisted sexual assault

Consider this when patient has no memory of events and time or other suspicious circumstances. Urine or blood testing may be appropriate.

REFERENCES

1. Kerr A. Domestic violence. Treat it seriously. Aust Fam Physician, 1989; 18: 1362 - 9.
2. Hazelwood R, Warren J, Dietz P. Compliant victims of the sexual sadist. Aust Fam Physician, 1993; 22: 474 - 9.
3. Knowlden S, Helman T. How to treat domestic violence. Aust Dr Weekly, 1989; 21 April: I - VIII.
4. Hegarty K, Hindmarsh E, Gillies M. Domestic violence in Australia. Med J Aust, 2000; 175: 363 - 7.
5. Stark E. et al. *Wife Abusing in the Medical Setting - an Introduction to Health Personnel.* Rockville, MA: National Clearing House for Domestic Violence, 1981: 7.
6. Western Australian Domestic Violence Task Force. *Break the Silence*. Report to the Western Australian Government. Perth: Government Printer, 1986: 26 - 35, 162 - 3.

7. Rodriguez MA et al. The factors associated with disclosure of intimate partner abuse to clinicians. J Fam Pract, 2001; 50(4): 338 - 44.
8. Tierney LM et al. *Current Medical Diagnosis and Treatment* (45th edn). New York: The McGraw-Hill Companies, 2006: 758 - 9.
9. Mein JK, Palmer CM, Shand M et al. Management of acute adult sexual assault. MJA, 2003; 178: 226 - 30.

CHAPTER 99. BASIC ANTENATAL CARE

An examiner, no lover of females, thrust a femur into her hand.

‘How many of these have you got?’ he demanded.

‘Five.’

‘How do you come to that conclusion?’ he asked contemptuously.

‘I have two of my own, the one in my hand, and the two of my unborn child.’

ANONYMOUS ANECDOTE

BASIC ANTENATAL CARE

Pregnancy and childbirth are very important and emotional events in the lives of women and their families. Their care during and after pregnancy is one of the most satisfying aspects of the work of the family doctor, who generally chooses breadth of knowledge rather than depth of knowledge. The changing trend towards specialisation has meant a change of role for the city practitioner and now shared obstetric care is a commonly practised routine. The quality of care that can be given in family practice is often superior to that offered in the hospital antenatal clinic, partly because of the continuing personal care offered by the family doctor.[1]

Antenatal care presents preventive medicine opportunities par excellence and is the ideal time to develop an optimal therapeutic relationship with the expectant mother. The opportunities for anticipatory guidance should be seized and education about the multitude of possible sensations during pregnancy (such as heartburn, backache, leg cramps, various fears and anxieties) should be addressed. In other words, an optimal communication system should be established between the expectant couple and the health care system.

Early diagnosis of high-risk pregnancy is important but of little value unless followed by normal attendance for antenatal care.

The information presented here is a basis for the shared care strategy where family doctors share basic antenatal care with consultants and have a ready referral strategy for high-risk pregnancies.

The basic aim of antenatal care is to assess the risk of harm to mother and baby and apply the appropriate level of surveillance to minimise or eradicate harmful effects.

PRECONCEPTUAL CARE

Preconceptual care is to be commended to the woman contemplating pregnancy and her family doctor is well placed to provide general health care and screening as well as genetic counselling.

General advice should include optimal nutrition and diet, weight control, regular exercise and discouragement of smoking, alcohol and drugs. Listeria infection is a problem if contracted, with foetal mortality being 30 - 50%. Protection is afforded by good personal and food hygiene. For those who ask, advise avoidance of unpasteurised dairy products, soft cheeses, cold meats and raw seafoods, chilled ready to eat foods and takeaway foods.

Folic acid 0.5 mg tablets is now generally recommended to commence 3 months prior to conception, continuing to 12 weeks post conception. Examination should include blood pressure, cardiac status, urinalysis and cervical smear.

Rubella serology should be estimated and, if required, immunisation 3 months prior to conception should be initiated. Test for seroconversion 3 months later. Vaccination should be avoided in early pregnancy.

Ask about a history of varicella and, if necessary, consider serology and vaccination.

Genetic counselling based on past obstetric or family history, advanced maternal age or other factors should be considered. This applies especially to Down syndrome, neural tube defect, congenital heart disease, cystic fibrosis and fragile X syndrome.

Checkpoint summary: advice to patients

- Stop smoking.
- Stop alcohol and other social drugs.
- Reduce or stop caffeine intake.
- Review current medications with your GP.
- Follow a healthy diet.
- Avoid being overweight.
- Take folic acid for 3 months before conception.
- Have a good exercise routine
- Ensure rubella immunity.
- Have a regular breast check and Pap smear
- Eat freshly cooked and prepared food.
- Consider genetic and family history.
- Consider health insurance cover.

THE INITIAL VISIT

It is important to book the patient into hospital at the first visit and into the antenatal outpatient department if appropriate. It is mandatory to make an accurate estimation of the expected due date - the use of ultrasonography can help.

History checkpoints[2]

- Confirm the pregnancy by the menstrual history and by urine or serum human chorionic gonadotrophin (HCG) if necessary.
- Previous obstetric history:
  - Gestation, length of labour, mode of delivery, birthweight of each baby.
  - Consider previous problems: foetal or neonatal abnormalities or deaths; preterm or growth-retarded infants.
  - Abortions: determine if there has been any termination of pregnancies or first or second trimester spontaneous abortions.
- Medical history:
  - Check for past evidence of diabetes, tuberculosis, anaemia, rubella, rheumatic fever, heart or renal disease, jaundice, depression, transfusions and rhesus status.
- Family history:
  - Features to consider are multiple pregnancies, hypertension and diabetes.
  - If any of these pertain to first-degree relatives, consider a glucose screening or tolerance test.
- Psychosocial history:
  - This is very important and includes an assessment of the emotional attitude.
- Drug history:
  - Includes intake of nicotine, alcohol, aspirin, illicit drugs, OTC drugs and prescribed drugs.
- Important points to consider:
  - Establish date of confinement (see obstetric calendar in Figure 99.1).



The calculation is made from the first day of the last menstrual period

January	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	**January**
October	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	1	2	3	4	5	6	7	***November***
February	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28				**February**
November	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	1	2	3	4	5				***December***
March	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	**March**
December	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	1	2	3	4	5	***January***
April	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30		**April**
January	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	1	2	3	4		***February***
May	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	**May**
February	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	1	2	3	4	5	6	7	***March***
June	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30		**June**
March	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	1	2	3	4	5	6		***April***
July	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	**July**
April	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	1	2	3	4	5	6	7	***May***
August	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	**August**
May	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	1	2	3	4	5	6	7	***June***
September	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30		**September**
June	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	1	2	3	4	5	6	7		***July***
October	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	**October**
July	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	1	2	3	4	5	6	7	***August***
November	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30		**November**
August	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	1	2	3	4	5	6		***September***
December	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	**December**
September	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	1	2	3	4	5	6	7	***October***

Or: (approximately) subtract 3 from months and add 7 to days e.g. 19/8/89
+ 7 − 3
26/5/90 Or: Naegele's rule—add 7 days, 9 months

Figure 99.1 Obstetric calendar to determine expected due date

- If maternal age > 37 years, consider feasibility of amniocentesis or chorionic villus sampling or other relevant tests (Down syndrome).
- Consider unusual causes for severe nausea and vomiting (e.g. hydatidiform mole, cerebral tumour).
- Investigate possible exposure to rubella.
- If vaginal bleeding: if Rh negative, send blood sample for Rh antibodies - if absent, give one ampoule anti-D gammaglobulin within 72 hours of first bleed.

Physical examination

During the initial examination assess the patient's general physical and mental status. Examine the following:

- general fitness, colour (? anaemia)
- basic parameters: height, weight, blood pressure, pulse, urinalysis (protein and glucose)
- head and neck: teeth, gums, thyroid
- chest: including breasts/nipples
- abdomen: palpate for uterine size and listen to foetal heart (if indicated)
  Perform the four classic techniques of palpation (applies to later visits):
  1. Fundal palpation
  2. Lateral abdominal palpation
  3. Pawlik palpation
  4. Deep pelvic palpation
- legs: note oedema or varicose veins

Speculum examination: perform a Pap smear and swab (if indicated by abnormal vaginal discharge).

Pelvic examination (optional): confirm uterus size and period of gestation by bimanual palpation.

ANTENATAL SCREENING

The recommendations for antenatal screening of asymptomatic women are the source of de-bate with EBM calling into question traditional tests. Table 99.1 reflects the recommendations of the RANZ College of Obstetricians and Gynaecologists, who do not recommend the screening of all pregnant women for gestational diabetes, or for asymptomatic bacteriuria.[3, 4] Debate continues about screening for syphilis, group B Streptococcus in late pregnancy, thyroid function, hepatitis B and C and ultrasound.

Table 99.1 Routine antenatal screening[*]

Recommended[3]	Consider
First visit:	
• full blood count	• HCV serology (if significant risk)
• blood group and antibody screen	• Haemoglobin electrophoresis (if indicated)
• rubella antibody status	
• cervical cytology	
• HBV serology	*Subsequent visits:*
• HIV serology (after counselling)	• 18-week morphology scan
• syphilis serology	
• urine culture sensitivity	
Discuss:	
• ultrasound 18 - 20 weeks	
Subsequent:	
• oral glucose challenge test 28 weeks: if abnormal →OGTT	
• rubella antibodies 30 and 36 weeks (Rh-negative mother)	
• FBC - 36 weeks	

[*] For prenatal diagnosis of genetic abnormalities: Discuss availability of maternal serum screening and nuchal transparency for Down syndrome.

VISITS DURING PREGNANCY

- The box below is a summary of standard antenatal practice.
- The average number of visits is twelve but the need for this number is being questioned, with some authorities recommending as few as six visits.

A common routine schedule

- Initial in first trimester: 8 - 10 weeks
- Up to 28 weeks: every 4 - 6 weeks
- Up to 36 weeks: every 2 weeks
- 36 weeks - delivery: weekly

A systematic review of seven RCTs found no difference in the detection of pre-eclampsia, urinary tract infection, low birthweight or maternal mortality when a schedule of reduced antenatal visits was compared with the traditional routine.[5] However, a sensible approach would be to plan visits according to need and circumstances in a flexible way.

For each visit record:

- weight gain
- blood pressure
- urinalysis (protein and sugar) - see Table 99.2

Table 99.2 Causes of proteinuria in pregnancy

Urinary tract infection
Contamination from vaginal discharge

Pre-eclampsia toxaemia
Underlying chronic renal disease

- uterine size/fundal height
- foetal heart (usually audible with stethoscope at 25 weeks and definitely by 28 weeks)
- foetal movements (if present)
- presentation and position of foetus (third trimester)
- presence of any oedema

Record day of first foetal movements (ask patient to write down the dates)

- primigravida: 17 - 20 weeks
- multigravida: 16 - 18 weeks

Fundal height

The relative heights of the uterus fundus are shown in Figure 99.2. The uterus is a pelvic organ until the twelfth week of pregnancy. After this time it can be palpated abdominally. At about 20 - 22 weeks it has reached the level of the umbilicus and reaches the xiphisternum between 36 and 40 weeks. Palpation of the fundal height is affected by obesity and tenseness of the abdominal wall.



Figure 99.2 Fundal height in normal pregnancy (in weeks); the height of the fundus is a guide to the period of gestation. Nulliparas experience lightening at about 36 weeks when the fundal height usually reverts to the 34 week level

MANAGEMENT OF SPECIFIC ISSUES

Nutrition advice

A healthy diet is very important and should contain at least the following daily allowances:

1. Eat most:

    - fruit and vegetables (at least 4 serves)
    - cereals and bread (4 - 6 serves)

2. Eat moderately:
    - dairy products - 3 cups (600 mL) of milk or equivalent in yoghurt or cheese
    - lean meat, poultry or fish - 1 or 2 serves (at least 2 serves of red meat per week)

3. Eat least:
    - sugar and refined carbohydrates (e.g. sweets, cakes, biscuits, soft drinks)
    - polyunsaturated margarine, butter, oil and cream

Bran with cereal helps prevent constipation of pregnancy.

If the ideal diet is followed, iron, vitamin and calcium supplements should not be necessary. Do not diet to lose weight. It is usual to gain about 12 kg during pregnancy.

Smoking, alcohol and other drugs[6]

Encourage patients to avoid all street drugs, alcohol, tobacco and caffeine (ideally). If they find this impossible, encourage the following daily limitations:

- 1 standard drink of alcohol
- 1 cup of coffee or 2 cups of tea

Other household members should also stop smoking as passive smoking may be harmful to mother and child.

There is convincing evidence that promotion of smoking cessation programs during pregnancy is effective, with improved outcomes, including reduction in preterm birth rates and low birthweight rate.[7] The fetal alcohol syndrome is a leading cause of mental retardation so it is best to abstain.

Breastfeeding

Mothers-to-be should be encouraged to breastfeed. Give advice and relevant literature. They can be directed to a local nursing mothers' group for support and guidance if necessary.

Antenatal classes

Referral to therapists conducting such classes can provide advice and supervision on antenatal exercises, back care, posture, relaxation skills, pain relief in labour, general exercises and swimming. Enrolment with the partner is recommended.

Normal activities

Mothers should be reassured that pregnancy is a normal event in the life cycle and that normal activities should be continued. Housework and other activities should be performed to just short of getting tired. The importance of getting sufficient rest and sleep should be emphasised.

Sex in pregnancy[8]

Coitus should be encouraged during pregnancy but with appropriate care, especially in the 4 weeks before delivery. Restriction would only seem necessary if there has been an adverse obstetric history and there are major complications in the current pregnancy.

The couple should be encouraged to be loving to each other and communicate their feelings freely, as the need for affection and physical contact is important. Coital techniques can be modified as the pregnancy progresses - posterior entry and the female superior position are quite suitable.

Travel

Pregnant women should avoid standing in trains. They should avoid international air travel after 28 weeks and travel after 36 weeks is usually not permitted. Patients should be counselled to wear a seat belt during car travel.

Psychosocial and emotional stress

Antenatal visits provide an ideal opportunity to become acquainted with the ‘real’ person and explore issues that help the patient. Provide whole person understanding with appropriate help and reassurance where necessary. Areas to be explored include support systems, attitudes of patient and partner to the pregnancy, sexuality, expectation of labour and delivery, financial issues, and attitudes of parents and in-laws.

Weight gain in pregnancy

Although a standard weight gain is given as 12 kg over 40 weeks of pregnancy, it is common for some women in Australia to gain up to 20 kg without adverse effects.[2]

Normal weight gain is minimal in the first 20 weeks, resulting in a 3 kg weight gain in the first half of pregnancy. From 20 weeks onwards there is an average weight gain of 0.5 kg per week. From 36 weeks the weight gain usually levels off.[2]

Foetal movement chart

If daily foetal movements exceed 10 and the regular pattern has not changed significantly, then usually the foetus is at no risk. However, if the movements drop to fewer than 10 per day, the patient should be referred to hospital for foetal monitoring.

Possible exposure to rubella

When contact with a possible case of rubella occurs during pregnancy it is essential to establish the immune status of the patient. If she is already immune no further action is necessary. If her immune status is unknown, perform a rubella IgG titre and IgM and repeat the IgG and IgM titres in 2 - 3 weeks.

Vaginal bleeding in early pregnancy[9]

This is a common problem in the first trimester in particular. At least 10% of normal pregnancies will have an episode and about 15% of recognised pregnancies will miscarry. If the bleeding is light to moderate and the pain mild or absent the question is ‘Can a viable pregnancy continue or is there an ectopic pregnancy or a threatened miscarriage?’

- < 6 weeks: Do serial quantitative HCG levels, which should double every 2 days (ultrasound usually unhelpful). If rise is too slow it means a non-viable pregnancy (? in tube or uterus). If HCG > 1500 IU/L transvaginal ultrasound is used to show gestational sac.
- 6 - 8 weeks: Ultrasound will define an intrauterine pregnancy by excluding an ectopic.
- > 8 weeks: Normal ultrasound reassuring since miscarriage rate is only 3%.

Note: rest is inappropriate for threatened miscarriage

A small bleed between 18 - 24 weeks indicates cervical ‘weakness’ and warrants a speculum or vaginal examination plus foetal assessment.

Threatened miscarriage

If a threatened miscarriage occurs, check the blood group and test for rhesus antibodies in maternal serum. If the mother is Rh-negative and no antibodies are detected, give one ampoule of anti-D gammaglobulin intramuscularly. Assess her pelvis to rule out an ectopic pregnancy and, if indicated, perform pelvic scanning to confirm viability of the foetus or the presence of an extrauterine gestation.

Pregnancy sickness[10]

- Nausea and vomiting occur in more than 50% of women
- Almost always disappears by the end of first trimester
- Mild cases can be dealt with by explanation and reassurance; it is preferable to avoid drug therapy if possible
- Simple measures:
    - small frequent meals
    - a fizzy soft drink, especially ginger drinks, may help
    - avoid stimuli such as cooking smells
    - take care with teeth cleaning
    - avoid oral iron
- Medication (for severe cases):
    - pyridoxine 50 - 100 mg bd
    - if still ineffective add metoclopramide 10 mg tds

Hyperemesis gravidarum

This is severe vomiting in pregnancy, which may result in severe fluid and electrolyte depletion.

Association

- Normal complication
- Hydatiform mole
- Multiple pregnancy
- Urinary infection

Management

- Test urine - MCU (micro-culture of urine) ketones: if the admit to hospital
- Ultrasound examination
- Bed rest
- Ultrasound examination
- Electrolytes, area, LFTs
- Nil orally
- Fluid & electrolyte replacement
- Pyridoxine 50 - 100 mg daily IV/oral
- Metoclopramide 10 mg IV ↑ 10 mg tds (if necessary)
- Return to oral intake

Heartburn[6]

Gastro-oesophageal reflux is a major source of discomfort to women in the latter half of pregnancy. Non-pharmacological treatment such as frequent small meals, avoidance of bending over and elevation of the head of the bed are the mainstays of treatment. Smoking, alcohol and caffeine (coffee, chocolate, tea) intake should be avoided. Regular use of antacids is effective (e.g. alginate/antacid liquid - Gaviscon, Mylanta Plus - 10 - 20 mL) before meals and at bedtime. H_2-receptor antagonists may be necessary.

Cramps

Pregnant women are more prone to cramp. If it develops they should be advised simply to place a pillow at the foot of the bed so that plantar flexion of the feet is avoided during sleep. Prolonged plantar flexion is the basis of the cramps. Tonic water prior to sleeping is helpful. There is no evidence that calcium supplements help cramps during pregnancy.[11]

Varicose veins

These can be troublesome as well as embarrassing. Wearing special supportive pantyhose (not elastic bandages) is the most comfortable and practical way to cope, in addition to adequate rest.

Haemorrhoids

Haemorrhoids in the later stages of pregnancy can be very troublesome. Emphasising the importance of a high-fibre diet to ensure regular bowel habit is the best management. Painful haemorrhoids may be eased by the application of packs soaked in warm saline or perhaps haemorrhoidal ointments containing local anaesthetic.

Dental hygiene

Dental problems can worsen during pregnancy so special care of teeth and gums, including a visit to the dentist, is appropriate. Continuation of cleaning with a softer brush is recommended.

Back pain

Back pain, especially low back pain, is common during pregnancy and special back care advice can help women cope with this problem, which can become debilitating. Advice with lifting, sitting, and resting, using a firm mattress and avoiding high-heeled shoes will help.

Physical therapy administered by a skilled therapist can be extremely effective for pregnant patients but certain safety rules should be followed:

- First trimester: use normal physical therapy and advise exercises.
- Second trimester: use supine side lying rotation and sitting techniques only; advise exercises.
- Third trimester: avoid physical therapy (if possible); encourage exercises.

Guidelines for treatment

- Keep mobilisation and manipulation to a minimum.
- Use stretching and mobilisation in preference to manipulation.
- Safeguard the SIJs in the last trimester.
- Encourage active exercises as much as possible.
- Avoid medications wherever possible.
- Give trigger point injections (5 - 8 mL 1% lignocaine) around the SIJs if necessary.

Exercise guidelines

Advise the patient that walking is an excellent exercise. For additional exercise activity:

- exercise at a mild to moderate level only
- avoid overheating and dehydration
- allow for a long warm-up before exercise and a long cool-down
- choose low-impact or water exercise
- stop if there are adverse symptoms (e.g. any pain, bleeding, faintness, undue distress)
- avoid scuba diving and sky diving

Carpal tunnel syndrome

Splinting of the hand and forearm at night might be beneficial. If desperate, an injection of corticosteroid into the carpal tunnel can be very effective (check drug category for risk relative to dates). Sometimes operative division of the volar carpal ligament is necessary. Most problems subside following delivery.

Hypotension

This is due to increased peripheral circulation and vensus pooling. If bleeding eliminated, advise to avoid standing suddenly and hot baths. It may cause syncope. Fainting may occur when the women lies on her back in the latter half of pregnancy (supine hypotension).

Pruritus

Generalised itching (pruritis gravidarum) is usually associated with cholestasis due to oestrogen sensitivity in the third trimester. Order LFTs and, if not serious, reassure and prescribe a soothing skin preparation, eg. aqueous cream ± glycerol.

Obesity

Obesity is associated with increased obstetric morbidity, including difficult labour and potential anaesthetic risks. Encourage weight loss diet with the aid of a dietitan.

Breathlessness of Pregnancy[14]

Consider physiological breathless of pregnancy in a woman with unexplained dyspnoea. It starts in the second trimester, is constant and aggravated by exercise and emotional stress. No special treatment is needed nor helpful. The breathlessness usually settles 6 - 8 weeks after delivery.

Mineral supplements in pregnancy

Iron

Iron is not routinely recommended for pregnant women who are healthy, following an optimal diet and have a normal blood test. Those at risk (e.g. with poor nutrition, Vegan diet) will require supplementation.

Folic acid

Folic acid is advised for all women con-templating pregnancy, starting 3 months prior to conception and continuing until 12 weeks after conception. Dose: 0.5 mg (o) daily.[12] In those at risk (e.g. previous neural tube defect and lustong of epilepry) the dose is 5 mg per day.[13]

Advice on when to seek medical help

- If contractions, unusual pain or bleeding occur before term
- If the baby is less active than usual
- If the membranes rupture (with fluid loss)
- The onset of regular contractions 5 - 10 minutes apart

REFERENCES

1. Barker JH. General Practice Medicine. Edinburgh: Churchill Livingstone, 1984: 76 - 89.
2. Fung P, Morrison J. Obstetric share-care. Aust Fam Physician, 1989; 18: 479 - 84.
3. Oats JJN. Routine antenatal screening: a need to evaluate Australian practice (editorial). Med J Aust, 2000; 172: 311 - 2.
4. Hunt JM, Lumley J. Are recommendations about routine antenatal care in Australia consistent and evidence-based? Med J Aust, 2002; 176: 255 - 61.
5. Carroli G, Villar J, Piaggio G et al. WHO systematic review of randomised controlled trials of routine antenatal care. Lancet, 2001; 357: 1565 - 70.
6. Smibert J. The principles of antenatal care. Aust Fam Physician, 1978; 9: 1087 - 94.
7. Lumley J, Oliver S, Waters E. Interventions for promoting smoking cessation during pregnancy (Cochrane review). In: the Cochrane Library, Issue 1, Oxford: Update Software, 2001.
8. Beischer NA, Mackay EV. Obstetrics and the Newborn. Sydney: Saunders, 1986.
9. Peat B. Antenatal care: common issues facing GPs in shared care. Med Today, 2001; June: 81 - 5.

10. Humphrey M. Common conditions in an otherwise normal pregnancy. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 116 - 20.
11. Hammer I et al. Calcium treatment of leg cramps in pregnancy. Acta Obstet Gynaecol Scand, 1981; 60: 345 - 7.
12. Goh J, Erian M. An update on obstetric practice. Mod Med Aust, 1996; 8: 20 - 5.
13. MRC Vitamin Study Research Group. Prevention of neural tube defects. Lancet, 1991; 338: 131 - 7.
14. Burdon J. Respiratong medicine. CHECK Program, Unit 395. Melbourne RACGP, 2005: 17 - 18
15. Birth defects Part A; 76: 73 - 77.

CHAPTER 103. MEN'S HEALTH: AN OVERVIEW

Call it the ‘M-factor’, call it maleness, call it what you like - but from infancy onwards, in every age group, males are more likely to die than females.

DR ANDREW PATTISON 2001[1,2]

MEN'S HEALTH: AN OVERVIEW

In recent years increasing attention has been focused on men's health, mainly because it became evident that the average male's lifestyle was slowly killing him. As doctors we are beginning to understand that a great proportion of male ill-health is related to behavioural and social factors. Being male has been described by Dr Ian Ring of the Queensland Health Department ‘as a health hazard’.[3]

An important statistic is the constant discrepancy in average life expectancy (ALE) between the sexes. At present in Australia, the ALE is 76 years for males compared with 81 years for women.[1,4]

Since the beginning of the 20th century and even further back this discrepancy has continued. In 1900 the ALE for males was 55.2 years and 58.8 for females.[4] This increased to a difference of 6 years for most of the last century. However, the significant increase in ALE for both sexes has been encouraging.

Men have a significantly greater incidence of medical conditions, such as cardiovascular disease, accidental death, suicide, obesity, alcoholism, HIV and hypertension. The following comparative statistics for Australian society highlight this difference.

MEN'S HEALTH AT A GLANCE[1,2]

- Average life expectancy: males 76 years, females 81 years, and the discrepancy begins in infancy or earlier.
- Up to 14 years, boys are at least *twice* as likely to die from accidental injury (e.g. motor vehicle accidents [MVAs] and drowning).
- In the 15 - 24 year age group, males are *three* times more likely to die in MVAs and *four* times more likely to suicide. The overall death rate is 3.65 times higher than for females.
- In the 25 - 65 year age group, males are *four* times more likely to die from coronary artery disease, *three* times more likely to die in MVAs, *four* times more likely to suicide, *four* times more likely to die in other accidents, and *twice* as likely to die from cancer. The overall death rate is two times that of females.
- The figures are worse in the poorer socioeconomic classes. Low-income males are nearly *three* times more likely to state that their overall health is poor compared to men with higher income.



Figure 103.1 Reprinted with permission of Ron Tandberg

- At least four out of five heroin overdose deaths occur in males.
- Aboriginal males' life expectancy is 17 years less than that of non-Aboriginal males. In the 35 - 45 year age group, the death rate is *eleven* times that of non-Aboriginals.
- Workplace deaths - 93% occur in males (who constitute 56% of the workforce).
- 46% of Australian marriages end in divorce. The majority of these are initiated by women.
- 90% of those convicted for acts of violence are males: 80% of the victims are males.
- In Australian schools, 90% of children with documented behavioural problems are males. (This list has been reproduced with the permission of Andrew Pattison.)

These statistics are very revealing and reflect attitudes to lifestyle. Men smoke more, drink more and indulge in greater risk-taking behaviour in general. Even two-thirds of pedestrians killed on the streets are male.[3] The main causes of death by age group for Australian males is summarised in Table 103.1.[5]

Table 103.1 Main causes of death by age group for Australian males

Years	No. 1	No. 2	No. 3
1 - 14	Non-traffic accidents	Traffic accidents	Cancer
15 - 24	Traffic accidents	Suicide	Non-traffic accidents
25 - 44	Suicide	Cancer	Non-traffic accidents
45 - 54	Cancer	Circulatory diseases	Suicide
55 - 64	Cancer	Circulatory diseases	Suicide
65 - 74	Circulatory diseases	Cancer	Respiratory diseases
75+	Circulatory diseases	Cancer	Respiratory diseases

The question has been asked ‘Is the Y chromosome some kind of hidden killer or are men just too reckless - or dumb - to look after themselves?’ [3]

A study across 75 species, including spiders, reptiles, birds, fish and mammals found males had shorter life spans than females in virtually every instance. Ian Ring postulates that 'when you add tobacco, alcohol and social pressure to succeed onto what is already a biological inferiority and a natural aggression to protect, you get a deadly mix'.[3]

PROSTATIC DISEASE

Ageing men find disorders of the prostate almost inevitable, as they become aware of lower urinary tract symptoms (LUTS). The usual cause of benign prostatic hyperplasia is now being managed better with α-blocking agents, such as tamsulosin, which is delaying the almost inevitable resection of the prostate - a procedure that provokes considerable anxiety in men. Cancer of the prostate is the second most common cause of death from cancer in men and the third overall including both sexes, yet its management remains controversial and at times confusing, especially the screening issues. Although it almost matches death from breast cancer in women, there is a huge discrepancy in health funding between men's and women's health. More attention needs to be given to the prevention and management of prostate and testicular cancer.

ERECTILE DYSFUNCTION AND SEXUALITY

The new era of a 'liberalised' and 'sensationalist' press has placed this issue in the limelight. There is a pressure and focus on the sexual performance of men as never before and this carries the issue of performance anxiety and the various manifestations of sexual unfulfilment, which some claim is a significant factor in the disturbingly high suicide rate of young men.

Then we have the issue of young men discovering their sexuality and working through the confusion and anxiety caused by the realisation of their gayness. These men need understanding and support by a profession and society that is emerging from a culture of ignorance.

The new treatments for erectile dysfunction have brought mixed blessings but they have certainly helped many couples in their relationship and helped the self-esteem of many men. However, we are faced with the juggling act of affordability, government subsidisation, adverse effects and good outcomes. The advent of the phosphodiesterase inhibitors has brought a new and improved dimension to this management.

SEX-LINKED INHERITED DISORDERS

Males bear the burden of X-linked recessive gene disorders which always manifest as there is no normal gene on the additional X chromosome (as there is in the female) to counteract the action of the abnormal gene.

Occasionally a gene can be carried on an autosome but manifests only in one sex. An example is frontal baldness, manifesting as an autosomal dominant disorder in males but as a recessive disorder in females.

Examples of X-linked disorders significantly affecting males include:

- haemophilia A and B
- glucose-6-phosphate dehydrogenase deficiency
- Duchenne muscular dystrophy
- retinitis pigmentosa
- Hunter's syndrome
- ocular albinism

SUMMARY

There is an increasing emphasis and interest in men's health. The recently opened men's health clinics have been very successful and will heighten the interest in the health area. However, to make progress we need to redefine our concept of masculinity. Andrew Pattison has summarised the important tasks ahead in a powerful way:[2]

- men and boys need to be more aware of their maleness
- self-esteem is important at every stage of our lives
- men and boys should try and be in touch with their feelings
- men and boys need to be more in tune with their bodies
- men and boys should look closely at how they communicate with others
- health is our personal responsibility

REFERENCES

1. Pattison A. *The M factor* (2nd edn). Sydney: Simon & Schuster, 2001.
2. Pattison A. Men and their health: the M factor. Curr Ther, 2001; August: 9 - 13.
3. Smith R. The gender trap. Time (South Pacific edn), 1994; 12 December: 56 - 61.
4. Australian Institute of Health and Welfare. Australia's Health 2004. < http://www.aihw.gov.au >
5. Australian Bureau of Statistics. *Causes of Death.* Canberra: ABS, 2006.

CHAPTER 104. SCROTAL PAIN

Acute scrotal pain in infancy and adolescence should be regarded as torsion of the testis until proved otherwise.

SCROTAL PAIN

Scrotal pain in males can occur at all age groups but the child or adolescent with acute scrotal pain often poses a diagnostic challenge. Serious problems include testicular torsion, strangulation of an inguinoscrotal hernia, a testicular tumour and a haematocele, all of which require surgical intervention.

KEY FACTS AND CHECKPOINTS

- Torsion of the testis is the most common cause of acute scrotal pain in infancy and childhood.
- Torsion is also a feature of young men less than 25 years.
- Testicular pain can be referred to the abdomen.
- Torsion of the testis should form part of the differential diagnosis in a boy or young man who is vomiting and has intense pain in the lower abdominal quadrant inguinal region.
- The clinical picture of epididymo-orchitis can mimic torsion of the testis so closely that, in most children, the diagnosis should be made only at surgical exploration.[1]
- An abnormality predisposing to torsion of the testis is usually present bilaterally; the opposite testis should also be fixed to prevent torsion.
- Torsion must be corrected within 4 - 6 hours to prevent gangrene of the testis.
- Suspect self-correcting testicular torsion in repeated episodes of severe spontaneously resolving pain. Refer for orchidopexy.
- Suspect abscess formation if epididymo-orchitis does not settle with a reasonable course of antibiotics. Surgical drainage may be necessary.
- A varicocele can cause testicular discomfort - examine the patient in the standing position.

THE CLINICAL APPROACH

History

It is important to determine whether there were any pre-existing predisposing factors or history of trauma.

Key questions

- Have you noticed any burning of urine or penile discharge?
- Have you had an injury to your scrotal region such as being struck by a baseball, cricket ball or falling astride something?
- Have you travelled overseas recently?
- Have you been aware of a lump in your testicle or groin?
- Have you had an illness lately and have you noticed swelling of the glands in your neck or near your ear?
- Do you have back pain or have you injured your back?

Examination

Both sides of the scrotum must be examined and contrasted. Inguinal and femoral hernial orifices, the spermatic cord, testis and epididymis must be checked on both sides. The patient should be examined standing and supine. The scrotum and its contents are examined systematically starting with the skin, which may include sebaceous cysts or rarely may exhibit thickening, sinuses or ulcers with inflammatory disorders such as filariasis and tuberculosis. A painful testis should be elevated gently to determine if the pain improves.

Investigations

Investigations that may help diagnose the painful testis in particular include:

- blood cell count
- urine analysis: microscopy and culture
- *Chlamydia* antigen detection tests
- ultrasound
- technetium-99m scan

ACUTE SCROTAL PAIN IN CHILDREN AND ADOLESCENTS

This problem is more likely to be encountered in the adolescent. A list of causes is presented in Table 104.1. Infants, however, can also have torsion of a testis or a testicular appendage, such as the hydatid of Morgagni.

Table 104.1 Causes of scrotal pain or swelling

Torsion of the testis
Torsion of a testicular appendage
Epididymo-orchitis
Mumps orchitis
Acute hydrocele
Idiopathic scrotal oedema
Haematoma/haematocele
Neoplasm
Henoch-Schönlein purpura
Strangulated inguinoscrotal hernia
Referred pain
Scrotal skin conditions
Varicocele

Clinical problem

A 15-year-old teenager presents with relatively acute onset of pain in his lower right abdomen and scrotum. He has vomited several times. On examination the right testicle is tender, red and swollen.

Discussion

The two main differential diagnoses are acute epididymo-orchitis (which requires little more than conservative treatment) or torsion of the testis, which demands emergency surgical intervention (Figure 104.1). Less commonly the problem would be a haematoma or an acute hydrocele mimicking testicular torsion. This patient, however, must be regarded as having torsion of the testis. Early operation with torsion is imperative because, if the testis is deprived of its blood supply for more than a few hours, infarction is inevitable and excision becomes necessary. Excluding mumps, no youth under the age of 18 years should be diagnosed as suffering from acute epididymo-orchitis until the testis has been exposed at operation and torsion excluded.



Figure 104.1 Illustration of torsion of the testis and an appendage: the 'black' hydatid of Morgagni is the one most likely to undergo torsion

Torsion of the testis v epididymo-orchitis

With torsion of the testicle there is pain of sudden onset, described as a severe, aching, sickening pain in the groin that may be accompanied by nausea and vomiting. With epididymo-orchitis the attack usually begins with malaise and fever and is often associated with a urinary infection. The testicle soon becomes swollen and acutely tender; however, elevation of the scrotum usually relieves pain in this condition while tending to increase it with a torsion. A comparison of the clinical presentations is given in Table 104.2.

Table 104.2 Clinical presentations of torsion of testis and acute epididymo-orchitis

	Torsion of testis	**Epididymo-orchitis**
Typical age	Early teens, average range	Young adults
	5 - 15 years	Elderly
Onset	Usually sudden but can be gradual	Gradual
Severity of pain	Very severe	Moderate
Associated symptoms	Vomiting	Fever
	Groin pain	± Dysuria
	Possibly abdominal pain	
Examination of scrotum	Very tender and red Testis high and transverse Scrotal oedema	Swollen, tender and red; can be tender on rectal examination
	Possibly an acute hydrocele	Possibly an acute hydrocele
Investigations	Technetium-99m scan (if available, time permits and diagnosis doubtful)	Leucocytosis Possibly pyobacteria of urine

Radiology as a diagnostic aid

Ultrasound, particularly colour Doppler, is useful in distinguishing a cystic scrotal lump (such as a hydrocele) from a solid tumour. Its use to distinguish between a torsion and epididymo-orchitis is controversial as it cannot reliably detect changes that are diagnostic of early torsion. Since the investigation can involve unnecessary delay in treatment it is generally not recommended. A technetium-99m scan can differentiate between the two conditions: in torsion the testis is avascular while it is hyperemic in epididymo-orchitis.

Time factor in surgical intervention

The optimal time to operate for torsion of the testis is within 4 - 6 hours of the onset of pain. About 85% of torsive testes are salvageable within 6 hours but by 10 hours the salvage rate has dropped to 20%.[2]

At surgery the testicle is untwisted and if viable an orchidopexy is performed. A gangrenous testicle is removed (Figure 104.2). The opposite testis should be fixed by orchidopexy.



Figure 104.2 Torsion of the testis resulting in gangrene after 12 hours from onset of pains. The testis was excised and the other normal testis 'anchored'

TORSION OF A TESTICULAR APPENDAGE

Vestigial remnants to the testis or the epididymis are present in 90% of the male population.[1] Torsion of the testicular appendage, the pedunculated hydatid of Morgagni, has a similar presentation to that of torsion of the testis but is less severe (see Figure 104.1). It can be diagnosed by the appearance of a dark blue nodule at the upper pole of the testis (provided that it is not masked by an associated hydrocele). Surgical exploration may be needed to distinguish this from torsion of the testis.

SCROTAL PAIN AT VARIOUS AGES

Acute epididymo-orchitis

Apart from mumps, acute epididymo-orchitis is usually caused by sexually transmitted pathogens in young males and by urinary tract pathogens in the older males. In older men it usually follows urinary tract obstruction and infection or instrumentation of the lower genitourinary tract.

Investigations

Blood cell count:	Leucocytosis

Urine microscopy and culture:	pyuria, bacteria and possibly *Escherichia coli*. A sterile culture suggests *Chlamydia* infection[3]
Tests for *Chlamydia*:	PCR kits
Ultrasound:	can differentiate a swollen epididymis from testicular tumour

Treatment[4]

- Bed rest
- Elevation and support of the scrotum
- Analgesics
- Antibiotics

  Sexually acquired: ceftriaxone 250 mg IM or ciprofloxacin 500 mg (o) as single dose plus doxycycline 100 mg (o) 12 hourly for 10 - 14 days

  Associated with urinary infection: amoxycillin/clavulanate 875/125 mg (child 22.5 mg/kg) (o) 12 hourly for 14 days *or* trimethoprim 300 mg (o) daily (child 6 mg/kg) for 14 days *or* cephalexin 500 mg (child 12.5 mg/kg) (o) 6 hourly for 14 days *or* (if resistance to above) norfloxacin 400 mg (o) 12 hourly for 14 days

Orchitis

Acute orchitis is invariably due to mumps and occurs during late adolescence. It is usually unilateral (Figure 104.3) but may be bilateral.



Figure 104.3 Mumps orchitis with a swollen tender testicle

Chronic orchitis may be due to syphilis, tuberculosis, leprosy or various helminthic infections such as filariasis. The majority are tuberculous in origin.

TESTICULAR NEOPLASM

Testicular tumours can occur at all ages but are more common in young men aged 20 - 30 years (teratoma) and 25 - 40 years (seminoma). Sometimes they can mimic an acute inflammatory swelling and present with acute pain.

STRANGULATED INGUINOSCROTAL HERNIA

It is possible that a supposed testicular torsion is found to be a strangulated inguinoscrotal hernia, usually an indirect inguinal hernia extending into the scrotum. It can be detected by careful palpation of the base (neck) of the scrotum.

TRAUMA AND HAEMATOCELES

A diffuse haematoma into the scrotum which causes no significant problems can follow surgery to the inguinal area, a blow to this area or a fracture of the pelvis. These conditions cause extravasation of blood distally. However, a haematocele of the tunica vaginalis can be either acute or an ‘old clotted haematocele’ following injury, such as a blow to the testis, or the drainage of a hydrocele.[5] Sometimes it can arise spontaneously. All types of haematoceles require surgical exploration to exclude testicular rupture or a tumour.

Trauma to the scrotum may produce urethral injury and extravasation of urine into the scrotum. This problem requires urgent surgery.

PROBLEMS OF SCROTAL SKIN

Sebaceous cysts are common and may be infected and require drainage. Fournier's gangrene (idiopathic gangrene of the scrotum) is a rare form of acute fulminating cellulitis affecting the scrotal skin. It usually develops suddenly and without any apparent cause. Gangrene of the scrotal skin appears early if the infection is not quickly checked with broad spectrum antibiotics. The end result is sloughing of the scrotal coverings, leaving the testes exposed.[5]

IDIOPATHIC SCROTAL OEDEMA

This uncommon and unusual condition usually occurs in boys aged 5 - 10 years. Scrotal swelling and redness begins gradually and spreads. Palpation reveals normal non-tender testes but torsion needs to be excluded in some instances of the swollen red scrotum. Idiopathic scrotal oedema is believed to be allergic in origin, either localised (e.g. insect bite) or globalised as part or urticaria. It sometimes results from exposure to cold water. The oedema is bilateral.

Practice tips

- Acute scrotal pain in infancy and adolescence should be regarded as torsion of the testis until proved otherwise.
- A history of recurrent transient pain (with or without swelling of the testis) in a young person means recurrent torsion. Urgent referral is essential.
- A pitfall is the phenomenon of ‘testis redux’ in which the descended testis undergoes torsion, is pulled into the superficial inguinal pouch by the cremasteric reflex and then becomes fixed by oedema.
- The development of an acute hydrocele should be regarded with suspicion.
- Beware of the strangulated inguinoscrotal hernia presenting as a testicular torsion.
- Consider dissecting aneurysm in an older person presenting with testicular pain.

REFERRED PAIN

Pain can be referred to the scrotal region from ureteric colic and quite commonly from disorders of the thoracolumbar spine, notably a disc disruption at the T12 - L1 level involving the L1 nerve root. The pain therefore may be referred or radicular.

WHEN TO REFER

- Any suspicion of torsion of the testis
- Sudden onset of acute scrotal pain at any age
- A history of recurrent transient testicular pain in a young man
- Presence of a tender testicular lump
- Presence of a haematocele surrounding the testis

Note: Referral should be most urgent, using the critical 4 - 6 hours guideline.

REFERENCES

1. Hutson J, Beasley S, Woodward A. Jones Clinical Paediatric Surgery. Oxford: Blackwell Scientific Publications, 2003; 185 - 8.
2. Wijesinha S. Torsion of the testis. Update, 1997; 2: 212 - 8.
3. Berger RE. Urethritis and epididymitis. Semin Urol, 1983; 1: 139.
4. Spicer J (Chair). Therapeutic Guidelines: Antibiotic (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 94 - 5.
5. Fry J, Berry H. Surgical Problems in Clinical Practice. London: Edward Arnold, 1987: 87 - 8.

CHAPTER 105. INGUINOSCROTAL LUMPS

He has a rupture, he has sprung a leake.

BEN JONSON (1573 - 1637), THE STAPLE OF NEWS

INGUINOSCROTAL LUMPS

Lumps in the groin are common to both sexes but males are likely to have a greater variety of swellings in this area and several may be associated with scrotal lumps.

LUMPS IN THE GROIN

The commonest swellings encountered in the groin or inguinal area are hernias (also known as 'ruptures') and enlarged lymph nodes. The diagnosis of a hernia is usually straightforward but it must be differentiated from other swellings, including Malgaigne's bulgings - these are not true hernias but diffuse swellings in both inguinal regions seen in people with poor lower abdominal musculature.[1] Table 105.1 lists the differential diagnoses of groin lumps.

Table 105.1 Differential diagnoses of a groin mass

Hernia - femoral, inguinal
Malgaigne's bulgings
Lipoma
Undescended testis
Spermatic cord swelling - encysted hydrocele, lipoma
Lymph node - localised, generalised
Haematoma (post femoral artery puncture)
Neoplasm - lipoma, others
Psoas abscess
Vascular anomalies
• saphenous varix
• femoral aneurysm

HERNIAS

The commonest types of hernias in the groin are inguinal, femoral and a combination of the two. Rare hernias in the region are obturator, Spigelian (low abdominal), preperitoneal inguinal and prevascular femoral. The basic parts of a hernia are shown in Figure 105.1 and important anatomical landmarks in Figure 105.2. An indirect inguinal hernia is a hernia through the deep inguinal ring, originating lateral to the inferior epigastric vessels, following the path of the processus vaginalis, and can traverse the whole length of the inguinal canal (Figure 105.3). In the male it closely approximates the spermatic cord and may enlarge as it passes through the superficial inguinal ring into the scrotuman inguino-scrotal hernia.


neck
peritoneum
sac
coverings
contents

Figure 105.1 Basic components of a hernia


deep inguinal ring
anterior superior
iliac spine
inferior
epigastric
artery
superficial
ring
femoral nerve
femoral artery
femoral vein
pubic tubercle
scrotum

Figure 105.2 Key landmarks in the left inguinal region: the deep inguinal ring lies above the mid-inguinal point (between the ASIS and the public tubercle); the femoral artery lies below this point


indirect
inguinal
hernia

Figure 105.3 Left indirect inguinal hernia: it emerges lateral to the inferior epigastric artery and passes into the scrotum medial to the pubic tubercle

Because of their narrow neck and oblique path in the inguinal canal, such hernias are often irreducible and are prone to lead to strangulation of entrapped bowel.

A direct inguinal hernia originates medial to the inferior epigastric vessels and protrudes through the posterior wall of the inguinal canal, and is therefore separate from the spermatic cord (Figure 105.4). It is almost always seen in men and rarely descends into the scrotum.[2] Due to a wider neck, strangulation and obstruction are most unusual. It must be emphasised that the distinction between a direct and an indirect inguinal hernia can be very difficult and the two may occur together.



Figure 105.4 Left direct inguinal hernia: it emerges medial to the inferior epigastric artery and bulges forward

A femoral hernia herniates through the femoral ring (also known as the femoral canal), which is the medial component of the femoral sheath. The hernia tends to bulge forwards and then upwards as it becomes larger. The neck is lateral to the pubic tubercle (Figure 105.5).



Figure 105.5 Left femoral hernia: its neck is lateral to the pubic tubercle and it lies below the inguinal ligament

Femoral hernias are often small, usually occur in females and may be unnoticed by the patient. They are particularly liable to produce bowel obstruction or strangulation.[2]

Guidelines with hernias

Acquired hernia

- Always examine the scrotum and both sides
- Frequently bilateral
- Result from muscular weakness
- Commonest - direct inguinal and femoral
- Predisposing factors
    - age (more common with increasing age)
    - obesity
    - pregnancy
- Precipitating factors (related to above factors)
    - increased intra-abdominal pressure
        - difficulty of micturition
        - straining at stool (constipation)

        - chronic cough (e.g. bronchitis)
        - straining or lifting heavy objects
    - nerve damage (e.g. post appendicectomy)
- complications
    - intestinal obstruction (Table 105.2)
    - incarceration
    - strangulation
    - sliding

Table 105.2 Symptoms and signs of hernial obstruction

Colicky abdominal pain
Nausea and vomiting
Constipation and failure to pass flatus
Abdominal distension
High-pitched tinkling bowel sounds
Local tenderness and swelling of the hernia
No expansile cough impulse

Clinical features

The main symptoms and signs:[1]

- lump
- discomfort or pain
    - a dragging pain
    - worse after standing or walking
    - referred to testicle (indirect inguinal)
- testicular pain - referred or with compression of the spermatic cord
- expansile impulse on coughing

Note:

- A femoral hernias is easily missed in obese patients.
- Larger femoral hernia are often irreducible.
- Always attempt reduction in the recumbent position (direct hernias usually reduce easily).
- In over 50% of strangulated obturator hernias, pain is referred along the geniculate branch of the obturator nerve to the knee.[1]

Treatment of hernias

Surgery[3]

All symptomatic hernias require repair and all femoral hernias should be repaired. Obstructed and strangulated hernias require urgent surgery. The risk of strangulation is greatest with femoral hernias, moderate with indirect inguinal hernias and least with direct inguinal hernias.

Conservative

Asymptomatic inguinal hernias in patients with associated medical conditions, and who pose a significant operative risk, can be treated conservatively. A suitable truss to control a small inguinal hernia is a rat-tailed spring truss with a perineal band to prevent slipping.[1] Such a truss must be used with care and patients well instructed in its proper use. Trusses must always be applied over the inguinal canal with the patient lying flat and with the hernia reduced. Difficult reduction can be aided by a warm moist towelette.

HERNIAS IN CHILDREN

The most common hernias in infants are inguinal hernias and umbilical hernias.

Umbilical hernias

Features

- Soft round skin-coloured lump in umbilicus
- May increase in size in first few months
- Not painful, not tender to palpate, easily reducible
- Swelling disappears when child asleep
- Usually gradually disappears
- Most disappear by age 12 months
- Larger ones usually disappear by age of 4 years
- Consider hypothyroidism

Management

- Explanation and reassurance
- No treatment required
- Do not tape or strap (may cause strangulation and does not help)
- Refer for repair if still present at age 4 years

Inguinal hernia

Features

- More common in premature infants and boys
- Present with groin lumps - may be intermittent sightings (e.g. when crying)
- May cause intermittent pain or discomfort
- If irreducible is strangulated

Note: Bowel strangulation is a possibility

Management

Rules for surgical intervention:

- General rule is ASAP, especially in infants and for irreducible ones
- Reducible herniae - the ‘6 - 2’ rule
    - birth - 6 weeks: surgery within 2 days
    - 6 weeks - 6 months: surgery within 2 weeks
    - over 6 months: surgery within 2 months

SCROTAL LUMPS

The scrotum contains the testes and distal parts of the spermatic cords, covered by layers of fascia and the dartos muscle. The testes are invested with tunica vaginalis derived from the peritoneal cavity during their descent.[1]

Disorders of the scrotum may be acute or chronic and bilateral or unilateral. Lumps may be cystic, solid or otherwise, such as a varicocele, oedema and hernia. Solid lumps include a testicular tumour, epididymo-orchitis, and torsion of the testes. Cystic lumps include hydroceles, epididymal cysts and spermatoceles, and resolving extravasation. A comparison of scrotal lumps appears in Figure 105.6 and Table 105.3. Lumps in the scrotum usually develop from deeper structures, particularly the testes and their coverings, rather than scrotal skin.[1] Refer to Figure 105.6 for a comparison of scrotal lumps.


vas deferens
epididymis
testis
scrotum
Normal
firm mass
Testicular tumour
? secondary hydrocele
Swelling of the epididymis
(separate from testis)
(e.g. chronic epididymis or cyst)
blood
Haematocele
dilated veins
Varicocele
serous fluid
Hydrocele

Figure 105.6 Basic comparison of scrotal lumps

Table 105.3 Features of scrotal lumps

	Possible clinical setting	Position	Palpation	Trans-illumination
Hydrocele	Any age Primary or secondary • tumour • infection • torsion	Confined to scrotum Anterior: surrounds testis except posteriorly	Smooth, pear-shaped Lax or tense Testis impalpable, non-tender	Yes
Cyst of epididymis Epididymal cysts and spermatoceles clinically similar	Asymptomatic or dragging sensation	Behind and above testis	Smooth and tense Multilocular swelling Testis easily palpable Appears separate from testis	Yes
Chronic epididymo-orchitis	Non-specific Tuberculosis Chlamydia (Occasional associated small hydrocele)	Behind and above testis	Firm swelling Hard and craggy Normal testis	No
Varicocele	Dragging discomfort	Usually left-sided Along line of spermatic cord Above testis	Soft, like bunch of worms or grapes Collapses when patient supine and testis elevated Testis often smaller	No
Carcinoma	Young men 20 - 40 Painless lump Loss of testicular sensation	In body of testis Usually felt anteriorly May be hydrocele	Enlarged firm testis Feels heavy if large Normal epididymis (palpable)	No

The cardinal sign of a true scrotal mass is that it is possible to palpate it from above, i.e. get above the lump.

The patient usually presents with pain or a lump.

Examination of the scrotum

The scrotum should be examined with the patient supine and then standing. The left testis usually hangs lower than the right. On inspection note any sebaceous cysts in the scrotal skin (common), scabies if there are very pruritic nodules, and scrotal oedema, which causes taut pitting skin. Careful palpation will elicit the relevant structures in the scrotum. Gently palpate each testis and epididymis between the thumb and the first two fingers. The spermatic cord is palpable as it enters the scrotum after passing through the superficial ring and the testis and epididymis are readily palpable.

After palpation test for translucency of any swelling in a darkened room by shining the beam of a strong torch from behind the scrotum through the swelling. Transilluminable swellings that light up with a red glow include hydroceles and cysts of the epididymis. Swellings that contain blood or other tissue, such as testicular tumours and most hernias, do not illuminate.

Unilateral scrotal swelling

It is important to determine whether the lump is inguinoscrotal or scrotal. It is scrotal if it is possible to get above the lump. If it is not possible to get above the lump then it is a large inguinal hernia or a combined hernia and hydrocele (Figure 105.7). This palpation should be co-ordinated with the cough impulse. The next feature to determine is whether the testis and/or epididymis can be palpated or whether they are obscured by a swelling such as a hydrocele.



Figure 105.7 Difference between a true scrotal swelling and an inguinoscrotal swelling

Small testes

Testes are considered small in adults if less than 3.5 cm long. Small firm testes, 2 cm long or less, are a feature of Klinefelter's syndrome. Small soft testes indicate atrophy, which may follow mumps orchitis, oestrogen therapy, androgen deficiency or anti-androgen therapy, hypo-pituitarism, cirrhosis and other related conditions.

HYDROCELE

A hydrocele is a collection of clear amber fluid in the tunica vaginalis and can be primary or secondary. If a hydrocele develops it is im-portant to rule out intrascrotal disease, such as a tumour or infection. Ultrasound examination of the scrotum is helpful in assessing the state of the testis in the presence of a hydrocele. Hydroceles may be symptomless or cause dragging discomfort in the scrotum and groin.

Hydroceles in the neonate

Hydroceles present at birth are invariably communicating (failed closure of the processus vaginalis) or (less commonly) non-communicating, where the tunica vaginalis contains fluid. Transillumination will prove that it is cystic but if the diagnosis is in doubt perform an ultrasound. Hydroceles may vary in size from day to day and can appear quite large. They do not extend proximal to the external inguinal ring so it is possible to palpate above them. There is no impulse on crying or straining. Of the 5% of those born with a hydrocele, most will resolve spontaneously within 12 months. If the hydrocele is very large or persists beyond 12 months, surgical intervention should be considered.

Treatment of a primary hydrocele in adults

Surgery is the most effective long-term treatment. A primary hydrocele can be treated by simple aspiration but the fluid usually reaccumulates and there is risk of bleeding or infection with repeated procedures.[4] However, aspiration followed by injection of a sclerosant agent (e.g. dilute aqueous phenol or sodium tetradecyl sulphate [STD]) can prevent fluid accumulation and after two or three times can often cure the problem. This sclerotherapy may be complicated by pain and inflammatory reaction to the sclerosant.

Method

1. Inject LA into the scrotal skin down to the sac.
2. Insert an 18- or 19-gauge intravenous cannula through this site into the sac and remove the stilette, leaving the soft cannula in the sac (Figure 105.8).



Figure 105.8 Aspiration of hydrocele

3. Remove the serous fluid initially by free drainage, possibly aided by manual compression on the sac and then by aspiration with a 20 mL syringe.
4. Record the volume.
5. Inject 2.5% sterile aqueous phenol or STD into the empty sac (10 mL for 200 mL of fluid removed, 15 mL for 200 - 400 mL and 20 mL for over 400 mL).

If the hydrocele re-forms (10% of cases) the procedure can be repeated after 6 weeks.

Encysted hydrocele of the cord

This is a localised fluid-filled segment of the processus vaginalis within the spermatic cord. It is palpable as a cystic lump in the upper scrotum above the testis. It characteristically moves down with traction on the testis. No treatment is usually needed.

EPIDIDYMAL CYSTS

These are common and often multiple. Usually found in middle aged/elderly men. The majority of epididymal cysts contain a clear colourless fluid. If the cysts communicate with the vasa efferentia a spermatocele filled with whitish fluid containing spermatazoa may form.

Epididymal cysts may be asymptomatic or they may cause discomfort and cosmetic embarrassment and should be excised. Fertility may be impaired in patients undergoing bi-lateral cyst excision.

Aspiration and injection of sclerosant agents can also be used for epididymal cysts.[4]

VARICOCELES

A varicocele is a varicosity of the veins of the pampiniform plexus (Figure 105.6). It is seen in 8 - 10% of normal males and occurs on the left side in 98% of affected patients, due to a mechanical problem in drainage of the left renal vein. A relationship with infertility has been observed but its nature is controversial.

Most varicoceles are asymptomatic and incidental findings. They can cause a dragging discomfort in the scrotum. Investigation is usually not necessary but an ultrasound is useful where the diagnosis is doubtful or a neoplasm is suspected. Treatment is indicated if it is symptomatic or for infertility. Firm-fitting underpants may relieve discomfort.[1] Surgical treatment is by venous ligation, above the deep inguinal ring. Ligation is indicated if there is any reduction in the size of the left testis.

HAEMATOCELES

These can be either acute, resulting from trauma such as a fall astride, sports injury or tapping of a hydrocele, or chronic, where there is no obvious history of injury. Haematoceles are anterior to the testis and not trans-illuminable. Surgical drainage is required with acute injury where there is a possibility of testicular rupture (associated urethral injury has to be considered); and a tumour has to be excluded with the chronic type. Pressure atrophy of the testis can occur with injury and is much more common if early drainage is not instigated.

SPERM GRANULOMAS

- Firm tender lumps
- Post-vasectomy-at cut end of vas

Treatment

- Leave to resolve
- Consider NSAIDs.
- Refer for excision if symptomatic and enlarging.

TESTICULAR TUMOURS[5,6,7]

A mass that is part of the testis, and solid, is likely to be a tumour. Malignant testicular tumours account for about 1-2% of malignant tumours in men. They mainly affect fit young men and represent the commonest neoplasm in men aged 15 - 44 years in Australia. (Table 105.4). 95% of testicular tumours arise from the germ cells and for practical purposes are classified into

- seminomas 40%
- non-seminoma germ cell tumours (NSGCT) 60% [6]

Table 105.4 Testicular tumours[1,4,5]

Tumour	Incidence (%)	Peak incidence (years)
Seminoma	40	25 - 40
Teratoma	32	20 - 35
Mixed seminoma/teratoma	14	20 - 40
Lymphoma	7	60+

Other tumours (e.g. interstitial - Leydig, gonadoblastoma)	uncommon	variable

Clinical features

- Young men aged 15 - 40 years
- Painless lump in body of testis (commonest feature)
- Up to 15% experience pain
- Loss of testicular sensation
- Associated presentations (may mask tumour)
  - hydrocele
  - varicocele
  - epididymo-orchitis
  - swollen testis with trivial injury
  - gynaecomastia (teratoma)

Risk factors

Those at high risk include a family history and a history of:

- Cryptorchidism (x5 risk)
- orchidopexy
- testicular atrophy
- previous testicular cancer

Golden rules

- All solid scrotal lumps are malignant until proved otherwise and must be surgically explored.
- Beware of hydroceles in young adults.
- Tumours can mimic acute epididymo-orchitis - the so-called ‘inflammatory’ or ‘flash fire’ presentation.[3]

Metastases

Testicular tumours spread by direct infiltration via the lymphatics and the bloodstream. Metastases typically occur in the para-aortic nodes so it is advisable to palpate carefully from the umbilical area upwards. However, they are best detected by a CT scan of the abdomen and chest. Metastases also occur in the neck, liver and chest.

Investigations

Investigations to aid diagnosis include:[3]

- ultrasound of the testis: can detect and diagnose with considerable precision underlying testicular lumps plus any invasion of the tunica
- tumour markers: alpha-fetoprotein and beta human chorionic gonadotrophin - indicates teratomas
- lactate dehydrogenase - monitors secondary spread

Investigations for staging include:

- chest X-ray
- CT scanning of abdomen, pelvis and chest for node involvement (any spread is direct to the para-aortic nodes)

Note: Avoid scrotal needling biopsy because of the risk of tumour implantation in the scrotal wall. Avoid scrotal incisions for surgery.

Treatment

The initial treatment is orchidectomy through on inguin incissionwith inguinal division of the spermatic cord. Specialised treatment that depends on the staging of the tumour gives good results for seminoma, which is very sensitive to radiotherapy. The results for teratoma and NSGCTs in general are not as satisfactory as for seminoma.

Prognosis is good for most testicular tumours with 5 year cure rates of 90 - 95%.[3] A comparison of the testicular tumours is summarised in Table 105.5.

Table 105.5 Comparison of the common testicular cancers[6]

	Seminoma	Non-seminoma (NSGCT)
Typical age	25 - 40 years	< 35 years
Incidence	40%	60%
Growth rate	Slow	Rapid
Nature	Solid	Mixed - solid + cystic
Stage at presentation	90% - stage 1	60% - stage 1
Tumour markers		
α-FP	Never	Common
β-HCG	Occasional	Common
Treatment	Inguinal orchidectomy + radiotherapy	Stage 1: orchidectomy Relapse: chemotherapy
Sensitive to chemotherapy	+ + +	+ + +
Sensitive to radiotherapy	+ + +	±
Prognosis	Stage 1: 98% 5 year survival Overall: > 85%	Stage 1: 75% cure by surgery Otherwise varies

Follow-up[7]

CT scans of the chest, abdomen and pelvis are performed regularly every few months in the first 2 years and less often after that.

Serum tumour markers are checked at each visit.

Screening and testicular self-examination

Studies of testicular cancer have shown the benefits of early detection.[8] However, studies to date indicate that there is insufficient evidence to screen routinely for testicular cancer in asymptomatic patients considering also that it is uncommon (1:265 men). It is recommended for those at high risk. This screening includes colour Doppler ultrasound and tumour markers. Evidence also indicates that, to date, there is little evidence to show that those performing testicular self-examination are more likely to detect early stage tumours or have better survival than those who do not.[9]

UNDESCENDED TESTES (CRYPTORCHIDISM)

A testis that is not in the scrotum may be ectopic, absent, retractile or truly undescended. The incidence is 2 - 4% in full-term males, 20% in premature male births and 1% at 1 year. More than two-thirds of undescended testes are located in the superficial inguinal pouch; that is, they are palpable in the groin.

Undescended testis

An undescended testis is one that cannot reach the bottom of the scrotum despite manual manipulation. After the indirect inguinal hernia it is the most common problem in paediatric surgery. The testis is usually normal but may become secondarily dysplastic if left outside the scrotum.

A truly undescended testis is one that has stopped in the normal path of descent and can occupy intra-abdominal, inguinal canal, emergent (just outside the external ring), high scrotal and mid-scrotal positions (Figure 105.9).[4] The cause of maldescent is most probably mechanical.



Figure 105.9 Undescended testis: arrested in the line of descent

Retractile testis

A retractile testis is one that can be manipulated into the scrotum irrespective of the position in which it is first located. It is a common condition. The testis can be present in the scrotum under circumstances such as a warm bath but retracted out of the scrotum when cold. Cremasteric contraction is absent in the first few months after birth and is maximal between 2 and 8 years.[1]

Ectopic testis

An ectopic testis is one that has left the normal path of descent and cannot be manipulated into the scrotum. It can be found in the perineum, upper thigh (femoral), base of the penis (prepubic), anterior abdominal wall or in the superficial inguinal pouch (Figure 105.10). True ectopic testes form only about 5% of all undescended testes.



Figure 105.10 Undescended testis: ectopic

Ascending testis

An 'ascending' testis is one that was in the scrotum in infancy but subsequently moved back to the groin because the spermatic cord failed to elongate at the same rate of body growth.

Examination[10]

The examination of the testes should take place in a warm room and relaxed environment. Begin by placing one finger on each side of the neck of the scrotum to prevent the testes from being retracted out of the scrotum by the other hand. The scrotum is then carefully palpated for a testis. If impalpable the fingertips of one hand are placed just medial to the anterior superior iliac spine and moved firmly towards the pubic tubercle where the other hand waits to entrap the testis should it appear. The diagnosis then depends on carefully determining the range of movement.

The problem of non-descent

- Testicular dysplasia
- Susceptible to direct violence (if in inguinal region)
- Risk of malignant change (seminoma) is 5 - 10 times greater than normal

Optimal time for surgery

The optimal time for orchidopexy is 6 - 12 months of age.[10] It is considered to be satisfactory as long as testis is in the scrotun by 2 years. The production of spermatozoa is adversely affected in undescended testes from the age of 2 years onwards.[8] Exploration for the uncommon impalpable testis is worthwhile: 50% salvage rate, while in the other 50% either there is no testis or an abnormal and potentially neoplastic testis is removed.[10]

The advantages of early orchidopexy are summarised in Table 105.6.

Table 105.6 Advantages of early orchidopexy (1 year)[4]

Provides optimal chance of fertility

Corrects indirect inguinal hernias (coexists in 90%)
Reduces risk of trauma
Reduces risk of torsion
Reduces psychological consequences
Probably lessens the risk of malignancy (seminoma)

Hormone injections

Injections of chorionic gonadotrophic hormones are generally not recommended. They are ineffective except for cases of borderline retractile testes.

VASECTOMY

Dangerous features (beware)

- Umarried
- Young < 35 years old
- No children
- Emotional crisis
- Spouse not involved

Post operative

- Avoid strenous activity and sex from 4-7 years.
- Use alternative cotraception until negative semen analysis.
- First semen analysis at 3 months.
- Inform not immediately infertile.
- Need to ejaculate 20 times.
- Prone to chronic orchidalgia (orchalgia).

Practice tip

Inguinal scrotal lumps

General rules for optimal times for surgical repair in children:

- inguinal hernia - ASAP
- umbilical hernia - aged 4 years (most resolve)
- femoral hernia - ASAP
- undescended testes - 6 to 18 months
- hydrocele - after 12 months if still large most resolve
- varicocele - leave and review

Complications

- Haematoma 2 - 3%
- Wound infection approx. 2%
- Congestive epidedymitis (painful testis/epidedymis for 2 weeks) treat with NSAIDs
- Sperm granuloma 15 - 40%
- Epididymal apsts
- Formatur of sperm antibiotics ~ 50%. May be problem if subsequently requested for vasectormy reversal
- Chronic orchalgia ≅ 1% = post-vasectomy syndrome

- Failure rate 2-3/1000
- Psychological and psychosexual problems

Other facts

- No difference in ejaculate appearance (sperm maker up 1% speculate volume)
- No known association with prostate and testicular cancers

REFERENCES

1. Fry J, Berry H. *Surgical Problems in Clinical Practice*. London: Edward Arnold, 1987: 79 - 92.
2. Davis A, Bolin T, Ham J. *Symptom Analysis and Physical Diagnosis* (2nd edn). Sydney: Pergamon Press, 1990: 212 - 20.
3. Hunt P, Marshall V. *Clinical Problems in General Surgery*. Sydney: Butterworths, 1991: 329 - 42.
4. Bullock N, Sibley G, Whitaker R. *Essential Urology*. Edinburgh: Churchill Livingstone, 1989: 287 - 318.
5. Messing EM. Testicular seminoma. In: *Current Therapy in Genitourinary Surgery* (2nd edn). New York: Resnick & Kursh, 1992: 168.
6. Stockler M, Boyer M. How to treat: Testis cancer Australian Doctor, 2003; 21 February, 1-VIII
7. Boyer M.J. Diagnosis and treatment of testicular cancer. Current therapeutics, 2001: 26 - 9.
8. Bolse J, Vogelzang NJ, Goldman A et al. Impact of delay in diagnosis on clinical stage of testicular cancer. Lancet, 1981; 2: 970 - 2.
9. RACGP. *Guidelines for Preventative Activities in General Practice* (5th edn). Melbourne: RACGP, 2005: 36.
10. Hutson J, Beasley S, Woodward A. *Jones Clinical Paediatric Surgery*. Oxford: Blackwell Scientific Publications, 20 03: 181 - 4.

CHAPTER 106. DISORDERS OF THE PENIS

Ironically there is no organ about which more misinformation has beenperpetuated than the penis.

William Masters & Virginia Johnson, Human Sexual Response

DISORDERS OF THE PENIS

The most common penile disorders are those of psychosexual dysfunction and STIs, but there are many other problems and these are most often related to the foreskin.

DISORDERS AFFECTING THE FORESKIN AND GLANS

Phimosis

Phimosis is tightness of the foreskin (prepuce), preventing its free retraction over the glans penis. The foreskin can be adherent to the glans penis even up to 5 - 6 years of age. It gradually separates until it becomes non- adherent, usually by the age of 6. Forcible retraction of the foreskin of any boy should be discouraged.

True congenital narrowing of the preputial orifice is rare. If the foreskin cannot be retractedby the age of 7 years and is causing symptoms such as balanitis, thencircumcision is recommended. Ballooning of the foreskin during micturition can be a feature, and if *persistent consider consequent vesico-ureteric reflux*. However, mild ballooning iscommon in the process offoreskin separation from the glans. It should resolve as long as urine drains freely and no pain. It rarely causes urinary retention.

Treatment

Inflammatory phimosis can be treated by local corticosteroid cream (e.g. 0.05% betamethasone valerate cream qid for 4 weeks applied generously to the tight shiny part of the foreskin where inner skin meets the outer skin). If unsuccessful, can change to a higher potency steroid cream.True scarring requires circumcision.[1] Some patients with true phimosis may have problems once they start to have intercourse. They require circumcision. Generally, circumcision is rarely needed.[1]

Paraphimosis

In paraphimosis the foreskin is retracted, swollen and painful (Figure 106.1). This is because it has been pulled back over the glans, trapped behind it, and cannot be pulled forwards again.This problem occursin older boys and the elderly, especially if a mild degree of phimosis is already present. Typically it occurs when the penis is erect or after catheterisation.



Figure 106.1 Paraphimosis showing retracted, swollen, oedematous foreskin

Management

Urgent manual reduction should be attempted first. It is usually performed without anaesthesia but a penile block (never use adrenaline in LA or general anaesthesia) or a generous application of 2% lignocain gel or EMLA cream (5 - 10 minutes beforehand) may be appropriate.

Note: Do not apply ice.

Method 1

The glans penis and oedematous tissue distalto the constricting ring of foreskin are gently squeezed for several minutes to reduce the oedema. Using a lubricating jelly, manual reduction can then be performed by trying to advance the prepuce over the glans with the index fingers while gently compressing the glans with both thumbs (Figure 106.2).



Figure 106.2 Acute paraphimosis: method of manual reduction

Method 2

- Take hold of the oedematous part of the glans in the fist of one hand and squeeze firmly. A gauze swab or cool towelette will help to achieve a firm grip (Figure 106.3).



Figure 106.3 Acute paraphimosis: squeezing with a swab method

- Exert continuous pressure until the oedema passes under the constricting collar to the shaft of the penis.
- The foreskin can then usually be pulled over the glans.

Method 3

Immediate referral is necessary if manual reduction methods fail. As an emergency a dorsal slitincision can be made in the constricting collar ofskin under local or general anaesthetic (Figure 106.4). The incision allows the foreskin to be advanced and reduces the swelling. Circumcision should be performed some days later when the inflammation has settled.[2]



Figure 106.4 Acute paraphimosis: dorsal slit incision in the constricting collar of skin

Balanitis (balanoposthitis)

Balanitis is inflammation of the foreskin which usually affects the glans penis and tissues behind the foreskin (balanoposthitis). The inflammation may simply be redness or irritation of the glans and foreskin or bacterial infection. This quite common problem may be due to *Candida albicans*infection, but in infants may be caused by wet nappies and in the elderly by diabetes alone or with an associated organism. Severe cases may have purulent discharge with spreading cellulitis and require systemic antibiotics.

Men presenting with balanitis should be screened for:

- diabetes
- Reiter's syndrome (especially if asymptomatic)

Treatment

- Mild cases may be treated with gentle saline bathing, a barrier cream or hydrocortisone 1% cream to tip of penis, and then application of an antibacterial ointment, e.g. fucidin 4times daily, if more severe
- Take swabs for culture
- Careful washing behind foreskin

If yeasts present:

- topical nystatin or miconazole or clotrimazole cream

If trichomonads present:

- metronidazole or tinidazole (oral treatment)

If bacteria present (usually cellulitis):

- appropriate antibiotic using a narrow longish nozzle (e.g. chloramphenicol or chlortetracycline)
- oral antibiotics may be needed

Balanitis xerotica obliterans

Thickening of the foreskin with skin pallor suggests balanitis xerotica obliterans. It results in progressive phimosis, typically in late childhood, usually 10 - 12 years. It may respond to corticosteroid cream if it is mild, but circumcision is usually indicated.

FORESKIN HYGIENE

The normal foreskin in infants and children does not need special care and should not be retracted for cleaning from birth to 5 years of age when the foreskin can be retracted in 90% of boys. From the age of 6 or 7, males can practise proper hygiene by gentlyretracting the foreskin and washing the area as often as washing behind the ears. However, it is not a significant issue until puberty when adhesions and smegma have reduced.

Basic rules

- The foreskin should only be retracted by its owner
- The foreskin should not be retracted forcibly
- Once retraction has been demonstrated, encourage daily retraction and gentle cleansing in the bath or shower

Instructions to patients

- During a shower or bath slide the foreskin back towards your body (Figure 106.5).



Figure 106.5 Foreskin hygiene: sliding foreskin back for washing

- Wash the end of the penis and foreskin withsoap and water.
- After washing the area, dry the end of yourpenis and foreskin and then replace the foreskin.
- If the foreskin has a tendency to become irritated and smelly, slide the foreskin back sufficiently to allow free urination.

LUMPS UNDER THE FORESKIN OF CHILDREN

Consider:

- smegma cysts/deposits (yellow - white lumps beneath prepuce) - they are normal and can discharge a white ooze

- dermoid (ventral and midline within skin)

CIRCUMCISION

Apart from abnormalities of the foreskin and religious reasons, circumcision for social reasonsis generally discouraged. A policy statement from the Paediatric and Child Health Division of the RACP does not usually recommend circumcision. Arguments against circumcision are that it is not natural, it is unnecessary, it carries a small but significant risk of morbidity and mortality, and is associated with meatal stenosis. Circumcision is now performed less frequently and most practitioners do not appear to favour it for social reasons. Proponents, however, argue that it reduces the risk of periurethral bacterial colonisation, urinary infection, systemic infections such as septicaemia, carcinoma of the penis, and STIs including HIV. Indications for circumcision include phimosis, paraphimosis (occasionally), recurrent balanitis, balanitis xerotica obliterans and serious urinary tract anomaly. Boys with hypospadiasshould never be circumcised, since the foreskin may be a vital source of skin for subsequent repair.[2]

Complications of circumcision:

- bleeding
- infection (local/septicaemia)
- ulceration of glans/meatus
- meatal stenosis
- penile deformity

Absolute contra indications:

- hypospadius and other congenital abnormalities
- chordee (painful dorsolateral curvature of the penis during erection which interferes with sexual intercourse)
- buried penis
- sick ‘unstable’ infants
- FH bleeding
- inadequate experience

Using a dorsal slit of the foreskin as an alternative to routine circumcision produces a cosmetically unacceptable result and should only be used as an emergency measure.

Frenuloplasty

A congenitally tight frenulum may lead to a tear during intercourse. Repeated bleeding occurs. Division of the frenulum and suturing in the opposite direction is preferable to circumcision.

THE ‘BURIED’ PENIS

In this condition the penis looks small or the foreskin quite huge and tight. There is failure of skin fixation at the base of the penis with possible excessive fatpad at the base. Erections do not stretch the foreskin appropriately. The condition does not invariably resolve spontaneously. Early referral to a paediatric surgeon is advisable.

DISORDERS AFFECTING THE URETHRAL MEATUS

Meatal stenosis

Meatal stenosis or stricture may be congenital or acquired. It may beacquired in the circumcised child due to abrasion and ulceration of the tip of the glans. The incidence can be reduced by the application of Vaseline on the glans after circumcision.[1] Uncommon causes are direct trauma during circumcision and irritation from ammoniacal dermatitis. Meatal ulceration predisposesto meatal stenosis. It usually presents as pain during micturition oras slight bleeding on thenapkin. In the child with mild meatal stenosis gentle dilation can be applied. Severe cases require surgical correctionby meatotomy.

Catheter trauma is the usual cause in adults.

Hypospadias

Hypospadias is a condition where the urethra opens on the underside or ventral aspect of the penis. It occurs in 1 in 300 males.[1] Congenital hypospadias may be glanular (most common), coronal, penile or perineal.[3] The ventral shaft of the penis is angulated and shortened, i.e. chordee.

Hypospadias may cause the stream of urine to be deflected downwards or splash or drip back along the penile shaft. Unless it is glanular, surgical repair is usually advised, using the available foreskin. Chordee is corrected at thesame time to allow eventual successful sexual intercourse. These boys should not undergo routinecircumcision.

Epispadias

Epispadias is where the urethra opens at the base of the penis, on its dorsal aspect. It occurs in 1 in 30 000 males.[1] Most patients are incontinent of urine because of a deficient bladder neck.

PENILE WARTS

Penile warts are usually fleshy, papillomatous multiple outgrowths, commonly found around the coronal sulcus, the adjacent prepuce and the meatus(Figure 106.6). They are caused by human papilloma virus and usually transmitted sexually. Look for warts within the meatus by allowing gentle dilation of the distal urethra with mosquito forceps. Treatment includes keeping the affected areas cool and dry and applying with extreme care 25% podophyllinsolution in compound benzoin tincture or podophyllotoxin 0.5% paint to each wart 12 hourly for 3 days,[4] followed by 4-day break (repeat weekly until disappearance) or imiquimod 5%cream to each wart, 3 times per week for 4 to 12 weeks. Tenper cent recur at 3 months.



Figure 106.6 Penile (genital) warts caused by the human papilloma virus

PEARLY PENILE PAPULES

These are very small regular round lumps that appear on the corona ofthe glans of the penis. They are common and are often first noticed by adolescent males who should be reassured that they are normal variants. They must not be treated as penile warts.



Figure 106.7 Pearly penile papules

PENILE ULCERS

A common cause related to sexual activity is trauma to the frenulum if it is congenitally tight. Such traumatic ulcers may be slow to heal and the frenulum may needsurgical division. The ulcers may resemble a venereal ulcer (e.g. syphiliticchancre, herpes simplex or AIDS). Another important (although rare) cause is carcinoma of the penis. Various causes are listed in Table 106.1.

Table 106.1 Causes of penile lesions

Non-ulcerative
Balanitis
• *Candida albicans*
• Reiter's syndrome
• diabetes mellitus
• poor hygiene
Skin disease
• psoriasis
• lichen planus
Venereal warts
Ulcerative
Trauma (tender)
Carcinoma (non-tender)
Herpes simplex (tender)

Syphilis (non-tender)
Chancroid (tender)
Behçet's syndrome

CARCINOMA OF THE PENIS

Carcinoma of the penis is rare, occurring at a rate of 1 in 100 000 of the male population.[2] There is an association with the non-circumcised person, thetheory being that smegma may be carcinogenic.

Carcinoma usually starts as a nodular warty growth (or ulcer) on the glans penis or in the coronal sulcus. Initially it may resemble a venereal wart. The presentingsymptom may be a bloodstained or foulsmelling discharge as the lesion is usually hidden by the foreskin. It is usually seen in elderly patients with poor hygiene.[3] Associated lymphadenopathy, which is present in 50%of patients on presentation, may be infective or neoplastic.

PRIAPISM[5]

Priapism is a persistent painful erection not associated with appropriate sexual stimulation. The corpora cavernosa are engorged and painful, but the corpus spongiosum and glans remain flaccid.[2] The cause is usually poor venous drainage, but 10% reflect post-traumatic excess arterial inflow. Penile Doppler ultrasound distinguishes the differences. Venous priapism should be regarded as an emergency;if prolonged it may lead to venousthrombosis, resulting in impotence. Radiological embolisationmay be required.

Venous priapism is usually associated with intracavernosal injection for impotence with prostaglandin or papaverine. Haematological disorders such as sickle cell anaemia and leukaemia, metastatic malignant infiltration, spinal cord injuries, IV drug users and drugs such as anticoagulants, marijuana, phenothiazines and some antihypertensives are uncommon causes. Some cases are idiopathic.

Management[5] is an urgent blood film to exclude polycythemia and leukaemia; then, under local anaesthesia, repeated saline flushing and aspiration of thick blood from the ipsilateral corpora cavernosa with a 16-gauge needle through the glans penis.If resolution is incomplete, use a very slow injection of 10 mL of saline containing aramine 1 mg into the corpus cavernosum, followed by massage. A poorresponse at 1 hour may require a second aramine injection, carefully monitoring for hypertension. Venous bypass surgery is rare ifdrainage can be established.

PEYRONIE'S DISORDER

Peyronie's disorder is a fibrotic process, sometimes associated with Dupuytren's contracture, that affects the shaft of thepenis and results in discomfort and deformity on erection. It affects 4% of men over 40 years, and one in three have penile pain and bleeding upon erection. Typically, the patient presentswith painful ‘crooked’ erections. There is abnormal curvature of the erect penis. The penile deformity may prevent satisfactory vaginal penetration. On examination a non-tender hard plaque may be palpable inthe shaft of the penis. Mild cases require reassurance and daily vitamin E tablets 200 mg tds for 6 months to reduce discomfort.[5] The problem may increase, remain static or spontaneously lessen over 1 - 2 years. Occasionally, surgicaltreatment by penile tuck isindicated if the patient's erection is so deformed that sexual intercourse is difficult, or rarely by penile implant if the patient is impotent.[2]

‘FRACTURED’ PENIS

A ‘fractured’ penis describes sudden rupture of the penile erectile tissue during intercourse, resulting in a snapping sensation with severe pain. Themanagement is urgent urological consultation for surgical repair. The disruption can affect the corpus spongiosium (has a better prognosis) or the corpora cavernosa, in which case permanentimpotence is a possible complication.

HAEMATOSPERMIA

Haematospermia, which is blood in the semen, presents as a somewhat alarming symptom. It is sometimes encountered in young adults and middleaged men. The initial step is to determine that the blood is actually in the semen and not arising from warts inside the urethral meatus or from the partner.

It usually occurs as an isolated event but can be secondary to urethral warts or prostatitis, or with prostatomegaly or prostatic tumour (especially in elderly patients). If a micro-urine shows no accompanying haematuria, and prostatic specific antigen and blood pressure are normal, reassurance and a 6-week review is appropriate as spontaneous cessation of haematospermia is the rule.

REFERENCES

1. Hutson J, Beasley S, Woodward A. *Jones Clinical Paediatric Surgery*. Oxford: Blackwell Scientific Publications, 2003: 195 - 8.
2. Bullock N, Sibley G, Whitaker R. *Essential Urology*. Edinburgh: Churchill Livingstone, 1989: 287 - 99.
3. Hunt P, Marshall V. *Clinical Problems in General Surgery*. Sydney: Butterworths, 1991: 365 - 8.
4. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 97 - 8.
5. Lawson P. Difficult male urological problems. Urology Seminar Proceedings. Box Hill Hospital, 1997: 2 - 4.

CHAPTER 107. DISORDERS OF THE PROSTATE

Mind and body, like man and wife, do not always agree to die together.

CHARLES COLTON 1780 - 1832

DISORDERS OF THE PROSTATE

The main function of the prostate gland is to aid in the nutrition of sperm and keep the sperm active. It does not produce any hormones so there is usually no alteration in sexual drive following prostatectomy.

PROSTATITIS

Prostatitis embraces a group of conditions with voiding discomfort and pain in the prostate referred to the perineum, low back, urethra and testes. It typically affects men aged 25 - 50 years. Prostatitis usually occurs in the absence of identifiable bacterial growth, when it is termed non-bacterial prostatitis. The prostate may develop acute or chronic bacterial infection.

Bacterial prostatitis is usually caused by *Escherichia coli* (commonest), *Streptococcus faecalis, Pseudomonas* or *Staphylococcus*. Some chronic infections have been shown to be associated with *Chlamydia trachomatis*.[1]

Prostatodynia means the presence of symptoms typical of prostatitis but without objective evidence of inflammation or infection (Table 107.1).

Table 107.1 Classification of prostatitis syndromes

	Prostatic pain	Prostatic rectal examination	Positive bladder culture	Positive prostatic secretion culture
Acute bacterial prostatitis	Yes	Very tender, swollen	Yes	Yes
Chronic bacterial prostatitis	Often	Normal or indurated	Occasionally	Low counts
Non-bacterial prostatitis	Often	Normal	Occasionally	No
Prostatodynia	Often	Normal	No	No

It is preferable to use the term ‘prostatitis syndromes’ to embrace the four terms used in Table 107.1.

Clinical features of acute bacterial prostatitis

Symptoms

- Fever, sweating, rigors
- Pain in perineum (mainly), back and suprapubic area
- Urinary frequency, urgency and dysuria
- Variable degrees of bladder outlet obstruction (BOO)
- ± Haematuria

DxT: dysuria + fever + perineal pain = acute prostatitis

Signs

- Fever
- Rectal examination: prostate exquisitely tender, swollen, firm, warm, indurated

Complications

- Abscess
- Recurrence
- Epididymo-orchitis
- Acute retention
- Bacteraemia/septicaemia

Chronic bacterial prostatitis

Chronic bacterial prostatitis is diagnosed by a history of mild irritative voiding with perineal, scrotal and suprapubic pain. Ejaculatory pain can occur. The gland may be normal on clinical examination or tender and boggy. It should be suspected in men with recurrent UTI (refer Table 107.2).

Table 107.2 Features of chronic bacterial prostatitis

Difficult to treat
Relapsing infection
Perineal pain
Some leucocytes in expressed prostatic secretion

Investigations

- Fractional urine specimens and expressed prostatic secretions (EPS) obtained after prostatic massage can show excess white cells.
- Culture of the urine or ejaculate may be negative or give low counts.
- Prostatic stones (demonstrated by plain X-ray or transrectal ultrasound) may prevent successful treatment.
- Prostatic specific antigen (PSA): elevation occurs with inflammation and may cause confusion with cancer.

Treatment

Acute bacterial prostatitis[2]

amoxycillin (or ampicillin) 2 g IV 6 hourly +

gentamicin 5 mg/kg/day as a single daily dose

until there is substantial improvement, when therapy may be changed to an appropriate oral agent, based on the sensitivity of the pathogen(s) isolated, for the remainder of 14 days[2]

For milder infection, oral treatment with amoxycillin/potassium clavulanate, trimethoprim or norfloxacin is suitable. Urinary retention or abscess formation almost always requires endoscopic deroofing for drainage.

Chronic bacterial prostatitis

Treatment of this condition is made difficult by uncertainty in differentiating it from non-bacterial prostatitis as cultures may grow low counts of what may be normal flora. Avoid overtreatment with antibiotics. Reassurance is important and it is worth suggesting frequent ejaculation and hot baths. Massage only for recalcitrant cases.

doxycycline 100 mg (o) daily for 1 month *or*

trimethoprim 300 mg (o) daily for 1 month *or*

norfloxacin 400 mg (o) 12 hourly for 1 month *or* ciprofloxacin 500 mg (o) 12 hourly for 1 month[2]

Non-bacterial prostatitis

This is the commonest by far. It is often recurrent and each episode can last several months. Antibiotic therapy is inappropriate. The symptoms may reflect retrograde passage of urine into prostatic tissue with urate crystallisation. Emphasise good voiding habits.[2] Avoid straining at the end of micturition. Encourage normal sexual activity and use stress management. Prescribe allopurinol 300 mg daily for at least 3 months for each episode as it reduces prostatic urate deposition and relieves symptoms.[3]

Prostatodynia

Perform a thorough genitourinary tract investigation. Some patients have urethral sphincter spasm and may respond to diazepam or the alpha-blocking agent minipress 0.5 mg bd. Psychological counselling may be appropriate. Occasionally alcohol and caffeine induce flares of prostatodynia, which may often be regarded as synonymous with non-bacterial prostatitis.

Very rare causes of prostatitis include tuberculosis, gonorrhoea, parasites and fungi.

LOWER URINARY TRACT SYMPTOMS (LUTS)

These symptoms can be grouped as voiding symptoms (obstructive) or storage symptoms (irritative). Irritative symptoms may be caused by a bladder problem only. Obstructive symptoms are usually caused by the prostate (which can also cause irritative symptoms). The old term ‘prostatism’ is ill defined and best dropped.

Voiding (obstructive) symptoms

- hesitancy
- weak stream
- postmicturition dribble
- incomplete emptying/urinary retention
- straining

Storage (irritative) symptoms

- urgency
- urge incontinence
- frequency
- nocturia
- dysuria
- suprapubic pain

BLADDER OUTLET OBSTRUCTION

Symptoms of BOO are present in most men after the age of 60 years. The commonest cause is benign prostatic obstruction. Benign prostatic hypertrophy is a histological diagnosis which, strictly, should not be used for symptoms. Only 10 - 15% require surgery for relief of obstructive symptoms.[1] Bladder outlet obstruction can also be caused by bladder neck obstruction and urethral sphincter spasm (Figure 107.1).


bladder
transitional zone of prostate
peripheral zone of prostate
verumontanum
compressed peripheral zone forming the capsule
hypertrophy of transitional zone
carcinoma developing in peripheral zone of prostate

1. Normal prostate

2. Benign prostatic hyperplasia

3. Carcinoma of prostate

4. Bladder neck obstruction

Figure 107.1 Diagrammatic comparison of bladder outlet obstruction

DxT: poor urine flow + straining to void + frequency = BOO

Clinical features of BOO

- Hesitancy
- Frequency of micturition
- Urgency
- Nocturia
- Slow interrupted flow
- Terminal dribbling
- Acute retention (the presenting problem in 15% of patients)
- Retention with overflow incontinence (less common)
- Haematuria from ruptured submucosal prostatic veins can occur
- Rectal examination usually detects an enlarged prostate

Note: Small prostate glands can also cause BOO.

Investigations

These include:

- urine culture
- kidney function
- PSA
- prostatic needle biopsy (guided by transrectal ultrasound) if carcinoma suspected
- voiding flow rate of < 10 - 15 mL/sec tends to confirm that the symptoms reflect obstruction and not bladder irritability (Figure 107.2)



Figure 107.2 A visual scale of urinary flow. The numbers signify the flow rate in millilitres per second. When assessing a patient with voiding dysfunction, ask him to indicate which stream is closest to his own. A flow rate below 15 mL/s suggests obstruction and below 10 mL/s significant obstruction

5

Complications of prostatic obstruction

- Retention
- Urinary infection
- Bladder calculus formation
- Uraemia

Self-help advice for patients with mild symptoms (after cancer excluded)

- Avoid certain drugs, especially OTC cough and cold preparations.
- Reduce caffeine.
- Avoid fluids before bedtime.
- Urinate when you need to (do not hang on).
- Wait 30 seconds after voiding and try to void again.

Medical treatment

Patients with mild symptoms may be helped with alpha-adrenergic blocking drugs such as phenoxybenzamine, tamsulosin, terazosin and prazosin to inhibit contraction of the muscle in the bladder neck and urethra.[1] Typical doses are tamsulosin 0.4 mg daily and prazosin 0.5 mg bd, or 1 mg nocte after commencing with 0.5 mg nocte. Prazosin can be increased to a maximum of 2 mg bd. Symptoms are not improved by increasing beyond this dose. The 5-alpha-reductase inhibitors (e.g. finasteride) reduce prostatic volume. Urine flow improves by 3 months, plateauing at 6 months, but not to the same degree as with surgery. They are not very effective for symptomatic control. Systematic reviews reveal that both alpha-adrenergic blockers and 5-alpha-reductase inhibitors reduce symptoms compared with placebo and that alpha-adrenergic blockers were more effective that 5-alpha-reductase inhibitors.[4] Both drugs are used long-term. They do not summate.

Surgical management

The most effective treatment for obstruction is transurethral resection of the prostate (TURP), or laser ablation (Figure 107.3) Transurethral incision of the prostate (TUIP) gives identical results with small glands. Open prostatectomy accounts for less than 1% of benign prostatic surgery today.



Figure 107.3 The process of prostatectomy

Permanent springs (stents) placed in the prostatic urethra under local anaesthesia are an option for the very frail patient. Microwave and heat treatment are new methods but less effective.

Absolute indications for prostatectomy include deterioration in kidney function, the development of upper tract dilatation, retention (following drainage and assessment) and bladder stones. Eighty per cent of patients have surgery for bothersome symptoms.

Postoperative guidelines for the patient

- There may be urgency even to the point of incontinence for a few days.
- Bleeding can occur intermittently for 3 weeks, so increase fluid intake.
- Erectile function is usually unchanged (neither better nor worse) but loss of erection probably occurs in 5% of patients.
- Avoid intercourse for 3 weeks.
- Orgasms continue but there is usually no emission with ejaculation. The semen is ejaculated back into the bladder.
- If obstructive problems recur early there may be a stricture.

Note:

- Persisting postoperative frequency bothers about 15% of patients.
- 10 - 15% reveal unsuspected carcinoma.

Drugs and LUTS

Certain drugs can cause LUTS due to the effect of the drug on the bladder. It is very important for the family doctor to enquire about these drugs when evaluating such a patient. The problem is mainly an adverse effect of drugs with anticholinergic activity.

Anticholinergics

- Atropine and hyoscine compounds, e.g.
  - isopropamide
  - mazindol
  - phenothiazines
  - dicyclomine
  - propantheline
  - other belladonna alkaloids
- Antidepressants
  - especially tricyclic compounds
- Anti-parkinson agents, e.g.
  - amantadine

  - benzhexol
  - benztropine
  - biperiden
  - orphenadrine
  - procyclidine

Beta-adrenoceptor agonists

- Ephedrine
- Salbutamol
- Terbutaline
- OTC preparations (mainly for coughs and colds - e.g. sympathomimetics including ephedrine)

CARCINOMA OF THE PROSTATE

Prostate carcinoma is the commonest maligancy in men and the third commonest cause of death from malignant disease in Australia. It is rare before the age of 50 years. By the age of 80 years 80% of men have histologic carcinoma within the gland but most are dormant.[6] The lifetime risk of the diagnosis of prostate cancer is 1 in 12. The risk increases with age and is greater for those with a close relative diagnosed before 70 years. Prostatic carcinoma may be asymptomatic, even when it has extended beyond the prostate. It usually commences in the peripheral part of the gland. There are dramatic racial differences in the frequency that tend to change with migration, indicating that prostate cancer reflects environmental influences (and possibly dietary fat).

Clinical features

Unsuspected carcinomas are often detected by the tumour marker PSA or histologically after TURP. Clinical prostatic carcinoma presents typically with rapidly progressive symptoms of lower urinary tract obstruction or of metastatic spread, especially to bone (pelvis and vertebrae).[4] Symptoms include BOO 70%, acute retention 25%, back pain 15%, haematuria 5% and uraemia 5%.[1] Other symptoms include tiredness, weight loss and perineal pain.

Digital rectal examination (DRE) may reveal a nodule (50% are not carcinoma). Locally advanced cancer typically reveals a hard, nodular and irregular gland. The tumour may be large enough to obliterate the median sulcus. The borders may lack definition. On the other hand, with cancer, the prostate may feel normal.

Signs of abnormal prostate (DRE):

- hard lump
- asymmetry
- induration
- loss of median sulcus

Investigations to detect carcinoma

Blood analysis

- PSA:
  - can be elevated without cancer
  - must be tested with DRE
  - is prostate specific, not prostate cancer specific
  - can be ‘normal’ (< 4 ng/mL) in 15 - 25% of cancers7
  - levels > 10 ng/mL are strongly suggestive of cancer
  - levels > 20 ng/mL suggest metastatic disease
- Age-specific PSA reference range:

  - 40 - 49 years 0 - 2.5 ng/mL
  - 50 - 59 years 0 - 3.5 ng/mL
  - 60 - 69 years 0 - 4.5 ng/mL
  - 70 - 79 years 0 - 6.5 ng/mL
- Free/total PSA ratio: Refinements to the PSA test are the ‘free/total PSA ratio’ and the rate of rise of PSA over time. The specificity and sensitivity limitations of the PSA marker are well known, especially in the 4 - 20 ng/mL ‘grey zone’ where there is significant overlap between BPH and prostate cancer. The F/T PSA ratio helps to discriminate between these in patients with moderately raised PSA levels.

Biopsy[7]

Consider biopsy guided by transrectal ultrasound if the DRE is positive or if the PSA is elevated.

Investigations to stage disease

- Radionuclide bone scan

Screening

The value of screening all asymptomatic men is debatable. If it has a place it would consist of a PSA test and DRE annually from 50 - 70 years if life expectancy exceeds 10 years.[7] As a screening procedure DRE, PSA or transabdominal ultrasound is not currently recommended by the RACGP and other authorities.[8] Results of definitive studies, due for completion in 2008, are awaited.

Treatment of prostate cancer[7]

Patients over 70 years with no symptoms may be observed (‘watchful waiting’) with regular DRE and PSA measurement. The treatment depends on the age of the patient and the stage of the disease. Repeating the PSA in 3 months is a useful strategy.

For tumours that are potentially curable, radical prostatectomy or local radiotherapy are the options. Metastatic carcinoma is currently not curable.

External beam radiotherapy

The cure rate is 10% less than with surgery but the difference does not become apparent for 10 years. The main adverse reactions are faecal urgency and diarrhoea together with urine frequency. Impotence is common 2 years after radiotherapy.

Brachytherapy

In this, a form of radiotherapy, the X-rays come from tiny radioactive seeds inserted directly into the tumour.

Radical prostatectomy

- Usually recommended for patients under 70 years with a PSA < 20 ng/mL.
- Long-term cure rates of 80% must be balanced with urine incontinence rates (sufficient to need a pad) of about 10% and frequently impotence (> 70%).

If the PSA is < 0.1 ng/mL at the end of 2 years the patient can be considered cured.

Hormone manipulation

For metastatic or locally advanced disease, androgen deprivation is the cornerstone of treatment, the options being:

- bilateral orchidectomy *or*
- daily antiandrogenic tablets e.g.
  - cyproterone acetate (Androcur)
  - flutamide (Eulexin)

  - bicalutamide (Cosudex) *or*
- luteinising hormone releasing hormone (LHRH) agonists: depot injections of LHRH analogues e.g.
  - goserelin (Zoladex)
  - leuprorelin acetate (Lucrin, Eligard)

Treatment combinations for small volume metastatic prostate carcinoma *may* prolong life e.g.:

- orchidectomy plus flutamide
- LHRH agonists plus flutamide or bicalutamide - LHRH agonists cause an initial surge of testosterone so a preliminary antiandrogenic agent is advised to prevent a flare in the cancer.

Complementary medicine

Among the disorders most widely promoted as benefiting from natural remedies are those of the prostate. Herbal remedies that have been widely used include saw palmetto (*Seronea repeus*), stinging nettle (*Urtica dioica*), pygeum Africanum and cernilton-r.

Considerable media attention has been given in Europe and Australia to the benefits of willowherb (*Epilobium*) but no trials have been conducted to test efficacy.

Saw palmetto has been used widely, especially in Germany, with studies confirming efficacy and safety in benign prostatic hypertrophy when compared with finasteride4 but not as efficacious as alpha-1-antagonists.

Benign hyperplasia has been treated with isoflavone phytoestrogens. A weak oestrogen agonist effect in males may antagonise the growth-promoting effects of androgens on the prostate. Epidemiological data indicates that in countries such as Japan where isoflavone diets are prevalent, prostatic enlargement occurs less with ageing. The most widely used source of phytoestrogen is soy protein. It is also present in lentils, chick peas, some variations of beans and alfalfa sprouts.

There is evidence from some epidemiological studies that selenium and leucopenes (from tomato and tomato products) in the diet protects against carcinoma of the prostate.[9]

REFERENCES

1. Bullock N, Sibley G, Whitaker Q. *Essential Urology*. Edinburgh: Churchill Livingstone, 1989: 287 - 99.
2. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therapeutic Guidelines Ltd, 2006: 104 - 5.
3. Persson BE, Ronquist G, Ekblom M. Ameliorative effect of allopurinol on nonbacterial prostatitis: a parallel doubleblind controlled study. J Urol, 1996; 155: 961 - 4.
4. Barton S ed. *Clinical Evidence*. London: BMJ Publishing Group, 2001: 588 - 98.
5. Millard R. Benign prostatic hyperplasia: recent advances in treatment. Mod Med Supplement, 1998; 41(7); 7 - 8.
6. Kumar PJ, Clark ML. *Clinical Medicine* (5th edn). London: Saunders, 2003: 663 - 5.
7. Stricker P, Phelps K. *PSA for the General Practitioner*. Sydney: Prostate Cancer Foundation of Australia, 2004: 1 - 24.
8. RACGP. *Guidelines for Preventative Activities in General Practice* (6th edn). Melbourne: RACGP, 2005: 56.
9. Yoshizawa K et al. Study of prediagnostic selenium level in toenails and the risk of advanced prostatic cancer. J Natl Cancer Inst, 1998; 90: 1219 - 24.

CHAPTER 108. THE SUBFERTILE COUPLE

You notice that the tabetic has the power of holding water for an indefinite period. He is also impotent - in fact, two excellent properties to possess for a quiet day on the river.

DR DUNLOP 1913, *Teaching at Charing Cross Hospital*

THE SUBFERTILE COUPLE

Infertility is defined as the absence of conception after a period of 12 months of normal unprotected sexual intercourse.[1] Interestingly, normal fertility is defined as 'achieving a pregnancy within 2 years of regular sexual intercourse' and applies to about 90% of couples.[2] Infertility can be a very distressing and emotional problem for the couple, who need considerable care, empathy and relatively rapid investigation of their problem. In assessing a couple with the problem of subfertility (this term is a preferable way of describing the condition to the patients), it is appropriate to involve both partners in the consultation. In determining the cause of the subfertility, three basic fertility parameters should be investigated.[3]

- The right number of sperm have to be placed in the right place at the right time.
- The woman must be ovulating.
- The tubes must be patent and the pelvis sufficiently healthy to enable fertilisation and implantation.

KEY FACTS AND CHECKPOINTS

- Infertility affects 10 - 15% of all cohabitating couples.[4]
- This incidence increases with age.
- After the age of 32 fertility decreases by 1.5% per year.
- About 15% of couples who do not use contraception fail to achieve a pregnancy within 12 months.[5]
- More than 10% remain unsuccessful after 2 years.[5]
- About 50% of couples will seek medical assistance.[4]
- The main factors to be assessed are ovulation, tubal patency and semen analysis.
- About 40 - 50% of couples have an identifiable male factor. Male infertility affects 5% of men.
- The technique of intracytoplasmic sperm injection (ICSI) has revolutionised the treatment of male infertility.
- Female factors account for about 45%: tubal problems account for about 20% and ovulatory disorders about 20%. Polycystic ovary syndrome (PCOS) is the most common cause of ovulatory dysfunction.[6]
- The initial investigation for the man is semen analysis on two occasions. The initial investigation for the woman is a basal body temperature chart followed by midluteal progesterone measurement.
- About 15% of cases have no apparent explanation.
- A significant number (25%) have combined male and female problems.
- Current specialised treatment helps the majority of subfertile couples to achieve pregnancy.[7]

PHYSIOLOGICAL FACTORS[7, 8]

Male fertility

Fertility in the male requires:

- normal hypothalamic function producing gonadotrophin-releasing hormone (GnRH)
- normal pituitary function producing the gonadotrophic hormones follicle stimulating hormone (FSH) and luteinising hormone (LH)
- normal seminiferous tubule and Leydig cell function
- normal sperm transport and delivery

Facts about sperm viability

- The maximum number of viable sperm is found in the ejaculate after a 48 hour abstinence.
- After entering receptive cervical mucus, sperm are capable of fertilising an egg for at least 48 hours.
- Sperm survive for less than 30 minutes in the vagina.

Female fertility

Fertility in the female requires:

- normal function of the ovulatory cycle, which requires:
    - normal hypothalamic - pituitary function producing the hormones GnRH, FSH and LH
    - normal ovarian function with follicular response to FSH and LH (Figure 108.1)



Figure 108.1 The normal ovulatory cycle: the midcycle peak of LH and FSH is at 14 days and ovulation occurs shortly afterwards

-
    - appropriate prolactin levels (which are normally low); excessive prolactin secretion (hyperprolactinaemia) causes anovulation
- normal tubal transport and access of the ovum to incoming sperm
- receptive cervical mucus
- normal uterus to permit implantation of the fertilised ovum

Probabilities of pregnancy

About 50% of normal couples, having unprotected intercourse at least twice a week, will probably achieve pregnancy in 6 months, 80% in 1 year and 90% in 2 years.[7]

CAUSES OF INFERTILITY

Significant causes of infertility are summarised in Table 108.1 and illustrated in Figure 108.2.

Table 108.1 Significant causes of infertility

Female factors
Ovulation disorders
• ovarian failure (e.g. oocyte ageing)
• stress
• PCOS
• weight-related ovulation disorders
• hyperprolactinaemia
• other endocrine disorders

• idiopathic eugonadotropic anovulation Tubal disease • PID • endometriosis • previous ectopic pregnancy • previous tubal ligation • previous peritonitis Uterine and cervical abnormalities • congenital • acquired Endometriosis
Male factors Reduced sperm production • congenital cryptorchidism (maldescent) • inflammation (e.g. mumps orchitis) • antispermatogenic agents - chemotherapy - drugs - irradiation - heat • idiopathic • Klinefelter's syndrome (46 XXY) Hypothalamic pituitary disease • hypogonadotropic disorder Disorders of coitus • erectile dysfunction • psychosexual ejaculatory failure • retrograde ejaculation - genitourinary surgery - autonomic disorders (e.g. diabetes) - congenital abnormalities Ductal obstruction
Couple factors Joint subfertility Psychosexual dysfunction


GnRH
hypothalamus
anterior pituitary
FSH
LH
prolactin
FSH
LH
FSH
LH
excess prolactin inhibits ovulation
ovum
oestrogens progesterone
testosterone
ovulation
tubal disorders
uterine disorders
cervical-vaginal factors
potency-sperm dysfunction
vas blockage
testicular dysfunction
? adequate coitus
Female factors
Male factors
Combined factors

X indicates possible disorder causing infertility

Figure 108.2 The major factors involved in subfertility. Adapted from Kumar and Clarke.

A DIAGNOSTIC APPROACH

It is important to see both partners, not just the woman.

History

The following basic facts should be ascertained.

The man

- Sexual function
- Previous testicular problems/injury (e.g. orchitis, trauma, undescended testes)
- Medical problems: diabetes, epilepsy, tuberculosis, kidney disorders
- Past history (PH) of STIs
- PH of mumps
- PH of urethral problems
- Genitourinary surgery (e.g. hernia)
- Recent severe febrile illness
- Occupational history (exposure to heat, pesticides, herbicides)
- Drug intake (possible adverse effects from)
    - alcohol
    - chemotherapy
    - anabolic steroids
    - aminoglycoside antibiotics
    - sulphasalazine
    - cimetidine/ranitidine
    - colchicine
    - spironolactone
    - antihypertensive agents
    - narcotics

  - phenytoin
  - nitrofurantoin
  - nicotine
  - marijuana

The woman

- Evidence of previous fertility
- Onset of menarche
- Menstrual history
- Symptoms of ovulation
- Symptoms of endometriosis
- PH of STIs and pelvic infection
- Previous IUCD
- PH of intra-abdominal surgery (e.g. appendicitis, ovarian cyst)
- PH of genitourinary surgery, including abortions
- Obstetric history
- Body weight: eating disorders (anorexia, obesity)
- Drug intake
  - alcohol
  - smoking, especially > 20/day
  - oral contraception
  - anabolic steroids
  - major tranquillisers

Combined history

- Frequency and timing of intercourse
- Adequate penetration with intercourse, and ejaculation
- Use of lubricants
- Attitudes to pregnancy and subfertility
- Expectations for the future

Physical examination

A general assessment of body habitus, general health, including diabetes mellitus, and secondary sexual characteristics should be noted in both man and woman. Urinalysis should be performed on both partners.

The man

- Secondary sexual characteristics; note any gynaecomastia
- Genitalia
  - size and consistency of the testes
    - normal size 3.5 - 5.5 cm long; 2 - 3.5 cm wide
    - small testes < 3.5 cm long
    - Klinefelter's: 2 (or less) cm long (typical)
  - palpate epididymis and vas (present and non-tender is normal)
  - evidence of varicocele (controversial role)
  - PR: check prostate
  - note penis and location of urethra (always retract the foreskin for examination)

The woman

- Secondary sexual characteristics
- Thyroid status
- Genitalia and breasts
- Vaginal and pelvic examination
    - assess uterus and ovaries (normal - present, mobile and non-tender)
    - the adnexae (any masses)

Investigations

These are usually performed after referral but the family doctor should organise initial investigations to assess where to refer (e.g. andrologist, endocrinologist, gynaecologist).

Initial investigations

Male - semen analysis

It is advisable to obtain two samples at least 80 - 90 days apart as the cycle of production of sperm is about 80 days. It requires a complete ejaculation, preferably by masturbation, after at least 3 days sexual abstinence. Use a clean, dry widemouthed bottle; condoms should not be used. Semen should be kept warm and examined within 1 hour of collection.[2, 8]

- Normal values:
    - volume > 2 mL (average 2 - 6 mL)
    - concentration > 20 million sperm/mL
    - motility > 40% after 2 hours
    - normal forms > 20%
    - velocity > 30 microns/s

Female - ovulation status

- Educate about temperature chart and cervical mucus diary, noting time of intercourse (take temperature with thermometer under tongue before getting out of bed in the morning). Now considered to be of low value.
- Midluteal hormone assessment (21st day of cycle) - i.e. serum progesterone. This is the most commonly used first line test for ovulation (refer to Table 108.2).

Table 108.2 WHO classification of ovulatory dysfunction[6]

Group 1	Hypothalamic - pituitary failure	Low FSH, LH, e.g. anorexia nervosa
Group 2	Hypothalamic - pituitary dysfunction	Normal FSH, e.g. PCOS
Group 3	Ovarian failure	High FSH, e.g. ovarian failure

Subsequent investigations

Diagnostic laparoscopy - direct visualisation of corpus luteum, tubes; check tubal patency by insulfating blue dye from the cervix through the tubes to the peritoneal cavity.

Further investigations (if necessary)

Male

If azoospermia or severe oligospermia:

- serum FSH level (if 2.5 times normal, indicates irreversible testicular failure) - this is the most important endocrine test in the assessment of male infertility
- LH and inhibin (a low inhibin also indicates irreversible failure)
- antisperm antibodies (in semen or serum)

- sperm function tests
- chromosome analysis: 46 XXY or 46 XXY/46 XY

Female

(Other investigations may be necessary.)

- Thyroid function tests: ? hypothyroidism
- Serum prolactin, FSH, LH, androgen levels
- Sonohysterogram
- Endometrial biopsy
- Transvaginal ultrasound
- Hysteroscopy/laparoscopy
- CT of the pituitary fossa
- Chlamydia (cervical culture)

Note: Ovulation or its absence is best demonstrated by luteal progesterone.

Essential investigations are outlined in Table 108.3.

Table 108.3 Essential investigations of the subfertile couple[2]

First-line
Basal body temperature chart and cervical mucus diary
Semen analysis
Serum progesterone (mid-luteal - day 21) in female
Transvaginal ultrasound
Rubella immune status (female)

Management principles[4]

- Both partners should be involved in management decisions since fertility is a couple's problem.
- Infertility can cause considerable emotional stress, including the taking or placing of blame by one partner or the other, and subsequent guilt feelings; hence sensitive and empathetic support is essential. This may include marital counselling.
- Since recent advances have helped this problem so much, there is no place for guesswork or for empirical therapy and early referral is necessary.

POLYCYSTIC OVARY SYNDROME

PCOS is a common chronic anovulatory disorder affecting 5 - 10% of women of reproductive age. It should not be confused with the common ultrasonographic diagnosis of cystic ovaries with no clinical features. The classic clinical features are infertility, oligomenorrhoea, hirsutism (70%) and obesity (50%).

PCOS has a strong hereditary basis and can begin in the pubertal years. There are variations of the syndrome, including normal menstruation (20%) and abnormal uterine bleeding (50%).

General features of PCOS[9]

Patients have some of the following (rare to have all):

- ovarian dysfunction - infertility, oligomenorrhoea, anovulation, tendency to miscarry, endometrial cancer
- androgen excess - hirsutism and acne
- obesity

- metabolic syndrome - insulin resistance, impaired glucose tolerance, hyperlipidaemia, atherosclerosis, hypertension, tendency towards type 2 diabetes
- ovarian abnormalities - multiple small follicles, stromal expansion

Aetiology

The exact cause is unknown. There is a hereditary factor and hormonal dysfunction which cause more egg-containing cysts to form in the ovaries without ova reaching maturation. Some experts claim that the principal underlying disorder is insulin resistance with hyper-insulinaemia.

Diagnosis/investigations

- Raised LH level with normal FSH levels (LH/FSH ratio > 2)
- Raised serum testosterone level
- Transvaginal ultrasound (usually 10 - 20 follicles < 8 mm)
- Possible endometrial biopsy

Suggested screening for all women with PCOS[9]

- Smoking history
- Blood pressure
- Blood glucose levels
- Oral glucose tolerance test
- Fasting plasma insulin
- Fasting serum HDL and LDL levels

Management strategies[9]

First-line

- Weight reduction and exercise (the key - this alone may restore normal ovarian function)
- PCOS support group

Potential

- Screening and treat if necessary:
  - glucose intolerance and diabetes mellitus
  - hyperlipidaemia
  - hypertension
- Primary treatment of insulin resistance
  - metformin
  - thiazolidenediones (glitazones)
- Ovulation induction - clomiphene/gonadotrophins
- Assisted conception
- Laparoscopic ovarian diathermy (failed medical treatment)

COUNSELLING THE SUBFERTILE COUPLE[10]

The counselling of subfertile couples has to be adapted to the level reached by the couple along the infertility pathway. The needs of each couple may be very different depending on their emotional nature, their lifestyle, moral, religious and ethical beliefs. However, their suffering can run very deep and deserves attention, time and opportunities for free expression of feelings and concerns.

The medical counselling model developed by Colagiuri and Craig[10] (Figure 5.1 at p. 34) is very useful as it empowers patients to make their own decision through facilitation as opposed to the directive and advisory medical model.

The couple are provided initially with accurate and appropriate information. Anxiety is alleviated by reassurance and by dispelling myths such as their problem being caused by an unfavourable position for intercourse, leakage of excess semen from the vagina or previous use of the pill.

The facilitation process enables the couple to ventilate any feelings of guilt, anxiety, fear, anger and sexuality. The style of questioning should aim to explore the influence that the problem has had on the couple and then the influence they have over it.

These processes then lead to decision making by the couple about further management strategies. A graph of emotional responses to the infertility (Figure 108.3) can be used to help the couple explore their current and past emotional responses to their problem. Apart from helping them realise that their problem is not unique, it provides opportunities for ventilation of important feelings that can act as a basis for counselling.



Figure 108.3 Emotional responses to infertility. Colagiuri and Craig[7] reprinted with permission.

TREATMENT

If the problem has been identified, specific treatment needs to be prescribed by the consultant.

- Anovulation can be treated with ovulation induction drugs such as clomiphene, bromocriptine, gonadotrophins or GnRH.
- Endometriosis can be treated medically or surgically (peritubal adhesions).
- Male problems - little can be done (including testosterone and vitamins) to enhance semen quality. Corticosteroids may help if sperm antibodies are present. Consider invitro fertilisation (IVF) and related technology for male factor infertility, particularly intracytoplasmic sperm injection (ICSI) into the occyte.
- Artificial insemination.
- Donor insemination.
- Severe tubal disease - use IVF and embryo transfer (IVF-ET).
- Unexplained subfertility - consider gamete intrafallopian transfer (GIFT), a modification of IVF. This method, in which eggs and sperm are placed into the Fallopian tubes, is best used in the treatment of infertility of unknown aetiology and carries a pregnancy rate of about 30% per couple.[3]

Evidence-based RCTs

The conclusion as published in Clinical Evidence[11] has summarised known benefits of infertility treatment (to date) as follows:

- ovarian disorders - clomiphene, laparoscopic ovarian ‘drilling’

- tubal infertility - IVF
- endometriosis - intrauterine insemination with ovarian stimulation
- male infertility - intrauterine insemination, donor insemination, Fallopian tube sperm infusion
- unexplained infertility - intrauterine insemination

WHEN TO REFER

A family doctor should perform the initial investigations of a couple with infertility, including temperature chart, semen analysis and hormone levels, to determine whether it is a male or female problem and then organise the appropriate referral.

REFERENCES

1. Kumar PJ, Clark ML. Clinical Medicine (2nd edn). London: Bailliere Tindall, 1990: 792 - 4.
2. Stern K. How to treat: Infertility. Aust Doctor 2002; 12 July: I - VIII.
3. Wood C, Kovacs G. Infertility: All Your Questions Answered. Melbourne: Hill of Content, 1996.
4. Jequier AM. Infertility. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 273 - 8.
5. Smith H. Female infertility. Mod Med Aust, 1990; August: 98 - 106.
6. Illingworth P, Lahoud R. Investigation of the infertile couple 1. Medical Observer, 2006; 21 April: 27 - 30.
7. Knowlden S, Helman T. How to treat: Infertility. Aust Dr Weekly, 1989; 25 August: I - VII.
8. DeKrester D. Female infertility. Mod Med Aust, 1990; July: 98 - 109.
9. Norman RJ et al. Metformin and intervention in polycystic ovary syndrome. Med J Aust, 2001; 174: 580 - 3.
10. Craig S. A medical model for infertility counselling. Aust Fam Physician, 1990; 19: 491 - 500.
11. Duckitt K. Infertility and subfertility. In: Barton S ed. Clinical Evidence. London: BMJ Publishing Group, 2001, issue 5: 1279 - 94.

CHAPTER 109. SEXUAL HEALTH

The functional form of impotence fills the coffers of the quacks, and swells the list of suicides.

RUTHERFORD MORRISON (1853 - 1939)

SEXUAL HEALTH

Family doctors are often asked to provide advice and help for sexual concerns and are continually challenged to detect such problems presenting in some other guise. Since we deal with so much illness, including debilitating problems, and prescribe so many drugs we must be aware and sensitive to the possible implications of their various effects on sexual health.

Sexual disorders can be considered in three major groups: sexual dysfunction, sexual deviation, and gender role disorders. This chapter largely confines itself to a discussion of sexual dysfunction.

SEXUAL DYSFUNCTION

Sexual dysfunction in men refers to persistent inability to achieve normal sexual intercourse while in women it refers to a persistent lack of sexual satisfaction.[1]

Several studies have demonstrated that sexual concerns and problems are common with a prevalence ranging from 10 - 70% of the population.[2] Difficult problems are summarised in Table 109.1. These studies have also indicated that patients are certainly willing to discuss their sexuality and wish their family doctors to become involved in counselling and management of their problems. Between 25% and 30% of sexual difficulties have an organic cause, while the remainder are emotional or psychogenic in origin.[3] The unique place of general practice and the family doctor provides ideal opportunities to address the sexual concerns of patients as the family doctor often has considerable insight into the family dynamics and first-hand perspective of the individuals involved.

Table 109.1 Sexual dysfunction: difficult problems

Sexual desire
• low libido
Sexual arousal
• erectile impotence
• failure of arousal in women
Sexual orientation/activity
• homosexuality
• fetishism
Orgasm
• premature ejaculation
• retarded ejaculation
• orgasmic dysfunction in women
Male problems
• low libido
• erectile difficulties
• premature ejaculation
• failure to ejaculate, or retarded ejaculation

Female problems

- low libido
- failure of arousal
- vaginismus
- orgasmic difficulties
- dyspareunia

The most common problem influencing an effective outcome is difficulty in communication between doctor and patient, which prejudices effective history taking and counselling. The problem is not contentrelated, much of which is based on commonsense, but the ubiquitous problem of communication.

If, as a practitioner, you counsel on the assumption that astounding ignorance about sexuality still exists in our society, you will be amazed at the results and at how relatively simple it is to help so many confused people who often have unrealistic expectations of their partners and themselves.

OPPORTUNISTIC SEXUALITY EDUCATION

The family doctor has many opportunities to provide education in sexuality throughout the lifelong care of the patient and it is wise to have a strategy that matter of factly incorporates enquiries and information about sexual health.

Examples include:

- antenatal and postnatal care
- contraceptive requests
- parents concerned about their children's sex play
- serious illness - medical and surgical
- adolescent problems
- menopause problems

PRESENTATION OF SEXUAL CONCERNS

Although some patients may present directly with a complaint of sexual dysfunction, many will be less direct and use some other pretext or complaint as a ‘ticket of entry’ for their sexual concerns (Table 109.2). Despite a seemingly terse approach the issue must be recognised and treated with considerable importance. This may mean scheduling an appropriate time to discuss the concerns.

Table 109.2 How sexual issues may present in family practice[2]

Minor non-sexual complaint - ‘entry ticket’
Specific sexual concern
Marital or relationship problem
Non-sexual problem (as perceived by the patient)
Sexual enquiry and counselling as part of illness management
Sexual enquiry as part of total health check-up
Infertility
Menopausal problems

Sometimes patients are unaware of an association between their medical problem and underlying sexual issues.[2]

Doctors may recognise such an association and initiate a tactful psychosocial history that includes questions about sexuality. Examples are chronic backache, pelvic pain, vaginal discharge, tiredness, insomnia and tension headache.

THE EFFECT OF ILLNESS ON SEXUAL FUNCTION

Doctors seldom enquire about the impact of an illness on the sexual function of patients and their partners and tend to be unaware of the sexual needs of elderly people (Table 109.3). It is most appropriate to enquire about these issues in our patients - e.g. the postmyocardial infarction patient, the postprostatectomy patient, the patient taking antihypertensives or other drugs (Table 109.4), and the post-mastectomy or posthysterectomy patient. Diabetes deserves special attention as 27 - 55% of diabetic men reported some erectile difficulties.[2]

Table 109.3 Medical conditions affecting sexual performance

Cardiovascular
• previous myocardial infarction
• angina pectoris
• peripheral vascular disease
• hypertension and its treatment
Respiratory
• asthma
• COPD
Endocrine
• diabetes mellitus
• hypothyroidism
• hyperthyroidism
• Cushing's syndrome
Neurological
• multiple sclerosis
• neuropathy
• spinal cord lesions
• Parkinson's disease
Musculoskeletal
• arthritis
Depression
Kidney
• kidney failure
Urological problems
• prostatectomy
• phimosis
• Peyronie's disorder
• priapism
Hepatobiliary
• cirrhosis

Surgical
• vaginal repair
• hysterectomy
• others
Trauma
• motor vehicle accidents
Cancer
Other
• Klinefelter's syndrome

Table 109.4 Drugs affecting sexual arousal and function

Male	Female
Alcohol	Alcohol
Narcotics	Narcotics
Marijuana	Marijuana
Anticholinergics	Combined oral contraceptive
Antiepileptics	CNS depressants
Antihypertensives	Antihypertensives (selected)
Antihistamines	Antiepileptics
Benzodiazepines	
Disulfiram	
Psychotherapeutic drugs	
Oestrogens	
Cytotoxic drugs	

TAKING A SEXUAL HISTORY

It is important to be alert for psychiatric disorders and situational factors and not to predict a person's sexual disposition. Avoid being too formal or too familiar but aim to display a wise, matter-of-fact, empathic, commonsense rapport. Tactfully explore the patient's attitude to sexuality and examine the relationship. Ideally, it is best to see a couple together if the problem is occurring within a steady relationship. As a practitioner, you have to be comfortable with your own sexuality and learn to be relaxed, confident and understanding when dealing with sexual concerns.

Enquiry about possible child sexual abuse is an important part of the history.

Probing questions for a suspected sexual problem

- Do you have any trouble passing urine or any vaginal discharge (women)?
- Are you sexually active?
- What is the physical side of your marriage/relationship like?
- Do you have any pain or discomfort during intercourse?
- Is your relationship good?
- Do you communicate well? Generally? Sexually?

- Do you have any difficulties in your sexual relationships?
- What is your sexual preference?
- Are you attracted to men, women or both?
- Have you experienced the ‘coming out’ process?
- What drugs are you taking?
- Do you take recreational drugs (e.g. alcohol, marijuana, nicotine)?

Specific questions about sexuality

- Do you get aroused/turned on? What turns you on?
- Do you look forward to making love?
- Do you spend much time on love play?
- Does lovemaking make you feel happy and relaxed?
- Do you worry about getting pregnant (women)?
- What do you do about contraception?
- Do you worry about getting an STI?
- Do you worry about getting AIDS?
- How often do you reach a climax during lovemaking?
- How often do you have intercourse, or sexual activity without intercourse?
- Do you ‘come’ together?

Female

- Do you have enough lubrication? Are you wet enough?
- Do you find intercourse uncomfortable or painful?

Male

- Do you have trouble getting a full erection?
- How long does it take you to ‘come’ after you insert your penis?
- Do you ‘come’ too quickly?

Background history for an admitted problem

- Can you think of any reasons why you have this problem?
- What sex education did you have as a child? At home or at school?
- Were your parents happily married?
- Were sexual matters something that could be discussed in the home?
- Did you come from a religious family?
- Did you receive warnings or prohibitions as a child?
- What was the family attitude to masturbation, extramarital sex, menstruation, contraception, etc.?
- What is your attitude to masturbation?
- Were you fondled or sexually abused by an adult, especially a member of the family?
- Were there healthy shows of affection such as touching or hugging between family members?
- Did you have any upsetting sexual experiences during childhood and adolescence?
- What was your first sexual experience like?

Examination

The routine medical examination should include the basics such as urinalysis, BP measurement, genital examination and neurological where indicated. A careful vaginal and pelvic examination should be an opportune educational experience for the patient and an exercise in preventive medicine.

Investigations

No particular routine tests are recommended. Tests for male impotence are outlined on p.1120. Tests that may help exclude significant causes of low libido are those for diabetes, liver dysfunction, thyroid dysfunction and endocrine dysfunction. Endocrine dysfunction tests include prolactin, free testosterone, FSH, LH and oestradiol estimations. Other investigations may include pelvic ultrasonography, colposcopy or laparoscopy.

EXPLORING SEXUAL MYTHS

The acceptance in part or in total of many sexual myths that have prevailed in our society may have affected the relationship of a couple, especially in the context of the modern trend towards openness in discussing sexuality. It is worthwhile to help patients identify whether any of these myths have influenced their concerns by exploring common myths and their significant consequences to the individual or couple.

Sexual myths that could be explored include:[2]

- men need sex, women need love
- men need more sex than women
- men must be the instigators
- men know all about it
- sex = intercourse
- in this enlightened age everyone understands sexual issues

Sexual myths in the male[4]

- A hard erection is essential for good sex.
- A man should not show his feelings.
- A real man is always horny and ready for sex.
- As a person gets older there is no change in sexual interest, response or performance.
- As a person gets older there is a loss of interest in sex.
- Sexual performance is what really counts.
- Men are responsible for their partner's sexual pleasure.
- Sex must lead to orgasm.
- A man and his partner must reach orgasm simultaneously.

BASIC SEXUAL COUNSELLING

The family doctor can learn to be an effective sex counsellor. Sex counselling can be emotionally demanding and, while good interviewing skills, interest, support and basic advice are important, additional skills are needed to be an effective counsellor.

The fundamental methods involve:

- good communication and allowing a ‘comfortable’ exchange of information
- giving the patient ‘permission’ to talk openly about sexual matters
- providing basic ‘facts of life’ information
- dispelling sexual myths, correcting other misunderstandings
- gentle guidance for appropriate insight
- de-emphasising the modern-day obsession with performance and orgasm and emphasising the value of alternative forms of sexual expression (e.g. caressing, kissing, and manual and oral stimulation)
- reducing the patient's anxiety
- bolstering self-images affected by feelings of rejection, avoidance, guilt, resentment or incompetence

- reassuring the patient that he or she is normal (where appropriate)

Inappropriate doctor behaviour is presented in Table 109.5.

Table 109.5 Sexual counselling: inappropriate doctor behaviour

Overfamiliarity
Being too formal
Being too talkative
Blunt questioning
Being judgmental
Making assumptions about the other's sexuality
Imposing one's own beliefs and standards
Dogmatism
Tackling problems beyond one's experience

An interesting realisation after counselling families in sexuality is that most of the problems are not difficult and often spring from basic ignorance of normal sexual function; it's simply a matter of setting the record straight. The greatest hurdle is ‘getting started’ with delineating the problem. Once that barrier is crossed, satisfactory results appear to follow.

Another significant realisation is that sexual problems can be grossly underestimated. Human beings generally have a basic craving for intimacy, touching, stroking and loving sex. Apparently ‘good’ harmonious relationships can lack this type of intimacy, which may lead to various psychosomatic manifestations.

Ideally the family doctor should undertake a course in sexual counselling to promote confidence in the counselling process. Patients can be taught basic methods (where appropriate) such as sensate focus, squeeze or stop - start techniques for premature ejaculation, self-exploration using Kegel's exercises, fantasy conditioning with videos, and behaviour modification. Complex problems, especially those involving impotence, infertility and sexual deviations or perversions, demand referral to an expert.

THE PLISSIT COUNSELLING MODEL

The PLISSIT counselling model developed by Annon[5] can be used to build the skills needed to deal with sexual problems, especially if there is a psychological element.[3]

The mnemonic PLISSIT stands for:

- P is for permission giving
- LI is for limited information
- SS is for specific suggestion
- IT is for intensive therapy

‘Permission giving’ allows patients to talk about sex, ask questions, feel guilty and so on. Their problems are shared with a reflective listening confidant.

Most medically trained people can probably provide the limited information required about sexual physiology and behavioural patterns.[3] ‘Specific suggestion’ provides ideas for self-help and may include key reference books, and relevant audiotapes or videotapes (See Further reading). Videos can certainly arouse interest, ideas and motivation for a renewal of sexual activity. With a little support and permission, the patient can take simple action to remedy or improve a problem.

Intensive therapy, whether psychiatric or emotional, calls for deeper involvement and can be a dangerous area for the inexperienced. Referral to the appropriate practitioner is usually advisable.[3]

Further reading

Recommended books

- Comfort A. *The Joy of Sex.* London: Mitchell Beazley, 1987.
- Zilbergeld B. *Men and Sex. A Guide to Sexual Fulfilment.* Medindie SA: Souvenir Press, 1979.
- Castleman M. *Making Love* (mainly for men.) Penguin.
- Crooks R, Baur K. *Our Sexuality.* Menlo Park, CA: Benjamin/Cummings Publishing Co., 1984.
- Williams W. *It's Up to You - a Self-Help Book for the Treatment of Erectile Problems.* Sydney: Williams & Wilkins, 1989.
- Rickard-Bell R. *Loving Sex: Happiness in Mateship.* Sydney: Wypikaninkie Publications, 1992.
- Gochros, Fischer. *Treat Yourself to a Better Sex Life.* Simon & Schuster.
- Kitzinger S. *Women's Experience of Sex.* Penguin, 1993.
- Phelps K. *Confronting Sexuality.* Sydney: Harper Collins, 1993.
- Heiman J, Lo Piccolo J. *Becoming Orgasmic: a Sexual Growth Program for Women.* Sydney: Simon & Schuster, 1988.

Recommended videos

- *The lovers' guide I and II.* Andrew Stanway.
- *The language of love.*

ANALOGOUS ROLES OF THE PENIS AND CLITORIS

An explanation of the analogous roles of the penis and clitoris (proposed by Cohen and Cohen) is a very useful strategy for educating patients and helping them to understand the relationship of intercourse and penile and clitoris stimulation with orgasm. The simple model (Figure 109.1) can be shown to patients to explain, for example, why some women are unable to achieve orgasm by intercourse alone, especially using the conventional missionary position.[2, 6] It can readily be explained that clitoral stimulation in women is analogous to penile stimulation in men. Such information is very helpful for women and also to men who may perceive themselves as inadequate lovers. The use of such explanatory aids greatly facilitates the educational process and makes it more 'comfortable' for all concerned.



Figure 109.1 Analogous structures in male and female genitalia. Reproduced with permission from G. and M. Cohen, Canadian Family Physician, 852, 31: 767 - 71.
[2]

FEMALE ORGASMIC DIFFICULTIES

It is necessary to determine whether the woman has been anorgasmic or can experience orgasms from other activities such as masturbation, manual or oral stimulation, even though she is non-orgasmic during intercourse.

The use of the Cohen model (Figure 109.1) is very helpful in emphasising the importance of clitoral stimulation.

Therapy includes:

- sensate focus exercises[7]
- advice on the most appropriate positions for intercourse
- permission to use
    - sexual aids: books, magazines
    - visual tapes
    - self-stimulation

DYSPAREUNIA

Painful intercourse is a source of considerable distress both physically and psychologically for the sufferer and also for her partner. It may be one of the ‘hidden agenda’ presentations with a vague complaint such as ‘I am uncomfortable down below’. Tact and sensitivity is very important in management.

The patient needs to be encouraged to talk freely during history taking with an opportunity to express what they believe is the basic problem. Montgomery claims that most cases (about 80%) of dyspareunia have a physical cause and careful physical examination is mandatory. In particular it is helpful to keep in mind vulval vestibular syndrome, which has subtle physical signs (refer to p. 1031). Important causes are listed in Table 109.6. A common problem encountered is the presence of painful scar tissue following an episiotomy, especially after the first vaginal delivery, so it is worth asking about this potential problem because some women are reluctant to raise the issue.

Table 109.6 Important causes of dyspareunia

Pain worse on insertion
• physiological - inadequate lubrication
• vaginitis in chronic candidiasis
• vulvar dermatoses
• postnatal perineal scarring
• incompletely ruptured hymen
• vulvar vestibular syndrome (vestibulitis)
• vulvovaginal atrophy (e.g. postmenopausal)
• vaginismus
Pain worse on deep penetration
• endometriosis
• PID
• pelvic adhesions
• ovarian and uterine tumours
• postnatal

Management of dyspareunia involves treating the organic cause and giving appropriate advice about the use of lubricants, including oestrogen creams.

Vaginismus

Vaginismus is the involuntary contraction of muscles around the introitus (outer third of vagina) in response to and preventing the possibility of penetration. It is often related to the vulval vestibular syndrome. In some instances it may be voluntary.

It is not an uncommon cause of unconsummated marriages, usually associated with fears of internal damage, pregnancy, learned negative attitudes to sex and part sexual trauma. The problem usually responds well to brief therapy. This includes explanation of the anatomy and physiology with the use of a hand-held mirror during examination.

The patient or couple can use a lubricant such as baby oil, KY jelly, Vaseline or oestrogen cream to make intercourse comfortable. The couple will benefit from a sensate focus program, which includes the most comfortable position for intercourse, controlled by the woman usually in a superior position. Otherwise, progressive vaginal dilation can be practised using lubricated fingers or graduated dilators.

ERECTILE DYSFUNCTION

Erectile dysfunction (impotence) is the inability to achieve or maintain an erection of sufficient quality for satisfactory intercourse. It doesn't refer to ejaculation, fertility or libido. Patients often use the term to refer to a problem of premature ejaculation, hence careful questioning is important.

Erectile dysfunction is a common problem. US data shows the prevalence to be 39% of males at 40 years and 67% of males aged 70.[8]

The most effective and practical approach to the man with erectile dysfunction is to determine the response to an intrapenile injection where prostaglandin E_1 is preferred to papaverine.

Causes of erectile dysfunction

- Psychogenic: related to stress, interpersonal or intrapsychic factors (e.g. depression, marital disharmony, performance anxiety)
- Neurogenic: disorders affecting the parasympathetic sacral spinal cord (e.g. multiple sclerosis); it usually develops gradually
- Vascular
- Diabetes
- Hormone disorder
  - androgen deficiency (e.g. testicular disease)
  - hypothyroidism
  - hyperprolactinaemia (rare) → impotence and loss of libido due to secondary testosterone deficiency
- Drug-induced
  - alcohol
  - cocaine
  - nicotine (four times the risk by age 50)
  - pharmaceutical preparations
- Ageing
- Unknown

History

The nature of the onset of erectile dysfunction is very important and this includes the nature of the relationship. Of particular importance is a drug history, including alcohol, nicotine, street drugs and pharmaceutical agents, particularly antihypertensives (beta-blockers and thiazide diuretics), antidepressants, antipsychotics and H_2-receptor antagonists. Ask about nocturnal and early morning erections.

Examination

This should include a rectal examination and examination of the vascular and neurological status of the lower limbs and the genitalia, especially the testicles and penis. Check the cremaster and bulbocavernosus reflexes.

Investigations

First-line blood tests:

- free testosterone ? androgen deficiency
- thyroxine ? hypothyroidism
- prolactin ? hyperprolactinaemia
- luteinising hormone
- glucose

Other blood tests to consider:

- LFTs, especially GGT (alcohol effect)
- kidney function tests

Nocturnal penile tumescence

This is an electronic computerised test used to detect and measure penile erections during REM sleep. Normally, there are 3 - 5 spontaneous erections lasting 20 - 35 minutes. The test helps to differentiate between psychogenic (normal studies) and organic (poor function). A very simple screening test is to use the snap gauge device.

Dynamic tests of penile function[7]

These tests include injections of drugs into the corpus cavernosum (which is the simplest method) to assess function. If the patient does not have overt psychogenic impotence and the diagnosis is uncertain, the response to intracavernosal injections of prostaglandin E (PGE) can be tested. A good response to PGE indicates that the patient has psychogenic or neurogenic impotence (e.g. due to pelvic nerve division during colon resection). Responses at higher doses indicate an incomplete organic disorder (e.g. partial arterial occlusion, venous leak, or diabetic neuropathy - early). Total failure to respond suggests arterial occlusion or an idiopathic disorder of the corpora cavernosa.

Management

This should comprise appropriate patient education including a videotape of the specific recommended treatment and technique. The partner should be included in the discussions and general management process with an emphasis on bolstering the couple's self-image, which may have been affected by feelings of rejection or avoidance.

Psychogenic disorders

These will involve psychotherapy and sex behavioural modification as outlined under sexual counselling. Referral to a consultant may be appropriate.

Hormonal disorders

- Testosterone for androgen deficiency: primary testicular disease (e.g. Klinefelter's syndrome) or gonadotrophin deficiency
  Stepwise trial
  1. Oral: testosterone undecanoate (Andriol)
  2. IM: testosterone enanthate (Primoteston Depot) or testosterone esters (Sustanon)
  3. Subcutaneous implantation: testosterone implants (last 5 - 6 months)
- Thyroxine for hypothyroidism

- Bromocriptine for hyperprolactinaemia

Oral medication

PDE-5 inhibitors

The phospho-diesterase type 5 (PDE-5) inhibitors are the first line oral medication (Table 109.7). They are about 70% effective but not very effective for neurogenic ED. They do not initiate an erection but enhance whatever erection the man is capable of having. Sexual stimulation is necessary. They are contraindicated if the patient has unstable angina, recent stroke or myocardial infarction. Use with nitrates should be avoided and a nitrate should never be taken within 24 hours of use. PDE-5 inhibitors have the potential for side-effects, especially headache. In some cases combining them with alcohol or fatty foods can delay their onset of action.[9]

Table 109.7 Phospho-Diesterase 5 Inhibitors[9]

	Sildenafil (Viagra)	Tadalafil (Cialis)	Vardenafil (Levitra)
Dosage (mg)	25, 50, 100	10, 20	5, 10, 20
Usual starting dose (1 hour pre i/c)	50 mg	10 mg	10 mg
Onset of action	30 - 60 min	1 - 2 hours	30 - 60 min
Alcohol effect	Yes	Possibly	Possibly
Nitrate contraindication	Yes	Yes	Yes
Class side-effects	Headache, nasal stuffiness, facial flushing, dyspepsia		
Specific s/effects	Blue vision	Myalgia, back pain	

Other oral agents

- Apomorphine (Uprima)
- Trazodole
- Yohimbine

Intrapenile injection

- Alprostadil intracavernosal injections:
    - self-administered after supervised teaching (use a penile model if available)
    - start with a lower dose
    - maximum of three a week
    - if prolonged erection > 2 hours take 2 pseudoephedrine tablets - repeat at 31/2 hours if necessary (provided not hypertensive)

The co-operation of the partner is essential and urological back-up must be arranged.

Transurethral alprostadil (Muse)

Urethral pellet: initial dose 250 µg

Vacuum constriction

Vacuum constriction devices may have a place in management.

Surgery

- Malleable penile prosthesis
- Inflatable penile prosthesis (Figure 109.2)



Figure 109.2 An inflatable prosthesis, showing positioning of the components

- Vascular surgery where appropriate

PREMATURE EJACULATION

Premature ejaculation is a common problem that may not be clearly described by the patient so a careful history is necessary to define the problem. Both patient and partner may complain about the problem.

There are many approaches to treatment but they are aimed either at prolonged ejaculatory control or at satisfactory sexual activity without preoccupation with ejaculation and anticipation of better control with time and experience.

The standard management strategies to enhance ejaculatory control include a combination of three techniques:[7]

- graded sensate focus
- Masters' and Johnson's squeeze technique[10]
- Semans' ‘stop - start’ technique[11]

The tricyclic antidepressant, clomipramine, has been variously reported to be very effective (e.g. 25 - 50 mg/day).[12] The SSRIs have also been reported as effective using fluoxetine 20 mg daily, but are still being evaluated. A new agent, specifically to treat premature ejaculation, the serotonin transport inhibitor, dapoxetine (taken 2 - 3 hours before i/c) is also being evaluated for approval.

GENDER IDENTITY CONCERNS

Sexual orientation or gender identity is a very important issue that GPs should handle competently and sensitively. It is incumbent of the general practitioner to be aware of the special health concerns of people of homosexual persuasion and practice an empathic understanding of their specific organic and psychosocial concerns. It would be true to say that in the past we have been handicapped by ignorance, prejudice, physician homonegativity and social homophobia - an issue probably reinforced by the AIDS epidemic of the 80s. Studies have revealed that significant numbers of gay men did not reveal their sexual orientation to their GP even when presenting with an STI.[13, 14]

People experiencing uncertainty about their gender identity often present with high levels of distress, anxiety and guilt.

Gay men and women may have higher rates of depression, suicide, alcoholism, smoking and organic illness than their heterosexual counterparts.[14] They are also at increased risk of family rejection and being victims of violence.

DSM IV refers to a ‘gender identity disorder’ and outlines diagnostic criteria, including problems of ‘discomfort’.[15]

Definitions and facts

Sexual orientation refers to adult stable sexual attractions, desires and expression towards other men and women and which cannot be determined until emerging adulthood.[13] Homosexuality is the condition in which the libido is directed towards one of the same sex. Lesbians are women where primary emotional and sexual relationships are with other women. The term is derived from a community of women living on the Greek Island of Lesbos around 600 BC.

Gay is the comparable term for men but is generally used to describe both lesbians and gay men and perhaps bisexual people of both sexes. The term ‘gay’ is a more respectful and acceptable term, which seems to have gained more positive recognition from the ‘mardi gras’ gay festivals.

The estimated incidence of homosexuality has varied from 2% to 13% over the past 60 years but the most recent general estimation lies between 3% and 10%, with a higher incidence in males.[13, 14] At the present time there is no proven theory of the causation although it is no longer considered to be one's conscious individual choice or preference but rather a complex innate sociobiological determinant. Genetic and hormonal hypotheses have not been verified to date. Interestingly, gender identity develops in the first four years of life.[16]

It is clear that no one factor causes an individual to become homosexual or heterosexual. In 1948 Kinsey concluded that sexual orientation fell on a continuum from exclusive homosexuality to exclusive heterosexuality at opposite ends, and this perspective remains today. It is noteworthy that Freud determined that homosexual orientation appeared stable in most adults and resistant to modification through analysis.

Recurring problems for gay people

- Health service providers often insensitive to their needs
- Prejudice and discrimination in rural communities
- The threat or reality of violence
- Isolation and loneliness
- Relationships, stress and anxiety
- Homophobia leading to reduced sense of self-esteem and self-worth
- Safety of sexual activity
- Concerns regarding infectious diseases: hepatitis A and B, C, HIV enteric infections, other STIs in gay men
- Concerns regarding fertility, safe insemination practices and raising children in lesbians

Issues and concerns commonly raised

The following are typical of some of the questions and issues posed to the author over several years in family practice:

- How normal or freakish is it to feel like this?
- I hate to think of what my parents will think.
- How should I best approach them?
- How on earth can I tell my spouse (and kids)?
- Should I try out an opposite sex affair?
- Is it advisable to go ahead with the marriage?
- Can I be changed to be ‘normal’?
- What about hypnosis or hormones or a little operation?
- My religious beliefs are a worry. The conflict is devastating.
- What can I do to be safe?
- Is there anyone or a group I can contact about this?

Managing this unfamiliar health area was difficult and it highlights the value of sexual counselling courses, although a listening ear, an acceptance of our patient's natural feelings and commonsense supportive advice is the basis of counselling. The PLI component of the PLISSIT model works very well in general practice. As a rule it is wise to resist giving much specific advice but rather listen and then guide. The appropriate answers to the above questions are that it is important to accept oneself as they are, to let sexuality evolve, not to rush into experimental heterosexual affairs, be sceptical about treatments to change orientation as there is no evidence to date that any are successful, find close trusting friend/friends to share concerns and inform the closest family members as soon as possible and then others. It is appropriate to refer to those with expertise for more intensive therapy.

Developmental stages of the ‘coming out’ process[17]

Coleman has proposed identifiable stages of homosexual identity formation, with each stage having unique health challenges for both the patient and the health provider, especially stress, depression and suicide risk in stage 1:

1. pre-coming out
2. coming out
3. exploration and experimentation
4. first relationships
5. integration

Some guiding management principles[13]

- Don't assume that all patients are heterosexually orientated - use neutral terms.
- Ask good opening questions:
  - ‘How would you see your sexual orientation? Straight, gay or bisexual?’
  - ‘What has been hard for you in your sexual development?’
- The great majority of openly gay men, lesbian women and bisexual persons will experience few or no problems with their sexual orientation.[13]
- A prime reason for dissatisfaction with being homosexual and desiring heterosexuality is feeling lonely and isolated. These people can be counselled successfully through training in social skills and assertiveness.
- Be aware of significant stress-related mental problems and somatisation disorders in people during the ‘pre-coming out’ and ‘coming out’ stages.
- Studies have shown that about 30 - 40% of gay teens have attempted or seriously considered suicide - many times above the national average.
- There is no evidence that reorientation strategies are successful.
- Don't assume stereotyped sexual behaviour and therefore unsafe sex or risk of STIs.
- Anal intercourse only occurs in about one-third of male homosexual encounters.
- It is necessary to differentiate homosexual orientation from transgender activities.
- In counselling it is important that the view of individuals as being either homosexual or heterosexual is not reinforced and that the person as an individual is the prime focus of attention.
- ‘In understanding and managing sexual orientation concerns, the central issue is to appreciate that homosexual individuals cover the same wide spectrum of humanity as heterosexual persons and the division of people into ‘heterosexual’ or ‘homosexual’ groups is arbitrary.’ [13]

THE SMALL PENIS SYNDROME

In general practice it is not uncommon to counsel men and adolescent males for anxiety, sometimes pathological, about the relatively small size of their penis and its possible impact on sexual adequacy. Some males appear to be preoccupied with the size of their penis, especially when reaching the sexually active phase of their life. It is a manifestation of abnormal body image perception.

This attitude is related to the myth that a man's sexual performance depends on the size of his penis. The patient may present with minor (often trivial) non-sexual complaints as a ‘ticket of entry’ into the consulting room or perhaps as a manifestation of anxiety or depression related to preoccupation with penile size.

Measurement

Irrespective of physique or facial configuration most men are concerned about penile size.[15] However, as for all parts of the body, there is considerable variation in size and shape of the penis.

The average adult penis, when measured from the symphysis pubis to the meatus, is 7.5 - 10.5 cm (3 - 4 inches) long when flaccid (Table 109.8).[18, 19] The erect penis has an average length of 15 cm (6 inches) with a range of slightly more than 2.5 cm (1 inch).[18, 19] This increase in size is not necessarily related to the original flaccid state.

Table 109.8 Average penile size

		Flaccid	Erect
Length	centimetres	7.5 - 10.5	12 - 18
	inches	3 - 4	5 - 7
Circumference	centimetres	6 - 10	8 - 12
	inches	2.5 - 4	4 - 6

Masters and Johnson[20] point out that a penis that is larger in its flaccid state does not increase in length proportionately during erection.

Psychological factors

Virility and performance are not related to the size of the penis.[18] Orgasm in the female does not depend on deep vaginal penetration. Penile size was found to have little relationship to a partner's satisfaction from sexual intercourse. The vagina, which is 10 cm (4 inches) long in the unstretched state, tended to accommodate itself to the size of the penis.

Counselling

Counselling the male with fears about sexual inadequacy related to penile size is based on providing reassuring information about the preceding anatomical and physiological facts. The reasons for the patient's concerns should be explored. It should be pointed out that the feeling of inadequacy often follows comparisons with unreal images of macho men portrayed in the media.

If a potential problem is suspected, a useful strategy is to raise the issue subtly by using the third person; in a casual matter-of-fact manner, say something like ‘It's interesting how many men worry about such things as their performance and the size of their penis’.

It is important to emphasise that there is no way of physically enlarging a penis, and this includes regular masturbation and coitus. Furthermore, it should be explained that size has absolutely no relationship with physical serviceability or with the capacity to satisfy a partner.

SEXUALITY IN THE ELDERLY

The sexual needs of the elderly in our society tend to be ignored or misunderstood. While sexual activity and sexual interest generally decline with age, our elderly are not asexual and their sexuality has to be recognised and understood. They have the same needs as younger people - namely, the need for closeness, intimacy and body contact.[21] The same studies have shown that significant numbers of elderly people continue to enjoy both sexual interest and activity throughout their lives. Their activity is determined by factors such as marital status, knowledge about sexuality, prior patterns of sexual expression, privacy and physical health.

A common problem is that termination of sexual activity stems from the belief that people feel they are ‘over the hill’ and have a performance anxiety. This applies particularly to people who have invariably experienced orgasm with intercourse and then start failing to maintain this pattern.

Many women require additional lubrication and need advice about the use of oestrogen cream or lubricating jelly. Testosterone cream has been reported to be beneficial for elderly women with vulvar dryness and fissuring.

The application of the PLISSIT model applies to the elderly with an emphasis initially on permission.

REFERENCES

1. Kumar PJ, Clark ML. *Clinical Medicine* (2nd edn). London: Bailliere Tindall, 1990: 997 - 8.
2. Cohen M, Cohen G. The general practitioner as an effective sex counsellor. Aust Fam Physician, 1989; 18: 207 - 12.
3. Richardson JD. Sexual difficulties: a general practice speciality. Aust Fam Physician, 1989; 18: 200 - 4.
4. Williams W. *It's Up to You - a Self-Help Book for the Treatment of Erectile Problems*. Sydney: Williams & Wilkins, 1985: 16 - 34.
5. Annon JS. *Behavioural Treatment of Sexual Problems. Brief Therapy*. Hagerstown, MA: Harper & Rowe, 1976: 45 - 119.
6. Hite S. *The Hite Report: a Nationwide Study of Human Sexuality*. New York: Dell Publishing Co, 1976: 229.
7. Ross MW, Channon-Little LD. *Discussing Sexuality*. Sydney: MacLennan & Petty, 1991: 42 - 66.
8. Feldman HA et al. Impotence and its medical and psychological correlates: Results of the Massachusetts male aging study. J Urol, 1994; 151: 54 - 61.
9. Lowy M, Baker M. Erectile dysfunction. Australian Doctor, 2005; 11 February: 27 - 34.
10. Masters WH, Johnson VE. *Human Sexual Inadequacy*. Boston: Little, Brown & Co., 1970.
11. Semans JH. Premature ejaculation. A new approach. South Med J, 1956; 49: 353.
12. McMahon CG. Ejaculatory dysfunction. Curr Ther, 1996; 37(3): 49 - 73.
13. Ross MW, Channon-Little LD, Rosser R. *Sexual Health Concerns* (2nd edn) Sydney: MacLennan & Petty, 2002: 161 - 80.
14. Harrison AE. Primary care of lesbian and gay patients: educating ourselves and our students. Fam Med, 1996; 28: 10 - 23.
15. *Diagnostic and Statistical Manual* (4th edn). Washington, DC: American Psychiatric Association, 2000.
16. Newman L. Gender identity issues. Australian Doctor, 2005; 10 June: 33 - 40.
17. Coleman E. Development stages of the comingout process. Am Behavioural Sci, 1974; 25: 469 - 82.
18. Green R. *Human Sexuality*. Baltimore: Williams & Wilkins, 1975: 22 - 3.
19. Katchadourian HA, Lunde DT. *Fundamentals of Human Sexuality*. New York: Holt, Rinehart & Winston, 1975: 44.
20. Masters WH, Johnson VE. *Human Sexual Response*. Boston: Little, Brown & Co., 1966: 191 - 3.
21. Cohen M. Sex after sixty. Can Fam Physician, 1984; 30: 619 - 24.

CHAPTER 110. SEXUALLY TRANSMITTED INFECTIONS

> He who immerses himself in sexual intercourse will be assailed by premature ageing, his strength will wane, his eyes will weaken, and a bad odour will emit from his mouth and his armpits, his teeth will fall out and many other maladies will afflict him.
>
> Moses ben Maimon (1135 - 1204), *Mishneh Torah*

SEXUALLY TRANSMITTED INFECTIONS

STIs are a group of communicable infections, usually transmitted by sexual contact. Their incidence has been of widespread significance during the past 30 years and they are a major public health problem in all countries.

The STIs have developed a high profile in modern society with the advent of HIV infection, hepatitis B, *Chlamydia trachomatis* as a major cause of PID, the emergence of penicillin-resistant gonorrhoea and the increasing frequency of the human papilloma (wart) virus infection with its association with carcinoma of the cervix. STIs are summarised in Table 110.1.

Table 110.1 Sexually transmitted infections: causative organisms and treatment[3, 4]

STI	Causative organism/s	Treatment
Bacterial		
Gonorrhoea	*Neisseria gonorrhoeae*	Ceftriaxone IM + doxycycline or azithromycin
Chlamydia urethritis Non-specific urethritis	*Chlamydia trachomatis Ureaplasma urealyticum Mycoplasma hominis*	Azithromycin (o)
Cervicitis and PID	*Neisseria gonorrhoeae* Chlamydia trachomatis mixed 'vaginal' flora	Mild: doxycycline + metronidazole or tinidazole + azithromycin (o) + ceftriaxone (IM) (if *N. gonorrhoeae*)
		Severe: add cephalosporins (IV use in hospital)
Syphilis	*Treponema pallidum*	Benzathine penicillin: best to refer
Bacterial vaginosis	*Gardnerella vaginalis* other anaerobes	Metronidazole or clindamycin 2% cream
Granuloma inguinale (Donovanosis)	*Calymmatobacterium granulomatis*	Azithromycin
Chancroid	*Haemophilus ducreyi*	Ciprofloxacin or azithromycin: best to refer
Lymphogranuloma venereum	*Chlamydia trachomatis*	Doxycycline: best to refer
Viral		
AIDS	HIV-1, HIV-2	Refer to Internet site and specialist
Genital herpes	Herpes simplex virus	Aciclovir, famciclovir or valaciclovir
Genital warts	Human papilloma virus	Podophyllotoxin paint or imiquimod cream
Hepatitis	HBV, HCV	Interferon, anti-viral agent
Molluscum contagiosum	Pox virus	Various simple methods (e.g. deroofing with needle)
Fungal		
Vaginal thrush (possible) Balanoposthitis	*Candida albicans*	Any antifungal preparation
Protozoal		

Vaginitis, urethritis Balanoposthitis	*Trichomonas vaginalis*	Tinidazole or metronidazole
Arthropods		
Genital scabies	*Sarcoptes scabiei*	Permethrin 5% cream
Pediculosis pubis	*Phthirus pubis*	Permethrin 1% lotion

KEY FACTS AND GUIDELINES

- In Western society most patients with STIs are in the 15 - 30 year age group.1
- Gonorrhoea and syphilis are no longer the commonest STIs.
- Chlamydial infection, hepatitis B, human papilloma virus and genital herpes are now common infections.
- Not all STIs are manifest on the genitals.
- Not all genital lesions are STIs.
- The 5% rule2
  - 5% of urethritis (STI) in males is lower UTI
  - 5% of lower UTI in females is urethritis (STI)
- *C. trachomatis* is now the commonest cause of urethritis.
- *Chlamydia* typically causes dysuria in men but may be asymptomatic. It usually causes no symptoms in women.
- Gonorrhoea may cause no symptoms, especially in women.
- STIs such as donovanosis, lymphogranuloma venereum and chancroid occur mainly in tropical countries. Donovanosis is common in Indigenous Australians.
- The presentation of STI in children, especially vaginitis, should alert practitioners to consider sexual abuse.
- HIV infection, which is predominantly sexually transmitted, should be considered in any person at risk of STI as well as IV drug users. It must be appreciated that it can present as an acute febrile illness (similar to Epstein-Barr mononucleosis) before going into a long asymptomatic ‘carrier’ phase.
- Nucleic acid amplification tests such as PCR on urine and other specimens have been a great advance in management of STIs.

COLLECTION OF SPECIMENS

It is mandatory to collect the appropriate specimens before treatment, because of the epidemiological implications.

Material requirements (obtainable from laboratories):

- standard MSU jar
- standard swabs
- stiff wire swabs
- glass slides
- teflon-coated slides for viral or chlamydial microscopy
- transport media (three types)

- Stuart's (or Amies or similar)
- chlamydial
- viral

Your laboratory will advise on the most appropriate test kits and methods of collection.

PRESENTING CONDITIONS[2]

Most STIs fit into one (or sometimes more) of the easily definable categories of clinical presentation:

- urethritis - discharge and/or dysuria
- vaginitis - discharge + irritation + odour + dyspareunia
- cervicitis/PID (possible symptoms)
    - pelvic pain/lower abdominal pain (PID)
    - backache (PID)
    - mild discharge
    - mucopurulent cervical discharge
    - dyspareunia (PID)
    - dysuria
- ulcer
- lump
- pruritus
- rash with
    - secondary syphilis
    - HIV infection
    - hepatitis B

VAGINITIS

Vaginitis is presented in more detail in Chapter 96.

The common pathogens are:

- *Candida albicans* ??vaginal thrush
- *Trichomonas vaginalis*
- *Gardnerella vaginalis* ??bacterial vaginosis

Of the three common pathogens, only *Trichomonas* is considered to be sexually transmitted and the only vaginitis requiring routine treatment of partners.

Gardnerella is more a marker of the condition than a true pathogen. Bacterial vaginosis (also termed anaerobic vaginosis) is really an altered physiological state rather than an infection or inflammation. The hallmark of the condition is the absence of lactobacilli. It is important to note that anaerobic vaginosis is frequently asymptomatic and found by accident when vaginal swabs are made for other purposes. In these circumstances treatment is not warranted.

Collection of specimens

Make two slides:

- one smear for air-drying and Gram stain
- one wet film preparation, under a cover slip, for direct inspection for the
    - pseudohyphae of *Candida*
    - 'clue cells' of *Gardnerella*
    - motile *Trichomonas*

Treatment (in summary)[1, 2]

- *Candida* on Gram stain: any antifungal preparation e.g. clotrimazole 500 mg vaginal tablets statim and clotrimazole 1% cream daily (or nystatin) for symptomatic relief
- *Gardnerella* on Gram stain: e.g. metronidazole 400 mg (o) bd for 7 days or clindamycin 2% cream Aci-jel topically bd
- *Trichomonas* on wet film: e.g. tinidazole or metronidazole 2 g (o) statim and treat partner

URETHRITIS

The important STIs that cause urethritis are gonorrhoea and *Chlamydia,* which is three times more common than gonorrhoea.2 Non-specific urethritis (NSU) is commonly due to *C. trachomatis* but may also be caused by *Ureaplasma, Mycoplasma homminis* and other unknown organisms.

Symptoms of urethritis

In males

The main symptoms (if present) are:

- a burning sensation when passing urine (dysuria)
- a penile discharge or leakage (clear, white or yellow)

Sometimes there is no discharge, just pain. Sometimes the infection is asymptomatic. Most often the symptoms are trivial with *Chlamydia*. Although a creamy pus-like discharge is typical of gonorrhoea, and a less obvious milky-white or clear discharge typical of *Chlamydia*, it is often difficult to differentiate the causes from the discharge. In some males the only complaint is spots on the underpants or dampness under the foreskin. Epididymo-orchitis in the young male should be presumed to be a complication of an STI urethritis.

In females

Gonorrhoea often causes no symptoms but can produce vaginal discharge or dysuria or PID. *Chlamydia* usually causes no symptoms but may cause vaginal discharge, dysuria or PID. NSU is the commonest form of PID, which can result in infertility.

Gonococcal infection of anus and throat

In both sexes, gonorrhoea may infect the anus or oropharynx. Anorectal gonorrhoea may be asymptomatic or may present as a mucopurulent anal discharge (a feeling of dampness) and anal discomfort.

Oropharangeal gonorrhoea may be asymptomatic or present as a sore throat or dysphagia.

Collection of specimens

Take two swabs:

- standard swab for *Gonococcus* (into the urethral meatus): place into Stuart's transport medium
- wire swab for *Chlamydia* (2 - 3 cm into the urethra and endocervical canal in women and twist around for minimum15 seconds) after wiping away frank pus and exudate (Figure 110.1). Place into *Chlamydia* transport medium



Figure 110.1 Taking a urethral swab

Note: A wire swab and dedicated *Chlamydia* transport medium are appropriate for *Chlamydia* diagnosis, as sap from wooden swabs and chemicals from some plastics kill *Chlamydia*. Culture is now less important.

In males PCR or ligase chain reaction (LCR) are the tests that usually provide the diagnosis of chlamydial infection.5 The PCR *Chlamydia* urine test (95% specific) is the preferred test (in both sexes) on the first catch urine specimen (don't urinate for 4 hours, then first 10 mL passed in ordinary MSU jar). These two tests are also used to diagnose gonorrhoea. However, urine PCR is less reliable in women for gonorrhoea than an endocervical specimen.

Special notes

- Testing for *Chlamydia* and *Gonococcus* can be performed on a PAP smear.
- It is possible for women to do self-collecting from tampons.
- Take an MSU in males who have dysuria but no discharge. *or* Take urethral swabs from females who have dysuria but not frequency. The presence of large numbers of coliform in a urethral swab culture is suggestive of bacterial cystourethritis (lower UTI).
- If there are gonococci on Gram stain, treat with azithromycin and doxycycline; if no gonococci on Gram stain, treat with azithromycin; and if microscopy unavailable give both antibiotics.

CHLAMYDIA URETHRITIS

Incubation period

Symptoms appear 1 - 2 weeks after intercourse, although the incubation period can be as long as 12 weeks or as short as 5 days (compare with incubation period of gonorrhoea - about 2 - 3 days).

It is an underdiagnosed disorder since many cases are asymptomatic; hence the value of screening.[6] Woman can carry *Chlamydia* silently for 12 months or more.

Treatment[5, 6]

Azithromycin 1 g (o) single dose (preferred) *or* Doxycycline 100 mg (o) 12 hourly for 7 days

A second course may be required if the symptoms persist or recur (about one in five cases).

Second-line treatment is erythromycin 500 mg qid for 7 days. All sexual partners, even if asymptomatic, need to be treated in the same way. If a female partner has proven cervicitis the treatment must be as for PID. Sexual intercourse must be avoided until the infection has cleared up in both partners. The importance of compliance must be stressed.

Prevention

Using condoms for vaginal and anal sex provides some protection.

Screening guidelines for higher risk

- All sexually active females < 25 years
- All sexually active teenagers, especially females, Aboriginals and Torres Strait Islanders
- Those with a pattern of inconsistent or no condom usage:
  - men who have anal sex with men
  - 6 - 12 months post-infection

GONORRHOEA

Incubation period

Gonorrhoea has a short incubation period of 2 - 3 days and symptoms usually appear 2 - 7 days after vaginal, anal or oral sex. The incubation period can be as long as 3 weeks.

Other manifestations of gonorrhoea

- Epididymo-orchitis and prostatitis (males)
- Urethral stricture is not uncommon in males

Treatment

If there is infection with penicillin-resistant gonococci (PPNG) due to β-lactamase (penicillinase) production (a problem prevalent in South-East Asia and eastern Australia), the following should be used.[3, 6] The organism is becoming resistant to quinolone.

ceftriaxone 250 mg IM (dissolved in lignocaine

1 - 2 mL 1%) as a single dose (preferred)

or

plus either

azithromycin 1 g (o) as a single dose

or

doxycycline 100 mg (o) bd for 10 days

Where PPNG prevalence is low:

amoxycillin 3 g (o) as a single dose +

probenecid 1 g (o)

plus

azithromycin 1 g (o) as a single dose (if chlamydia has not been ruled out)

If the above antibiotics are inappropriate (e.g. pregnancy):

erythromycin 500 mg (o) bd for 10 days

or

roxithromycin 300 mg (o) daily for 10 days

Sexual partners must be examined and treated and sexual intercourse must be avoided until the infection has cleared. Follow-up culture (after 4 weeks) is advisable, especially in females.

PREVENTION

Using condoms for vaginal, anal and oral sex provides good protection. Sexually active men and women (especially those at risk) need at least annual checks.

CERVICITIS

Cervicitis is often a forerunner to PID. If there is cervicitis only (mucopus at the cervix without uterine pain or tenderness) treat as for urethritis (the likely organisms are *C. trachomatis* or *N. gonorrhoeae*) - ciprofloxacin plus doxycycline.

PELVIC INFLAMMATORY DISEASE

PID is covered in more detail in Chapter 93. It is not always an STI. The intrauterine device is also a common cause. Often multiple pathogens are involved in the infection.

Common pathogens are *N. gonorrhoeae* and *C. trachomatis*. Swabs from the cervical os frequently underestimate the organisms involved and thus treatment needs to be directed to all possible pathogens.

Mucopurulent cervicitis is now known to be an early sign of PID, usually due to *Chlamydia*.[2]

Specimen collection

Cervical and urethral swabs for urethritis for *N. gonorrhoeae* and *C. trachomatis*.

Treatment

Therapy for PID is deliberately vigorous because the major aim is to prevent infertility and the consequent need for IVF in the long term. The detailed treatment is outlined on pp. 1000 - 1.

Summary[2, 3, 6]

Mild to moderate infection:

a 14-day course of doxycycline 100 mg bd

plus

azithromycin 1 g as a single dose

plus

metronidazole or tinidazole

If gonorrhoea: ceftriaxone 250 mg IM, single dose Severe PID: hospitalise for IV therapy

ULCERS

STI causes of genital ulcers are presented in Table 110.2. Most genital ulcers are herpes - any small genital ulcer which is superficially ulcerated, scabbed, red-edged, multiple and painful is invariably herpes.

Table 110.2 STI causes of anogenital ulcers

	Pain	Specimen collection
Common		
Herpes simplex virus	Yes	Scraping for direct immunofluorescence Swab for antigen detection by PCR and culture into viral transport medium
Uncommon		
Treponema pallidum (primary chancre)	No	Exudate for dark ground microscopy and serum for leutic screen (reagin or treponemal tests)
Haemophilus ducreyi	Yes	Scraping for Gram stain and special culture
Calymmatobacterium granulomatis (granuloma inguinale)	No	Scraping for special stains

Syphilis is uncommon and may get over-looked, especially with anal chancres.

Chancroid is almost always an imported infection.

GENITAL HERPES

The incubation period is usually 3 - 6 days but can be longer. A firm microbiological diagnosis is recommended. Swab for culture or PCR.

HSV 1 and 2 account for over 90% of cases.

Symptoms

With the first attack there is a tingling or burning feeling in the genital area. A crop of small vesicles then appears; these burst after 24 hours to leave small, red, painful ulcers. The ulcers form scabs and heal after a few days. The glands in the groin can become swollen and tender, and the patient might feel unwell and have a fever.

The first attack lasts about 2 weeks.

Males

The virus usually affects the shaft of the penis, but can involve the glans and coronal sulcus, and the anus (Figure 110.2).



Figure 110.2 Usual sites of vesicles/ulcers in males

Females

Vesicles develop around the opening of, and just inside, the vagina and can involve the cervix and anus (Figure 110.3). Passing urine might be difficult, and there can be a vaginal discharge. In about 25%, the cervix is the only site of lesions and these cases may be asymptomatic. Th is occult presentation is a problem and a high index of suspicion is needed. The problem could be a simple tingling around the saddle area or a scratch-like lesion on epithelium.



Figure 110.3 Usual sites of vesicles/ulcers in females

In both sexes, it can affect the buttocks and thighs. A serious but uncommon complication, especially in females, is the inability to pass urine.

Specimen collection

Take a swab from a vesicle for culture, direct immunofluorescence or PCR.

Transmission

It can be caught by direct contact through vaginal, anal or oral sex. Rarely is it transferred to the genitals from other areas of the body by the fingers. Sometimes in stable relationships it appears spontaneously. A new type-specific herpes serology (EIA) is available for testing partners of someone with known HSV 2 infection.[5]

Recurrence

Half of those who have the first episode have recurrent attacks; the others have no recurrence. Fortunately, attacks gradually become milder and less frequent, last 5 - 7 days and usually stop eventually. Recurrences after many months or years can be precipitated by menstruation, sexual intercourse, masturbation, skin irritation and emotional stress. Condom use reduces the risk by about 50%.

Treatment of genital herpes

Antimicrobial therapy

Topical treatment

The proven most effective topical therapy is topical aciclovir (not the ophthalmic preparation). Other topical preparations provide relief but do not alter significantly the course of the infection; they should be applied as soon as the symptoms start.

Alternatives:

10% silver nitrate solution applied with a cotton bud to the raw base of the lesions, rotating the bud over them to provide gentle debridement. Repeat once or twice. This promotes healing and spreading. *or* 10% povidone - iodine (Betadine) cold sore paint on swab sticks for several days.

Pain relief can be provided in some patients with topical lignocaine.

Oral treatment

For the first episode of primary genital herpes (preferably within 24 hours of onset):

aciclovir 200 mg 5 times a day for 5 days or

until resolution of infection[6]

or

famciclovir 125 mg (o) bd for 5 days

or

valaciclovir 500 mg (o) bd for 5 days

This appears to reduce the duration of the lesions from 14 days to 5 - 7 days.[2] The drugs are not usually used for recurrent episodes, which last only 5 - 7 days. A 5-day course of any of the drugs can be used for a rather severe recurrence. Very frequent recurrences (six or more attacks in 6 months) benefit from continuous low-dose therapy for 6 months (e.g. valaciclovir 500 mg (o) once daily).

Supportive treatment (advice to the patient)

- Rest and relax as much as possible. Warm salt baths can be soothing.
- Icepacks or hot compresses can help.
- Painkillers such as aspirin or paracetamol give some relief.
- If urination is painful, pass urine under water in a warm bath.
- Keep the sores dry; dabbing with alcohol or using warm air from hairdryer can help.
- Leave the rash alone after cleaning and drying; do not poke or prod the sores.
- Wear loose clothing and cotton underwear. Avoid tight jeans.

Counselling

'A chat beats medicine for herpes'. Since genital herpes is distressing and recurrent, patients are prone to feel stressed and depressed and can be assisted by appropriate counselling and support. Sexual abstinence should be practised while lesions are active. Consider referral to a self-help/support group.

Prevention

Spread of the disease can be prevented by avoiding sexual contact during activity of the lesions. Condoms offer some protection (not absolute) and patients should wash their genitals with soap and water immediately after sex. Condoms should always be used where a partner has a history of this infection.

SYPHILIS

In Australia syphilis usually presents either as a primary lesion or through chance finding on positive serology testing (latent syphilis).

It is important to be alert to the various manifestations of secondary syphilis. The classification and clinical features of syphilis are presented in Table 110.3 (see also Chapter 30).

Table 110.3 Classification and clinical features of syphilis

Type	Time period	Infectivity	Clinical features
Acquired			
Early (within first 2 years of infection)			
Primary	10 - 90 days, average 21	Infectious	Hard chancre Painless Regional lymphadenopathy
Secondary	6 - 8 weeks after chancre	Infectious	Coarse non-itchy maculopapular rash Constitutional symptoms (may be mild) Condylomata lata Mucous membrane lesions
Early latent	months to 2	Infectious	No clinical features but positive serology

	years		
Late (after the 2nd year of infection)			
Late latent	2 years plus	Non-infectious	
Tertiary (now rare)		Non-infectious	Late benign: gummas *or* Cardiovascular *or* Neurosyphilis
Congenital			
Early	within first 2 years of life	Infectious	Stillbirth or failure to thrive Nasal infection: ‘snuffles’ Skin and mucous membrane lesions
Late	after second year of life	Non-infectious	Stigmata (e.g. Hutchinson's teeth) Eye disease CNS disease Gummas

Transmission

- Sexual intercourse (usual common mode)
- Transplacental to fetus
- Blood contamination: IV drug users
- Direct contact with open lesions

Management

The management of syphilis has become quite complex and referral of the patient to a specialist facility for diagnosis, treatment and follow-up is recommended.

Recommended antimicrobial therapy

Early syphilis (primary, secondary or latent) of not more than one year's duration:[6]

benzathine penicillin 1.8 g IM as a single dose

or

procaine penicillin 1 g IM daily for 10 days

For patients hypersensitive to penicillin:

doxycycline 100 mg (o) 12 hourly for 14 days

or

erythromycin 500 mg (o) 6 hourly for 14 days

Note:

- Sex should be avoided until ulcers healed.
- Sexual contacts in the past 3 months should have treatment.

Late latent syphilis: more than 1 year or indeterminate duration:

benzathine penicillin 1.8 g IM once weekly for

3 doses, or procaine penicillin

Cardiovascular and neurosyphilis and congenital syphilis are also treated with penicillin but require special regimens.

LUMP

Common pathogens:

- wart (papilloma) virus - condylomata acuminata, venereal ‘warts’, anogenital warts
- molluscum contagiosum (pox) virus

Uncommon:

- *Treponema pallidum* - condylomata lata

Physiological:

- Fordyce cysts which are enlarged ectopic sebaceous glands in the mucosa are a differential diagnosis of a genital lump

Diagnosis

Warts and molluscum contagiosum have a distinctive appearance and are readily diagnosed by inspection (Figure 110.4). Removal for diagnosis is usually not required. Condylomata acuminata are multiple lesions that resemble warts superficially but are covered by abundant exudate. They occur in secondary syphilis and leutic screen is positive.



Figure 110.4 General appearance of molluscum contagiosum and genital warts

Treatment of warts[6]

Counselling and support are necessary. Not all genital warts are sexually transmitted.

Warts may be removed by chemical or physical means, or by surgery. Treatment needs to be individualised. For small numbers of readily accessible warts the simplest treatment is:[6]

- cryotherapy weekly until resolved *or*
- podophyllotoxin 0.5% paint or 0.15% cream (a more stable podophyllin preparation)
  - apply bd with plastic applicator for 3 days
  - repeat in 4 days and then weekly for 4 - 6 cycles if necessary

Note:

The normal surrounding skin should be spared as much as possible. Avoid this treatment in pregnancy on cervical, meatal or anorectal warts. *or*

- topical imiquimod 5% (Aldara) cream applied by the patient three times a week to each wart for up to 16 weeks (usually 12 - 24 weeks).

All females (including partners of males with warts) should be referred to a specialised clinic where colposcopy is available, because of the causal link of warts to cervical cancer.

Treatment of molluscum contagiosum

These lesions often resolve spontaneously. There are many treatment choices to provoke resolution.

These include:

- deroofing aseptically with a needle or sharppointed stick and expressing the contents (recommended)
- lifting open the tip with a sterile needle inserted from the side and applying 10% povidone - iodine (Betadine) solution
- liquid nitrogen (for a few seconds)
- application of 25% podophyllin in tincture of benzoin compound
- application of 30% trichloroacetic acid
- destruction with electrocautery or diathermy

ITCH

Common pathogens:

- *Sarcoptes scabiei* (scabies)
- *Phthirus pubis* (pubic lice)
- *C. albicans*
  - vulvovaginitis - females
  - balanitis - males

Non-STI itchy rashes on genitals include dermatitis and psoriasis.

DIAGNOSIS

Scabies: inspection on scraping and microscopy. Inspection: scabies is diagnosed by a very itchy, lumpy rash. It is rare to find the tiny mites, but it may be possible to find them in the burrows, which look like small wavy lines.

Pubic lice: inspection for moving lice and nits (eggs) on hair shaft.

C. albicans: swab for Gram stain and *Candida* culture.

TREATMENT

Scabies[5]

Permethrin 5% cream if > 2 months of age. Apply to whole body from jawline down (include every flexure and area), leave overnight, then wash off. Wash clothing and linen after treatment and hang them in sun. *or* Benzyl benzoate 25%, left for 24 hours before washing off.

The whole family and close contacts must be treated regardless of symptoms which can take weeks to develop. One treatment is usually sufficient. It can be repeated in a week if necessary.

Note: Persistence of the itch after treatment is common. If the itch has not abated after 7 days, re-treat. After this, reassurance is usually all that is required. Also prescribe a topical antipruritic (e.g. crotamiton cream for 3 - 5 days and an oral antihistamine for the itch).

Pubic lice

Permethrin 1% lotion: apply to pubic hair and surrounding area, leave for 10 minutes and then wash off. *or* Pyrethrins 0.165% with piperonyl butoxide 2% in foam base; apply as above.

Shaving pubic hair is also effective. Bed clothes and underwear should be washed normally in hot water after treatment and hung in the sun to dry. Repeat the treatment after 7 days. Sometimes a third treatment is necessary. Sexual contacts and the family must be treated (young children can be infested from heavily infested parents). Where the lice or the nits are attached to eyelashes, insecticides should not be used: apply Vaseline liberally to the lashes.

Candidiasis

Topical imidazole (e.g. clotrimazole 1% applied 2 - 3 times daily).

EXTRAGENITAL STIS

Viral hepatitis

Sexual activity is a factor in the transmission of hepatitis B (in particular), hepatitis A (where faecal - oral contact is involved), hepatitis C (probably occasionally) and hepatitis D.[1]

Hepatitis B[1]

In Western societies, sexual transmission of HBV is a common mode of spread and there is a higher prevalence in homosexual men and prostitutes. HBV prevalence in homosexual men is correlated with insertive and receptive anogenital contact and oroanal contact.

There is no specific therapy for hepatitis B, so prevention is important. Interferon α-2 can be used for complications such as chronic active hepatitis.

Prevention[1]

Several prevention strategies are available; they include:

- immunisation
- prevention of infection in health care establishments
- management of exposure (needlestick injuries etc.)
- management of infants of mothers who are hepatitis B carriers
- condoms, which offer some reduction of risk of sexual transmission
- personal hygiene

Immunisation

Immunisation should be encouraged in hepatitis B marker-free people at risk of acquiring this infection. At-risk groups include sexual partners of carriers, institutional individuals, all homosexuals, prostitutes and drug addicts. Some health workers are exposed to risk.

Management of exposure

Sexual partners of acute cases and chronic carriers who are negative for surface antigen (HBsAg) and antibody can be offered hepatitis B immunoglobin, and routine hepatitis B immunisation should be commenced.

Hepatitis C

Although there have been doubts about the potential for sexual transmission of hepatitis C, Tedder et al. in 19917 demonstrated evidence for the sexual transmission of HCV but the epidemiological evidence is not strong.

HIV INFECTION

HIV infection (colloquially called AIDS, although this represents only the severe end of the disease spectrum) is predominantly transmitted by IV drug use in the community. In Australia about 80% of HIV cases are related to IV drug use and the rest are mainly sexually transmitted.[8] The important risk factors in infected men are receptive anal intercourse and multiple sexual partners.

Sexual transmission to women[1]

Although the heterosexual partners of infected men are at risk of infection, spread to and from women has been relatively uncommon in developed countries, but now appears to be increasing significantly. In central Africa, heterosexual spread is an important means of transmission. Genital ulcerative diseases such as syphilis and genital herpes may be associated with an increased risk of heterosexual transmission.

HIV infection is considered in more detail in Chapter 28.

THE FULL STI CHECK-UP[7, 9]

Family doctors may be consulted by a prostitute or other sexually active female requesting a thorough check-up. Such people require certificates from time to time and may not have access to a public STI clinic. The visit should provide an opportunity for counselling and education about her health risks.

The screening program includes:

- full sexual history
- physical examination: genital appearance, skin, breasts, oropharynx, lymph nodes, abdomen, careful anogenital examination
- investigations (guide only):
    - Pap smear: 6 - 12 monthly
    - first catch urine for PCR
    - endocervical swabs for *Chlamydia* and gonorrhoea 1 - 3 monthly (depending on risk)
    - high vaginal swab and ‘wet film prep’ for vaginal pathogens 1 - 3 monthly
    - HIV antibody test (with informed consent) - not ordered more often than every 3 months
    - syphilis screening test: RPR/VDRL (as for HIV)
    - hepatitis B screening: if negative, organise hepatitis B vaccination
    - rubella IgG as baseline test: if negative, advise rubella vaccination

Consider:

- Throat swabs for gonorrhoea (if oral sex without condoms)
- Urethral swab for gonorrhoea and *Chlamydia* if urinary symptoms
- anorectal swab for gonorrhoea if sexual history indicates need

WHEN TO REFER

- Syphilis
    - probably all suspected or confirmed cases but certainly for suspected tertiary syphilis
    - HIV-positive patients, and
    - suspected treatment failure
- Pubic lice and scabies
    - unresolved rash or itch despite apparently appropriate treatment
- Genital warts
    - urethral or cervical warts
    - associated cervical HPV changes on cytology
    - refractory warts
- Gonorrhoea or non-specific urethritis
    - if complications, pelvic spread or extragenital problems develop, or symptoms persist after two courses of antibiotics.

CONTACT TRACING

It is important to contact partners of those presenting with more serious STIs. The Government's guidelines recommend that for patients with asymptomatic infection (e.g. *Chlamydia*, gonorrhoea), partners in the previous 12 months should be contacted and they should refer sexual contact partners within 30 days of the onset of symptoms. A discreet letter sent to the contact is advisable.[10]

Practice tips

- Do not presume that the patient or his/her partner has acquired an STI outside their relationship.
- The itch of scabies or pubic lice can be distressing: prescribe the topical antipruritic (crotamiton cream) and/or an oral antihistamine.
- Reassure the patient that the itch will gradually subside over a few weeks (especially with scabies). This allays anxiety that may lead to overzealous self-medication.
- Make every attempt to confirm or exclude genital herpes, using the appropriate investigations.
- Use aciclovir or similar antivirals for first episodes of genital herpes and when recurrences are either frequent or painful.
- Twelve golden rules of management are presented in Table 110.4.

Table 110.4 STIs: Twelve golden rules of management (Venereology Society of Victoria)

1 An STI can only be diagnosed if the possibility is considered.
2 An adequate sexual history is paramount.
3 A proper history and careful examination must precede laboratory investigations.
4 Remember the sexual partner(s)!
5 Treatment consists of the appropriate antibiotic in correct dosage for an adequate period of time.
6 A patient concerned about STIs is probably an 'at risk' patient.
7 Counselling and education are fundamental to STI management.
8 Penicillin will not cure NSU.
9 *Not* all vaginal discharges are thrush.
10 Multiple, painful genital ulcers are most often due to herpes simplex.
11 Prompt, accurate treatment of PID is necessary to preserve fertility.
12 Remember the three Cs - Consent, Confidentiality and Counselling - of HIV antibody testing.

GOOD COMMUNICATION

Note: Tests for STIs, including the HIV antibody test, should only be performed with the patient's knowledge and consent and after adequate counselling. An appointment should be made to give results person to person, irrespective of the results.

ACKNOWLEDGMENT

Part of this text, on hepatitis B, is reproduced from the *Handbook on Sexually Transmitted Diseases*. ‘Commonwealth of Australia copyright, reproduced by permission’. Professor John Turnidge has given permission to adopt his categories of presenting conditions.

FURTHER READING

National management guidelines for sexually transmissable infections. Melbourne: Venerology Society of Victoria Inc, 2002. < http://www.mshc.org.au >.

REFERENCES

1. *NH&MRC Handbook on Sexually Transmitted diseases*. Canberra: Department of Community Services & Health, 1990: 1 - 55.
2. Turnidge J. *Sexually Transmitted Diseases*. CHECK Programme. Melbourne: RACGP, 1989, Unit 210/211.
3. Donovan B. Management of sexually transmissable infections. Medicine Today, 2006; 7(1): 63 - 5.
4. Department of Health and Ageing. Communicable Diseases Intelligence 2005, 29: 417 - 33.
5. Waddell R. *Sexually Transmitted Diseases*. CHECK Programme. Melbourne: RACGP, May 2002, Issue 363.
6. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 13). Melbourne: Therepeutic Guidelines Ltd, 2006: 93 - 116.
7. Tedder RS, Gilson RJC, Briggs M et al. Hepatitis C virus: evidence for sexual transmission. Br Med J, 1991; 302: 1299 - 1302.
8. National Centre in HIV Epidemiology and Clinical Research. HIV/AIDS and Related Diseases in Australia. Annual Surveillance Report 1997.
9. Bradford D. *Sexually transmitted disease*. CHECK Programme. Melbourne: RACGP, 1993, Unit 252/253: 7 - 8.
10. Wines N, Daylan L. Nongonococcal urethritis management in general practice. Med Today, 2000; November: 33 - 6.

CHAPTER 111. A DIAGNOSTIC AND MANAGEMENT APPROACH TO SKIN PROBLEMS

The skilful doctor knows by observation, the mediocre doctor by interrogation, the ordinary doctor by palpation.

CHANG CHUNG-CHING (c. AD 170-196)

A DIAGNOSTIC AND MANAGEMENT APPROACH TO SKIN PROBLEMS

The diagnosis of skin problems depends on astute clinical skills based on a systematic history and examination and, of course, experience. If the diagnosis is in doubt it is appropriate to refer the patient to a skilled cooperative consultant, as the referral process is an excellent educational opportunity for the GP. Another opinion from a colleague/s in a group practice is also very educative. At least, cross-referencing the skin lesion with a colour atlas facilitates the learning process.

TERMINOLOGY OF SKIN LESIONS

Primary lesions

- *Macule.* Circumscribed area of altered skin colour < 1 cm diameter (Figure 111.1).
- *Patch.* Macule of > 1 cm diameter (Figure 111.1).
- *Papule*. Palpable mass on skin surface < 0.5 cm diameter (Figure 111.2).
- *Maculopapule.* A raised and discoloured circumscribed lesion.
- *Nodule*. Circumscribed palpable mass > 0.5 cm diameter (Figure 111.2).
- *Plaque*. A flat-topped palpable mass > 1 cm diameter.
- *Wheal*. An area of dermal oedema (can be any size).
- *Angio-oedema*. A diffuse area of oedema extending into subcutaneous tissue.
- *Vesicle*. A fluid-filled blister < 0.5 cm in diameter (Figure 111.3).
- *Bulla*. A vesicle > 0.5 cm diameter (Figure 111.3).
- *Pustule.* A visible collection of pus in the skin < 1 cm diameter.
- *Abscess.* A localised collection of pus in a cavity > 1 cm diameter.
- *Furuncle*. A purulent infected hair follicle; includes:
    - folliculitis (small furuncles)
    - boils (larger furuncles)
- *Carbuncle*. A cluster of boils discharging through several openings.
- *Purpura*. Bleeding into the skin (dermis) appearing as multiple haemorrhages. May be macular or papular.
- *Petechiae*. Purpuric lesions 2 mm or less in diameter.
- *Ecchymosis*. Larger purpuric lesion.
- *Haematoma*. A swelling from gross bleeding.
- *Telangiectasia*. Visible dilatation of small cutaneous blood vessels.
- *Comedo*. A plug of keratin and sebum in a dilated pilosebaceous gland.
- *'Blackhead'*. An open comedo.
- *'Whitehead'.* A closed comedo.
- *Erythema*. Redness of the skin due to increased vascularity.
- *Milium*. Tiny white cyst containing keratin, from occlusion of pilosebaceous gland.
- *Papilloma*. Warty projection above the skin surface.



Figure 111.1 Macule and patch



Figure 111.2 Papule and nodule



Figure 111.3 Vesicle and bulla

Secondary lesions

- *Scales*. An accumulation of excess keratin that presents as flaking.
- *Crust*. Superficial dried secretions (serum and exudate).
- *Ulcer*. A circumscribed deep defect with loss of all the epidermis and part or all of the dermis (Figure 111.4); they usually heal with scarring.
- *Erosion*. A skin defect with complete or partial loss of the epidermis; they heal without scarring (Figure 111.4).
- *Fissure*. A linear split in the epidermis and dermis (Figure 111.4).
- *Atrophy*. Thinning or loss of epidermis and/or dermis with loss of normal skin markings.
- *Sclerosis*. Thickening of the dermis with induration of subcutaneous tissue; resembles a scar but may arise spontaneously (e.g. scleroderma).
- *Scar*. A healed dermal lesion where normal structures are replaced by fibrous tissue.
- *Hypertrophic scar.* Rises above the skin surface.
- *Atrophic scar.* Settles below the skin surface.
- *Keloid*. Overgrowth of dense fibrous tissue extending beyond the original wound.
- *Excoriation*. Scratch marks causing an erosion or an ulcer (loss of epidermis).
- *Acanthosis*. Thickening of skin without accentuation of skin markings.
- *Lichenification*. Thickening secondary to chronic scratching or rubbing (in dermatitis).
- *Callus*. Localised hypertrophy of the stratum corneum.
- *Exfoliation*. Loss of epidermal keratin as large scales or sheets.
- *Keratoderma*. Thickening of skin especially stratum corneum.



Figure 111.4 Ulcer, erosion and fissure

Defining terms

Terms that are continually referred to in skin disease include:

- seborrhoea = yellow-brown and waxy
- nummular = coin-like }(interchangeable)
- discoid = disc-like }(interchangeable)
- annular = ring-like
- circinate = circular
- arcuate = curved
- reticulate = net-like
- pityriasis = (pityron = bran): fine branlike scaly desquamation or powdery
- guttate = 'dew drop'
- rosea = rose-coloured
- morbilli-form = like measles
- morphoea = circumscribed scleroderma or skin infiltrate
- livido = cyanotic discoloration
- lichen = any papular skin disorder resembling lichens
- verrucous = rough and warty

A DIAGNOSTIC APPROACH

The diagnostic approach of Robin Marks[1] presented here helps to achieve order in the midst of confusion. He describes the importance of simplifying the diagnostic process by being a 'lumper' rather than a 'splitter'. Most common dermatological problems fall into one of seven categories (Table 111.1).

Table 111.1 Common dermatological conditions

Infections	
bacterial	• impetigo
viral	• warts
	• herpes simplex, herpes zoster
	• pityriasis rosea
	• exanthemata
fungal	• tinea
	• candidiasis
	• pityriasis versicolor
Acne	
Psoriasis	

Atopic dermatitis (eczema)	
Urticaria acute and chronic papular	• pediculosis
	• scabies
	• insect bites
Sun-related skin cancer	
Drug-related eruptions	

A problem that does not fit into one of these seven groups is either an unusual condition or an unusual presentation of a common condition and probably merits a consultant's opinion.

History

The three basic questions are:[1]

1. Where is the rash and where did it start?
2. How long have you had the rash?
3. Is the rash itchy?

Note: The split into three time zone groups (Table 111.2) is very useful. This question leads on to the next question regarding itch, as the patient is unlikely to tolerate an itchy eruption.

Table 111.2 How long has the rash been present?

Acute (hours - days)	urticaria
	atopic dermatitis
	allergic contact dermatitis
	insect bites
	drugs
	herpes simplex/zoster
	viral exanthemata
Acute → chronic (days - weeks)	atopic dermatitis
	impetigo
	scabies
	pediculosis
	drugs
	pityriasis rosea
	psoriasis
	tinea
	Candida
Chronic (weeks - months)	Psoriasis
	atopic dermatitis
	tinea
	pityriasis versicolor
	warts
	cancers

	skin infiltrations (such as granulomata, xanthomata)

If so, is it mild, moderate or severe? The nature of the itch is very helpful diagnostically. A severe itch is one that wakes the patient at night and leads to marked excoriation of the skin, while a mild itch is one that is only slightly upsetting for the patient and may not be noticeable for significant periods during the day.

Three questions the doctor must consider

1. Could this be a drug rash?
2. Has this rash been modified by treatment?
3. Do any contacts have a similar rash?

Further questions for the patient

- Do you have contact with a person with a similar eruption?
- What medicines are you taking or have taken recently?
- Have you worn any new clothing recently?
- Have you been exposed to anything different recently?
- Do you have a past history of a similar rash or eczema or an allergic tendency (e.g. asthma)?
- Is there a family history of skin problems?

The nature of itching [1]

The characteristics of the itch are very useful in dividing up the diagnoses: an eruption that is not itchy is unlikely to be scabies and one that is very itchy is unlikely to be a skin tumour (Table 111.3).

Table 111.3 Is the rash itchy?

Very	urticaria
	atopic dermatitis
	scabies, pediculosis
	insect bites
	chickenpox (adults)
	dermatitis herpetiformis
	Grover's disease
Mild to moderate	tinea
	psoriasis
	drugs
	pityriasis rosea
	Candida
	stress itching/lichen simplex
Often not	warts, tinea
	impetigo, psoriasis
	cancers
	viral exanthemata
	seborrhoeic dermatitis

However, nothing is absolute and variations to the rule will occur - tinea, psoriasis and pityriasis versicolor are sometimes itchy and sometimes not. Chickenpox can vary from being intensely itchy, especially in adults, to virtually no itching.

Relieving or aggravating factors of the itch provide useful diagnostic guidelines; for example, Whitfield's ointment applied to an itchy eruption for a provisional diagnosis of ringworm would make the itch worse if it were due to eczema.

Table 111.4 Appearance of individual lesions

Epidermal	atopic dermatitis
	psoriasis
	tinea
	pityriasis rosea
	impetigo, herpes, warts
	cancers
	scabies
	solar keratoses
Dermal	urticaria
	insect bites, pediculosis, scabies
	drugs
	skin infiltrations
	viral exanthemata



Figure 111.5 Epidermal skin lesion



Figure 111.6 Dermal skin lesion

Physical examination [1]

Examine the skin in good light, preferably natural light, and ensure that any make-up is removed.

There are two basic stages in the physical examination of a rash. The first is an assessment of the characteristics of the individual lesion and the second is the distribution or pattern of the lesions.

Characteristics of the individual lesion

The single most important discriminating feature is whether it involves the dermis alone or the epidermis as well (Table 111.4). If the lesion involves the epidermis there will be scaling, crusting, weeping, vesiculation or a combination of these (Figure 111.5). If the dermis alone is involved the lesion is by definition a lump, a papule or a nodule (Figure 111.6). No lesion ever involves the epidermis without involving the dermis as well.

Other characteristics of individual lesions which must be sought are the colour, the shape and the size. It is important to feel the skin during the physical examination and to note the consistency of the lesion: Is it firm or soft? The activity of the lesion may also be useful: does it have a clearing centre and an active edge?

Distribution of the lesions

The clinician must decide whether the lesions are localised or widespread. If they are widespread, are they distributed centrally, peripherally, or both? (Table 111.5). Diagnosis is often helped when the skin lesions are in a specific area (Table 111.6 and Figures 111.7, 111.8). Itchy papules on the penis associated with a widespread pruritus is likely to be scabies. However, care has to be taken because many misdiagnoses are made instinctively on the distribution (e.g. anything in the flexures is dermatitis or anything on the feet is tinea).

Table 111.5 Distribution of the rash

Widespread	atopic dermatitis
	psoriasis scabies
Central trunk (initially at least)	drugs urticaria tinea versicolor
Peripheral	herpes zoster pityriasis rosea viral exanthemata atopic dermatitis
	herpes zoster tinea
	psoriasis
	warts insect bites

Table 111.6 Specific areas affected

Face	rosacea
	impetigo atopic dermatitis psoriasis photosensitive (e.g. drugs) herpes simplex acne cancers viral exanthemata

Scalp	psoriasis
	seborrhoeic dermatitis
	pediculosis
	tinea
	chickenpox
Flexures	atopic dermatitis
	psoriasis
	seborrhoeic dermatitis
	tinea
	Candida
	pediculosis
Mouth	aphthous ulcers
	herpes simplex
	Candida
	measles
Nails	psoriasis
	tinea
Penis	scabies

Tables 111.1 to 111.6 have been prepared by Dr Robin Marks[1] and are reproduced with his permission.

Another feature of an eruption which should be sought on examination is whether the lesions are all at the same stage of evolution.

It is necessary to perform a complete physical examination as well. There is, after all, no such thing as a skin disease but rather disease affecting the skin. The clinician must always bear this in mind when managing patients complaining of a skin eruption. Disease does not affect the skin in isolation and it is unforgivable to look only at the skin and ignore the patient as a whole.

Note: In every case examine the mouth, scalp, nails, hands and feet.

DIAGNOSTIC TOOLS

Appropriate diagnostic tools include:

- a magnifying lens
- a diascope, which is a glass slide or clear plastic spoon that is used to blanch vascular lesions in order to determine their true colour
- a ‘maggylamp’, which is a hand-held fluorescent light with an incorporated magnifier; the device allows shadow-free lighting and magnification
- a dermatoscope - very valuable in the diagnosis of pigmented tumours but it does require skill and familiarity to achieve effective use
- Wood's light
- swabs for culture
- skin biopsy (even with psoriasis, etc.)


Eyelids
seborrhoeic dermatitis/blepharitis
atopic dermatitis
xanthelasma
allergic contact dermatitis
psoriasis
Face
cancers
acne
rosacea
systemic lupus erythematosus
atopic dermatitis
pityriasis alba
impetigo
keratoacanthoma
solar keratoses
seborrhoeic dermatitis
leprosy (Hansen's disease)
Lips
herpes simplex
Candida chelitis
leucoplakia
Scalp
tinea capitis
psoriasis
seborrhoeic dermatitis
pediculosis
Ears
seborrhoeic dermatitis
allergic contact dermatitis
solar keratosis
Nasolabial folds
perioral dermatitis
seborrhoeic dermatitis
Folliculitis
sycosis barbae (S. aureus)
tinea barbae
Chin
perioral dermatitis
seborrhoeic dermatitis
rosacea

Figure 111.7 Typical sites on the face affected by the skin conditions indicated


Cubital fossa
atopic dermatitis
Hands
occupational dermatoses
granuloma annularae
dyshydrosis
contact dermatitis
tinea mannum
secondary syphilis
Chest
seborrhoeic dermatitis
acne
pityriasis versicolor
Grover's disease
drug exanthem
tinea
Axillae
hidradenitis
erythrasma
contact dermatitis
seborrhoeic dermatitis
tinea
psoriasis
Nails
tinea
candidiasis
psoriasis
Trunk
pityriasis versicolor
pityriasis rosea
guttate psoriasis
drug exanthem
scabies
Elbows (dorsum)
psoriasis
dermatitis herpetiformis
Wrists
contact dermatitis (bands)
lichen planus
solar keratoses
Knees (dorsum)
psoriasis
dermatitis herpetiformis
Legs
keratitis pilaris
dermatofibroma
erythema nodosum
necrobiosis lipoidica
nummular dermatitis
stress dermatitis
Dorsum of foot
granuloma annularae
lichen simplex chronicus
contact dermatitis
atopic dermatitis
Groin
tinea cruris
candidiasis
erythrasma
seborrhoeic dermatitis
contact dermatitis
Toes
tinea pedis
contact dermatitis
erythrasma
candidiasis
Perianal
psoriasis
seborrhoeic dermatitis
warts
Candida albicans
Popliteal fossa
atopic dermatitis
Soles of feet
plantar warts
tinea
hyperhidrosis
pitted keratolysis
callus
corn
pustular psoriasis
secondary syphilis
hand, foot and mouth disease
juvenile plantar dermatosis

Figure 111.8 Typical regional location of various skin conditions

* Same conditions apply to chest and trunk

OFFICE TESTS AND DIAGNOSTIC AIDS

Wood's light

Wood's light examination is an important diagnostic aid for skin problems in general practice. It has other uses, such as examination of the eye after fluorescein staining. (A new, low-cost, small ultraviolet light unit called 'the black light' is available.)

Method

Simply hold the ultraviolet light unit above the area for investigation in a dark room.

Limitations of Wood's light in diagnosis

Not all cases of tinea capitis fluoresce because some species that cause the condition do not produce porphyrins as a by-product. See Table 111.7 for a list of the skin conditions that do fluoresce. Wood's light is really only useful for hair-bearing areas.

Table 111.7 Skin conditions that produce fluorescence in Wood's light

Tinea capitis	green/bright yellow (in hairs)
Erythrasma	coral pink
Pityriasis versicolor	pink-gold
Pseudomonas spp.	yellowish-green
Porphyria cutanea tarda	red (urine)

Porphyrins wash off with soap and water, and a negative result may occur in a patient who has shampooed the hair within 20 hours of presentation. Consequently, a negative Wood's light reading may be misleading. The appropriate way of confirming the clinical diagnosis is to send specimens of hair and skin for microscopy and culture.

Skin scrapings for dermatophyte diagnosis

Skin scrapings is an excellent adjunct to diagnosis of fungal infections. Requirements are a scalpel blade, glass slide and cover slip, 20% potassium hydroxide (preferably in dimethyl-sulphoxide) and a microscope. Skin scrapings can also be sent for microscopy and culture.

Clinical indications:

1. tinea (superficial dermatophyte infection)
2. pityriasis versicolor
3. *Candida*

Method

- Scrape skin from the active edge.
- Scoop the scrapings onto the glass microscope slide.
- Cover the sample with a drop of potassium hydroxide.
- Cover this with a cover slip and press down gently.
- Warm the slide and wait at least 5 minutes for ‘clearing’.

Microscopic examination

- Examine at first under low power with reduced light.
- When fungal hyphae are located, change to high power.
- Use the fine focus to highlight the hyphae (Figure 111.9).

Note: Some practice is necessary to recognise hyphae.

Other uses of microscopy

Detection of the scabies mite: the burrow of the scabies mite is found (can be difficult!) and the epidermis is decisively scraped with a no. 15 scalpel blade, and transferred to a slide after adding a drop of liquid paraffin. The mite is very distinctive.

Patch testing

Patch testing is used to determine allergens in allergic contact dermatitis. Read in 48 hours.

Biopsies

Punch or shave biopsies can be useful (see Figures 116.25 and 116.26 at pp. 1213 - 4).



Figure 111.9 Diagrammatic representation of microscopic appearance of fungal hyphae

Hair[2]

Send hair samples for microscopy and root analysis. There are two main tests:

1. the hair pull test (refer p. 1231)
2. the hair pluck test

In the hair pluck test (used for tinea capititis and hair shaft disorder), a tuft of about 20 hairs are drawn out, forceps applied close to the skin of the scalp and rotated slightly prior to sharp extraction. The hair should be collected on a glass slide for counting and analysis.

Terminology of topical skin preparations

Bases or **vehicles** A mixture of powders, water and greases (usually obtained from petroleum). The relative blending of these compounds determines the nature of the base (e.g. lotion, cream, ointment, gel or paste).

Moisturiser An agent that increases the water content of the stratum corneum. Classifiedas:

- emollient
- humectant, and
- occlusive, e.g. white soft paraffin (Vaseline)

Emollient A topical preparation of emulsified oils and fatty acids that is softening or soothing to the skin. It replaces natural oils in the stratum corneum. It also acts as a skin moisturiser and is therefore used on dry skin or dermatoses related to dry skin (e.g. atopic dermatitis). Examples are:

- mineral or vegetable oil, e.g. peanut oil 5% cream
- sorbolene cream
- aqueous cream

Humectant A chemical-containing agent that attracts and retains water due to its hygroscopic or osmotic properties. Examples are:

- urea 10% cream
- glycerol 10% cream

Astringent A topical agent that has styptic or binding properties with an ability to stop secretions from skin or tissues. An example is aluminium acetate solution (Burow's solution); the aluminium acetate acts as a protein precipitator and is a very effective soothing agent and antipruritic.

Keratolytic An agent that softens or breaks up keratin. Examples are:

- urea 10% - for xerosis or keratosis pilaris
- urea 20% - cracked palms and soles
- salicylic acid 2-10%
- benzoic acid
- alpha-hydroxy acids, e.g. lactic acid, propylene glycol

Antipruritic agent One that relieves itching. Examples are:

- menthol (0.25%)
- phenol (0.5%)
- coal tar solution (2 - 10%)
- camphor (1 or 2%)

Lotion A suspension of an insoluble powder in water. Modern lotions use an emulsifying agent, which eliminates the need to shake the lotion. An example is calamine lotion (zinc oxide 5, calamine 15, glycerine 5, water to 100).

Paints and tinctures Rapidly drying liquid preparations which are very useful for intertriginous areas, especially between the toes and natal cleft. 'Tincture' is the preparation when alcohol is the vehicle. Example: podophyllin in tinct. benz. co. (for genital warts).

Cream A suspension of a powder in an emulsion of oil and water with the addition of an emulsifying agent.

Ointment A suspension of a substance in an oily vehicle.

Gels Substances with a greaseless, watermiscible base.

Pastes Similar to ointments in composition but are more viscid. They consist of an ointment to which another agent such as starch has been added.

Emulsions Mixtures of two immiscible liquids, one being dispersed throughout the other in small droplets.

Table 111.8 Guidelines for choosing a topical vehicle

Disorder	Topical vehicle
Acute inflammation: erythema, weeping	wet dressing solution lotions
Subacute inflammation: erythema, scaling	creams gels (for hairy areas)
Chronic inflammation: scaling, dryness, thickening	ointments impregnated tapes

TRADITIONAL CHEMICALS USED IN EXTEMPORANEOUS PREPARATIONS[3]

- *Salicylic acid:* produces painless destruction of epithelium, thereby facilitating absorption. Consider its use for psoriasis, neurodermatitis, tinea of palms and feet, seborrhoeic dermatitis.
- *Resorcinol:* a topical keratolytic with bactericidal and fungicidal properties. Consider its use for psoriasis, acne, rosacea, seborrhoeic dermatitis.
- *Tar:* the two commonly found tar preparations are used for their anti-inflammatory, soothing and antimitotic properties. Consider their use for psoriasis, atopic dermatitis, seborrhoeic dermatitis and neurodermatitis.
- *Menthol* and *phenol:* added to various preparations for their soothing and cooling effects. Consider for use in pruritic problems such as varicella, urticaria and atopic dermatitis.
- *Sulphur:* of benefit in dermatoses, due mainly to its keratolytic properties. The other actions of sulphur are

scabicidal, parasiticidal and fungicidal. Consider its use for acne, rosacea, seborrhoeic dermatitis, psoriasis and tinea.

- *Dithranol:* for psoriasis.

SELECTION OF CORTICOSTEROID PREPARATIONS

- Class I and class II preparations are appropriate for most problems.
- Creams or lotions are used for ‘weeping’ lesions, the face, flexures and hair-bearing areas.
- Use ointments for dry and scaly skin.
- Use ointments and occlusive vehicles for dry and chronic skin surfaces.
- Ointments should not be used on weeping surfaces.
- Stubborn dermatoses such as psoriasis respond better to preparations under occlusion, such as plastic wrap applied overnight with appropriate securing in place.
- Use a gel or lotion for the scalp.
- For *Candida* infection (e.g. complicating seborrhoeic napkin dermatitis), mix 1% hydrocortisone in equal quantities with an antifungal preparation such as nystatin.
- Use the weakest strength for chronic dermatoses but treat severe acute dermatoses aggressively.
- Steroids not needed if no inflammation present.

Topical corticosteroids for chronic dermatoses

Guidelines for long-term use (examples):[4]

- face - hydrocortisone 1%
- flexures - hydrocortisone 1%
- trunk - betamethasone valerate 0.2%
    - triamcinolone acetonide 0.02%
- elbows/knees - betamethasone dipropionate 0.05%
- palms/soles - methyprednisolone aceponate 0.1%

Cautions

- Avoid high-potency preparations on the face, in flexures and on infants.
- Corticosteroids can mask or prolong an infection.
- Long-term use can cause striae and skin atrophy, perioral dermatitis, ‘steroid acne’ and rosacea.
- Excessive use of more potent preparations can cause adrenal suppression; predispositions include use > 2 weeks, and use on thinner skin such as face, genitalia and intertriginous areas.
- Avoid sudden cessation: alternate with an emollient or a milder preparation.

The relative clinical potency of topical corticosteroids is given in Table 111.9.

Table 111.9 Potency ranking of the most commonly used topical corticosteroid preparations in Australia and New Zealand

Generic name	Formulation
Group I Mild	
Hydrocortisone 0.5%	Cream
Hydrocortisone acetate 0.5%	Cream
Hydrocortisone 1%	Cream
Hydrocortisone acetate 1%	Cream, ointment
Group II Moderately potent	

Betamethasone valerate 0.02%	Cream, ointment
Betamethasone valerate 0.05%	Cream, ointment, gel
Clobetasone butyrate 0.05%*	Cream, ointment
Triamcinolone acetonide 0.02%	Cream, ointment
Triamcinolone acetonide 0.05%	Cream, ointment
Group III Potent	
Betamethasone valerate 0.1%	Cream, ointment, scalp lotion
Betamethasone diproprionate 0.05%	Cream, ointment, lotion
Fluocinolone acetonide 0.025%*	Cream, gel, ointment
Fluclorolone acetonide 0.025%*	Cream
Fluorcortolone hexanoate 0.25%*	Cream
Hydrocortisone butyrate 0.1%*	Cream, lipocream
Methylprednisolone aceponate 0.1%	Cream, ointment
Mometasone furoate 0.1%	Cream, ointment, lotion
Triamcinolone acetonide 0.1%	Cream, ointment
Group IV Super-potent	
Betamethasone diproprionate 0.05% (enhanced)	Cream, ointment
Clobetasol propionate 0.05%*	Cream, ointment, scalp application
Halcinonide 0.1%	Cream
Preparations containing other agents	
Triamcinolone acetonide 0.01% +	
neomycin, gramicidin, nystatin	Cream, ointment
Betamethasone valerate 0.1% + gentamicin	Cream, ointment
Hydrocortisone 1% + clioquinol 1%	Cream
Hydrocortisone 1% + clioquinol 3%	Cream
Hydrocortisone 1% + clotrimazole	Cream

* Not available in Australia.

SKIN TIPS

- Do no harm. Introduce the mildest possible preparation to alleviate the problem.
- Creams tend to be drying, lotions even more.
- Ointments tend to reduce dryness and have greater skin penetration. If wet - use a wet dressing (wet soaks and a lotion). If dry - use an ointment (salve).
- Occlusive dressings with plastic wraps permit more rapid resolution of stubborn dermatoses.
- Most toilet soaps are alkaline and are very drying; they should not be used on dry skin or dermatitis with dry skin. Soap substitutes include neutral soaps (Dove, Neutrogena), superfatted soaps (Oilatum) and non-soap cleanser (Cetaphil).
- Soaps should be used only in the axillae and groin and on the feet of people with dry and irritated skin.
- Bath additives can be useful for dermatoses such as psoriasis, atopic dermatitis and for pruritus. For some people it may be better not to add it to the bath (diluting effect; accident from slipping) but to massage the oil into the dry itchy skin after the bath.

- Always give careful instructions to the patient regarding application of preparations: use a prepared handout if available.
- Alter the treatment according to the response.
- Explain the costs involved, especially where a preparation is expensive.
- Avoid combination creams unless clear evidence of secondary infection proven by culture and sensitivity.

RULES FOR PRESCRIBING CREAMS AND OINTMENTS

How much cream? [5]

On average, 30 g of cream will cover the body surface area of an adult. Ointments, despite being of thicker consistency, do not penetrate into the deeper skin layers so readily, and the requirements are slightly less. Pastes are applied thickly, and the requirements are at least 3 - 4 times as great as for creams.

The 'rule of nines', used routinely to determine the percentage of body surface area affected by burns (Figure 111.10 on p. 1149), may also be used to calculate the amount of a topical preparation that needs to be prescribed.

For example:

- if 9% of the body surface area is affected by eczema, approximately 3 g of cream is required to cover it
- 9 g of cream is used per day if prescribed three times daily
- a 50 g tube will last 5 - 6 days

Table 111.10 Suitable quantities of skin preparations for specific body areas[6] (twice daily application for 1 week)

	Creams and ointments		
	Corticosteroids	**Others**	**Lotions**
Face and neck	15-30 g	15-30 g	100 mL
Both hands	15-30 g	25-50 g	200 mL
Scalp	15-30 g	50-100 g	200 mL
Both arms	30-60 g	100 g	200 mL
Both legs	100 g	100-200 g	200 mL
Trunk	100 g	400 g	500 mL
Groins and genitalia	15-30 g	15-25 g	100 mL

One gram of cream will cover an area approximately 10 cm × 10 cm (4 square inches), and this formula may be used for smaller lesions.

Table 111.10 provides guidelines for approximate weekly quantities of skin preparations required to cover specific areas of the body.



Figure 111.10 ‘Rule of nines’ for body surface area

Some general rules

Remember:

- to use cream or lotions for acute rashes
- to use ointments for chronic scaling rashes
- that a thin smear only is necessary
- that 30 g
  - will cover the adult body once
  - will cover hands twice daily for 2 weeks
  - will cover a patchy rash twice daily for 1 week
- that 200 g will cover a quite severe rash twice daily for 2 weeks

REFERENCES

1. Marks R. A diagnosis in dermatology. Aust Fam Physician, 2001; 30(11): 1028 - 32.
2. Marks R, Sinclair R. *A Guide to the Performance of Diagnostic Procedures used in the Management of Common Skin Diseases*. Melbourne: Skin & Cancer Foundation Publication, 2002: 25 - 6.
3. Kelly B. Extemporaneous preparations. Aust Fam Physician, 1993; 22: 842 - 4.
4. Marley J (Chair). *Therapeutic Guidelines* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2004: 35 - 6.
5. Gambrill J. How much cream? Aust Fam Physician, 1982; 11: 350.
6. George CF et al. London: British National Formulary, Number 31, 1996; 451 - 6.

CHAPTER 112. PRURITUS

It is easy to stand a pain, but difficult to stand an itch.

CHANG CH'AO 1676

PRURITUS

Pruritus (the Latin word for itch) is defined simply as the desire to scratch.

It is one of the most important dermatological symptoms and is usually a symptom of primary skin disease with a visible rash. However, it is a subjective symptom and diagnostic difficulties arise when pruritus is the presenting symptom of a systemic disease with or without a rash. An associated rash may also be a manifestation of the underlying disease.

The broad differential diagnoses of pruritus are:

- skin disease
- systemic disease
- psychological and emotional disorders

PHYSIOLOGY[1]

Itch arises from the same nerve pathway as pain, but pain and itch are distinct sensations. The difference is in the intensity of the stimulus. Unrelieved chronic itch, like unrelieved pain, can be intolerable and cause suicide. There are many similarities: both are abolished by analgesia and anaesthesia; subdued by counter-irritation, cold, heat and vibration; and referred itch occurs just like referred pain. Antihistamines which act on the H_1 receptor are often ineffective, suggesting that histamine is not the only mediator of itch.[1]

LOCALISED PRURITUS

Pruritus may be either localised or generalised. Localised itching is generally caused by common skin conditions such as atopic dermatitis (Table 112.1). Scratch marks are generally presented. Pruritus is a feature of dry skin. An intense localised itch is suggestive of scabies, also known as ‘the itch’.

Table 112.1 Primary skin disorders causing significant pruritus

Atopic dermatitis (eczema)
Urticaria
Dermatitis herpetiformis
Grover's disease
Scabies
Pediculosis
Asteatosis (dry skin)
Lichen planus
Chickenpox
Contact dermatitis
Insect bites

Itching of the anal and vulval areas is a common presentation in general practice.

A careful examination is necessary to exclude primary skin disease; a detailed history and examination should be undertaken to determine if one of the various systemic diseases is responsible.

The history may provide a lead to the diagnosis - the itching of polycythaemia may be triggered by a hot bath which can cause an unusual prickling quality that lasts for about an hour.[2] On the other hand, the itching may be caused by a primary irritant such as a 'bubble bath' preparation.

GENERALISED PRURITUS

Pruritus may be a manifestation of systemic disease. It can accompany pregnancy, especially towards the end of the third trimester (beware of cholestasis), and disappear after childbirth. These women are then prone to pruritus if they take the contraceptive pill.[3]

Systemic causes are summarised in Table 112.2 and a summary of the diagnostic strategy model is given in Table 112.3.

Table 112.2 Systemic conditions that can cause pruritus

Pregnancy
Chronic kidney failure
Liver disorders
• cholestatic jaundice, e.g.
- carcinoma of head of pancreas
- primary biliary cirrhosis
- drugs: chlorpromazine, antibiotics
• hepatic failure
Malignancy
• lymphoma: Hodgkin's lymphoma
• leukaemia, esp. chronic lymphatic leukaemia
• multiple myeloma
• disseminated carcinoma
Haematological disorders
• polycythaemia rubra vera
• iron-deficiency anaemia
• pernicious anaemia (rare)
• macroglobulinaemia
Endocrine disorders
• diabetes mellitus
• hypothyroidism
• hyperthyroidism
• carcinoid syndrome
• hyperparathyroidism
Malabsorption syndrome
• gluten sensitivity (rare)

Tropical infection/intestinal parasites
• filariasis
• hookworm
Drugs
• alkaloids
• opiates. cocaine
• quinidine
• chloroquine
• CNS stimulants
Senile pruritus
Polyarteritis nodosa
Irritants
• fibreglass
• others
Psychological and emotional causes:
• anxiety/depression
• psychosis
• parasitophobia

Table 112.3 Generalised pruritus: diagnostic strategy model

Q. Probability diagnosis A. Psychological/emotional[3] Old dry skin (senile pruritus) Atopic dermatitis (eczema)
Q. Serious disorders not to be missed A. Neoplasia • lymphoma/Hodgkin's • leukaemia: CLL • other carcinoma Chronic kidney failure Primary biliary cirrhosis
Q. Pitfalls A. Pregnancy Tropical infection/infestation Polycythaemia Generalised sensitivity (e.g. fibreglass, bubble bath)
Q. Seven masquerades checklist A. Depression Diabetes Drugs

Anaemia	iron deficiency
Thyroid disorders	hyper and hypo
Spinal dysfunction	-
UTI	-
Q. Is the patient trying to tell me something?	
A. Quite likely: consider anxiety, parasitophobia.	

Guidelines

- The prevalence of itching in Hodgkin's lymphoma is about 30%. The skin often looks normal but the patient will claim that the itch is unbearable.[2]
- Pruritus can be the presenting symptoms of primary biliary cirrhosis and may precede other symptoms by 1 - 2 years.[3] The itch is usually most marked on the palms and soles.
- Pruritus can occur in both hyperthyroidism and hypothyroidism, especially in hypothyroidism where it is associated with the dry skin.

Investigations to consider

- Urinalysis
- Pregnancy test
- FBE and ESR
- Kidney function tests
- Liver function tests
- Thyroid function tests
- Random blood sugar
- Stool examination (ova and cysts)
- Chest X-ray
- Lymph node biopsy (if present)
- Immunological tests for primary biliary cirrhosis

Treatment

The basic principle of treatment is to determine the cause of the itch and treat it accordingly. Itch of psychogenic origin responds to appropriate therapy, such as amitriptyline for depression.[1]

If no cause is found:

- apply cooling measures (e.g. airconditioning, cool swims)
- avoid rough clothes; wear light clothing
- avoid known irritants
- avoid overheating
- avoid vasodilatation (e.g. alcohol, hot baths/showers - keep showers short and not too hot)
- treat dry skin with appropriate moisturisers, e.g. propylene glycol in aqueous cream
- topical treatment
    - emollients to lubricate skin
    - local soothing lotion such as calamine, including menthol or phenol (avoid topical antihistamines)
    - pine tar preparations (e.g. Pinetarsol)
    - crotamiton cream
    - consider topical corticosteroids
- sedative antihistamines (not very effective for systemic pruritus)
- non-sedating antihistamines during day

- antidepressants or tranquillisers (if psychological cause and counselling ineffective)

PRURITIC SKIN CONDITIONS

Scabies

Scabies is a highly infectious skin infestation caused by a tiny mite called *Sarcoptes scabiei*. The female mite burrows just beneath the skin in order to lay her eggs. She then dies. The eggs hatch into tiny mites, which spread out over the skin and live for only about 30 days. The mite antigen, in its excreta, causes a hypersensitivity rash. Diagnosis is by microscopic examination of skin scrapings.

Clinical features

- Intense itching (worse with warmth and at night)
- Erythematous papular rash
- Usually on hands and wrists
- Common on male genitalia (see Chapter 110) (Figure 112.1)



Figure 112.1 Genital scabies causing severe pruritus, showing bruising on the upper thighs from intense scratching

- Also occurs on elbows, axillae, feet and ankles, nipples of females (Figure 112.3)



Figure 112.3 Typical distribution of the scabies rash

Spread

The mites are spread from person to person through close personal contact (skin to skin), including sexual contact. They may also be spread through contact with infested clothes or bedding, although this is uncommon. Sometimes the whole family can get scabies. The spread is more likely with overcrowding and sexual promiscuity.

Norwegian scabies

While the majority of cases have a relatively small number of mites (as few as 15), infestation with thousands or millions will cause the crusted condition of Norwegian scabies. Diagnosis is made on a scraping which reveals vast numbers of lesions. It may be encountered in nursing homes. Treatment is with ivermectin 200 μg/kg (o) as single dose plus topical treatment.[4]

Treatment

For all ages (except children under 2 months) permethrin 5% cream (preferable) *or* benzyl benzoate 25% emulsion (dilute with water if under 10 years)

- Apply to entire body from jawline down (including under nails, in flexures and genitals)
- Leave permethrin overnight, then wash off thoroughly
- Leave benzyl benzoate for 24 hours
- Avoid hot baths or scrubbing before application
- Treat the whole family at the same time even if they do not have the itch
- Wash clothing and bedclothes as usual in hot water and hang in sun
- One treatment is usually sufficient but repeat scabicide treatment in 1 week for moderate and severe infections
- For children under 2 months use sulphur 5% cream daily for 2 - 3 days or crotamiton 10% cream daily for 3 - 5 days



Figure 112.2 Scabies mite (*Sarcopes scabiei*)

Dermatitis herpetiformis

This extremely itchy condition is a chronic subepidermal vesicular condition in which the herpes simplex-like vesicles erupt at the dermoepidermal junction. The vesicles are so pruritic that it is unusual to see an intact one on presentation.

Some consider that it is always caused by a glutensensitive enteropathy. Most patients do have clinical coeliac disease.

Clinical features

- Most common in young adults
- Vesicles mainly over elbows and knees (extensor surfaces)
- Also occurs on trunk, especially buttocks and shoulders (Figure 112.4)



Figure 112.4 Typical distribution of dermatitis herpetiformis

- Vesicles rarely seen by doctors

- Presents as excoriation with eczematous changes
- Masquerades as scabies, excoriated eczema or insect bites
- Typically lasts for decades
- Associated with gluten-sensitive enteropathy
- Skin biopsy shows diagnostic features

Treatment

- Gluten-free diet
- Dapsone 100 mg (o) daily (usually dramatic response)

Lichen planus

Lichen planus is an epidermal inflammatory disorder of unknown aetiology characterised by pruritic violaceous flat-tipped papules, mainly on the wrists and legs. If in doubt diagnosis is confirmed by biopsy. One differential diagnosis is a lichen planus-like drug eruption, e.g. antihypertensives, anti-malarials.

Clinical features

- Young and middle aged adults
- Small, shiny, lichenified plaques
- Symmetrical and flat-tipped
- Violaceous
- Flexor surfaces: wrists, forearms, ankles
- Can affect oral mucosa - white streaks or papules or ulcers
- Can affect nails and scalp

Management

- Explanation and reassurance
- Usually resolves over 6 to 9 months, leaving discoloured marks without scarring
- Recurrence rare
- Asymptomatic lesions require no treatment
- Topical moderately potent corticosteroids (may use occlusive dressing)
- Intralesional corticosteroids for hypertrophic lesions

Pruritus ani

The generalised disorders causing pruritus may cause pruritus ani. However, various primary skin disorders such as psoriasis, dermatitis, contact dermatitis and lichen planus may also cause it, in addition to local anal conditions. It is covered in more detail in Chapter 36.

Pruritus vulvae

Refer to Chapter 97.

Tinea cruris

Also known as Dhobie itch and jock itch, tinea cruris is a common infection of the groin area in young men (Figure 112.5), usually athletes, that is commonly caused by a tinea infection, although there are other causes of a groin rash (Table 112.4). The dermatophytes responsible for tinea cruris (Dhobie itch) are *Trichophytonrubrum (60%), Epidermophyton floccosum* (30%) and *Trichophyton mentagrophytes.*[5] The organisms thrive in damp, warm, dark sites. The feet should be inspected for evidence of tinea pedis. It is transmitted by towels and other objects, particularly in locker rooms, saunas and communal showers.



Figure 112.5 Trinea cruris (also known as Dhobie itch and jock itch) in a young man caused by *Trichophytonrubrum*

Table 112.4 Common causes of a groin rash (intertrigo)

Simple intertrigo
Skin disorders
• psoriasis
• seborrhoeic dermatitis
• dermatitis/eczema
Fungal
• *Candida*
• tinea
Erythrasma
Contact dermatitis

Clinical features of tinea cruris

- Itchy rash
- More common in young males
- Strong association with tinea pedis (athlete's foot)
- Usually acute onset
- More common in hot months - a summer disease
- More common in physically active people
- Related to chafing in groin (e.g. tight pants, and especially nylon ‘jock straps’)
- Scaling, especially at margin
- Well-defined border (Figure 112.6)



Figure 112.6 Dermogram for tinea cruris

If left untreated the rash may spread, especially to the inner upper thighs, while the scrotum is usually spared. Spread to the buttocks indicates *T. rubrum* infection.

Diagnostic aids

- Skin scrapings should be taken from the scaly area for preparation for microscopy (see p. 1145).
- Wood's light may help the diagnosis, particularly if erythrasma is suspected.

Management of tinea cruris

- Soak the area in a warm bath and dry thoroughly.
- Apply an imidazole topical preparation (e.g. miconazole or clotrimazole cream); rub in a thin layer bd for 3 - 4 weeks on the rash and 2 cm beyond the border.
- Apply tolnaft ate dusting powder bd when almost healed to prevent recurrence.
- If itch is severe, a mild topical hydrocortisone preparation (additional) can be used.[5]
- If weeping: Burow's solution 1 in 40 compresses to dry the area.
- For persistent or recurrent eruption, use oral griseofulvin for 6 - 8 weeks, or terbinafine for 2 - 4 weeks.

Candida intertrigo

Candida albicians superinfects a simple intertrigo and tends to affect obese or bedridden patients, especially if incontinent.[6]

Clinical features

- Occurs equally in men and women
- Erythematous scaly rash in groin
- Less well-defined margin than tinea (Figure 112.7)


indefinite border with satellite macular lesions at the edge

Figure 112.7 Dermogram for candidiasis of crural area

- Associated satellite lesions
- Yeast may be seen on microscopy

Treatment

- Treat underlying problem.
- Apply an imidazole preparation such as miconazole 2% or clotrimazole 1%.
- Continue treatment for 2 weeks after symptoms resolve.
- Use Burow's solution 1 in 40 compresses to dry a weeping area.
- Use short-term hydrocortisone cream for itch or inflammation (long-term aggravates the problem).

Erythrasma

Erythrasma, a common and widespread chronic superficial skin infection, is caused by the bacterium *Corynebacterium minutissimum*, which can be diagnosed by coral pink fluorescence on Wood's light examination. Itch is not a feature.

Clinical features

- Superficial reddish-brown scaly patches
- Enlarges peripherally
- Mild infection but tends to chronicity if untreated
- Coral pink fluorescence with Wood's light
- Common sites: groin, axillae, submammary, toe webs (Figure 112.8)



Figure 112.8 Typical sites of erythrasma

Treatment

- Erythromycin or tetracycline (oral)
- Topical imidazole

Asteatotic eczema (winter itch)

This often unrecognised problem, which can be very itchy, is a disorder of the elderly. It is a form of eczema that typically occurs on the legs of the elderly (Figure 112.9), especially if they are subjected to considerable scrubbing and bathing. Other predisposing factors include low humidity (winter, central heating) and diuretics. The problem may be part of a malabsorption state.



Figure 112.9 Asteatotic eczema (winter itch) showing the ‘crazy paving’ pattern with a pruritic scaling erythematous eruption on the legs of a 74 year old man

Clinical features

- Dry skin
- Fine scaling and red superficial cracking
- ‘Crazy paving’ appearance
- Occurs on legs, especially shins
- Also occurs on thighs, arms and trunk

Treatment

- Avoid scrubbing with soaps
- Use aqueous cream and a soap substitute
- Apply topical steroid diluted in white soft paraffin

‘Golfer's vasculitis’ (summer leg rash)[7]

This is a term used to describe an erythematous pruritic rash that appears on the legs after prolonged exercise such as golf or hiking, usually during summer months.

The rash is a red blotchy flat to slightly raised eruption on the lower leg. It is more common over age 50. It usually clears spontaneously within 3 days.

Grover's disease

Also known as transient acantholytic dermatosis, Grover's disease produces small, firm, intensely pruritic, reddish-brown, warty papules with minimal scale, mainly on the upper trunk. It usually occurs in middle-aged to elderly men (typically 70 - 80 years). Trigger factors include heat, sweating, fever, occlusion, especially on photodamaged skin. Diagnosis is by biopsy. Treatment is to relieve the itch until spontaneous resolution occurs. Effective treatments include topical (preferable) or oral corticosteroids and ultraviolet light.[3]

Lichen simplex

Lichenification is a form of dermatitis caused by repeated scratching or rubbing, which results in epidermal thickening. Lichen simplex is the term used when no primary dermatological cause can be found.

Urticaria

Urticaria is a common condition that mainly affects the dermis. It can be classified as acute (minutes to weeks) or chronic (lasting more than 2 months). It can also be classified as diffuse wheal-like or papular (hives).

The three characteristic features of diffuse urticaria are:

- transient erythema
- transient oedema
- transient itch

Classification according to site

1. *Superficial*: affecting superficial dermis = urticaria; occurs anywhere on body, especially the limbs and trunk.
2. *Deep*: affecting subcutaneous tissue = angiooedema; occurs anywhere but especially periorbital region, lips and neck.

Checklist of causes[3]

- Allergies (acute allergic urticaria is dramatic and potentially very serious)

  - azo dyes
  - drugs: penicillin and other antibiotics
  - foods: eggs, fish, cheese, tomatoes, others
- Infections: bacteria, parasites, protozoa, yeasts, viruses
- Pharmacological
  - drugs: penicillin, aspirin, codeine
  - foods: fish, shellfish, nuts, strawberries, chocolate, artificial food colourings, wheat, soy beans
  - plants: nettles, others
- Systemic lupus erythematosus
- Physical
  - cholinergic: response to sweating induced by exercise and heat (e.g. young athletes)
  - heat, cold, sunlight
- Insect stings: bees, wasps, jellyfish, mosquitoes
- Pregnancy (last trimester), other hormonal
- Unknown (idiopathic) - 80%; possible psychological factors

Investigations

- Full blood examination - look for eosinophilia of parasites
- ANF and DNA binding tests - consider SLE
- Challenge tests

Treatment

- Avoid any identifiable causes.
- Avoid salicylates and related food preparations (e.g. tartrazine).
- Consider elimination diets.
- Use oral antihistamines (e.g. cyproheptadine) or a non-sedating one (e.g. cetirizine, loratadine, fexofenadine).
- Consider adding a H_2 antagonist, e.g. ranitidine 150 mg bd.
- Give short course of systemic corticosteroids if severe (e.g. prednisolone 50 mg once daily for 10 - 14 days).[9]
- Use topical soothing preparation if relatively localised (e.g. crotamiton 10%, or phenol 1% in oily calamine).
- Lukewarm baths with Pinetarsol or similar soothing bath oil.

Flea bites

Fleas (Figure 112.10) cause itchy erythematous maculopapular lesions. They are usually multiple or grouped in clusters, occuring typically on the arms, forearms, leg and waist (where clothing is tight). Treat the source of infestation.



Figure 112.10 Flea

Bed bug bites[8]

The common bed bug (*Cimex lectularis*, Figure 112.11) is now a major problem related to international travel. It travels in baggage and is widely distributed in hotels, motels and backpacker accomodation. Clinically bites are usually seen in children and teenagers. The presentation is a linear group of three or more bites (along the line of superficial blood vessels), which are extremely itchy. They appear as maculopapular red lesions with possible wheals. The lesions are commonly found on the neck, shoulders, arms, torso and legs. A bed bug infestation can be diagnosed by identification of specimens collected from the infested residence.



Figure 112.11 Bed bug

Management

- Clean the lesions.
- Apply a corticosteroid ointment.
- A simple anti-pruritic agent may suffice.
- Call in a licensed pest controller.

Control treatment is basically directed towards applying insecticides to the crevices in walls and furniture.

WHEN TO REFER

- Lichen planus
- Dermatitis herpetiformis
- Norwegian scabies

REFERENCES

1. Walsh TD. *Symptom Control*. Oxford: Blackwell Scientific Publications, 1989: 286 - 94.
2. Hunter JAA, Savin JA, Dahl MV. *Clinical Dermatology* (3rd edn). Oxford: Blackwell Scientific Publications, 2002: 291.

3. Fry L et al. *Illustrated Encyclopedia of Dermatology*. Lancaster: MTP Press, 1981: 313 - 15, 485 - 8.
4. Marley J (Chair). *Therapeutic Guidelines: Dermatology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2004: 224 - 5.
5. Gin D. Tinea infection. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 512.
6. Cowen P. Candidiasis, cutaneous. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 89 - 91.
7. Australas J. Golfers Vasculitis: An Unusual Presentation. Dermatology, 2005; 46: 11 - 14.
8. Bed bugs: < http://www.bedbug.org.au >
9. Levine M, Lexchin J, Pellizzari R. *Drugs of Choice: a Formulary for General Practice*. Ottawa: Canadian Medical Association, 1995: 15 - 33.

CHAPTER 113. COMMON SKIN PROBLEMS

The power of making a correct diagnosis is the key to all success in the treatment of skin diseases; without this faculty, the physician can never be a thorough dermatologist, and therapeutics at once cease to hold their proper position, and become empirical.

LOUIS A. DUHRING (1845 - 1913)

COMMON SKIN PROBLEMS

Skin disorders are very common in general practice. According to Fry, 13% of the population (UK)[1] will be treated for skin disorders each year, with the most common conditions being acute infections, dermatitis (eczema), warts, urticaria, pruritus, acne and psoriasis. According to Bridges-Webb,[2] 13% of the problems managed in Australian general practice will be skin problems, with the most common problems (in order of frequency) being dermatitis/eczema, solar/hyper keratosis, laceration, malignant neoplasm, bruise, skin ulcer, dermatophytosis, boil/carbuncle, naevi and warts.

This chapter focuses on the common dermatoses.

DERMATITIS/ECZEMA

The terms ‘dermatitis’ and ‘eczema’ are synonymous and denote an inflammatory epidermal rash, acute or chronic, characterised by vesicles (acute stage), redness, weeping, oozing, crusting, scaling and itch.

Dermatitis can be divided into exogenous causes (allergic contact, primary irritant contact, photoallergic and phototoxic) and endogenous, which implies all forms of dermatitis not directly related to external causative factors. Endogenous types are atopic, nummular (discoid), vesicular hand/foot (dyshidrotic), pityriasis alba, lichen simplex chronicus and seborrhoeic.

Dermatitis is really a multifactorial disorder of uncertain aetiology that is not always associated with allergy. It can have a wide range of presentations.

The meaning of atopy

The term ‘atopic’ refers to a hereditary background or tendency to develop one or more of a group of conditions, such as allergic rhinitis, asthma, eczema, skin sensitivities and urticaria. It is not synonymous with allergy.

An estimated 10% of the population are atopics, with allergic rhinitis being the most common manifestation.[3]

ATOPIC DERMATITIS

Features of classic atopic dermatitis:[4]

- itch
- usually a family history of atopy
- about 3% of infants are affected, signs appearing between 3 months and 2 years.
- often known trigger factors (Table 113.1) are evident

Table 113.1 Trigger factors for atopic dermatitis

Sweating
Sand (e.g. in sandpits)
Extremes of hot and cold
Rapid temperature changes

Soap and water/frequent washing, especially in winter

Chlorinated water

Bubble baths

Infection (viral, bacterial, fungal)

Allergy

Stress/emotional factors

Skin irritants

- wool (e.g. sheepskin covers)
- brushed nylon or silk clothing
- chemical disinfectants
- detergents
- petrochemical products
- pollens

Scratching and rubbing

Perfumes

Poor general health

Foodstuffs (consider)

- cow's milk
- beef
- chicken
- nuts
- eggs
- food colourants
- oranges
- wheat

Note: The relationship to food is controversial and uncertain. Rast testing is misleading. Consider eliminating the foodstuffs and reintroducing one at a time. If sensitive, the children go bright red a few minutes after feeding.

- dust mite allergy is not always obvious, especially for periorbital rash
- lichenification may occur with chronic atopic dermatitis
- flexures usually involved Figure 113.1



Figure 113.1 Atopic dermatitis in the flexures of the knees - a typical location

- dryness is usually a feature

Criteria for diagnosis

- Itch
- Typical morphology and distribution
- Dry skin
- History of atopy
- Chronic relapsing dermatitis

Distribution

The typical distribution of atopic dermatitis changes as the patient grows older. In infants the rash appears typically on the cheeks of the face, the folds of the neck and scalp. And exknsor surfaces of the limits It may then spread to the Hexanes of limbs and groins. The change from infancy through to adulthood is presented in Figure 113.2.





Figure 113.2 Relative distribution of atopic dermatitis in (a) infants, (b) children and (c) typical distibution of atopic dermatits in adults

During childhood a drier and thicker rash tends to develop in the cubital and popliteal fossae and on the hands and feet, which may be dry, itchy, fissured and painful. The face often cleans.

Prognosis

It is generally correct that children tend to 'grow out of' the problem as the function of their oil and sweat glands matures. The skin becomes less dry, less overheated and irritable.[5] About 60% of patients have virtually normal skin by 6 years and 90% by puberty.[5]

Treatment

Advice to parents of affected children:

- Avoid soap and perfumed products. Use a bland bath oil in the bath and aqueous cream as a soap substitute.
- Older children should have short, tepid showers.
- Avoid rubbing and scratching - use gauze bandages with hand splints for infants.
- Avoid overheating, particularly at night.
- Avoid sudden changes of temperature, especially those that cause sweating.
- Wear light, soft, loose clothes, preferably made of cotton. Cotton clothing should be worn next to the skin.
- Avoid wool next to the skin.
- Avoid dusty conditions and sand, especially sandpits.
- Avoid contact with people with 'sores', especially herpes.
- Keep the skin moist.
- Consider dust mite strategies:
- dust mite covers (premium grade) for bedding
- wash linen in hot water > 55°
- consider replacing fabric on chairs and carpet

Education and reassurance

Explanation, reassurance and support are very important. Emphasise that atopic dermatitis is a superficial disorder and will not scar or disfigure under normal circumstances. The child should be treated normally in every respect. Coun-selling is indicated where family stress and psychological factors are contributing to the problem.

Medication

Note: Corticosteroid creams (for acute phase) and ointments (for chronic phase) are the basis of treatment.

Mild atopic dermatitis

- Soap substitutes, such as aqueous cream
- Emollients (choose from)
  - aqueous cream
  - sorbolene alone or with 10% glycerol
  - paraffin creams (e.g. Dermeze) - good in infants
  - bath oils (e.g. Alpha-Keri)
  - moisturising lotions (e.g. QV) in summer
- 1% hydrocortisone ointment (if not responding to above)
  - short term for flares

Moderate atopic dermatitis

- As for mild
- Topical corticosteroids (twice daily)
  - vital for active areas
  - moderate strength (e.g. fluorinated) to trunk and limbs
  - weaker strength (e.g. 1% hydrocortisone) to face and flexures
  - use in cyclic fashion for chronic cases (e.g. 10 days on, 4 days off)
- Non-steroidal alternative: pimecrolimus (Elidel) cream bd; best used when eczema
- flares - then cease Oral antihistamines at night for itch

Severe dermatitis

- As for mild and moderate eczema
- Potent topical corticosteroids to worse areas (consider occlusive dressings)
- Consider hospitalisation
- Systemic corticosteroids (rarely used)
- Allergy assessment if unresponsive

Weeping dermatitis (an acute phase)

This often has crusts due to exudate. Burow's solution diluted to 1 in 20 or 1 in 10 can bed to soak affected areas. Saline (1 teaspoon to 500 mL water) dressings can also be used: soak old sheets till damp and lay on the areas.

General points of dermatitis management

Acute weeping	→ wet dressings (saline or Burow's)
Acute	→ Creams
Chronic	→ ointments, with or without occlusion
Lichenified	→ ointments under occlusion
Infection	→ antibiotics, eg mupirocin 2% topaid o dralky usesponme to topical alone.

OTHER TYPES OF ATOPIC DERMATITIS

Nummular (discoid) eczema

- Chronic, red, coin-shaped plaques
- Crusted, scaling and itchy
- Mainly on the legs, also buttocks and trunk
- Often symmetrical
- Common in middle-aged patients
- May be related to stress
- Tends to persist for months

Treatment as for classic atopic dermatitis.

Pityriasis alba

- These are white patches on the face of children and adolescents
- Very common mild condition
- More common around the mouth and on cheeks
- Can occur on the neck and upper limbs, occasionally on trunk
- It is a subacute form of atopic dermatitis
- Full repigmentation occurs eventually

Treatment

- Reassurance
- Simple emollients
- Restrict use of soap and washing
- May prescribe hydrocortisone ointment (rarely necessary)

Lichen simplex chronicus

- Circumscribed thick plaques of lichenification
- Often a feature of atopic dermatitis
- Caused by repeated rubbing and scratching of previously normal skin
- Due to chronic itch of unknown cause
- At sites within reach of fingers (e.g. neck, forearms, thighs, vulva, heels, fingers)
- May arise from habit

Treatment

- Explanation
- Refrain from scratching
- Fluorinated corticosteroid ointment with plastic occlusion

Dyshidrotic dermatitis (pompholyx)

- Typically aged 20 - 40 years
- Itching vesicles on fingers
- May be larger vesicles on palms and soles
- Commonly affects sides of digits and palms
- Lasts a few weeks
- Tends to recur
- Possibly related to stress

- Often triggered by high humidity

Treatment

- As for atopic dermatitis
- Potent fluorinated corticosteroids
- topically - use under occlusion (e.g. damp cotton gloves)
- Oral corticosteroids may be necessary
- Wet dressings/ soaks & sertere

Asteatotic dermatitis

This is the common very itchy dermatitis that occurs in the elderly, especially in the winter with a dry 'crazy paving' pattern, especially on the legs (see page 1147).

'Asteatotic' means without moisture.



Figure 113.3 Pompholyx showing the typical vesicular dermatitis along the borders of the fingers. Look for associated inflammatory tinea of the feet which can precipitate it. *Photo courtesy Robin Marks*

Cracked (fissured) hands/fingers

This common cause of disability is usually due to dermatitis of the hands, or a very dry skin. It is usually part of the atopic dermatitis problem and it is important to consider allergic contact dermatitis.

Management of hand dermatitis

Hand protection:

- Avoid domestic or occupational duties that involve contact with irritants and detergents.
- Wear protective work gloves; cotton-lined PVC gloves.
- Avoid toilet soaps - use a substitute (e.g. Dove, Neutrogena).
- Cetaphil lotion is a useful soap substitute.
- Apply emollients (e.g. 2% salicylic acid in white soft paraffin at night, or Neutrogena Norwegian formula hand cream).
- If necessary:
  - hydrocortisone 1% ointment (not cream), *or*
  - stronger fluorinated preparation (e.g. Advantan fatty ointment)

Cracked heels

Cracked painful heels are a common problem, especially in adult women. It is a manifestation of very dry skin.

Treatment

- Soak the feet for 15 minutes in warm water containing an oil such as Alpha-Keri or Derma oil.
- Pat dry, then apply an emollient foot cream (e.g. Nutraplus - 10% urea).

Contact dermatitis

Acute contact (exogenous) dermatitis can be either irritant or allergic.

Features

- Itchy, inflamed skin
- Red and swollen
- Papulovesicular
- May be dry and fissured

Irritant contact dermatitis

Caused by primary irritants such as acids, alkalis, detergents, soaps, oils, solvents. This is irritation, not allergy.

Allergic contact dermatitis

Caused by allergens that provoke an allergic reaction in some individuals only - most people can handle the chemicals without undue effect. It is immunologically mediated. This allergic group also includes photo-contact allergens. Contact dermatitis is due to delayed hyper-sensitivity, sometimes with a long time of days to years. It is common in industrial or occupational situations where it usually affects the hands and forearms.

Common allergens

- Ingredients in cosmetics (e.g. perfumes, preservatives)
- Topical antibiotics (e.g. neomycin)
- Topical anaesthetics (e.g. benzocaine)
- Topical antihistamines
- Plants: rhus, grevillea, primula, poison ivy
- Metal salts (e.g. nickel sulphate, chromate)
- Dyes
- Perfumes
- Rubber/latex
- Epoxy resins and glues/acrylates
- Coral

Note:The skin of mangoes cross-reacts with rhus and grevillea.

Clinical features[4]

- Dermatitis ranges from faint erythema to ‘water melon’ face oedema
- Worse in periorbital region, genitalia and hairy skin; least in glabrous skin (e.g. palms and soles)

Note:Can be delayed onset.

- Think of rhus, grevillea or poison ivy allergy if puffy eyes

Diagnostic hallmarks[4]

- Site and shape of lesions suggest contact
- Linear lesions a feature

- Allergic causes may be found by patch testing
- Improvement when off work or on holiday

Diagnosis

- Careful history and examination
- Consider occupation, family history, vacation or travel history, clothes (e.g. wetsuits, new clothes, Lycra bras), topical applications (e.g. medicines, cosmetics)
- Refer to a dermatologist for patch testing

Management

- Determine cause with vigor and remove it
- Wash with water (only) and pat dry (avoid soap)
- If acute with blistering, apply Burow's compresses
- Oral prednisolone for severe cases (start with 25 - 50 mg day for adults for 1-2 week keen gradually veduce over 1-2 week)
- Topical corticosteroid cream
- Oral antibiotics for secondary infection

Chronic phase

Use fragrant free moisturers regularly eg.

- glycerol 10% on sabolene cream *or*
- White soft paraffen *or*
- a proprietary eniblleint hand cream.

SEBORRHOEIC DERMATITIS

Seborrhoeic dermatitis is a very common skin inflammation that usually affects areas abundant in sebaceous glands or intertriginous areas. It is therefore common in hair-bearing areas of the body, especially the scalp and eyebrows. It can also affect scalp the face, neck, axillae and groins, eyelids (blepharitis), external auditory meatus and nasolabial folds. The presternal area is often involved (Figure 113.4).



Figure 113.4 Seborrhoeic dermatitis in an adult showing typical position on the chest

There are two distinct clinical forms: seborrhoeic dermatitis of infancy; and the adult form, which mainly affects young adults.

Studies have indicated that it may be caused by a reaction to the yeast Pityrosporum ovale. It may be associated with HIV infection and Parkinson's disease.

A feature of seborrhoelic dermatitis is that unbile atopic dermatitis it is not itchy. Seborrholic scale is greary and yellowrsh unlike the silvery scale prsoreases.

Principles of treatment

- Topical sulphur, salicylic acid and tar preparations are first-line treatment: they kill the yeast.
- Ketoconazole cream is most effective.
- Ketoconazole shampoo for scalp twice weekly for 4 weeks.
- Topical corticosteroids are useful for inflammation and pruritus and best used in combination. Avoid corticosteroids if possible.

Seborrhoeic dermatitis of infancy

This rash may be known as 'cradle cap' if it affects the scalp, or nappy rash/diaper dermatitis if it involves the napkin area.

It can be difficult to differentiate the rash from that of atopic dermatitis but seborrhoeic dermatitis tends to appear very early (before atopic dermatitis), even in the first month of life and mostly within the first 3 months, when androgen activity is most prevalent. The different features are summarised in Table 113.2 and the distribution is presented in Figure 113.5

Table 113.2 Differential diagnosis of seborrhoeic dermatitis and atopic dermatitis in infancy

	Seborrhoeic dermatitis	**Atopic dermatitis (eczema)**
Age of onset	mainly within first 3 months	usually after 2 months
Itchiness	nil or mild	usually severe
Distribution	scalp, cheeks, folds of neck, axillae, folds of elbows and knees	starts on face elbow and knee flexures
Typical features	cradle cap red and yellow greasy scale	vesicular and weeping becomes dry and cracked
Napkin rash	common prone to infection with Candida	less common
Other features	may become generalised	may become generalised



Figure 113.5 Seborrhoeic dermatitis: typical distribution in infants

Seborrhoeic dermatitis usually appears as red patches or blotches with areas of scaling. This becomes redder when the baby cries or gets hot. Cradle cap may appear in the scalp. A flaky, scurf-like dandruff appears first, and then a yellowish, greasy, scaly crust forms. This scurf is usually associated with reddening of the skin.

The dermatitis can become infected, especially in the napkin area, and this may be difficult to treat. If untreated, it often spreads to many areas of the body. It is said that cradle cap and nappy rash 'may meet in the middle'.

Treatment

Simple basic methods:

- Keep areas dry and clean.
- Bathe in warm water, pat areas dry with a soft cloth.
- Keep skin exposed to air as much as possible.
- Avoid toilet soaps for washing (use emulsifying ointment or Cetaphil lotion).
- Rub scales of cradle cap gently with baby oil, then wash away loose scales.
- Change wet or soiled nappies often.
- For mild areas on body, apply a thin smear of zinc cream.

Medication

Scalp

Infants:

- 1% sulphur and 1% salicylic acid in sorbolene cream
  - apply overnight to scalp, shampoo off next day with a mild shampoo
  - use 3 times a week
- Egozite cradle cap lotion (6% salicylic acid)
  Older children and adults:
  - zinc pyrithione 1% or selenium sulphide 2.5% shampoo *or*
  - ketoconazole 1 - 2% or miconazole 2% shampoo

Face, flexures and trunk

select from:

- Ketoconazole 2% cream, once or twice daily

- 2% sulphur and salicylic acid in aqueous or sorbolene cream
- hydrocortisone 1% (irritation on face and flexures)
- betamethasone 0.02 - 0.05% (if severe irritation on trunk)
- desonide 0.0. 5% lotion, bd tds for faceleyelids and for weeping cream.

Napkin area

- Mix equal parts 1% hydrocortisone with nystatin or ketoconazole 2% or clotrimazole 1% cream (or use Hydrozole cream)

Prognosis

Most children are clear by 18 months (rare after 2 years).

Adult seborrhoeic dermatitis

Clinical features:

- any age from teenage onwards
- the head is a common area: scalp and ears, face, eyebrows, eyelids (blepharitis), nasolabial folds (Figure 113.6)


scalp, especially anterior margin
eyebrows
eyelids (blepharitis)
wings of nose
nasolabial folds
external auditory meatus (otitis externa)

Figure 113.6 Seborrhoeic dermatitis: facial distribution in adults

- other areas: centre of chest, centre of back, scapular area, intertriginous areas, especially perianal (Figure 113.7)


external auditory meatus
scalp → dandruff
eyebrows
eyelid margins
behind ears
nasolabial folds
beard area
presternal
interscapular
axilla
submammary intertrigo
umbilicus
groin
perianal area (pruritus ani)

Figure 113.7 Seborrhoeic dermatitis: possible distribution in adults

- red rash with yellow greasy scale
- secondary candidiasis infection common in flexures
- dandruff a feature of scalp area
- worse with stress and fatigue
- It is a chronic recurring condition

Treatment

Scalp

Options:

1. salicylic acid 2% + sulphur 2% in aqueous cream overnight - shampoo off next day using selenium sulphide or zinc pyrithione shampoo (apply 3 times a week)
2. ketoconazole shampoo; immediately after using medicated shampoo (leave 5 minutes), twice weekly
3. betamethasone lotion to scalp (if very itchy)

Face and body

- Wash regularly using bland soap
- Salicylic acid 2% + sulphur 2% (° tar) in aqueous cream *or*
- Ketoconazole 2% cream (very effective); apply once daily for 4 weeks
- Hydrocortisone 1% bd (if inflamed and pruritic)

Note: Oral antifungaloare not recommanded at all.

PSORIASIS

Psoriasis is a chronic skin disorder of unknown aetiology which affects 2 - 4% of the population. It appears most often between the ages of 10 and 30 years, although its onset can occur any time from infancy to old age. It has a familial predisposition although its mode of inheritance is debatable. If one parent is affected there is a 25% chance of developing it; this rises to 65% if both parents are affected.[3]



Figure 113.8 Psoriasis showing the typically raised pink plaques surmounted with a silvery scale

Posriasis is now regarde a disorder involving activa of helper Talls in the SK cyto kines are them relea and cause skin cells multiply faster, leade to theckening of the skin and overocale the new biological of intervene in this proee.

Capillary dilatation explains the redness.

Types of psoriasis	Differential diagnosis
Infantile	Seborrhoeic dermatitis, atopic dermatitis
Plaque (commonest)	Seborrhoeic dermatitis, discoid eczema, solar keratoses, Bowen's disorder
Guttate	Pityriasis rosea, secondary syphilis, drug eruption
Flexural	Tinea, Candida intertrigo, seborrhoeic dermatitis
Scalp (sebopsoriasis)	Seborrhoeic dermatitis, tinea capitis
Nail	Tinea, idiopathic onycholysis
Pustular (palmoplantar)	Tinea, infected eczema
Exfoliative	Severe seborrhoeic dermatitis

Factors that may worsen or precipitate psoriasis

- Infection, especially group A Streptococcus
- Trauma or other physical stress
- Emotional stress
- Sunburn
- Puberty/menopause
- Drugs

  - antimalarials (e.g. chloroquine)
  - beta-blockers
  - lithium
  - NSAIDs
  - oral contraceptives

The typical patient

- Older teenager or young adult
- Possible family history
- Onset may follow stress, illness or injury
- Rash may appear on areas of minor trauma - the Koebner phenomenon
- Rash improves on exposure to sun but worse with sunburn
- Worse in winter
- Itching not a feature
- Lesions are most unlikely to appear on the face

Arthropathy

About 5% can develop a painful arthropathy (fingers, toes or a large joint) or a spondylo-arthropathy (sacroiliitis).[7]

The rash

The appearance depends on the site affected. The commonest form is plaque psoriasis, which begins with red lesions that enlarge and develop silvery scaling. The commonest sites are the extensor surfaces of the elbows and knees; then the scalp, sacral areas, genitals and nails are affected (Figure 113.9).



Figure 113.9 Psoriasis: typical skin distribution

Diagnosis

Psoriasis is a clinical diagnosis but biopsy may be needed for confirmation. No laboratory test (including blood testing) is available.

Principles of management

While realising there is no cure for psoriasis the aim of treatment is to relieve discomfort, slow down the rapid skin cell division and work in consultation with a specialist to achieve these aims.[7]

- Provide education, reassurance and support.
- Promote general measures such as rest, and holidays, preferably in the sun.
- Advise prevention, including avoidance of skin damage and stress if possible.
- Tailor treatment (including referral) according to the degree of severity and extent of the disease.

Treatment options[5,8]

1. Topical therapy
    - dithranol (anthralin): concentrations range from 0.05 to 2% (max.)
        - effective on difficult thick patches
          Note:Facts about dithranol
        - Stains light-coloured hair purple so don't use on scalp.
        - Start in low concentration and build up according to tolerance and response.
        - Use in strengths 0.05% (children), 0.1%, 0.25%, 0.5% and 2.0% (max.).
        - Can use a higher strength of 0.25% to start but only for short contact therapy (30 minutes before shower).
        - Irritates skin, causing a burning sensation.
        - Don't use it on face, genitalia or flexures.
    - tar preparations: effective but also messy to use
    - corticosteroids: the mainstay of therapy for small plaques; use 1% hydrocortisone on more sensitive areas (genitals, groin, axillae, face) but use stronger types elsewhere (e.g. betamethasone dipropionate)
      Note:Avoid using corticosteroids alone as they promote instability - best to combine with tar or calcipotriol.
    - calcipotriol ointment
        - easy to use
        - apply twice daily for a minimum of 6 weeks
        - tends to irritate face and flexures
        - caution with hypercalcaemia
        - wash hands after use
        - used for chronic stable plaques (not for severe extensive rash)
    - bland preparations and emollients: these can be used for dryness, scaling and itching - liquor picis carbonis (LPC) and menthol (or salicylic acid) in sorbolene base

    Recommended topical regimens

    - Combined method for mild to moderate psoriasis (a good starting method)
        - Dithranol 0.1%, salicylic acid 3%, LPC (tar) 10% in white soft paraffin (preferable) or sorbolene cream.
        - Leave overnight (warn about dithranol stains - use old pyjamas and sheets).
        - Review in 3 weeks then gradually increase strength of dithranol to 0.25%, then 0.5% then 1%. Can cut down frequency to 2 - 3 times per week.
        - Shower in the morning and then apply a topical fluorinated corticosteroid.
    - Short contact anthralin therapy (SCAT method)
        - apply dithranol 2% in sorbolene cream
        - wash off in shower after 10 - 15 minutes
    - General adjunctive therapy

    - tar baths (e.g. Pinetarsol or Polytar)
    - tar shampoo (e.g. Polytar, Ionil-T)
    - sunlight (in moderation)

2. Systemic therapy
    - methotrexate: can have dramatic results in severe cases
    - cyclosporin (hospital use only)
    - corticosteroids (for erythrodermic psoriasis only) - oral use may unstabilise
    - psoriasis on withdrawal
    - acitretin - a vitamin A derivative effective in severe intractable psoriasis (never use in females of child-bearing age)
3. Biological agents this new day of drup are directed at the T-call dysfunctum, ie immune respouse nowdipens.
4. Physical therapy
    - phototherapy (ultraviolet light, UV-B): needs careful supervision
    - UV-B plus coal tar (Goeckerman regimen): reserved for severe psoriasis;
        a. tar bath
        b. UV-B
        c. 2 - 5% crude coal tar
    - Ingram regime: also for severe psoriasis (best as inpatient)
        a. tar bath
        b. UV-B
        c. 0.1% - 0.5% dithranol in Lassar's paste
    - photochemotherapy (PUVA = psoralen + UV-A) - reserved for non-responders to UV-B treatment or other therapies. A variation is REPUVA = retinoid + psoralen + UV-A
    - intralesional corticosteroids - an excellent and effective treatment for isolated small or moderate-sized plaques that can be readily given by the family doctor

Method of injection

Mix equal parts of triamcinolone acetonide 10 mg/mL, or similar corticosteroid, and plain LA or N saline and, using a 25 - gauge or 23-gauge needle, infiltrate the psoriatic plaque intradermally to cover virtually all the plaque. A small plaque can be covered by a single insertion while a larger plaque may require separate insertions (Figure 113.10).



Figure 113.10 Intralesional corticosteroid injection technique for psoriatic plaque (requiring double injection - small plaques need only one infiltration)

Other practice tips

Chronic stable plaque psoriasis

- Apply stronger fluorinated corticosteroid (II - III class) cover with Duoderm Thin and leave for 7 days (other occlusive dressings can be used)
  plus
- Tar (as an alternative to dithranol), e.g. crude coal tar 1 - 4% plus salicylic acid 3 - 5% in sorbolene cream. It can be left on overnight and washed off in the morning *or*
- Calcipotriol 0.005% ointment or cream, apply bd

Resistant localised plaque

- Intralesional injection with corticosteroid diluted 50:50 with N saline or local anaesthetic

Maintenance for milder stabilised plaque psoriasis

- Salicyclic acid 3%, LPC 8% in sorbolene

Scalp psoriasis

- Apply 20% urea cream for 4 days, then
- Tar preparations for the scalp, steroid lotions or 5 - 10% salicylic acid in mineral oil

Genital psoriasis

- Apply non fluorinated corticosteroids

Flexural psoriasis

Note that fissuring (e.g. inframammary, natal cleft) is a feature.

- Apply topical steroids - fluorinated if possible.

NAPPY RASH[5]

Nappy rash (or diaper dermatitis) is an in-flammatory contact dermatitis occurring in the napkin area and can be a common presentation of mild or moderate underlying skin disease. It is found in children up to 2 years old and has a peak incidence from 9 - 12 months.[9]

Most children will develop nappy rash at some stage of infancy with an estimated 50% having it to a significant extent. The commonest type is irritant dermatitis, but consider also:

- Candida albicans (invariably present although often not obvious)
- seborrhoeic dermatitis
- atopic dermatitis
- psoriasis

Causes of nappy rash

The main predisposing factor in all types is dampness due to urine and faeces. It is far less common since the use of disposable nappes. Other causes or aggravating factors are:

- a tendency of the baby to eczema
- a tendency of the baby to seborrhoeic dermatitis
- infection, especially Candida (thrush)
- rough-textured nappies
- detergents and other chemicals in nappies
- plastic pants (aggravates wetness)
- excessive washing of the skin with soap
- too much powder over the nappy area (avoid talcum powders)

- teething aggravates (punched out lesions)

Irritant dermatitis

This is a type of contact dermatitis with the erythema and scaling conforming to the napkin area. The flexures are usually spared. It is related to the activity of faecal proteases and lipases and probably not to the activity of ammonia (from urea) as previously promoted. The problem can vary from mild erythema to a severe blistering eruption with ulceration. Ultra-bsorbent disposable nappies appear to be better than cloth nappies.[10] Diarrhoea is a causative factor of irritant dermatitis. If the eruption extends further than the points of contact with the nappy an underlying skin disease such as seborrhoeic or atopic dermatitis must be suspected. Psoriasis always involves the skin folds.

Seborrhoeic dermatitis

This affects mainly the flexures of the natal cleft and groin. It is important to look for evidence of seborrhoeic dermatitis elsewhere, such as cradle cap and lesions on the face and axillae.

Atopic dermatitis

Atopic dermatitis can involve the napkin area. Pruritus is a feature and the child may be observed scratching the area. There may be evidence of atopic dermatitis elsewhere, such as on the face.

Candidiasis (Monilia nappy rash)

Superinfection of intertrigo or napkin derm-atitis will result in a diffuse, red, raw, shiny rash that will involve the flexures and extend beyond the napkin area as ‘satellite lesions’. Candida tends to invade the skin folds and the foreskin of male babies.

Uncommon causes

Psoriatic nappy rash

This presents as a non scaling eruption, pri-marily on the napkin area, but can extend to the flexures, trunk and limbs. The edge of the rash is sharply demarcated. The typical psoriatic scale is absent. It tends to occur in the first weeks of life. There is usually a family history.

Infections

Bacterial infections to consider include staphylococcal folliculitis, impetigo and perianal streptococcal dermatitis. Culture of the lesion will reveal the cause.

Impetigo

If there is Staphylococcus superinfection, bullae and pus-filled blisters will be present.

Histiocytosis X (Letterer-Siwe syndrome)

There is a similar rash to seborrhoeic dermatitis but the lesions are purpuric. In this serious syndrome the child is very ill and usually lymphadenopathy and hepatosplenomegaly may be found.

Zinc deficiency

May be more common than realised.

Management of nappy rash

Basic care (instructions to patients):

1. Keep the area dry. Change wet or soiled nappies frequently and as soon as you notice them. Disposable nappies are helpful.

2. After changing, gently remove any urine or moisture with diluted sorbolene cream or warm water.
3. Wash gently with warm water, pat dry (do not rub) and then apply any prescribed cream or ointment to help heal and protect the area. Vaseline or zinc cream applied lightly will do. Stoma adhesive powder is an excellent protector.
4. Expose the bare skin to fresh air wherever possible. Leave the nappy off several times a day, especially if the rash is severe.
5. Do not wash in soap or bath too often - once or twice a week is enough.
6. Avoid powder and plastic pants.
7. Use special soft nappy liners that help protect the sensitive skin.
8. Thoroughly rinse out any bleach or disinfectants.

Medical treatment

Some principles:

- The cornerstone of treatment is prevention.
- Emollients should be used to keep skin lubricated (e.g. mixture of zinc oxide and castor oil or Vaseline).
- A mild topical corticosteroid is the treatment of choice.
- It is usual to add an antifungal agent.
- Be careful of excessive use of corticosteroids, especially fluorinated steroids.
- If infection is suspected, confirm by swab or skin scraping.

Treatment

Atopic dermatitis	1% hydrocortisone
Seborrhoeic dermatitis	1% hydrocortisone + ketoconazole cream
Candida	topical nystatin at each nappy change
Widespread (Candida present)	1% hydrocortisone cream mixed in equal quantities with nystatin or clotrimazole cream (apply qid after changes)
Psoriatic dermatitis	tar preparation, or 1% hydrocortisone
Impetigo	topical mupirocin; oral antibiotics if severe

FACIAL RASHES

Common facial skin disorders include acne, rosacea, perioral dermatitis and seborrhoeic dermatitis. These conditions must be dis-tinguished from lupus erythematosis (discold LE is more common).

ACNE

Acne is inflammation of the sebaceous (oil) glands of the skin(Figure 113.11). At first there is excessive sebum production due to the action of androgen. These glands become blocked (blackheads and whiteheads) due to increased keratinisation of the sebaceous duct. The bacteria in the sebum produce lipases with the resultant free fatty acids, thus provoking inflammation.



Figure 113.11 Facial acne showing typical nodulocystic acne, a distressing problem which has been improved greatly by the development of isotretinoin

Types of acne[11]

- *Infantile*. Occurs in the first few months of life, mainly on the face. Affects mainly boys and is a self-limiting minor problem. Reassurance only is required in most cases. Some are severe and may scar.
- *Adolescent*. The most common type, occurring around puberty. Acne is rare under 10 years; ages 13 - 16 years are commonest and it is worse in males aged 18 - 19 years. It is slightly less common in girls and worse around 14 years with premenstrual exacerbations.
- *Cosmetica*. In females, associated with the prolonged use of skin care products (e.g. moisturiser, foundation cream and heavy make-up).
- *Oil*. Occurs mainly on the legs of workers exposed to petroleum products.
- *Drug induced*. Especially steroide.

Taking a history

Enquire about use of skin preparations - therapeutic or cosmetic, exposure to oils, possible diet relationships and drug intake.

Drugs that aggravate acne:[11]

- corticosteroids
- chloral hydrate
- iodides or bromides
- lithium
- antiepileptics (e.g. phenytoin)
- quinine
- oral contraceptives

Management

Support and counselling

Adolescents hate acne; they find it em-barrassing and require the sympathetic care and support not only of their doctor but also of their family. It should not be dismissed as a minor problem.

Education

People with acne should understand its pathogenesis and be given leaflets with appropriate explanations. Myths must be dispelled:

- It is not a dietary or infectious disorder.
- It is not caused by oily hair or hair touching the forehead.
- Ordinary chemicals (including chlorine in pools) do not make it worse.
- Blackheads are not dirt, and will not dissolve in hot, soapy water.

Reassure the patient that acne usually settles by the age of 20 years.

General factors

- Diet is considered not to be a factor but if there is a causal relationship with any foods (e.g. chocolate), avoid them.
- Special soaps and overscrubbing are unhelpful. Use a normal soap and wash gently.
- Avoid oily or creamy cosmetics and all moisturisers. Use cosmetics sparingly.
- Avoid picking and squeezing blackheads.
- Exercise, hair washing and shampoos are not of proven value.
- Ultraviolet light such as sunlight may help improve acne.

Principles of treatment

1. Unblock the pores (follicular ducts) with keratolytics such as sulphur compounds, salicylic acid (5 - 10%); with benzoyl peroxide (2.5%, 5% or 10%) or retinoic acid (tretinoin) 0.01% gel, cream (0.025%, 0.05% or 0.1%), lotion or liquid; or with adapalene (Differin) cream or gel.
2. Decrease bacteria in the sebum with systemic antibiotics - tetracyclines or erythromycin - or with topical antibiotics such as clindamycin and erythromycin.
3. Decrease sebaceous gland activity with oestrogens, spironolactone, cyproterone acetate, or isotretinoin

Note:Oral isotretinoin is teratogenic.

Recommended treatment regimens

Topical regimens

Suitable for mild to moderate acne

Basic starting regimen (benzoyl peroxide and tretinoin)

- Use tretinoin 0.01% gel or 0.05% cream or isotretinoin 0.05% gel or adapalene 0.1 % cream or gel: apply each night (tretinoins are photosensitive).
- After 2 weeks, add benzoyl peroxide 2.5% or 5% gel or cream once daily (in the morning). That is, maintenance treatment is:
  - tretinoin or isotretinoin at night
  - benzoyl peroxide gel mane
- Maintain for 3 months and review.

Alternative regimens, if recalcitrant

- Clindamycin HCl 600 mg in 60 mL of 70% isopropyl alcohol (e.g. Clindatech). Apply with fingertips twice daily. Alternative bases for clindamycin, especially if the alcohol is too drying or irritating:
  - Cetaphil lotion (Alcon) 100 mL, or
  - Dermatech liquid 100 mL

  Clindamycin is particularly useful for pregnant women and those who cannot tolerate antibiotics or exfoliants.[11]

- Erythromycin 2% gel.
- Azelaic acid, apply bd.
- Adapalene 0.1% cream or gel.

Oral antibiotics

Use for inflammatory acne: (moderate to severe papulopustular) ± trunk involvement

- tetracycline 1 g per day or doxycycline 100 mg per day or minocycline 100 mg per day
- use half this dose if mild
- reduce dosage according to response e.g. tetracycline 250 mg per day or every second day or
- doxycycline 50 mg per day or minocycline 50 mg nocte
- reserve minocycline for resistant cases
- erythromycin or cotrimoxazole are alternatives
- give a minimum 12 week trial; 6 months is standard. Avoid using antibiotics alone.

Other therapies

Severe cystic or recalcitrant acne (specialist care)

- spironolactone
- isotretinoin (Roaccutane)
- dapsone

Females not responding to first-line treatment

- combined oral contraceptive pill, e.g. ethinyloestradiol/cyproterone acetate (Diane-35 ED)

Note:Response to any acne treatment occurs in about 8 weeks or longer.

ROSACEA

Rosacea is a common persistent eruption of unknown aetiology. It is typically chronic and persistent with a fluctuant course.

Clinical features

- Mainly 30 - 50 years
- Usually females of Celtic origin'the course of the celts'
- On forehead, cheeks, nose and chin (Figure 113.13)



Figure 113.13 Rosacea: typical facial distribution

- ‘Flushing and blushing’ (often precedes the rash)
- Fluctuates from day to day
- Periorbital and perioral areas spared
- Vascular changes - erythema and telangiectasia
- Inflammation - papules and pustules

Complications

- Blepharitis
- Conjunctivitis, rarely keratitis and corneal ulcer
- Associated rhinophyma in some cases



Figure 113.12 Rosacea: typical appearance with erythema, papules and pustules

Management

- Apply cool packs if severe.
- Avoid factors that cause facial flushings (e.g. excessive sun exposure, heat, alcohol, spicy foods, hot drinks - tea and coffee).

Systemic antibiotics

- Tetracycline (first choice)
- e.g. 500 - 1000 mg daily, then 250 mg daily when controlled, for a total of 8 - 10 weeks. Repeated for recurrences: avoid maintenance. If using doxycycline, start with 100 mg then 50 mg daily, weane slowly.
- Erythromycin (second choice)
- Metronidazole 200 mg bd for 10 days for resistant cases

Topical agents

For mild erythema and inffammatory lesons:

- 2% sulphur in aqueous cream tds (milder cases) or (for more severe cases) metronidazole gel bd or azelaic cid gel or clinda mycin 15 solution bd or erythromycin 2% gel bd

Hydrocortisone 1% cream is effective, but steroids are best avoided and strong topical steroids should not be used because of severe rebound vascular changes.

Laser therapy

Telangectasia, erythema and rhinophymas respond to laser therapy.

Rhinophyma

This is due to hypertrophy of the nasal sebaceous glands. There is no specific association with alcohol. May be associated with rosacea. Carbon dioxide laser therapy is the treatment of choice. Shave excision is also effective.

PERIORAL DERMATITIS

Clinical features:

- acne-like dermatitis of lower face
- usually young women
- around mouth and on chin, sparing adjacent perioral area (Figure 113.14 and 113.5); also in periorbital skin in men.



Figure 113.14 Perioral dermatitis: typical distribution

- frequently begins at the nasolabial folds
- multiple small red macules and papules
- on a background of erythema and scaling
- burning and irritation
- may be related to pregnancy and oral contraception
- related to the use of creamy cinsmetu products.
- may be related to repeated topical corticosteroid (especially fluorinated) use (may be a rebound on ceasing it)



Figure 113.15 Perioral dermatitis: this eruption (which has been linked with the use of fluorinated steroid creams on the face) frequently begins in the nasolabral fold. Papules and pustules occur on a background of erythema and scaling

Treatment

- Tetracycline, doxycycline or minocycline for 6 - 8 weeks (e.g. doxycycline or minocycline 100 mg daily).
- Discontinue any topical corticosteroid therapy.(femds to flare initially)and all, creamy preparations cleansers moistures and make up.

TINEA

Tinea, or ringworm infections, are caused mainly by three major classes of dermatophytic organisms that have the ability to invade and proliferate in keratin of the skin, nails and hair.

It is most useful to perform skin scrapings and microscopy to look for encroaching septate hyphae.[12] Confirm the diagnosis by fungal culture. Tinea cruris is presented on page 1146.

Tinea pedis (athlete's foot)

Tinea pedis is usually caused by Trichophyton rubrum and is the commonest type of fungal infection in humans. Candida intertrigo and interdigital maceration (alone urihout secondary finca) are important differential diagnoses.

Symptoms

The commonest symptoms are itchiness and foot odour. Sweat and water make the top layer of skin white and soggy. There is scaling, maceration and fissuring of the skin between the fourth and fifth toes and also third and fourth toes

Management

Advice to the patient:

- Keep your feet as clean and dry as possible.
- Carefully dry your feet after bathing and showering.
- After drying your feet, use an antifungal powder, especially between the toes.
- Remove flaky skin from beneath the toes each day with dry tissue paper or gauze.
- Wear light socks made of natural absorbent fibres, such as cotton and wool, to allow better circulation of air and to reduce sweating. Avoid synthetic socks.
- Change your shoes and socks daily.
- If possible, wear open sandals or shoes with porous soles and uppers.
- Go barefoot whenever possible.
- Use thongs in public showers such as at swimming pools (rather than bare feet).

Medication

Clotrimazole 1%, ketoconazole 2% or miconazole 2% cream or lotion, applied after drying, bd or tds for 2 - 3 weeks. If severe and spreading, prescribe oral griseofulvin (see next column) or terbinafine after confirming the diagnosis by fungal culture.

TINEA CORPORIS

Tinea corporis (ringworm infection of the body) is usually caused by *T. rubrum* (60%) or *Microsporum canis*.[11] Related to cats and dogs but a potent source of facial tinea is the guinea pig (can present as pustular folliculitis).

Clinical features

- Spreading circular erythematous lesions (Figure 113.16)



Figure 113.16 Multiple ringworm (tinea corporis): this 12-year-old boy presented with a two-week history of an increasing number of pruritic scaling erythematous lesions on his face, neck and upper chest. Fungal scrapings confirmed the cause as Microsporum canis.*Photo courtesy Robin Marks*

- Slight scaling or vesicles at the advancing edge
- Central areas usually normal
- Mild itch
- May involve hair, feet and nails

Management

- Clotrimazole 1% or miconazole 2% cream or ketoconazole 2% cream, applied bd for 3 - 4 weeks
- Oral terbinafine or griseofulvin if no response or if widespread

TINEA MANUUM

Tinea manuum is ringworm infection of the hand, usually presenting with scaling of the palms and palmar aspects of the fingers. It also commonly presents on the doisu of the hand and is typically undateal.Dif-ferential diagnoses are atopic dermatitis and contact dermatitis of the hands.

Clinical features

- Usually unilateral
- Spreading edge
- Erythematous; fine scaling
- May be associated with tinea pedis

Treatment

- Topical clotrimazole 1%, ketoconazole 2% or miconazole 2% cream for 6 weeks or, if resistant, terbinafine 250 mg or griseofulvin 500 mg daily for 6 weeks.

TINEA CAPITIS

In Australia tinea capitis is usually due to *M. canis* acquired from cats and dogs.

Clinical features

- Usually in children (rare after puberty)

- Patches of partial alopecia
- Scaly patches
- Small broken-off hair shafts
- Hairs fluoresce yellow-green with Wood's light (not invariably, e.g. with Trichophyton tonsurans infection)

Treatment

- Griseofulvin (o) - 6 week course
  Adults: 500 - 1000 mg daily
  Children: 10 mg/kg/day (max. 250 mg)

KERION

Kerion of the scalp and beard area may present like an abscess - tender and fluctuant. Usually occurs on the scalp, face or limbs. A fungal cause is possible if the hairs are plucked out easily and without pain (if painful and stuck, bacterial infection is likely).

Tinea incognito

This is the term used for unrecognised tinea infection due to modification with corticosteroid treatment. The lesions are enlarging and persistent, especially on the groins, hands and face.

The sequence is initial symptomatic relief of itching, stopping the ointment or cream and then relapse.

Tinea unguium (toenails and fingernails)

Refer to Chapter 119.

PITYRIASIS VERSICOLOR (TINEA VERSICOLOR)[13]

Pityriasis versicolor is a superficial yeast infection of the skin (usually on the trunk) caused by Pityrosporum orbiculare (Malassezia furfur). The old name, tinea versicolor, is inappropriate because the problem is not a dermatophyte infection. It occurs worldwide but is more common in tropical and subtropical climates.

There are two distinct presentations:

1. reddish brown, slightly scaly patches on upper trunk
2. hypopigmented area that will not tan, especially in suntanned skin

The term ‘versicolor’ means variable colours.

Clinical features

- Mainly young and middle-aged adults
- Brown on pale skin or white on tanned skin (Figure 113.17)



Figure 113.17 Pityriasis versicolor showing the hypopigmented scaly patches on a suntanned skin mainly affecting the trunk

- Trunk distribution (Figure 113.18)



Figure 113.18 Pityriasis versicolor: typical truncal distribution (corresponding area on back)

- Patches may coalesce
- May involve neck, upper arms, face and groin
- Slight scaling when scratched indicates active infection
- Scales removed by scraping show characteristic short stunted hyphae with spores on microscopy
- often recurrent especially in summer.

Differential diagnosis

Seborrhoeic dermatitis of trunk (more erythematous), pityriasis rosea, vitiligo, pityriasis alba (affects face).

Treatment [5,13]

- Selenium sulphide (Selsun yellow shampoo). Wash area, leaving on for 5 - 10 minutes, then wash off. Do this daily for 2 weeks (at night), then every second day for 2 weeks, then monthly for 12 months. Shampoo scalp twice weekly. and/or

- Clotrimazole, miconazole or ketoconazole cream/lotion nocte for 10 days or
- Terbinafine 1% cream twice daily[12] for 2 weeks Or
- econazole 1% solutions to wet SKIN, leave over night, for 3 mghts.
- Sodium thiosulphate 25% solution bd for 4 weeks (wash off after 10 minutes) or (for persistent or recurrent problems)
- Ketoconazole 200 mg daily for 7 - 10 days or 400 mg single dose[0]
- Itraconazole 200 mg (0) once every 30 to 90 days

Note:

- Ketoconazole may be hepatotoxic. Always perform LFTs first (do not use long-term). Griseofulvin is inappropriate because the rash is not a fungal infection.
- Warn patients that the white patches will take a long time to disappear and that cure does not equate with disappearance.

DRY SKIN

Disorders associated with scaling and roughness of the skin include:

- atopic dermatitis - all types (e.g. pityriasis alba, nummular eczema, asteatotic dermatitis)
- ageing skin
- psoriasis
- ichthyotic disorders
- keratosis pilaris

Itching may be a feature of dry skin (but is not inevitable).

Aggravating factors:

- low humidity (e.g. heaters, air-conditioners)
- frequent immersion in water
- heat/hot water
- toilet soaps
- swimming in chlorinated pools.

Management

- Ensure humidification if there is central heating.
- Reduce bathing.
- Bathe or shower in tepid water.
- Use a soap substitute (e.g. Dove or Neutrogena/Cetaphil lotion).
- Pat dry - avoid vigorous towelling.
- Rub in baby oil after bathing (better than adding oil to the bath).
- Avoid wool next to the skin (wear cotton).
- Use emollients (e.g. Alpha-Keri lotion, QV skin lotion).
- Use moisturisers (e.g. Nutraplus, Calmurid).

SUNBURN

Sunburn is normally caused by UV-B radiation which penetrates the epidermis and superficial dermis, releasing substances such as leucotrienes and histamines, which cause redness and pain. Severe sunburn may develop on relatively dull days because thin clouds filter UV-B poorly.Bewase of soariums and the med day sum.

Clinical presentations:

Minor sunburn	Mild erythema with minimal discomfort for about 3 days.
Moderate	Moderate to severe erythema within a few hours; worse the following day - red, hot and moderately painful. Settles in 3 - 4 days with some desquamation.
Severe	Classic signs of inflammation - redness, heat, pain and swelling. Skin develops vesicles and bullae. Systemic features develop with very severe burns (e.g. fever, headache, nausea, delirium, hypotension). May require IV fluids.

Differential diagnosis

- General photosensitivity: consider drugs (e.g. thiazide diuretics, tetracyclines, sulphonamides, phenothiazines, griseofulvin, NSAIDs, isotretinon)
- Acute systemic lupus erythematosus may present as unexpectedly severe sunburn
- Photocontact dermatitis

Treatment

Hydrocortisone 1% ointment or cream for severe sunburn, early. Repeat in 2 - 3 hours and then the next day. Hydrocortisone is not useful after 24 hours and should be used for unblistered erythematous skin, not on broken skin.

Oral aspirin eases pain. Oil in water baths or bicarbonate of soda paste may help and wet applications such as oily calamine lotions may give relief.

Prevention

Avoid direct exposure to summer sunlight from 10 am to 2 pm (or 11 am to 3 pm daylight saving time). Use natural shade - beware of reflected light from sand or water and light cloud. Use a sunscreen with a minimum of SPF 15 or more. Wear broadbrimmed hats and protective clothing.

PHOTOAGEING/WRINKLES

Prevention

- Stop smoking
- Avoid add, dry and windy condutions
- Avoid exposure to the sun.
- Use an SPF 15 or more sunscreen during the day.
- Avoid soaps with perfuine and alcohol.
- Wash with a ‘neutral’ mild soap (e.g. Neutrogena, max. twice daily) and pat dry.
- Apply a simple moisturiser immediately after the bath.

Treatment (options)

- Optimal nuttition det in fruits and vegetable
- Alpha hydroxy and preparalens eg. Elucent cream.
- Tretinoin (Retin-A) cream
  apply once daily at bedtime (on dry skin)

test for skin irritation by gradual exposure (e.g. 5 minutes at first, wash off, then 15 minutes until it can be left on overnight)

- Topical olive oil.
- Lac-Hydrin (USA)

  12% solution may be effective alternative; other lactic acid preparations may be useful
- Botulinum toxin injections

SWEATING AND ODOUR DISORDERS

Hyperhidrosis (excessive sweating)[14]

This problem is usually idiopathic and prolonged.

Features

- Affects axilla, Groin soles and palms
- Onset usually around puberty
- Tends to improve after age 25 years
- Family history
- ± Bromhidrosis (malodorous perspiration)
- Usually independent of climate
- Exacerbated by stress

Secondany /Pathological causes

- Fever
- Thyrotoxicosis
- Acromegaly
- Diabetes mellitus
- Phaeochromocytoma
- Drugs: alcohol/narcolic withdrawal
- Some neurological conditions (e.g. Parkinson's)

Treatment[5]

- Reduce caffeine intake
- Avoid known aggravating factors
- First use an antiperspirant deodorant (spray or roll on each morning) in axilla (suitable for palms and soles)

  or
- Aluminium chloride hexahydrate 20% solution or spray - apply to affected areas at night when the area is dry (best for palms and soles)

Additional treatments

- Iontophoresis (in specialist centres) to hands and feet
- Injection of botulinum toxin into dermis of affected area. prorem electiveness for axilla and palms.
- (Trial of) probanthine aluminium chloride 20% in alcohol solution to localised area
- Wedge resection of axillary sweat glands

Palmar hyperhidrosis

Treatment

- Aluminium chloride 20% in alcohol solution (Driclor)
- Iontophoresis

Axillary hyperhidrosis

Treatment

- Explanation and reassurance
- See treatment of general hyperhidrosis
- Aluminium chloride 20% in alcohol solution (Driclor); apply nocte for 1 week, then 1 - 2 times weekly or as necessary

Surgery

Wedge resection of a small block of skin and subcutaneous tissue from axillary vault. Define sweat glands with codeine starch powder. The area excised is usually about 4 cm × 2.5 cm.

Body odour/bromhidrosis

Cause: poor hygiene, excessive sweating and active skin bacteria

Main focus: axilla and groin

Precautions: consider uraemia, vaginitis

Treatment method

- Scrub body, especially groins and axillae, with deodorant soap at least morning and night
- Try an antibacterial surgical scrub
- Keep clothes clean, launder regularly
- Choose suitable clothes - natural fibres (e.g. cotton), not synthetics
- Use an antiperspirant deodorant
- Alternative soap - pine soap
- Diet: avoid garlic, fish, asparagus, onions, curry
- Reduce caffeine (coffee, tea and cola drinks), which stimulates sweat activity
- Consider a sugar-free diet
- Shave axillary hair
- Axillary wedge resection for excessive perspiration

Foot odour (smelly and sweaty feet)

Includes pitted keratolysis secondary to hyperhidrosis (common in teenagers).

Treatment (with options)

- Education and reassurance
- Wear cotton or woollen socks
- Aluminium chloride 20% in alcohol solution (Driclor, Hidrosol) or Neat Feet - apply nocte for 1 week, then 1 - 2 times weekly as necessary
- Shoe liners (e.g. ‘Odor eaters’), charcoal inner soles
- Apply undiluted Burow's solution after a shower or bath
- Formalin 1 - 5% soaks every second night
- Iontophoresis
- The teabag treatment (if desperate)
  - prepare 600 mL of strong hot tea (from two teabags left in water for 15 minutes)

  - pour the hot tea into a basin with 2 L of cool water
  - soak the feet in this for 20 - 30 minutes daily for 10 days, then as often as required

Skin disorders of feet

Two conditions commonly seen in teenagers are pitted keratolysis and juvenile plantar dermatosis.

Pitted keratolysis

This malodorous condition known as ‘stinky feet’ or ‘sneakers feet’, usually seen in 10 - 14 year olds, is related to sweaty feet. It has a ‘honeycomb’ pitted appearance. Treatment includes keeping the feet dry and using an ointment such as Whitfield's or an imidazole or sodium fusidate to remove the responsible Corynebacterium organism. Change to all-leather shoes with charcoal liners. Use a drying agent to decrease sweating.

Juvenile plantar dermatosis

‘Sweaty sock dermatitis’ is a painful condition of weight-bearing areas of the feet. The affected skin is red, shiny, smooth and often cracked. It is rare in adults. The treatment is to change to leather or open shoes and to cotton socks. A simple emollient cream gives excellent relief.

PRICKLY HEAT (MILIARIA/HEAT RASH)

- Keep the skin dry and cool (e.g. fan, air-conditioner).
- Dress in loose-fitting cotton clothing.
- Reduce activity.
- Avoid frequent bathing and overuse of soap.
- Duluti topeal sterod cream bd.

Treatment

Lotion: salicylic acid 2%, menthol 1%, chlorhexidine 0.5% in alcohol *or*

Egozite (infants), Isophyl (adults)

If severe: hydrocortisone + clotrimazole cream

Prevention

- Ego Prickly Heat powder

CHILBLAINS AND FROSTBITE

Chilblains (perniosis)

These are localised inflammatory reactions, caused by prolonged exposure to cold, usually on toes and fingers but also on heels, nose, ears and thighs (horse riding).

Precautions

- Think Raynaud's phenomenon.
- Protect from trauma and secondary infection.
- Do not rub or massage injured tissues.
- Do not apply heat or ice.
- Wear warm gloves and socks.

Differences between chilblains and Raynaud's phenomenon

- Chilblains are intermittent without a pattern.
- Itchy at the onset.
- Patchy appearance (can be more generalised).
- Raynaud's has two or three phases including a ‘dead white’ phase with line of demarcation.
- It is significant if it extends to the MCP joints.

Physical treatment

- Elevate affected part
- Warm gradually to room temperature

Drug treatment

- Potent topical corticosteroid (see Table 111.9 at p.1148)
- Apply glyceryl trinitrate vasodilator spray or ointment or patch (use plastic gloves and wash hands for ointment)

Other treatment

- Rum at night
- Nifedipine 20 mg tds or cr 30 mg once daly.
- LIVB therapy weekly for 4-6 weeks prior to cold weather

Frostbite

Treatment depends on severity.

Precautions

- Watch for secondary infection, tetanus, gangrene.

Physical treatment

- Elevate affected limb
- Rewarm in water just above body temperature 40°C (104°F) or use body heat (e.g. in axillae)
- Avoid thawing or refreezing
- Surgical debridement
- Don't debride early (wait until dead tissue dried)
- Don't drink alcohol or smoke
- For blistering, apply warm water compresses for 15 minutes every 2 hours

Drug treatment

- Analgesics

REFERENCES

1. Fry J. Common Diseases (4th edn). Lancaster: MTP Press Ltd, 1985: 337 - 9.
2. Bridges-Webb C. Morbidity and treatment in general practice. Med J Aust: Supplement, 1992; 19 October: 26 - 7.
3. Berger P. Skin Secrets. Sydney: Allen & Unwin, 1991: 93 - 170.
4. Brown P. Dermatitis/eczema. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 142 - 4.
5. Marley (Chair). Therapeutic Guidelines: Dermatology. (version.2)Melbourne: Therapeutic Guidelines Ltd,

2004:148 - 339.

6. ibid 82-90.
7. Brown M. Psoriasis. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 430 - 2.
8. Buxton PK. ABC of Dermatology. London: British Medical Association, 1989: 40.
9. Gallachio V. Nappy rash. Aust Fam Physician, 1988; 17: 971 - 2.
10. Aldridge S. Nappy rash. Aust Paed Review, 1991; 2(1): 2.
11. Berger P. Acne - pathogenesis and treatment. Mod Med Aust, 1990; September: 83 - 9.
12. Gin D. Tinea infections. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 511 - 13.
13. Hunter JAA, Savin JA, Dahl MV. Clinical Dermatology. Oxford: Blackwell Scientific Publications, 1989: 162 - 3.
14. Hunter JAA, Savin JA, Dahl MV. Clinical Dermatology (3rd edn). Oxford: Blackwell Scientific Publications, 2002: 159 - 60.

CHAPTER 114. ACUTE SKIN ERUPTIONS

They say love's like the measles - all the worse when it comes late in life.

DOUGLAS JERROLD (1803 - 57)

ACUTE SKIN ERUPTIONS

The sudden appearance of a rash, which is a common presentation in children (Chapter 83), usually provokes patients and doctors alike to consider an infectious aetiology, commonly of viral origin. However, an important cause to consider is a reaction to a drug.

A knowledge of the relative distribution of the various causes of rashes helps with the diagnostic methodology. Many of the eruptions are relatively benign and undergo spontaneous remission. Fortunately, the potentially deadly rash of smallpox is no longer encountered.

Serious eruptions that demand accurate diagnosis and management include:

- primary HIV infection
- secondary syphilis
- Stevens-Johnson syndrome
- purpuric eruption of meningococcal septicaemia, typhoid, measles, other septicaemia

A list of important causes of acute skin eruptions is presented in Table 114.1.

Table 114.1 Important causes of acute skin eruptions

Maculopapular • Measles • Rubella • Scarlet fever • Viral exanthem (fourth syndrome) • Erythema infectiosum (slapped face syndrome or fifth syndrome) • Roseola infantum (sixth syndrome) • EBM (primary or secondary to drugs) • Primary HIV infection • Secondary syphilis • Pityriasis rosea • Guttate psoriasis • Urticaria • Erythema multiforme (may be vesicular) • Drug reaction • Scabies • Ross River and Barmah Forest infection
Maculopapular vesicular • Varicella • Herpes zoster • Herpes simplex

• Eczema herpeticum • Impetigo • Hand, foot and mouth disease • Drug reaction
Maculopapular pustular • *Pseudomonas* folliculitis • *Staphylococcus aureus* folliculitis • Impetigo
Purpuric (haemorrhagic) eruption • Purpura (e.g. drug-induced purpura, severe infection) • Vasculitis (vascular purpura) - Henoch-Schönlein purpura - polyarteritis nodosa

THE DIAGNOSTIC APPROACH

The diagnostic approach to skin eruptions presupposes a basic knowledge of the causes, and then a careful history and physical examination should logically follow.

The history should include:

- site and mode of onset of the rash
- mode of progression
- drug history
- constitutional disturbance (e.g. pyrexia, pruritus)
- respiratory symptoms
- herald patch?
- diet - unaccustomed food
- exposure to irritants
- contacts with infectious disease
- bleeding or bruising tendency

The examination should include:

- skin of whole body
- nature and distribution of rash, including lesion characteristics
- soles of feet
- nails
- scalp
- mucous membranes
- oropharynx
- conjunctivae and the lymphopoietic system (? lymphadenopathy ? splenomegaly)

Laboratory investigations may include:

- a full blood examination
- syphilis serology
- Epstein-Barr mononucleosis test
- HIV test

- rubella haemagglutination tests (×2)
- viral and bacterial cultures

ACUTE SKIN ERUPTIONS IN CHILDREN

The following skin eruptions (some of which may also occur in adults) are outlined in childhood infectious diseases (Chapter 83).

- measles (p. 902)
- rubella (p. 903)
- viral exanthem (fourth syndrome) (p. 904)
- erythema infectiosum (fifth syndrome) (p. 904)
- roseola infantum (sixth syndrome) (p. 905)
- Kawasaki's disorder (p. 907)
- varicella (p. 899)
- impetigo (p. 910)



Figure 114.1 Pityriasis rosea in a 10-year-old child showing the salmon pink scaly eruption following skin cleavage lines - the ‘Christmas tree’ pattern

PITYRIASIS ROSEA

Pityriasis rosea is a common but mild acute inflammatory skin disorder. Although a viral agent (possibly human herpesvirus 7) is suspected to be the cause, no infective agent has been demonstrated.

Clinical features

- Any age, mainly young adults (aged 15 - 30 years)
- Preceding herald patch (1 - 2 weeks); can have 2 - 3, but none in 20% (can be mistaken for ringworm)
- Oval, salmon-pink or copper-coloured eruption
- Coin-shaped patches with scaly margins 1 - 2 cm

- Follows cleavage lines of skin (Figure 114.2)


widespread and symmetrical, in old 'bathing trunk' distribution
follows cleavage lines of skin
'Christmas tree pattern'
oval salmon-pink maculopapules
herald patch

Figure 114.2 Pityriasis rosea: typical distribution

- On trunk ('T-shirt' distribution)
- Occurs also on upper arms, upper legs, neck, face (rare) and axillae
- Patients not ill
- Itch varies from nil to severe (typically minor itching)
- Scale is on inner aspect of active border

Differential diagnosis

Herald patch	tinea corporis
Generalised rash	seborrhoeic dermatitis (slower onset), guttate psoriasis, drug eruption (Table 114.2), secondary syphilis

Table 114.2 Drugs that cause eruptions suggestive of pityriasis rosea[1]

Main drugs
• gold salts
• penicillamine
• captopril
Others
• arsenicals
• barbiturates
• bismuth
• clonidine
• metoprolol
• metronidazole

Prognosis

A mild self-limiting disorder with spontaneous remission in 2 - 10 weeks (average 2 - 5). It does not appear to be contagious.

Management

- Explain and reassure with patient education handout.[2]
- Bathe and shower as usual, using a neutral soap (e.g. Neutrogena).
- Use a soothing bath oil (e.g. QV bath oil).
- For bothersome itch: apply mild topical corticosteroid ointment or calamine lotion with 1% phenol or menthol 1% in aqueous cream.
- For severe itch use a potent topical corticosteroid or oral corticosteroids.
- UV therapy is good but, like psoriasis, sunburn must be avoided. Expose rash to sunlight or UV therapy (if florid) three times a week.

SECONDARY SYPHILIS

The generalised skin eruption of secondary syphilis varies and may resemble any type of eruption from psoriasiform to rubelliform to roseoliform. The rash usually appears 6 - 8 weeks after the primary chancre.

Features of the rash

- Initially faint pink macules
- Then becomes maculopapular
- Can involve whole of body (Figure 114.3)



Figure 114.3 Secondary syphilis: typical features

- Palms and soles involved
- Dull red in colour and round
- More prolific on flexor surfaces
- Symmetrical and relatively coarse
- Asymptomatic

Associations (possible)

- Mucosal ulcers: ‘snail track’
- Lymphadenopathy
- Patchy hair loss
- Condylomata lata

Treatment

- As for primary syphilis (Chapter 110)

PRIMARY HIV INFECTION

A common manifestation of the primary HIV infection is an erythematous, maculopapular rash, although other skin manifestations such as a roseolalike rash and urticaria can occur.

Clinical features

- Symmetrical
- May be generalised
- Lesions 5 - 10 mm in diameter
- Common on face and/or trunk
- Can occur on extremities including palms and soles (Figure 114.4)



Note: mimics glandular fever

Figure 114.4 Primary HIV infection: typical features

- Non-pruritic

If such a rash accompanied by an illness like glandular fever occurs, HIV infection should be suspected and specific tests ordered.

GUTTATE PSORIASIS

Guttate psoriasis is the sudden eruption of small (less than 5 mm), very dense, round red papules of psoriasis on the trunk (Figure 114.5 and Figure 114.6). The rash may extend to the limbs.



Figure 114.5 Guttate psoriasis: small, drop-like lesions, mainly on trunk



Figure 114.6 Guttate psoriasis in a child showing the small dense, round erythematous papules of psoriasis on the trunk

It is usually seen in children and adolescents and often precipitated by a streptococcal throat infection. The rash soon develops a white silvery scale. It can persist for up to 6 months. It may undergo spontaneous resolution or enlarge to form plaques.

Treatment includes UV light as well as tar preparations.

EPSTEIN-BARR MONONUCLEOSIS

The rash of EBM is almost always related to antibiotics given for tonsillitis. The primary rash, most often non-specific, pinkish and maculopapular (similar to that of rubella), occurs in about 5% of cases only. The secondary rash, which can be extensive and sometimes has a purplish-brown tinge, is most often precipitated by one of the penicillins (Figure 114.7):

- ampicillin 90 - 100% association
- amoxycillin 90 - 100% association
- penicillin up to 50%


rubelliform rash
pink to purple maculopapular raised rash
extensor surfaces on limbs

Figure 114.7 Epstein-Barr mononucleosis: typical rash induced by penicillin, amoxycillin or ampicillin

DRUG ERUPTIONS

A rash is one the most common side-effects of drug therapy, which can precipitate many different types of rash; the most common is toxic erythema (Table 114.3). Most drug-evoked dermatoses have an allergic basis with the eruption appearing approximately 10 days after administration, though much sooner if previously sensitised.[3] The most common drugs that cause skin eruptions are summarised in Table 114.4.

Table 114.3 Most common types of drug eruptions (after Thomas)[3]

	Relative frequency %
Toxic erythema	45
Urticaria/angioedema	25
Erythema multiforme	7
Eczematous dermatitis	5
Fixed drug reaction	3
Photosensitivity	3

Others • acne form • psoriasiform • pigmentation • erythema nodosum • toxic epidermal necrolysis • vasculitis/purpuric • pigmentary • exfoliative

Table 114.4 Most common drugs that cause skin eruptions

Antimicrobials	• penicillin/cephalosporins • sulphonamides • tetracyclines • nitrofurantoin • streptomycin • griseofulvin • metronidazole • antiretroviral agents • trimethoprim • dapsone
Diuretics	• thiazides • frusemide
Anticonvulsants	• carbamazepine • phenytoin • lamotrigine
Tranquillisers	• phenothiazines • barbiturates • chlordiazepoxide
Anti-inflammatory and analgesics	• gold salts • aspirin/ salicylates • codeine/ morphine • pyrazolones (e.g. BTZ) • other NSAIDs
Hormones	• combined oral contraceptive • stilboesterol • testosterone
Others	• phenolphthalein • serum • amiodarone

	• cytotoxic drugs • quinidine/quinine • bromides and iodides • sulphonylureas • allopurinol • warfarin • amphetamines

The most important fact to realise about drug reactions is that their appearances are so variable - they may mimic almost any cutaneous disease and, in addition, create unique appearances of their own.

When taking a history it is appropriate to enquire about medications or chemicals that may be overlooked such as aspirin, vitamins, toxins, laxatives and medicated toothpaste.

Toxic erythema

The maculopapular erythematous eruption is either morbilliform or scarlatiniform. It is more pronounced on the trunk than on the limbs and face but may become confluent over the whole body (Figure 114.8).



Figure 114.8 Toxic erythema: maculopapular erythematous scarlatiniform eruption caused by amoxycillin

Drugs that typically cause toxic erythema include:

- antibiotics
    - penicillin/cephalosporins
    - sulphonamides
- thiazides
- carbamazepine
- barbiturates
- allopurinol
- gold salts

Photosensitivity

Several antibiotics increase the sensitivity of the skin to UV light and may lead to a rash with a distribution according to sunlight exposure. The photosensitive rash may be erythematous, resembling sunburn; eczematous; or vesicular.

Typical drugs:

- tetracyclines
- sulphonamides/sulphonylureas
- thiazides and frusemide
- phenothiazines
- retinoids
- amiodarone
- griseofulvin
- antihistamines, especially promethazine
- antimalarials
- psoralens

Fixed drug eruption

The mechanism of fixed drug eruption is unknown. The most commonly affected areas are the face, hands and genitalia. The lesions, which are usually bright red but can have other characteristics, are fixed in site and appearance within hours of the drug's administration.

Typical drugs:

- phenolphthalein
- tetracyclines
- penicillins
- sulphonamides
- salicylates
- the oral contraceptive pill
- barbiturates
- chlordiazepoxide
- quinine

Treatment of any drug reaction

The important aspect of management is to recognise the offending agent and withdraw it. The rash should be treated according to its nature.

There is a therapeutic impulse to prescribe antihistamines but they should be reserved for the treatment of urticarial drug eruptions. They may actually delay healing in purpuric, erythematous and vesiculobullous reactions. Antihistamines may act as allergens and show cross-sensitivity with phenothiazines, sulphonamides and topical antihistamines.

Table 114.5 lists drugs with the highest skin reaction rates.

Table 114.5 Drugs with the highest skin reaction rates[4]

• penicillin and derivatives
• sulphonamides[*]
• trimethoprim[*]
• thiazide diuretics

- allopurinol*
- dapsone*
- NSAIDs, esp. piroxicam*
- nevirapine*, abacavir*
- barbiturates
- quinidine
- anticonvulsants (phenytoin, lamotrigine*)
- blood products
- gold salts

*Severe reactions

ERYTHEMA MULTIFORME

Erythema multiforme is an acute eruption affecting the skin and mucosal surfaces.

Clinical features:

- mainly in children, adolescents, young adults
- symmetric
- erythematous papules
- mainly backs of hands, palms and forearms (Figure 114.9)


raised
macules
of concentric
rings
usually
mouth and
extremities
flexor sides
of forearms

Figure 114.9 Erythema multiforme: typical distribution

- also on feet and toes, mouth
- occasionally on trunk and genitalia
- polymorphic
- vesicles and bullae may develop
- self-limiting (up to 2 weeks)

Stevens-Johnson syndrome

A very severe and often fatal variant. Sudden onset with fever and constitutional symptoms.

Causes and associations

Erythema multiforme is a vasculitis affecting the skin and mucosa. Herpes simplex virus (usually type 1) is the commonest known association.

Associations include:

- unknown 50%
- herpes simplex virus 33%
- other infections
    - *Mycoplasma* pneumonia
    - tuberculosis
    - *Streptococcus*
- connective tissue disorders (e.g. SLE)
- neoplasia
    - Hodgkin's lymphoma
    - myeloma
    - carcinoma
- deep X-ray therapy
- drugs
    - barbiturates
    - penicillin
    - sulphonamides
    - phenothiazines
    - phenytoin

Treatment

- Identify and remove cause (e.g. withdraw drugs).
- Symptomatic treatment (e.g. antihistamines for itching).
- Refer severe cases.

ERYTHEMA NODOSUM

Erythema nodosum is characterised by the onset of bright red, raised, tender nodules on the shins. It is an acute, inflammatory, immunological reaction in the subcutaneous fat. The nodules may appear on the thighs and the arms. Adult females are typically affected. An arthritic reaction can affect the ankles and knees.

Causes/associations

- Sarcoidosis (commonest known cause)
- Infections
    - streptococcal infections
    - viral infections, e.g. hepatitis B
    - tuberculosis
    - leprosy
    - chlamydia infection
    - fungal infections
- Inflammatory bowel disorders, e.g. Crohn's

- Drugs
    - sulphonamides
    - tetracyclines
    - oral contraceptives
    - bromides and iodides
- Malignancy, e.g. lymphoma, leukaemia
- Unknown (about 40%)

Investigations

Tests include FBE, ESR, chest X-ray (the most important) and Mantoux test.

Treatment

Identify the cause if possible. Rest and analgesics for the acute stage. Systemic corticosteroids speed resolution if severe episodes.

Prognosis

There is a tendency to settle spontaneously over 3 - 8 weeks. The lesions may recur.

HERPES ZOSTER

Herpes zoster (shingles) is caused by reactivation of varicella zoster virus (acquired from the primary infection of chickenpox) in the dorsal root ganglion. The term comes from the Greek *herpes* (to creep) and *zoster* (a belt or girdle). Shingles is from the Latin *cingere* (to gird) or *cingulum* (a belt). In most instances the reason for reactivation is unknown, but occasionally it is related to an underlying malignancy, usually leukaemia or a lymphoma, to immunosuppression, or to a local disease or disturbance of the spine or spinal cord, such as a tumour or radiotherapy.

The incidence is 3.4 cases per 1000 population per year. A person of any age can get herpes zoster but it is more common in people over 50 years.

Clinical features

The main features are:

- the condition is preceded by several days of radicular pain with hyperaesthesia
- unilateral patchy rash in one or two contiguous dermatomes (Figure 114.10)



Figure 114.10 Herpes zoster (shingles) involving the L2 nerve root in a 63-year-old woman presenting with low back and groin pain. Calamine lotion has been applied to soothe the discomfort

- intense erythema with papules in affected skin
- later crusting and separation of scabs after 10 - 14 days, often with depigmentation, and
- regional lymphadenopathy

Distribution

Any part of the body may be affected, but thoracic and trigeminal dermatomes are the most common. It follows the distribution of the original varicella rash (worse on the face and trunk).

Cranial nerve involvement

The trigeminal nerve - 15% of all cases:

- ophthalmic branch - 50% affects nasociliary branch with lesions on tip of nose and eyes (conjunctivae and cornea)
- maxillary and mandibular - oral, palatal and pharyngeal lesions

The facial nerve: lower motor neurone facial nerve palsy with vesicles in and around external auditory meatus (notably posterior wall) - the Ramsay Hunt syndrome.

Complications

- Rare: meningoencephalitis
- Uncommon: motor paralysis
- Common:
  - postherpetic neuralgia; increased incidence with age and debility, with duration greater
  - than 6 months:
    - less than 50 years 1%
    - 50 - 59 years 7%
    - 60 - 69 years 21%
    - 70 - 79 years 30 - 50%
  - the neuralgia resolves within 1 year in 70 - 80% of these patients but in others it may persist for years
  - eye complications of ophthalmic zoster including keratitis, uveitis and eyelid damage

Management

- Appropriate detailed explanation and reassurance. Dispel myths: namely, that it is not a dangerous disease, the patient will not go insane nor die if the rash spreads from both sides and meets in the middle.
- Explain that herpes zoster is only mildly contagious but children can acquire chickenpox after exposure to a person with the disorder. It is advisable to avoid contact with infants and young children who have never had chickenpox and avoid contact with the immunocompromised and those undergoing chemotherapy. Consider giving varicella zoster immunoglobulin to those immunocompromised contacts who have no history of varicella.
- Treating the rash: Instruct the patient to avoid overtreating the rash, which may become infected. Calamine lotion may be soothing but removal of the calamine can be painful. For a hot painful rash to remove crusts and exudate bathe the lesions with saline three times daily. A drying lotion (e.g. menthol in flexible collodion) is most soothing and suitable. Cover the lesions in a light non-adherent padded dressing.
- Oral medication[5]
  - Analgesics such as aspirin or paracetamol with or without codeine should be first-line therapy.
  - Antiviral therapy may reduce the duration of the disease in all people. Optimal indications - patients:
    - treated within 72 hours of onset of vesicles
    - who are > 50 years of age
    - who are immunocompromised
    - with acute severe pain
    - with involvement of special areas (e.g. eye, perineum

  Dose:[5] aciclovir 800 mg 5 times daily for 7 days *or* famciclovir 250 mg 8 hourly for 7 days *or* valaciclovir 1000 mg 8 hourly for 7 days Prevention: Vaccine ‘Zostavac’ when available.

Postherpetic neuralgia

This pain is usually severe, varying in quality from paroxysmal stabbing pain to burning or aching. Spasms of pain upon light brushing of the skin is a feature.

Treatment is difficult and a careful ‘trial and error’ approach can be used until appropriate evidence from scientific trials establishes the optimal treatment.

Treatment options[5]

- Simple analgesics
- TENS as often as necessary (e.g. 16 hours/day for 2 weeks)[5] *plus*
- Oral medication
  - tricyclic antidepressants e.g. amitriptyline 10 - 50 mg (o) nocte *or* desipramine 25 - 50 mg (o) nocte This is a starting trial dose, particularly in elderly patients
  - Carbamazepine (for lancinating pain) 50 - 100 mg (o) bd initially, increasing the dose gradually to avoid drowsiness (up to 400 mg bd) *or*
  - Gabapentin 300 mg (o) daily (nocte) initially, increasing as tolerated to tds
- Topical medication
  - capsaicin 0.025% (Capsig) cream; apply 4 times a day (application of local anaesthetic cream 20 minutes beforehand may prevent a burning sensation) for 6 weeks[6] *or*
  - lignocaine 5% ointment or 10% gel *or*
  - lignocaine 5% patch to painful area

Evidence-based medicine

Systematic reviews confirm that oral antiviral agents (as above) are beneficial for preventing postherpetic neuralgia while tricyclic antidepressants and gabapentin are beneficial for relieving the established condition.[7]

Physical treatments

- Local corticosteroid and anaesthetic injections.
- Nerve blocks (e.g. supraorbital nerve).
- Excision of painful skin scar. If the neuralgia of 4 months or more is localised to a favourable area of skin, a most effective treatment is to excise the affected area, bearing in mind that the scar tends to follow a linear strip of skin. This method is clearly unsuitable for a large area.

Method

- Mark out the painful area of the skin.
- Incise it with its subcutaneous fat, using an elongated elliptical excision (Figure 114.11).



Figure 114.11 Postherpetic neuralgia: example of type of excision for a severe problem

- Close the wound with a subcuticular suture or interrupted sutures.

HERPES SIMPLEX

Herpes simplex is a common infection caused by the large DNA herpes simplex virus (HSV) which can cause a vesicular rash anywhere on the skin or mucous membranes (Figure 114.12). There are two major antigenic strains of HSV:

- *HSV 1*, which commonly involves the lips and oral mucosa
- *HSV 2*, which basically affects the genitalia (common in adolescents and young adults)



Figure 114.12 Acute eruption of herpes simplex on the face - a recurrent problem in this person

Epidemiology

HSV has a worldwide distribution and is spread orally or genitally by infected secretions. Primary HSV infection is usually a disease of childhood, characteristically causing acute gingivostomatitis in a preschool child (p. 909).

Table 114.6 summarises the major manifestations of HSV and the possible complications.

Table 114.6 Herpes simplex virus: manifestations and complications

Examples of manifestations
• Herpes labialis (synonyms: fever blisters, cold sores)
• Keratoconjunctivitis, including dendritic ulcer
• Genital infection
• Other areas of skin such as buttocks
Complications
• Eczema herpeticum
• Erythema multiforme (3 - 14 days postinfection), often recurrent
• Myeloradiculopathy with genital herpes
• Pneumonia
• Encephalitis

Recurrent infection

Recurrences range from weeks to months and appear due to reactivation rather than reinfection. The cause is not clear but there are several known precipitating factors. These are fever, sunlight, respiratory infections, menstruation, emotional stress, local trauma and, with genital lesions, sexual intercourse.

Fatalities

HSV infections can be potentially fatal. Reactivated HSV can cause a focal destruction encephalitis. The untreated case fatality rate is as high as 70%, but this can be greatly reduced with the use of aciclovir. Neonates exposed to HSV can develop fatal disseminated infection. In compromised patients the infection can be severe.

Diagnosis

If the clinical picture is uncertain, immuno-fluorescence of, or culture from, vesicle fluid can aid diagnosis.

Genital herpes

See pp. 1130 - 1.

Herpes labialis (classic cold sores)

The objective is to limit the size and intensity of the lesions. At the first sensation of the development of a cold sore:

- apply an ice cube to the site for up to 5 minutes every 60 minutes (for first 12 hours)
- topical applications include idoxuridine 0.5% preparations (Herplex D liquifilm, Stoxil topical, Virasolve) applied hourly *or* saturated solution of menthol in SVR *or* povidone - iodine 10% cold sore paint: apply on swab sticks 4 times a day until disappearance *or* 10% silver nitrate solution: apply the solution carefully with a cotton bud to the base of the lesions (deroof vesicles with a sterile needle if necessary); may be repeated[8] *or* aciclovir 5% cream 5 times daily for 4 days *or* penciclovir 1% cream for 4 days

Note: Corticosteroids are contraindicated.

Oral medication

For a severe primary attack:

- aciclovir 200 mg 5 times daily or 800 mg twice daily for 7 - 10 days or until resolution or famciclovir or valaciclovir

Topical zinc treatment

- Zinc sulphate solution 0.025 - 0.05%, apply 5 times a day for cutaneous lesions
- Use 0.01 - 0.025% for mucosal lesions9

Prevention

If exposure to the sun precipitates the cold sore, use a 15 or more sun protection lip balm, ointment or solastick. Zinc sulphate solution can be applied once a week for recurrences. Oral aciclovir 200 mg bd (6 months) can be used for severe and frequent recurrences.[9]

Advice to the patient

Herpes simplex is contagious. It is present in saliva and can be spread in a family by the sharing of drinking and eating utensils and toothbrushes, or by kissing. It is most important not to kiss an infant if you have an active cold sore.

FOLLICULITIS[10]

Folliculitis, which is infection in and around hair follicles, can be superficial or deep.

Superficial folliculitis

This usually presents as mild itchy pustules on an erythematous base on any part of hair-bearing skin, particularly in hot weather in a patient who is often a chronic carrier of *S. aureus.* A swab supports diagnosis. Management involves removal of the cause and the application of an antiseptic wash, such as triclosan 1%, chlorhexidine or povidone - iodine. Occasionally oral flucloxacillin may also be required.

Bacterial folliculitis[3]

A generalised acute erythematous maculopapular rash can be a manifestation of bacterial folliculitis, typically caused by *S. aureus, Pseudomonas aeruginosa*, or by fungal folliculitis due to *Pityrosporum orbiculare* or other dermatophytes.

Pseudomonas folliculitis can cause confusion, the typical features being:

- rapidly spreading rash
- mainly on trunk, buttocks and thighs, especially axillae and groin
- itchy
- small pustules surrounded by circular red-purple halo
- follows immersion in a hot spa bath or tub

Treatment is based on the sensitivity of the cultured organisms (e.g. ciprofloxacin). Many cases resolve spontaneously in 1 - 2 weeks.

Folliculitis of trunk from spa baths

'Hot tub folliculitis' is caused by *P. aeruginosa* (usually) in poorly chlorinated water maintained at temperatures 37 - 40°C.

Treatment is with ciprofloxacin 500 mg (o) bd for 7 days.

Folliculitis of groin (pseudofolliculitis)

Folliculitis of the groin area is common in women who shave. It tends to recur.

Management

- Use tea tree (melaleuca) lotion daily for folliculitis.
- Prior to shaving apply 'tea tree wash'.
- Change shaving habits: avoid close shaving; shave less often; shave in direction of hair growth and use good quality blades.
- If persistent, use povidone - iodine or chlorhexidine (Hibiclens) solution or triclosan 1%.
- If severe, use mupirocin 2% (Bactroban) ointment.

Deep folliculitis

Deep forms are usually very tender and painful. Examples are styes, boils (furuncles) and carbuncles.

Boil (furuncle)[10]

This is an *S. aureus* infection of a hair follicle and may occur in any hair-bearing site. The painful red nodule enlarges, becomes fluctuant and develops a necrotic centre, which discharges a core of thick yellow pus tinged with blood (with associated relief of pain).

Treatment (adults)

- (According to swabs) di(flu)cloxaciilin 500 mg or cephalexin 500 mg (orally, 6 hourly for 5 - 7 days) or roxithromycin 300 mg (o) daily or erythromycin 500 mg (o) 12 hourly for 7 days

Boils - recurrent

- Obtain swabs
- 3% hexachlorophane body wash daily
- Mupirocin to the lesions and nares
- Antibiotics (as above) - according to swabs

Carbuncle

This is a cluster of small abscesses involving a group of adjoining hair follicles. Common sites are the back of the neck, the shoulders, buttocks or over the hips. Treatment is as for a boil.

Stye of eye

- Apply heat with direct steam from a thermos (Figure 51.8 at p. 559) onto the closed eye or by a hot compress (helps spontaneous discharge).
- Perform lash epilation to allow drainage (incise with a D_{11} blade if epilation doesn't work).
- Only use topical antibiotic ointment (e.g. chloramphenicol) if infection is spreading locally, and systemic antibiotics if distal spread noted by preauricular adenitis.

REFERENCES

1. Hunter JAA, Savin JA, Dahl MV. *Clinical Dermatology* (3rd edn). Oxford: Blackwell Scientific Publications, 2002: 64.
2. Murtagh J. *Patient Education* (4th edn). Sydney: McGraw-Hill, 2005: 235.
3. Thomas RM. Drug eruptions. Med Int, 1988; 49: 2038 - 42.
4. Marley J (Chair). *Therapeutic Guidelines: Dermatology* (Version 2). Melbourne: Therapeutic Guidelines Ltd; 2004: 117 - 46.
5. Marley J (Chair). *Therapeutic Guidelines: Analgesics* (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2002: 309 - 11.
6. Bernstein JE, Korman NJ, Bickers DR et al. Topical capsaicin treatment of chronic postherpetic neuralgia. J Am Acad Dermatol, 1989; 21: 265 - 70.
7. Lancaster T, Warehain D, Yaphe J. Postherpetic neuralgia. In: Barton S ed. *Clinical Evidence* (Issue 7). London: BMJ Publishing Group, 2002: 739 - 45.
8. Pollack A. Treatment of herpes simplex: a practice tip. Aust Fam Physician, 1982; 11: 952.
9. Russo GJ. *Herpes Simplex and Herpes Zoster*. Glendale: Audio Digest, Family Practice, 1991; 39: 38.
10. Marley J (Chair). *Therapeutic Guidelines: Dermatology* (Version 2). Melbourne: Therapeutic Guidelines Ltd; 2004: 218 - 9.

CHAPTER 115. SKIN ULCERS

An ulcer, which occurring at any of the vital parts of the body secretes a copious quantity of pus and blood, and refuses to be healed, even after a course of proper and persistent medical treatment, is sure to have a fatal determination.

SUSHRUTA-SAMHITE (5th CENTURY BC)

SKIN ULCERS

An ulcer is a localised area of necrosis of the surface of the skin or mucous membrane. It is usually produced by sloughing of inflamed necrotic tissue. Ulcers of the skin are common, particularly on the legs and feet, on areas exposed to the sun, and on pressure areas such as the sacrum in older people.

The national morbidity survey (UK) showed that 2 - 3 per 1000 patients per annum consulted their general practitioner with ‘chronic ulcers of the skin’.

KEY FACTS AND CHECKPOINTS[1]

- The great majority of leg ulcers are vascular in origin due to arterial insufficiency or venous hypertension.
- If clinical findings don't provide the diagnosis, ordering the ankle brachial index (ABI) is essential if pulses are not palpable to exclude arterial disease. Duplex Doppler ultrasound is the key investigation for venous disease.
- Most ulcers are multifactorial - 1. venous + obesity + immobility (often with osteoarthritis) + poor compliance or 2. venous + arterial + trauma + infection.
- In treatment correct as many factors impairing wound healing as possible.
- Correctly applied graduated compression is the mainstay of treatment for venous ulceration but inappropriate compression can exacerbate arterial ulcers.
- Multi layered bandages is the gold standard.
- High compression is better than low compression.
- Elastic bandages are better than non-elastic bandages.
- Venous surgery has a limited role in management since most ulcers heal by compression alone.
- Moist or occlusive dressings provide a physiological environment for wound healing.
- Adequate wound debridement is essential to remove necrotic material and slough, and enable healing to commence and progress.
- Hypergranulation tissue can delay wound healing. It should be removed and strategies include surgical removal, topical silver nitrate, hypertonic saline dressings and pressure bandaging.
- Diabetes and rheumatoid arthritis (vasculitis) are also common conditions associated with leg ulceration.
- Accurate diagnosis of the ulcer is vitally important for management decisions.
- Bacterial swabs are unhelpful because all chronic ulcers will become colonised with both Gram-positive and Gram-negative bacteria.

A CLINICAL APPROACH

It is useful to keep in mind the various causes of ulcers (Table 115.1). The commonest causes or types are venous and ischaemic ulcers of the leg, pressure ulcers (decubitus) and trauma. It is important not to misdiagnose malignant ulcers, including ‘Marjolin's ulcer’, which is SCC developing in unstable chronic scars or ulcers (e.g. burns, venous ulcers, tropical ulcers) of longstanding duration. Amelanotic melanoma is a specific trap.

Table 115.1 Types and causes of skin ulcers

Traumatic

Decubitus (related to trauma)
Vascular • Venous - varicose veins - post thrombophlebitis • Arterial insuffiency • Skin infarction (thrombolitic ulcer) • Vasculitis - rheumatoid arthritis, SLE, scleroderma
Infective • Tropical ulcer • Tuberculosis • Mycobacterium ulcerans • Post cellulitis • Chronic infected sinus
Malignant • Squamous cell carcinoma • Marjolin's ulcer (SCC) • Basal cell carcinoma (rodent ulcer) • Malignant melanoma • Ulcerating metastases **Neurotrophic**
• Peripheral neuropathy (e.g. diabetes) • Peripheral nerve injuries (e.g. leprosy)
Haematological • Spherocytosis • Sickle cell anaemia
Miscellaneous • Artefactual • Pyoderma gangrenosum (diagnosis of exclusion) • Insect and spider bites

History

A careful history helps determine the cause of the ulceration. Relevant history includes previous deep venous thrombosis or pulmonary embolism, diabetes, rheumatoid arthritis, inflammatory bowel disease, chronic skin ulcers and arterial insufficiency, including a history of intermittent claudication and ischaemic rest pain.

A drug history is important, considering especially beta-blockers and ergotamine, which can compromise the arterial circulation, corticosteroids and NSAIDs, which affect healing, and nifedipine, which tends to aggravate ankle oedema.

Examination of the ulcer [2]

Any ulcer should be assessed for the following characteristics:

- site
- shape
- size
- edge - consider consistency
- floor
- base
- discharge
- surrounding skin
    - colour (? signs of inflammation)
    - sensitivity
- mobility in relation to underlying tissue
- regional lymph nodes

Site of ulcer

Venous ulcers typically occur on the medial side of the leg in relation to incompetent perforating veins in the traditional gaiter area (Figure 115.1). Ischaemic ulcers tend to occur on the lateral side and anterior part of the leg.



Figure 115.1 Area typically affected by varicose eczema and ulceration (the ‘gaiter’ area)

Trophic ulcers, which are associated with neuropathy, occur on parts subject to repeated pressure and trauma, such as the ‘ball’ of the foot or the pulps of the fingers.

Solar-induced ulcers, such as SCCs and BCCs, occur on such parts exposed to the sun. It should be noted if the ulcer is related to old scars, including burns and chronic ulcers.

Size, shape and edge

The classic appearances of various ulcers are presented in Figure 115.2. These are general guidelines only. Infective ulcers due to *Mycobacterium* species, and bed sores, tend to have an undermined edge while a trophic ulcer is punched out and typically round in surface shape. A raised firm ulcer edge may indicate malignancy.


(a)
edge
floor
base
(b)
Edge
Example
punched out
trophic ulcer
arterial ulcer
undermined
bed sore
everted
squamous cell carcinoma
rolled
basal cell carcinoma

Figure 115.2 (a) Parts of an ulcer (b) Types of ulcer Reproduced with Permission of Pergamon Press: Davis et al. Symptom Analysis and Physical Diagnosis (2nd edn), p. 309

Floor of the ulcer

The floor or base of the ulcer provides useful clinical information. A dry or extended base or necrotic eschar in the floor implies ischaemia. Venous ulcers on the other hand are often superficial and tend to have fibrinous exudate and ooze, sometimes purulent fluid.

Colour guide:

- black - necrosis
- yellow - slough
- red - granulation
- pink - epithelium

Investigations

The following should be considered, according to the clinical findings:

- full blood count
- ESR
- random blood sugar
- rheumatoid factor tests
- duplex Doppler ultrasound
- swab for specific organisms
- biopsy, especially if SCC suspected (be careful of biopsy if melanoma: amelanotic melanomas are a trap)

LOWER LIMB ULCERATION

The most common causes of lower limb ulceration are venous disease, arterial disease and diabetes.

Differentiating between leg ulcers (85%) and foot ulcers (15%) is very important since they present two very different problems.[3] According to the survey by Stacey, venous disease is present in two-thirds of leg ulcers, while arterial disease occurs in 28% (Table 115.2). Ulceration on the foot frequently has an arterial aetiology (72%), with many of these patients also having diabetes, whereas venous disease is present in only 6%.[3]

Table 115.2 Causes of chronic ulceration of the leg and foot[3]

	%
The leg	
Venous disease	52
Mixed venous and arterial disease	15
Arterial disease	13
Others	20
The foot	
Arterial disease	72
Mixed venous and arterial disease	2
Venous disease	4
Others	22

The differential characteristics are presented in Table 115.3.

Table 115.3 Comparison of typical features of venous and arterial ulceration of the leg

	Venous	Arterial
Site	Around ankle and lower third of leg (gaiter area)	Distal to ankle
	Just above medial and lateral malleoli	Over pressure points on toes, side of foot, metatarsal heads
Pain	Nil to mild	Usually moderate to severe
Oedema	Usually present	Usually absent
Ulcer features	Ragged' edge	‘Punched out’
	Often superficial	Often deep, involving deep fascia
	Ooze + + +	Dry
Associated limb features	Varicosities	‘Cold’ extremities
	Leg warm, red, oedematous	Ischaemic changes
	Varicose dermatitis	Diminished or absent peripheral pulses
	Haemosiderin deposits	Thin, shiny, dry skin
	Atrophie blanche	
History	Limb oedema	Peripheral vascular disease - claudication, rest pain
	Past DVT	Diabetes
	Failed graft	

		Smoker
ABI	> 0.9	< 0.5 - 0.8

Based on a table prepared by Dr Denise Findlay; reproduced with permission

A general examination, including the leg, is very important. This includes examining the venous drainage (pp. 720 - 1), the arterial pulses and the sensation of the leg, and checking for the presence of diabetes.

Appropriate investigations (if required) include:

- full blood count
- blood sugar
- duplex Doppler ultrasound for arterial circulation

To swab or not to swab

A routine ulcer swab is not considered to be of significant value. If specific organisms such as *Mycobacterium ulcerans* are suspected, then cultures are necessary.

Measurement of ankle/brachial pressure index

To plan management of a leg ulcer, it is ideal to determine the blood flow with hand-held Doppler ultrasound. Measure ankle and brachial systolic pressures and then determine the ABI, which is the ankle pressure divided by the brachial pressure. Typical levels are:[4]

- normal > 0.9 (venous ulcer)
- ischaemic < 0.5 (arterial ulcer)
- claudicant 0.5 - 0.9 (mixed arterial - venous ulcer, significant ischaemia if < 0.8)

The ability to determine the cause of the ulceration and thus treat accordingly, especially with pressure dressings, has been a major advance in management. An ABI < 0.8 warrants caution in applying any compression. < 0.4 demands urgent referral.

Mild compression may be used if AB 0.6 - 0.8.

VENOUS ULCERATION

Venous ulceration (synonyms: ‘varicose’, ‘stasis’ and ‘gravitational’ ulcers) accounts for the majority of leg ulcers. Chronic venous insufficiency is one of the most common medical problems in the elderly, with an estimated incidence of 5.9%.[5]

The problem is invariably secondary to deep venous thrombophlebitis. The subsequent chronic venous hypertension produces trophic changes such as hyperpigmentation (Figure 115.3), fibrotic thickening, induration and oedema. The end point of this process is ulceration, which affects 3% of those with varicose veins and 30% of those with trophic changes.[6]



Figure 115.3 Venous ulceration in an elderly patient with post phlebitic varicose ‘eczema’. Venous pigmentation, dermatitis, atrophy and calcification of the subcutaneous tissues are present. Photo courtesy Terry Devine

Clinical features of venous ulcers[7]

- Occur in same area as venous eczema
- Shallow (but can reach periosteum)
- More common medial than lateral
- Sometimes circumferential
- Granulating floor sometimes with surrounding cellulitis
- Notoriously slow in healing
- Generally not tender but can be painful
- Associated pain is usually relieved by raising the leg

On examination, superficial varicosities are usually but not invariably present. Pitting oedema may be present early but with time fibrosis and firm induration develop. Other clinical features include dermatitis (eczema), punctuate capillary proliferation, haemosiderin, hyperpigmentation and ‘atrophie blanche’ (porcelain white scar with rim of telangectasia).[8]

ARTERIAL (ISCHAEMIC) ULCERS

Ischaemic ulcers are generally localised to the most peripheral areas below the ankle joint (Figure 115.4), such as the tips of the toes and the point of the heel, or to pressure points such as the heels, malleoli or head of the first metatarsal.



Figure 115.4 Ischaemic 'arterial' ulcers in an elderly woman with a long history of intermittent claudication and a recent history of nocturnal ischaemic rest pain in the feet. The ulcers healed after reconstruction for arterial obstruction. *Photo courtesy Terry Devine*

Clinical features:

- painful
- punched out
- minimal granulation tissue

Management is directed towards reperfusion.

MANAGEMENT OF VENOUS LEG ULCERS

A major advance in the management of venous ulcers has been the finding that wounds heal better in an occluded or semi-occluded state.[8] A moist environment also aids healing.

Principles of optimal management

- Explanation about the cause, and promotion of patient compliance
- Promoting clean granulation tissue to permit healing
- Meticulous cleansing and dressing (avoid soaps and sensitising preparations)
- Prevention and control of infection - antibiotics if cellulitis (cephalexin or erythromycin)
- Firm elastic compression bandage - use a minimal stretch bandage from base of toes to just below the knee. The degree of compression depends on the blood flow and is proportional to it
- Bed rest with elevation (if severe, 45 - 60 minutes twice a day and at night): ensure legs are elevated higher than the heart
- Encourage early ambulation
- Appropriate modification of lifestyle including weight reduction, smoking cessation (NB)
- Good nutrition includes a healthy balanced diet with ample protein and complex carbohydrates

Note: Firm compression is the single most important factor in the healing of venous ulcers.[6] Options include elastic stockings, elastic bandages, Unna's boots and legging orthoses.[9]

Cleansing/debridement agents

There are many cleansing agents, including:

- N saline, benzoyl peroxide and Intra Site Gel.

As a rule, avoid antiseptics which destroy cells, although Cadexomer iodine, which is a slow-release form of iodine, is non-toxic to tissues, reduces bacterial load and clears odorous slough. A good combination is normal saline cleansing followed by Intra Site Gel for debridement. Hydrogels such as Intra Site Gel, which have been found to be effective at debridement (including black necrotic areas), have generally replaced enzyme dressings.

Medicated occlusive bandages

There are several suitable occlusive paste bandages for ambulant patients, which ideally should be left on for 7 - 14 days. All contain zinc oxide with the indicated additives - Calaband (calamine), Ichthaband (ichthammol), Quinaband (clioquinol + calamine), Tarband (coal tar), Viscopaste and Zincaband (zinc oxide only).

Patch testing for an allergic response should be performed for a few days beforehand.

Pitfalls and other factors to be considered

- Treat the primary cause by surgery or other means (e.g. varicose veins, vascular insufficiency).
- If oedema, elevate legs and prescribe diuretics. An ulcer will not heal in the presence of significant ankle oedema.
- Be careful of allergy to local applications (e.g. zinc).
- Be careful of irritation from local applications (e.g. antibiotics). Antibiotic-impregnated dressings are not generally recommended.
- Avoid heavy packing of the wound.
- Consider graft ing (pinch skin or split thickness).
- Consider oxpentifylline 400 (Trental) for chronic occlusive arterial disease.

Post-healing and prevention of ulcers

- Encourage preventive measures, such as regular walking, good nutrition, no smoking, elevation of leg when resting, great care to avoid trauma.
- Apply emollients for varicose eczema.
- Wear a compression grade elastic stocking for varicose ulcers (e.g. Jobst Fast-Fit, Tensor Press).

A recommended treatment routine for a leg ulcer is presented inTable 115.4. It is desirable (for the outpatient) to leave the dressings and bandages in place for 1 week, perhaps 2 weeks, depending on the state of the dressing.

Table 115.4 A recommended leg ulcer treatment method

Clean with normal saline
• If slough, apply Intra Site Gel
• Dressing: non-adherent paraffin gauze
• Padding bandage (e.g. Velband)
• Occlusive paste bandage - Icthaband or Viscopaste (7 - 14 days), from base of toe to just below knee, plus compression bandage or Compression bandage (e.g. Elastocrepe or Eloflex) to just below knee
• Consider Tubigrip stockinette cover

Unna's boots

Unna's boots, which are boot-like dressings made of layers of gauze and Unna's paste (a mixture of 15% zinc oxide in a glycogelatin base), have an important place in the management of severe ulceration despite the discomfort and inconvenience for the patient. They appear to provide excellent fixed compression and to encourage healing, and one particular study showed their superiority to elastic stockings (7-3 weeks compared to 18-4 weeks).[10]

Principles of management for chronic ulcers are summarised in Table 115.5.

Table 115.5 Principles of management of chronic ulcers[11]

Ulcer type	Major management principles
Venous	Control venous insufficiency • compression bandage • improve calf muscle pump action (ambulation, exercises) • vertical leg drainage
Arterial	Vascular assessment for surgical intervention
Mixed venous/arterial	Vascular assessment for surgical intervention
Pressure	Eliminate or reduce pressure

DECUBITUS ULCERS (PRESSURE SORES)

Pressure sores tend to occur in elderly immobile patients, especially those who are unconscious, paralysed or debilitated. The cause is skin ischaemia from sustained pressure over a bony area, particularly the heels, sacrum, hips and buttocks. Poor general health, including anaemia, is a predisposing factor. Clinical features:

- preliminary area of fixed erythema at pressure site
- relatively sudden onset of necrosis and ulceration
- ulcer undermined at edges
- possible rapid extension of ulcers
- necrotic slough in base

Management

Prevention

- Good nursing care including turning patient every 2 hours.
- Regular skin examinations by the nursing and medical staff.
- Special care of pressure areas, including gentle handling.
- Special beds, mattresses (e.g. air-filled ripple) and sheepskin to relieve pressure areas.
- Good nutrition and hygiene.
- Control of urinary and faecal incontinence.
- Avoid the donut cushion.
- Avoid soaps

Treatment of ulcer

Use above measures, plus:

- clean base with saline solution (applied gently via a syringe) or Intra Site Gel
- general guidelines for dressings:
  - deep ulcers - alginates (e.g. Tegagel, Kaltostat)
  - shallow ulcers - hydrocolloids (e.g. Duoderm, CGF, Cutinova Hydro)
  - dry or necrotic ulcers - hydrogels (e.g. Intrasite, Solosite)
  - heavy exudative ulcers - foams (e.g. Lyofoam)
- give vitamin C, 500 mg bd
- give antibiotics for spreading cellulitis (otherwise of little use)
- healing is usually satisfactory but, if not, surgical intervention with debridement of necrotic tissue and skin graft ing may be necessary; this is very effective if the patient can cope

Undressing wounds

Removal of dressings from ulcerated wounds is very important. The contact layer should be removed slowly to prevent detachment of fragile epithelial surface cells and trauma to healthy granulation tissue.[12]

Role of honey

Honey has been advocated for centuries for healing ulcers. A particular type, Medihoney, is marketed. It provides moist healing and has anti-bacterial properties. Care has to be taken with over-moisturisation and maceration. Its role is still somewhat controversial as is the application of sugar, cromoglycate powder, maggots and hyperbaric oxygen.

TROPHIC ULCERS

Trophic ulcers are due to neuropathy causing loss of sensation (invariably diabetic) and usually follow an injury of which the patient was unaware.

A feature is a deep punched-out lesion (similar to ischaemic ulcers) over pressure points. A common site is the ball of the foot under the first metatarsal head, but the heel or a bunion may also be affected.

The ulcers may extend to the bone and into joints. They are prone to secondary infection.

Treatment is based on controlling the diabetes and clearing infection with appropriate antibiotics, but referral for surgical management is usually essential.

DERMATITIS ARTEFACTA AND NEUROTIC EXCORIATIONS

These self-inflicted ulcerated or erosive skin lesions have a psychological basis.

Dermatitis artefacta

In this condition, patients deny self-trauma and may have deep-seated psychological problems; or they may be malingering or manipulative for secondary gain.

Neurotic excoriations

These lesions, which are usually identical to the artefactual lesions, are caused by patients who admit to scratching, picking or digging at their skin. It occurs at times of stress and treatment is seldom successful. Treatment consists of counselling with CBT, a trial of antidepressants and topical antipruritics such as:

- coal tar solution (liquor picis carbonis) and menthol in sorbolene cream *or*
- menthol (0.5%) or phenol (1.0%) in aqueous cream

REFERENCES

1. Kelly R. Leg ulcers and wound healing. In: Dermatology Conference Notes. Melbourne: Combined Alfred Hospital/Skin and Cancer Foundation, 2002: 29.
2. Davis A, Bolin T, Ham J. *Symptom Analysis and Physical Diagnosis* (2nd edn). Sydney: Pergamon Press, 1990: 380 - 9.
3. Stacey MC. Chronic venous ulcers. Sydney: Med Observer, 1991; 29 March: 31 - 2.
4. Marley J (Chair). *Therapeutic Guidelines: Dermatology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2004: 237 - 47.
5. Beauregard S, Gilchrest BA. A survey of skin problems and skin care regimens in the elderly. Arch Dermatol, 1987; 123: 1638 - 43.
6. Fry J, Berry HE. *Surgical Problems in Clinical Practice*. London: Edward Arnold, 1987: 115 - 17.

7. Buxton P. *ABC of Dermatology*. London: Br Med J, 1989; 34 - 9.
8. Fitzpatrick JE. Stasis ulcers: update on a common geriatric problem. Mod Med Aust, 1990; June: 81 - 8.
9. Vernick SH, Shapiro D, Shaw FD. Legging orthosis for venous and lymphatic insufficiency. Arch Phys Med Rehab, 1987; 68: 459 - 61.
10. Hendricks WM, Swallows RT. Management of stasis leg ulcers with Unna's boots versus elastic support stockings. J Am Acad Dermatol, 1985; 12: 90 - 8.
11. Findlay D. Wound management and healing in general practice. Annual Update Course notes. Melbourne: Monash University, 1996; 13 - 16.
12. Rowland J. Pressure ulcers: a literature review and a treatment scheme. Aust Fam Physician, 1993; 22: 1819 - 27.

CHAPTER 116. COMMON LUMPS AND BUMPS

It will never get well if you pick it.

AMERICAN PROVERB

COMMON LUMPS AND BUMPS

Lumps and bumps are very common pre-sentations and the skin a very common site for neoplastic lesions. Most of these lesions only invade locally, with the notable exception of malignant melanoma. Pigmented skin tumours thus demand very careful consideration, although only a very few are neoplastic. The optimum time to deal with the problem and cure any skin cancer is at its first presentation. The family doctor thus has an important responsibility to screen these tumours and is faced with two basic decisions: the diagnosis and whether to treat or refer.

Most skin lumps are benign and can be left in situ, but the family doctor should be able to remove most of these lumps if appropriate and submit them for histological verification. The main treatment options available in family practice are: biopsy, cryotherapy, curette and cautery, excision or intralesional injections of corticosteroid.[1] A list of common and important lumps is presented in Table 116.1.

Table 116.1 Important lumps and their tissue of origin[3]

Skin and mucous membranes
• fibroepithelial polyp (skin tag)
• Epidermoid (sebaceous) cyst
• Implantation cyst
• Sebaceous hyperplasia
• Mucocele
• Hypertrophic scar and keloid
• warts and papillomas
• pox virus lumps
- molluscum contagiosum
- orf
- milker's nodules
• Seborrhoeic keratoses
• Granuloma annularae
• dermatofibroma
• solar keratosis/actinic keratosis
• keratoacanthoma
Malignant tumours
• basal cell carcinoma
• Squamous cell carcinoma
• Bowen's disorder
• Malignant melanoma
• Kaposi's sarcoma
• Secondary tumour

Subcutaneous and deeper structures

- Lipoma
- neurofibroma
- soft fibroma
- lymph node

Musculoskeletal

- Ganglion

SKIN CANCER

The three main skin cancers are the non-melanocyctic skin cancers - basal cell carcinoma (BCC) and squamous cell carcinoma (SCC) - and melanoma. The approximate relative incidence is BCCs 80%, SCCs 15 - 20%, and melanomas less than 5%.[2] The incidence of non-melanotic skin cancer is approximately 800 new cases per 100 000 population per year, and 25 per 100 000 for melanoma. About 80% of skin cancer deaths are due to melanoma and the rest mainly due to SCC.[2]

A DIAGNOSTIC APPROACH TO THE LUMP

As with any examination, the routine of look, feel, move, measure, auscultate and transilluminate should be followed.

The lump or lumps can be described thus:

- number
- site
- shape - regular or irregular
- size (in metric units)
- position
- consistency (very soft, soft, firm, hard or stony hard)
- solid or cystic
- mobility
- surface or contour
- special features
    - attachments (superficial/deep)
    - exact anatomical site
    - relation to anatomical structures
    - relation to overlying skin
    - colour
    - temperature (of skin over lump)
    - tenderness
    - pulsation (transmitted or direct)
    - impulse
    - reducibility
    - percussion
    - fluctuation (? contains fluid)
    - bruit
    - transilluminability
    - special signs
        - slipping sign

        - emptying sign of cavernous
        - haemangioma
    - spread
        - local
        - lymphatic
        - haematogenous
    - regional lymph nodes
    - ? malignancy (is it primary or secondary?)

Relation of the lump to anatomical structures[3]

The question ‘In what tissue layer is the lump situated?’ needs to be addressed.

- Is it in the skin? The lump moves when the skin is moved (e.g. epidermoid cyst).
- Is it in subcutaneous tissue? The skin can be moved over the lump. The slipping sign: if the edge of the lump is pushed, the swelling slips from beneath the finger (e.g. lipoma).
- Is it in muscle? The lump is movable when the muscle is relaxed but on contraction of the muscle this movement becomes limited.
- Is it arising from a tendon or joint? Movement of these structures may cause a change in the mobility or shape of the tumour.
- Is it in bone? The lump is immobile and best outlined with the muscles relaxed.

FIBROEPITHELIAL POLYPS

Synonyms: skin tags, acrochordon, benign squamous papilloma.

Clinical features:

- benign skin overgrowth
- increased incidence with age
- commonest on neck, axillae, trunk, groins
- no malignant potential
- can be irritating or unsightly to patient

Management

- Can leave or remove
- Snip off with scissors or bone forceps (Figure 116.1) *or*


(a)
skin tag
bone forceps
The base is crushed flush with the skin until the tag gives way
(b)
Bandaid is applied to wound
defect (usually bloodless)

Figure 116.1 Removal of skin tag using bone forceps

- tie base with fine cotton or suture material *or*
- diathermy base *or*
- apply liquid nitrogen (Figure 116.2)



Figure 116.2 Removal of skin tag by liquid nitrogen: a cotton bud soaked in liquid nitrogen is applied to the forceps, which grasp the tag firmly

These methods do not require local anaes-thetic.

EPIDERMOID (SEBACEOUS) CYST

Synonyms: 'pilar' cysts, keratinous cyst wens, epidermoid cysts, sebaceous cysts (similar in appearance).

Clinical features:

- firm to soft regular lump (usually round)
- fixed to skin but not to other structures (Figure 116.3a)



(a)
punctum with keratin
cyst attached to skin
sebaceous material
(b)
cyst outline
excised ellipse of skin

Figure 116.3 (a) Configuration of a sebaceous cyst (b), (c) Standard dissection of a large sebaceous cyst

- move with the skin
- found in hair bearing skin mainly on scalp - then face, neck, trunk, scrotum
- contains sebaceous material
- usually fluctuant
- may be a central punctum containing keratin
- tendency to inflammation

Management

If before puberty - think of polyposis coli

Can leave if small and not bothersome.

Surgical removal methods

There are several methods of removing epi-dermoid cysts after infiltrating local anaesthetic over and around the cyst. These include:

- *Method 1: Incision into cyst* Make an incision into the cyst to bisect it, squeeze the contents out with a gauze swab and then avulse the lining of the cyst with a pair of artery forceps or remove with a small curette.
- *Method 2: Incision over cyst and blunt dissection* Make a careful skin incision over the cyst, taking care not to puncture its wall. Free the skin carefully from the cyst by blunt dissection. When it is free from adherent subcutaneous tissue, digital pressure will cause the cyst to 'pop out'.
- *Method 3: Standard dissection* Incise a small ellipse of skin to include the central punctum over the cyst (Figure 116.3b). Apply forceps to this skin to provide traction for dissection of the cyst from the adherent dermis and subcutaneous tissue. Ideally, forceps should be applied at either end. The objective is to avoid rupture of the cyst. Inserting curved scissors (e.g. McIndoe's scissors), free the cyst by gently opening and closing the blades (Figure 116.3c). Bleeding is not usually a problem. When the cyst is removed, obliterate the space with subcutaneous Vicryl, Dacron, PDS or Monocryl (all preferable) or catgut. The skin is sutured with a vertical mattress suture to avoid a tendency to inversion of the skin edges into the slack wound. Send the cyst for histopathology.

Treatment of infected cysts

Incise the cyst to drain purulent material. When the inflammation has resolved completely the cyst should be removed by method 1 or method 3 (above).

IMPLANTATION CYST

Synonym: implantation dermoid.

Clinical features:

- small cystic swelling
- may be tender
- usually follows puncture wounds
- especially on finger pulp (e.g. hairdressers, sewers)
- contains mucus

Management

- Incision removal (similar to epidermoid cyst)

MUCOCELE

A mucous retention cyst.

Clinical features:

- a benign tumour
- cyst containing mucus
- appears spontaneously
- common on lips and buccal mucosa
- smooth and round
- yellow or blue colour

Management

- Incision removal

HYPERTROPHICSCAR

A hypertrophic scar is simply a lumpy scar caused by a nodular accumulation of thickened collagen fibres. It does not extend beyond the margins of the wound and regresses within a year.

Keloid

A keloid is a special type of hypertrophic scar that extends beyond the margins of the wound.

Clinical features:

- firm, raised, red-purple, skin overgrowth
- common on ear lobes, chin, neck, shoulder, upper trunk
- hereditary predisposition
- follows trauma, even minor (e.g. ear piercing)
- may be burning or itchy and tender

Management

- Prevention (avoid procedures in keloid-prone individuals)
- Intradermal injection of corticosteroids in early stages (2 - 3 months) or X-ray treatment of surgical wounds within 2 weeks of operation.[4]

WARTS AND PAPILLOMAS

Warts are skin tumours caused by the human papilloma virus (HPV). The virus invades the skin, usually through a small abrasion, causing abnormal skin growth. Warts are transmitted by direct or fomite contact and may be auto-inoculated from one area to another.[5]

Clinical features:

- average incubation period - 4 months
- increased incidence in children and adolescents
- peak incidence around adolescence
- occurs in all races at all ages
- about 25% resolve spontaneously in 6 months[5] and 70% in 2 years
- present as various types

Types of warts

These include common warts, plane warts, filiform warts (fine elongated growths, usually on the face and neck), digitate warts (finger-like projections, usually on scalp), genital and plantar warts (Figure 116.5).



Figure 116.5 Configuration of various types of warts

Common warts

Skin-coloured tumours with a rough surface, found mainly on the fingers, elbows and knees.

Plane warts

Skin-coloured, small and flat, occurring in linear clusters along scratch lines (Figure 116.4). Mainly occur on the face and limbs. Difficult to treat because they contain very few virus particles. Prone to Koebner phenomenon, which is seeding when a scratch passes through a plane wart.



Figure 116.4 Plane warts on the dorsum of the hand

Treatment options for warts

Topical applications:[5]

- salicylic acid - e.g. 5 - 20% in flexible collodion (apply daily or bd) or 16 - 17% salicylic acid + 16 - 17% lactic acid
- formaldehyde 2 - 4% alone or in combination
- podophyllotoxin 0.5% for anogenital warts - it is good on mucosal surfaces but does not penetrate normal keratin
- cytotoxic agents (e.g. 5-fluorouracil: very good for resistant warts such as plane warts and periungual warts)
- the immunomodulator, imiquimod

Cryotherapy

Carbon dioxide (- 56.5?C) or liquid nitrogen (- 195.8?C) destroys the host cell and stimulates an immune reaction.

Note:

- Excessive keratin must be pared before freezing.
- Results often disappointing.

Curettage

A most common treatment; some plantar warts can be removed under LA with a sharp spoon curette. The problem is a tendency to scar so avoid over a pressure area such as the sole of the foot.

Electrodissection

A high-frequency spark under LA is useful for small, filiform or digitate warts. A combination of curettage and electrodissection is suitable for large and persistent warts.

Vitamin A and the retinoids

- Topical retinoic acid (e.g. tretinoin 0.1% cream - Retin-A) is effective on plane warts
- Systemic oral retinoid, acitretin (Neotigason) for recalcitrant warts (with care)

Medication

- Consider cimetidine.

Specific wart treatment

The method chosen depends on the type of wart, its site and the patient's age.

- *Plantar warts*: refer to Chapter 68.
- *Genital warts*: podophyllotoxin 0.5% paint or imiquimod (see Chapter 110).
- *Filiform and digitate warts*: liquid nitrogen or electrodissection.
- *Plane warts*: liquid nitrogen; salicylic acid 20% co (e.g. Wartkil); consider 5-fluorouracil cream or Retin-A.
- *Common warts*: a recommended method:
    1. Soak the wart/s in warm soapy water.
    2. Rub back the wart surface with a pumice stone.
    3. Apply the paint (only to the wart; protect the surrounding skin with Vaseline). The paint: formalin 5%, salicylic acid 12%, acetone 25%, collodion to 100%.[4]
       Do this daily or every second day.
       Carefully remove dead skin between applications. or (preferable applications) (adult) 16% salicylic acid, 16% lactic acid in collodion paint (Dermatec, Duofilm), apply once daily (children) 8% salicylic acid, 8% lactic acid in collodion
- *Periungual warts (fingernails)*: consider 5-fluorouracil or liquid nitrogen with care. Always use a paint rather than ointment or paste on fingers.

POX VIRUS LUMPS

Skin tumours can be caused by pox viruses, some of which result from handling infected sheep, cows and monkeys and other animals such as deer. Hence, they are usually found in sheep shearers, farmers and zookeepers.

Molluscum contagiosum

This common pox virus infection can be spread readily by direct contact, including sexual contact (p. 1124). The incubation period is 2 - 26 weeks.

Clinical features:

- common in school-age children
- single or multiple (more common)
- shiny, round, pink-white papule (Figure 116.6)



Figure 116.6 Molluscum contagiosum with the round pink pearly appearance and central punctum

- hemispherical up to 5 mm
- central punctum gives umbilical look
- can be spread by scratching

Management

They are difficult to treat. Avoid using the bath - they spread to other body parts and those sharing the bath. Showering is preferable. There is a case for simply reassuring the family and waiting for spontaneous resolution.

Treatment options

- Liquid nitrogen with care (a few seconds) then dry dressings for 2 weeks
- Pricking the lesion with a pointed stick soaked in 1% or 2.5% phenol
- Application of 15% podophyllin in friar's balsam (compound benzoin tincture)
- Application of 30% trichloracetic acid
- Destruction by electrocautery or diathermy
- Ether soap and friction method
- Lifting open the tip with a sterile needle inserted from the side (parallel to the skin) and applying 10% povidone-iodine (Betadine) solution (parents can be shown this method and continue to use it at home for multiple tumours)
- If more localised, covering with a piece of Micropore or Leucosilk tape - change every day after showering (may take a few months). This method also prevents spread
- For large areas, aluminium acetate (Burow's solution 1:30) applied bd can be effective

Note: The extract of the Cantharis beetle (prepared as Canthrone) if available is reportedly very effective.

Orf

Orf is due to a pox virus and presents as a single papule or group of papules on the hands of sheep-handlers after handling lambs with contagious pustular dermatitis. The papules change into pustular-like nodules or bullae with a violaceous erythematous margin. It clears up spontaneously in about 3 - 4 weeks without scarring and usually no treatment is necessary.

Practice tip

Rapid resolution (days) can be obtained by an intralesional injection of triamcinolone diluted 50:50 in normal saline.[6]

Milker's nodules

In humans 2 - 5 papules appear on the hands about 1 week after handling cows' udders or calves' mouths. The papules enlarge to become tender grey nodules with a necrotic centre and surrounding inflammation (Figure 116.7). The patient can be reassured that the nodules are a self-limiting infection and spontaneous remission will occur in 5 - 6 weeks without residual scarring. One infection gives lifelong immunity.



Figure 116.7 Milker's nodule in a person who milks cows showing the grey nodule with the necrotic centre

Practice tip

Intralesional corticosteroid injection (as for orf).

SEBORRHOEIC KERATOSES

Synonyms: seborrhoeic wart, senile wart, senile keratoses (avoid these terms).

Clinical features:

- very common
- there is a variety of subtype
- increasing number and pigmentation with age > 40 years
- sits on skin, appears in some like a ‘sultana’ pressed into the skin (i.e. well-defined border)
- has a ‘pitted’ surface (Figure 116.8)



Figure 116.8 Seborrhoeic keratosis in a 70-year-old man. The large pigmented warty mass appears to sit on top of the skin. Photo courtesy Robin Marks

- may be solitary but usually multiple
- common on face and trunk, but occurs anywhere
- usually asymptomatic
- usually causes patients some alarm (confused with melanoma)

Management

- Usually nil apart from reassurance
- Does not undergo malignant change
- Can be removed for cosmetic reasons
- Light cautery to small facial lesions
- Freezing with liquid nitrogen (especially if thin) decolours the tumour
- 10% (or stronger) phenol solution applied carefully - repeat in 3 weeks
- Apply trichloroacetic acid to surface: instil gently by multiple pricks with a fine-gauge needle, twice weekly for 2 weeks
- May drop off spontaneously
- If diagnosis uncertain, remove for histopathology

STUCCO KERATOSES

This subtype of seborrhoek keratoses comprises multiple non-pigmented (often white) small friable keratoses over the lower legs. They can be treated with a topical keratolytic such as 3 - 5% salicyclic acid in sorbolene.

GRANULOMA ANNULARAE

Granuloma annularae are a common benign group of papules arranged in an annular fashion.

Clinical features:

- most common among children and young adults
- firm papules grouped in a ‘string of pearls’ pattern (Figure 116.9)



Figure 116.9 Granuloma annularae: this pearly papular tumour was long standing on this finger. After taking a small biopsy 20 mg of methylpredisolone acetate was injected into the tumour

- dermal nodules
- may be associated with minor trauma
- associated with diabetes
- usually on dorsum or sides of fingers (knuckle area), backs of hands, the elbows and knees

Management[7]

- Check urine/blood for sugar
- Give reassurance (they usually subside in a year or so)
- Cosmetic reasons:
  - first-line: potent topical corticosteroids ± occlusion, apply bd for minimum of 4 - 6 weeks
  - if ineffective: intradermal injection into the extending outer margin of triamcinolone 10% or similar corticosteroid (dilute equal volume with N saline); can repeat at 6 weekly intervals if effective

DERMATOFIBROMA

Synonyms: sclerosing haemangioma; histio-cytoma.

This is a common pigmented nodule arising in the dermis due to a proliferation of fibroblasts, believed to develop as an abnormal response to minor trauma including insect bites. The nodule gives a characteristic button-like feel and dimpling when laterally compressed (pinched) from the side with the fingers.

Clinical features:

- usually multiple
- firm, well-circumscribed nodules
- oval, 0.5 - 1 cm in diameter
- freely mobile over deeper structures
- slightly raised in relation to skin
- mainly on limbs, especially legs
- may itch
- mainly in women

- variable colour, pink or brown, tan or grey or violaceous
- characteristic ‘dimple’ sign on pinching margins

Treatment

- Reassurance
- Simple excision if requested

SOLAR KERATOSIS

Solar keratoses (actinic keratoses) are reddened, adherent, scaly thickenings occurring on light-exposed areas, with a potential for malignant change, especially on the ears.

Clinical features:

- sun-exposed fair skin
- mainly on face, ears, scalp (if balding), forearms, dorsum of hands (especially) (Figure 116.10)



Figure 116.10 Solar keratoses showing the reddened scaly thickenings on sun-exposed areas. Biopsy of one of the lesions proved squamous cell carcinoma

- dry, rough, adherent scale
- discomfort on rubbing with towel
- scale can separate to leave oozing surface
- a small proportion undergo malignant change

Terminology

Solar, keratosis = sun spots

Solar lentinges = age spots or liver spots

Management

- Reduced exposure to sunlight
- Can disappear spontaneously
- Liquid nitrogen if superficial (don't freeze unless inflamed i.e. red) *or*
  5-fluorouracil 5% cream daily for 3 - 4 weeks *or*

imiquimod, once daily 3 times a week up to 16 weeks

- Surgical excision for suspicious and ulcerating lesions
- Biopsy if doubtful

KERATOACANTHOMA

Keratoacanthomas (KA), which are rapidly evolving tumours of keratinocytes, occur singly on light-exposed areas. The major problem is differentiation from SCC, especially if on the lip or ear. The relative growth rates of three types of skin tumours are shown in Figure 116.11.



Figure 116.11 Relative growth rates of three types of skin tumours: keratoacanthoma, squamous cell carcinoma and basal cell carcinoma

Clinical features:

- rapidly growing lesion on sun-exposed skin
- raised crater with central keratin plug (Figure 116.13 and Figure 116.12)



Figure 116.13 A typical keratoacanthoma



Figure 116.12 Keratoacanthoma: this tumour, with its central plug appeared suddenly on the face of a 63-year-old man. It may be confused with squamous cell carcinoma. Surgical excision is appropriate treatment

- grows to 2 cm or more
- arises over a few weeks, remains static, then spontaneously disappears after about 4 - 6 months; can leave a big scar
- can be confused with SCC

Management

- Remove by excision - perform biopsy
- If clinically certain - curettage/diathermy
- Treat as SCC (by excision) if on lip/ear

The recommended treatment is surgical excision and histological examination. Ensure a 2 - 3 mm margin for excision. Most patients will not tolerate a tumour for 4 - 6 months on an exposed area such as the face while waiting for a spontaneous remission. Also, if it is an SCC, a potentially lethal cancer has remained in situ for an unnecessarily long period.

SEBACEOUS HYPERPLASK

Sebaceous hyperplasia presents as, single or multiple nodules on the face, especially in older people. The nodules are small, yellow-pink, slightly umbilicated and are found in a similar distribution to basal cell carcinomas for which that may be mistaken. There is no need for surgical excision.

BASAL CELL CARCINOMA

Clinical features:

- most common skin cancer (80%)
- age: usually > 35 years
- more frequent in males
- mostly on sun-exposed areas: face (mainly), neck, upper trunk, limbs (10%) (Figure 116.15)



Figure 116.15 Typical areas in which basal cell carcinomas occur

- may ulcerate easily = 'rodent ulcer'
- slow-growing over years
- has various forms: nodular, pigmented, ulcerated, etc.
- stretching the skin demarcates the lesion, highlights pearliness and distinct margin
- does not metastasise via lymph nodes or bloodstream
- local spread is a problem
- can spread deeply if around nose, eye or ear

Clinical types

1. Cystic nodular - translucent or pale grey
2. Ulcerated - nodular BCC that has necrosed centrally
3. Pigmented - usually spotted, may be all black
4. Superficial - erythematous scaly patch, may be misdiagnosed as eczema or psoriasis
5. Morphoeic (fibrotic) - scar-like, poorly defined margin
6. Common: pearly edge, telangiectasia, ulcerated (Figure 116.14)



Figure 116.14 Basal cell carcinoma showing a pearly nodular appearance with telangiectatic vessels. *Photo courtesy Robin Marks*

Management

- Excision (3 mm margin) is best
- If not excision, do biopsy before other treatment
- Radiotherapy is an option, especially in frail people
- Moh's micrographic surgery - a form of surgical treatment suitable for large or recurrent tumours or those in a site when maximal normal tissue needs to be preserved.
- Photodynamic therapy - response rate is > 90% for nodular and superficial BCCs.

Note: Avoid cryotherapy; imiquimod may be an option. To biopsy a BCC, do a shave biopsy, not a punch biopsy.

SQUAMOUS CELL CARCINOMA

SCC is an impozrtant malignant tumour of the epidermis; it is also found on sun-exposed areas, especially in fair-skinned people. It tends to arise in premalignant areas such as solar keratoses, burns, chronic ulcers, leucoplakia and Bowen's disorder, or it can arise de novo.

Note: Although BCC and SCC are related to cumulative sun exposure they are not always found in sun-exposed areas.

Clinical features:

- usually > 50 years
- initially firm thickening of skin, especially in solar keratosis
- surrounding erythema
- the hard nodules soon ulcerate (Figure 116.16)



Figure 116.16 Squamous cell carcinoma. This recurrent non-healing lesion on the index finger of a 58-year-old man had raised hard edges and was fixed to tendon and bone. Treatment was by surgical amputation of the finger

- occurs on the hands and forearms and the head and neck (see Figure 116.17)



Figure 116.17 Common sites of squamous cell carcinoma

- ulcers have a characteristic everted edge
- capable of metastases and may involve regional nodes
- SCCs of ear, lip, oral cavity, tongue and genitalia are serious and need special management

Management

- Early excision of tumours < 1 cm with a 4 mm margin (in most cases), to deep fat level.
- Referral for specialised surgery and/or radiotherapy if large, in difficult site or lymphadenopathy.
- SCCs of the ear and lip, which have considerably more malignant potential, can be excised by wedge excision.
- There is no alternative to surgery if the SCC is over cartilage - central nose or helix

NB: Surgery is the treatment of choice for most tumours, cryotherapy and curettage is not.

BOWEN'S DISORDER

Bowen's disorder begins as a slowly enlarging, sharply demarcated, thickened red plaque, especially on the lower legs of females. It may resemble solar keratosis or a patch of psoriasis. It remains virtually unchanged for months or years. It may become very crusty, ulcerate or bleed. It has a potential for malignant change since it is a full thickness SCC in situ.

Management

- Biopsy first for diagnosis
- Wide surgical excision if small
- Skin grafting may be required
- Cryotherapy by double freeze thaw technique
- Imiquimod is considered promising (awaiting trials)

Note: Biopsy a single patch of suspected psor-iasis or dermatitis not responding to topical steroids.

LUMPS ON EARS

Lumps on ears, especially on the helix, demand close attention. SCCs which arise here have up to 17 times the ability to metastasise and demand early wedge resection

Causes of ear lumps include:

- solar keratosis
- dasal cell carcinoma
- squamous cell carcinoma
- keratoacan throw
- gonty tophi
- chondrodermatitis nodulans helicus

Chondrodermatitis nodularis helicus

This lump, which is not a neoplasm, presents as a painful nodule on the most prominent part of the helix on ante helix of the ear. (Figure 116.18). It is seen more often in men while it is found more often on the ante helix in women. It Is caused by seen damage. It is to logically a thickened epidermis overlies inflamed cartilage. It looks like a small corn, is tender, and affects sleep if that side of the head lies on the pillow. If cryotherapy fails, wedge resection under local anaesthetic is an effective treatment.



Figure 116.18 Typical sites of chondrodermatitis nodularis helicus

MALIGNANT MELANOMA

These are usually enlarging pigmented lesions with an irregular notched border. Refer to Chapter 117 on pigmented skin lesions.

SECONDARY TUMOUR

These complex tumours may metastasise from the lung, melanoma or bowel and may arise in surgical scars (e.g. for carcinoma of the breast).

KAPOSI'S SARCOMA

Kaposi's sarcoma presents as brownish-purple papules on the skin and mucosa. Apart from the well-known presentation in immuno-compromised individuals, it is seen as a primary tumour mostly in elderly men of central or eastern European origin.

LIPOMA

Lipomas are common benign tumours of mature fat cells situated in subcutaneous tissue.

Clinical features:

- soft and may be fluctuant
- well defined; lobulated (Figure 116.19)



Figure 116.19 Lipoma: the 66-year-old woman had a longstanding soft, fluctuant rubbery lump of 18 years. It was surgically removed for cosmetic reasons

- rubbery consistency
- may be one or many
- painless
- most common on limbs (especially arms) and trunk
- can occur at any site

Management

- Reassurance about benign nature
- Removal for cosmetic reasons or to relieve discomfort from pressure

Surgical excision

Many lipomas can be enucleated using a gloved finger, but there are a few traps: some are deeper than anticipated, and some are adjacent to important structures such as large nerves and blood vessels. Others are tethered by fibrous bands, and can recur. Recurrence is also possible if excision is incomplete.

Caution: Lipomas on back (don't shell out early) If > 5 cm consider referral

Note: Ultrasound is good at assessing depth of lipok.

Method

Principle: cut, squeeze, 'pop'

1. Outline the extent of the lipoma and note its anatomical relationships.
2. Infiltrate the area with 1% lignocaine with adrenaline (include the deepest part of the lipoma).
3. Make a linear incision (Figure 116.21a) in the overlying skin, preferably in a natural crease line for about two-thirds of its length. The lipoma should bulge through the wound. For large lipomas, incise an ellipse of skin (Figure 116.21b).



Figure 116.21 (a) Linear incision for small lipomas; (b) elliptical incision for large lipomas; (c) gloved-finger dissection to bring the lipoma to the surface; (d) blunt scissors dissection to free the lipoma from tethering fibrous bands

4. Insert a gloved finger between the skin and fatty tumour to determine whether it will shell out. It is important to seek the outer edge of each lobule, dissect it and bring it to the surface (Figure 116.21c).
5. If necessary, insert curved scissors and use a blunt opening action to free any fibrous bands tethering the lipoma (Figure 116.21d).
6. Ensure that all the fatty tissue is removed. Send it for histological examination.
7. Use a gauze swab to control bleeding and remove debris from the dead space.
8. Close the dead space with interrupted catgut sutures.
9. Close the skin with interrupted or subcuticular sutures.

NEUROFIBROMA

These benign tumours are firm (sometimes soft) painless subcutaneous lumps aligned length-wise in the long axis of the limb in relation to peripheral nerves (Figure 116.20). The lumps are more mobile from side to side than along the long axis. Some are tender to pressure with associated pain and paraesthesia on the nerve distribution.



Figure 116.20 Neurofibroma. This mobile firm subcutaneous lump was tender to firm pressure.

BURSAE

Bursae are cystic sacs between the skin and an underlying bony prominence or sacs of gelatinous fluid that separate and aid gliding of adjacent tendous and ligaments.

GANGLION

Ganglia are firm cystic lumps associated with joints or tendon sheaths.

Clinical features:

- deep subcutaneous lumps
- around joints or tendon sheaths (Figure 116.22)



Figure 116.22 Ganglion of wrist - firm, immobile and translucent. It was eventually treated by aspiration of gelatinous fluid followed by infusion of 40-mg methyl prednisolone acetale

- mostly around wrists, fingers, dorsum of feet
- immobile, fixed to deep tissues
- translucent
- contain viscid gelatinous fluid
- associated with arthritis and synovitis
- may disappear spontaneously
- recurrences common

Management

- Can be left - wait and see
- Do not ‘bang with a Bible’
- Needle aspiration and steroid injection *or* surgical excision (can be difficult)
- Suture compression technique: a larger gauge catgut suture is inserted through the middle of the ganglion and firmly tied over it. Side pressure may express the contents through the needle holes. Remove the knot 12 days later.

Injection treatment of ganglia

Ganglia have a high recurrence rate after treatment, with a relapse rate of 30% after surgery. A simple, relatively painless and more effective method is to use intralesional in-jections of long-acting corticosteroid, such as methylprednisolone acetate.[8]

Method

1. Insert a 21-gauge needle attached to a 2 mL or 5 mL syringe into the cavity of the ganglion.
2. Aspirate some (not all) of its jelly-like contents, mainly to ensure the needle is in situ.
3. Keeping the needle exactly in place, swap the syringe for an insulin syringe containing up to 0.5 mL of steroid.
4. Inject 0.25 - 0.5 mL (Figure 116.23).



Figure 116.23 Injection treatment of ganglion

5. Rapidly withdraw the needle, pinch the overlying skin for several seconds and then apply a light dressing.
6. Review in 7 days and, if still present, repeat the injection using 0.25 mL of steroid.

Up to six injections can be given over a period of time, but 70% of ganglia will disperse with only one or two injections.[8]

SOME PREFERRED THERAPEUTIC OPTIONS

Liquid nitrogen therapy

Ideally, liquid nitrogen is stored in a special, large container and decanted when required into a small thermos flask or spray device.

The easiest method of application to superficial skin tumours (Table 116.2) is via a ball of cotton wool rolled rather loosely on the tip of a wooden applicator stick. The ball of cotton wool should be slightly smaller than the lesion, to prevent freezing of the surrounding skin.

Table 116.2 Superficial skin tumours suitable for cryotherapy

Warts (plane, periungual, plantar, anogenital)
Skin tags
Seborrhoeic keratoses
Molluscum contagiosum
Sebaceous hyperplasia
Solar keratoses

Method (basic steps)

1. Inform the patient what to expect.
2. Pare excess keratin with a scalpel.
3. Use a cotton wool applicator slightly smaller (not larger - see Figure 116.24a) than the lesion.



Figure 116.24 Application of liquid nitrogen: (a) applicator too large; (b) correct size and approach of applicator; (c) correct size but wrong position of applicator

4. Immerse it in nitrogen until bubbling ceases.
5. Gently tap it on the side of the container to remove excess liquid.
6. Hold the lesion firmly between thumb and forefinger.
7. Place applicator vertically (Figure 116.24b) on tumour surface.
8. Apply with firm pressure: do not dab.
9. Freeze until a 2 mm white halo appears around the lesion.

Explain likely reaction to patient, such as the appearance of blisters (possibly blood blisters). The optimal time for retreatment of warts is in 2 - 3 weeks (not longer than 3 weeks).

Biopsies

There are various methods for taking biopsies from skin lesions. These include scraping, shaving and punch biopsies, all of which are useful but not as effective or safe as excisional biopsies.

Shave biopsies

This simple technique is generally used for the tissue diagnosis of premalignant lesions and some malignant tumours, but not melanoma.

Method

1. Infiltrate with LA.
2. Holding a number 10 or 15 scalpel blade horizontally, shave off the tumour just into the dermis (Figure 116.25)



Figure 116.25 Shave biopsy

3. Diathermy may be required for haemostasis.

The biopsy site usually heals with minimal scarring.

Punch biopsy

This biopsy has considerable use in general practice, where full-thickness skin specimens are required for histological diagnosis. (Good quality disposable biopsy punches are available from Derma Tech Laboratories.)

Method

1. Clean the skin.
2. Infiltrate with LA.
3. Gently stretch the skin between the finger and thumb to limit rotational movement.
4. Select the punch (4 mm is the most useful size) and hold it vertically to the skin.
5. Rotate (in a clockwise, screwing motion) with firm pressure to cut a plug about 3 mm in depth (Figure 116.26). Remove the punch.



Figure 116.26 Punch biopsy

6. Use fine-toothed forceps or a tissue hook to grip the outer rim of the plug.
7. Exert gentle traction and undercut the base of the plug parallel to the skin surface, using fine-pointed scissors or a scalpel.
8. Place the specimen in fixative.
9. Secure haemostasis by firm pressure or by diathermy.
10. Apply a dry dressing or a single suture to the defect.

Steroid injections into skin lesions

Suitable lesions for steroid injections are:

- plaque psoriasis
- granuloma annulare
- hypertrophic scars (early development)
- keloid scars (early development)
- alopecia areata
- lichen simplex chronicus
- necrobiosis lipoidica
- hypertrophic lichen planus
- orf and milker's nodules

Triamcinolone is the appropriate long-acting corticosteroid (10 mg/mL). It may be diluted in equal quantities with saline.

Method

1. The steroid should be injected into the lesion (not below it).
2. Insert a 25- or (preferably) 27-gauge needle, firmly locked to a small insulin-type 1 mL syringe, into the lesion at the level of the middle of the dermis (Figure 116.27).



Figure 116.27 Injection of corticosteroid into mid-dermis

3. High pressure is required with some lesions (e.g. keloid).
4. Inject sufficient steroid to make the lesion blanch.
5. Several sites will be needed for larger lesions, so preceding LA may be required in some instances. Avoid infiltration of steroid in larger lesions: use multiple injections.

Elliptical excisions

Small lesions are best excised as an ellipse. Generally, the long axis of the ellipse should be along the skin tension lines identified by natural wrinkles.

The intended ellipse should be drawn on the skin (Figure 116.28). The placement will depend on such factors as the size and shape of the lesion, the margin required (usually 2 - 3 mm) and the skin tension lines.



Figure 116.28 Ellipse excision

General points

- The length of the ellipse should be three times the width.
- This length should be increased (say, to four times) in areas with little subcutaneous tissue (dorsum of hand) and high skin tension (upper back).
- A good rule is to obtain an angle at each end of the excision of 30° or less.
- These rules should achieve closure without 'dog ears'.

Excisions on the face

It is important to select optimal sites for elliptical excisions of tumours of the face. As a rule it is best for incisions to follow wrinkle lines and the direction of hair follicles in the beard area. Therefore, follow the natural wrinkles in the glabella area, the 'crow's feet' around the eye, and the nasolabial folds (Figure 116.29). To determine non-obvious wrinkles, gently compress the relaxed skin in different directions to demonstrate the lines.



Figure 116.29 Recommended lines for excisions on face Adapted From J.S. Brown, *Minor Surgery: A Text And Atlas*. London: Chapman & Hall, 1986

For tumours of the forehead make horizontal incisions, although vertical incisions may be used for large tumours of the forehead. Ensure that you keep your incisions in the temporal area quite superficial, as the frontal branch of the facial nerve is easily cut.

WHEN TO REFER

Referral should be considered for:

- uncertainty of diagnosis
- suspicion of melanoma
- tumours larger than 1 cm
- recurrent tumours, despite treatment
- incompletely excised tumours, especially with poor healing
- doubts about appropriate treatment
- recommended treatment beyond skills of practitioner
- frequent multiple tumours, eg. organ transplant patients
- squamous cell carunoma on the lip or ear
- infiltrating or scar like morphoeic BCC, particularly those on the nose or around the nasal labial fold
- cosmetic concerns, such as lesions in the upper chest and upper arms where keloid scarring is a potential problem.

KEY POINT

- Uncomplicated small tumours are best removed by an elliptical excision with a 3 mm margin for BCC and a 4 mm margin for SEC.
- Caution should be used in the management of tumours on the face, including the ears and lips.

REFERENCES

1. Paver R. The surgical management of cutaneous tumours in general practice. Mod Med Aust, 1991; March: 43 - 51.
2. Marks R. Skin cancer. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 469 - 72.
3. Davis A, Bolin T, Ham J. *Symptom Analysis and Physical Diagnosis* (2nd edn). Sydney: Pergamon Press, 1990: 302 - 6.
4. de Launey WE, Land WA. *Principles and Practice of Dermatology* (2nd edn). Sydney: Butterworths, 1984: 280 - 1.
5. Berger P. Warts: how to treat them successfully. Mod Med Aust, 1990; August: 28 - 32.
6. Reddy J. Intralesional injection for orf: a practice tip. Aust Fam Physician, 1993; 22: 65.
7. Marley V (Chair). *Therapeutic Guidelines: Dermatology* (Version 1). Melbourne: Therapeutic Guidelines Ltd, 2004; 253 - 4.
8. La Villa G. Methylprednisolone acetate in local therapy of ganglion. Clin Therapeutics, 1986; 47: 455 - 7.

CHAPTER 117. PIGMENTED SKIN LESIONS

> The skin calls for the faculty of close observation and attention to detail.
>
> Louis A. Duhring (1845 - 1913), valedictory address, University of Pennsylvania Medical School

PIGMENTED SKIN LESIONS

The management of pigmented skin lesions is a constant concern for all practitioners and requires careful evaluation based on the natural history of these lesions and the increasing incidence of malignant melanoma in particular.

Most pigmented lesions are benign and include simple moles or melanocytic naevi, seborrhoeic keratoses, freckles and lentigines. Reassurance is all that is necessary in the management of these problems.

However, one-third of all melanomas arise in preexisting naevi, many of which are dysplastic, and it is the recognition and removal of such naevi that is so important in the prevention of melanoma.[1]

Malignant melanoma is doubling in incidence each decade, which is an alarming statistic considering the public education programs about the hazards of sun exposure. Of equal interest is the fact that the cure rate for melanoma is also increasing, reflecting earlier diagnosis and treatment. The most important factor in management is early detection.[2]

A classification of pigmented skin lesions is given in Table 117.1.

Table 117.1 Classification of pigmented skin lesions

Non-melanocytic • Pigmented basal cell carcinoma • Seborrhoeic keratoses (p. 1205) • Solar keratoses (p. 1206) • Dermatofibroma (p. 1206) • Pyogenic granuloma • Foreign body granuloma • Talon noir (black heel) • Tinea nigra • Becker's naevus **Melanocytic**
Non-melanoma • Freckles • Lentigines • Naevi **1** congenital **2** acquired - junctional→compound→intradermal - dysplastic - halo - blue - Spitz

Melanoma

1 Lentigo maligna (Hutchinson's melanotic freckle)

2 Superficial spreading melanoma

3 Nodular melanoma

4 Acral lentiginous melanoma

KEY FACTS AND CHECKPOINTS

- The incidence of melanoma is greatest in white Caucasians and increases with proximity to the equator.
- The early diagnosis and treatment of melanoma profoundly affects the prognosis.
- Melanoma is extremely rare before puberty.
- Currently the greatest rate of increase is in men > 55 years.
- Most people have 5 - 10 melanocytic naevi on average.
- Multiple dysplastic naevi carry a higher risk of malignant change, which may occur in young adults. Such patients require regular observation (with photography).

PYOGENIC GRANULOMA

Synonyms: granuloma, granuloma telangiect-aticum, acquired haemangioma.

A pyogenic granuloma is a vascular lesion (without pus) due to a proliferation of capillary vessels. It is considered to be an abnormal reaction to minor trauma.

Clinical features:

- common in children and young adults
- usually on hands and face
- bright red ‘raspberry’-like lesion
- raised, sometimes pedunculated
- friable - bleeds easily
- Beware of misdiagnosing it for a nodular melanoma.

Management

It must be distinguished from amelanotic melanoma or anaplastic SCC. Shave biopsy with electrocautery of base. The specimen must be sent for histological examination. They are prone to recur.

TALON NOIR (‘BLACK HEEL’)

Talon noir is a black spotted appearance on the heel and is common in sportspeople. A similar lesion (probably smaller) is often found on the other heel.

‘Black heel’ is formed by small petechiae caused by the trauma of the sharp turns required in sport: shearing stress on the skin of the heel produces superficial bleeding. The diagnosis can be confirmed by gentle paring of the callus to reveal the multiple small petechial spots in the epidermis; these are then pared away. If there is doubt about the diagnosis (malignant melanoma is the main differential diagnosis), the lesion should be excised.

TINEA NIGRA

Tinea nigra is characterised by solitary black macular lesions on the palm or sole. The simple technique of taking skin scrapings to reveal fungal elements will allow easy differentiation from malignant melanoma.

BECKER'S NAEVUS

Becker's naevus is a faint, brown, diffuse pigmented area with a component of coarse hairs and is usually found on the shoulder and upper trunk. It occurs mainly in boys around puberty. It is not a birthmark, it is benign and reassurance is appropriate.

FRECKLES

Freckles are small, brown flat macules (usually < 0.5 cm), coloured by excessive epidermal melanin without any increase in the number of naevus cells (melanocytes). They occur mainly on light-coloured skin and tend to darken in summer and fade in winter. Cosmetic improvement can be achieved through the use of sunscreens.

LENTIGINES

Lentigines are small rounded, brown to black macular areas ranging from 1 mm to 1 cm or more across. They are very common and may appear in childhood as a few scattered lesions, often on areas not exposed to the sun. In the elderly, lentigines often develop on sun-damaged skin, usually on the backs of the hands (so-called ‘liver spots’) and on the face.

Unlike freckles they have increased numbers of melanocytes.

Management

- Treatment is usually unnecessary. Liquid nitrogen or excision can be used for cosmetically unacceptable lesions. Sunscreens are needed to prevent further darkening of existing lesions.
- Otherwise, rather than use ‘fade cream’, use fresh lemon juice. Squeeze lemon juice (1/2 lemon) into a small bowl and apply the juice to the spots daily. Continue for 8 weeks.
- Apply tretinoin 0.05% cream daily at night, if necessary.
- Laser for severe cases.



Figure 117.1 Pyogenic granuloma showing bright red friable tumour on face. It followed a puncture from a spiky plant in the garden. *Photo courtesy Robin Marks*

CONGENITAL MELANOCYTIC NAEVI

These moles are present at birth and are sometimes large.

Clinical features:

- variable colour - brown to black
- sometimes hairy and protruding
- increased risk of malignant change (especially in larger ones)

COMMON ACQUIRED NAEVI

These are the common moles for which an opinion is so often sought. The moles are localised benign proliferation of naevus cells. There may be a sharp increase in numbers during pregnancy. New lesions appear less frequently after the age of 20 years. The types are junctional, compound and intradermal. Naevi in children are usually the junctional type with proliferating naevus cells clumped at the dermoepidermal junction. With time the naevus cells ‘move’ into the dermis. A compound naevus has both junctional and dermal elements. With maturation all the naevus cells move into the dermis. Refer to Figure 117.3.



Figure 117.3 Comparison of melanocyte (naevus cell) distribution in various common acquired naevi

Clinical features

Junctional

- Usually < 5 mm
- Circular-shaped macules
- May be slightly elevated
- Colour usually brown to black
- May be ‘fuzzy’ border

Most naevi of the palms, soles and genitals are junctional but there is no evidence to support the traditional view that naevi in these sites have more malignant potential.[2]

Compound

- Dome-shaped, slightly raised pigmented nodules
- Up to 1 cm in diameter
- Colour varies from light to dark brown/black, but lighter than junctional naevi (Figure 117.2)
- Most are smooth but surface can be rough or verrucoid

- Larger ones may be hairy, especially after puberty
- Become 'flesh'-coloured in time



Figure 117.2 Compound naevus. This pigmented lesion which was benign has been present since birth. At puberty there is often a rapid increase in size and colour. *Photo courtesy Robin Marks*

Intradermal

- Look like compound but less pigmented
- Often skin-coloured
- May evolve to pink or brown senile nodules or to soft pedunculated tags

Malignant potential of common acquired melanocytic naevi

- *Junctional:* have significant potential to undergo malignant change (as long as junctional activity is present)
- *Compound:* very rarely undergo malignant change
- *Intradermal:* these are totally benign lesions

Management

- Provide appropriate reassurance.
- Observe.
- If lesion is changing or there is uncertainty, perform surgical excision (2 - 3 mm margin) for histopathology.

HALO NAEVUS

A halo naevus consists of a depigmented halo around a central melanocytic naevus (Figure 117.4). It is the result of an autoimmune reaction. The central naevus gradually involutes. It tends to occur around puberty. Multiple halo naevi are often seen on the trunk of adolescents.



Figure 117.4 Halo naevus in a child. The central lesion is usually a benign pigmented naevus. *Photo courtesy Robin Marks*

Caution: A halo can occur around a melanoma.

Management

Measure the lesion. Reassure and do nothing, as it usually disappears over the next few years; if doubtful at all, remove and obtain histological diagnosis.

BLUE NAEVUS

A blue naevus presents as a solitary slate-grey-blue dermal lesion. Blue naevi usually present in childhood and adolescence on the lower back and buttocks and the limbs, especially dorsa of the hands and feet. Malignant change is rare. often excised for cosmetic reasons.

SPITZ NAEVUS (BENIGN JUVENILE MELANOMA)

Spitz naevi are also called benign juvenile melanomas or spindle cell naevi.

Clinical features:

- solitary pigmented or erythematous nodules
- often appear in children, usually 4 - 8 years
- develop over 1 - 3 months
- well circumscribed, dome-shaped lesions

Management

Surgical excision is treatment of choice (because of rapid growth and best ‘reassurance’ policy).

DYSPLASTIC MELANOCYTIC NAEVI

These are large irregular moles which appear predominantly on the trunk in young adults (Figure 117.5). They can be familial or sporadic and are markers of an increased risk of melanoma, rather than necessarily being premalignant lesions. Even so, melanoma may arise within these lesions more frequently than would be expected by random chance.[3]

They are considered to be intermediate between benign naevi and melanoma.



Figure 117.5 Dysplastic melanocytic naevi. These lesions may have ill-defined borders and irregular pigmentation. *Photo courtesy Robin Marks*

Clinical features:

- age: adolescence onwards
- large > 5 mm (variable size)
- most common on trunk
- irregular and ill-defined border
- irregular pigmentation
- background redness
- variable colours - brown, tan, black, pink, red
- variation of colours within the naevus
- most are stable and do not progress to melanoma

DYSPLASTIC NAEVUS SYNDROME

The presence of multiple, large, irregular pigmented naevi, mainly on the trunk, presents a difficult management problem, especially if there is a family history of melanoma. The lifetime risk of melanoma may approach 100% for such patients.

Management

Use a follow-up program (similar to excised early melanoma) of 6 monthly review for 2 years (3 monthly if family history of melanoma) and yearly there after, provided no lesions become malignant during the first 2 years. During this time the patient and family should become well versed in surveillance. Apart from measurement, good professional quality photographs of areas of the body or specific lesions of concern may also be helpful.

Any suspicious lesions should be excised for histological examination.

Advice to patients

To decrease your chances of getting a melanoma, you should protect yourself from the sun. These rules should be followed:

- Try to avoid direct sunlight when the sun is at its strongest (from 10 am to 2 pm).
- Always wear a broad-brimmed hat and T-shirt in the sun.

- Use a factor 15 or more sunscreen on exposed skin and renew the sunscreen regularly.
- Sunbaking might give you a good tan but it is also going to increase your chances of getting a melanoma, so you should avoid it.

MELANOMA

The early diagnosis of melanoma is vital to outcome. Thickness of a melanoma when it is removed is the major factor determining prognosis: it is vital to detect melanomas when they are in the thin stage and look like an unusual freckle.

In Australia, only about 30% of melanomas develop in pre-existing melanocytic naevi (moles).[2, 3] The majority arise in apparently normal skin. Initially the tumour tends to spread laterally in many cases and it should be removed at this stage when it is easily cured. An irregular border or margin is suggestive of the tumour.

Clinical features

- Typical age range 30 - 50 years (average 40 years)
- can occur anywhere on the body - more common
    - lower limbs in women
    - upper back in men
- often asymptomatic
- can bleed or itch

Change

The sign of major importance is a recent change in a ‘freckle’ or mole:

- change in size - at edge or thickening
- change in shape
- change in colour - brown, blue, black, red, violet, white, including combinations
- change in surface
- change in the border
- bleeding or ulceration
- other symptoms (e.g. itching)
- development of satellite nodules
- lymph node involvement

Types of melanoma

Lentigo maligna[3]

Lentigo maligna (Hutchinson's melanotic freckle) is a slow-growing form of intraepidermal melanoma which occurs on areas exposed to light (usually the face), predominantly in the elderly (Figure 117.6). If allowed to remain it may become invasive and the prognosis will be similar to that for other invasive melanomas. These lesions have all the variations in size, shape and colour of superficial melanomas. Lentigo maligna should be excised.



Figure 117.6 Lentigo maligna in an 72-year-old man. Excision is recommended as it is a form of intraepidermal melanoma. *Photo courtesy Robin Marks*

Superficial spreading melanoma

Like lentigo maligna, the initial growth is in a lateral or radial intraepidermal manner, rather than in an invasive downward or vertical manner (Figure 117.7). It exhibits a striking colour variation. It accounts for 70% of melanomas. It can be detected early by biopsy - shave is preferred to punch (less sampling error). Excisional biopsy is preferable.



Figure 117.7 Superficial spreading melanoma with an irregular border which has altered and variable colours. Require excision. *Photo courtesy Robin Marks*

Nodular melanoma

Nodular melanoma, which accounts for 20% of melanomas, has no radial growth phase. It is typically found on the trunk and limbs of young to middle-aged individuals (Figure 117.8). It may have a ‘blueberry’-like nodule. Prognosis is determined by thickness at the time of excision.



Figure 117.8 Nodular melanoma on the back. It has no radial growth phase and because it grows vertically can be readily misdiagnosed. The ABCD rule often does not apply but this lesion shows variable colors and an irregular border. *Photo courtesy Robin Marks*

The early nodular melanoma problem[4, 5]

Nodular melanoma can present a diagnostic dilemma since the ABCD rule often does not apply. The mnemonic EFG, standing for 'elevated', 'firm' and 'growing for more than 1 month', is more appropriate. Early melanomas tend to be symmetrical, non-pigmented, even in colour, small diameter and grow vertically.

They are often mistaken for a haemangioma or a pyogenic granuloma. If one is suspicious, refer early to a specialist/specialist clinic.

Acral lentiginous melanoma

These typically occur on palms, soles and distal phalanges (Figure 117.9). They have a poorer prognosis than other types. They occur mainly in dark-skinned races.



Figure 117.9 Acral lentiginous melanoma. A 30-year-old man presented with a 'mole' on his toe that had become 'lumpier'. This type of melanoma, which occurs on the distal areas of the limbs, begins as a spreading pigmented, macule before developing into a nodule surrounded by a pigmented halo (as shown). *Photo courtesy Robin Marks*

Variations

Amelanotic melanomas are flesh-coloured papules that increase in size or change shape. These lesions can be extremely difficult to diagnose and the poor prognosis associated with them is due to late diagnosis rather than an increased malignancy.

The features and associations of melanoma subtypes are presented in Table 117.2.

Table 117.2 Features and associations of melanoma subtypes (after Kelly)[6]

Melanoma subtype	Frequency %	Radial growth phase	Location	Average age	Occupation profile
Superficial spreading	70	+	Trunk (back), limbs (legs)	Middle-aged	Indoor worker
Nodular	20	-	Trunk, limbs	Middle-aged	Indoor worker
Lentigo maligna	7.5	+	Head, neck	Elderly	Outdoor worker
Acral lentiginous	2.5	+	Palms, soles mucosae	Not known	Not known

Reproduced with permission of J. Kelly[4]

Prognosis

Determinants of prognosis include:[3]

- thickness (Breslow classification)
- level or depth (worse in level 4 or 5) (Figure 117.10)
- site (worse on head and neck, trunk)
- sex (worse for men)
- age (worse > 50 years)
- amelanotic melanoma
- ulceration



Figure 117.10 Assessment of tumour level: the levels of melanoma invasion. Produced by permission J. Kelly[4]

Vertical growth is associated with invasion and the prognosis worsens with depth. The chance of cure is greater than 90% if a melanoma is removed when it is less than 0.75 mm thick.3 If the lesion is allowed to invade to a thickness of 4 mm or more, the likelihood of a cure is reduced to less than 30%.[3]

The influence of tumour thickness on 5-year survival rates is shown in Table 117.3.

Table 117.3 The influence of tumour thickness on 5-year survival rates (after Kelly)[6]

Range of tumour thickness (mm)	Level (Clark)	5-year survival rates %
0	I	
< 0.76	II	95

0.76 - 1.5	III	70 - 98
1.51 - 4.0	IV	55 - 85
> 4.0	V	30 - 60

Reproduced with permission of J. Kelly[6]

Staging is based on the tumour level (depth) shown in Figure 117.10:

- level I - confined to the epidermis (in situ)
- level II - tumour cells extend into the superficial (papillary) dermis
- level III - tumour cells fill up the superficial dermis
- level IV - tumour cells extend into the deeper (reticular) layer
- level V - invasion of subcutaneous tissue

DIFFERENTIAL DIAGNOSIS

There are several common skin lesions that may be mistaken for melanoma.[1] They are:

- haemangioma (thrombosed)
- dermatofibroma (sclerosing haemangioma)
- pigmented seborrhoeic keratosis
- pigmented BCC
- junctional and compound naevi
- blue naevi
- dysplastic naevi
- lentigines

FACILITATING EARLY DIAGNOSIS OF MELANOMA

An adequate light source without shadows is essential. Refer to p. 1143 for use of the ‘maggylamp’ and the dermatoscope.

Clinical examination of the skin[1]

It is important to examine the entire skin and not just the lesion presented by the patient. Comparison of pigmented skin lesions is very helpful in differentiating between benign and malignant. One satisfactory routine is:

- Starting at the head, examine the hairline, backs of ears, neck, back, and backs of the arms. Pull down the underwear to expose the buttocks; examine the backs of the legs.
- With the patient facing you, examine the anterior hairline, the front of the ears, the forehead, cheeks and neck, moving downwards to the anterior chest. Move bra straps as required to achieve complete coverage. Then examine the abdomen, pulling down the underwear to examine as far as the pubic hairs.
- Then examine the anterior surfaces of the legs. The ‘maggylamp’ is very useful for this examination.

After scanning the entire skin surface and comparing and contrasting naevi, specific lesions may be examined with the dermatoscope. Compare suspicious lesions with similar lesions elsewhere on the patient's skin.

Applying the ABCDE system[1]

- **A** = **A**symmetry Melanoma is almost always asymmetrical. Most non-melanoma lesions are symmetrical, oval or round.
- **B** = **B**order The border of the melanoma is usually well defined, especially in the more malignant, compared

with the dysplastic naevus, which is almost always indistinct with a fading out ‘shoulder’ effect. The border of the melanoma is irregular while most benign lesions have a regular edge.

- **C** = **C**olour The classic blue-black colour is helpful but the *variety* of colours present in most melanomas is the most helpful. Magnification usually visualises greys, whites, violets, reds, oranges and shades of brown interspersed in the darker blue-black pigmentation. Early melanomas developing in dysplastic naevi tend not to have this deep pigmentation.
- **D** = **D**iameter The majority of melanomas when first seen are at least 7 mm in diameter, especially if arising from a pre-existing naevus. However, it is possible to diagnose small nodular melanomas < 5 mm.
- **E** = **E**levation Elevation indicates invasion and is a sign of more advanced disease.

DIAGNOSIS BY EXCLUSION

In the diagnostic process consider the lesions outlined in the differential diagnosis and check out the various characteristics. Haemangiomas may have an emptying sign when pressed with a finger. Pigmented BCCs can be difficult if they are fully pigmented but this is uncommon. The characteristic pearly-grey look and the telangiectasia are usually still visible on magnification with the ‘maggylamp’. The most useful feature of dysplastic naevi is that they are usually multiple and lesions for comparison can generally be found elsewhere. Dysplastic naevi also have greater breadth and height and often a darker nodule in the centre - the ‘target’ sign.

PITFALLS IN DIAGNOSIS OF MELANOMA

- Nodular melanoma
- Small melanoma
- Amelanotic melanoma
- Regressing melanoma
- Rapidly growing melanoma

MANAGEMENT POINTS FOR NAEVI AND MELANOMAS

- Do not inject local anaesthetic directly into the lesion.
- Incisional biopsy of a melanoma or suspicious mole is best avoided.
- Accurate clinical diagnosis, with the definitive treatment performed in one stage, is optimal, rather than excision biopsy with follow-up surgery.

MANAGEMENT TIPS[1]

- The solitary dysplastic naevus has no significant malignant potential.
- Multiple excision of naevi is not justified.
- If in doubt, perform excision biopsy with a margin of 2 - 3 mm. Refer to Figure 117.11 for the protocol for a suspicious lesion.
- If melanoma is strongly suspected, referral to a consultant is necessary.
- Beware of the pigmented BCC - it is easily missed but it usually has a shiny surface.
- Do not freeze a pigmented lesion - take a biopsy.


Suspicious pigmented lesion
local excision biopsy 2–3 mm margin to mid fat layer
histopathology
benign naevus
dysplastic naevus
dermatofibroma
pigmented BCC
(no further surgery)
malignant melanoma
melanoma in situ (lentigo maligna)
re-excision 5 mm margin
level 1–2 < 1.5 mm depth
re-excision 1 cm margin
level 3–5 > 1.5 mm depth
re-excision 2 cm margin

Figure 117.11 Plan for management of a suspicious pigmented lesion

GUIDELINES FOR EXCISION MARGINS[5, 7]

- Suspicious lesion - margin 2 - 3 mm
- Melanoma *in situ* - margin 5 cm
- Melanoma < 1.5 mm thick - margin 1 cm
- Melanoma 1.5 - 4 mm thick, minimum margin 1 cm and maximum 2 cm
- Melanoma > 4 mm thick, minimum margin 2 cm, maximum 3 cm

COUNSELLING

An encouragingly positive and supportive approach can realistically be taken for most patients, as the overall survival for melanoma in Australia is more than 80%.[1]

Even with tumours greater than 4 mm thickness, 50% of patients will survive.

FOLLOW-UP

Follow-up tends to be based on the tumour thickness:[1]

- 1 mm - 6 monthly review for 2 years
- 1 - 2 mm - 4 monthly for 2 years, 6 monthly for next 2 years, then yearly for 10 years
- > 2 mm - review by both specialist and GP, regularly, for 10 years

The first sign of metastasis is usually to the lungs, so a yearly chest X-ray is advisable.

REFERENCES

1. McCarthy W. The management of melanoma. Aust Fam Physician, 1993; 22: 1177 - 86.
2. Lauritz B. Pigmented skin lesions. In: Monash University Medical School's Annual Update Course for GPs. Course Proceedings. Melbourne: Monash University, 1992: 33 - 4.
3. Marks R. Skin cancer. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 469 - 72.
4. Kelly J. Nodular melanoma - no longer as simple as ABC. Aust Fam Physician, 2003, 32(9): 702 - 9.

5. Marley J (Chair). *Therapeutic Guidelines* (Version 2) Melbourne: Therapeutic Guidelines Ltd, 2004: 347 - 50.
6. Kelly J. Malignant melanomas - how many have you missed? Med J Aust, 1996; 164: 431 - 6.
7. Australian Cancer Network. *Clinical Practice Guidelines. The Management of Cutaneous Melanoma*. NH&MRC, 1999. < http://www.health. gov/nhmrc/publications >.

CHAPTER 118. HAIR DISORDERS

> Look it's only death. It's not like losing your hair.
>
> Harold Brodkey upon learning he had AIDS, New Yorker Magazine 1994

HAIR DISORDERS

The hair of our scalp is referred to as our crowning glory and the threat of hair loss in both sexes provokes extraordinary anxiety bordering on grief in some people. It behoves us as medical practitioners to treat the patient presenting with 'I'm losing my hair' with appropriate support and understanding. Likewise, the problem of hirsutism provokes similar anxiety and concerns about body image.

THE NORMAL HAIR GROWTH CYCLE

An understanding of the process of normal hair growth is necessary to comprehend and evaluate hair disorders. Each follicle progresses quite independently through regular cycles of growth and shedding (see Figure 118.1).


telogen club hair
dermal papilla
epithelial cells
new anagen hair
Anagen
Catagen
Telogen
telogen club hair
dermal papilla
epithelial cells
new anagen hair
Anagen
Catagen
Telogen

Figure 118.1 The normal hair cycle

The three phases of follicular activity are:[1]

1. Anagen phase - the active growth phase of hair production.
    - The dermal papilla stimulates division of epithelial cells that produce the hair shaft.
    - The hair shaft grows 1 cm per month.
    - It lasts for about 3 - 5 years on the scalp (average 1000 days).

  - It lasts 1 - 2 months on eyebrows and eyelashes.
  - It varies between individuals.

2. Catagen phase - a short transition phase from active growth to inactivity (the involutional stage).
    - The base of the hair becomes club-shaped.
    - It lasts for only about 2 weeks.
3. Telogen phase - the resting (dormant) phase of the cycle at the end of which the club hair with its non-pigmented bulb is shed.
    - This lasts 2 - 4 months (average 100 days).
    - The percentage of hairs in telogen varies from site to site - 10% in scalp to 60 - 80% in pubic hair.[2]
    - The hair is anchored in the follicle but does not grow longer.
    - The follicle then re-enters anagen.

Thus every 3 - 5 years every hair on the scalp is shed and replaced.

Some facts on hair numbers[1]

- Hair growth is asynchronous (i.e. continuous production and shedding).
- Humans produce 1 km of hair a month.
- About 50 - 100 hairs are shed daily without a reduction in density.
- The scalp contains, on average, 100 000 hair follicles.
- The hair follicle is subject to melanocytic activity.
- At least 25% of hair must be shed before a noticeable loss of density occurs.
- Hair loss counts consistently above 100 per day indicate excessive hair loss.
- Significant hair loss tends to block the shower drain or be visible all over the pillow.

KEY FACTS AND CHECKPOINTS

- Alopecia is a generic term for hair loss.
- Hair loss (alopecia) generates considerable anxiety and the fear of total hair loss should be addressed with the patient and a realistic prognosis given.
- Androgenic alopecia is the most common cause of human hair loss, affecting 50% of men by age 40 and up to 50% of women by age 60.[2]
- In telogen effluvium, the traumatic event has preceded the hair loss by about 2 months (peak loss at 4 months).
- Although severe stress could precipitate alopecia areata, day-to-day stressors are not considered to be a trigger. Stress seems to be a consequence of alopecia rather than the cause of it.[3]
- Hair loss can be patchy or diffuse where it involves the entire scalp.
- Patchy loss - alopecia areata and trichotillomania.
- Generalised loss - telogen effluvium, systemic disease, drugs (see Table 118.1).

Table 118.1 Causes of diffuse hair loss

• Androgenetic alopecia
• Telogen effluvium
• Postpartum telogen effluvium
• Alopecia areata (diffuse type)
• Drugs - cytotoxics and others
• Hypothyroidism

- Nutritional
  - iron deficiency
  - severe dieting
  - zinc deficiency
  - malnutrition
- Post-febrile state
- Anagen effluvium

- Alopecia areata has a poor prognosis if it begins in childhood, if there are several patches and there is loss of eyebrows or eyelashes.
- Scarring alopecia can be an indicator of lupus erythematosus or lichen planus.

ALOPECIA AREATA, ALOPECIA TOTALIS AND ALOPECIA UNIVERSALIS

Alopecia areata is a disorder of the hair follicle causing a sudden onset of localised or diffuse hair loss. It is thought to be an autoimmune disorder which has a genetic susceptibility (20% have a positive family history). The hairs are affected during the growth phase, resulting in cessation of anagen.

DxT: patch of complete hair loss + clean scalp + exclamation-mark hairs = alopecia areata

Clinical features

- Complete hair loss (small patch or diffuse)
- Pigmented hairs often lost first
- Clean normal scalp
- No or minimal inflammation
- Exclamation-mark hairs, especially around the periphery (Figure 118.2)



Figure 118.2 Exclamation-mark hair (a feature of alopecia areata)

Associations

- Loss of facial (beard in men) or body hair
- Nail changes - dystrophy or pitting

Alopecia areata[3]

The course taken may be as follows:

Localised patches

Small patches may recover spontaneously (usually 80%), while others depend on the extent of loss and general factors.

Treatment (options) includes:

- topical potent class III steroids (especially in children) bd, 12 weeks
- topical irritants (e.g. dithranol 0.5% ointment initially, increasing in strength to 2% applied once daily)
- intralesional steroids - triamcinolone 10 mg/mL or betamethasone acetate/phosphate 5.7 mg/mL bd (caution needed on face - risk of cutaneous atrophy)
- topical minoxidil 5% - 1 mL bd (for 4 or more months) only when hair growing
- topical immunotherapy with DNCB antigen

Extensive area (> 50% loss)

- Treatment
- Counselling
- Alopecia areata support group
- Use of cosmetic aids (e.g. wigs, hair pieces, scarves) and camouflage
- Topical steroids not recommended - ineffective
- Topical immunotherapy under specialist care may help
- Psoralen and ultraviolet A (PUVA) phototherapy may also help
- Systemic steroids only for active and progressive cases
  Prednisolone 0.75 mg/kg (o) daily for 3 - 4 weeks, then taper off over 2 months

A solitary patch of hair loss will recover within 6 months in 33% of cases, while 50% of cases will recover over a year and 33% will never recover. Unfortunately, most patients (86% in one study) relapse.[3]

Alopecia totalis

- This is where the alopecia extends over the total scalp.
- There is, at best, a 50% chance of hair recovery in a fit adult but only a very small chance in childhood.

Alopecia universalis

- This is where the alopecia involves the eyebrows and eyelids as well. Recovery is rare.

SCARRING ALOPECIA[1]

In this condition, the hair follicles are damaged and if the follicular openings cannot be seen with a magnifying lens, regrowth of hair cannot be expected. Thus, the process is irreversible and a scalp biopsy is essential to determine the diagnosis. Apart from obvious causes such as trauma, severe burns, a carbuncle and scalp ringworm with kerion, the causes of scarring alopecia are as follows:

- lichen planopilaris - this is a variant of lichen planus that produces small follicular papules in the scalp that tend to heal with scarring and destruction of hair follicles; treatment, which is difficult, includes corticosteroids, antimalarials and etretinate[1]

- discoid lupus erythematosus - this gives a similar picture to the former and tends to be treated in a similar manner
- folliculitis decalvans - a chronic folliculitis of the scalp, probably as a response to staphylococci on the scalp; treated with longterm tetracyclines
- pseudopelade - a slowly progressive, noninflammatory scarring condition causing patchy areas of hair loss without any obvious preceding skin disease

TELOGEN EFFLUVIUM

Telogen effluvium, which is increased shedding of hairs in the telogen phase, is one of the most common causes of diffuse hair loss and can be triggered by a variety of stressors.

It is worth noting that follicular matrix cells have a high metabolic rate second only to haematological tissue and stress can result in shunting into premature telogen with cessation of anagen.[1]

An obligatory delay occurs between the ‘insult’ or precipitating event and the onset of hair shedding because the hair follicle cycles through catagen and telogen - approximately 2 - 4 months when the club hairs with white bulbs of telogen are shed.

Greater than 25% of hair must be lost before there is a perceptible thinning and in this disorder up to 50% loss is common.

DxT: stressful event + 2 - 3 months gap to diffuse hair loss + ‘white bulbs’ = telogen effluvium

Patients usually complain of large clumps of hairs with white bulbs coming out with gentle tugging on combing or shampooing (this can exceed 150 hairs per day compared with the normal average of 50 - 100 hairs).

‘Stress’ precipitants of telogen effluvium

- Any severe stress
- Childbirth (common)
- High fever
- Weight loss, especially crash dieting
- Trauma - surgical or accidental
- Oral contraceptive pill (OCP) cessation
- Malnutrition
- Haemorrhage

Course of telogen effluvium

If uncomplicated, spontaneous recovery can be expected in 6 months so reassurance with explanation is all that is required. If it persists longer than 6 months, consider the chronic idiopathic form or an unmasked androgenetic alopecia. However, if there is concern about non-recovery and the stress factors are corrected, topical minoxidil for a minimum of 4 months is an option.[3] Referral to a specialist is advised for relapsing episodes or incomplete recovery.

Chronic telogen effluvium[4]

This occurs usually in perimenopausal and postmenopausal women. It may be primary and idiopathic or secondary to hypothyroidism, hyperthyroidism, malnutrition or cancer. The feature is episodes of dramatic hair shedding that recover but recur weeks to months later and last up to several days. It does not result in obvious balding - it is self-limiting and does not usually need treatment.

ANAGEN EFFLUVIUM

This is hair loss during the anagen phase and is typically seen in association with cancer chemotherapy and radiotherapy to the scalp, which results in immediate metabolic arrest. Anagen hair shafts are identified by their long and pigmented hair bulb. The follicle may remain in anagen, leading to a quick recovery, or move into telogen, thus delaying growth by about 3 months.[3]

DRUG-INDUCED ALOPECIA

Drugs are a very important cause of alopecia (Table 118.2). They may cause telogen effluvium, anagen effluvium or accelerate androgenetic alopecia.

Table 118.2 Drug-induced causes of alopecia generally from prolonged use[2,3]

• Cytotoxic agents/thallium
• Amphetamines
• Anticoagulants: heparin, warfarin
• Antiepileptics: phenytoin, sodium valproate, carbimazole
• Antigout agents: allopurinol, colchicine
• Antihelminthic agents: albendazole/mebendazole
• Anti-inflammatories: indomethacin, gold, penicillamine, salicylate
• Anti-parkinson: levodopa, bromocriptine
• Antithyroid agents/thyroxine/iodine
• Cardiovascular agents:
- Amiodarone
- Statins, clofibrate
- Selected ACE inhibitors
- Selected β-blockers
• Cimetidine
• Gentamicin
• Hormones: OCP, androgens, danazol
• Interferon
• Lithium
• Methysergide
• Vitamin A derivatives/retinoids: roaccutane

Drugs tend to cause telogen effluvium but cancer chemotherapy, radiation to the scalp, thallium/mercury/arsenic and colchicine in high dosage cause anagen effluvium. Acceleration of androgenetic alopecia is caused by hormone therapy, namely the oral contraceptive pill, danozol, testosterone and anabolic steroids.

ANDROGENETIC ALOPECIA (MALE PATTERN BALDNESS)

This is the most common form of alopecia which is age related and is genetically determined in addition to being androgen dependent. The key androgen is dihydrotestosterone which is produced from testosterone by 5α reductase. The hair loss is reasonably predictable, with some men losing hair quickly and others more slowly in a familial pattern. In others it is unpredictable. The diffuse pattern loss seen in women tends to progress slowly. About 50% of women have significant hair loss by the age of 60. The typical male pattern is shown in Figure 118.3.



Figure 118.3 Typical male pattern baldness

Alopecia affects 30% of men by age 30 and 50% by age 50.

Androgenetic alopecia in women

- In women, the pattern of hair loss is different to men.
- Diffuse thinning occurs, usually on the top of the head (the crown). The front hairline usually remains but in some women this can recede with bitemporal loss (Figure 118.4).



Figure 118.4 Typical female pattern baldness

- Although hair loss can appear in men and women as early as the 20s, it may not appear before the age of 50 in women.
- Some women notice a short period of considerable hair loss but this may be followed by a long stable period of no loss. Total loss of hair rarely occurs in women.
- It may be unmasked after an episode of diffuse hair loss such as after childbirth or an acute illness.

Treatment for men

It is appropriate to counsel men to accept their balding as part of a natural ageing process. This includes cutting the hair short to make it look better cosmetically. If baldness is not acceptable, some options include wearing a toupee, a wig or other hair substitute or having a hair transplant operation. However, with hair transplantation the new hair is often just as likely to disappear as the original hair. The use of medicated oils and shampoos should be discouraged.

Available medications[3]

- Minoxidil 2% and 5%, 1 mL applied bd to the dry scalp for a minimum of 12 months. The results vary from good to no change. One study showed that one-third of men using 2% minoxidil for 12 months experienced ‘cosmetically significant hair growth’.

  However, a lifetime of expensive treatment is required in these people and hair loss resumes, with loss of that being maintained by the treatment, when it is ceased.
- Finasteride 1 mg tablet taken daily for a minimum of 2 years. This can halt the balding process in over 85% of men

and may initiate some modest regrowth in 65% over a 12-month period, with further improvement if it is continued over a longer period. However, it is expensive, is not a cure and balding resumes when it is discontinued.

Treatment for women

Physical treatments/hair styling

This includes the use of wigs, hair transplantation and camouflage. Wigs can be worn on the whole head or on the bald spot, or fibres can be interwoven into the remaining hairs.

Camouflage can be used either by having the existing hair bleached by a skilled hairdresser or by colouring the scalp the same colour as the hair. Mascara can be lightly brushed into the roots of the hair at receding hairlines or along gaps.

Medications available[2]

- Minoxidil 2% and 5% topical application bd for minimum of 12 months (to assess efficacy), or
- Spironolactone 200 mg daily for 6 months, or
- Cyproterone acetate 100 mg (o) daily (course of 10 tablets a month) usually with OCP
- If postmenopausal, cyproterone acetate 50 mg (o) daily in consultation with an endocrinologist

As for men, these drugs tend to prevent further loss and, if effective, need long-term use.

TRICHOTILLOMANIA (HAIR PULLING)

This is patchy hair loss caused by deliberate plucking or twisting of hair shaft s. It is reasonably common in young children, where it may be of little significance, simply being a ‘habit’. In older children and adults it may be an obsessive compulsive disorder often associated with stress.[1]

Clinical features

- Incomplete patchy alopecia
- Hairs of different length
- Hairs broken and twisted
- Strange pattern of loss
- Tends to occur on side of dominant hand
- Eyelashes or eyebrows may be involved

HAIR DISORDERS IN CHILDREN

Loose anagen (growing hair) syndrome[5]

This is a disorder of the hair follicle characterised by the ability to pluck anagen hairs painlessly from the scalp by gentle pulling. It presents as very thin, wispy new hair growth. It is an autosomal dominant trait.

Clinical features

DxT: fair females under 5 + thin wispy hair + easy loss with pulling = loose anagen syndrome

- Thin wispy hair with tatty ends
- More common in girls; can affect boys
- Onset in early childhood - usually < 5 years

- Large clumps easily pulled out at play
- Light microscopy of hair shafts aids diagnosis

Outcome

- Spontaneous improvement with age
- Usually normal hair by teens

Treatment

- Reassurance and explanation
- Gentle hair care

Traction alopecia

Traction alopecia is thinning of the hair seen in female children and young women due to very tight hairstyles, as with ponytails, excessive hair rolling and hair braiding in particular. The bald areas that show short broken hairs and sometimes scarring are found in areas of maximal tug. The most common type is ‘marginal alopecia’ found in the forehead and sides at the hair edges. Patients should be advised to cease the procedure. Rollers that cause traction can be replaced by those that heat.

Trichotillomania[5]

This occurs in children typically between 4 and 10 years, usually as a nocturnal habit and parents may be unaware of the hair pulling. The affected areas are usually on the anterior and temporal areas. The areas are never completely bald. A characteristic feature is an irregular-shaped area of incomplete patchy alopecia containing hairs of different length. The variable length is due to the fact that some hairs will not break with pulling while others will break at varying distances from the scalp surface. There may be associated follicular pustules. However, scrapings should be taken to exclude a particular type of tinea capitis (black dot ringworm) caused by *Trichophyton tonsurans*. The management is similar to thumb sucking or nail biting with a low-key approach. It does not imply a significant psychological problem.

Localised alopecia areata

This can also occur in children (refer pp. 1231 - 2). Most childhood cases resolve spontaneously but it can progress to total hair loss or recurrent alopecia can occur. Regrowth decades later is a possibility. Treatment with moderate-potency topical steroids for 12 weeks or so may help.

Tinea capitis

Dermatophyte infection that produces an area of incomplete, ‘unclean’ alopecia with various degrees of scaling and inflammation of the scalp surface. A boggy swelling (kerion) can develop in severe cases. Wood's light examination will be positive in only 50% of cases. Confirm diagnosis with scalp scrapings for microscopy and culture.

OFFICE PROCEDURES

The following procedures can be useful in determining diagnosis and prognosis.

The hair-pull test[1]

In this simple method, 50 - 100 hairs are grasped between thumb and forefinger and gently pulled proximal to distal. This is repeated 6 - 8 times and should yield a total of 2 - 5 telogen hairs, which can be analysed. More than eight hairs per pull is abnormal.

Trichogram

Twenty to 50 hairs are extracted by a short, sharp pull with artery forceps. The anagen-to-telogen ratio is calculated. The ratio is decreased in telogen effluvium because of the increased numbers of hairs in telogen.

Scalp biopsy

This differentiates between scarring and nonscarring alopecia and also between alopecia areata and trichotillomania.

Light microscopy of hair shafts

Light and/or electron microscopy is indicated if a hair shaft defect is suspected. Skin scrapings and hair samples are taken for fungal microscopy and culture.

HIRSUTISM

Hirsuties is growth in the female of excess, coarse, terminal pigmented hair in androgen-dependent sites, namely in a male sexual pattern (e.g. upper lip, beard area and back; Figure 118.5).



Figure 118.5 Areas prone to hirsutism

Most cases are due to idiopathic hirsutism, which may follow racial or familial factors, or to a variant of the polycystic ovarian syndrome. Other causes include adrenal hyperplasia, virulising adrenal tumours, Cushing's disease, virulising ovarian tumours and iatrogenic causes, particularly drugs (Table 118.3).

Table 118.3 Causes of virilisation[2]

• Prolactinoma
• Adrenal
- congenital hyperplasia
- Cushing's disease
• Gonal dysgenesis

- androgen therapy

- obesity

• Ovarian

- polycystic ovaries (PCOS)

- ovarian tumours

Routine investigation is not required unless there are other features of virulisation.

It is important to exclude adrenal or ovarian pathology.

Hirsutism of reasonably sudden onset is an important marker for serious underlying pathology.

Appropriate investigations include pelvic ultrasound to diagnose polycystic ovaries and serum testosterone and dehydroepiandrostenedione (DHEA), which indicates an adrenal cause of androgen excess if elevated.

For patients with severe hirsutism and regular menses, serum testosterone, DHEA and early morning 17-hydroxyprogesterone levels should be measured. If menses is irregular, measure follicle stimulating hormone, luteinising hormone and serum prolactin.

Management principles

- Exclude underlying adrenal or ovarian pathology.
- Support the patient to achieve an acceptable body image through appropriate advice and referral.
- Recommend appropriate cosmetic measures, which include bleaching (a good option), waxing or depilatory creams, or shaving.
- Do not pluck hairs, especially around the lips and chin-plucking stimulates hair growth but shaving appears to have no effect.
- Electrolysis may help for hair removal but it is expensive and requires multiple treatments from an experienced therapist (beware of scarring). It achieves temporary hair removal - significant regrowth can occur within 3 months.
- Laser epilation may also help but seems to be most suitable for dark hair on a light skin and provides temporary removal, lasting up to 6 months.

Pharmaceutical options

These include either inhibition of androgen action at the hair follicles (spironolactone and cyproterone acetate) or suppression of ovarian and/or adrenal androgen production (dexamethasone or oral contraception).

The following are options:[2]

- spironolactone 100 - 200 mg (o) daily (takes 6 - 12 months to respond), or
- cyproterone acetate 10 - 100 mg (o) daily for 10 days of each menstrual cycle plus an OCP

Eflornithine HCl (Vaniqa) cream for topical use (especially for facial hair) is a new preparation.

HYPERTRICHOSIS

Hypertrichosis is the increased growth of fine vellus or downy hair over the entire body. It is non-androgendependent hair and does not respond to anti- androgen therapy. The cause is generally unknown - it may be primary or constitutional where it is usually apparent prior to puberty and where it is evenly distributed over the back and limbs. Prepubertal hypertrichosis is commonly familial but a positive family history is not always determined.

An important cause of secondary hypertrichosis is drugs. Drugs that are implemented include phenytoin, minoxidil, cyclosporin, corticosteroids, including topicals, diazoxide and anabolic steroids.

It is also associated with an underlying neoplasia, such as carcinoma of the stomach, when it is termed hypertrichosis lanuginosa acquisita or 'malignant down'.

Other causes include anorexia nervosa, the menopause, starvation and a few rare syndromes (e.g. Cornelia de Lange syndrome).

SCALY SCALP DISORDERS

Dandruff

Dandruff (pityriasis capitis) is mainly a physiological process, the result of normal desquamation of scale from the scalp. It is most prevalent in adolescence and worse around the age of 20.

If it is persistent with heavy scaling, seborrhoeic dermatitis and scalp psoriasis, which is distinguished by palpable plaques, are likely causes.

Dandruff and seborrhoeic dermatitis are related conditions on a continuum of severity. They both have a common pathogenesis and are chronic recurring conditions presenting with scaling and a varying degree of itch. More importantly, they all respond to similar treatments.

Treatment of dandruff

Shampoos:

- zinc pyrithione (e.g. Dan-Gard, Head and Shoulders) or
- selenium sulphide (e.g. Selsun)

Method: massage into scalp, leave for 5 minutes, rinse thoroughly - twice weekly.

Treatment of persistent or severe dandruff

- Coal tar plus salicylic acid compound (Sebitar) shampoo, or
- Lonil T plus shampoo

Method: as above, followed by Sebi Rinse or ketoconazole (Nizoral) shampoo. If persistent, especially itching, and Nizoral shampoo ineffective, use a corticosteroid (e.g. betamethasone scalp lotion).

Dry hair

Management tips

- Don't shampoo every day.
- Use a mild shampoo (labelled for 'dry or damaged hair').
- Use a conditioner.
- Snip off the split or frayed ends.
- Avoid heat (e.g. electric curlers, hair dryers).
- Wear head protection in hot wind.
- Wear a rubber cap when swimming.

Oily hair

Management tips

- Shampoo daily with a 'shampoo for oily hair'.
- Massage the scalp during the shampoo process.
- Leave the shampoo on for at least 5 minutes.

- Avoid hair conditioners.
- Avoid overbrushing.
- Attend to lifestyle factors: relaxation and balanced diet are important.

REFERENCES

1. Cargnello J. I think I'm losing my hair. Aust Fam Physician, 1997; 26: 683 - 7.
2. Cargnello S, Sheil R. How to treat: Hair loss and hirsutism. Australian Doctor, 2003; 21 November: 31 - 8.
3. Marley J (Chair). *Therapeutic Guidelines: Dermatology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2004: 189 - 99.
4. Whiting DA. Chronic telogen effluvium: increased scalp hair shedding in middle aged women. J Am Acad Dermatol, 1996; 35: 889 - 906.
5. Smart JS ed. *Paediatric Handbook* (6th edn). Melbourne: Blackwell Science, 2000: 268 - 9.

CHAPTER 119. NAIL DISORDERS

A physician should wear white garments, put on a pair of shoes, carry a stick and an umbrella in his hand, and walk about with a mild and benignant look as a friend of all created beings. He should be cleanly in his habits and well shaved, and should not allow his nails to grow.

SUSHRUTA-SAMHITE (5TH CENTURY BC)

NAIL DISORDERS

Making a diagnosis of abnormal nails for a concerned patient can be quite simple for a few obvious conditions. However, in many cases the diagnosis can be elusive when we are not familiar with classic patterns that are seen so infrequently. The diagnostic process can be facilitated by learning the basic anatomy and function of the nail, as well as characteristic patterns, which are presented with the aid of diagrams in this chapter. There are, in fact, only a limited number of ways in which injury, infection and inflammation can present in a nail.[1]

The main nail problems encountered in general practice are trauma, onychomycosis, infection, ingrowing toenails, paronychia and psoriasis. Fungal nail infection and psoriasis are the commonest causes of nail dystrophy. Damage to the nail from trauma or disease results in nail dystrophy. The problem of nail changes due to onychotillomania, be it from excessive nail biting, picking or cleaning, should be suspected from the history and examination.

The examination should include a general inspection of the skin including in the webbing of the toes, looking for evidence of a skin disorder such as psoriasis, atopic eczema, lichen planus and tinea.

KEY FACTS AND CHECKPOINTS

- The growth rate of nails varies between individuals: fingernails average 0.5 - 1.2 mm per week while toenails grow more slowly.[2]
- An avulsed or totally dystrophic nail will take up to 9 months to regrow.
- It takes approximately 6 weeks to grow a new cuticle.
- Beau's lines associated with a severe acute illness take about 3 months to appear.
- Do not confuse chronic paronychia with onychomycosis. The former affects the nail folds, the latter mainly affects the distal nail.
- Nail clippings for culture and histology may be the only way to differentiate between nail dystrophy and onychomycosis.
- Not all white crumbly nails are caused by a fungus.
- Suspect melanoma in any subungual pigmented lesion. Do not confuse with subungual haematomas, which ‘grow out’ with the nail. They usually present as a longitudinal pigmented streak. Beware of amelanotic melanoma, which may mimic chronic paronychia or a pyogenic granuloma.2, 3 Any suspicion necessitates early referral.
- Various dermatoses and connective tissue disorders can affect the nails - psoriasis, lichen planus, lupus erythematosus, scleroderma, bullous pemphigoid, Darier's disease (keratosis follicularis).4
- Clubbing of the fingers is basically an abnormality of the fingertips - look for evidence of major pulmonary or cardiac disease.
- Significantly bitten or traumatised nails may be a symptom of a major anxiety disorder - explore psychogenic issues.

ANATOMY AND FUNCTION OF THE NAIL

The basic nail unit (apparatus) consists of the nail matrix, the proximal and lateral nail folds, the cuticle and the nail plate (Figure 119.1). The hard keratin of the nail plate is formed in the nail matrix. The matrix, which contains germinal epithelium, lies in an invagination of the epidermis (the nail fold) and is protected by a waterproof seal formed by the cuticle. The matrix runs from the proximal end of the nail fold to the cuticle.


(a)
proximal nail fold
cuticle
nail plate
nail bed
matrix
distal phalanx
(b)
lateral nail fold
nail plate
lunula
cuticle
proximal nail fold

Figure 119.1 Normal anatomy of the nail. Diagram of the nail apparatus: (a) lateral section view; (b) dorsal surface view

The nail bed closely approximates the distal phalanx. The nail has an important functional and cosmetic role. It enhances fine touch and motor skills for picking up fine objects, such as a pin from a flat surface, not possible without fingernails.

NAIL DISORDERS AND THEIR CAUSES[1]

Lifting of the nail plate (onycholysis)


psoriasis
trauma
fungal

Causes

- Trauma
- Factitious (self-induced)
- Tinea
- Psoriasis
- Photosensitivity, usually tetracyclines
- Others (e.g. warts, lichen planus)
- Hyperthyroidism

Thickening of the nail plate

Causes

- Trauma
- Tinea
- Psoriasis
- Onchogryphosis



Thinning of the nail plate

Causes

- Trauma - wear and tear (repeated water immersion)
- Artificial fingernails
- Lichen planus
- Peripheral vascular disease
- 20-nail dystrophy, usually children

Pitting of the nail plate

Causes

- Psoriasis
- Alopecia areata
- Atopic dermatitis



Longitudinal marks in the nail plate

Grooved

Causes

- Myxoid cyst
- Angiofibroma



Central single stripe

- Darier's disease
- Hereditary/congenital
- Mechanical trauma



Horizontal grooves in nail plate

Causes

- Beau's lines (from acute illness)
- Habit tic (picking cuticle of thumb nail)
- Atopic dermatitis



Horizontal single white lines or band

Causes

- Chemotherapy
- Arsenic poisoning
- Kidney failure

Lamellar splitting

- Trauma - repeated wetting and drying (e.g. housework)



Abnormal curvature

Spoon nails (koilonychia)

Causes

- Hereditary
- Trauma
- Iron deficiency
- Haemochromatosis



Over curvature

- Hereditary pincer nail

Clubbing

Causes

- Hereditary/congenital
- Lung disease (e.g. cancer), pulmonary fibrosis
- Heart disease (e.g. congenital cyanotic, SBE)
- Liver disease (e.g. cirrhosis)
- Gastrointestinal disorders (e.g. Crohn's disease, refer p. 480)



Splinter haemorrhages

Causes

- Minor trauma (e.g. manual workers)
- Bacterial endocarditis



Discolouration of the nail plate

- White
  - striate leuconychia (usually trauma)
  - liver cirrhosis
  - hypoalbuminaemia (total whiteout)
- Red
  - splinter haemorrhages
  - kidney failure
- Brown
  - fungal infection
  - cigarette staining
  - drugs - gold, chlorpromazine

    - Addison's disease
    - psoriasis (‘oil stain’ patches)
    - melanoma
- Black
    - haematoma
    - melanoma
    - racial pigmentation
    - naevus
    - minocycline
    - cytotoxic drugs (transverse bands)
- Green
    - *Pseudomonas*
    - *Aspergillus*
- Blue
    - antimalarials
    - argyria
- Yellow
    - yellow nail dystrophy (? respiratory illness)
    - fungal infections
    - psoriasis
    - slow growth
    - tetracyclines
- Half (pale proximal) and half (brown or red distal)
    - kidney failure (chronic)
    - liver cirrhosis
    - trauma (biting, splits, splinters)

Swelling of the proximal nail folds (paronychia)

- Manicure (retracting the cuticles)
- *Candida* infection
- Staphylococcal infection
- Herpetic whitlow

Swelling of the lateral nail folds

- Ingrowing toenails
- Retinoids
- Over-curvature of the nail plate

Tumours of the nail fold

Benign

- Myxoid (mucus) cyst
- Warts
- Periungual fibroma

Malignant

- Squamous cell carcinoma
- Melanoma

Destruction of the nail apparatus

- Lichen planus
- Melanoma
- Bowen's disorder
- Squamous cell carcinoma
- Trauma

ONYCHOLYSIS[1]

Onycholysis refers to the separation of the nail plate from the underlying nail bed and is a sign rather than a disease. This separation creates a subungual space with an air interface that gathers unwanted debris, such as dirt and keratin. It is usually seen in fingernails but it can develop in toenails from rubbing against shoes. Adverse local reactions to agents such as formaldehyde and resins in polishes or nail glues can distort nails.

Self-induced trauma is a common cause from obsessive manipulation, including meticulous cleaning and frequent manicuring.

The band of discolouration at the end of the separated nail is usually in a straight line compared with other causes such as psoriasis and tinea. Tinea may be distinguished from other causes by white or yellow streaks or ‘spears’ travelling proximally in the nail.

Greenish discolouration indicates invasion by *Pseudomonas pyocyanea* or *Aspergillus.*

A summary of causes of abnormal nails is presented in Table 119.1.

Table 119.1 Abnormal nails: diagnostic strategy model (modified)

Q. Probability diagnosis
A. Fungal infection: onychomycosis
Trauma to nail bed
Trauma from biting
Trauma from habit picking
Onychogryphosis
Paronychia
Psoriasis
Q. Serious disorders not to be missed
A. Melanoma
Iron deficiency: koilonychia
Liver disease: leuconychia
Endocarditis: splinter haemorrhages
Chronic kidney failure: white bands, half and half nail
Glomus tumour
Bowen's disorder
Q. Pitfalls (often missed)
A. Atopic dermatitis
Lichen planus

Pyogenic granuloma (usually with ingrowing toenails)[3]

Drug effects (e.g. tetracycline)

Pseudomonas infection

Connective tissue disorders (e.g. SLE)

Arsenic (mee's stripes)

Management of onycholysis[5]

First exclude psoriasis, tinea (check toe webbing) and trauma (check history).

Fingernails

- Keep nails as short as possible
- Avoid insertion of sharp objects under nails for cleaning
- Apply tape (micropore or similar) over free edge for months, until healed
- Avoid unnecessary soaps and detergents - wear gloves for housework, gardening etc.
- Keep hands out of water
- Use a mild soap and shampoo

Toenails

- Exclude fungal infection (clinical tinea pedis) - culture
- Improve footwear to avoid any rubbing

Pharmaceutical treatment

- Daily application of an imidiazole (e.g. clotrimazole) or terbinafine
- For *Pseudomonas* infection soak the nails in vinegar or Milton's solution and/or gentamicin sulphate cream

PSORIASIS OF NAILS

Psoriasis can have many manifestations, such as pitting, onycholysis, discolouration, splinter haemorrhages, distal subungual hyperkeratosis (which can resemble warts) and severe total nail dystrophy (often with arthropathy). Psoriasis can closely mimic onychomycosis, which should be excluded by fungal culture and histology before commencing presumed tinea therapy.

There is no effective topical therapy for psoriasis of the nails. Intralesional steroid injections, which are painful and require multiple treatments, can help. Successful treatment of the skin does not help the nails.

ONYCHOMYCOSIS (FUNGAL NAIL INFECTION)

Facts

- Affects 3 - 5% of population and 40% over 60 years.1
- Classified as superficial, distal or proximal.
- Toenails affected more commonly than fingernails.
- The most common form is distal lateral subungual caused by *Trichophyton mentagrophytes* var. *intertigitale* (typical of toe web space tinea and responds well to terbinafine) or by *Trichophyton rubrum* (common on sole of foot and more resistant).
- Superficial white onychomycosis is also common, and is usually confined to the toenails with small superficial white plaques with distinct edges and caused by *Trichophyton mentagrophytes* var. *intertigitale*.

- Total dystrophic onychomycosis - whole nail affected, thickened, opaque and yellow brown (caused by *Trichophyton* sp.).
- *Candida albicans* and other moulds are not a common factor.
- Diagnosis is by culture and histology of the distal nail plate clippings placed in formalin. Positive in 60 - 80% of cases.

Treatment [5]

The treatment of choice for all types of toenail tinea is:

- terbinafine 250 mg (o) daily for 12 weeks (cures 70 - 80% of cases): 6 weeks for fingernails

Tips

No improvement apart from the proximal part of the nail will be noticed after months because it takes 12 months or more for the toenail to grow.

Mark the base of the dystrophic nail with a scalpel blade or black ink to assess progress.

Topical treatment

- Amorolfine 5% (loceryl) nail lacquer 1 - 2 times weekly after filing (fingernails: 6 months; toenails: 9 months)

A systematic review found poor evidence for the effectiveness of topical therapy for onychomycosis.[6]

PARONYCHIA

Acute paronychia

This is a painful condition that is mainly due to bacterial infection, especially *Staphylococcus aureus.*

Management

Uncomplicated with localised pus:

- simple elevation of nail fold or puncture the fold - close to drain pus
- advice on hygiene
- antibiotics rarely necessary
- exclude diabetes

Complicated with subungual extension:

- small vertical incision alongside the nail, or
- removal of nail in part or totally
- exclude diabetes

Chronic paronychia

Features

- Painless
- A form of traumatic nail dystrophy
- Loss of cuticle fundamental to diagnosis

Causes

- Excessive manipulation of cuticles (e.g. manicurists)
- Occupational (e.g. chefs, housewives, nurses, fishmongers)

- Frequent contact with water, detergents and chemicals
- Habit tic - picking the nail fold

Note:

- Secondary infection with *Candida* common but not basic cause.
- The damaged cuticle permits access of water and grit to the nail matrix and causes inflammation of the proximal nail fold.

Management [4, 5]

- Culture organisms
- Exclude diabetes
- Basic nail care advice:
  - keep hands dry (avoid wet work if possible)
  - wear cotton-lined gloves when washing dishes (for max. of 15 minutes)
  - minimise contact with water, soap, detergents, lipid solvents and other irritants
  - never pick, push back or manicure cuticles
  - never insert anything beneath cuticle for cleaning
  - wear cotton gloves in garden
  - use a mild soap and shampoo

Medications [5]

For Candida (if cultured):

- tincture miconazole bd, or
- clotrimazole topical preparations

For *Staphylococcus* (if cultured):

- garamycin topical ointment

Topical medications to nail folds (especially if persistent exudate):

- 4% thymol in alcohol (SVR) qid, or
- 10% sulphacetamide in alcohol

Vaseline and/or steroid ointment can be applied frequently when it is dry and without exudate.

ATOPIC DERMATITIS

- Avoid irritants: special soaps, wear gloves for dishwashing
- Good nail hygiene
- Apply potent topical steroids to proximal nail fold

LICHEN PLANUS

This commonly presents in the nail with atrophy of one or more nails. The nail plate is predisposed to breaking and splitting as it thins. If advanced, a pterygium arising from the proximal nail fold grows as the nail matrix is destroyed and eventually total loss of the nail may occur.

Biopsy of the nail matrix is recommended before treatment.

Intralesional steroids are the treatment of choice if seen prior to destruction of the nail unit.

TWENTY-NAIL DYSTROPHY

This is an uncommon disorder seen in preschool or preadolescent children with thinning and roughening of all or almost all 20 nails. It may be a prodrome to psoriasis, lichen planus or alopecia areata but it tends to be a self-limiting condition and most cases resolve over 2 - 3 years.[4]

It is reported to respond well to the superpotent topical steroid, clobetazal.

BRITTLE NAILS

The nails break easily usually at their distal end. Brittle nails are age related and are usually caused by local physical factors, such as repeated water immersion, and exposure to chemicals, such as detergents, alkalis and nail polish removers and also hypothyroidism and digital ischaemia. Systemic causes such as deficiency of iron and vitamins are not considered to be a common factor.

Calcium does not contribute to the hardness of nails and calcium deficiency does not cause brittle nails.[4] No cosmetic applications appear to be helpful.

Management [4]

- Avoid excessive hydration and trauma
- Wear rubber gloves with cotton liners for wet work
- Massage Vaseline or nail creams (e.g. Eulactol or Neostrat) into the nail several times daily
- Nail polishes and hardeners (preferably without formalin) may give a good cosmetic result

NAIL APPARATUS MELANOMA

Features [5]

- Rare but potentially fatal
- Responsible for 2 - 3% of all melanoma
- All age groups but especially in seventh decade
- Affects all races in all climates
- Presents as a longitudinal pigmented streak in the nail (Figure 119.2)


longitudinal pigmented streak
pigmentation of proximal nail fold

Figure 119.2 Longitudinal pigmented streak sign of nail apparatus melanoma

- Hutchinson's sign (pigmentation of the proximal nail fold) may be present
- Usually diagnosed late
- Mortality > 50%
- Median survival < 12 months after diagnosis
- Early recognition may result in a cure

Management

- All cases require a longitudinal nail biopsy for diagnosis
- If confirmed, treatment is based on Breslow thickness and level of invasion
- Level 1 or in situ - removal of whole nail apparatus
- Invasive melanoma - amputation of distal phalanx

GLOMUS TUMOURS

Glomus tumours are those mysterious little tumours that occur beneath the nail plate and can give paroxysms of sharp pain.

They are small purplish lesions that are exquisitely tender to touch and activate with changes in temperature. They may not be visible but may be revealed when the nail bed is blanched with gentle pressure. Imaging may be needed for diagnosis. Treatment is surgical.

ONYCHOGRYPHOSIS

Onychogryphosis, or irregular thickening and overgrowth of the nail, is commonly seen in the big toenails of the elderly and appears to be related to pressure from footwear (Figure 119.3). It is really a permanent condition. Simple removal of the nail by avulsion is followed by recurrence some months later. Softening and burring of the nail gives only temporary relief, although burring sometimes provides a good result.



Figure 119.3 Onychogryphosis

The powder from burring can be used as culture for fungal organisms. Permanent cure requires ablation of the nail bed after removal of nail.

Two methods of nail ablation are:[8]

1. total surgical excision
2. cauterisation with phenol (with care)

INGROWING TOENAILS

Refer to section in Chapter 68.

PERIUNGUAL WARTS

These are similar to other warts and fortunately rarely spread under the nail to the nail bed. Caution is needed with treatment.

Cryotherapy must be used with caution to avoid damage to the nail matrix - otherwise it is usually safe but often ineffective. Consider wart paints under occlusion.

SUBUNGUAL HAEMATOMA [7]

The small, localised haematoma

There are several methods of decompressing a small localised haematoma under the fingernail or toenail that causes considerable pain. The objective is to release the blood by drilling a hole in the overlying nail with a hot wire or a drill/needle.

Method 1: The sterile needle

Simply drill a hole by twisting a standard disposable hypodermic needle (21- or 23-guage) into the selected site. Some practitioners prefer drilling two holes to facilitate the release of blood.

Method 2: The hot paper clip

Take a standard, large paper clip and straighten it. Heat one end (until it is red hot) in the flame of a spirit lamp. Immediately transfer the hot wire to the nail, and press the point lightly on the nail at the centre of the haematoma. After a small puffof smoke, an acrid odour and a spurt of blood, the patient will experience immediate relief (Figure 119.4).



Figure 119.4 Treatment of subungual haematoma. The point of a heated end of a paper clip is pressed lightly on the nail at the centre of the haematoma

Method 3: Electrocautery

This is the best method. Simply apply the hot wire of the electrocautery unit to the selected site (Figure 119.5). It is very important to keep the wire hot at all times and to be prepared to withdraw it quickly, as soon as the nail is pierced. It should be painless.



Figure 119.5 Electrocautery to subungual haematoma

Important precautions

- Reassure patients that the process will not cause pain; they may be alarmed by the preparations.
- The hot point must quickly penetrate, so go no deeper than the nail. The blood under the nail insulates the underlying tissues from the heat and, therefore, from pain.
- The procedure is effective for a recent traumatic haematoma under tension. Do not attempt this procedure on an old, dried haematoma as it will be painful and ineffective.
- Advise the patient to clean the nail with spirit or an antiseptic and cover with an adhesive strip to prevent contamination and infection.
- Advise the patient that the nail will eventually separate and a normal nail will appear in 6 - 9 months.

The large haematoma

Where blood occupies the total nail area, a relatively large laceration is present in the nail bed. To permit a good, long-term functional and cosmetic result it is imperative to remove the nail and repair the laceration (Figure 119.6).



Figure 119.6 Treating the diffuse haematoma. **(a)** The diffuse haematoma. **(b)** Sutures to laceration. **(c)** Fingernail as splint

Method

- Apply digital nerve block to the digit.
- Remove the nail.
- Repair the laceration with 4/0 plain catgut.
- Replace the fingernail, which acts as a splint, and hold this in place with a suture for 10 days.

MYXOID PSEUDOCYST

There are two types of digital myxoid pseudocysts (also known as mucous cysts) appearing in relation to the distal phalanx and nail in either fingers or toes (more common) (Figure 119.7). One type occurs in relation to, and often connecting with, the distal interphalangeal joint and the other occurring at the site of the proximal nail fold. The latter (more common) is translucent and fluctuant, and contains thick clear gelatinous fluid, which is easily expressed after puncture of the cyst with a sterile needle. Osteoarthritis of the DIP is associated with leakage of myxoid fluid into the surrounding tissue to form the cyst.



Figure 119.7 Myxoid pseudocyst: typical position of the cyst

Some pseudocysts resolve spontaneously. If persistent and symptomatic attempt:[7]

- repeated aspiration (aseptically) at 4 - 6 weekly intervals *or*
- cryosurgery *or*
- puncture, compression, then infiltration intralesionally with triamcinolone acetonide (or similar steroid)

They tend to persist and recur and, if so, refer to surgery for total excision of the proximal nail fold and/or ligation of the communicating stalk to the PIP.

REFERENCES

1. Sinclair R. There is something wrong with my nail. Aust Fam Physician, 1997; 26: 673 - 81.
2. Hunter JA, Savin JA, Dahl MV. *Clinical Dermatology.* Oxford: Blackwell Scientific, 1989: 70.
3. Reid C. Nail disorders. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 334 - 6.
4. Howard A. Common nail disorders - part 1. Aust Doctor, 1999; 30 July: i - iv.
5. Sinclair R. Treating common nail problems. Aust Fam Physician, 1997; 26: 949 - 2.
6. Crawford F, Hart D, et al. Topical treatments for fungal infections of the skin and nails of the foot. Cochrane Database Syst Rev, 2000(2); CD 001434.
7. Murtagh J. *Practice Tips.* Sydney: McGraw-Hill (4th edn), 2004: 90 - 1.
8. Marley J (Chair). *Therapeutic Guidelines*: *Dermatology* (Version 3). Melbourne: Therapeutic Guidelines Ltd, 2004: 277.

CHAPTER 120. ALCOHOL PROBLEMS

If you want to keep a dead man, put him in whisky; if you want to kill a live man put whisky in him.

THOMAS GUTHRIE (1803 - 73)

ALCOHOL PROBLEMS

Excessive drinking of alcohol is one of the most common and socially destructive problems in Australia. One survey found that 5% of Australian men and 1% of women were alcohol-dependent. It also showed that 86% of men and 79% of women drink alcohol, with 8.3% of the population drinking alcohol every day.[1]

Skinner refers to the prevalence of ‘alcoholism’ in North America as about 5 - 7% of the adult group, with a much larger group (20 - 30%) of individuals having drinking problems without major symptoms of alcoholism.[2] Abstainers represent 10 - 20% of populations.

EXCESSIVE AND HARMFUL DRINKING

Intoxication, regular excessive drinking and alcohol dependence are three dimensions of alcohol-related problems that should be assessed by the medical practitioner.

For men, excessive drinking is more than four standard drinks of alcohol a day. For women, drinking becomes a serious problem at lesser amounts - more than two standard drinks (SDs) a day. This level can also affect the fetus of the pregnant woman.

RISK OF HARM IN THE SHORT TERM (BASED ON STANDARD DRINKS)

The guidelines for risk of harm in the short term and in the long term are summarised in Tables 120.1 and 120.2.[3]

Table 120.1 Risk of harm in the short term

	Low risk (standard drinks)	At risk (standard drinks)	High risk (standard drinks)
Men (on any 1 day)	Up to 6 on any 1 day, no more than 3 days/week	7 - 10 on any 1 day	11 or more on any 1 day
Women (on any 1 day)	Up to 4 on any 1 day, no more than 3 days/week	5 - 6 on any 1 day	7 or more on any 1 day

Table 120.2 Risk of harm in the long term

	Low risk (standard drinks)	At risk (standard drinks)	High risk (standard drinks)
Men (on average per day)	Up to 4	5 - 6	7 or more
Overall weekly level	Up to 28 per week	29 - 42 per week	43 or more per week
Women (on average per day)	Up to 2	3 - 4	5 or more
Overall weekly level	Up to 14 per week	15 - 28 per week	29 or more per week

EXTENT OF THE PROBLEM

- Alcohol is estimated to have a harmful effect on about 1 in 10 people.
- At least 20 - 40% of acute general and psychiatric hospital admissions have an alcohol-related illness.

- About 20% plus of fatal traffic accidents involve alcohol.
- The author's study[4] identified alcohol dependence in 9.7% of the population studied and a further group of problem drinkers that included the 'explosive' or binge drinker (Figure 120.1). Problem drinkers represent about 15 - 20% of the population.



Figure 120.1 Prevalence of alcohol drinking patterns in the adult population (figures expressed as a percentage)

IDENTIFYING THE PROBLEM

As a profession we generally seem to be very slow in recognising the problem drinker; we need to train ourselves to have a sixth sense in early detection of the heavy or problem drinker.

Clinical pointers

Alcohol abuse should be suspected in any patient presenting with one or more of the physical or psychosocial problems presented in Table 120.3. Target areas for clinical scrutiny are outlined in Table 120.4. The facial features of the patient can be a helpful pointer, albeit of the more advanced drinker. These include:

- plethoric facies
- puffy 'greasy' facies
- telangiectasia
- rosacea + rhinophyma
- suffused ('bloodshot') conjunctivae
- prominent lower lip with chelitis of corners of mouth
- smell of stale alcohol or very 'minty' sweet breath (masking effect)

Table 120.3 Adverse clinical effects of alcohol

Psychological and social effects	Physical effects
• Concern about drinking by self, family or others	• Brain damage (if severe)
• Heavy drinking - more than six glasses per day	• Depression
• Early morning drinking	• Epilepsy

• Reaching for the bottle when stressed	• Wernicke-Korsakoff syndrome
• Regular hotel patron	• Insomnia - nightmares
• Skipping meals/poor diet	• Hypertension
• Cancelling appointments	• Heart disease
• Increased tolerance to alcohol	- arrhythmias
• Frequent drinking during working day	- cardiomyopathy
• Loss of self-esteem	- beri-beri heart disease
• Irritability	• Liver disease
• Devious behaviour	• Pancreatic disease
• Anxiety and phobias	• Dyspepsia (indigestion)
• Depression	• Acute gastritis
• Paranoia	• Stomach ulcers
• Stress	• Sexual dysfunction
• Relationship breakdown	• Hand tremor
• Marital problems	• Peripheral neuropathy
• Child abuse	• Myopathy
• Poor work performance	• Gout
• Absenteeism from work/loss of job	• Obesity
• Memory disturbances	• Other metabolic/endocrine effects
• Financial problems	- hyperlipidaemia
• Alcohol related accidents	- pseudo-Cushing's syndrome
• Driving offences	- osteoporosis
• Crime - violence and other offences	- osteomalacia
• Personal neglect	• Haemopoiesis
• Attempted suicide	- macrocytosis
• Pathological jealousy	- leucopenia
• Heavy smoking	- thrombocytopenia
• Behavioural problems in children	

Table 120.4 Target areas for clinical scrutiny

Young and middle-aged single males
Divorced or separated individuals
Alcoholic beverage trade: bar trade, hotel staff
Professionals: politicians, doctors and others
Travelling professions (e.g. sailors, salespeople, truck drivers)
Armed forces, especially returned servicepeople
Authors, journalists and related workers
Social club patrons (e.g. sporting) clubs

TAKING A DRINKING HISTORY

This requires tact and skill and it must be noted that many problem drinkers considerably understate the level of their intake.

Useful strategies

- Ask questions as part of a matter-of-fact enquiry into health risk factors, such as smoking and diet.
- Place the onus of denial on the patient by asking questions such as 'When did you last drink alcohol?' rather than 'Do you ever drink alcohol?'
- Record your patient's intake quantitatively in terms of standard drinks or grams of alcohol.
- Confirm the history by enquiring about the time spent drinking per day and expenditure on alcohol.

Useful questions

- When did you last drink alcohol?
- Do you like alcohol?
- What is your usual intake each day? Each week?
- What type of alcohol do you prefer to drink?
- Do you take a drink in the morning?
- Do you eat breakfast?
- When was the last time you felt nauseated or 'off-colour' in the morning?
- When do you get heartburn?
- Do you drink with your mates or family or at the club?
- How long do you usually go without alcohol?
- When was the last time you were drunk?
- When was the last time you cannot remember a drinking session?
- About how much alcohol can you take before it affects you?
- Has alcohol had any effects on you?
- Does it give you the shakes?
- Do you ever need to take alcohol to help you get to sleep?
- Do you need it to steady your nerves?

Questionnaires

There are several questionnaires that can be most helpful, assuming the patient is fully cooperative. Two or more positive replies for the CAGE questionnaire[5] are suggestive of a problem drinker.

1. Have you ever felt you should CUT down on your drinking?
2. Have people ANNOYED you by criticising your drinking?
3. Have you ever felt bad or GUILTY about your drinking?
4. Have you ever had a drink first thing in the morning to steady your nerves or get rid of a hangover? (an EYE-OPENER)

Another very practical questionnaire is AUDIT (The Alcohol Use Disorders Identification Test) developed by the WHO to facilitate the early detection of problem drinkers in primary care.

LABORATORY INVESTIGATIONS

The following blood tests may be helpful in the identification of excessive chronic alcohol intake:

- blood alcohol
- serum GGT: elevated in chronic drinkers (returns to normal with cessation of intake)

- MCV: > 96 fL

Other changes:

- abnormal liver function tests (other than GGT)
- carbohydrate deficient transferrin (quite specific - dependent on an enzyme induced by alcohol)
- HDLs elevated
- LDLs lowered
- serum uric acid elevated

MEASURING ALCOHOL INTAKE

One standard drink contains 10 g of alcohol, which is the amount in one middy (or pot) of standard beer (285 mL), two middies of low-alcohol beer or five middies of super-light beer. These are equal in alcohol content to one small glass of table wine (120 mL), one glass of sherry or port (60 mL) or one nip of spirits (30 mL) (Figure 120.2).



Figure 120.2 Standard drinks

- 1 stubby or can of beer = 1.3 standard drinks
- 1 standard light beer = 0.9 standard drinks
- 1 × 750 mL bottle of beer = 2.6 standard drinks
- 1 × 750 mL bottle of wine = 6.8 standard drinks

The 0.05 level

To keep below the 0.05 blood alcohol level drinking and driving limit, a 70 kg man should not exceed:

- 2 standard drinks in 1 hour
- 3 standard drinks in 2 hours
- 4 standard drinks in 3 hours

The rule is that one standard drink is eliminated per hour so it is important to spread drinking time.

ALCOHOL DEPENDENCE

Alcohol dependence is a syndrome in which an individual demonstrates clinically significant impairment or distress as manifested by three or more of the following, occurring at any time in the same 12 month period:

1. tolerance
2. withdrawal
3. drinking in larger amounts or for a longer period than intended
4. unsuccessful attempts to cut down or control drinking
5. a great deal of time spent in activities necessary to obtain, use or recover from the effects of alcohol
6. important social, occupational or recreational activities reduced or given up because of drinking
7. continued drinking despite knowledge of having persistent or recurrent problems caused by or exacerbated by drinking

Identification of the alcohol-affected person is complicated by the tendency of some to hide, underestimate or understate the extent of their intake.

In order to diagnose and classify alcohol-dependent people, the family doctor has to rely on a combination of the above features and parameters that includes clinical symptoms and signs, available data on quantity consumed, clinical intuition, personal knowledge of the social habits of patients, and information (usually unsolicited) from relatives, friends or other health workers.

A checklist of pointers to the adverse effects of chronic alcohol abuse is presented in Table 120.3. In a study by the author the outstanding clinical problems are the psychogenic disorders (anxiety, depression and insomnia) and hypertension.[6] Susceptibility to work and domestic accidents were also significant findings.

The challenge to the family doctor is early recognition of the alcohol problem. This is achieved by developing a special interest in the problem and a knowledge of the early clinical and social pointers, and being ever alert to the tell-tale signs of alcohol dependence.

APPROACH TO MANAGEMENT

The challenge to the family doctor is early recognition of the problem. There are specific target areas which should be considered carefully by the GP (Table 120.4). Several studies have shown that early intervention and brief counselling by the doctor are effective in leading to rehabilitation.[7] Some of the results are very revealing:

- Patients expect their family doctor to advise on safe drinking levels.[8]
- They will listen and act on our advice.[9]
- Treatment is more effective if offered before dependence or chronic disease has developed.[9, 10]

Of prime concern to the GP is the assessment of whether the patient is interested in changing his or her excessive drinking behaviour. The proposed model of change by Prochaska and Di Clemente helps identify the stage reached by the patient (Figure 120.3).[11]



Figure 120.3 Prochaska's and Di Clemente's[11] proposed model of change to facilitate the identification of behavioural stages and the provision of counselling for treating dependence on alcohol, tobacco and other drugs

Precontemplators are satisfied users who are either unconcerned about their drinking or have no desire to change.

However, if there is any evidence of ambivalence or concern about drinking then the opportunity exists for motivational interviewing techniques.

Patients tend to have little insight into their problem and often need the development of unpleasant sequelae to make them aware of their alcohol-related problem. Furthermore, patients are not likely to offer concern about their drinking problem spontaneously but are often receptive to the initiative coming from their doctor.

The family doctor is ideally placed to identify and treat the problem of alcohol because the individual who abuses alcohol will tend to surface at some point in the provision of primary health care.

Of particular concern are teenage and early adult drinking patterns, often influenced by environmental factors, including the home and sporting clubs. Fortunately, many young people are able to control their drinking as they mature, provided they survive the risk-taking behaviour period. However, those who remain single tend to adopt drinking as part of their lifestyle. When the duration of excessive drinking increases to 10 or 15 years, patients tend to present with classic alcohol-related diseases.[2] According to Skinner, alcohol dependency usually develops in individuals in their early twenties, yet most patients admitted to alcohol treatment programs have a history of heavy drinking of 10 - 20 years with associated alcohol-related morbidity.

ALCOHOL-SENSITISING DRUGS

Drugs that cause a most unpleasant reaction when taken with alcohol include the adversive agents disulfiram and calcium carbimide. Their role is controversial and they should be restricted to a cooperative and highly motivated patient who gives informed consent and when someone is available to supervise compliance.

The reaction causes nausea, vomiting, flushing, dizziness and dyspnoea.

Disulfiram is used as a 200 - 400 mg (o) daily dosage. However, there is poor quality evidence generally to support its use predominantly due to poor compliance and it has not shown any effect on abstinence.[11]

“ANTI-CRAVING” DRUGS

The two agents acamprosate (Campral) and naltrexone (Revia) have been shown to reduce cravings and improve the rate of continuous (total) abstinence, cumulative number of days abstinent or the time to first-drinking relapse, relative to placebo during the treatment periods.[12, 13] However, studies vary in out-comes but the general conclusion is that the drugs show a modest effect on the frequency of drinking and the likelihood of maintaining abstinence. Both agents need to be taken for 6 months or more (usually 9 months) and used in combination with counselling or attendance at Alcoholics Anonymous (AA) meetings.

Dosage:
- acamprosate 666 mg (o) tds (less if < 60 kg)[15]
- naltrexone 50 mg (o) once daily

Current evidence indicates that the combination of naltrexone and acamprosate is more effective in preventing relapse than the individual drugs.[15]

MANAGEMENT

The best results are obtained with hearly recognition of the problem and prompt intervention to resolve it; the patient's admission of the problem and a resolve to face it; firm support and interest by the medical team and appropriate support from family and friends. The patient should set their own goals and develop a contract with their GP.

A brief practical management plan[16]

Giving patients feedback about their level of alcohol consumption, presenting objective evidence of harm and setting realistic goals for reducing alcohol intake induces many to change their drinking behaviour.

A six-step management plan, which has been employed in a general-practice early intervention program, is as follows:

1. *Feed back* the results of your assessment and specifically the degree of risk associated with their daily alcohol intake and bout drinking. Emphasise any damage that has already occurred.
2. *Listen* carefully to their reaction. They will need to ventilate their feelings and may respond defensively.
3. *Outline the benefits* of reducing drinking:
    - save money
    - less hassles from family
    - sleep better
    - have more energy
    - be less depressed
    - lose weight
    - better physical shape
    - lessen the risk of:
        - hypertension
        - liver disease
        - brain disease
        - cancer
        - accidents
4. *Set goals* for alcohol consumption which you both agree are feasible. In most cases this will involve reduction to below certain ‘safe limits’.
    - For men: *no more than* 3 - 4 drinks 3 - 4 times per week (aim for less than 12 per week).
    - For women: *no more than* 2 - 3 drinks 2 - 3 times per week (aim for less than 8 per week). Pregnant women should not have more than 1 drink 2 - 3 times per week. Some authorities recommend total abstinence.
      These are the *upper low-risk limits* and are *not* amounts that patients should be recommended to drink if their intake is normally lower. For patients who have already experienced some physical damage or substantial psychosocial problems, it is best to advise a period of total abstinence. For patients who are physically dependent on alcohol, long-term abstinence is advisable.
5. *Set strategies* to keep below the upper safe limits:
    - Quench thirst with non-alcoholic drinks before having an alcoholic one.
    - Have the first alcoholic drink *after* starting to eat (avoid drinking on an empty stomach).
    - Switch to low-alcohol beer.
    - Take care which parties you go to: avoid constant parties and other high-risk situations.
    - Think of a good explanation for cutting down on your drinking.
    - Have a physical workout when bored or stressed.
    - Explore new interests - fishing, cinema, social club, sporting activity.
6. *Evaluate* progress by having patients monitor their drinking by using a diary; check that any abnormal blood test results are returning to normal. Make a definite appointment for follow-up and give appropriate literature such as *Alcohol and Health*. Obtain consent for a telephone follow-up. A useful minimum intervention plan is presented in Table 120.5.

Table 120.5 Minimum intervention technique plan (5 - 10 minutes)

1 Advise reduction to safe levels

2 Outline the benefits

3 Provide a self-help pamphlet

4 Organise a diary or other feedback system

5 Obtain consent for a telephone follow-up

6 Offer additional help (e.g. referral to an alcohol and drug unit or to a support group)

Follow-up (long consultation 1 week later)

Review the patient's drinking diary. Explore any problems, summarise, listen and provide support and encouragement. If appointment is not kept, contact the patient.

Specialist services

According to progress and the patient's wishes and consent, specialist treatment units, group therapy and attendance at meetings of Alanon or AA are potential sources of help to keep the alcohol-dependent person abstinent and coping.

WITHDRAWAL SYMPTOMS

Symptoms of a ‘hangover’ include headache, nausea, irritability, malaise and a mild tremor. Withdrawal from alcohol in a chronic problem drinker includes:

- agitation
- prominent tremor
- sweating
- insomnia
- seizures
- delirium tremens (DTs)

Treatment for moderate symptoms is based on diazepam. The aim is to prevent development of DTs. Maintain fluid, electrolytes and nutrition. Add vitamin B complex, including thiamine, because the patients are invariably thiamine deficient.

Recommended treatment for acute withdrawal symptoms[15]

- Diazepam 10 - 20 mg (o) every 2 hours (up to 120 mg (o) daily) titrated against clinical response (taper off after 2 days) in the hospitalised or well-supervised patient
- Thiamine 100 mg IM or IV daily for 3 - 5 days, then 100 mg (o) daily
- Vitamin B group supplement (o) or IM daily

Delirium tremens

DTs is a serious life-threatening withdrawal state. It has a high mortality rate if inadequately treated and hospitalisation is always necessary.

Clinical features:

- may be precipitated by intercurrent infection or trauma
- 1 - 5 days after withdrawal (usually 3 - 4 days)
- disorientation, agitation
- clouding of consciousness

- marked tremor
- visual hallucinations (e.g. spiders, pink elephants)
- sweating, tachycardia, pyrexia
- signs of dehydration

Treatment[14]

- Hospitalisation
- Correct fluid and electrolyte imbalance with IV therapy
- Treat any systemic infection
- Thiamine (vitamin B_1 100 mg IM or IV daily for 3 - 5 days, then thiamine 100 mg (o) daily
- diazepam 5 mg by slow IV injection (over several minutes) every half hour until symptoms subside *or* diazepam 10 - 20 mg (o) every 2 hours (up to max. 120 mg daily) until symptoms subside

This dose is usually required for 2 - 3 days, then should be gradually reduced till finished. If psychotic features (e.g. hallucinations and delusions) add haloperidol 2.5 - 5 mg (o) bd

Note: Chlormethiazole is not recommended because of its potential to produce excessive CNS depression and dependency.[14]

ALCOHOL OVERDOSE[15]

Overdose is potentially fatal. The average lethal blood alcohol concentration is about 0.45 - 0.5%. Death from a lower concentration may occur with other sedative drugs. Alcohol withdrawal may begin at 0.1%. The rate of decline is usually 0.015% to 0.02% per hour. Treatment of overdose is supportive and symptomatic. No stimulants should be given. Overdose may cause hypoglycaemia and metabolic acidosis.

HANGOVER

A type of acute drug toxicity causing headache, nausea and fatigue.

Preventive advice:

- Drink alcohol on a full stomach.
- Select alcoholic drinks that suit you: avoid champagne.
- Avoid fast drinking - keep it slow.
- Restrict the quantity of alcohol.
- Dilute your drinks.
- Avoid or restrict smoking while drinking.
- Drink three large glasses of water before retiring.

Treatment

- Drink ample fluids especially water because of relative dehydration effect of alcohol.
- Take two paracetamol tablets for headache.
- Drink orange juice or tomato juice, with added sugar.
- A drink of honey in lemon juice helps.
- Coffee and tea (preferable) are suitable beverages.
- Have a substantial meal but avoid fatty food.

REFERENCES

1. Alcohol and other drug use in Australia. Australian Institute of Health and Welfare. Australian Government. Canberra, 2004: 23 - 25.
2. Skinner HA. Early detection and basic management of alcohol and drug problems. Aust Alcohol Drug Rev, 1985; 4: 243 - 9.
3. NH&MRC. *Australian Alcohol Guidelines: Health Risk and Benefits*. Canberra: NH&MRC, 2001.
4. Murtagh JE. Stigmata of alcohol dependence. Aust Fam Physician, 1985; 14: 204 - 5.
5. Mayfield D, McLeod G, Hall P. The CAGE questionnaire. Am J Psych, 1974; 131: 1121 - 3.
6. Murtagh JE. Alcohol abuse in an Australian community. Aust Fam Physician, 1987; 16: 20 - 5.
7. NH&MRC. Guidelines Preventive Interventions in Primary Health Care: Cardiovascular Disease and Cancer. No. 6. *Alcohol Overuse*. Canberra: NH&MRC, 1996.
8. Cockburn J, Killer D, Campbell E, Sanson-Fisher RW. Measuring general practitioners' attitudes towards medical care. J Fam Pract, 1987; 3: 192 - 9.
9. Wallace P, Cutler S, Haines A. Randomised controlled trial of general practitioner intervention in patients with excessive alcohol consumption. Br Med J, 1988; 297: 663 - 8.
10. Saunders JB. The WHO project on early detection and treatment of harmful alcohol consumption. Aust Drug Alcohol Rev, 1987; 6: 303 - 8.
11. Prochaska JO, Di Clemente CC. Towards a comprehensive model of change. In: Miller WRJ, Heath N, eds. *Treating Addictive Behavior*. New York: Plenum, 1986: 3 - 27.
12. Garbutt JC et al. Pharmacological treatment of alcohol dependence. A review of the evidence. JAMA, 1999; 281: 1318 - 25.
13. NSW Government Printer, Managing drug and alcohol problems in primary care. NPS, 2002; 22: 3 - 6.
14. Tedeschi M. Naltrexone and acamprosate. Aust Fam Physician, 2001; 30(5): 447 - 50.
15. Dowden J (Chair). *Therapeutic Guidelines: Psychotropic* (Version 5). Melbourne: Therapeutic Guidelines Ltd, 2003: 99 - 105.
16. Saunders JB, Roche AM. One in six patients in your practice. NSW medical education project on alcohol and other drugs. A drug offensive pamphlet. Sydney, 1989: 1 - 6.

CHAPTER 121. ALLERGIC DISORDERS INCLUDING HAY FEVER

> The nostril membrane is so irritable that light dust, contradiction, an absurd remark - anything - sets me sneezing and I can be heard in Taunton with a favourable wind, a distance of six miles. Turn your mind to this little curse. If consumption is too powerful for physicians, at least they should not suffer themselves to be outwitted by such little upstart disorders as the hay fever.
>
> SYDNEY SMITH 1835, LETTER TO DR HOLLAND

ALLERGIC DISORDERS INCLUDING HAY FEVER

Allergic disorders affect approximately 20% of the population. The most common allergies are those associated with IgE-mediated (immediate, or type 1) hypersensitivity, such as allergic rhinoconjunctivitis (hay fever), atopic dermatitis (eczema) and allergic asthma.[1]

Less commonly encountered but of increasing clinical importance in the community are IgE-mediated allergies to foods such as peanuts and/or other nuts and seafoods (crustaceans or molluscs), which may cause urticaria, angioedema, anaphylaxis and even death. Peanuts are one of the most common causes of food-induced anaphylaxis in adults. A clinically significant cross-reactivity between peanuts and other legumes is uncommon, but allergy to tree nuts such as almonds and walnuts may occur in up to 50% of those with peanut allergy.[2] Another special case is the oral allergy syndrome in which people with some degree of seasonal allergy to grass pollens or birch pollen suffer oral itch and swelling when they come into contact with certain fruits. This problem may be alleviated by desensitisation to pollens.[2]

Natural rubber latex allergy, associated with the introduction of universal precautions to decrease transmissible infections, is an increasingly important cause of Type 1 hypersensitivity, affecting particularly medical and paramedical personnel. Patients who have had multiple operations or procedures are another high-risk group. Diagnosis is suggested by history and confirmed by specific skin tests or the detection of serum specific IgE. The development of urticaria on contact with latex is highly suggestive of underlying type 1 hypersensitivity. An interesting association between latex allergy and sensitivity to fruit is recognised, most commonly banana, kiwifruit or avocado.[2]

ATOPY

Atopy refers to those 40% of people who have an inherited tendency for an exaggerated IgE antibody response to common environmental antigens.[1] There will be a positive response to one or more allergen skin-prick tests, and usually a family history of allergic disorders. Of those who are atopic, one-half to onethird manifest an allergic disorder, most commonly allergic rhinitis, asthma, atopic dermatitis or allergic gastroenteropathy.

COMMON ALLERGENS CAUSING IMMEDIATE HYPERSENSITIVITY

It is helpful to consider important allergen exposure during history taking. Table 121.1 lists sources of common allergens.

Table 121.1 Sources of common allergens[1]

Inhalants
Pollens, domestic animals, house dust mites, mould spores, cockroaches
Foods
Peanuts, fish, shellfish, milk, eggs, wheat
Other
Drugs, latex, insect venoms, occupational

Inhalant allergies

Allergic rhinoconjunctivitis and asthma are the main manifestations. The history provides a strong pointer to the causative allergen. If symptoms are seasonal, pollen allergy is most likely; perennial symptoms indicate an allergy to dust mites, household pets or moulds. Certain activities which precipitate symptoms may also provide a clue - these include mowing lawns, dusting and vacuuming.

Food allergy and intolerance[3]

Food allergy usually manifests in infancy and childhood with symptoms ranging from severe urticarialtype reactions to gastrointestinal symptoms such as anorexia, nausea, vomiting and spitting up of food, colic, diarrhoea and failure to thrive.

Foods that commonly cause allergic reactions include milk and other dairy products, eggs and peanuts. Other foods implicated include oranges, soya beans, nuts, chocolate, fish, shellfish and wheat. The allergic reactions are not to be confused with non-immunological food intolerance such as lactose intolerance.

A food intolerance is an adverse reaction to a specific food or food ingredient. It is regarded as a food allergy if the reaction is immune based. Food allergies can be simply classified as:

- immediate reactions - occurring within 2 hours
- delayed reactions - occur up to 24 hours after ingestion

IgE-mediated food reactions[3]

These are immediate reactions and relatively easy to diagnose.

Features

- Typically in infants and toddlers
- Due to release of mast cell mediators
- Produce flushing or blotchiness/pallor (if severe)
- Urticaria - facial or generalised (Figure 121.1)



Figure 121.1 Acute urticaria in a child caused by a sensitivity to aspirin

- Feeling of intense fear (angor animi)
- Angioedema

- Vomiting (immediate of soon after)
- ± naphylaxis with wheezing etc.
- Death can occur
- Big three foods - cow's milk, egg, peanuts
- Also: soy beans, fish, especially shellfish, various fruits and vegetables
- Cow's milk can cross-react with goat's milk and soy protein
- Frequently resolves by 3 - 5 years

Management

- Document diet, symptoms, past history, family history
- Refer for specialist allergy advice
- Advise avoidance of suspected food
- IgE reactions investigated with skin testing

Non-IgE-mediated food reactions[3]

These are less common and are usually delayed, occurring within 24 - 48 hours of food ingestion but some reactions may be immediate. The real explanation is not clear.

Features

- Gastrointestinal symptoms (e.g. vomiting, diarrhoea, abdominal colic)
- May be malabsorption, weight loss, failure to thrive (rare)
- Aggravation of atopic dermatitis
- Severe reactions possible
- Main foods - cow's milk, soy proteins
- Uncommon after 3 years of age

Management

- Elimination of suspected food, then formal food challenge
- Skin testing usually not helpful

Food intolerances[4]

In general practice it is common to see a variety of food intolerances that are not immune (allergic) or psychologically based. The food constituents represent an important group. The intolerances can be grouped as:

- fructose intolerance from excessive ingestion of fruit juices and soft drinks
- lactase deficiency from milk
- histamine-related reactions from strawberries, tomatoes
- chemical triggers
    - aspirin, tartrazine, sodium metabisulphite - triggering rhinitis and asthma
    - aspirin, tartrazine, benzoic acid - triggering chronic urticaria
- others - salicylates (manufactured or natural), amines, preservatives and colourings

Note:

- Foods with high-content natural salicylate - dried fruits, pineapple, apricots, oranges, cucumbers, grapes, honey, olives, tomato sauce, wines, tea, herbs.
- Foods likely to contain tartrazine (food colouring) - bottled sauces, cakes (from shops), coloured fizzy drinks, fruit cordial, custard, coloured sweets, ice-cream and lollies, jam.

Symptoms

There can be a variety of symptoms:

- irritability, behavioural problems
- gastrointestinal (e.g. infantile colic, diarrhoea, irritable bowel syndrome)
- respiratory - rhinitis, asthma
- headache/migraine

Management

- Document diet, reactions, past history, family history
- Elimination diet and controlled food challenge
- Referral for specialist allergy advice
- Radioallergosorbent (RAST) testing is indicated where skin testing contraindicated

Peanut allergy

Peanuts are one of the most common causes of food-induced anaphylaxis in adults. The diagnosis is confirmed by demonstration of peanut-specific IgE by either skin-prick tests with peanut extract or RAST testing. Reaction to peanuts usually begins within minutes of ingestion, the first symptoms being oropharyngeal itching or burning. Flushing, urticaria, wheeze, stridor, angioedema and collapse may follow.[5] The combination of peanut allergy and asthma is dangerous, as evidenced by fatal or near fatal reactions in young children.[6] The key to management is avoidance of peanut-containing foods. Desensitisation is currently not recommended. Those at risk should carry an anaphylaxis kit (Table 121.2).

Table 121.2 Adult anaphylaxis kit[3]

Epi Pen (0.3 mg adrenaline 1:1000 IM injection)
Inject into outer thigh muscle at first sign of swelling of throat or tongue, or other reaction (e.g. breathlessness)
Medihaler Epi MDI (adrenaline metered aerosol spray)
Spray 10 - 20 times in milder reactions only (e.g. local lip tingling or swelling)
Oral antihistamines
e.g. 10 mg loratadine tablets (× 2) Take 1 tablet after adrenaline injection
Prednisolone 25 mg tablets (× 2)
Take immediately after adrenaline

Latex allergy

The clinical manifestations of type 1 hypersensitivity reactions to latex protein are wide-ranging, from urticaria to life-threatening anaphylaxis and death. It is believed that some episodes of intraoperative anaphylaxis are due to the patient - who has been sensitised to latex - reacting after mucosal contact with gloves worn by operating staff. Some institutions now provide latex-free operating suites in response to this serious problem.[7]

Latex allergy is thus a significant problem for at-risk people, including health care workers, patients with spina bifida or other spinal cord abnormalities, and those who have had multiple operations. The greatest risk is contact with “dipped” rubber products (e.g. gloves, condoms, balloons). Some hard rubber products may not pose a risk.

Symptoms

Contact dermatitis (type 4 hypersensitivity), urticaria, worsening atopic dermatitis, allergic rhinoconjunctivitis, asthma, allergy to multiple fruits and possibly anaphylaxis.

Diagnosis

Skin-prick tests (dangerous and best left to experts) are more sensitive than blood tests at this stage but carry a risk of anaphylaxis. Measurement of serum specific IgE is safe although less sensitive. Contact allergy (type 4) is identified by patch testing.

Management

Health care workers who are allergic can never again wear latex gloves.[8]

TESTS FOR SPECIFIC IGE

Skin-prick tests

This is the preferred method as results can be read at the first consultation, provided the high-quality allergen preparations are used. A positive test alone may be of no diagnostic significance if the patient is asymptomatic to the specific allergens. A negative test is very useful for excluding IgE-mediated allergy.

Detection of serum specific IgE

A number of tests including RAST tests[1] and ELISA tests measure allergen-specific IgE in the serum. They are no more accurate than skin testing, are expensive and do not provide an immediate result.

Indications include: history and skin tests not matching, extensive eczema, dermographism, infants and very young children, immunotherapy work-up, antihistamine use in past 48 hours.

MANAGEMENT PRINCIPLES[1]

Allergen avoidance

If relevant from history and skin-prick testing, special attention should be paid to reducing exposure to house dust mites and mould, to pet selection and specific food avoidance. Change of occupation and environment may be necessary for some people.

Pharmacotherapy

Drugs are used to alleviate symptoms where avoidance methods have failed or are impractical. Examples include antihistamines (H_1-receptor and H_2-receptor antagonists), adrenaline (emergency use), sodium cromoglycate, corticosteroids, some anticholinergics and sympathomimetics.

Immunotherapy (desensitisation)

This involves repeated administration of small increasing doses of allergen by subcutaneous injection. This is the treatment of choice for severe wasp or bee venom allergy and for resistant allergic rhinoconjunctivitis where a single causative allergen can be identified. Patients should be observed for at least 45 minutes and adequate resuscitation facilities are essential.

Management of specific allergic disorders

- Asthma - see Chapter 123 (p. 1275)
- Atopic dermatitis - see Chapter 113 (p. 1158)
- Urticaria - see Chapter 112 (p. 1156)
- Anaphylaxis and angioedema - see Chapter 130 (pp. 1342 - 3)

RHINITIS

Refer to p. 634.

The classification of rhinitis can be summarised as:

- seasonal allergic rhinoconjunctivitis = hay fever
- perennial rhinitis
  - allergic (usually due to house dust mites)
  - non-allergic = vasomotor
    - eosinophilic
    - non-eosinophilic

ALLERGIC RHINITIS

Allergic rhinitis may be seasonal or perennial. It can be classified as either intermittent (lasting for < 4 days of the week or < 4 weeks) or persistent (lasting > 4 days of the week or > 4 weeks). The severity of symptoms is classified as either mild or moderate/severe.[9] Its prevalence varies from 5 - 20% with a peak prevalence in children and young adults up to 20%.[10] The symptoms are caused by release of powerful chemical mediators such as histamine, serotonin, prostaglandins and leukotrienes from sensitised mast cells.[10]

Seasonal allergic rhinoconjunctivitis (hay fever)

This is the most common type of allergic rhinitis and is due to a specific allergic reaction of the nasal mucosa, principally to pollens. The allergens responsible for perennial allergic rhinitis include inhaled dust, dust mite, animal dander and fungal spores.

Most cases of hay fever begin in childhood with one-half having the problem by the age of 15 and 90% of eventual cases by the age of 30.[11] Approximately 20% suffer from attacks of asthma.

While patients with hay fever tend to have widespread itching (nose, throat and eyes), those with perennial rhinitis rarely have eye or throat symptoms but mainly sneezing and watery rhinorrhoea. Nasal polyps are associated with this disorder (refer to p. 635).

Management

Management consists of four main areas:

1. appropriate explanation and reassurance
2. allergen avoidance
3. pharmacological treatment
4. immunotherapy

Advice to the patient

- Keep healthy, eat a well-balanced diet, avoid ‘junk food’ and live sensibly with balanced exercise, rest and recreation. If your eyes give you problems, try not to rub them, avoid contact lenses and wear sunglasses.
- Avoid using decongestant nose drops and sprays: although they soothe at first, a worse effect occurs on the rebound.
- Avoidance therapy: avoid the allergen, if you know what it is (consider pets, feather pillows and eiderdowns).
- Sources of the house dust mite are bedding, upholstered furniture, fluffy toys and carpets. Seek advice about keeping your bedroom or home dust-free, especially if you have perennial rhinitis.
- Pets, especially cats, should be kept outside.
- Avoid chemical irritants such as aspirin, smoke, cosmetics, paints and sprays.

Allergen avoidance

This is difficult during the spring pollen season, particularly where patients are living in high-pollen (e.g. country farming) areas, or spending considerable time outdoors in the course of work or sporting and recreational activities.

Pharmacological therapy[12]

Therapy can be chosen from:

1. antihistamines
    - oral (not so effective for vasomotor rhinitis)
    - intranasal spray (rapid action)
    - ophthalmic drops
2. decongestants (oral or topical)
3. sodium cromoglycate
    - intranasal: powder insufflation or spray
    - ophthalmic drops for associated conjunctivitis
4. corticosteroids
    - intranasal (not so effective for noneosinophilic vasomotor rhinitis)
    - oral (very effective if other methods fail)
    - ophthalmic drops for allergic conjunctivitis

Immunotherapy

Consider hyposensitisation/immunotherapy when specific allergens are known and conventional response is inadequate. Immunotherapy to grass pollen is generally very effective and should be considered in moderate to severe springtime hay fever.

Antihistamines

Oral antihistamines are the first line of treatment for seasonal hay fever and are generally effective where symptoms are intermittent, or when they can be used prophylactically before periods of high pollen exposure. The newer generation, so-called ‘non-sedating’ antihistamines that do not cross the blood - brain barrier are used in preference to the first-generation drugs, although some degree of sedation may occur even with these. A list of non-sedating antihistamines is presented in Table 121.3. It is claimed by some that the new topical preparations, levocabastine and azelastine, as intranasal sprays, are rapidly effective for an exacerbation of symptoms. If sedation is desirable (e.g. overnight) a sedating antihistamine can be used.

Table 121.3 Less-sedating antihistamines (oral regimens)

Generic name	Onset	Dosage
Fexofenadine	Rapid	60 mg bd
Loratadine	Very rapid	10 mg daily

Oral decongestants

Oral sympathomimetics, either used alone or in combination with antihistamines (where they may help reduce drowsiness), may be of value particularly where nasal discharge and stuffiness are major symptoms. Side-effects include nervousness and insomnia. They should be used cautiously in patients with hypertension, heart disease, hyperthyroidism, glaucoma and prostatic hypertrophy.

Examples:

- pseudoephedrine HCl 60 mg (o) tds (max. 240 mg/day), or 120 mg controlled release (o) bd.

Intranasal therapy[12]

Intranasal decongestants should be used for limited periods only (i.e. less than a week) or intermittently (3 - 4 doses per week) because of the potential problems with rebound congestion and rhinitis medicamentosa. They are often of particular value during the first week of treatment with intranasal corticosteroids (where the onset of action is delayed several days), improving nasal patency and allowing more complete insufflation of the corticosteroids. Adverse reactions similar to those of oral decongestants may occur.

Intranasal sodium cromoglycate acts by preventing mast cell degranulation and is effective without serious side-effects. The capsule variety must be used (the spray form requires 1 - 2 hourly dosage to be effective); it is useful in perennial allergic rhinitis but is not as effective as intranasal corticosteroids for springtime hay fever.

Intranasal corticosteroid sprays are the most effective agents for treating seasonal allergic rhinitis. Side-effects are minimal and adrenal suppression is not a problem with normal usage. Patients should be informed that these medications will not give immediate relief (often taking 10 - 14 days to have peak effect) and must be used continuously throughout the hay fever season.

Intranasal antihistamines group - includes azelastine and levocabastine - are effective at relieving itching and sneezing.

Table 121.4 lists intranasal preparations for rhinitis.

Table 121.4 Intranasal preparations for rhinitis

	Brand name	Dosage	Comments
Sodium cromoglycate	Rynacrom powder (capsules)	Insufflate 1 capsule, qid	Compliance a problem
	Rynacrom nasal spray	2% Spray 4 - 6 times daily	
		4% Spray 2 - 4 times daily	
Beclomethasone dipropionate 50 µg/spray	Beconase hayfever	100 µg spray each nostril bd or tds	
Budesonide	Budamax nasal Rhinocort nasal	1 - 2 sprays each nostril daily	
Fluticasone propionate 50 µg/spray	Beconase Allergy 24 hour aqueous	2 sprays each nostril daily reducing to 1 spray	
Mometasone furonate	Nasonex	2 sprays per nostril daily	
Ipratropium bromide	Atrovent	1 - 2 sprays per nostril tds prn	Useful for vasomotor rhinitis and profuse rhinorrhoea Care needed with elderly
Azelastine	Azep	1 spray each nostril bd	Antihistamine
Levocabastine 0.05%	Livostin	2 sprays each nostril bd	Antihistamine, max. 8 weeks
Various sympathomimetics (e.g. phenylephrine)		2, 3 or 4 times daily (max. 7 days)	Short-term use only Care with elderly, prostatic hypertrophy

Ophthalmic preparations

Sodium cromoglycate eyedrops are usually very effective for springtime conjunctivitis. They can be used as necessary (there is no dosage limit) and are most helpful when used prophylactically before periods of high pollen exposure. Decongestant eyedrops may also be helpful (care with narrow angle glaucoma), while corticosteroid eyedrops are reserved for resistant allergic conjunctivitis and should be used with care to exclude infection and glaucoma. Antihistamine eyedrops antazoline and levocabastine are yet another option.

Other treatments

Corticosteroids (oral)

These can be very effective where other treatments or methods have failed. A 6 - 10 day short course can be used. An example of a 6-day ‘rescue course’ is prednisolone 25, 25, 20, 15, 10, 5 mg daily doses.

Ipratropium bromide (Atrovent)[12]

The nasal preparation of this topical anti-cholinergic is often very effective when rhinorrhoea is the major problem.

Table 121.5 summarises recommended steps in management.

Table 121.5 Summary of recommended treatment steps for rhinitis[9]

Allergic Rhinitis
• Patient education
• Allergen avoidance (if possible)
Mild cases
• Less sedating antihistamines including levocabastine nasal spray ±
• Pseudoephedrine
Moderate to severe
• Inhaled corticosteroids (the most effective)
• Sodium cromoglycate (Opticrom) eyedrops
• Oral corticosteroids (if topicals ineffective)
• Immunotherapy if applicable

WHEN TO REFER

- Where surgical intervention is required, such as with nasal obstruction from polyps, bulky nasal turbinates and deviated septum
- For immunotherapy

Practice tips

- Avoid long-term use of topical decongestant nasal drops.
- Avoid topical antihistamine preparations.
- Prescribe sodium cromoglycate eyedrops for the hay fever patient with itchy eyes.
- Be careful of severe systemic reactions that can occur with intradermal skin testing and with immunotherapy. Resuscitation facilities should be available.

REFERENCES

1. Loblay RH. Allergies (type 1). In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 12 - 15.
2. O'Hehir R. Update in allergic diseases. Update Course for GPs handbook. Melbourne: Monash University, 1996: 19 - 20.
3. Smart J. *Paediatric handbook* (6th edn). Melbourne: Blackwell Science, 2000: 208 - 9.
4. Oates K, Currow K, Hu W. *Child Health: A Practical Manual for General Practice.* Sydney: MacLennan & Petty, 2001: 150 - 2.
5. Douglas R, O'Hehir R. Peanut allergy. Med J Aust, 1997; 166: 63 - 4.
6. Sampson HA, Mendleson L, Rosen JP. Fatal and near fatal anaphylactic reactions to food in children and adolescents. N Engl J Med, 1992; 327: 380 - 4.

7. Katelaris CH et al. Prevalence of latex allergy in a dental school. Med J Aust, 1996; 164: 711 - 14.
8. Walls RS. Latex allergy: a real problem. Med J Aust, 1996; 164: 707.
9. Dowden J (Chair). *Therapeutic Guidelines: Respiration* (Version 3). Melbourne: Therapeutic Guidelines Ltd, 2005.
10. Scoppa J. Rhinitis (allergic and vasomotor). In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 450 - 1.
11. Fry J. *Common diseases* (4th edn). Lancaster: MTP Press, 1985: 134 - 8.
12. Sharp A, Murtagh J. Hay fever (seasonal allergic rhinitis). Aust Fam Physician, 1995; 24: 1899 - 1900.

CHAPTER 122. ANXIETY

The lives of type A, coronary-prone personalities are dominated by time. All is rush. Extreme cases are easily diagnosed as those men who flush the lavatory before they have finished urinating.

ANONYMOUS

ANXIETY

Anxiety is an uncomfortable inner feeling of fear or imminent disaster. The criterion for anxiety disorder as defined by the ICHPPC2 (WONCA, 1985)[1] is:

> generalised and persistent anxiety or anxious mood, which cannot be associated with, or is disproportionately large in response to a specific psychosocial stressor, stimulus or event.

Anxiety is a normal human emotion and most of us experience some temporary degree of anxiety in our lives as a normal reaction to stress and misfortune. Table 122.1 presents the scale formulated by psychologists to quantify life's main stresses. However, some people are constantly anxious to the extent that it is abnormal and interferes with their lives. They suffer from an anxiety disorder, a problem that affects 5 - 10% of the population.

Table 122.1 Life change and stress survey of recent experiences

Life event	Life change units
Death of spouse	100
Divorce	73
Marital separation	65
Jail term	63
Death of close family member	63
Personal injury or illness	53
Marriage	50
Fired at work	47
Marital reconciliation	45
Retirement	45
Change in health of family member	44
Pregnancy	40
Sex difficulties	39
Business readjustment	39
Death of close friend	37
Change to different line of work	36
Mortgage (large)	31
Change in responsibility at work	29
Son or daughter leaving home	29
Outstanding personal achievement	28
Spouse begins or stops work	26
Begin or end school	26
Trouble with boss	23

Change in work hours or conditions	20
Change in residence	20
Change in schools	20
Mortgage or loan (modest)	17
Change in sleeping habits	16
Vacation	13
Christmas	12

Adapted from the *Journal of Psychomatic Research*, 1967: 11: 216.

The symptoms of anxiety, which are psychological or physical in manifestation, can vary enormously from feeling tense or tired to panic attacks.

CLASSIFICATION OF ANXIETY

The following list represents approximately the categories of anxiety disorders recognised by the DSM-IV-TR:[2,3]

- generalised anxiety disorder
- anxiety disorder due to a medical condition
- panic attack
- panic disorder, with or without agoraphobia
- agorophobia without history of panic disorder
- specific phobia
- social phobia
- obsessive compulsive disorder
- post-traumatic stress disorder
- acute stress disorder

GENERALISED ANXIETY DISORDER

Generalised anxiety comprises excessive anxiety and worry about various life circumstances and is not related to a specific activity, time or event such as trauma, obsessions or phobias. There is an overlap between generalised anxiety disorder and other anxiety disorders.

General features:

- persistent unrealistic and excessive anxiety
- worry about a number of life circumstances for 6 months or longer

Diagnostic criteria for generalised anxiety disorder

Three or more of:

- irritability
- restless, ‘keyed up’ or ‘on edge’
- easily fatigued
- difficulty concentrating or ‘mind going blank’
- muscle tension
- sleep disturbance

Clinical features

Psychological

- Apprehension/fearful anticipation
- Irritability
- Exaggerated startle response
- Sleep disturbance and nightmares
- Impatience
- Panic
- Sensitivity to noise
- Difficulty concentrating or 'mind going blank'

Physical

- Motor tension
    - muscle tension/aching
    - tension headache
    - trembling/shaky/twitching
    - restlessness
    - tiredness/fatigue
- Autonomic overactivity
    - dry mouth
    - palpitations/tachycardia
    - sweating/cold clammy hands
    - flushes/chills
    - difficulty swallowing or 'lump in throat'
    - diarrhoea/abdominal distress
    - frequency of micturition
    - difficulty breathing/smothering feeling
    - dizziness or lightheadedness

Note: Psychological disturbances are also referred to as disturbances of vigilance and scanning.

Symptoms and signs according to systems

• *Neurological*	dizziness, headache, trembling, twitching, shaking, paraesthesia
• *Cardiovascular*	palpitations, tachycardia, flushing, chest discomfort
• *Gastrointestinal*	nausea, indigestion, diarrhoea, abdominal distress
• *Respiratory*	hyperventilation, breathing difficulty, air hunger
• *Cognitive*	fear of dying, difficulty concentrating, 'mind going blank', hypervigilance

Diagnosis of anxiety

The diagnosis is based on:

- the history - it is vital to listen carefully to what the patient is saying
- exclusion of organic disorders simulating anxiety by history, examination and appropriate investigation
- exclusion of other psychiatric disorders, especially depression

Main differential diagnoses

Note that this conforms to the seven masquerades list (refer to Table 122.2):

- depression

- drug and alcohol dependence/withdrawal
- benzodiazepine dependence/withdrawal
- hyperthyroidism
- angina and cardiac arrhythmias
- iatrogenic drugs
- caffeine intoxication

Table 122.2 Significant differential diagnoses of anxiety

Psychiatric disorders • Depression • Drug and alcohol dependence/withdrawal • Benzodiazepine dependence/withdrawal • Schizophrenia • Acute or chronic organic brain disorder • Presenile dementia
Organic disorders • Drug-related - amphetamines - bronchodilators - caffeine excess - ephedrine - levodopa - thyroxine • Cardiovascular - angina - cardiac arrhythmias - mitral valve prolapse • Endocrine - hyperthyroidism - phaeochromocytoma - carcinoid syndrome - hypoglycaemia - insulinoma • Neurological - epilepsy, especially complex partial seizures - acute brain syndrome • Respiratory - asthma - acute respiratory distress - pulmonary embolism

Important checkpoints

Five self-posed questions should be considered by the family doctor before treating an anxious patient:[1]

- Is this hyperthyroidism?
- Is this depression?
- Is this normal anxiety?
- Is this mild anxiety or simple phobia?
- Is this moderate or severe anxiety?

Management of anxiety

The management applies mainly to generalised anxiety as specific psychotherapy is required in other types of anxiety. Much of the management can be carried out successfully by the family doctor using brief counselling and support.[2] CBT, in which maladaptive thinking, feelings, perceptions and related behaviours are identified, assessed, challenged and modified, can be of considerable benefit.[4]

Management principles

- The aim is to use non-pharmacological methods and avoid the use of drugs if possible.
- Give careful explanation and reassurance
    - explain the reasons for the symptoms
    - reassure the patient about the absence of organic disease (can only be based on a thorough examination and appropriate investigations)
    - direct the patient to appropriate literature to give insight and support (see Further Reading)
- Provide practical advice on ways of dealing wiThthe problems.
- Advise on the avoidance of aggravating substances such as caffeine, nicotine and other drugs.
- Advise on general measures such as stress management techniques, relaxation programs and regular exercise and organise these for the patient (don't leave it to the patient).
- Advise on coping skills, including personal and interpersonal strategies, to manage difficult circumstances and people (in relation to that patient).
- Provide ongoing supportive psychotherapy.

Further Reading

Herbert Benson	*The Relaxation Response*, Collins, London, 1984.
Dale Carnegie	*How to Stop Worrying and Start Living*, rev. edn, ed. Dorothy Carnegie, Angus and Robertson, Sydney, 1985.
Ainslie Mears	*Relief without Drugs*, Fontana, Glasgow, 1983.
Norman Peale	*The Power of Positive Thinking*, Cedar, London, 1982.
Claire Weekes	*Peace from Nervous Suffering*, Angus and Robertson, London, 1972.
Claire Weekes	*Self-help for your Nerves*, Angus and Robertson, London, 1976.

Patient education material

The author has found the following handout material to be invaluable in helping to manage less severe cases of generalised anxiety.[5]

Self-help

It is best to avoid drugs if you can; instead look at factors in your lifestyle that cause you stress and anxiety and modify or remove them (if possible). Be on the lookout for solutions. Examples are changing jobs and keeping away from people or situations that upset you. Sometimes confronting people and talking things over will help.

Special advice

Be less of a perfectionist: do not be a slave to the clock; do not bottle things up; stop feeling guilty; approve of yourself and others; express yourself and your anger. Resolve all personal conflicts. Make friends and be happy. Keep a positive outlook on life, and be moderate and less intense in your activities.

Seek a balance of activities, such as recreation, meditation, reading, rest, exercise and family/social activities.

Relaxation

Learn to relax your mind and body: seek out special relaxation programs such as yoga and meditation.

Make a commitment to yourself to spend some time every day practising relaxation. About 20 minutes twice a day is ideal, but you might want to start with only 10 minutes.[6]

- Sit in a quiet place wiThyour eyes closed, but remain alert and awake if you can. Focus your mind on the different muscle groups in your body, starting at the forehead and slowly going down to the toes. Relax the muscles as much as you can.
- Pay attention to your breathing: listen to the sound of your breaThfor the next few minutes. Breathe in and out slowly and deeply.
- Next, begin to repeat the word ‘relax’ silently in your mind at your own pace. When other thoughts distract, calmly return to the word ‘relax’.
- Just ‘let go’: this is a quiet time for yourself, in which the stresses in body and mind are balanced or reduced.

Medication

Doctors tend to recommend tranquillisers only as a last resort or to help you cope wiTha very stressful temporary period when your anxiety is severe and you cannot cope without extra help. Tranquillisers can be very effective if used sensibly and for short periods.

Pharmacological treatment

Acute episodes

The following drugs are recommended for patients who have intermittent transient exacerbations not responding to other measures:[4]

- diazepam 2 - 5 mg(o) as a single dose repeated bd as required *or*
- diazepam 5 - 10 mg (o) nocte *or*
- oxazepam 15 - 30 mg (o) as a single dose repeated bd as required

Special notes:

- Recommended (if necessary) for up to 2 weeks, then taper off to zero over next 4 weeks. Reassess in 7 days.
- Oxazepam is preferred in women patients; middle-aged men generally respond well to short-term diazepam.
- Be wary of drug-seeking behaviour (e.g. unfamiliar patients, especially if they request a specific benzodiazepine).
- Consider beta-blockers in patients with sympathetic activation such as palpitations, tremor or excessive sweating (e.g. propranolol 10 - 40 mg (o) tds).[4] They do not relieve the mental symptoms of anxiety, however.

Long-term treatment[4]

If non-pharmacological treatment is ineffective for persisting disabling anxiety, the drugs of choice are:

- venlafaxine (modified release) 75 mg (o) mane pc increasing gradually to 225 mg daily *or*
- buspirone 5 mg (o) tds4 and continue for several weeks aft er symptoms subside
    - mean effective dose is 20 - 25 mg daily

  - response takes 7 - 10 days
  - does not appear to cause sedation
- paroxetine 10 mg (o) daily, increasing to 40 mg/day

PANIC ATTACK

A panic attack is defined as a discrete period of intense fear or discomfort in which 4 (or more) of the following symptoms develop abruptly and reach a peak within 10 minutes:

- shortness of breaTh(dyspnoea) or smothering sensations
- dizziness, unsteady feelings or faintness
- palpitations or accelerated heart rate (tachycardia)
- trembling or shaking
- sweating
- choking
- nausea or abdominal distress
- depersonalisation or derealisation
- numbness or tingling sensations (paraesthesias)
- flushes (hot flashes) or chills
- chest pain or discomfort
- fear of dying
- fear of going crazy or of doing something uncontrolled

Organic disorders that simulate a panic attack are hyperthyroidism, phaeochromocytoma and hypoglycaemia.

Management

Reassurance, explanation and support (as for generalised anxiety). This is the mainstay of treatment. Patients should be taught breathing techniques to help control panic attacks.

If hyperventilating, breathe in and out of a paper bag.

Cognitive behaviour therapy

This aims to reduce anxiety by teaching patients how to identify, evaluate, control and modify their negative, fearful thoughts and behaviour. If simple psychotherapy and stress management fails then patients should be referred for this therapy.

Patients' fears, especially if irrational, need to be clearly explained by the therapist, examined rationally and challenged, then replaced by positive calming thoughts.[2]

Pharmacological treatment[4]

Acute episodes, i.e. the panic attack:

- diazepam 5 mg (o) *or*
- oxazepam 15 - 30 mg (o) *or*
- alprazolam 0.25 - 0.5 mg (o) *or*
- paroxetine 20 - 60 mg (o)

PANIC DISORDER WITH OR WITHOUT AGORAPHOBIA[4]

There are separate diagnostic criteria for panic disorder depending on whether or not there is associated agoraphobia (Table 122.3). Treatment of an acute episode is described under panic attack.

Table 122.3 Criteria for panic disorder with or without agoraphobia (DSM-IV-TR)

A The presence of both:

1 Recurrent panic attacks in which the onset of the attack is not associated wiTha situational trigger (i.e. occurring spontaneously out of the blue).

2 At least 1 of the panic attacks has been followed by 1 month (or more) of 1 (or more) of the following:

• persistent concern about having additional attacks

• worry about the implications of the attack or its consequences (e.g. losing control, having a heart attack, going crazy)

• a significant change in behaviour related to the attacks

B The presence or absence of agoraphobia.

C The panic attacks are not due to the direct physiological effects of a substance, medication, general medical condition or another mental disorder.

Prophylaxis

The evidence base regarding effectiveness indicates the number-to-treat to get one person panic-free is 3 for CBT and 5 for medication.[4] The initial approach for management should include psychological measures such as CBT. Antidepressants are the basis of first-line pharmacotherapy in panic disorder.

First-line therapy[4]

Approved antidepressants are the SSRIs, for example:

- paroxetine 10 mg (o) daily at first, increasing to 40 mg daily *or*
- sertraline 25 mg (o) daily initially, increasing to maximum 200 mg (increase for these agents is as according to tolerability and patient response)

The antidepressants may need to be continued for 6 - 12 months and then reviewed. Response may not be apparent for several weeks.

Second-line therapy

The following can be used:

- alprazolam 0.25 - 6 mg (o) daily in 2 to 4 divided doses
- tricyclic antidepressants (e.g. imipramine)

PHOBIC DISORDERS

In phobic states the anxiety is related to specific situations or objects. Patients avoid these situations and become anxious when they anticipate having to meet them. A list of specific phobias is presented in Table 122.4.

Table 122.4 Phobias

Name of phobia	Fear or aversion of
Acrophobia	height
Aerophobia	draughts
Agoraphobia	open spaces
Aichmophobia	sharp objects
Ailurophobia	cats
Algophobia	pain

Androphobia	men
Anthophobia	flowers
Anthropophobia	people
Apiphobia	bees
Aquaphobia	water
Arachnophobia	spiders
Astraphobia	lightning
Aviatophobia	flying
Bacteriophobia	bacteria
Bathophobia	depth
Belonephobia	needles
Brontophobia	thunder
Cancerophobia	cancer
Cardiophobia	heart disease
Claustrophobia	closed spaces
Cynophobia	dogs
Demonophobia	demons
Dromophobia	crossing streets
Equinophobia	horses
Genophobia	sex
Gynophobia	women
Haptephobia	being touched
Herpetophobia	creeping, crawling things
Homophobia	homosexuals
Hypsophobia	falling
Hypnophobia	going to sleep
Iatrophobia	doctors
Musophobia	mice
Mysophobia	dirt, germs, contamination
Necrophobia	death
Neophobia	anything new
Noctiphobia	night
Numerophobia	numbers
Nyctophobia	darkness
Ochlophobia	crowds
Ophidiphobia	snakes
Pyrophobia	fire
Scotophobia	blindness
Sociophobia	social situations
Taphophobia	being buried alive

Theophobia	God
Trypanophobia	injections
Xenophobia	strangers
Zoophobia	animals

The three main types of phobic states are:

- specific phobias
- agoraphobia
- social phobias (social anxiety disorder)

The 10 most common phobias (in order) are spiders, people and social situations, flying, open spaces, confined spaces, heights, cancer, thunderstorms, deaThand heart disease.[7]

Specific phobias

These are common among normal children and include fear of specific things such as snakes, spiders, thunder, darkness, dogs and heights. The problem is seldom encountered in practice and there is usually no call for drug therapy.

Agoraphobia

Avoidance includes the many situations involving the issues of distance from home, crowding or confinement. Typical examples are travel on public transport, crowded shops and confined places. The patients fear they may lose control, faint and suffer embarrassment.

The condition is commonly associated wiThdepression, obsessions, marital and family disharmony, or drug and alcohol abuse.[8]

Social phobias

Social phobias include anxiety-provoking social gatherings when the person feels subject to critical public scrutiny (e.g. canteens, restaurants, staff meetings, speaking engagements). The sufferer may be a shy, self-conscious, premorbid personality.[2] Social phobias, including performance anxiety and symptoms, are oft en related to sympathetic overactivity.

Management

The basis of treatment for all phobic disorders is psychotherapy that involves behaviour therapy and cognitive therapy.

Pharmacological treatment[4]

This should be used only if non-pharmacological measures fail.

- Agoraphobia wiThpanic: use medications as for panic attacks.
- Social phobia: if problematic, a trial of a newer antidepressant agent is recommended.[4] Antidepressants wiThevidence for modest treatment efficacy are moclobemide, paroxetine or phenelzine.
- Social phobia wiThperformance anxiety: propranolol 10 - 40 mg (o) 30 - 60 minutes before the social event or performance.
- Specific phobia: pharmacotherapy is not recommended.

OBSESSIVE COMPULSIVE DISORDER (OCD)

Anxiety is associated wiThobsessive thoughts and compulsive rituals.

The obsessions are recurrent and persistent intrusive ideas, thoughts, impulses or images that are usually resisted by the patient (e.g. a religious person having recurrent blasphemous thoughts).

Compulsions are repetitive, purposeful and intentional behaviours conducted in response to an obsession to prevent a bad outcome for the person (e.g. excessive washing of the genitals).

Mild obsessional or compulsive behaviour can be regarded as normal in response to stress.

Management[4]

Optimal management is a combination of psychotherapeutic, particularly CBT, and pharmacological treatment, namely:

- cognitive behaviour therapy for obsessions
- exposure and response prevention for compulsions
- medication:
    - any of the SSRIs e.g. paroxetine 20 - 60 mg/day *or*
    - clomipramine 50 - 75 mg (o) nocte increasing gradualy to 150 - 250 mg (o) nocte[4]

BODY DYSMORPHIC DISORDER

The person wiThthis disorder has an exaggerated preoccupation wiThan imagined defect in appearance (see p. 495).

Patients may be helped by counselling and psychotherapy.

Consider SSRIs in the context of OCD or psychotropics in the context of a psychotic disorder.

POST-TRAUMATIC STRESS DISORDER

This term describes the various symptoms and behaviour that follow a psychologically distressing event or experience outside the range of usual human experience (e.g. violent crime such as an armed hold-up, warfare, or natural disasters such as bushfires). The symptoms usually develop immediately aft er the event but can be delayed for months or years.

Typical symptoms and features:

- recurrent and intrusive distressing recollections
- recurrent distressing dreams of the event
- acting or feeling as if the event were recurring
- intense distress on exposure to resembling events
- persistent avoidance of events that symbolise or resemble the trauma
- increased arousal symptoms (e.g. insomnia, hypervigilance, exaggerated startle response, poor concentration, moodiness)

Treatment

This is difficult and involves counselling, the basis of which is facilitating abreaction of the experience by individual or group therapy.[4] The aim is to allow the patient to face up openly to memories. Persistent symptoms are an indication for referral.

Pharmacological treatment

There is no specific indication for drugs but medication can have benefit in the treatment of panic attacks, generalised anxiety or depression.[4] Long-term use of benzodiazepines is not recommended but short-term use for their antianxiety and hypnotic effects may be appropriate for the very anxious patient.

HYPERVENTILATION

Hyperventilation syndrome can be a manifestation of anxiety. The main symptoms are:

- lightheadedness, faintness or dizziness
- breathlessness
- palpitations
- sweating
- dry mouth with aerophagy
- agitation
- fatigue and malaise

Other symptoms include paraesthesia of the extremities, perioral paraesthesia and carpopedal spasm.

Management

- Reassure.
- Encourage patients to identify the cause and then control their rate and depth of breathing.
- First aid management is to raise the carbon dioxide level by rebreathing from a paper (not plastic) bag or from cupped hands (if a bag is unavailable).

ADJUSTMENT DISORDER WITH ANXIOUS MOOD

This term is reserved for patients who present with anxiety symptoms within 3 months of response to an identifiable psychosocial stressor. It is the most common presentation of anxiety symptoms and should be regarded as a separate entity to a generalised anxiety disorder.[4]

The symptoms are in excess of the normal expected reaction to the stressor but have persisted for less than 6 months following the removal of the stressor. The basic treatment is non-pharmacological - counselling, relaxation and stress management. A short-term course of drug treatment can be used in severe or persisting cases, e.g.

- diazepam 2 - 5 mg (o) daily or bd[4] *or*
- oxazepam 15 - 30 mg (o) daily or bd (up to 14 days)

SOMATISATION DISORDER

This is a chronic condition wiTha history of numerous unsubstantiated physical complaints beginning before the age of 30. The usual criteria includes four pain symptoms, two gastrointestinal symptoms, one sexual symptom and one pseudoneurological symptom (e.g. lump in throat, double vision, localised weakness), all wiThno adequate physical explanation. There is persistent refusal to be reassured that there is no explanation for the symptoms. There is associated impaired social, occupational and family functioning.

Management involves skillful counselling, explanation for symptoms, searching for and treating comorbid conditions (e.g. depression, anxiety) and CBT. It is preferable to be managed by a single supportive doctor. It is not malingering.

ANXIETY IN CHILDREN

Anxiety disorders can occur in childhood. Panic attacks are not uncommon. Children are generally more responsive to non-pharmacological approaches. Separation anxiety disorders for real, threatened or imagined separation can occur; if severe and persistent, treatment (with care) wiThimipramine is recommended.[4]

BENZODIAZEPINE USAGE

The use of benzodiazepines as anxiolytics should be restricted and they should be used discretely. Markus et al.[9] recommend reserving benzodiazepines to the following clinical situations:

1. Self-perpetuating anxiety following a precipitating event and not responding to non-pharmacological management. Give a short course for 2 weeks.
2. Situational anxiety affecting lifestyle (e.g. plane travel, dental appointments). Intermittent use only.
3. Emergency short-term use for agoraphobia or panic attacks.

They should not be used to treat depression, obsessional neuroses or chronic psychoses and should be used with caution in bereavement and crisis situations.

Problems associated with benzodiazepine use include:[1]

- impaired alertness, oversedation
- dependence
- increased risk of accidents
- adverse effects on mood and behaviour
- interaction with alcohol and other drugs
- potential for abuse and overdose
- risks during pregnancy and lactation
- muscle weakness
- sexual dysfunction
- diminished motivation
- lowered sense of competency
- lower self-esteem

Benzodiazepine withdrawal syndrome

This syndrome is usually relatively delayed in its onset and may continue for weeks or months.[1] Withdrawal features include rebound anxiety, depression, confusion, insomnia and seizures.

There are several strategies to help the consenting patient to stop benzodiazepines, ranging from stopping completely to very gradual withdrawal. An effective method is to withdraw the drug very slowly while providing counselling and support, including referral to a self-help group. Antidepressants can be substituted if there is evidence of depression, while beta-blockers may help the withdrawal syndrome if other measures have failed.

WHEN TO REFER

- If the diagnosis is doubtful
- If drug and alcohol dependence or withdrawal complicate the management
- If depression or a psychosis appears to be involved
- Failure of response to basic treatment

Practice tips

- Be careful not to confuse depression with anxiety.
- A depressive disorder can be the cause of anxiety symptoms.
- For anxiety, especially with cardiovascular symptoms (palpitations and/or flushing), always consider the possibility of hyperthyroidism and order thyroid function tests.
- Always try non-pharmacological measures to manage anxiety whenever possible.

- Be careful with the use of benzodiazepines: aim at short-term treatment only.

Tips to beat stress

- Get enough sleep and rest.
- Listen to music.
- Do things that you enjoy.
- Look at positives.
- Develop strategies to laugh.
- Go to the movies or a show weekly.
- Consider a pet.
- Your job is what you do (not who you are).
- Have regular chats wiThclose friends.
- Exercise 30 minutes, 4 to 5 times a week.
- Learn to meditate.
- Avoid interpersonal conflicts.
- Learn to accept what you cannot change.

REFERENCES

1. Wilkinson G. *Anxiety: Recognition and Treatment in General Practice*. Oxford: Radcliffe Medical Press, 1992: 5 - 62.
2. Vine RG, Judd FK. Anxiety disorders and panic states. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 43 - 5.
3. Gelder M. Diagnosis and management of anxiety and phobic states. Med Int, 1988; 45: 1857 - 61.
4. Dowden J (Chair). *Therapeutic Guidelines: Psychotropic* (Version 5). Melbourne: Therapeutic Guidelines Ltd, 2003: 147 - 61.
5. Murtagh J. *Patient Education* (4th edn). Sydney: McGraw-Hill, 2005: 169.
6. Beattie R. Anxiety: patient education. Aust Fam Physician, 1985; 14: 901.
7. Anonymous author. Making it to the top ten. *Sun*, 1990; 3 March: 67.
8. Fryer AJ. Agoraphobia. Mod Problems Pharmaco-psychiatry, 1987; 22: 91 - 126.
9. Markus AC, Murray Parker C, Tomson P et al. *Psychological Problems in General Practice*. Oxford: Oxford University Press, 1989: 12 - 43.

CHAPTER 123. ASTHMA

After I'd written to you yesterday I had an attack of asthma and an incessantly running nose, which forced me to tramp about, lighting cigarettes at every tobacconist etc. And worse was to come: I went to bed about midnight, feeling all right after spending a long time inhaling smoke, but 3 or 4 hours later came the real attack of the summer.

MARCEL PROUST 1901, LETTER TO HIS MOTHER

ASTHMA

Asthma is defined as ‘a chronic inflammatory disorder of the airways in which many cells and cellular elements play a role. In susceptible individuals this inflam-mation causes recurrent episodes of wheezing, breathlessness, chest tightness and coughing, particularly at night or in the early morning. These episodes are usually associated with widespread but variable airflow obstruction that is often reversible either spontaneously or with treatment. The inflammation also causes an associated increase in airway responsiveness to a variety of stimuli’.[1] It can also be defined as a cough or wheeze associated with heightened airway responsiveness to inhaled airway irritants such as histamine.[2] Asthma is a common and potentially fatal disorder; it is now regarded as an inflammatory disorder of the airways, which become hyperactive in the asthmatic patient.

Chronic asthma is an inflammatory disease with the following pathological characteristics:

- infiltration of the mucosa with inflammatory cells (especially eosinophils)
- oedema of the mucosa, thickening of the basement membrane
- damaged mucosal epithelium
- hypertrophy of mucus glands with increased mucus secretion
- smooth muscle constriction (Figure 123.1)



Figure 123.1 Airway changes in asthma: (a) normal airway; (b) airway in asthma

KEY FACTS AND CHECKPOINTS

- Asthma continues to be underdiagnosed and undertreated.[3] It is increasing worldwide.
- It has an unacceptable mortality rate of approximately 5 per 100 000 of the population.
- About one child in four or five has asthma (usually in a mild form).
- It tends to develop between the ages of 2 and 7 years, but can develop at any age.
- Most children present with a cough.
- Most children are free from it by puberty.
- At least one in seven adolescents has asthma.
- About one adult in eight has or has had asthma.
- The focus of management should be on prevention; an acute asthmatic attack represents failed treatment.
- Measurement of function is vital as ‘objective measurement is superior to subjective measurement’.
- Spirometry is the key investigation.

- Doubling the radius of the airway increases the flow rate 16 times.
- The earlier steroid therapy is introduced, the better the outcome.
- Inhaled corticosteroids are the cornerstone of asthma treatment.
- Avoid concomitant medication that may exacerbate asthma (e.g. beta-blockers, aspirin, NSAIDs).
- New aerosols, notably hydrofluoroalkanes, have non-CFC propellants leading to increased lung deposition and thus requiring overall lower dosage.

CAUSES OF ASTHMA

No single cause for asthma has been found, but a variety of factors may trigger an attack. These include specific factors such as viruses, allergens and non-specific factors such as temperature or weather changes and exercise. A checklist of trigger factors includes:

- infections, especially viral URTIs
- allergies (e.g. to animal fur, feathers, grass pollens, fungal allergies, mould)
- allergy to house dust, especially the dust mites and cockroaches
- cigarette smoke, other smoke and fumes
- sudden changes in weather or temperature
- occupational irritants (e.g. wood dust, synthetic sprays, chemicals)
- drugs (e.g. aspirin, NSAIDs, beta-blockers - oral, parenteral or topical)
- complementary medicines (e.g. echinacea, ‘royal jelly’)
- certain foods and food additives may trigger asthma
  - metabisulphites/sulphite preservatives/food colouring agents (e.g. tartrazine)
  - monosodium glutamate
  - seafood
  - nuts especially peanuts
- exercise, especially in a cold atmosphere
- emotional upsets or stress

Additional points

- Patients with asthma must never smoke.
- Atopic patients should avoid exposure to furred or feathered domestic animals if they have problems.
- About 90% of children with atopic symptoms and asthma demonstrate positive skin-prick responses to dust mite extract. Total eradication of house dust mite from the home is difficult.

Clinical features

The classic symptoms are:

- wheezing
- coughing (especially at night)
- tightness in the chest
- breathlessness

Note: Asthma should be suspected in children with recurrent nocturnal cough and in people with intermittent dyspnoea or chest tightness, especially after exercise.

Severe symptoms and signs are presented in the section on dangerous asthma on page 1278.

Examination[4]

Physical signs may be present if the patient has symptoms at the time of examination.

The absence of physical signs does not exclude a diagnosis of asthma. If wheeze is not present during normal tidal breathing it may become apparent during a forced expiration. Wheeze does not necessarily indicate asthma.

Absence of wheeze in a breathless person is a serious sign.

Investigations

- Measurement of peak expiratory flow rate (PEFR): demonstrates variation in values over a period of time.
- Spirometry: a value of < 75% for FEV_1/VC ratio indicates obstruction. It is the more accurate test and recommended for those who can perform it (i.e. most adults and children > 6 years).
- Measurement of PEFR or spirometry before and after bronchodilator: has a characteristic improvement for (a short acting β agonist - SAβA) > 15% in FEV_1. and PEF.
- Inhalation challenge tests: airway reactivity is tested in a respiratory laboratory to inhaled histamine, methacholine or hypertonic saline. Sometimes useful to confirm diagnosis.
- Mannitol inhalation test.
- An exercise challenge may also be helpful.
- Allergy testing may be appropriate.
- Chest X-ray: not routine but useful if complications suspected or symptoms not explained by asthma.

Six big advances in the management of asthma

1. The realisation that asthma is an inflammatory disease. Therefore the appropriate first- or second-line treatment in moderate to severe asthma is inhaled sodium cromoglycate (especially in children) or corticosteroids (ICS).
2. The regular use of spirometry.
3. The use of spacers attached to inhalers/puffers.
4. Improved and more efficient inhalers.
5. Combined long-acting relievers and preventers including combinations of long acting (agonists (LABA) and ICS.
6. Leucotriene antagonists and ante IgE agent.

Reasons for suboptimal asthma control are presented in Table 123.1.

Table 123.1 Reasons for suboptimal asthma control[1, 3]

Poor compliance
Procrastination in introducing optimal therapy
Failure to prescribe preventive medications, particularly inhaled corticosteroids for chronic asthma
Using bronchodilators alone and repeating these drugs without proper evaluation Reliance on inappropriate alternate therapies
Patient fears
• inhaled or oral corticosteroids
• concern about aerosols and the ozone layer
• overdosage
• developing tolerance
• embarrassment
• peer group condemnation
Doctor's reluctance to
• use corticosteroids
• recommend obtaining a mini peak flow meter

- recommend obtaining a spacer

TREATING INFLAMMATORY AIRWAYS DISEASE

If inflammatory airways disease is undertreated there is the risk of fixed irreversible airways obstruction from submucosal fibrosis. One of the most common mistakes in medical practice is to fail to introduce inhaled corticosteroids for the management of patients with moderately severe asthma.

MEASUREMENT OF PEAK EXPIRATORY FLOW (PEF)

Patients with moderate to severe chronic asthma require regular measurement of PEF, which is more useful than subjective symptoms in assessing asthma control. This allows the establishment of a baseline of the 'patient's best', monitors changes, and allows the assessment of asthma severity and response to treatment.

Spirometry including FEV_1 is the gold standard. Peak flow meters are not a substitute for spirometry. There is considerable variation between users and instruments. However, they have a place in helping patients self-manage their asthma by serially recording their PEFR and comparing this with the best peak flow.

LARGE VOLUME SPACERS[2]

Some people who have trouble using metered dose inhalers (MDIs) can have a special Ilspacer' fitted onto the mouthpiece of the inhaler. One puff of the MDI is put in the spacer. The patient breathes in from its mouthpiece, taking one deep inhalation, then 1 - 2 very deep breaths, or 4 - 6 normal breaths (especially in children). This method is useful for adults having trouble with the MDI and for younger children (older than 3 years). Spacers are very efficient, overcome poor technique and cause less irritation of the mouth and throat (Figure 123.2).



Figure 123.2 Using a spacer device. Rules: children - single puffs, then inhale; adults - single puff, single breath

They allow increased airway deposition of inhalant and less oropharyngeal deposition.

Note: It is recommended to dip plastic spacers into ordinary household detergent and dry in sunlight (no rinsing, no wiping) every 10 days or at least monthly.

Small volume spacers

Children under 5 - 6 years and or 20 kg can use an MDI and a small volume valved spacer (Aero Chamber, Breath-A-Tech) with a face mask.

MANAGEMENT PRINCIPLES

Aims of management:

- Absent or minimal daytime symptoms and no nocturnal symptoms. Restore normal airway function. (>80% of predicted)
- Maintain best possible lung function at all times - keep asthma under control.
- Reduce morbidity.
- Control asthma with the use of regular anti-inflammatory medication and relieving doses of β_2-agonist when necessary.

Long-term goals:

- Achieve use of the least drugs, least doses and least side-effects.
- Reduce risk of fatal attacks.
- Reduce risk of developing irreversible abnormal lung function.

DEFINITION OF CONTROL OF ASTHMA

- No cough, wheeze or breathlessness most of the time
- No nocturnal waking due to asthma
- No limitation of normal activity
- No overuse of β_2-agonists
- No severe attacks
- No side-effects of medication

THE SIX STEP ASTHMA MANAGEMENT PLAN

The National Asthma Campaign of Australia has developed this plan, which can be summarised for both patient and doctor in the following point form. An important underlying theme for the plan is careful attention to educating the patient and family.

1. *Assess the severity of the asthma* (Table 123.2)

Table 123.2 Asthma severity classification for an untreated newly diagnosed asthma patient[5]

Severity/grade	Status before treatment	Lung function FEV1 or PEF (% predicted)	Recommended β-agonist	Estimated starting daily dose range of inhaled ICS required to achieve good control
Intermittent	Episodic Symptoms < weekly Night symptoms < 2 per month Mild occasional symptoms with exercise	≥ 80%	SAβA prn	Regular ICS not required
Mild persistent	Symptoms > weekly, not every day Night symptoms > 2 per month Symptoms regularly with exercise	≥ 80%	SAβA prn	< 250 µg beclomethasone < 400 µg budesonide < 250 µg fluticasone < 160 µg ciclesonide
Moderate	Symptoms every day Night symptoms	60 - 80 %	LAβA + SAβA	250 - 500 µg

persistent	> weekly Several known triggers apart from exercise		prn	beclomethasone 400 - 800 µg budesonide 250 - 500 µg fluticasone 160 - 320 µg ciclesonide
Severe persistent	Symptoms every day Wakes frequently at night with cough/wheeze Chest tightness on waking Limitation of physical activity	< 60%	LAβA + SAβA prn	> 500 µg beclomethasone > 800 µg budesonide > 500 µg fluticasone > 320 µg ciclesonide

   - Establish the PEFR with spirometry.
   - Assess severity when patient is stable.
   - Divide into four grades of severity: intermittent/episodic, mild persistent, moderate persistent, severe persistent.

2. *Achieve best lung function*
   - Prescribe drug therapy to keep PEFR at best and to minimise symptoms.
   - Maintain the ‘best’ PEFR.
   - If PEF remains below predicted PEF level, treatment with high-dose inhaled steroids achieves optimal lung function in about 66% of patients[5] (Appendix V).
3. *Avoid trigger factors*
   - Note any domestic or occupational triggers.
   - Triggers can be inhaled or ingested.
   - If an allergen is clearly identified, avoid it (e.g. get rid of the cat, don't smoke, house dust mite avoidance strategies).
4. *Maintain best lung function with optimal medication*
   - Organise an optimal medication program.
   - Consider inhaled medications, monitored with PEF, with as few drugs, doses and side effects as possible.
   - Ensure the patient understands the difference between ‘preventer’ and ‘reliever’ medications.
5. *Develop an individualised, written action plan* (prepare an easy-to-follow action plan)
   - This must cover three points:
     - recognition that asthma is deteriorating
     - patient initiates own extra medication
     - getting access to medical attention
6. *Educate and review regularly*
   Check and review the patient regularly (both clinically and with spirometry) and provide continuing care, even when the asthma is mild. Examine inhaler technique.

PATIENT (AND FAMILY) EDUCATION

This aspect is vital and patients can be referred to an asthma education resource centre. However, the family doctor should be continually educating and encouraging the patient to follow the six-step asthma manage-ment plan.

Asthmatics tend to use ‘denial’ as a coping mechanism and are generally ‘non-attenders’ when well.

Prevention of attacks is the best treatment, and all asthmatics and their families should aim to know the disorder very well and become expert in managing it.

Know your asthma (advice for patients)

- Read all about it.
- Get to know how severe your asthma is.
- Try to identify trigger factors such as tobacco smoke and avoid them. There is an 80% increase in asthma incidence in children whose parents smoke.
- Become expert at using your medication and inhalers. A big problem is incorrect inhaler technique (35% of patients).
- Use your inhalers correctly and use a spacer if necessary.
- Know and recognise the danger signs and act promptly.
- Have regular checks with your doctor and get your peak flow rate checked by a spirometer.
- Have physiotherapy: learn breathing exercises.
- Keep fit and take regular exercise.
- Keep to ideal weight.
- Work out a clear management plan and an action plan for when trouble strikes.
- Get urgent help when danger signs appear.
- Keep at your best with suitable medications.
- Always carry your bronchodilator inhaler and check that it is not empty (learn about the water flotation test).

PHARMACOLOGICAL AGENTS TO TREAT ASTHMA

Simple classification

- Reliever = brochodilator
- Preventer = anti-inflammatory
- Symptom controller = long-acting β_2-agonist

It is useful to teach patients the concept of the ‘preventer’ and the ‘reliever’ for their asthma treatment. The pharmacological treatment of asthma is summarised in Table 123.3.

Table 123.3 Pharmacological treatment of bronchial asthma

			Vehicle of administration				
	Generic types	**Examples**	**Nebulising solution**	**Oral**	**Aerosol (metered dose inhalation)**	**Dry powder (inhalation)**	**Injection**
Bronchodilators							
1. β_2-adrenoceptor agonists	Salbutamol	Ventolin					
	Salmeterol	Serevent					
	Terbutaline	Bricanyl					
	Eformoterol	Foradile					

	Adrenaline	
2. Anticholinergics	Ipratropium bromide	Atrovent
3. Methylxanthines	Theophylline	Brondecon
		Nuelin
	Aminophylline	
Mast cell stabilisers	Sodium cromoglycate	Intal Intal forte
		Intal forte
	Nedocromil sodium	Tilade
Corticosteroids	Beclomethasone	QVAR (50, 100)
		Becloforte
		Becotide
	Budesonide	Pulmicort
	Ciclesonide	Alvesco
	Fluticasone	Flixotide
	Prednisolone	
	Hydrocortisone	Solu-cortef
Leucotriene	Montelukast	Singulair
antagonists	Zafirlukast	Accolate

'Preventer' drugs or anti-inflammatory agents

These medications are directed towards the underlying abnormalities - bronchial hyper-reactivity and associated airway inflammation.

Corticosteroids

Inhaled (ICS)

Types	beclomethasone budesonide ciclesonide (single daily dose) fluticasone
Dose range	400 - 1600 μg (adults); aim to keep below 500 μg children and 1000 μg (adults)
Availability	▪ metered dose inhaler ▪ Turbuhaler ▪ Autohaler ▪ Accuhaler
Frequency	Once or twice daily (helps compliance)
Side-effects	▪ oropharyngeal candidiasis, dysphonia (hoarse voice) - less risk with once daily ciclesonide ▪ bronchial irritation: cough ▪ adrenal suppression (doses of 2000 μg/daily; sometimes as low as 800 μg)

Note: Rinse mouth out with water and spit out after using inhaled steroids.

ICSs have a flat dose - response curve so it may not be necessary to prescribe above ICS doses considered high - beclomethasone or budesonide 1000 μg/day or fluticasone 500 μg/day. For newly diagnosed patients with mild-to-moderate asthma 'start low and step up prn' (e.g. 250 - 400 μg/day).[4]

Oral

Prednisolone is used mainly for exacerbations. It is given with the usual inhaled corticosteroids and bronchodilators.

Dose	Up to 1 mg/kg/day for 1 - 2 weeks
Side-effects	these are minimal if drug is used for short periods

Long-term use: osteoporosis, glucose intolerance, adrenal suppression, thinning of skin and easy bruising.

Oral corticosteroids can be ceased abruptly.

Cromolyns

These are sodium cromoglycate (SCG) and nedocromil sodium. SCG is available as dry capsules for inhalation, metered dose aerosols and a nebuliser solution. The availability of the metered aerosol and spacer has helped the use of SCG in the management of asthma in children. Adverse effects are uncommon; local irritation may be caused by the dry powder. Systemic effects do not occur.

Nedocromil is used for frequent episodic asthma in children over 2 years of age for the prevention of exercise induced asthma and the treatment of mild to moderate asthma in some adults. The initial dose is 2 in-halations qid. Adverse effects are uncommon.

Leucotriene antagonists

These new drugs, which include montelukast and zafirlukast, are very useful for seasonal asthma, aspirin-sensitive asthma and reduce the need for inhaled steroids or offer an alternative for those who cannot tolerate ICSs or have trouble using an inhaler. Favourable evidence is based on a small number of trials only mostly in children but same adults benefit.[6] Montelukast is taken as a 5 or 10 mg chewable tablet once daily.

'RELIEVER' DRUGS OR BRONCHODILATORS

The three groups of bronchodilators are:

- the β_2-adrenoceptor agonists (β_2-agonists)
- methylxanthines - theophylline derivatives
- anticholinergics

β_2-agonists

The inhaled route of delivery is the preferred route and the vehicles of administration include metered dose inhalation, a dry powder, and nebulisation where the solution is converted to a mist of small droplets by a flow of oxygen or air through the solution.

Oral administration of β_2-agonists is rarely required. The inhaled drugs produce meas-urable bronchodilation in 1 - 2 minutes and peak effects by 10 - 20 minutes. The traditional agents such as salbutamol and terbutaline are short-acting preparations. The new longer-acting agents (LAβA) include salmeterol and formoterol.

Theophylline derivatives

These oral drugs may have complementary value to the inhaled agents but tend to be limited by side-effects and efficacy.

Omalizumab

This anti-IgE agent is marketed for SC injection in patients > 12 years with moderate to severe allergic asthma treated by ICS and who have raised serum IgE levels.

Starting treatment

Current treatment supports the initial treatment (summarised in Table 123.2) of a SAβA with low to moderate doses of ICS with estimated equivalent doses shown in the table.

Initiate therapy with sufficient achieve best lung function promptly.

Wean inhaled corticosteroid to the minimum dose needed to maintain dequate asthma control.

Fixed dose combination medication[5,7]

- Inhaled corticosteroids
  +
  LAβA
- fluticasone + salmeterol (Seretide)
  MDI: 50/25; 125/25; 250/25 μg
  dose: adults, 2 inhalations bd
  children 4 - 12, 2 inhalations bd 50/25
  Accuhaler: 100/50; 250/50; 500/50
  dose: adults, 1 inhalation bd,
  children 4 - 12, 1 inhalation 100/50 bd
- budesonide + eformoterol (Symbicort) Turbuhaler 100/6; 200/6; 400/12; 1 - 2 inhalations bd

INDICATIONS FOR PREVENTIVE THERAPY[4]

Guidelines for introducing preventive asthma therapy in adults and children include any of the following:

- requirement of β_2-agonist > 3 - 4 times each week or > 1 canister every 3 months (excluding pre exercise)
- symptoms (non-exercise) > 3 - 4 times per week between attacks
- spirometry showing reversible airflow obstruction during asymptomatic phases
- asthma significantly interfering with physical activity despite appropriate pre-treatment
- asthma attacks > every 6 - 8 weeks
- infrequent asthma attacks but severe or life-threatening

PROPHYLACTIC AGENTS

This term is reserved for those medications that are taken prior to known trigger factors, particularly for exercise-induced asthma.

Exercise-induced asthma (options)

- β_2-agonist inhaler (puffer); two puffs 5 minutes immediately before exercise last 1 - 2 hours. LAβA such as salmeterol and eformeterol are more effective.
- SCG or nedocromil, two puffs.
- Combination β_2-agonist + SCG (5 - 10 minutes beforehand).
- Montelukast long (less in children ± 2years) (0) daily or 1 - 2 hours beforehand)
- Paediatricians often recommend a non-drug warm-up program as an alternative to medication.

A general management plan for chronic asthma is summarised in Figure 123.3.


Very mild asthma
• inhaled β2-agonist (on demand)
All other grades of severity
Regular use of:
• inhaled corticosteroid
or
a cromolyn
plus
• inhaled β2-agonist
plus
Prophylactic use prior to exercise or allergen challenge of:
• inhaled β2-agonist
and/or
• a cromolyn
First-line therapy
if control inadequate
Medium to high dose ICS plus:
• inhaled LAβA
(consider combined) ICS + LAβA
• theophylline derivation
or
• leukotriene antagonist
For exacerbations:
• oral prednisolone (intermittent use)
Second-line therapy
consider anti IgE agent
—mauzumab
—omalizumab

Figure 123.3 Management plan for chronic asthma - Adapted from Seale JP, Asthma[2] with permission of MIMS Australia, a Division of Medimedia Australia Pty Limited

Practice tip

- For breakthrough asthma or persistent poorly controlled asthma with poor compliance switch to combined medi-cation (e.g. Seretide MDI or Accuhaler).

CORRECT USE OF THE ASTHMA MDI (PUFFER)

Did you know that:

- faulty inhaler technique occurs in at least one-third of users?
- 90% of the medication sticks to the mouth and does not reach the lungs?
- it is the inhalation effort - not the pressure from the aerosol - that gets the medication to the lungs?
- it is important to instruct patients properly and check their technique regularly?

The two main techniques

The open-mouth technique and the closed-mouth technique are the main methods, and both are effective but the closed mouth technique is preferred. Both techniques are suitable for most adults. Most children from the age of 7 can learn to use puffers quite well.

The open-mouth technique (Figure 123.4)



Figure 123.4 Using the metered dose inhaler: the closed-mouth technique

Instructions for patients:

1. Remove the cap. Shake the puffer vigorously for 1 - 2 seconds. Hold it upright (canister on top) to use it (as shown).
2. Place the mouthpiece between your teeth but do not bite it and close your lips around it.
3. Breathe out slowly and gently to a comfortable level.
4. Tilt your head back slightly with your chin up.
5. Just as you then start to breathe in (slowly) through your mouth, press the puffer firmly, once. Breathe in as far as you can over 3 - 5 seconds. (Do not breathe in through your nose.)
6. Remove the puffer from your mouth and hold your breath for about 10 seconds; then breathe out gently.
7. Breathe normally for about 1 minute, then repeat the inhalation if you need to.

Common mistakes

- Holding the puffer upside down
- Inspiring through the nose after an initial puff from the MDI
- Pressing the puffer too early and not inhaling the spray deeply
- Pressing the puffer too late and not getting enough spray
- Doing it all too quickly: not breathing in slowly and holding the breath
- Squeezing the puffer more than once during a single breath
- Not breathing in deeply
- Spacer valve not functioning

Extra points

- The usual dose of standard MDI is one or two puffs every 3 - 4 hours for an attack.
- If you do not get adequate relief from your normal dose, you should contact your doctor.
- It is quite safe to increase the dose, such as to 4 - 6 puffs.
- If you are using your inhaler very often, it usually means your other asthma medication is not being used properly. Discuss this with your doctor.

AUTOHALER

The Autohaler is a breath-activated MDI which can improve lung deposition in patients with poor inhaler technique.

THE TURBUHALER

The Turbuhaler is a dry powder delivery system that is widely used as an alternative to the metered dose inhaler. It is a breath-activated device.

Other dry powder devices are the Accuhaler and Diskhaler.

SPACERS VERSUS NEBULISERS

Both MDIs via a spacer and dry powder inhalers are at least as effective as a nebuliser for treating acute exacerbations in both adults and children.[8]

Summary of devices

- Breath-actuated MDIs: Autohaler®
- Breath-actuated dry powder inhalers: Accuhaler®, Aerolizer®, Diskhaler®, Rotahaler®, Spinhaler®, Turbuhaler®
- Large-volume spacer: Nebuhaler®, Volumatic®
- Small-volume spacer: Aerochamber®, Breath-A-Tech®

DANGEROUS ASTHMA

Failure to recognise the development of a severe attack has cost the lives of many asthmatics. The severe attacks can start suddenly (even in mild asthmatics) and catch people by surprise.

High-risk patients

People who have experienced one or more of the following are more likely to have severe attacks:

- a previous severe asthma attack
- previous hospital admission, especially if to intensive care
- hospital attendance in the past 12 months
- long-term oral steroid treatment
- carelessness with taking medication
- night-time attacks, especially with severe chest tightness
- recent emotional problems
- frequent SAβA use

Early warning signs of severe asthma or an asthma attack:

- symptoms persisting or getting worse despite adequate medication
- increased coughing and chest tightness
- poor response to two inhalations
- benefit from inhalations not lasting 2 hours
- increasing medication requirements
- sleep being disturbed by coughing, wheezing or breathlessness
- chest tightness on waking in the morning
- low PEF readings

Dangerous signs

- Marked breathlessness, especially at rest
- Sleep being greatly disturbed by asthma
- Asthma getting worse quickly rather than slowly, despite medication
- Feeling frightened
- Difficulty in speaking; unable to say more than a few words
- Exhaustion and sleep deprivation
- Drowsiness or confusion
- Chest becoming 'silent' with a quiet wheeze, yet breathing still laboured
- Cyanosis
- Chest retraction
- Respiratory rate greater than 25 (adults) or 50 (children)
- Pulse rate > 120 beats/min
- Peak flow < 100 L/min or < 40% predicted FEV_1
- Oximetry on presentation (S_{aO_2}) < 90%

Asthma action plans

Examples of action plans for patients are presented below.

Action plan

If you are distressed with severe asthma:

- call an ambulance and say 'severe asthma attack' (best option) *or*
- call your doctor *or*
- if you are having trouble finding medical help, get someone to drive you to the nearest hospital

Keep using your bronchodilator inhaler continuously if you are distressed.

The following is an example of an asthma action plan that the patient keeps on a card for easy reference.

The acute severe asthma attack

Summary (adult dosage):[4, 5]

- continuous nebulised salbutamol (or terbutaline) if nebuliser available (or 12 puffs of β_2-agonist inhaler, with spacer, using one loading puff at a time followed by 4 - 5 normal tidal breaths)

Ipratropium bromide may be mixed with β_2-agonist for concurrent nebulisation.

Asthma Action Plan

Name ____________________

Contact Dr ____________ Tel ____________

Hospital tel ____________________

1. Sit upright and stay calm.
2. Take 4 separate puffs of a reliever puffer (one puff at a time) via a spacer device. Just use the puffer on its own if you don't have a spacer.
   Take 4 breaths from the spacer after each puff.
3. Wait 4 minutes. If there is no improvement, take another 4 puffs.
4. If little or no improvement **CALL AN AMBULANCE IMMEDIATELY (DIAL 000** and/or 112 from mobile phone) and state that you are having an asthma attack. Keep taking 4 puffs every 4 minutes until the ambulance arrives.

See your doctor immediately after a serious asthma attack.

- parenteral β $_2$-agonist (e.g. salbutamol 500 μg IM, SC)
- corticosteroids, e.g. prednisolone 50 mg (o) statim then daily until resolved *or* hydrocortisone 250 mg IV or IM, 6 hourly
- oxygen 8 L/min by face mask
- monitor PEF

For imminent cardiorespiratory arrest:

- adrenaline 0.5 mg 1:1000 SC, IM or 1:10,000 IV

Guidelines for spacer use in severe asthma[5]

- Frequency - every 20 minutes (first hour)
- One puff actuation at a time
- 4 - 5 normal breaths each time
- Age < 6 years
  - 6 puffs - salbutamol
  - 2 puffs - ipratropium
- Age > 6 years
  - 12 puffs - salbutamol
  - 4 puffs - ipratropium
- For moderate asthma use salbutamol only

The management of severe asthma is presented on pp. 938 and 1344.

ASTHMA IN CHILDREN

The prevalence of asthma is increasing in childhood and the management (especially in infants) is always a concern for the family doctor. The aim of treatment is to enable children to enjoy a normal life, comparable with that of non-asthmatic children with the least amount of medication and at minimum risk of adverse events. Maintainence should be determined by symptom con and lung function, especially clinical crilers since PEF- unreliat.

Key checkpoints

- Bronchodilators, inhaled or oral, are ineffective under 12 months.
- The delivery method is a problem in children and Table 123.3 gives an indication of what systems can be used at various levels.
- In the very young (e.g. 1 - 2 years old), a spacer with a face mask such as Aero Chamber or Breath-A-Tech can deliver the aerosol medication.
- The PEF rate should be measured in all asthmatic children older than 6 years. Children under 6 years generally cannot cope with the meters and those with mild asthma don't usually need PEF measurement.
- The Turbuhaler is usually not practical under 7 - 8 years.

Practice tips

- Reassure the patient that 6 - 10 inhaled doses of a β2-agonist is safe and appropriate for a severe attack of asthma.
- It is important to achieve a balance between undertreatment and overtreatment.
- Beware of patients, especially children, manipulating their peak flow.
- Get patients to rinse out their mouth with water and spit it out after inhaling corticosteroids.

- Patients who are sensitive to aspirin/ salicylates need to be reminded that salicylates are present in common cold cure preparations and agents such as Alka-Seltzer.[3]
- Aspirin-sensitive asthma usually manifests late in life with associated rhinitis. It crosssensitises with NSAIDs.
- Possible side-effects of inhaled drugs can be reduced by always using a spacer with the inhaler, using the medication qid rather than bd, rinsing the mouth, gargling and spitting out after use, and using corticosteroid sparing medications.

Prophylaxis in children

The non-steroidal medications, montelukast (oral) and SCG and/or nedocromil sodium by inhalation, are the prophylactic drugs of choice in childhood chronic asthma of mild to moderate severity.

If there is no clinical response to these agents in 4 weeks, consider use of inhaled corti-costeroids, but the risks versus benefits must always be considered. Any dose equal to or greater than 400 μg in children can have side-effects, including growth suppression and adrenal suppression. Aim for a maintenance of 100 - 400 εg, which keeps the child symptom-free. Once this stage is reached, consider stopping treatment or changing to the non-steroidal options.

Leukotriene antagonists taken orally aged 6 years and above is another option.

Guidelines for the management of asthma in children are summarised in Table 123.4. Delivery systems for children are presented in Table 123.5Table 123.5.

Table 123.4 Delivery systems for asthma in children

			Age in years	
Vehicle of administration	**Less than 2**	**2 - 4**	**5 - 7**	**8 and over**
MDI alone			*	
MDI + small volume spacer + face mask				
MDI + large volume spacer				
Nebuliser/air compressor/face mask				
Dry powder inhalers (e.g. Turbuhaler, Rotahaler)			*	
Breath-activated device			*	

* Possible in some individual children

Table 123.5 Stepwise interval management plan for children[5, 9]

Grade of asthma	Therapeutic agents
Mild - infrequent episodic:	SAβA prn
• attacks not severe	
• > 6 - 8 weeks apart	
Moderate - frequent episodic:	SAβA prn
• attacks <6 weeks apart	*and*
average every 4 - 6 weeks	• montelukast especially 2 - 5 yo: 4 mg (o) nocte
• attacks more troublesome	6 - 14 yo: 5 mg (o) nocte
	or

	• cromolyn *or* • ICS - minimum effective dose e.g. beclomethasone 100 - 200 µg/day budesonide 200 - 400 µg/day
Severe - persistent asthma:	SAβA prn
• symptoms most days	*and*
• nocturnal asthma > 1 per week	• ICS (as above)
• multiple ED visits	• consider combination LAβA + ICS Add • theophylline CR (sprinkles) • ipratropium bromide (nebuliser) • oral prednisolone (when required)

WHEN TO REFER

- If you are doubtful about the diagnosis
- For problematic children
- For advice on management when asthmatic control has failed or is difficult to achieve

REFERENCES

1. Global strategy for asthma management and preventing global initiative for asthma (GINA) updated 2004 < http://www.ginasthma.com >
2. Seale JP. Asthma. In: *MIMS Disease Index*. Sydney: IMS Publishing, 1991 - 92: 59 - 65.
3. Rees J, Price J. *ABC of Asthma* (2nd edn). London: BMJ Publishing Group, 1989: 1 - 34.
4. Improve asthma control with six-step manage-ment plan. NPS News, 23; 2002: 1 - 6.
5. Rewden (chair). *Therapeutic guidelines: Respiratory* (Version 3), Melbourne: Therapeutic guidelines Ltd., 2005: 37-70.
6. Ducharme F, Hicks GC. Anti-leukotriene agents compared to inhaled corticosteroids in the management of recurrent and/or chronic asthma. In: The Cochrane Library, Issue 1, 2002. Oxford: Update Software.
7. Worsnop C. Combination inhalers for asthma. Aust Prescr, 2005; 28: 26 - 8.
8. Cates CJ et al. Holding chambers versus nebulisers for beta-agonist treatment of acute asthma. In: The Cochrane Library, Issue 2, 2002. Oxford: Update Software.
9. Bochner F (chair). *Australian Medicines Handbook*. Adelaide: Australian Medicines Handbook Pty Ltd, 2006: 719 - 22.

CHAPTER 124. CHRONIC OBSTRUCTIVE PULMONARY DISEASE

> Tobacco drieth the brain, dimmeth the sight, vitiateth the smell, hurteth the stomach, destroyeth the concoction, disturbeth the humors and spirits, corrupteth the breath, induceth a trembling of the limbs, exsiccateth the windpipe, lungs and liver, annoyeth the milt, scorcheth the heart, and causeth the blood to be adusted.
>
> TOBIAS VENNER (1577 - 1660), *VIA RECTA AD VITAM LONGAM*

CHRONIC OBSTRUCTIVE PULMONARY DISEASE

Chronic obstructive pulmonary disease (COPD) is a respiratory disease characterised by airflow obstruction that is not fully reversible. The airflow limitation is generally both progressive and associated with an abnormal inflammatory response of the lungs to noxious airborne agents, especially cigarette smoke. As outlined on page XXX, it incorporates features of emphysema and airway obstruction with airway wall thickening and narrowing.

COPD typically affects middle-aged and older people with the usual age of onset in the fifth and sixth decades. It is the fourth leading cause of death and the third leading burden of disease in Australia, affecting 12.4% of Australians between 45 and 70 years.

Cigarette smoking is undoubtedly the major cause of both chronic bronchitis and emphysema, although only 10 - 15% of smokers develop the diseases.[1]

Figure 124.1 illustrates the influence of smoking on lung function.



Figure 124.1 Decline of FEV_1 with age. Adapted from Fletcher and Peto [2]

FACTORS IN CAUSATION

- Cigarette smoking (usually 20/day for 20 years or more)[3]
- Air pollution (outdoor and indoor)
- Airway infection
- Occupation: related to cadmium, silica, dusts
- Familial factors: genetic predisposition
- $Alpha_1$-antitrypsin deficiency (emphysema)
- Bronchial hyperresponsiveness

DIAGNOSIS AND MANAGEMENT OF COPD

The COPDX Plan guidelines[4] developed by the Australian Lung Foundation and the Thoracic Society of Australia and New Zealand provide an appropriate framework for diagnosis and management. The key recommendations are Confirm diagnosis, Optimise function, Prevent deterioration, Develop a self-management plan and manage eXacerbations.

C - Confirm diagnosis and assess severity

Symptoms

- Breathlessness
- Cough main symptoms
- Sputum production
- Chest tightness
- Wheezing
- Airway irritability
- Fatigue
- Anorexia with advanced disease
- Weight loss

Consider the diagnosis of COPD in all smokers and ex-smokers older than 35 years. The diagnosis of COPD rests on the demonstration of airflow and obstruction.

The sensitivity of the physical examination for detecting mild to moderate COPD is poor.[5]

Signs

The signs vary according to the nature of the disease and the presence of infection. Signs may be completely absent in the early stages of COPD and there may be wheezing only with chronic bronchitis and dyspnoea with chronic airflow limitation.

Signs may include:

- tachypnoea
- reduced chest expansion
- hyperinflated lungs
- hyper-resonant percussion
- diminished breath sounds ± wheeze
- ‘pink puffer’ - always breathless
- ‘blue bloater’ - oedematous and central cyanosis
- signs of respiratory failure
- signs of cor pulmonale

The diagnosis is usually clinical with a history of increasing dyspnoea and sputum production in a lifetime smoker. It is unwise to make a diagnosis of chronic bronchitis and emphysema in the absence of cigarette smoking unless there is a family history suggestive of alpha$_1$-antitrypsin deficiency.[7]

Investigations

Pulmonary function tests

Spirometry remains the gold standard for diagnosing, assessing and monitoring COPD. The PEFR is not a sensitive measure.

COPD is defined as post-bronchodilator FEV_1/FVC of < 0.70 ($< 70\%$) and $FEV_1 < 80\%$ predicted.

Chest X-ray

This can be normal (even with advanced disease) but characteristic changes occur late in disease. May exclude lung cancer > 1 cm.

Blood gases

- May be normal.
- $PaCO_2 \uparrow$; $PaO_2 \downarrow$ (advanced disease)

ECG

This may show evidence of cor pulmonale.

Sputum culture

If resistant organism suspected

FBE

- To identify anaemia and polycythemia
- Haemoglobin and PVC may be raised.

O - Optimise function

The principle goals of therapy are to stop smoking, to optimise function through relief of symptoms with medication and pulmonary rehabilitation, and to prevent or treat aggravating factors and complications.

Table 124.1 is a useful consultation checklist mnemonic.[6]

Table 124.1 SMOKES, a consultation checklist for chronic obstructive pulmonary disease[6]

S smoking cessation
M medication - inhaled bronchodilator, vaccines (influenza, pneumococcus), corticosteroids (if indicated)
O oxygen - is it needed?
K komorbidity - cardiac dysfunction, sleep apnoea, osteoporosis, depression, asthma
E exercise and rehabilitation
S surgery - bullectomy, lung volume reduction surgery, single-lung transplantation

Long-term treatment

Advice to patient

- If you smoke, you must stop (persuading the patient to stop smoking is the key to management). The only treatment proven to slow the progression of COPD is smoking cessation.[3] Nicotine replacement therapy should be considered.
- Avoid places with polluted air and other irritants, such as smoke, paint fumes and fine dust.
- Go for walks in clean, fresh air.
- A warm dry climate is preferable to a cold damp place (if prone to infections).
- Get adequate rest.
- Avoid contact with people who have colds or flu.
- Optimal diet - reduce weight if necessary.

Physiotherapy

Refer to a physiotherapist for chest physiotherapy, breathing exercises and an aerobic physical exercise program.

Drug therapy[7]

In the long-term treatment of COPD, bronchodilators are recommended for the relief of wheezing and shortness of breath. These include short-acting β $_2$-agonists (salbutamol, terbutaline) and short-acting anticholinergic drugs (ipratropium bromide); long-acting β $_2$-agonists (eformoterol, salmeterol); long-acting anticholinergic drugs (tiotropium); and corticosteroids.

The preferred route of administration of bronchodilator is by inhalation but it requires correct device technique.

Inhaled drugs can be administered by MDIs, dry-powder devices or nebulisers. The evidence suggests that an MDI and spacer are as effective as a nebuliser, but the appropriate method depends on patient needs and preference.

The usefulness of a bronchodilator for an individual can only be assessed by a therapeutic trial, accepting either objective improvement in lung function or improvement in symptom control as end points.

Short-acting bronchodilator therapy

Most studies suggest that short-acting β $_2$-agonists and ipratropium bromide are equally efficacious in patients with COPD. Table 124.2 shows a plan for initial treatment with short-acting bronchodilators.[8] If patients do not respond adequately to one of these bronchodilators then they should be given a trial with a second bronchodilator. It is also appropriate to consider a trial of a combination of the two classes of bronchodilator with objective monitoring of response.

Table 124.2 A plan for initial treatment with short-acting bronchodilators

COPD severity	FEV, % predicted	Suggested treatment
Mild	60 - 80%	Intermittent bronchodilator as needed before exercise
Moderate	40 - 50%	Regular combined therapy, e.g. salbutamol + ipratropium
Severe	< 40%	Add long-acting bronchodilator ± inhaled glucocorticosteroids

Modified from GOLD[12]

Use the following by inhalation:[7]

- salbutamol 100 - 200 μg, up to 4 times daily *or*
- terbutaline 500 μg, up to 4 times daily *or*
- ipratropium bromide 40 - 80 μg, up to 4 times daily

For patients with poor inhalation technique, the use of a large-volume spacer improves lung deposition of the aerosol.

For patients who are unable to use an MDI or a spacer, a nebuliser should be used, with the following doses:

salbutamol or terbutaline 2.5 to 5 mg and/or

ipratropium bromide 250 to 500 μg by nebuliser, up to 4 times a day

Long-acting bronchodilator therapy

Long acting β $_2$-agonists can be used in patients who remain symptomatic despite treatment with combinations of short-acting bronchodilators and those with frequent exacerbations. Used regularly they are more effective and convenient than use of short-acting bronchodilators (evidence level I).[9] Long-acting anticholinergic therapy[1] with tiotropium bromide (taken by inhalation) has been proven to reduce the frequency of exacerbations with COPD compared with short-acting anticholinergic drugs. The choice of drug can be determined by the patient's response to a trial of the drug, the drug's adverse effects and cost including PBS listing.

For treatment with long-acting bronchodilator use the following by inhalation:[7]

- tiotropium bromide 18 μg once daily *or*
- eformoterol 12 μg once daily *or*
- salmeterol 50 μg twice daily

Corticosteroids[7]

Only 10 % of patients with stable COPD benefit in the short term from corticosteroids. There are no distinguishing clinical features to predict in advance which patients may respond. The aim of treatment is to reduce exacerbation rates and slow the decline of the disease. Note that there may be possible coexisting asthma with COPD. Benefits are not seen in patients who continue to smoke.

Inhaled corticosteroids

Guidelines for prescription include:

- documented evidence of responsiveness to inhaled corticosteroids, including functional status
- those with an FEV_1 ? 50% predicted
- two or more exacerbations requiring oral steroids in 12 months
- Begin treatment with the following by inhalation:
- beclomethasone dipropionate 300 - 400 µg, twice daily *or*
- budesonide 800 µg, twice daily *or*
- fluticasone propionate 250 - 375 µg, twice daily

Reduce the doses gradually to the minimum dose that maintains subjective benefit. Warn patients about the risk of osteoporosis and advise them to rinse their throat and mouth with water and spit out after inhalation. Patients using an MDI should be advised to use a spacer.

Oral corticosteroids are not recommended for maintenance therapy in COPD, although they may be needed in patients with severe COPD where corticosteroids cannot be withdrawn following an acute exacerbation. A stepwise approach to management is outlined in Figure 124.2.


COPD with dyspnoea
smoking cessation
short-acting bronchodilator
(inhalation verses nebuliser)
*
combination
β agonist + ipratropium
*
long-acting β agonist or tiotropium
*
inhaled corticosteroid
*
refer

Figure 124.2 Stepwise approach to management of COPD

P - PREVENT DETERIORATION[4]

Reducing risk factors for COPD is a priority, and smoking is the most important and prime target if this continues to be a problem. Stopping is the only measure that slows the progression of COPD. Reinforce patient education programs.

Annual influenza vaccination

Influenza vaccination reduces the risk of exacerbations, hospitalisation and death. It should be given in early autumn to all patients, especially those with moderate to severe COPD.

Pneumococcal vaccination

Vaccination to prevent invasive bacteraemic pneumococcal pneumonia is recommended. It should be given at 5 yearly intervals.

Long-term oxygen therapy (LTOT)

LTOT reduces mortality in COPD. Long-term continuous therapy given for at least 15 hours a day (as close as possible to 24 hours a day) prolongs life in hypoxaemic patients - those who have PaO_2 consistently <55 mm Hg (7.3 KPa; SpO_2 88%) when breathing air. At assessment for ongoing therapy the patient's condition must be stable and the patient must have stopped smoking at least 1 month previously. Flow should be set at the lowest rate needed to maintain a resting PaO_2 of 60 mm Hg. A flow rate of 0.5 - 2.0 L/min is usually sufficient. There is no clear-cut evidence about the effectiveness of intermittent ambulatory domiciliary oxygen therapy, but patients with hypoxaemia during sleep may require nocturnal oxygen therapy.

Check current smoking status

Smoking cessation clearly reduces the rate of decline of lung function. General practitioners and pharmacists can help smokers quit. Brief counselling is effective and every smoker should be offered at least this intervention at every visit.

Refer to strategies in Chapter 22, pages 202 - 4, including effectiveness of treatment for nicotine dependence.

Antibiotics

Current evidence does not support long-term antibiotic use to prevent exacerbations, but they should be used in exacerbations with an increase in cough, dyspnoea, sputum volume or purulence.

Corticosteroids

No medication has yet been shown to prevent the long-term decline in lung function. Inhaled corticosteroids are indicated for patients with a documented response or those who have severe COPD with frequent exacerbations.

Mucolytic agents

Mucolytic agents may reduce the frequency and duration of exacerbations (evidence level I). Mucolytic therapy should be considered for patients with a chronic cough productive of sputum. Oral mucolytics include potassium iodide, bromhexine, N-acetylcysteine, ambroxal and glycerol guaiacolate. Compounds containing codeine should be avoided.

Antitussive agents

Regular use of antitussives in stable COPD is contraindicated.

Regular review

Regular review with objective measures of function and medication review is recommended in anticipated of reducing complications, frequency and severity of exacerbations and admissions to hospital. Regular schedules for follow-up visits are appropriate.

Lung surgery[7]

The options are bullectomy, lung volume reduction surgery[8] and transplantation. Patients should be referred for consideration for bullectomy if they have a single large bulla on CT scan associated with breathlessness and an $FEV_1 < 50\%$ predicted. Other lung surgery should be considered in patients with severe COPD who remain breathless with marked restriction of their activities of daily living, despite maximal therapy. Emphysema mainly involving the upper lobes with $PaCO_2 < 55$ mm Hg and FEV > 20% predicted are some factors required for lung volume reduction surgery.

D - DEVELOP SUPPORT NETWORK AND SELF-MANAGEMENT PLAN[4]

COPD imposes a considerable handicap with heavy psychosocial issues, including fears about the outcome of the disease, on patients and carers.

Respiratory physician referral

Early referral to a respiratory physician is appropriate in order to clarify the diagnosis, consider other therapies, consider long-term home oxygen and facilitate organisation of pulmonary rehabilitation.

Pulmonary rehabilitation

One highly effective strategy is pulmonary rehabilitation, which aims to increase patient and carer knowledge and understanding, reduce carer strain and develop positive attitudes toward self-management and exercise. Integrated programs include education, exercise, behaviour modification and support, which are more effective than separate components.

The support team and multidisciplinary care plans

As the patient's primary health care provider the GP is uniquely placed to identify smokers and help them quit, facilitate early diagnosis and coordinate the support team. The support team can enhance the quality of life and reduce morbidity for the COPD patient. The support team can include a nurse/respiratory educator, physiotherapist, occupational therapist, social worker, clinical psychologist, speech pathologist, pharmacist and dietician.

Government and community support services such as HomeCare, home maintenance, exercise programs, Meals on Wheels and support groups can be galvanised to provide support.

Self-management plans

Patients should be encouraged to take appropriate responsibility for their own management. The primary care team, supported by Extended Primary Care item numbers, should develop systems to identify those with more severe COPD who might benefit from more focused education and training in self-management skills.

Psychological issues

The issues facing the COPD patient include fear, stress, sleep disturbance, anxiety, panic and depression. Pro-active management, including optimal care of these problems as they arise, will facilitate coping. Management also focuses on symptom control and maximising the quality of life. These patients have to face palliative care at the end stage and ethical issues have to be handled sensitively.

Referral to in-patient care

The management plan should include the identification of clinical markers indicating more intensive hospital treatment.

Indications include:[4]

- rapid rate of onset of acute exacerbation
- inability to cope at home
- inability to walk between rooms when previously mobile
- severe breathlessness ? inability to eat or sleep
- inadequate response to ambulatory treatment

- altered mental status suggestive of hypercapnia
- significant comorbidity, e.g. cardiac disease
- new arrhythmia
- cyanosis

X - MANAGE EXACERBATIONS[4]

Diagnosis is symptomatically the acute onset over minutes to hours of at least two of the following:

- increasing dyspnoea including use of accessory muscles at rest
- reduced effort tolerance
- tachypnoea (> 25 breaths/min)
- increased fatigue
- increased cough and sputum
- increased wheezing

Fever may be present, but fever and chest pain are uncommon symptoms. One-third have no identifiable cause, but infections and heavy pollutants can cause the exacerbation. Consider investigating with pulse oximetry, chest X-ray and sputum culture.

Patients should be treated with a bronchodilator, preferably with a large volume spacer. If nebulisers are used they should be driven by compressed air (to avoid the problem of potentially adverse oxygenation). Systemic glucocorticoids reduce the severity and shorten recovery.

TREATMENT IN SUMMARY[7]

Bronchodilators

Initial therapy by inhalation:

- salbutamol 100 μg MDI, up to 8 to 10 inhalations, repeat as required *or*
- terbutaline 500 μg DPI, 1 to 2 inhalations, repeat as required *or*
- ipratropium bromide 20 μg MDI, up to 4 - 6 inhalations, repeat as required

If control is inadequate, combine salbutamol or terbutaline with ipratropium bromide.

If a nebuliser is used (which is usually the case upon hospitalisation), use salbutamol 2.5 - 5 mg, terbutaline 2.5 - 5 mg or ipratropium bromide 250 - 500 μg, as required.

Oxygen therapy[7]

Controlled oxygen delivery - 28% via Venturi mask or 2 L/min via nasal prongs should be commenced if the patient is hypoxaemic (oxygen saturation < 92% with pulse oximetry). Maintain the arterial oxyhaemoglobin saturation at 90%. It is important to obtain a direct measurement of arterial blood gases to confirm the degree of hypoxaemia and if hypercapnia or acidosis is present.

Note that patients with severe COPD are prone to hypercapnia if they breathe high oxygen concentrations. Supplemental oxygen should be kept to a minimum. If hypercapnia develops assisted ventilation may be required.

Corticosteroids

Corticosteroids should be used routinely for severe exacerbations: use

- prednisolone or prednisone 30 to 50 mg (0), daily

If oral medication cannot be tolerated, use:

- hydrocortisone 100 mg IV 6 hourly (or equivalent dose of alternate corticosteroid)

Conversion to oral corticosteroid should occur as soon as practicable.

The duration of oral corticosteroid therapy for exacerbations of COPD is not well established; however, courses of 7 to 14 days are commonly used.

A tapering dose schedule is not required for short courses.

Antibiotics[7]

The use of antibiotics for exacerbations is not routinely indicated as many episodes are due to viral infections. There are some patients who obviously have repeated exacerbations due to bacterial infection (usually *Haemophilus influenza, S. pneumoniae or Moraxella catarrhalis*) where antibiotics have been proven to be beneficial, reducing the risk of mortality by 77%.[10]

Practice Tips

- COPD patients should be referred early for rehabilitation. Contact your respiratory physician or hospital for help.
- Pulmonary rehabilitation programs benefit most patients with pulmonary disease.
- Rehabilitation teams are interdisciplinary, usually comprising a rehabilitation physician, physiotherapist, occupational therapist, social worker and dietician.
- Patients with COPD commonly present in the fifth decade with productive cough or an acute chest illness.[3]
- Diagnosis can only be established by objective measurement using spirometry with FEV_1 being the preferred parameter.
- Non-invasive positive pressure ventilation (NPPV) reduces mortality and hospital stay in patients with acute failure; it is also an effective weaning strategy for patients who require intubation.[10]
- The only treatment proven to slow the progression of COPD is smoking cessation.
- It is very difficult at times to distinguish COPD from the persistent airflow limitation of chronic asthma in older patients (see Table 49.3, p. 526).
- COPD is reportedly ‘massively’ underdiagnosed - screening with good quality spirometry is important.[13]

The indication for antibiotic treatment is:

- increased cough and dyspnoea *together with*
- increased sputum volume and/or purulence

When indicated, use:

- amoxicillin 500 mg (o), 8 hourly for 5 days *or*
- doxycycline 200 mg (o) as 1 dose on day 1, then 100 mg (o) daily for a further 5 days

A stepwise approach is summarised in Figure 124.3.


COPD
acute exacerbation
↑ short-acting bronchodilators
consider combination
β agonist + ipratropium
• check smoking
• check inhaler device/technique
severe
oral corticosteroids
consider
antibiotics (if indicated)
oxygen (if hypoxaemic)
*
refer

Figure 124.3 Management plan for acute exacerbation of COPD

EVIDENCE UPDATE[11, 12]

Evidence indicates that that regular treatment with long-acting ß $_2$-agonists is more effective than treatment with short-acting agents (evidence level I) and is associated with improved quality of life (evidence level II), although they are more costly and do not significantly improve lung function. Current recommended guidelines based on the severity of disease are summarised in Table 124.3.

Table 124.3 COPD therapy according to severity of disease[11, 12]

Stage of COPD	Treatment
0 At risk	Avoidance of risk factors, esp. smoking Influenza and pneumococcal vaccination ? *Haemophilus influenza* vaccination
1 Mild	*Add* short-acting bronchodilator
2 Moderate	*Add* one or more bronchodilators, including long-acting bronchodilator *Add* pulmonary rehabilitation
3 Severe	*Add* inhaled glucocorticosteroids
4 Very severe	*Add* long-term oxygen (if chronic respiratory failure) Consider surgical referral

REFERENCES

1. Tierney LM et al. *Current Medical Diagnosis and Treatment* (41st edn.). New York: McGraw and Lange, 2002: 290.
2. Fletcher C, Peto R. The natural history of chronic airflow obstruction. BMJ, 1977: 1645 - 48.
3. COPD. NPS News, 1999; 5: i - iv.

4. McKenzie DK, Firth PA, Burdon JG, Town GI. The COPDX Plan: Australian and New Zealand Guidelines for the management of chronic obstructive pulmonary disease 2003. Supplement MJA, 17 March 2003; vol. 178.
5. Badgett RC, Tanaka DV, Hunt DL et al. Can moderate chronic obstructive pulmonary disease be diagnosed by history and physical findings alone? Am J Med 1993; 94: 188 - 96.
6. Gibson PG. Management of chronic obstructive pulmonary disease. Australian Prescriber, 2001; 24(6): 152.
7. Dowden J (Chair). Writing Group. *Therapeutic Guidelines: Respiratory* (Version 3). Melbourne: Therapeutic Guidelines Ltd, 2005.
8. Snell GI, Solin P, Chin W. Lung volume reduction surgery for emphysema. Med J Aust, 1997; 167: 529 - 32.
9. Dahl R, Greefhorst LA, Nowak D et al. Inhaled formoterol dry powder versus ipratropium bromide in COPD. Am J Respir Crit Care Med 2001; 164: 778 - 84.
10. Antibiotics for COPD exacerbations. The Cochrane Database of Systematic Reviews 2006; Issue 2, online 19 April.
11. Abramson MJ, Crockett AJ, Frith PA, McDonald CF. COPDX: an update of guidelines of chronic COPD with a review of recent evidence. MJA 2006; 184: 342 - 5.
12. NHLBI/WHO Workshop Report. *Global Initiative for Chronic Obstructive Lung Disease (GOLD): Global Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease*. Bethesda, MD: National Institute of Health - National Heart, Lung and Blood Institute, 2005. Refer: http://www.goldcopd.org.
13. Petty TL. The history of COPD. International J COPD, 2006; 1: 3 - 14.

CHAPTER 125. EPILEPSY

> The fit makes the patient fall down senseless; and without his will or consciousness presently every muscle is put in action; as if all the powers of the body were exerted to free itself from some great violence. In these strong and universal convulsions, the urine, excrements and seed, are sometimes forced away, and the mouth is covered with foam, which will be bloody, when the tongue is bit, as it often is in the agony.
>
> WILLIAM HEBERDEN (1718 - 1801)

EPILEPSY

Epilepsy is defined as a ‘tendency to recurrence of seizures’. It is a symptom, not a disease. A person should not be labelled ‘epileptic’ until at least two attacks have occurred.[1] Epilepsy is common and affects about one person in 50. Both sexes are about equally involved, and it seems to run in some families. Famous people who have had epilepsy include Julius Caesar, Thomas Edison and Handel.

The most important factors in the man-agement of epilepsy are the accurate diagnosis of the type of seizures; identification of the cause and appropriate investigation; the use of first-line drugs as the sole therapy of some weeks; and adjustment of the dose, according to clinical experience and plasma levels, to give maximum benefit.

To be accurate in diagnosing seizures the diagnosis must be based on:

- the observation of a witness to the seizures
- a general and neurological examination
- an EEG, although this has considerable limitations
- a CT scan or preferably MRI (especially if the EEG is focal and a tumour is suspected)

Long-term ambulatory EEG recording now provides more information and, coupled with video monitoring, it will make a permanent record of the seizure which can be reviewed at will. The CT scan or MRI scan is necessary to exclude a focal cause (such as a cyst, tumour, malformation or abscess) which might be treatable by surgery. The MRI scan now shows developmental migration disorders.

The scans will identify mesotemporal sclerosis (an abnormality in the hippocampus due to birth hypoxia), thereby making some ‘idiopathic’ seizures into secondary seizures from a known cause.

Epilepsy usually starts in early childhood.

Optimal management includes adequate psychosocial support with education, coun-selling, advocacy and appropriate referral.

An underlying organic lesion becomes more common in epilepsy presented for the first time in patients over the age of 25 years and thus more detailed investigation is required.[1]

Secondary causes of seizures include:

- trauma - head injuries
- post cerebral surgery
- metabolic (e.g. calcium and sodium electrolyte disturbances, hypoglycaemia, uraemia, hepatic failure)
- drugs and other toxins (e.g. alcohol, amphetamines including withdrawal)
- infections (e.g. meningitis)
- vascular - cerebrovascular, arteritis
- hypoxia
- degenerative diseases
- sleep deprivation

TYPES OF EPILEPTIC SEIZURES/SYNDROMES

Epileptic seizures are classified in general terms as generalised and partial (Table 125.1). Partial seizures are about twice as common as generalised seizures and usually due to acquired pathology.[2]

Table 125.1 Classification of epileptic seizures

1 Generalised seizures
Convulsive
tonic - clonic (previously called grand mal)
clonic
Non-convulsive
tonic (drop attacks)
atonic (drop attacks)
absence (petit mal)
atypical absence
myoclonic
2 Partial seizures (non-convulsive)
Simple partial (consciousness retained)
with motor signs (Jacksonian)
with somatosensory symptoms
with psychic symptoms
Complex partial (consciousness impaired)
3 Secondary generalised seizures (convulsive)
Simple partial seizures evolving to tonic - clonic seizures
Complex partial seizures evolving to tonic - clonic seizures
Simple partial seizures evolving to complex partial seizures and then to tonic - clonic seizures

Generalised seizures

The epileptic discharge affects both cerebral hemispheres simultaneously from the outset. The seizure may be primary (no focus) or sec-ondary due to acquired cerebral pathology.

The main features are:

- abrupt impairment or loss of consciousness
- possible bilateral symmetrical motor events

Types

- Tonic - clonic (formerly called grand mal) seizure: this is the classic convulsive seizure with muscle jerking (p. 831)
- Tonic seizure: stiffness only, often with a ‘drop’ (hallmark of Lennox Gastaut syndrome)
- Clonic seizure: jerks only
- Atonic seizure: loss of tone, and ‘drops’
- Absence seizure (formerly called petit mal): involves loss of consciousness with no or only very minor bilateral muscle jerking, mainly of the face[2] (p. 599)
- Myoclonic seizure: bilateral discrete muscle jerks, which may be very severe, and loss of consciousness

Partial seizures

In partial seizures the epileptic discharge begins in a localised focus of the brain and then spreads out from this focus. The clinical pattern depends on the part of the brain affected:

- simple partial seizures: consciousness is retained (p. 599)
- complex partial seizures: consciousness is clouded so that the patient does not recall the complete seizure (p. 598).

Both these types of partial seizure can evolve into a bilateral tonic - clonic seizure; this is termed a secondary generalised seizure and is usually due to diffuse brain pathology.[1]

Investigations

Standard minimum investigations are:

- serum calcium magnesium and electrolytes
- fasting glucose
- EEG (usually with sleep deprivation)
- syphilis serology

Other tests may include:

- chest X-ray
- ECG (? ↑ QT internal)
- brain scan
- video EEG (limited mainly to frequent seizures or to diagnostic dilemmas)
- MRI
- CT scanning (if MRI unavailable)

A patient presenting with the first seizure after the age of 25 years will require more detailed investigation.

The single unprovoked seizure: to treat or not

Starting drugs after a single afebrile seizure can be a difficult decision. It basically depends on the EEG findings. If not treated the risk of recurrence over a 3-year period is 40% with a risk of approximately 25% if treated.[3]

APPROACHES TO MANAGEMENT

- An accurate diagnosis of the seizure type is essential.
- An underlying brain disease has to be investigated and treated.
- Treatment is based on drugs and lifestyle management
- A decision has to be made about whether drug therapy is appropriate. Pharmacotherapy should be offered if the patient has had two or more seizures within 6-12 months.[3, 4]
  Most seizures require long-term antiepileptic (anticonvulsant) drug therapy aimed at suppressing the underlying seizure activity in the hope that it may subside, so that 'cure' ultimately occurs and treatment may be ceased. A summary of antiepileptic drugs is presented in Table 125.2.
- The choice of drug depends on the seizure type, on consideration of the age and sex of the patient and on efficacy in relation to toxicity.
- Treatment should be initiated with one drug and pushed until it controls the events or causes side-effects, irrespective of the medication blood level. The disorder can usually be controlled by one drug provided adequate serum or plasma concentrations are reached.[4]
- If a maximum tolerated dosage of this single drug fails to control the seizures, replace it with an alternative agent with a different action. Add the second drug and obtain a therapeutic effect before removing the first drug.
- Special attention should be given to the adverse psychological and social effects of epilepsy. Emotional and social suport is important and advice about epilepsy support groups is appropriate.

Table 125.2 Antiepileptic drugs

The following antiepileptic drugs are used
Benzodiazepines
• clonazepam
• diazepam
• midazolam
• nitrazepam
Carbamazepine
Tetracosactin
Phenobarbitone and related drugs
• methylphenobarbitone
• primidone
Phenytoin
Sodium valproate (valproate)
Succinimides
• ethosuximide
New drugs (mainly as ‘added on’ therapy)
Clobazam
Felbamate
Gabapentin
Lamotrigine
Levetiracetam
Oxcarbazepine
Pregabilin
Sulthiame
Tiagabine
Topiramate
Vigabatrin

Drug therapy

The following are important specific consider-ations:

- The basic aim is to prevent seizure recurrence preferably with monotherapy and to minimise adverse effects.
- It is best to select the most effective recommended drug for a specific seizure type (Table 125.3).
- Young women prefer carbamazepine to phenytoin because of the adverse effects of gingival hypertrophy and hirsutism of phenytoin.
- Each drug has specific adverse effects (Table 125.4) while all drugs tend to be sedating, especially phenobarbitone and its derivatives.
- Twice daily dosage is usually practical.
- Phenytoin should be increased in small increments (25 - 50 mg) above a dose of 300 mg daily.
- Phenytoin or carbamazepine will bring about control in at least 80% of patients with tonic - clonic seizures.[1] Do not use these drugs together as they have a similar action.

Table 125.3 Recommended selection of antiepileptics in epilepsy[3, 4]

Type of seizure	First-line therapy	Second-line therapy (select from)
Tonic-clonic	Valproate Carbamazepine	Phenytoin Oxcarbazepine Lamotrigine Gabapentin Tiagabine Topiramate Phenobarbitone
Absence (petit mal)	Ethosuximide Valproate (especially if associated with other seizure type)	Clobazam Clonazepam Lamotrigine
Myoclonic	Sodium valproate	Clobazam Clonazepam Lamotrigine
Simple partial (Jacksonian) and Complex partial	Carbamazepine	Phenytoin Valproate Topiramate Gabapentin Lamotrigine Tiagabine Phenobarbitone
Infantile spasms	Tetracosactin	Clonazepam Valproate Nitrazepam

Table 125.4 Commonly used antiepileptics: usual oral dose and adverse reactions

	Usual starting adult dose	Average satisfactory adult dose (mg/kg/day)	Therapeutic plasma concentration range (μmol/L)	Significant adverse reactions
Carbamazepine	100 bd, increasing gradually to control (max. 2 g/day) Usual range: 400 mg - 1.2 g daily	30	25 - 50 free level 6 - 13	Anorexia, nausea, vomiting, dizziness, drowsiness, skin rashes, tinnitus, ataxia, diplopia Small risk of spina bifida in fetus Drug interactions (e.g. COC, warfarin, other anticonvulsants)
Clonazepam	0.25 mg bd increasing to control	0.1 - 0.2	0.08 - 0.24	Drowsiness, fatigue, muscle weakness, ataxia, dizziness

				Respiratory problems
				Interacts with alcohol
Ethosuximide (used only in absence seizures)	20 - 30 mg/kg/day in 2 divided doses	30	300 - 700	Anorexia, nausea, vomiting
	usual starting dose: 250 mg bd		(all free)	Diarrhoea
				Drowsiness, ataxia, headache
Lamotrigine	25 mg/day increasing to 100 - 200 mg/day	1.5 - 5	-	Beware of blood dyscrasias
				Drowsiness, skin rashes, nausea, headache, dizziness, ataxia
				Beware of severe rashes: introduce slowly
Phenobarbitone	60 - 240 mg daily at bed time	2 - 4	45 - 130	Drowsiness
	usual dose: 60 - 120 mg			Dizziness, ataxia
				Skin rashes
				Mood changes (e.g. excitable)
				Interacts with warfarin, COC, other anticonvulsants
Phenytoin	5 mg/kg/day in 2 divided doses	5 - 6	40 - 80	Drowsiness, fatigue, mental confusion, ataxia, nystagmus, slurred speech, anorexia, dizziness, nausea, vomiting
	average adult: 300 mg		free level	Skin reactions
			4 - 8	Gum hypertrophy, hirsutism
				Fetal malformations (e.g. cleft lip and palate, congenital heart disorders)
Sodium valproate	500 mg daily for 7 days then bd, increasing to achieve control (up to 2 - 3 g/day)	20 - 30	300 - 750	Drowsiness, tremor, hair loss
		Standard dose: 500 mg mane 1000 mg nocte	free level	Platelet effect
			30 - 75	Risk of neural tube defects in fetus
				Hepatic failure
				Interacts with other anticonvulsants
				To be avoided in pregnancy
Vigabatrin	500 mg daily for 7 days then bd increasing to 1 - 2 g bd, according to response	30 - 40	Not relevant	Drowsiness
				Behavioural disturbances

				Dizziness
Gabapentin	300 mg bd or tds, increasing to 3.6 g daily (max)	30 - 50	-	Drowsiness, ataxia, gastrointestinal, dizziness

Cessation of drug therapy

It is important to review the need for antiepileptics every 12 months. They may be ceased if the patient has been free of seizures for at least 3 years particularly if epileptiform activity has disappeared from the EEG (best under specialist supervision). Up to 60% of children have a mild self-limiting condition and can settle after medication is withdrawn.

Adverse drug reactions

Patients should be warned about significant side-effects:

- nausea, dizziness, ataxia, visual disturbance or excessive tiredness/fatigue indicate excessive dosage of carbamazepine or phenytoin
- most drugs can cause a rash
- gingival hyperplasia is a classic effect of phenytoin
- hirsutism can occur with phenytoin while hair loss can occur with sodium valproate
- sodium valproate has rare but potentially serious toxic effects
  - liver toxicity
  - dysmorphogenic effects (specifically spina bifida) on foetus during pregnancy (therefore it is a risk in women of child-bearing age, especially if inadvertent pregnancy occurs due to pill failure related to antiepileptic interaction)
- LFTs should be performed every 2 months for 6 months after starting sodium valproate;[1] liver toxicity is much more common in the under 2 years old age group

PATIENT EDUCATION

Points worth emphasising:

- Most patients can achieve complete control of seizures.
- Most people lead a normal life - they can expect to marry, have a normal sexual life and have normal children.
- Patients need good dental care, especially if they are taking phenytoin.
- A seizure in itself will not usually cause death or brain damage unless in a risk situation such as swimming, or prolonged.
- Patients cannot swallow their tongue during a seizure.
- Take special care with open fires.
- Encourage patients to cease intake of alcohol. Intoxication is very harmful.
- Adequate sleep is important. Deprivation is harmful.
- Avoid fatigue
- Advise showering in preference to bathing.

Driving

One has to be very careful about driving. Most people with epilepsy can drive but each case has to be considered individually. Applicants for learner's licence need to be seizure-free for 2 years, with an annual medical review for 5 years following receipt of the licence. Restrictions range from 1 month to 2 years, depending on the circumstances of the seizures. For a new patient the usual rule is suspension of driving until 3 months seizure free.

Employment

People with epilepsy can hold down most jobs, but if liable to seizures they should not work close to heavy machinery, in dangerous surroundings, at heights (such as climbing ladders) or near deep water. Careers are not available in some services, such as the police, military, aviation (pilot, traffic controller) or public transport (e.g. bus driver).

Sport and leisure activities

Most activities are fine, but epileptics should avoid dangerous sports such as scuba diving, hang-gliding, parachuting, rock climbing, car racing and swimming alone, especially surfing.

Table 125.5 outlines contraindications for sporting activities. These apply to patients who suffer from very frequent seizures, especially the complex partial seizures with prolonged postictal states.[4]

Table 125.5 Sporting activities: contraindications[5]

Absolute contraindications
• flying and parachuting • motor racing • mountain and rock climbing • high diving • scuba diving • underwater swimming, especially competitive • hang-gliding • Abseiling
Relative contraindications
• aiming sports such as archery and pistol shooting • contact sports such as boxing, rugby, football, including soccer, where heading the ball is involved • competitive cycling for children with absence epilepsy • bathing and swimming • gymnastics, especially activities such as trampolining and climbing on bars • ice skating and skiing • javelin throwing

Avoid trigger factors

- Fatigue
- Lack of sleep
- Physical exhaustion
- Stress
- Excess alcohol
- Prolonged flashing lights if photosensitive (e.g. video games - this applies to those with a proven response to a proper EEG with photic stimulation)

EPILEPSY IN CHILDREN

Refer to Chapter 54, on faints, fits and funny turns (p. 594).

Photosensitive epilepsy in children

Some children suffer from photosensitive epilepsy related to exposure to computer and video games. There is some evidence that such children may not have seizures if they keep one eye covered. If television provokes seizures, strategies such as watching it with ambient lighting and using the remote control rather than approaching the set will minimise the problem.

EPILEPSY IN THE ELDERLY

The elderly require the same diagnostic approach as any other age group. However, they are sensitive to side-effects of drugs and may require a lower dose than younger patients. It must be remembered that antidepressant or major tranquilliser drugs can precipitate a generalised seizure.

PREGNANCY AND EPILEPSY[6]

Pregnancy is associated with a 30% increase in seizures. Although the outcome is successful for more than 90% of epileptic women, there is a slightly increased risk of prematurity, low birth weight, mortality, defects and intervention. About 45% of women have an increased number of seizures, due mainly to a fall in antiepileptic drug levels.

All antiepileptic drugs are potentially teratogenic, with different drugs being related to different defects: phenytoin has been related to cleft lip and palate and congenital heart disease, while sodium valproate (in particular) and carbamazepine have been associated with spina bifida. All antiepileptic drugs are expressed in breast milk but in such reduced concentrations as not to preclude breastfeeding.

PITFALLS IN MANAGEMENT OF EPILEPSY[3]

Misdiagnosis

The main pitfall associated with seizure disorders and epilepsy is misdiagnosis. It should be realised that not all seizures are generalised tonic - clonic in type. The most common misdiagnosed seizure disorder is that of complex partial seizures (an underdiagnosed disorder) or the tonic or atonic seizures.

The diagnosis of epilepsy is made on the history rather than the EEG so a very detailed description of the events from eyewitnesses is important.

The features of complex partial seizure (**described** on p. 598) have many variations, the commonest being a slight disturbance of perception or consciousness. The complex partial seizure may evolve to a generalised tonic - clonic seizure. A simple partial seizure may also do this.

In tonic - clonic seizures the patient may become momentarily rigid or fall to the ground and perhaps have one or two jerks only.

Misdiagnosing behavioural disorders

It is important to differentiate between a fit and a behavioural disorder, but it can be difficult. About 20% of apparently intractable ‘seizures’ are considered to be non-epileptic (pseudo-seizures, i.e. emotionally based).[7] Ancillary testing, especially with video EEG recording, can help overcome these diagnostic problems but the differentiation may be difficult as the most common situation for pseudoseizures is in the person who has real fits.

Overtreatment

Polypharmacy

Polypharmacy may be counterproductive for the patient and the seizure disorder. This is especially applicable to drugs with a high incidence of side-effects. If a patient is taking several medications, management of the case needs questioning and perhaps reconsidering with a consultant's help.

Seizure control may be improved by reducing polypharmacy. When initiating treatment it is best to select one drug and increase its dose until its maximum recommended level, the onset of side-effects or appropriate control. If control is not obtained, the drug should be replaced with an alternative agent but a crossover period is essential. Monotherapy is preferred but combination therapy is often acceptable.

Prolonged treatment

The question should be asked at some stage 'Does this patient really need medication?'. Some patients are kept on antiepileptics for too long without any attempt being made to wean them off medication or to transfer them onto antiepileptics less prone to side-effects. Patients should not be left on inappropriate drugs especially if side-effects and drug interactions are a problem.

Drug interactions

Drug interactions with antiepileptics should always be kept in mind. The most serious of all is the interaction with the oral contraceptive pill because pregnancy can occur. Erythro-mycin and carbamazepine interact.

MANAGEMENT OF STATUS EPILEPTICUS

Definition

Status epilepticus is the situation in which a patient suffers from two or more generalised seizures without regaining consciousness be-tween seizures, or suffers from continuous partial seizures.[3]

Focal status

- A high index of suspicion is needed to diagnose
- Oral medication usually adequate
- Avoid overtreatment

Generalised status

Absence attack (petit mal)

- Hospitalisation
- IV diazepam

Tonic - clonic (dangerous!)[3]

- Ensure adequate oxygenation: attend to airway (e.g. Guedel tube); give oxygen
- Diazepam 0.05 mg/kg/minute IV until the seizures cease or respiratory depression begins (beware of respiratory depression and other vital parameters) usually 10 - 20 mg bolus in adult *or*
- clonazepam 1 - 2 mg IV statim then 0.5 - 1 mg/min IV until seizures cease or respiratory depression begins *or*
- midazolam 0.05 - 0.1 mg/kg IV (max. 10 - 15 mg) or 0.15 mg/kg IM followed by
- phenytoin 1000 mg IV over 20 - 30 minutes

Other drugs to consider:

- phenobarbitone; thiopentone; paraldehyde; valproate (can use rectally)

Note: Midazolam is suitable for all types of seizures and can be given IM or intranasally.

Diazepam can be given rectally. In an adult, 10 mg is diluted in 5 mL of isotonic saline and introduced via the nozzle of the syringe into the rectum. The dose in children is 0.5 mg/kg.

Intranasal midazolam (0.3 - 0.5 mg/kg/dose) up to maximum of 10 mg may be given in children.

Practice tips

- The EEG has considerable limitations in diagnosis. It is diagnostic in less than 50%,[1] although more diagnostic if conducted under sleep deprivation. An accurate eyewitness account of the seizure is the most reliable diagnostic aid.
- During evaluation look for evidence of neurofibromatosis.
- Beware of interactions between antiepileptics and the oral contraceptive pill.
- Interactions between erythromycin and carbamazepine can cause toxicity.
- Always aim to achieve monotherapy.
- An important toxic reaction can occur with combined phenytoin and carbamazepine.
- Patients should not drive while medication is being adjusted, particularly if weaning is being attempted.
- The development of sophisticated surgical techniques means that surgery can be used in selected patients with poor control. Evaluation for surgery is a very specialised area.

DOs and DON'Ts for the onlookers of a seizure

- Don't move the person (unless necessary for safety).
- Don't force anything into the person's mouth.
- Don't try to stop the fit.
- Do roll the person on to his or her side with the head turned to one side and chin up.
- Do call for medical help if the seizure lasts longer than 10 minutes or starts again.
- Do remove false teeth and help clear the airway once the fit is over.

WHEN TO REFER[2]

Specialist referral is advisable under the following circumstances:

- uncertainty of diagnosis
- at onset of seizure disorder to help obtain a precise diagnosis
- when seizures are not controlled by apparent suitable therapy ? wrong drug ? suboptimal dose ? progressive underlying disorder
- when the patient is unwell, irrespective of laboratory investigation
- when a woman is considering pregnancy (preferable) or has become pregnant: to obtain therapeutic guidance during a difficult phase of management
- for assessment of the prospects for withdrawing treatment after some years of absolute seizure control

REFERENCES

1. Scott AK. Management of epilepsy. In: Central Nervous System. London: BMJ Publishing Group, 1989: 1 - 2.
2. Eadie MJ. Epilepsy. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 175 - 8.
3. Tiller J (Chair). Therapeutic Guidelines: Neurology (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2002: 91 - 102.
4. Bochner F (Chair). *Australian Medicines Hand book*. Adelaide: Australian Medicines. Handbook Pty ltd. 2006; 595 - 610.
5. Cordova F. Epilepsy and sport. Aust Fam Physician, 1993; 22: 558 - 62.
6. Kilpatrick C. Epilepsy poses special problems. Aust Dr Weekly, 1993; 19 February: 48.
7. Theodore HR. Neurotrek. A PC based system for automated recording of epileptic seizures and similar events. Patient Management, 1992; 16: 15 - 16.

CHAPTER 126. HYPERTENSION

The greatest danger to a man is that someone will discover hypertension and some fool will try to reduce it.

JOHN HAY 1931

HYPERTENSION

Hypertension is a serious community disorder and the most common condition requiring long-term drug therapy in Australia. It is a silent killer because most people with hypertension are asymptomatic and unaware of their problem. Epidemiological studies have demon-strated the association between hypertension and stroke, coronary heart disease, renal disease, heart failure and atrial fibrillation. Treatment may be lifelong, hence the need for careful work-up.

- Target organs (including some specific examples) that can be damaged by hypertension include the heart (failure, LVH, ischaemic disease), the kidney (renal insufficiency), the retina (retinopathy), the blood vessels (peripheral vascular disease, dissecting aortic aneurysm) and the brain (cerebrovascular disease).
- Deaths in hypertensive patients have been shown to be due to stroke 45%, heart failure 35%, renal failure 3% and others 17%.[1]
- Factors increasing chances of dying in hypertensive patients are: male patient; young patient; family history; increasing diastolic pressure.[1]

DEFINITIONS AND CLASSIFICATION

For adults aged 18 years and older hypertension is:

- diastolic pressure > 90 mmHg and /or
- systolic pressure > 140 mmHg

- The various categories of BP are arbitrarily defined according to BP values for both diastolic and systolic readings (Table 126.1).[2,3]

Table 126.1 Definition and classification of blood pressure (mmHg)[2, 3]

Category	Systolic	Diastolic
Optimal	< 120	< 80
Normal	120 - 129	80 - 84
High Normal	130 - 139	85 - 89
Grade 1 hypertension (mild)	140 - 159	90 - 99
Grade 2 hypertension (moderate)	160 - 179	100 - 109
Grade 3 hypertension (severe)	≥ 180	≥ 110
Isolated systolic hypertension	≥ 140	< 90
Borderline	140 - 149	< 90

- Isolated systolic hypertension is that of ≥ 140 mmHg in the presence of a diastolic pressure < 90 mmHg
- Hypertension is either essential or secondary (Table 126.2).

Table 126.2 Classification of hypertension

Essential (90 - 95%)
Secondary (approximately 5 - 10%)
Kidney (3 - 4%) • glomerulonephritis • reflux nephropathy > • kidney artery stenosis • other renovascular disease • diabetes Endocrine • primary aldosteronism (Conn's syndrome) • Cushing's syndrome • phaeochromocytoma • oral contraceptives • other endocrine factors Coarctation of the aorta Immune disorder (e.g. polyarteritis nodosa) Drugs Pregnancy

- Essential hypertension is the presence of sustained hypertension in the absence of underlying, potentially correctable renal, adrenal or other factors.
- Malignant hypertension is that with a diastolic pressure > 120 mmHg and exudative vasculopathy in the retinal and renal circulations.
- Refractory hypertension is a BP > 140/90 mmHg despite maximum dosage of two durgs for 3 - 4 months.

When a patient's systolic and diastolic blood pressures fall into different categories, the higher category should apply.

RISK STRATIFICATION AND CALCULATION OF CARDIOVASCULAR RISK [2,3]

Treatment of hypertension is generally indefinite and it is important to establish the risk status and the prognosis prior to commencing therapy, especially if hypertension is an isolated factor. The WHO-15th recommendation is that decision of the management of patients with hypertension should be based on BP alone but also on the presence or absence of other risk factors, including important factors such as age, diabetes and smoking. Cardiovascular risk should be stratified according to the BP level and the presence of:

- absence cardiovascular risks factors
- associated clinical conditions
- target organ damage (Table 126.3)

Table 126.3 Factors influencing prognosis in hypertension

Risk factors for cardiovascular disease used for stratification	Associated clinical condition (ACC)	Target organ damage (TOD)

• Levels of systolic and diastolic BP	• Cerebrovascular disease	• Left ventricular hypertrophy
• Men > 55 years	- ischaemic stroke	• Microalbuminuria and /or proteinuria and/or eGFR < 60 ml/minute
• Women > 65 years	- cerebral haemorrhage	
• Smoking	- transient ischaemic attack	
• Diabetes mellitus	• Heart disease	• Ultrasound or angiographic evidence of atherosclerotic disease
• Dyslipidaemia (total cholesterol > 6.5 mmol/1, > 250 mg/dl or LDL-cholesterol > 4.0 mmol/l, > 155 mg/dl or HDL-cholesterol M <1.0, W <1.2 mmol/l, M < 40, W < 48 mg/dl)	- myocardial infarction	
	- angina	
	- coronary revascularization	• Hypertensive retinopathy (grade II or more)
	- congestive heart failure	
	• Kidney disease	
	- diabetic nephropathy	
• Family history of premature cardiovascular disease (at age < 55 years M, < 65 years W)	- kidney impairment	
	- proteinuria (> 300 mg/24 h)	
	• Peripheral vascular disease	
• Abdominal obesity (abdominal circumference M ≥ 102 cm, W ≥ 88 cm)	• Advanced retinopathy	
	- haemorrhages or exudates	
	- papilloedema	
• C-reactive protein ≥ 1 mg/dl		
	Other factors effecting prognosis	
	• Excessive	

	alcohol intake • Sedentary lifestyle • High risks socio-economic group • High risk ethnic group

A Practical approach to stratify total cardiovascular risk is proposed in table 126.4. The terms low, moderate (medium), high and very high added risk are calibrated to indicate an absolute 10 year risk of cardiovascular disease of <15%, 15 - 20%, 20 - 30% and > 30% respectively (based on Framingham criteria). For example, low risk Indicates starting treatment and monitoring, high risk indicates treating immediately.

Table 126.4 Stratification of cardiovascular risk to quantify prognosis[2]

Other risk factors and disease history	Normal BP	High normal BP	Mild hypertension	Moderate hypertension	Severe hypertension
No other risk factors	Average risk	Average risk	Low risk	Moderate risk	High risk
1 or 2 risk factors but not diabetes	Low risk	Low risk	Moderate risk	Moderate risk	Very high risk
3 or more risk factors or target organ damage or diabetes mellitus	Moderate risk	High risk	High risk	High risk	Very high risk
Associated clinical conditions	High risk	Very high risk	Very high risk	Very high risk	Very high risk

Risk estimation can be determined by referring to various cardiovascular risk tables on the website (and that appear at the end of chapter 128, 'Diabetes'). A commonly used tool in Australia is the modified New Zealand cardiovascular risk calculator (http://www.nzgg.org.nz or http://www.heartfoundation.com.au).

It is important to collaborate with patients in decision making, and thus discussing cardiovascular risk assessment and blood pressure level should be the starting point when discussing the risks and benefits of treatment.

SECONDARY HYPERTENSION

Secondary hypertension may be suggested in patients below 40 years by the history (Table 126.5), physical examination, severity of hypertension or the initial laboratory findings. Table 126.5 Clinical features suggesting secondary hypertension[4]

Table 126.5 Clinical features suggesting secondary hypertension[4]

Clinical features	Likely cause
Abdominal systolic bruit	Kidney artery stenosis
Proteinuria, haematuria, casts	Glomerulonephritis
Bilateral kidney masses with or without haematuria	Polycystic disease
History of claudication and delayed femoral pulse	Coarctation of the aorta
Progressive nocturia, weakness	Primary aldosteronism (check serum potassium)
Paroxysmal hypertension with headache, pallor, sweating, palpitations	Phaeochromocytoma

The most common causes of secondary hypertension are various renal diseases, such as renovascular disease, chronic glomer-ulonephritis, chronic pyelonephritis (often associated with reflux nephropathy) and analgesic nephropathy.[1] There will often be no physical findings to suggest the existence of such renal diseases, but an indication will generally be obtained by the presence of one or more abnormalities when the urine is ex-amined. Clinical pointers include proteinuria, an abnormal urine sediment, general atheroma, smokers and abdominal bruit.

Physical findings that may suggest secondary hypertension include epigastric bruits (indicating possible renal artery stenosis) and abdominal aortic aneurysm. Less common findings include abdominal flank masses (polycystic kidneys), delayed or absent femoral pulses (coarctation of the aorta), truncal obesity with pigmented striae (Cushing's syndrome), and tachycardia, sweating and pallor (phaeochromocytoma).

Further investigation will be required to confirm or reveal secondary hypertension.

Conn's syndrome: clinical features

- Weakness due to hypokalaemia
- Polyuria and polydypsia
- Na ↑, K ↓, alkalosis
- Aldosterone ↑ (serum and urine)
- Plasma renin ↓

Phaeochromocytoma: clinical features

Paroxysms or spells of:

- hypertension
- headache (throbbing)
- sweating
- palpitations
- pallor/skin blanching
- rising sensation of tightness in upper chest and throat (angina can occur)

Investigation: 24 hour urinary free catecholamines ↑ (VMA)

Kidney artery stenosis

Atherosclerotic kidney artery stenosis accounts for the majority of cases while fibromuscular dysplasia remains an important cause. Doppler ultrasound is a highly sensitive and specific investigation.

DETECTION OF HYPERTENSION[1]

Hypertension can only be detected when BP is measured. Therefore every reasonable opportunity should be taken to measure BP.

Diagnosis should not be made on the basis of a single visit. Initial raised BP readings should be confirmed on at least two other visits within the space of 3 months; average levels of 90 mmHg diastolic or more, or 140 mmHg systolic or more, are needed before hypertension can be diagnosed. This will avoid the possibility of an incorrect diagnosis, committing an asymp-tomatic normotensive individual to unjustified, lifelong treatment.

Measurement[2]

BP varies continuously and can be affected by many outside factors. Care should therefore be taken to ensure that readings accurately represent the patient's usual pressure. The essential steps in this process are outlined below.

Office BP measurement guidelines[2]

- Allow the patient to sit quickly for several minutes.
- Use a validated calibrated device.
- Take at least two measurements spaced by 60 seconds.
- Use a standard bladder 12-13 x 35 cm), but a larger one for big arms.
- Have the cuff at the heart level.
- Deflate the cuff slowly (2 mmHg/s).
- Measure BP also in standing position in elderly patients and diabetic patients.

Position

Patients should be seated with their bare arm supported and positioned at heart level. Any sleeve should be loose above the sphygmo-manometer cuff.

Cuff size and placement

A cuff size that will completely occlude the brachial artery is essential for accurate measurement. Cuffs that are too short or too narrow may give falsely high readings. The cuff's rubber bladder should have a width of at least 40% of the circumference of the patient's arm and a length at least double that. The commonly used cuffs are often shorter than this recommendation. Suitable cuffs are made by Trimline (PyMaH) and Accosan. Several sizes, including cuffs for children and the obese, should be available. The bladder of the cuff deteriorates over about 2 years and should be replaced at intervals.[5]

A preferable cuff placement is to have the tubes emerging from the bladder point proximally (Figure 126.1) thus leaving the cubital fossa free.[5]



Figure 126.1 Correct placement of the cuff

Equipment

Ideally, measurement should be taken with a reliable and properly maintained sphygmomanometer, Otherwise, a recently calibrated aneroid manometer or a calibrated electronic in-strument can be used. It is essential that all equipment is maintained regularly. All tubing connections should be airtight.

Palpation

Initially, systolic BP should be recorded by the palpatory method at the radial or brachial artery (the brachial artery is felt just medial to the biceps tendon in the cubital fossa). This will prevent low auscultatory systolic pressures caused by a ‘silent gap’.

Note this reading and add 30 mmHg to it as the upper level to which to inflate the cuff, while accurately measuring the

BP with the bell of the stethoscope over the brachial artery.

Taking the reading

While the BP is being measured, the cuff should be deflated at a rate no greater than 2mmHg for each beat. One of the commonest errors is to allow the column of mercury to fall too rapidly.

Pressure should be recorded to the nearest 2 mmHg (it should not end in 5). Parallax errors should be avoided when reading levels (Figure 126.2).[5, 6] Wait 60 seconds and repeat the BP measurement.



Figure 126.2 Correct eye level: the observer should be within 1 metre of the manometer, so the scale can be easily read to avoid a parallax error - the eye should be on the same horizontal level as the mercury meniscus

Recording

On each occasion when the BP is taken, two or more readings should be averaged. Wait at least 30 seconds before repeating the procedure. If the first two readings differ by more than 6 mmHg systolic or 4 mmHg diastolic, more readings should be taken.

Both systolic and diastolic levels should be recorded. For the diastolic reading the dis-appearance of sound (Phase 5) - that is, the pressure when the last sound is heard and after which all sound disappears - should be used.[7] This is more accurate than the muffling of sounds (phase 4) (Figure 126.3), which should only be used if the sound continues to zero.


Artery
Auscultatory sounds
Korotkoff's sounds
sphygmomanometer cuff
stethoscope
no blood flow
Systolic pressure
Phase 1 (sound first heard)
Phase 2 (sound increases)
partial flow
Phase 3 (sound decreases in intensity)
Phase 4 (sound becomes muffled)
Diastolic pressure
Phase 5 (sound finally disappears)
free flow

Figure 126.3 Illustration of blood pressure measurement in relation to arterial blood flow, cuff pressure and auscultation

Note: Measure sitting and standing BP is elderly and diabetic patients.

Heart rate and pulse

At the same time the BP is measured, the heart rate and rhythm should be measured and recorded. A high heart rate may indicate undue stress that is causing the associated elevated BP reading. An irregular heart rhythm may cause difficulty in obtaining an accurate BP reading.

BP modifying factors

Apprehension: Patient should be rested for at least 5 minutes and made as relaxed as possible.

Caffeine: Patients should not take caffeine for 4 - 6 hours before measurement.

Smoking: Patients should also avoid smoking for 2 hours before measurement.

Eating: Patients should not have eaten for 30 minutes.

Strategies for high initial readings

If the initial reading is high (DBP > 90 mmHg, SBP > 140 mmHg) repeat the measures after 10 minutes of quiet rest. The ‘white coat’ influence in the medical practitioner's office may cause higher readings so measurement in other settings such as the home or the workplace should be managed whenever possible.

Confirmation and follow-up[1]

Repeated BP readings will determine whether initial high levels are confirmed and need attention, or whether they return to normal and need only periodic checking. Particular attention should be paid to younger patients to ensure that they are regularly followed up.

Initial diastolic BP readings of 115 mmHg or more, particularly for patients with target organ damage, may need immediate drug therapy.

Once an elevated level has been detected, the timing of subsequent readings should be based on the initial pressure level, as shown in Table 126.6.

Table 126.6 Follow-up criteria for initial blood pressure measurement for adults 18 years and older[8]

Systolic (mm Hg)	Diastolic (mm Hg)	Action/recommended follow-up*
< 120	< 80	Recheck in 2 years.
120 - 139	80 - 89	Recheck in 1 year - lifestyle advice.
140 - 159	90 - 99	*Confirm within 2 months - lifestyle advice.
160 - 179	100 - 109	*Evaluate or refer within 1 month - lifestyle advice.
≥ 180	≥ 110	*Further evaluate and refer within 1 week (or immediately depending on clinical situation). If blood pressure has been confirmed at ≥ 180 mm Hg systolic and/or ≥ 110 diastolic mm Hg (after multiple readings and excluding 'white coat' hypertension), drug treatment should be commenced.

If systolic and diastolic categories are different, follow recommendations for shorter follow-up (e.g. BP 160/86 mm Hg evaluate or refer within 1 month).

*Note: Earlier initiation of drug therapy may be indicated for some patients.

If mild hypertension is found, observation with repeated measurement over 3 - 6 months should be followed before beginning therapy. This is because levels often return to normal.

Ambulatory 24 hour monitoring

This is not required for the diagnosis and follow-up of most hypertensive patients but in some patients with fluctuating levels, borderline hypertension or refractory hyper-tension (especially where the 'white coat' effect may be significant) ambulatory monitoring has a place in management. Studies have shown that this method provides a more precise estimate of BP variability than casual recordings. This has implications for the timing of drug therapy in individual patients.

Guidelines for ambulatory BP measurement:

- Unusual variability of office BP
- Marked discrepancy between office and house BP
- Resistance to drug treatment
- Suspected sleep apnoea

'White coat' hypertension

This group may comprise up to 25% of patients presenting with hypertension. These people have a type of conditioned response to the clinic or office setting and their home BP and ambulatory BP profiles are normal. They appear to be at low risk of cardiovascular disease but may progress to sustained hypertension. Ambulatory 24 hour monitoring has a place in managing these subjects.

EVALUATION

As well as defining the BP problem, the clinical evaluation for suspected hypertension should also determine:

- whether the patient has potentially reversible secondary hypertension
- whether target organ damage is present
- whether there are other potentially modifiable cardiovascular risk factors present and
- what co-morbid factors exist

MEDICAL HISTORY

The following should be included in the medical history of the patient:

History of hypertension

- Method and date of initial diagnosis
- Known duration and levels of elevated BP
- Symptoms that may indicate the effects of high BP on target organ damage, such as headache, dyspnoea, chest pain, claudication, ankle oedema and haematuria
- Symptoms suggesting secondary hypertension (Table 126.5)
- The results and side-effects of all previous antihypertensive treatment

Presence of other diseases and risk factors

- A history of cardiovascular, cerebrovascular or peripheral vascular disease, renal disease, diabetes mellitus or recent weight gain
- Other cardiovascular risk factors, including obesity, hyperlipidaemia, carbohydrate intolerance, smoking, salt intake, alcohol consumption, exercise levels and analgesic intake
- Other relevant conditions, such as asthma or psychiatric illness (particularly depressive illness)

Family history

Particular attention should be paid to the family history of hypertension, cardiovascular or cereb-rovascular disease, hyperlipidaemia, obesity, dia-betes mellitus, renal disease, alcohol abuse and premature sudden death.

Medication history

A history of all medications, including over-the-counter products, should be obtained because some can raise BP or interfere with anti-hypertensive therapy. Prohypertensive substances oral contraceptives include:

- Oral and depot contraceptives
- hormone replacement therapy
- steroids
- NSAIDs/ COX-$_2$ inhibitors
- nasal decongestants and other cold remedies
- appetite suppressants
- amphetamines
- monoamine oxidase inhibitors
- analgesics
- ergotamine
- cyclosporin
- tacrolimus
- carbenoxolone and licorice
- bupropion
- sibutramine

Alcohol intake[1]

Alcohol has a direct pressor effect that is dose-related. An assessment of the average daily number of standard drinks is important - more than two standard drinks (20 g alcohol) a day is significant.

PHYSICAL EXAMINATION

The approach to the physical examination is to examine possible target organ damage and possible causes of secondary hypertension. The main features to consider in the physical examination are illustrated in Figure 126.4.


Neck
carotid bruits
JVP
Blood pressure
two or more readings while sitting
measure in both arms
measure standing
Kidneys
kidney artery bruit best heard in epigastrium
? enlarged kidneys
measure BP
in thigh
Waist circumference
Height and weight
measure BMI
Fundi
? haemorrhages
exudates
AV nipping
papilloedema
Facies
plethora
alcohol abuse
Heart
rate and rhythm
apex beat
murmurs
S3 and S4 sounds
Lungs
crackles of LVF
wheezes in bronchospasm
measure PEFR
Abdomen
bruits
aortic dilatation/pulsation
liver
Femoral pulses
? radiofemoral delay
Urine
glucose
protein
casts
red and white cells
Periphery
pulses: absent or diminished
trophic changes

Figure 126.4 Physical examination of patient with hypertension: what to look for

The four grades of hypertensive retinopathy are illustrated in Figure 126.5.


vein
arteriole thickened, narrowed and more tortuous
± silver wiring
Grade 1
arteriovenous nipping
Grade 2
flame-shaped haemorrhages
soft 'cotton wool' exudates
Grade 3
papilloedema
Grade 4

Figure 126.5 The four grades of hypertensive retinopathy

LEG PULSES AND PRESSURE

To assess the remote possibility of coarctation of the aorta in the hypertensive patient, perform at least one observation comparing:

1. The volume and timing of the radial and femoral pulses

2. The BP in the arm and leg
3. Comparison of BP in the two arms

Blood pressure measurement in the leg

- Place the patient prone.
- Use a wide, long cuff at mid-thigh level.
- Position the bladder over the posterior surface and fix it firmly.
- Auscultate over the popliteal artery.

INVESTIGATIONS[2]

Routine tests

- Plasma glucose (preferably fasting)
- Serum total and high-density lipoprotein (HDL) cholesterol; fasting serum triglycerides
- Serum creatinie/eGFR
- Serum uric acid
- Serum potassium and sodium
- Haemoglobin and haematocrit
- Urinalysis (dipstick test and urinary sediment)
- Electrocardiogram

Recommended tests

- Echocardiogram
- Carotid (and femoral) ultrasound
- Postprandial plasma glucose (when fasting value ≥ 6.1 mmol/l or 110 mg/l)
- C-reactive protein (high sensitivity)
- Microalbuminuria (essential test in diabetes)
- Quantitative proteinuria (if dipstick +ve)
- Fundoscopy (in severe hypertension)

Other investigations, such as echo-cardiography, renal imaging studies (especially renal ultrasound), 24 hour urinary catechol-amines, aldosterone and plasma renin, are not routine and should be done only if specifically indicated. A chest X-ray may serve as a baseline against which to measure future changes. However, if a chest X-ray shows the heart is enlarged, it is more likely to represent chamber dilatation than increased ventricular wall thickness.[1] Specific renal studies now favoured include isotope scans, Doppler ultrasound studies of renal arteries and renal arteriography.

TREATMENT

A correct diagnosis is the basis of management. Assuming that the uncommon secondary causes are identified and treated, treatment will focus on essential hypertension.

Benefits of treatment

Trials have shown that lowering BP reduces:

- Cardiovascular and total mortality
- Stroke
- Coronary events

Benefits have been proven:

- in patients with systolic/diastolic hypertension
- in elderly patients with isolated systolic hypertension.

Management principles

- The overall goal is to improve the long-term survival and quality of life.
- Promote an effective physician - patient working relationship.
- Aim to reduce the levels to 140/90 mmHg or less (ideal).
- Undertake a thorough assessment of all cardiovascular risk factors.
- Instruct all patients in the use of non-drug treatment strategies and their potential benefits.
- In patients with mild-to-moderate hypertension and no target organ damage, consider ambulatory or home BP monitoring.
- Drug therapy should be given to those with:
  - sustained high initial readings (e.g. DBP 95 mmHg)
  - target organ damage
  - failed non-drug measures
- Make a careful selection of an antihypertensive drug and an appraisal of the side-effects against the benefits of treatment.
- Avoid drug-related problems such as postural hypotension.
- Avoid excessive lowering of BP-aim for steady and graduated control.
- Aim to counter the problem of patient non-compliance.
- Be aware of factors that may contribute to drug resistance.

PATIENT EDUCATION

Patient education should include appropriate reassurances, clear information and easy-to-follow instructions. It is important to establish patients' understanding of the concept of hypertension and its consequences by quizzing them about their knowledge and feelings.

Correction of patients' misconceptions[1]

Patients are likely to have several mis-conceptions about hypertension which may adversely affect their treatment.

For example, they might believe that:

- hypertension can be cured
- they do not need to continue treatment once their BP is controlled
- they do not have a problem because they do not have symptoms
- they need to take additional pills, or stop treatment in response to symptoms they believe are caused by high or low BP levels
- they need not take prescribed pills if they attend to lifestyle factors such as exercise and diet
- they can gauge their BP by how they feel

Tips for optimal compliance

- Establish a good caring rapport.
- Give patients a card of their history with BP readings.
- Give advice about pill-taking times.
- Set therapeutic goals.
- Establish a recall system.
- Provide patient education material.

On review:

- Ask if any pills were missed by accident.
- Attempt to reduce waiting time to a minimum (e.g. direct a patient to a spare room upon arrival).
- Review all cardiovascular risk factors.
- Enquire about any side-effects.

NON-PHARMACOLOGICAL (LIFESTYLE) MANAGEMENT MEASURES

If the average diastolic BP at the initial visit is 90 - 100 mmHg, and there is no evidence of end organ damage, non-pharmacological therapy is indicated for a 3 month period without use of antihypertensive drugs. Remember to remove, revise or substitute drugs which may be causing hypertension (e.g. NSAIDs, corticosteroids, oral contraceptives, hormone replacement therapy).

Behaviour intervention measures include:

- *Weight reduction* There is considerable evidence that weight loss and gain are linked to a corresponding fall and rise in BP. Hovell has estimated that for every 1 kg of weight lost, BP dropped by 2.5 mmHg systolic and 1.5 mmHg diastolic.[9] The BMI should be calculated for all patients and where required a weight loss program organised to reduce the BMI to between 20 and 25.
- *Reduction of excessive alcohol intake* [1] The direct pressor effect of alcohol is reversible. Drinking more than 20 g of alcohol a day raises BP and makes treatment of established hypertension more difficult. People with hypertension should limit their alcohol intake to one or two standard drinks (10 g) per day. Reduction or withdrawal of regular alcohol intake reduces BP by 5 - 10 mmHg.
- *Reduction of sodium intake* Some individuals seem to be more sensitive to salt restriction. Advise patients to put away the salt shaker and use only a little salt with their food. Reduction of sodium intake to less than 100 mmol sodium per day is advised. Special care should be taken with processed and takeaway foods.
- *Increased exercise* Regular aerobic or isotonic exercise helps to reduce BP.[10] Hypertensive patients beginning an exercise program should do so gradually. Walking is an appropriate exercise. Weights and other forms of isometric exercises should be avoided because they will significantly elevate BP in the hypertensive subject.
- *Reduction of particular stress* If avoiding stress or overwork is difficult, recommend relaxation and/or meditation therapy.
- *Other dietary factors* There is evidence that lactovegetarian diets and magnesium supplementation can reduce BP.[11, 12] A diet high in calcium, and low in fat and caffeine, may also be beneficial. Avoid licorice and licorice-containing substances.
- *Smoking* Smoking causes acute rises in BP but does not appear to cause sustained hypertension. However, the elimination of smoking is important as it is a strong risk factor for cardiovascular disease and continuing to smoke may negate any benefits of antihypertensive therapy.[1]
- *Management of sleep apnoea*

PHARMACOLOGICAL THERAPY

The benefits of drug therapy appear to outweigh any known risks to individuals with a per-sistently raised diastolic pressure > 95 mmHg. Although the ideal antihypertensive drug has yet to be discovered there are many effective antihypertensive drugs available from all the major classes of anti- hypertensive drugs. [13] Deciding which one to use first involves an assessment of the patient's general health, the medication's known side-effects, the simplicity of its administration and its cost. A useful plan is outlined in Figure 126.6.


D
Diuretic
B
β_2-adrenergic antagonist
Congestive heart failure
Coronary heart disease
A
ACE/AIIRA
C
Dihydropyridine Calcium-channel blocker
V & D
V = verapamil
D = diltiazepam
Metabolic risk (diabetes, lipids)

Figure 126.6 Common combinations of the therapeutic drug classes used for first-line therapy of hypertension

Various disorders such as diabetes, asthma, COPD, Raynaud's phenomenon, heart failure, and elevated serum urate and/or lipid levels may restrict the use of some classes of drugs.

When to treat

- Failed genuine non-pharmacological trial
- All cases of SBP 140 - 180 mmHg or DBP 90 - 110 mmHg

BP targets for adults[8]	**mmHg**
≥65 years (unless there is diabetes, kidney disease and /or proteinuria)	< 140/90
< 65 years and/or those with - diabetes - kidney impairment - proteinuria 0.25-1g.day	< 130/85
proteinuria > 1 g	< 125/75

Guidelines[14]

- Start with a single drug.
- A period of 4 - 6 weeks is needed for the effect to become fully apparent.
- If ineffective, consider increasing the drug to its maximum recommended dose, or add an agent from another compatible class, or substitute with a drug from another class.
- Use only one drug from any one class at the same time.
- A summary of first-line therapy options and the uses of the various pharmacological agents is shown in Table 126.7.

Table 126.7 First-line pharmacological options for the management of hypertension[6, 14]

Thiazide diuretic	**Beta-blocker**	**Drug class Calcium-channel antagonist**	**ACE inhibitor**	**Central-acting agent**	**Alpha-blocker**

Typical examples and starting dose (oral therapy)					
chlorthalidone 12.5 - 25 mg daily hydrochlorothiazide 12.5 mg daily indapamide (SR) 1.5 mg or 2.5 mg daily	atenolol 25 - 50 mg daily metoprolol 50 mg daily pindolol 5 mg daily propranolol 40 mg daily	amlodipine 2.5 mg daily diltiazem CD 180 mg daily felodipine SR 2.5 mg daily lercanidipine 10 mg daily nifedipine CR 30 mg daily verapamil SR 120 - 180 mg daily	captopril 6.25 mg bd enalapril 5 mg daily lisinopril 5 mg daily perindopril 2 mg daily ramipril 2.5 mg daily (others p. 1293) AIIRA irbesartan 150 mg daily losartan 50 mg daily (others p. 1294)	clonidine 50 μg, bd methyldopa 125 mg bd	prazosin 0.5 mg nocte terazosin 0.5 mg nocte *Note*: Labetalol (100 mg bd) is a combined α- and β-blocker
Recommended in					
Heart failure (mild) Older patients	Anxious patient Young patients Angina Postmyocardial infarction Migraine	Asthma Angina PVD Raynaud's phenomenon	Heart failure PVD Diabetes Raynaud's	Asthma Pregnancy	Asthma PVD Heart failure LUTS prostatism)
Contraindications					
Type 2 diabetics Hyperuricaemia Kidney failure	Asthma COPD History of wheeze Heart failure Heart block (2nd and 3rd degree) PVD Brittle Type I diabetes	Heart block 2nd and 3rd degree Heart failure (verapamil, diltiazem)	Bilateral kidney artery stenosis Pregnancy Hyperkalaemia	Liver disease (methyldopa)	Heart failure (mechanical obstruction) Orthostatic hypertension
Precautions					
Hypokalaemia Thiazides + ACE inhibitors Kidney failure	Avoid abrupt cessation with angina Use with verapamil Use with NSAIDs Use in smokers	With β-blockers and digoxin CCF	Chronic kidney disease Avoid K-sparing diuretics and NSAIDs	Depression	Elderly patients
Important side-effects					
Rashes Sexual dysfunction Weakness Blood dyscrasias Muscular	Fatigue Insomnia Vivid dreams Bronchospasm	Headache Flushing Ankle oedema Palpitations Dizziness Nausea	Cough Weakness Rash Dysgeusia	Sedation Dry mouth Bowel disturbances Fatigue	First dose syncope Orthostatic hypotension

cramps Hypokalaemia Hyponatraemia Hyperuricaemia Hyperglycaemia Lipid metabolism effect	Cold extremities Sexual dysfunction Lipid metabolism effect	Constipation (verapamil) Nocturia, urinary frequency	(taste) Hyperkalaemia First dose hypotension Angioedema	Orthostatic hypotension Sexual dysfunction Haemolytic anaemia (methyldopa)	Weakness Palpitations Sedation Headache

- Measure the BP at the same time each day.
- A good strategy is to get patients to self-measure.

Starting regimens

The traditional method has been to use stepwise therapy until ideal control has been reached, commencing with:

1. thiazide or thiazide-like diuretic (preferred)
    - Adjacent drug classes are useful combinations in that effects on blood pressure are additive and adverse effects are no more likely than with either drug used alone.
    - Verapamil (V) and diltiazem (D) generally should not be used with β2-antagonists.
    - Drug groups that are diametrically located on the diagram may be used together, but may not have fully additive effects.
    - Drugs on the left side should be combined in patients with hypertension and heart failure, and those on the right side are useful in patients with hypertension and coexisting coronary disease.
    - Drug groups on the lower panel lack metabolic effects and may be preferred combinations in the presence of diabetes or lipid abnormalities as well as hypertension.
    - Prazosin and other α-antagonists are also often used as monotherapy and may be combined with any of the above drug groups.
    - Small doses of centrally acting anti-adrenergic drugs (e.g. methyldopa, clonidine) can probably also be used with any of the other agents although data on their use in combination are scarce with the newer drug groups. Adapted from Management of Hypertension: A Consensus Statement, Med J Aust, 1994, 160: Supplement 21 March. © Copyright 1994. The Medical Journal of Australia. Reproduced with permission.
2. combination of diuretic and β-blocker or a vasodilator (ACE inhibitor or AIIR antagonist)

However, the newer classes of drugs can be used as first-line agents.

The following are useful combinations:[1, 4]

Initial agent	*Additional drugs*
diuretic	β-blocker or ACE inhibitor, or AIIRA
β-blocker	Diuretic Calcium antagonist (except verapamil and diltiazem)
β-blocker	Diuretic β-blocker
ACE inhibito AIIRA	Diuretic Calcium-channel antagonist
Calcium-channel antagonist Diuretic	β-blocker ACE inhibitor Central-acting agent ACE inhibitor

Relatively ineffective combinations[15]

- Diuretic and calcium antagonist
- β-blockers and ACE inhibitors

Undesirable combinations[15]

- More than one drug from a particular pharmacological group: beta-blockers and verapamil (heart block, heart failure); potassium-sparing diuretics and ACE inhibitors (hyperkalaemia).

Diuretics[4,8]

- Thiazides are good first-line therapy for hypertension.
- Hypokalaemia can be corrected with potassium-sparing diuretics or changing to another first-line drug.
- Loop diuretics are less potent as antihypertensive agents but are indicated where there is concomitant cardiac or renal failure and in resistant hypertension.
- Thiazides are less effective where there is renal impairment.
- Thiazides may precipitate acute gout.
- NSAIDs may antagonise the antihypertensive and natriuretic effectiveness of diuretics.
- A diet high in potassium and magnesium should accompany diuretic therapy (e.g. lentils, nuts, high fibre).
- Avoid use if significant dyslipidaemia.
- Indapamide has different properties to the thiazide and loop diuretics and has less effect on serum lipids.

Beta-blockers

- NSAIDs may interfere with the hypotensive effect of β-blockers.
- If BP is not reduced by one β-blocker it is unlikely to be reduced by changing to another.
- Verapamil plus a β-blocker may unmask conduction abnormalities causing heart block.
- In patients with ischaemic heart disease, or susceptibility to it, treatment must not be stopped suddenly - this can precipitate angina at rest.

Calcium-channel antagonists

- These drugs reduce BP by vasodilatation.
- The properties of individual drugs vary, especially their effects on cardiac function.
- The dihydropyridine compounds (nifedipine and felodipine) tend to produce more vasodilatation and thus related side-effects.
- Unlike verapamil or diltiazem (which slow the heart), dihydropyridine drugs can be used safely with a β-blocker.
- Verapamil is contraindicated in second and third degree heart block.
- Verapamil and diltiazem should be used with caution in patients with heart failure.
- These drugs are efficacious with ACE inhibitors, β-blockers, prazosin and methyldopa.

ACE inhibitors

Angiotensin-converting enzyme is responsible for converting angiotensin I to angiotensin II (a potent vasoconstrictor and stimulator of aldosterone secretion) and for the breakdown of bradykinin (a vasodilator). The available ACE inhibitors are captopril, enalapril, fosinopril, lisinopril, perindopril, quinapril, ramipril and trandolapril. ACE inhibitors are effective in the elderly; improve survival and performance status in cardiac failure; are protective of renal function in diabetes; and cardioprotective in postmyocardial infarction. For patients with normal renal function, the dose should not exceed 150 mg daily for captopril, 40 mg daily for enalapril or lisinopril, 10 mg daily for ramipril, and 8 mg daily for perindopril.

Disturbance in taste is usually transitory and may resolve with continued treatment. Cough, which occurs in about 15% of patients, may disappear with time or a reduction in dose but it often persists and requires a change in drug in some patients. Angioedema, a potentially lifethreatening condition, may occur in 0.1 - 0.2% of subjects. Like cough, it is a class effect and will mitigate against use of any ACE inhibitor in patients with such an adverse effect. The Heart Outcomes Prevention Evaluation (HOPE) trial demonstrated that the ACE inhibitor ramipril reduces the number of cardiovascular deaths, non-fatal myocardial infarctions, non-fatal strokes and instances of new onset heart failure in a high-risk population. The data also indicated that all diabetic patients with microalbuminuria or pre-existing vascular disease will benefit from ACE I treatment, even if normotensive.[14, 16]

Angiotension II receptor antagonists

These agents competitively block the binding of angiotension II to type I angiotension receptors and block the effects of angiotension more selectively than the ACE inhibitors. This reduces angiotension-induced vasoconstriction, sodium reabsorption and aldosterone release. They have a similar adverse renal profile and other adverse effects to the ACE inhibitors but cough does not appear to be a significant adverse effect. This group is used for mild-to-moderate hyper-tension alone or with other antihypertensive agents. The available AIIRAs are candesartan, eprosartan, irbesartan, losartan and telmisartan. They are useful alternatives for hypertensive patients who discontinue an ACE inhibitor because of cough and may be used in combination with thiazide diuretics.

Prazosin

A specific problem is the ‘first dose phenomenon’; this involves acute syncope about 90 minutes after the first dose, hence treatment is best initiated at bedtime. Prazosin potentiates β-blockers and works best if used with them. It is a useful first-line therapy in patients who are unsuitable for diuretic or β-blocker therapy (e.g. those with diabetes, asthma or hyperlipidaemia).

Vascular smooth muscle relaxants

(other than calcium-channel antagonists)

These include hydralazine, minoxidil and diazoxide, which are not used for first-line therapy but for refractory hypertensive states and hypertensive emergencies.

MILD HYPERTENSION[1]

Mild hypertension in adults is defined as a diastolic pressure (phase 5) persistently between 90 and 99 mmHg, without target organ damage.[14] This group includes those with ‘white coat’ hypertension.

Mildly hypertensive people have almost twice the risk of vascular disease compared with normotensive people, and evidence shows that morbidity and mortality rise with an increase in blood pressure.[1] The long-term risk in patients with ‘white coat’ hypertension has yet to be defined.

Patients treated appropriately have fewer strokes and pressure-related cardiovascular complications than those not treated. Drug treatment has potential side-effects which cannot be ignored; sometimes the risks from this form of treatment could exceed those if the patient remains untreated.[17] Therefore, the initial approach to the management of mild hypertension should be based on non-drug measures and should focus on behaviour change. In this way, overt side-effects are avoided and risk prevention measures enhanced.

Often the success of this approach is improved by using people with particular skills, such as dietitians. If after 6 months or more these methods have not succeeded, then drug therapy may be necessary.[17]

Practical guidelines for patients with persistent diastolic blood pressure between 90 and 99 mmHg are shown in Figure 126.7. This includes an assessment of 5 year absolute.


SBP > 140 mm Hg or
DBP > 90 mm Hg
on several occasions
Start lifestyle changes.
Assess 5-year absolute
cardiovascular risk.
low risk (10%)
moderate risk (10–15%)
High risk (> 15%)
or associated clinical conditions
or target organ disease
or high-risk ethnic groups
Maintain lifestyle changes for
6–12 months.
Consider drug treatment if BP
remains > 150/95 mm Hg.
Maintain lifestyle changes for
3–6 months.
Consider drug treatment if BP
remains > 140/90 mm Hg.
Maintain lifestyle changes,
begin drug treatment.

Figure 126.7 Decision tree for managing hypertension. As recommended by the National Heart Foundation

MODERATE HYPERTENSION[1]

If diastolic pressure is 100 - 109 mmHg then a more aggressive approach is necessary, particularly for patients with target organ damage. Appropriate non-pharmacological measures should be tried in all subjects, especially where success is possible (e.g. obesity, high alcohol intake). This will fail in the majority, who will require drug therapy. If there is poor response to initial therapy a second drug may be prescribed after a short time. The interval between the changes in treatment may also be reduced. Combination therapy may be more effective than maximum doses of any single agent.

SEVERE HYPERTENSION[4]

The BP of those with severe hypertension may be life-threatening. Patients with an average diastolic pressure of more than 110 mmHg should be checked immediately for hy-pertensive complications, particularly marked fundoscopic changes, proteinuria or cardiac failure. They are more likely to have an underlying cause (e.g. renovascular disease). Hospitalisation may be necessary for these patients.

In such cases the opinion of a specialist is important because of the likely risk of serious illness or death. A dihydropyridine calcium-channel blocker (e.g. nifedipine, amlodipine, felodipine) with early addition of a β-blocker or ACE inhibitor is suitable for urgent lowering of BP.

HYPERTENSIVE EMERGENCIES

A hypertensive emergency occurs when high BP causes the presenting cardiovascular problem. Typical presentations of hypertensive emer-gencies (which are rare) include hypertensive encephalopathy, acute stroke, heart failure, dissecting aortic aneurysm and eclampsia.

In all such cases referral to a specialist is essential and the patient should be hospitalised immediately for monitoring and treatment. Treatment must be individualised, mindful of the nature of the underlying problem and associated disorders.

Such treatment is the same as for severe hypertension. Otherwise, sodium nitroprusside IV in an intensive care setting is the optimal treatment.

Magnesium sulphate reduces the risk of eclampsia and maternal death in women with pre-eclampsia.

ISOLATED SYSTOLIC HYPERTENSION[1]

Isolated systolic hypertension is most frequently seen in elderly people.

Definition

SBP ≥ 140 mmHg with a DBP < 90 mmHg

Patients with isolated systolic hypertension should be treated in the same way as those with classic systolic/diastolic hypertension. Evidence that reducing isolated systolic BP decreases the mortality and morbidity risk has been demon-strated by the SHEP (systolic hypertension in the elderly) study.[18]

Non-pharmacological therapy should be commenced as is relevant to the patient. If drugs are used the systolic BP should be cautiously lowered to between 140 and 160 mmHg. The drugs of choice are diuretics, calcium-channel blocking agents and ACE inhibitors.

REFRACTORY HYPERTENSION

Refractory hypertension exists where control has not been achieved despite reasonable treatment for 3 - 4 months. A review of a possible secondary cause is appropriate. Consider also spurious factor, e.g. instrument factor s such as small cuff, isolated office hypertension.

Checklist of possible reasons[1]

- Drug-related causes: doses too low; inappropriate combinations; effects of other drugs (e.g. antidepressants, adrenal steroids, NSAIDs, sympathomimetics, nasal decongestants, ergotamine, oral contraceptives, psychotropics, cocaine
- Poor compliance with therapy
- Renovascular hypertension
- Nicotine, licorice, caffeine (strong coffee)
- Obesity
- Excessive alcohol intake and other causes of volum overload
- Excessive salt intake and other causes of volume overload
- Renal insufficiency and other undiagnosed causes of secondary hypertension
- Illicit substances (e.g. amphetamines, cocaine, anabolic steroids)
- Sleep apnoea

When adequate control is not possible and the cause is not obvious, the patient should be referred to a specialist. Measurement outside the clinic may help in the assessment of such people, as may 24 hour ambulatory monitoring.

HYPERTENSION IN CHILDREN AND ADOLESCENTS

The recording of BP should be part of the normal examination in children and used in their continuing care. BP should be measured in all children who are unwell. BP is less frequently measured in children for a number of reasons, such as the unavailability of an appropriately sized cuff or difficulty in measuring BP in the infant or toddler.

The children of parents with hypertension should be closely watched. Those at risk of secondary hypertension (e.g. renal or cardio-vascular disease, urological abnormalities and diabetes mellitus) should have routine measurements. Those children with visual changes, headache or recurrent abdominal pain or seizures, and those on drugs such as corticosteroids or the pill, should have their BP checked regularly.

Although secondary causes of hypertension are more common in children than in adults, young people are still more prone to developing essential rather than secondary hypertension. Renal parenchymal disease and renal artery stenosis are the major secondary causes.

The upper limits of normal BP for children in different age groups are:[1]

Age (in years)	*Arterial pressure (mmHg)*
14 - 18	135/90
10 - 13	125/85
6 - 9	120/80
5 or less	110/75

The proper cuff size is very important to avoid inaccurate readings and a larger rather than a smaller cuff is recommended. The width of the bladder should cover 75% of the upper arm. In infants and toddlers use of an electronic unit may be necessary. Although cessation of sound (phase 5) is the better reflection of true diastolic pressure, there is often no disappearance of sound in children and so estimation of the point of muffling has to be recorded.

Diagnostic evaluation and drug treatment for children are similar to those for adults. When a child is obese, reduction in weight may adequately lower BP. ACE inhibitors or calcium-channel blocking agents are preferable in the young hypertensive, with diuretics a second agent. ACE inhibitors should be avoided in postpubertal girls.

HYPERTENSION IN THE ELDERLY

BP shows a gradual increasing linear re-lationship with age.

Guidelines for treatment

- Isolated systolic hypertension is worth treating.[18]
- Older patients may respond to non-pharmacological treatment.
- Reducing dietary sodium is more beneficial than with younger patients.
- If drug treatment is necessary, commence with half the normal recommended adult dosage - 'start low and go slow'.
- Patients over 70 years and in good health should be treated the same as younger patients.
- A gradual reduction in BP is recommended.
- Drug reactions are a limiting factor.
- Drug interactions are also a problem: these include NSAIDs, antiparkinson drugs and phenothiazines.

Specific treatment

- *First-line choice*: indapamide (preferred) or thiazide diuretic (low dose)[1]; check electrolytes in 2 - 4 weeks: if hypokalaemia develops add a K-sparing diuretic rather than K supplements. Use a combination thiazide and K-sparing diuretic. Diuretics may aggravate bladder difficulties (e.g. incontinence).
- *Second line choice*: β-blockers (low dose) where diuretic cannot be prescribed or if angina.

Other effective drugs (especially for isolated systolic hypertension):

- ACE (or AIIRA) inhibitors (especially with heart failure)
- calcium-channel antagonists

Both these groups are generally well tolerated but constipation may be a problem with verapamil.

Renal function and electrolytes should be monitored when ACE inhibitors are started.

SPECIAL MANAGEMENT PROBLEMS

These conditions are summarised in Table 126.8.

Table 126.8 Choice of drugs in patients with coexisting conditions

	Diuretic	ACE inhibitors AIIRAs	Calcium-channel antagonists	β-blockers
Asthma/COPD				×
Bowel disease/constipation	×		×	
Bradycardia/heart block			Care	×
Cardiac failure	*	*	Care	×
Depression				×
Diabetes	×	*		×
Dyslipidaemia	×			×
Hyperuricaemia/gout	×	*	*	×
Impotence	×			×
Ischaemic heart disease			*	*
Peripheral vascular disease			*	×
Pregnancy	×	×		
Raynaud's phenomenon			*	×
Kidney artery stenosis		×	*	*
Kidney failure		Care		
Tachycardia (resting)	×		Care	*

*drug/s of choice. Calcium-channel antagonists need to be selected with care - some are suitable, others not.

Diabetes mellitus

Factors contributing to hypertension may be the same in type 2 diabetes and non-diabetic hypertensive patients. In both type 1 and type 2 diabetes, nephropathy can be a significant contributor.

The monitoring of patients for early signs of nephropathy with measurements of microalbumin excretion is helpful. Micro-albuminuria can also be detected in non-diabetic hypertensives where it appears to be a marker of cardiovascular disease. Diabetic autonomic neuropathy can cause orthostatic hypotension. Diabetics with persistent or sustained diastolic BP > 85 mmHg and proteinuria need treatment. The threshold for treatment of hypertension in the diabetic is lower than in the non-diabetic. Control of hypertension is an important factor in limiting progression of diabetic renal disease.

Treatment

- Use basic non-pharmacological treatments, especially weight reduction, if applicable.
- ACE inhibitors or AII blockers and calcium-channel blockers are useful first-line drugs because they do not affect insulin and diabetes control.and give renoprotective benefits.
- Other suitable drugs are prazosin, hydralazine and methyldopa.
- Diuretics such as indapamide added to an ACE inhibitor are effective but caution is required because they can aggravate glucose intolerance.

- Proteinuria and renal function need to be monitored.

Pregnancy

Hypertension in pregnancy can be either pre-eclampsia (pregnancy-induced hypertension) or essential hypertension. BP levels normally drop during the second trimester and rise in the third. A diastolic BP > 80 mmHg in late pregnancy is considered unacceptable. Preferred drugs to use are methyldopa, labetalol, and β-blockers. Diuretics and ACE inhibitors should not be used. Hypertensive pregnant women should be supervised in association with a specialist unit.

Surgical patients

Patients whose BP is under control before surgery should continue the same treatment. If oral medication is affected by the surgery, parenteral treatment may be needed to avoid rebound hypertension. This is a particular problem with clonidine and possibly methyldopa. Withdrawal of other drugs, such as β-blockers, may have adverse consequences.

Kidney disease

Kidney function is not adversely affected by the treatment of severe or malignant hypertension. Use a loop diuretic (e.g. frusemide) initially. β-blockers, calcium-channel antagonists, prazosin and methyldopa can be used, while caution is needed with ACE inhibitors, particularly if there is underlying renovascular disease. Kidney protection in diabetes requires strict BP control.

Heart failure

First-line treatment for associated hypertension includes ACE inhibitors and diuretics. Other suitable drugs are a hydralazine - nitrate combination and methyldopa. Calcium-channel antagonists should be used with care and verapamil and β-blockers should be avoided.

Ischaemic heart disease

Recommended drugs are β-blockers and calcium antagonists.[14] The non-dihydropyridine agents should be used with care with a β-blocker but the dihydropyridine agents are quite safe.

Obstructive pulmonary disease

Apart from β-blockers, all other routine antihypertensives can be used.

Impotence

It is prudent to avoid antihypertensives that are possibly associated with impotence (e.g. thiazide diuretics, methyldopa, resperine and β-blockers). Suitable agents to use are ACE inhibitors and calcium-channel blockers.

CAN HYPERTENSION BE OVERTREATED?[19]

Yes. Excessive BP reduction, particularly if acute, can seriously compromise perfusion in vital organs, especially if blood flow is already impaired by vascular disease. Careful monitoring of the patient, including standing BP measurement, is important.

One should avoid excessive BP reduction in the setting of acute stroke, where there is a tight carotid artery stenosis (particularly if symptomatic) and in the elderly subject (especially if there is postural hypotension).

It has been suggested that lowering the diastolic BP < 85 mmHg in particular subgroups (e.g. those with ischaemic heart disease) may raise the cardiovascular risk above that associated with a lesser reduction in

BP.[20] However, the relationship between cardio-vascular risk and blood pressure is a continuous one. Moreover, the SHEP study did not show any adverse effects of lowering diastolic BP in patients with pre-existing coronary heart disease. The safety of lowering diastolic BP to levels below 85 mmHg is being more formally addressed in the Hypertension Optimal Treatment (HOT) study.[21]

STEP-DOWN TREATMENT OF MILD HYPERTENSION[7]

This is an important concept that recognises that drug treatment need not necessarily be lifelong. If BP has been well controlled for several months to years it is often worth reducing the dosage or the number of drugs.

A ‘drug holiday’ (cessation of treatment) can be hazardous, however, because satisfactory control may be temporary and hypertension will re-emerge. Careful monitoring under such circumstances is mandatory.

WHEN TO REFER[4]

- Refractory hypertension - adequate control not possible and cause not obvious
- Suspected ‘white coat’ hypertension - for ambulatory BP monitoring
- Severe hypertension - diastolic BP > 115 mmHg
- Hypertensive emergency
- If there is evidence of ongoing target organ impairment
- If there is significant kidney impairment eGFR < 60 mL/mm
- If a treatable cause of secondary hypertension is found

Practice tips

- Hypertension should not be diagnosed on a single reading.
- At least two follow-up measurements with average systolic pressure > 140 mmHg or diastolic pressures > 90 mmHg are required for the diagnosis.
- Beware of using β-blockers in a patient with a history of wheezing.
- Add only one agent at a time and wait about 4 weeks between dosage adjustments.
- Excessive intake of alcohol can cause hypertension and hypertension refractory to treatment.
- If hypertension fails to respond to therapy, an underlying renal or adrenal lesion may have been missed.
- The low-pitched bruits of renal artery stenosis are best heard by placing the diaphragm of the stethoscope firmly in the epigastric area.
- Older patients may respond better to diuretics, calcium-channel antagonists and ACE inhibitors.
- Younger patients may respond better to β-blockers or ACE inhibitors.

REFERENCES

1. Sandler G. High blood pressure. In: Common Medical Problems. London: Adis Press, 1984: 61 - 106.
2. 1999 WHO - ISH guidelines for the management of hypertension. J Hypertension, 1999;17:151 - 83.
3. Practice guidelines for primary care physicians: 2003 ESH/ESC Hypertension guidelines; J Hypertension 2003; 21 (10): 1779 - 86
4. Stokes G. Essential hypertension. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 252 - 4.
5. Fraser A. Measurement of blood pressure. Aust Fam Physician, 1989; 18: 355 - 9.
6. ABC of Hypertension. London: British Medical Association, 1989: 1 - 50.
7. Bates B. A Guide to Physical Examination and History Taking (5th edn). Philadelphia: Lippincott, 1991: 284.

8. National Heart Foundation of Australia. Canberra. Hypertension Management Guide for Doctors. National Heart Foundation of Australia.
9. Hovell MF. The experimental evidence for weight loss treatment of essential hypertension. A critical review. Am J Public Health, 1982; April 72(4): 359 - 68.
10. Blair SN, Goodyear NN, Gibbons LW, Cooper KH. Physical fitness and incidence of hypertension in healthy normotensive men and women. JAMA, 1984; 252(4): 487 - 90.
11. Rouse IL, Beilin LJ. Vegetarian diet and blood pressure. Editorial review. J Hypertension, 1984; 2: 231 - 40.
12. Kestlefoot H. Urinary cations and blood pressure - population studies. Ann Clin Research, 1984; 16 Supp. (43): 72 - 80.
13. Jennings G, Sudhir K. Initial therapy of primary hypertension. Med J Aust, 1990; 152: 198 - 202.
14. Smith A (Chair). Therapeutic Guidelines: Cardio-vascular (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2003: 63 - 79.
15. Hypertension Guideline Committee, 1991 report. Hypertension: Diagnosis, Treatment and Maintenance. Adelaide: Research Unit RACGP (South Australian Faculty), 1991.
16. Yusef S et al. Effects of an angiotension-converting inhibitor, ramipril, on cardiovascular events in high risk patients. The Heart Outcomes Prevention Evaluation Investigations. N Engl J Med, 2000; 342: 145.
17. Guidelines for the treatment of mild hypertension. Memorandum from a WHO/ISH meeting. Endorsed by Participants at the Fourth Mild Hypertension Conference. Bulletin of the World Health Organization, 1986; 64(1): 31 - 5.
18. SHEP Co-operative research group. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension. JAMA, 1991; 265: 3255 - 64.
19. Vandongen R. Drug treatment of hypertension. Aust Fam Physician, 1989; 18: 345 - 8.
20. Cruickshank JM, Thorp JM, Zacharias FJ. Benefits and potential harm of lowering high blood pressure. Lancet, 1987; 1: 581 - 4.
21. The HOT study group. The Hypertension Optimal Treatment Study. Blood Pressure, 1993; 2: 6.

CHAPTER 127. DYSLIPIDAEMIA

The landmark Scandinavian Simvastatin Survival Study (4S) published in 1994, may well be remembered as the study that finally put to rest many of the apprehensions and misconceptions regarding lipid-lowering therapy.

DUFFY AND MEREDITH 1996[1]

DYSLIPIDAEMIA

Dyslipidaemia is the presence of an abnormal lipid/lipoprotein profile in the serum and can be classified as:

- predominant hyperglyceridaemia
- predominant hypercholesterolaemia
- mixed pattern with elevation of both cholesterol and triglyceride (TG)

Modern epidemiological studies have established the facts that elevated plasma cholesterol causes pathological changes in the arterial wall leading to CAD, and that lipid-lowering therapy results in reduction of coronary and cerebrovascular events with improved survival.

These studies, which can be summarised by their acronyms - 4S,[2] PLACI,[3] PLACII,[4] ACAPS,[5] KAPS,[6] REGRESS[7] and WOSCOPS[8] - all reinforce the benefits of lipid-lowering therapy for dyslipidaemia and the primary prevention of coronary heart disease.

A recent systematic review showed that statistic and n-3 fatty acids are the most favourable lipid- lowering interventions with reduced risks of overall and cardiac mortality. [9]

ESTABLISHED FACTS[10, 11, 12]

- Major risk factors for CAD include:
  - increased LDL cholesterol + reduced HDL cholesterol
  - ratio LDLC/HDLC > 4
- Risk increases with increasing cholesterol levels (90% if > 7.8 mmol/L)
- TG Levels > 10 mmol/L increases risk of pancreatitis
- Management should be correlated with risk factors
- 10% reduction of total cholesterol gives 20% reduction in CAD after 3 years

INVESTIGATIONS[10]

The following fasting tests are recommended in all adult patients 18 years and over:

- serum triglyceride level
- serum cholesterol level and HDLC and LDLC levels if cholesterol ≥ 5.5 mmol/L
- TFTs if overweight elderly female

Confirm an initial high result with a second test at 6 - 8 weeks. Patients requiring treatment are summarised in Table 127.1. Testing should occur at least every 5 years.

Table 127.1 Patients requiring treatment (National Heart Foundation and PBS guidelines)

Risk category	Initiate drug therapy if lipid level (mmol/L) is	Target levels
Patients with existing coronary heart disease	cholesterol > 4 mmol/L	cholesterol < 4

		mmol/L
Other patients at high risk with one or more of the following: - diabetes mellitus - familial hypercholesterolaemia - family history of coronary heart disease (first degree relative less than 60 years of age) - hypertension - peripheral vascular disease	cholesterol > 6.5 mmol/L or cholesterol > 5.5 mmol/L and HDL < 1 mmol/L	LDL cholesterol < 2 mmol/L HDL cholesterol > 1.0 mmol/L
Patients with HDL < 1 mmol/L	cholesterol > 6.5 mmol/L	Triglycerides < 2.0 mmol/L
Patients not eligible under the above: - men 35 - 75 years - postmenopausal women up to 75 years	cholesterol > 7.5 mmol/L or triglyceride > 4 mmol/L	
Other patients not included in the above	cholesterol > 9 mmol/L or triglyceride > 8 mmol/L	

Recommended treatment goals[12]

- Total cholesterol < 4.0 mmol/L
- LDLC < 2.5 mmol/L *
- HDLC > 1.0 mmol/L
- TG < 1.5 mmol/L

Treat all risk factors.

* < 2.0 in high rish patients

Non-pharmacological measures

- Dietary measures
  - keep to ideal weight
  - reduce fat intake, especially dairy products and meat
  - avoid ‘fast’ foods and deep fried foods
  - replace saturated fats with mono- or polyunsaturated fats
  - limit high-cholesterol foods (e.g. egg yolk, offal, fish roe)
  - use approved cooking methods, eg. Steaming, gulling
  - always trim fat off meat, remove skin from chicken
  - avoid biscuits and cakes between meals
  - eat fish at least twice a week
  - high-fibre diet, especially fruit and vegetables
  - increase complex carbohydrates
  - alcohol intake 0 - 2 standard drinks/day
  - drink more water
- Regular exercise
- Cessation of smoking
- Co-operation of family is essential
- Exclude secondary causes (e.g. kidney disease, type 2 diabetes, hypothyroidism, obesity, alcohol excess - especially → TG), specific diuretics

Checkpoints

- Diet therapy effective (TG ↓, LDLC ↓) within 6 - 8 weeks
- Continue at least 6 months before considering drug therapy in all but the highest-risk category

Pharmacological measures

The choice of the lipid-lowering agent depends on the pattern of the lipid disorder.[11, 12] See Table 127.2. Use the following agents in addition to diet.

Table 127.2 Lipid-lowering drugs

Drug	Dose (average)	Usage	Adverse effects	Safety monitoring
The statins	**Night dose**			
Atorvastatin	10-80 mg	↑ cholesterol	muscle pains; raised liver enzymes	liver enzymes: creatine kinase and ALT
Pravastatin	10-80 mg			
Simvastatin	10-80 mg			
Fluvastatin	20-80 mg			
Rosuvastatin	10-40 mg			
Bile acid binding resins				
Cholestyramine	8 g bd	↑ cholesterol	GIT dysfunction; drug interactions	
Colestipol	10 g bd			
Fibrates				
Gemfibrozil	600 mg bd	→triglycerides mixed hyperlipidaemia	GIT dysfunction; myositis; interaction with statins and warfarin	liver enzymes coagulation
Fenofibrate 160 mg daily				
Other agents				
Ezetimibe	10 mg daily	→cholesterol	arthralgia, myalgia, myositis, liver dysfunction	liver enzymes
Nicotinic acid	100 mg tds to 500 mg tds	→ cholesterol and triglycerides	flushing; raised glucose, urate and liver enzymes	glucose urate liver enzymes
Probucol	500 mg bd	→ cholesterol	GIT dysfunction; arrhythmias	liver enzymes ECG
Fish oils n - 3 fatty acids	2g daily	→ triglycerides	Minimal	bleeding time

Hypercholesterolaemia

Choose one of the following.

First-line agents

1. HMG-CoA reductase inhibitors ('statins')
   - e.g. atorvastatin 10 mg (o) nocte, increase to max. 40 mg/day
     or
     fluvastatin 20 mg (o) nocte, increase to max. 80 mg/day
     or

simvastatin 10 mg (o) nocte, increase to max. 40 mg/day
or
pravastatin 10 mg (o) nocte, increase to max. 40 mg/day

- adverse effects: GIT side-effects, myalgia, abnormal liver function (uncommon)
- monitor: measure LFTs (ALT and CPK) and CK as baseline
- repeat LFTs after 4 - 8 weeks, then every 6 weeks for 6 months

2. Ezetimibe 10 mg daily
3. Combination: ezetimibe + statin, eg. simvastatin
4. Bile acid sequestrating resins
    - e.g. cholestyramine 4 g daily in fruit juice increasing to maximum tolerated dose
    - adverse effects: GIT side-effects (e.g. constipation, offensive wind)
5. Fibrates : consider if above drugs not tolerated

Second-line agents

1. Nicotinic acid
    - nicotinic acid 250 mg (o) with food daily, increase to max. 500 mg tds
    - adverse effects: flushing, gastric irritation, gout
    - minimise side-effects with gradual introduction; take with food and aspirin cover
2. Probucol
    - probucol 500 mg (o) bd
    - problems: slow response, care with hepatic disease
3. Oestrogen[12]
    - oestradiol valerate 2 mg (o) mane ± medroxyprogesterone acetate

This hormone replacement therapy can reduce LDLC and is a physiological intervention in postmenopausal females, especially after hysterectomy. It has limited efficacy.

Resistant LDLC elevation

1. Combination statin + ezetimibe
2. Combined ‘statin’ and resin cholestyramine 4 - 8 g (o) mane plus a ‘statin’ (p. 1302)

Moderate to severe (isolated) TG elevation

gemfibrozil 600 mg (o) bd

or

fenofibrate 145 mg (o) daily

Note: Slow response; monitor LFTs; predisposes to gallstones and myopathy.

Alternatives

nicotinic acid

or

n-3 fish oil concentrate 6 g (o) daily in divided doses to max. 15 g/day

Note: Reduction in alcohol intake is essential.

Mixed hyperlipidaemia (→ TG + → LDLC)

- If TG < 4: a ‘statin’
- If TG > 4: gemfibrozil

Consider combination therapy,

e.g. fish oil + ‘statin’

gemfibrozil + resin

Note: Statin + gemfibrozil increases risk of myopathy and requires specialised supervision.

Special considerations

The decision to commence drug therapy should be based on at least two separate measurements at an accredited laboratory.

Be careful with β-blockers and diuretics affecting lipid levels.

Length of treatment

- Possibly lifelong (up to 75 years)

Follow-up investigations

- Serum lipids
- LFTs (ALT and CPK)
- Possibly CK

Special groups

Children

In general there is little justification for using lipid-regulating drugs in children, especially as the drugs have been shown to reduce heart disease within 2 - 5 years in adults.[11] Initial dietary advice and avoidance of smoking is recommended. Bile acid sequestrating resins are safe to use.

The elderly[11]

The role of lipid therapy remains unclear. Generally, elderly patients with established CHD should receive standard lipid manage-ment unless their general medical status is poor.

Pregnancy[12]

As a general rule the increase in cholesterol level associated with pregnancy subsides after delivery. Systematically absorbed lipid-lowering agents may be unsafe during pregnancy and should be avoided.

Complementary therapy

Claims have been made for the cholesterol lowering properties of policosanol (derived from sugar cane), fish oils, plant sterols, vitamin E, garlic and lecithin. The evidence from RCTs to date indicates a modest benefit from policosanol, fish oils (consuming a fish-based meal at least twice a week is recommended) and plant sterols but there is insufficient evidence to recommend vitamin E, garlic and lecithin.[13]

REFERENCES

1. Duffy SJ, Meredith IJ. Treating mildly elevated lipids. Curr Ther, 1996; 37(4): 49 - 58.

2. Scandinavian Simvastatin Survival Study Group. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). Lancet, 1994; 344: 1383 - 9.
3. Pitt B, Mancini BJ, Ellis SG et al. Pravastatin limitation of atherosclerosis in the coronary arteries (PLACI): reduction in atherosclerosis progression and clinical events. J Am Coll Cardiol, 1995; 26: 1133 - 9.
4. Byington RP, Furberg CD, Crouse JR et al. Pravastatin, lipids and atherosclerosis in the carotid arteries (PLACII). Am J Cardiol, 1995; 76: 54C - 59C.
5. Furberg CD, Adams HP, Applegate WB et al. for the Asymptomatic Carotid Plaque Study (ACAPS) Research Group. Effect of lovastatin and warfarin on early carotid atherosclerosis and cardio-vascular events. Circulation, 1994; 90: 1679 - 87.
6. Salonen R, Nyyssonen K, Porkkala-Sarataho E, Salonen JT. The Kuopio Atherosclerosis Prevention Study (KAPS): effect of pravastatin treatment on lipids, oxidation resistance of lipoproteins, and atherosclerosis progression. Am J Cardiol, 1995; 76: 34C - 39C.
7. Jukema JW, Bruschke AVG, van Boven A et al. Effects of lipid lowering by pravastatin on progression and regression of coronary artery disease in symptomatic men with normal to moderately elevated serum cholesterol levels. The Regression Growth Evaluation Statin Study (REGRESS). Circulation, 1995; 91: 2528 - 40.
8. Shepherd J, Cobbe SM, Ford I et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolaemia. N Engl J Med, 1995; 333:1301 - 7.
9. Studer M, Briel M et al. Effect of different antilipidemic agents and diets on mortality. Arch Intern Med, 2005; 165: 725 - 730.
10. Department Human Services and Health. Schedule Benefits. Canberra: Commonwealth of Australia, November 1996: 25 - 8.
11. National Heart Foundation (Australia). Guide to plasma lipids for doctors. Canberra: NHF, 1996.
12. Smith A (Chair). Therapeutic Guidelines: Cardiovascular (Version 4). Melbourne: Therapeutic Guidelines Ltd, 2003: 51 - 3.
13. Managing dyslipidaemia. NPS News, 2002; 20: 5 - 6.

CHAPTER 128. DIABETES MELLITUS: MANAGEMENT

Man may be the captain of his fate, but he is also the victim of his blood sugar.

WILFRED G. OAKLEY (1905 - 68)

DIABETES MELLITUS: MANAGEMENT

The main objectives for the GP in the optimal management of the diabetic patient, in order to prevent the development of cardiovascular disease and other complications, are:[1]

1. to achieve strict glycaemic control as measured by (most importantly) glycosylated haemoglobin (HbA_{1c}) and by blood glucose
2. to achieve blood pressure control (≤ 130/80 mmHg, supine)
3. to achieve control of blood cholesterol level

Note: Refer to the tables of estimation of cardiovascular risk (Table 128.6 and Table 128.7) that follow this chapter.

Table 128.6 Estimation of cardiovascular risk



How to use the Charts

- Identify the chart relating to the person's sex, diabetic status, smoking history and age.
- Within the chart choose the cell nearest to the person's age, blood pressure (BP) and TC:HDL ratio. When the systolic and diastolic values fall in different risk levels, the higher category applies.
- For Example, the lower left cell contains all non-smokers without diabetes who are less than 45 years and have a TC:HDL ratio of less than 4.5 and a BP of less than 130/80 mm Hg. People who fall exactly on a threshold between cells are placed in the cell indicating higher risk.

Certain groups may have CVD risk underestimated using these charts, see Table 128.7 for recommended adjustments.

Source: Reproduced with permission: New Zealand Guidelines Group. New Zealand Cardiovascular Guidelines

Handbook: Developed for primary care practitioners. Wellington: June 2005. www.nzgg.org.nz.

Table 128.7 Estimation of cardiovascular risk



Risk level: 5-year CVD risk (fatal and non-fatal)	Benefits: NNT for 5 years to prevent one event (CVD events prevented per 100 people treated for 5 years)		
	1 intervention (25% risk reduction)	2 interventions (45% risk reduction)	3 interventions (55% risk reduction)
30%	13 (7.5 per 100)	7 (14 per 100)	6 (16 per 100)
20%	20 (5 per 100)	11 (9 per 100)	9 (11 per 100)
15%	27 (4 per 100)	15 (7 per 100)	12 (8 per 100)
10%	40 (2.5 per 100)	22 (4.5 per 100)	18 (5.5 per 100)
5%	80 (1.25 per 100)	44 (2.25 per 100)	36 (3 per 100)

Based on the conservative estimate that each intervention: aspirin, blood pressure treatment (lowering systolic blood pressure by 10 mm Hg) or lipid modification (lowering LDL-C by 20%) reduces CV risk by about 25% over 5 years.

Note: Cardiovasular events are defined as myocardial infarction, new angina, ischaemic stroke, transient ischaemic attack (TIA), peripheral vascular disease, congestive heart failure and cardiovascular-related death.

NNT = Number needed to treat

Source: Reproduced with permission: New Zealand Guidelines Group. New Zealand Cardiovascular Guidelines Handbook: Developed for primary care practitioners. Wellington: June 2005. www.nzgg.org.nz.

MANAGEMENT PRINCIPLES

- Provide detailed and comprehensive patient education, support and reassurance.
- Achieve control of presenting symptoms.
- Achieve blood pressure control (≤ 130/80 mmHg supine).
- Emphasise the importance of the diet: good nutrition, adequate complex carbohydrates, restricted fats and sugars.
- Promptly diagnose and treat urinary tract infection.
- Treat and prevent life-threatening complications of ketoacidosis or hyperosmolar coma.
- Treat and prevent hypoglycaemia in those taking insulin and oral hypoglycaemic agents.
- Organise self-testing techniques, preferably blood glucose monitoring.

- Detect and treat complications of diabetes - neuropathy, nephropathy, retinopathy, vascular disease.

Metabolic Syndrome[2]

Beware of the deadly metabolic syndrome (syndrome X or insulin resistance syndrom).

- upper truncal obesity (↑ waist circumference) *plus* any 2 or more of the following
- ↑ triglycerides > 1.7 mmol/L
- ↓ HDL cholesterol < 1.030Ú : < 1.2 mmol/L 0♀
- fasting glucose ≥ 6.1 mmol
- BP ≥ 130/85

This syndrome is associated with increased risk for the development of Type 2 diabetes and atheroscleroic vascular disease.

Monitoring techniques

- Blood glucose estimation (fasting and post-prandial)
- Urine glucose (of limited usefulness)
- Urine ketones (for type 1 diabetes)
- Glycosylated haemoglobin (HbA_{1c}) (essential to know glycaemic control)
- Microalbuminuria (regarded as an early and reversible sign of nephropathy)
- Blood pressure
- Serum lipid levels
- Renal function (serum urea/creatinine)
- ECG

Control guidelines are summarised in Figure 128.1 and Table 128.1.



Figure 128.1 Control guidelines for diabetic management

Table 128.1 Suggested guidelines for glycaemic control (plasma glucose mmol/L)[1, 3]

	Ideal	Acceptable (fair)	Suboptimal or unacceptable
Before meals (fasting)	< 5.5	5.5 - 7	> 7.7
After meals (2 hours post-prandial)	< 7	7 - 10	> 11
HbA$_{1c}$ %[*]	< 7	< 8	> 11

[*] HbA$_{1c}$ is an index of the mean plasma glucose levels over the preceding 2 - 3 months (assume a reference range of 4.5 - 8%). The reference ranges vary in different laboratories.

Goals of Management[1]

All people with diabetes should be encouraged to maintain the following goals for optimum management of their diabetes:

• Blood glucose (fasting)	4 - 6 mmol/L
• HbA$_{1c}$	↑7%
• Cholesterol	< 4.0 mmol/L
• LDL cholesterol	< 2.5 mmol/L
• HDL cholesterol	↓ 1.0 mmol/L
• Blood pressure	< 130/80 mmHg without proteinuria
	↑ 125/75 mmHg with proteinuria (1 g/day)
• BMI	↑ 25 where practicable
• Urinary albumin excretion	< 20 µg/min timed overnight collection
	< 20 mg/L spot collection
• Albumin croatinine ratio	< 2.5 mg/mmol - men
	< 3.5 mg/mmol - women
• Cigarette consumption	Zero
• Alcohol intake	↑ 2 standard drinks (20) g/day (men)
	↑ 1 standard drink (10) g/day (women)
• Exercise	at least 30 minutes walking (or equivalent) 5 or more days/week (total 150 minutes/week)

Blood glucose monitoring at home

This can be done using visual strips or a glucose meter (glucometer). Patients should be advised about the most appropriate glucometer to obtain.

Method

- Obtain capillary blood by pricking the finger with a lancet.
- Place a large drop of blood to cover both colour strips (avoid smearing).
- At 60 seconds blot offexcess blood with tissue paper. (Time can vary from 30 to 60 seconds depending on the brand used.)
- The strip is read by comparing the colour with a colour chart or by using an electronic meter (glucometer).

How often and when?

- Type 1 diabetes:
  - twice a day (at least once)
  - four times a day (before meals and before bedtime) at first and for problems
  - may settle for 1 - 2 times a week (if good control)
- Type 2 diabetes:
  - twice a day (fasting and 2 - 3 hours post-prandial)
  - if good control - once a week or every 2 weeks

Note:

- Capillary blood glucose is approximately 7% higher than venous blood.
- Glucometer error is usually ± 5%.

Glycosylated haemoglobin

Glycosylated haemoglobin is abnormally high in diabetics with persistent hyperglycaemia and is reflective of their metabolic control. The major form of glycohaemoglobin is haemoglobin A_{1c}, which normally comprises 4 - 6% of the total haemoglobin.[4] Glycohaemoglobins have a long half-life and their measure reflects the mean plasma glucose levels over the past 2 - 3 months and hence provides a good method of assessing overall diabetic control. It should be checked every 3 - 6 months.

Type 1 diabetes

The three main objectives of the treatment of type 1 diabetes are:[4]

- Maintain good health, free from the problems of hyperglycaemia and hypoglycaemia.
- Achieve proper growth and maturation for children and protect the foetus and mother in a mother with type 1 diabetes.
- Prevent, arrest or delay long-term macrovascular and microvascular complications.

Insulin regimens for type 1 diabetes[2]

The most commonly used insulin injection preparations are the 'artificial' human insulins. Insulins are classified according to their time course of action:

- rapid-acting and short duration (ultra-short) - lispro insulin, insulin as apart
- short-acting - neutral (regular, soluble)
- intermediate-acting - isophane (NPH) or lente
- long-acting - ultralente, insulin detemir, insuin glargine
- pre-mixed short/intermediate - biphasic (neutral + isophane)

Starting insulin[2]

It is important to use the simplest regimen for the patient and to provide optimal education about its administration and monitoring. Full replacement of insulin is achieved by using 2, 3 or 4 injections per day.

1. The pre-mixed 2 injection system. Give twice daily, 30 minutes before breakfast and before evening meal (e.g. Mixtard 30/70, Humulin 30/70 - the most common)
   - Typical starting dose: 0.3 IU/kg/day - for a 70 kg person use 10 units bd
2. 3 injections per day
   - Short-acting insulin before breakfast and lunch
   - Intermediate- or long-acting insulin before evening meal
3. 4 injections (basal-bolus) system
   - Short-acting insulin before breakfast, lunch and dinner
   - Intermediate-acting insulin at bedtime

Insulin requirements often vary significantly even in the same individual under different lifestyle conditions. The new rapid-acting analogues can be taken with meals.

Methods of giving insulin injections

When

Get the patient to develop a set routine, such as eating meals on time and giving the injection about 30 minutes before the meal.

Where

Into subcutaneous tissue - the best place is the abdomen (Figure 128.2). It is advisable to keep to one area such as the abdomen and avoid injections into the arms, near joints and the groin. The injection should be given at a different place each time, keeping a distance of 3 cm or more from the previous injection. This reduces the risk of the development of lipodystrophy.



Figure 128.2 Method of giving insulin injections; use the abdomen

How

Pinch a large area of skin on the abdomen between the thumb and fingers and insert the needle straight in. After withdrawing the needle, press down firmly (do not rub or massage) over the injection site for 30 seconds.

Guidelines for the patient[5]

The proper injection of insulin is very important to allow your body, which lacks natural insulin, to function as normally as possible. You should be very strict about the way you manage your insulin injections and have your technique down to a fine art.

Common mistakes:

- poor mixing technique when mixing insulin
- wrong doses (because of poor eyesight)
- poor injection technique - into the skin or muscle rather than the soft, fatty layer
- not taking insulin when you feel ill

Drawing up the insulin

Make sure your technique is checked by an expert. You may be using either a single insulin or a mixed insulin. A mixed insulin is a combination of shorter and longer acting insulin and is cloudy.

Rules for mixing

- Always draw up clear insulin first.
- Do not permit any of the cloudy insulin to get into the clear insulin bottle.
- Do not push any of the clear insulin into the cloudy insulin bottle.

Golden rules

- Take your insulin every day, even if you feel ill.
- Do not change your dose unless instructed by your doctor or you are competent to do so yourself.

Problems

Injection sites should be inspected regularly because lipo-hypertrophy or lipo-atrophy can occur.

Type 2 diabetes[1, 6]

First-line treatment (especially if obese):

- diet therapy
- exercise program

Most symptoms improve within 1 to 4 weeks on diet and exercise.[3] Prescribe and ask about exercise at every visit. Aim for an average of 20-30 minutes a day. Suggest variations such as social type exercises. The secret to success is patient compliance through good education and supervision. The role of a diabetic education service, especially with a dietician, can be invaluable. If unsatisfactory control persists after 3 - 6 months, consider adding an oral hypoglycaemic agent (Table 128.2). These agents include insulin secretogogues, such as sulfonylureas and glitinides, which increase insulin production and insulin sensitisers, such as melformin and the glitazones which reduce insulin resistance. If glycaemic targets are not achieved on monotherapy usual practice is to combine a secretogogue and a sensitiser.

Table 128.2 Commonly prescribed oral hypoglycaemic agents[2]

Drug	Duration of action (hours)	Daily dose range	Notes
Sulfonylureas			Hypoglycaemia most common side-effect. Shorter acting sulphonylurea is preferred in elderly.
Gliclazide	12 - 24	40 - 320 mg	Longer acting potent ones cause troublesome hypoglycaemia in elderly.
Glipizide	6 - 12	2.5 - 40 mg	Others: weight gain (common), rash and GIT (rare).
Glibenclamide	16 - 24	2.5 - 20 mg	
Glimepiride	> 24	1 - 4 mg	
Biguanides			Usually reserved for obese but now first line.
Metformin	8 - 12	0.5 - 3 g	Side-effects: GIT disturbances (e.g. diarrhoea, a/n/v) Avoid in cardiac, renal and hepatic disease Lactic acidosis a rare but serious complication.
α-glucosidase inhibitors			
Acarbose	3	150 - 600 mg	Flatulence, skin rashes liver effects.
Glitinides			
Repaglinide	2 - 3	1.5 - 12 mg	Hypoglycaemia, GIT effects.
Thiazolidinediones (glitazones)			Caution with heart failure
Pioglitazone	24	15 - 45 mg	Oedema, weight gain.

Rosiglitazone	24	2 - 8 mg	Oedema, hepatic effects.

Consider metformin as the first-line agent for all patients with type 2 diabetes, irrespective of their weight, unless contraindicated. The usual starting dose is 500 mg once or twice daily. It has proven benefits over the sulphonylureas, expecially in those overweight. Other benefits include no significant weight gain, no hypoglycaemia and an improved lipid profile. If monotherapy does not provide adequate glycaemic control, a combination of metformin with a sulfonylurea is recommended.[5, 6]

When the first oral hypoglycaemics fail (secondary failure) one of the new agents can be added (e.g. acarbose, one of the glitazones). The glitazones can be used as monotherapy but are used more often in combination with melformin, sulfonylureas or insulin.[7]The classic symptoms of hyperglycaemia may be present but more commonly patients experience general disability. Approximately 30% of type 2 patients eventually require insulin even after years of successful oral therapy.

Starting insulin in type 2 diabetes[8, 6]

More than 30% of patients will eventually require insulin often after 10 - 15 years of successful oral therapy.[2] Before commencing insulin one should be assured that the patient's lifestyle activities are being adequately addressed and that oral medication (at recommended maximum dose) is appropriate. There is no clear-cut rule about when to start insulin for patients with HbA1c > 7%, but this can be as early as when drug therapy does not provide adequate control. A golden rule is 'don't delay initiating insulin' and then 'start low and go slow'.[9]

When commencing patients on insulin, reassure them that the injections are not as uncomfortable as finger pricks and that they will feel much improved with more energy. Short-term intensive treatment (approximately 2 weeks) can induce long-term improvement in glycaemic control to the extent of being off insulin for months to years.

It is appropriate to refer to your diabetic team for shared care at this point - when starting insulin.

Suggested stepwise approach[6, 8]

Step 1

- Continue oral agents: metformin ± sulfonylurea ± glitazone or acarbose
- Add 10 Units Isophane insulin at bedtime

Step 2

- Titrate insulin therapy according to fasting blood glucose (6 mmol/L).
- Increase insulin in about 4 - 5 U increments every 3 - 4 days (or more gradually).
- Cease glitazone or acarbose (if used).

Step 3

If larger doses of insulin are required (NPH or mixed regimen), gradually withdraw sulfonylurea, continue metformin and review.

Note: The combination of a glitazone and insulin has been shown to improve control of diabetes sometimes to the extent of being able to reduce insulin dosage. In patients with type 2 diabetes needing relatively high doses of insulin the addition of a glitazone is worth considering.

The importance of diet

Type 1 patients often require three meals and sometimes regular snacks each day. Type 2 patients usually require less food intake and restriction of total food intake.

Principles of dietary management

- Keep to a regular nutritious diet.

- Achieve ideal body weight.
- Reduce calories (kilojoules)
    - added sugar
    - dietary fat
- Follow the glycaemic index values (p. 80)
- Increase proportions of vegetables, fresh fruit, cereal foods.
- Special diabetic foods are not necessary.
- Qualitative diets, rather than quantitative diets (such as ‘exchanges’ or ‘portions’), are now used.

Patient education

The following handout is helpful to patients.

The importance of diet

All diabetics require a diet in which refined carbohydrate and fat intake is controlled. The objectives of the diet are:

- to keep to ideal weight (neither fat nor thin)
- to keep the blood sugar level as near normal as possible

This is achieved by:

- eating good food regularly (not skimping)
- spacing the meals throughout the day (three main meals and three snacks)
- following the healthy food pyramid (p. 78)
- cutting down fat to a minimum
- avoiding sugar and refined carbohydrates (e.g. sugar, jam, honey, chocolates, sweets, pastries, cakes, soft drinks)
- eating a balance of more complex carbohydrates (starchy foods such as wholemeal bread, potatoes and cereals)
- eating a moderate amount of protein
- eating a good variety of fruit and vegetables
- cutting out alcohol or drinking only a little

The importance of exercise[3]

Exercise is very beneficial to your health. Exercise is any physical activity that keeps you fit. Good examples are brisk walking (e.g. 2 km per day), jogging, tennis, skiing and aerobics. Aim for at least 30 minutes three times a week, but daily exercise is ideal. Go slow when you start and increase your pace gradually.

Good advice

- Exercise is important.
- Do not get overweight.
- A proper diet is a key to success.
- A low-fat, no-sugar diet is needed.
- Do not smoke.
- Minimise alcohol.
- Take special care of your feet.
- Self-discipline will help make your life normal.
- Join a diabetic-support organisation.

Psychosocial considerations

The psychological and social factors involving the patient are very influential on outcome. Considerable support and counselling is necessary to help both patient and family cope with the ‘distress’ of the diagnosis and the discipline required for optimal control of their blood glucose. Reasons for poor dietary compliance and insulin administration must be determined and mobilisation of a supportive multidisciplinary network (where practical) is most helpful. The general practitioner should be the pivot of the team. Joining a self-support group can be very helpful.

Foot care

Foot problems are one of the commonest complications that need special attention; so prevention is the appropriate approach. Pressure sores can develop on the soles of the feet from corns, calluses, ill-fitting footwear, and stones and nails. Minor injuries such as cuts can become a major problem through poor healing. Infection of the wound is a major problem. The patient's footwear must be checked.

Advice to the diabetic patient[1, 5]

1. Keep your diabetes under good control and do not smoke.
2. Check your feet daily. Report any sores, infection or unusual signs.
3. Wash your feet daily:
    - Use lukewarm water (beware of scalds).
    - Dry thoroughly, especially between the toes.
    - Soften dry skin, especially around the heels, with lanoline.
    - Apply methylated spirits between the toes to help stop dampness.
4. Attend to your toenails regularly:
    - Clip them straight across.
    - Do not cut them deep into the corners or too short across.
5. Wear clean cotton or wool socks daily; avoid socks with tight elastic tops.
6. Exercise the feet each day to help the circulation in them.

How to avoid injury

- Wear good-fitting, comfortable leather shoes.
- The shoes must not be too tight.
- Do not walk barefoot, especially out of doors.
- Do not cut your own toenails if you have difficulty reaching them or have poor eyesight.
- Avoid home treatments and corn pads that contain acid.
- Be careful when you walk around the garden and in the home.
- Do not use hot water bottles or heating pads on your feet.
- Do not test the temperature of water with your feet.
- Take extra care when sitting in front of an open fire or heater.

Control of hypertension

Studies have highlighted the importance of blood pressure control to reduce macrovascular and microvascular complications in diabetes patients.[12] Their blood vessels (both macro and micro) are more susceptible to hypertensive, namely > 130/85 mmHg, damage. Non-pharmacological measures should be tried first.

Preferred pharmacological agents are ACE or inhibitors and calcium channel blockers.[4, 5]

Current recommendations are for tight control of blood pressure to below a threshold of 130/80 mmHg in patients with uncomplicated diabetes.[3, 7]

Getting to target blood pressure (< 130/80 mmHg)[*]

Step 1: Diet, exercise, weight control

Step 2: ACEI or AII RA

Step 3: ACEI/AII RA and diuretic

Step 4: Beta-blocker

AII RA = angiotension iireceptor antagonist

* < 125/75 mmHg if proteinurea > 1 g/day present.

Control of dyslipidaemia[3, 9]

Mixed hyperlipidaemia is a common finding in patients with diabetes. Dyslipidaemia (especially hypercholesterolaemia) is an independent risk factor for the macrovascular complications of diabetes and proper control is important. Non-pharmacological measures should be tried first. The preferred agents are HMGCoA reductase inhibitors and resins for hypercholesterolaemia and fibrates and resins for mixed hyperlipidaemia.

Targets should be:

- total cholesterol - < 4 mmol/L
- triglycerides - < 1.5 mmol/L
- HDL cholesterol - ↓ 1 mmol/L
- LDL cholesterol - < 2.5 mmol/L

Management in summary[10]

The ABC of diabetic care is summarised in Table 128.3. The key to ongoing control of diabetes is to maintain the HbA1c below 7%. In patient review, the NH & MRC guidelines emphasise lifestyle review as step one.[3] A useful lifestyle evaluation mnemonic is NEAT:

- Nutrition - eat less, reach ideal weight, healthy low fat/simple carbohydrate diet
- Exercise - including ‘walk more’, interesting physical activities
- Avoid toxins - alcohol, tobacco, salt, sugar, illicit drugs
- Tranquillity - rest, recreation and stress reduction

Table 128.3 The ABCss of diabetes care[7]

Risk factor	Target
HbA1c	< 7%
BP	< 130/80 *
Cholesterol	< 4 mmol/ L **
Salicylates	Aspirin 75 - 150 mg/day
Smoking	Quit

* < 125/75 if proteinuria exists

** corresponding to LDL cholesterol <2.5 mmol/L

Antihypertensives and statins have an important role in management. A meta-analysis of the use of low-dose acetyl salicylic acid (75 - 150 mg/day) showed cardiovascular risk reduction in people with diabetes.[11] It is appropriate that a ‘type 2 tablet’ has been advocated for most people with diabetes.[12] The suggested pill mix is *metformin, an ACE inhibitor, a statin and aspirin*.

Metabolic complications of diabetes

Hypoglycaemia

Hypoglycaemia[2, 4] occurs when blood glucose levels fall to less than 3.0 mmol/L. It is more common with treated type 1 diabetes but can occur in type 2 diabetes patients on oral hypoglycaemic drugs, notably sulphonylureas (biguanides hardly ever cause hypoglycaemia). It is appropriate to ask often about symptoms of hypoglycaemia: 'recurrent hypoglycaemia begets hypoglycaemic unawareness'.

Clinical variations

1. Classic warning symptoms: sweating, tremor, palpitations, hunger, perioral paraesthesia. Usually treated with refined carbohydrate (e.g. glucose).
2. Rapid loss of consciousness, usually without warning - hypoglycaemic unawareness is less common.
3. Coma: stuporose, comatose or 'strange' behaviour.

For mild cases give something sweet by mouth, followed by a snack.

Treatment of hypoglycaemia requires one dose of carbohydrate.

Dose: (one of)

- 2 barley sugars
- 6 jelly beans
- glass of lemonade
- teaspoon of honey

Take one dose, and don't repeat unless still unwell 10 minutes later. Follow with a complex carbohydrate meal.

Treatment of severe cases or patient unconscious

Treatment of choice

- 10 - 25 mL 50% dextrose IV (instil rectally using the nozzle of the syringe if IV access difficult)

or

Alternative

- 1 mL glucagon IM

Admit to hospital if concerned (rarely necessary). Ascertain cause of the hypo-glycaemia and instruct the patient how to avoid a similar situation in the future.

Diabetic ketoacidosis

This life-threatening emergency requires intensive management. It usually occurs during an illness (e.g. gastroenteritis) when insulin is omitted.

Clinical features

- Develops over a few days, but may occur in a few hours in 'brittle' diabetics
- Preceding polyuria, polydipsia, drowsiness
- Vomiting and abdominal pain
- Hyperventilation - severe acidosis (acidotic breathing)
- Ketonuria

Management

- Arrange urgent hospital admission.
- Give 10 units rapid-acting insulin IM (not SC).
- Commence IV infusion of normal saline.

Hyperosmolar hyperglycaemia[2, 1]

Patients with this problem may present with an altered conscious state varying from stupor to coma and with dehydration.

The onset may be insidious over a period of weeks, with fatigue, polyuria and polydipsia. The key features are marked hyperglycaemia and dehydration without ketoacidosis. It occurs typically in uncontrolled type 2 diabetes, especially in elderly patients. Sometimes they have previously undiagnosed diabetes. There may be evidence of an underlying disorder such as pneumonia or a urinary infection. The essential findings are extreme hyperglycaemia and high plasma osmolarity. The condition has a high mortality - even higher than ketoacidosis.

Treatment

- IV fluids, e.g. normal to ½ normal saline, given slowly
- Insulin - relatively lower doses than acidosis

Lactic acidosis[2]

Patients with lactic acidosis present with marked hyperventilation 'air hunger' and confusion. It has a high mortality rate and must be considered in the very ill diabetic patient taking metformin, especially if renal function is impaired. The risk of lactic acidosis is low if the therapeutic dose of metformin is not exceeded. The investigations reveal blood acidosis (low pH), low bicarbonate, high serum lactate, absent serum ketones and a large anion gap. Treatment is based on removal of the cause, rehydration and alkalinisation with IV sodium bicarbonate.

Other issues in diabetic patients

Erectile dysfunction[2]

The prevalence of ED in men over 40 years may be as high as 50%. It may be caused by macrovascular disease, pelvic autonomic neuropathy or psychological causes. Those with organic-based ED may benefit from appropriate counselling and (if not taking nitrates) one of the phosphodiesterase inhibitors, starting with a low dose. The risk of cardiovascular disease needs to be evaluated.

Female dysfunction

Autonomic dysfunction may result in reduced vaginal lubrication with arousal in women, but not the degree of sexual dysfunction that affects men. Appropriate education, reassurance and the use of lubricants should be helpful.

Postural hypotension[2]

Autonomic neuropathy-related postural hypotension may be compounded by medication, including antihypertensives and antiangina agents. Persistent problems may be helped by graduated compression stockings to decrease venous pooling. If it continues to be a severe problem, the use of oral fludrocortisone may be helpful.

Gastroparesis

Symptoms of gastroparesis (due to autonomic neuropathy) with decreased gastric emptying include a sensation of fullness, dysphagia, reflux or recurrent nausea and vomiting, especially after meals. Treatment options include medication with domperidone, cisapride or erythromycin. A recent development is injections of botulinum toxin type A into the pylorus via gastroscopy to facilitate gastric emptying.

Diabetes and driving[1]

Diabetes may impair driving through medication causing hypoglycaemia or its complications including visual impairment. In 2003 revised medical standards, *Assessing Fitness to Drive*, were released for drivers of private and commercial vehicles. Patients are expected to provide details to the driver licensing authority and to their vehicle insurance company. There are specific legal obligations of medical practitioners and drivers. In general terms people controlled by diet alone have no restrictions for driving whereas those on insulin may obtain a conditional licence subject to annual or 2 yearly review. Further details can be found at < http://www.austroads.com.au/aftd/index.html >.

Contraception

The combined oral contraceptive pill is generally regarded as the most appropriate option for birth control in women not interested in permanent sterilisation. It is important in history taking to keep polycystic ovarian syndrome in mind.

The future[6]

- Use of immunosuppressants and immunomodulators for type 1 diabetes
- Glucagon-like peptide and amylin-like peptides for type 2 diabetes
- Continuous implantable venous glucose monitoring
- Combination ‘type 2 tablet’
- Inhaled insulin
- Transplantation:
  - combined kidney/pancreas
  - islet cells

TREATMENT ERRORS AND PITFALLS[3]

- Avoid prescribing oral hypoglycaemic agents prematurely. Allow a reasonable trial of diet and exercise for type 2 patients, especially if they are overweight.
- Review the need for continued oral therapy after 3 months of treatment.
- Glucose tolerance tests should be avoided if the diagnosis can be made on the basis of symptoms and fasting or random blood sugar (a glucose load carries a risk of hyperosmolar coma).
- Keep an eye on the development of ketones in type 1 patients by checking urinary ketones and, if present, watch carefully because diabetic ketoacidosis is a life-threatening emergency.

WHEN TO REFER[2]

- Type 1 diabetic patients for specialist evaluation and then 1 to 2 yearly review.
- Type 2 diabetic patients:
  - all young patients
  - those requiring education
  - those requiring insulin
  - those with complications
- For ophthalmological screening: every 1 to 3 years to inspect retina.
- Diabetics with treatable complications, including:
  - retinopathy
  - nephropathy
  - neuropathy

SHARED CARE

The management of the diabetic patient provides an ideal opportunity for shared care between a co-operative team comprising the patient, the general practitioner and the specialist diabetic team. The objective is to encourage patients to attend their own doctor for primary care and be less reliant on hospital outpatient services or the diabetic clinics. A well co-ordinated arrangement with good communication strategies provides optimal opportunities for the ongoing education of the patient, the general practitioner and the specialist diabetic team.

Practice tips

- For every diagnosed diabetic there is an undiagnosed diabetic, so vigilance for diagnosing diabetes is important.
- Follow-up programs should keep to a prepared format. A format that can be used is presented in Table 128.4 and

Table 128.5.

- Hyperglycaemia is a common cause of tiredness. If elderly type 2 diabetic patients are very tired, think of hyperglycaemia and consider giving insulin to improve their symptoms.
- The management of the diabetic patient is a team effort involving family members, a nurse education centre, podiatrists, domiciliary nursing service, general practitioner and consultant.
- If a diabetic patient (particularly type 1) is very drowsy and looks sick, consider first the diagnosis of ketoacidosis.
- Foot care is vital: always examine the feet when the patient comes in for review.
- Treat associated hypertension with ACE inhibitors or a calcium-channel blocker (also good in combination).
- Use a team approach and encourage joining special support groups (e.g. Diabetes Australia).
- Never let the sungo down on pus in a diabetic foot-admit to hospital.[6]
- If a foot ulcer hasn't healed in 6 weeks, exclude osteomyelitis. Order an MRI and investigate the vasculatuse.
- Prevention/detection of coronary heart disease should be an integral part of all consultation.

Table 128.4 Diabetes control: 4 - monthly review

- Discourage smoking
- Review symptoms
- Check weight/(BMI), BP, urine
- Review self-monitoring
- Review exercise and physical activity
- Review HbA1_c
- Review lipid levels

Table 128.5 Diabetes control: an annual review program[5]

1 History

- Smoking and alcohol use
- Symptoms of hypoglycaemia, hyperglycaemia
- Check symptoms relating to eyes, circulation, feet[*]

2 Examinations

- Weight, height, BMI
- Blood pressure - standing and lying
- Examine heart[*]
- Carotid and peripheral pulses[*]
- Eyes
    - visual acuity (Snellen chart)
    - ? cataracts
    - optic fundi (or ophthalmologist referral)[*]
    - ? diabetic retinal photography
- Tendon reflexes and sensation for peripheral neuropathy[*]
- Skin (general)
- Foot examination including footwear[*]
- Check injection sites
- Urine examination: albumin, ketones, glucose, nitrites

3 Review biochemical levels

[*] These items comprise a program for detection of long-term complications. They should be conducted annually, commencing 5 years after diagnosis.

REFERENCES

1. *Diabetes Management in General Practice*. Melbourne: Diabetes Australia & RACGP, 2005 - 6: 13 - 64
2. Shenfield G (Chair: expert group). *Therapeutic guidelines: Endocrinology* (Version 3). Melbourne: Therapeutic Guidelines Ltd, 2004
3. NH & MRC. Evidence-based guidelines for the management of type 2 diabetes mellitus. Canberra: NH & MRC, 2001
4. Cohen M. Management of insulin dependent diabetes. Aust Fam Physician, 1990; 19: 1201 - 14
5. Murtagh J. *Patient Education* (4th edn). Sydney: McGraw-Hill, 2005: 193 - 7
6. Newnham H. Diabetes in motion. Proceedings of the 27th Annual Update Course for General Practitioners. Monash University, 2005
7. Phillips P. Type 2 diabetes. CHECK Program, unit 401, 2005: 4 - 19
8. Reducing risk in type 2 diabetes. NPS News, 2005: 1 - VI
9. Goudswaard AN et al. Cochrane Database Syst Rev, 2004; (4): CD003418: pub 2
10. Evidence-based best practice guidelines. Management of type 2 diabetes. Wellington: New Zealand Guidelines Group, 2003. < http://www.nzgg.org.nz/guidelines/0036/Diabetes >
11. Colwell JA. Aspirin therapy in diabetes. Diabetes Care, 2001; 24: 62 - 3
12. Phillips P, Braddon J. The type 2 tablet. Evidence-based medication for type 2 diabetes. Aust Fam Physician, 2003; 32; 431 - 6

CHAPTER 129. CHRONIC HEART FAILURE

> In the year 1775 my opinion was asked concerning a family recipe for the cure of the dropsy. I was told that it had long been kept a secret by an old woman in Shropshire who had sometimes made cures after the more regular practitioners had failed - this medicine was composed of twenty or more different herbs and the active herb could be no other than the Foxglove.
>
> WILLIAM WITHERING (1741 - 99), ON THE USE OF FOXGLOVE (DIGITALIS) IN THE TREATMENT OF HEART DISEASE

CHRONIC HEART FAILURE

Heart failure occurs when the heart is unable to maintain sufficient cardiac output to meet the demands of the body for blood supply during rest and activity.

Chronic heart failure (CHF) remains a very serious problem with a poor prognosis. It has a 50% mortality within 3 years of the first hospital admission.[1] Overseas data indicates that 1.5% of the adult population have heart failure. The prevalence of CHF has been shown to increase from approximately 1% in those aged 50 to 59 years to over 5% in those 65 and older, to over 50% in those 85 years and older.[2] Australian research suggests that under-treatment with the all important ACE inhibitors continues to be a problem. A major goal of management of CHF is the identification and reversal where possible of underlying causes and/or precipitating factors.

DIAGNOSIS OF CHF

The classis symptom of CHF is dyspnoea on exertion but symptoms may be reported relatively late, mainly due to a sedentary lifestyle. Dyspnoea can progress as follows: exertional dyspnoea → dyspnoea at rest → orthopnoea → paroxysmal nocturnal dyspnoea.

Symptoms in summary

- Dyspnoea (as above)
- Irritating cough (especially at night)
- Lethargy/fatigue
- Weight change: gain or loss
- Dizzy spells
- Palpitations
- Ankle oedema

The irritating cough due to left ventricular failure can be mistaken for asthma, bronchitis or ACE-inhibitor-induced cough.

PHYSICAL EXAMINATION

The physical examination is very important for the initial diagnosis and evaluation of progress. The signs are as follows.

Signs

There may be no abnormal signs initially. It is helpful clinically to differentiate between the signs of right and left heart failure:

- Left heart failure:
    - Tachycardia
    - Low volume pulse
    - Tachypnoea
    - Laterally displaced apex beat

  - Bilateral basal crackles
  - Gallop rhythm (3rd heart sound)
  - Pleural effusion
  - Poor peripheral perfusion
- Right heart failure:
  - Elevated jugular venous pressure
  - Right ventricular heave
  - Peripheral/ankle oedema
  - Hepatomegaly
  - Ascites

Auscultation is important to identify adventitious sounds, a third heart sound and possible underlying valvular disease.

Systolic versus diastolic heart failure

The classic heart failure is systolic failure due to an inadequate pumping action of the heart. The ventricle is dilated and contracting poorly (left ventricular ejection fraction < 40%).[3]

However, diastolic heart failure, which is being recognised more widely, is due to impairment of left ventricular filling. At least one-third to half of heart failure presentations are due to diastolic heart failure (impaired ventricular relaxation).[2] It should be suspected in the elderly with hypertension and a normal heart size on chest X-ray who present with dyspnoea or pulmonary oedema.[4]

It is particularly common in elderly females.

Note that patients can have simultaneous systolic diastolic and failure.

The oedema of heart failure

Peripheral oedema appears initially on the lower legs and is ‘pitting’. To assess the presence of pitting, which is usually graded on a four-point scale, press firmly yet gently with the thumb for 5 - 10 seconds over the dorsum of the feet, behind each medial malleolus and over the shins. With increasing severity of failure the oedema extends proximally to involve the abdomen. In the recumbent position it may be apparent over the sacrum.[5]

Determining severity of heart failure

The severity of heart failure can be considered from three different perspectives: the severity of the symptoms, the degree of impairment of cardiac function and the severity of the congestive state. The severity of the symptoms or the degree of functional disability is usually described according to the New York Heart Association criteria (Table 129.1). The left ventricular ejection fraction provides an indication of cardiac function.

Table 129.1 New York Association classification of functional disability in heart failure

Class I (asymptomatic)	**No limitation**: cardiac disease present, but ordinary physical activity causes no symptoms such as fatigue, dyspnoea or palpitation, or rapid forceful breathing
Class II (mild)	**Slight limitation**: ordinary activity (moderate exertion) causes symptoms but patients comfortable at rest
Class III (moderate)	**Marked limitation**: symptoms with less than ordinary physical activity (mild exertion) although patients still comfortable at rest
Class IV (severe)	Unable to carry on any activity without symptoms; may have symptoms at rest

CAUSES OF HEART FAILURE

Causes of CHF can be classified under systolic heart failure (impaired ventricular contraction) and diastolic heart failure (impaired ventricular relaxation). Diagnosis is based on echocardiography.

Systolic heart failure

Ischaemic heart disease, including previous myocardial infarction, is the most common cause, accounting for approximately three-quarters. There is often a history of at least one myocardial infarction.[2] Essential hypertension is the other common cause.

Other causes:

- valvular heart disease, mainly aortic and mitral incompetence
- high output states, e.g. anaemia, hyperthyroidism, Paget's disease
- non-ischaemic idiopathic dilated cardiomyopathy
- viral cardiomyopathy
- alcoholic cardiomyopathy
- other cardiomyopathies - diabetic, familial
- persistent arrhythmias, especially atrial fibrillation
- other systemic illness, e.g. sarcoidosis, scleroderma, myxoedema

Diastolic heart failure

Obesity, hypertension and diabetes are significant risk factors. Common causes include ischaemic heart disease, systemic hypertension, aortic stenosis, atrial fibrillation (inadequate filling), hypertrophic cardiomyopathy and pericardial disease.

Investigations

The following should be considered:

- Echocardiography[2]
  The transthoracic echocardiogram is the investigation of choice to measure ventricular function. It can differentiate between systolic dysfunction and those with normal systolic function but abnormal diastolic filling. It gives information about left and right ventricular systolic and ventricular diastolic function, left and right ventricular size, volumes, thickness, structure and function. Similarly it provides other information about cardiac valves, congenital heart defects and pericardial disease.
- Electrocardiogram: to look for evidence of ischaemia, conduction abnormalities, arrhythmias and LV hypertrophy
- Chest X-ray: to look for:
  - cardiomegaly and interstitial oedema
  - upper lobe blood diversion
  - fluid in fissures
  - oedema in perihilar area with prominent vascular markings
  - small basal pleural effusions
  - Kerley B lines = raised pulmonary venous pressure
  - frank pulmonary oedema
- Spirometry/respiratory function testing: to detect associated airways dysfunction
- B type natriuretic peptide: an indicator of severity of CHF and prognosis

Peripheral markers

- FBE and ESR: anaemia can occur with CHF; severe anaemia may cause CHF
- Serum electrolytes: usually normal in CHF, important to monitor management
- Kidney function tests: congestive hepatomegaly gives abnormal LFTs
- Liver function tests: congestive hepatomegaly gives abnormal LFTs
- Urinalysis
- Thyroid function tests, especially if atrial fibrillation

- Viral studies: for suspected viral myocarditis

Specialised cardiac investigation (specialist directive)

- Coronary angiography - for suspected and known ischaemia
- Haemodynamic testing
- Endomyocardial biopsy
- Nuclear cardiology

TREATMENT OF HEART FAILURE

The treatment of heart failure includes determination and treatment of the cause, removal of any precipitating factors, appropriate patient education, general non-pharmaceutical measures and drug treatment.

Prevention of heart failure

The emphasis on prevention is very important since the onset of heart failure is generally associated with a very poor prognosis. Approximately 50% of patients with heart failure die within 5 years of diagnosis.[6]

The scope for prevention includes the following measures:[7]

- dietary advice (e.g. achievement of ideal weight, optimal nutrition)
- emphasising the dangers of smoking and excessive alcohol
- control of hypertension
- control of other risk factors (e.g. hypercholesterolaemia)
- early detection and control of diabetes mellitus
- early intervention of myocardial infarction to preserve myocardial function (e.g. thrombolytic therapy, stenting)
- secondary prevention after the occurrence of myocardial infarction (e.g. beta- blockers, ACE inhibitors and aspirin)
- appropriate timing of surgery or angioplasty for ischaemic or valvular heart disease

Treatment of causes and precipitating factors

Determination and treatment of the causes has been largely covered in the section on prevention. Precipitating factors that should be treated include:

- arrhythmias (e.g. atrial fibrillation)
- electrolyte imbalance, especially hypokalaemia
- anaemia
- myocardial ischaemia, especially myocardial infarction
- dietary factors (e.g. malnutrition, excessive salt or alcohol intake)
- adverse drug reactions (e.g. fluid retention with NSAIDs and COX-2 agents) tricyclic antidepressants, the calcium-channel blockers (verapamil and diltiazem), selected antiarrhythmic agents, e.g. quinidine, and the glitazones[8]
- infection (e.g. bronchopneumonia, endocarditis)
- thyrotoxicosis
- lack of compliance with therapy
- fluid overload

General non-pharmacological management

- Education and support
- Smoking: encourage no smoking
- Refer for a rehabilitation program with interdisciplinary care
- Encourage physical activity especially when symptoms absent or mild
- Rest if symptoms are severe

- Weight reduction, if patient obese
- Salt restriction: advise no-added-salt diet (< 2 g or 60 - 100 mmol/day)
- Water restriction: water intake should be limited to 1.5 L/day or less in patients
- with advanced heart failure, especially when the serum sodium level falls below 130 mmol/day[6]
- limit caffeine to 1 - 2 cups coffee/tea a day
- limit alcohol to 1 SD a day
- fluid aspiration if a pleural effusion or pericardial effusion is present
- daily weighing - check significant weight gain or loss

Other general measures[1]

- Optimise cardiovascular risk factors, e.g. BP, lipids, HbA_{1C}
- Monitor emotional factors including depression
- Regular review
- Vaccination: annual influenza, 5 yearly pneumococcal
- 2 yearly echo cardiography (or more) as indicated

Drug therapy of heart failure due to left ventricular systolic dysfunction

Any identified underlying factor should be treated.

Evidence from RCTs shows the beneficial results from ACE inhibitors,[2, 8] angiotensin II blockers, digoxin (improves outcome in people already receiving diuretics and ACE inhibitors), beta-blockers and spironolactone (in severe heart failure).

Atrial fibrillation should be treated with digoxin. Vasodilators are widely used for heart failure and angiotensin converting enzyme inhibitors (ACEI) are currently the most favoured vasodilator.

Note: Monitor and maintain normal potassium level in all patients.

ACE inhibitors improve prognosis in all grades of heart failure and should be employed as the initial therapy in all patients; except where they are contraindicated, e.g. renal artery stenosis, angioedema.

Diuretics have an important place in patients with fluid overload. As a rule they should be added to an ACEI to achieve euvolaemia. Diuretics should be used in moderation and excessive doses of a single drug avoided. In patients with systolic LV dysfunction they should not be used as monotherapy.[7] Close monitoring of weight, renal function and electrolytes is required. Loop diuretics such as frusemide, bumetanide or ethacrynic acid are commonly used, especially for heart failure of moderate severity.[7] Thiazide and related diuretics produce a gradual diuresis and are recommended for mild heart failure. Examples include hydrochlorothiazide, bendrofluazide, chlorthalidone or indapamide.

Digoxin has two indications for its use in CHF leading to improved symptoms, namely heart failure and atrial fibrillation to control ventricular rate and in heart failure and sinus rhythm to relieve symptoms not controlled by the three first-line drugs.

Initial therapy of heart failure

1. ACE inhibitor (start low, aim high) Dosage of ACE inhibitor: commence with ¼ to ½ lowest recommended therapeutic dose and then adjust it for the individual patient by gradually increasing it to the maintenance dose (Table 129.2).
2. Add a diuretic (if congestion): frusemide 20 - 40 mg (o) once or twice daily or hydrochlorothiazide 25 - 50 mg (o) daily (or other thiazide) or indapamide 1.5 - 2.5 mg(o) daily

Table 129.2 Some ACE inhibitors in common usage[7, 8]

ACE inhibitor	Initial daily dose	Usual maintenance daily dose

Captopril	6.25 mg (o) nocte	25 mg (o) tds
Enalapril	2.5 mg (o) nocte	10 mg (o) bd
Fosinopril	5 mg (o) nocte	20 mg (o) nocte
Lisinopril	2.5 mg (o) nocte	5 - 20 mg (o) nocte
Perindopril	2 mg (o) nocte	4 mg (o) nocte
Quinapril	2.5 mg (o) nocte	20 mg (o) nocte
Ramipril	1.25 mg (o) nocte	5 mg (o) nocte
Trandolapril	0.5 mg (o) nocte	2 - 4 mg (o) daily

ACE inhibitors

- ACE inhibitors are regarded as the agents of first choice because they correct neuroendocrine abnormalities and reduce cardiac load by their vasodilator action.
- Every effort should be made to up-titrate to the highest tolerated dose.
- The first dose should be given at bedtime to minimise orthostatic hypotension.
- If the ACEI is not tolerated (e.g. due to cough) consider an angiotensin II receptor antagonist as they have proven benefit in CHF.[9]
- In practice the usual initial treatment of heart failure is an ACE inhibitor plus diuretic. This combination optimises response and improves diuretic safety.
- Consider stopping any diuretic for 24 hours before starting treatment with an ACEI.
- Potassium sparing diuretics or supplements should not be given with (or at least used with caution) with ACEI because of the danger of hyperkalaemia.
- Kidney function and potassium levels should be monitored in all patients.

Some authorities are concerned about the over-reliance on diuretics and also about compliance as well as side-effects. Once the diuretic effect has been achieved, diuretics may be withdrawn and fluid restriction advised. The ACEI is then used alone.

Beta-blockers

Selective beta-blockers have been shown to prolong survival of patients with mild to moderate CHF taking ACE inhibitors who are stabilised. Start with extremely low doses (Table 129.3).

Table 129.3 Beta-blockers approved to treat heart failure

Beta-blocker	Initial daily dose	Target dose
Bisoprolol	1.25 mg (o) daily	10 mg (o) daily
Carvedilol	3.125 mg (o) bd	25 mg (o) daily
Metoprolol extended release	23.75 mg (o) daily	190 mg (o) daily

Heart failure (unresponsive to first-line therapy) - stepwise strategy[7, 10]

- ACE inhibitor +
- frusemide 40 - 80 mg (o) bd +
- a selective beta-blocker +
- digoxin (if not already taking it)[7]
    - 0.5 - 0.75 mg (o) statim (depending on renal function)
    - then 0.5 mg (o) 4 hours later
    - then 0.5 mg the following day

  - then individualise maintenance +
- spironolactone 12.5 - 25 mg (o) daily (monitor potassium and RFTs)

Severe heart failure[7, 10]

Seek specialist advice.

Hospital with bed rest.

- ACE inhibitor (o) to maximum tolerated dose +
- frusemide to max. 500 mg/day +
- spironolactone (low dose) 25 mg/day

  If poorly controlled, consider adding:
  - thiazide diuretic
  - spironolactone - doses up to 100 - 200 mg daily
  - a beta-blocker
  - digoxin
  - heparin (if confined to bed)

    If still uncontrolled consider other vasodilators:
    - isosorbide dinitrate and hydralazine

Consider cardiac transplantation for appropriate patients with end-stage heart failure (e.g. patients under 50 with no other major disease). Other surgical options include heart valvular surgery, coronary artery bypass surgery and surgical ventricular restoration (surgical reduction of an enlarged left ventricle).

A flow chart for the basic management of heart failure is presented in Figure 129.1.


ACE inhibitor
titrate upwards
+
diuretic
if needed to control congestive symptoms
selective β-blocker
titrate upwards
add digoxin if indicated
spironolactone
consider vasodilator
consider transplantation

Figure 129.1 A stepwise management approach to heart failure [10]

Diastolic heart failure[7, 8]

Management is based on treating the cause such as hypertension, ischaemia and diabetes. The basic treatment is with inotropic agents such as calcium antagonists (verapamil or diltiazem) and beta-blockers. If possible avoid diuretics (except for congestion), digoxin, nitrates/vasodilators, nifedipine. Excessive diuresis from overzealous diuretic therapy can cause severe consequences for cardiac output. ACE inhibitors can be used with caution.

Pitfalls in management

- The most common treatment error - excessive use of diuretics[4]
- Giving an excessive loading dose of ACE inhibitor
- Failure to correct remedial causes or precipitating factors
- Failure to measure left ventricular function
- Failure to monitor electrolytes and renal function

Acute severe heart failure

For the treatment of acute pulmonary oedema refer to page XXX under emergency care.

Device-based heart failure treatments[1]

The use of mechanical devices to treat patients with severe failure is gaining momentum. They include:

- implantable cardiac defibrillators
- biventricular pacemakers
- left ventricular assist devices (Definitive Ventrassist)

The evidence for the efficacy of these devices is good, but limitations include cost and infection. Biventricular pacing or cardiac resynchronisation therapy resynchronises cardiac contraction in patients with systolic CHF and left branch bundle block. Ventrassist is based on a continuous flow rotary blood pump that is surgically implanted in the abdominal wall and attached to the apex of the ventricle.

Practice tips

- Use a multidisciplinary team approach for the patient with CHF and refer early and readily for specialist advice.
- ACE inhibitors, if tolerated, are recommended for all patients with heart failure, whether symptoms are mild, moderate or severe.[1]
- The primary reason for prescribing an ACE inhibitor is to reduce the risk of death or hospitalisation.
- Diuretics are very effective in the presence of fluid overload, but should be used in combination with an ACE inhibitor - not as monotherapy.
- Drug treatment should include an ACE inhibitor and beta-blocker wherever possible.[1]
- Device-based heart failure therapy is an expanding field and includes three major groups of devices: biventricular pacemakers (cardiac resynchronisation therapy); implantable cardiac defibrillators and left ventricular assist devices.[1]

REFERENCES

1. Piterman L, Zimmet H, Krum H, Tonkin A, Yallop J. Chronic heart failure: optimising care in general practice. Australian Fam Physician 2005; 34(7): 547 - 53.
2. National Heart Foundation of Australia and the Cardiac Society of Australia and New Zealand. *Guidelines on the Contemporary Management of the Patient with Chronic Heart Failure in Australia*. 2002.
3. Krum H et al. Med J Aust 2001; 174: 439 - 44.
4. Kelly DT. Cardiac failure. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 97 - 9.
5. Davis A, Bolin T, Ham J. *Symptom Analysis and Physical Diagnosis* (2nd edn). Sydney: Permagon Press, 1990: 173.
6. Smith A (Chair: writing group). *Therapeutics Guidelines: Cardiovascular* (Version 4). Melbourne: Therapeutics Guideline Ltd, 2003: 141 - 59.
7. Kumar P, Clark M. *Clinical Medicines* (5th edn). London: Saunders 2003: 754.
8. A stepwise approach to heart failure management. NPS News, 2004; 36: i - iv.

9. Granger CB et al. Effects of candesartan in patients with chronic heart failure intolerant to ACE inhibitors: the CHARM - Alternative trial. Lancet 2003; 362: 772 - 6.
10. The National Collaborating Centre for Chronic Conditions. *NICE Guideline No. 5. Chronic Heart Failure*. National clinical guideline for diagnosis and management in primary and secondary care, 2003.

Further information: http://www.heartfoundation.com.au/downloads

CHAPTER 130. EMERGENCY CARE

When Elisha arrived, he went alone into the room and saw the boy lying dead on the bed. He closed the door and prayed to the Lord. Then he lay down on the boy, placing his mouth, eyes and hands on the boy's mouth, eyes and hands. As he lay stretched out over the boy, the boy's body started to get warm - the boy sneezed seven times and then opened his eyes.

II KINGS 4: 32 - 5 (A MIRACLE OR SUCCESSFUL ARTIFICIAL RESUSCITATION?)

DEFINITION OF THE EMERGENCY

Emergency: ‘An event demanding immediate medical attention’.

The general practitioner must be available and organised to cope with the medically defined emergency when it comes. Emergency care outside the hospital represents one of the most interesting and rewarding areas of medical practice. City doctors will have to modify their degree of availability, equipment and skills according to paramedical emergency services, while others, especially remote doctors, will need total expertise and equipment to provide optimal circumstances to save their patients' lives.

In dealing with a specific emergency, the doctor adopts a different approach. Instead of taking a history and performing an examination in the usual way, he or she replaces this with a technique of rapid assessment and immediate management. In fact, the diagnosis may be possible on the information available over the telephone.

An important yet obvious concept is that of ‘time criticality’, which implies that certain patients are at high risk of a critical outcome of deterioration if there is significant delay in appropriate management. This applies particularly to acute cornary syndromes.

KEY FACTS AND CHECKPOINTS

- The commonest emergency calls in a survey of a typical community[1] were accidents and violence (50.7%), abdominal pain (9.9%), dyspnoea (7.2%), chest pain (5.8%), syncope/blackout (5.2%), other acute pain (5.0%).
- The prevalence of emergency calls was 2.6 per 1000 population per week.
- The commonest specific conditions in this study were lacerations 19%, fractures 11%, injuries from transport accidents 11%, bronchial asthma 4%, ischaemic heart pain 3.5%, appendicitis 3%.
- The commonest causes of sudden death were myocardial infarction 67%, accidents 10%, cerebrovascular accidents 7%, pulmonary embolism 6%, suicide 4%.
- The main vital emergency procedures were cardiopulmonary resuscitation, intubation and ventilation, intravenous access including cutdown, intravenous (or rectal) dextrose and arrest of haemorrhage.

PRINCIPLES OF MANAGEMENT

The important principles of management of the emergency call can be summarised as follows:

1. The practitioner must be aware of life-threatening conditions.
2. The practitioner should be prepared mentally and physically:
   plan, equip and practise
3. Chest pain/collapse/myocardial infarction (collectively) represents the premium emergency call.
4. Beware of children with respiratory distress and traumatic injuries.
5. The most savable patients are those with blood loss. Hence IV fluids for intravascular volume expansion and blood availability are essential.
6. The necessary basic skills to cope with most emergencies involve ABC - airway, breathing, circulation.
7. Have the equipment and the skills to handle potentially HIV-contaminated body fluids.[2]

8. Stet 70% of cardiac arrests occur in the home so the availability of a portable defibrillator is important.

Vital basic skills

1. Rapid intravenous access: direct or cutdown.
2. Cardiopulmonary resuscitation, including upper airway relief, intubation and ventilation, treatment of cardiac arrhythmias and defibrillation.
3. Cricothyroidotomy.
4. Arrest of haemorrhage.
5. Knowledge of usage of common emergency drugs.

When to get up and go

The following symptoms and signs make attendance at the emergency mandatory:

- unconsciousness
- convulsions
- chest pain in an adult, especially associated with pallor and sweating
- pallor and sweating in any patient with pain, collapse or injury
- collapse, especially at toilet
- significant haemorrhage
- breathlessness, including bronchial asthma
- the agitated patient threatening homicide or suicide (take a police officer for company)
- serious accidents
- asthmatic patients

Don't forget the value of oxygen

Ideally, the doctor who attends emergency calls should carry an oxygen delivery unit or at least rely on the simultaneous arrival of an ambulance with resuscitation equipment. Most cases require a high flow rate of 8 - 10 L/minute.

Typical medical emergencies requiring oxygen:

- bronchial asthma
- acute pulmonary oedema
- acute anaphylaxis
- myocardial infarction
- cardiopulmonary arrest
- collapse

TWELVE GOLDEN RULES

Here are twelve important rules for the diagnostic approach to the emergency call.

1. Always consider the possibility of hypoglycaemia and opioid overdosage in the unconscious patient.
2. Consider intra-abdominal bleeding first and foremost in a patient with abdominal pain who collapses at toilet.
3. Acute chest pain represents myocardial infarction until proved otherwise.
4. Exclude acute epiglottitis in a child with a sudden onset of respiratory distress and pallor.
5. Consider the possibility of a ruptured intra-abdominal viscus in any person, especially a child, with persistent post-traumatic abdominal pain.
6. Always consider the possibility of acute anaphylaxis in patients with a past history of allergies.
7. Always consider the possibility of depression in a postpartum woman presenting with undifferentiated illness or

problems in coping with the baby.

8. Always consider ectopic pregnancy in any woman of child-bearing age presenting with acute abdominal pain.
9. If a patient is found cyanosed always consider upper airway obstruction first.
10. Beware of the asthmatic who is cyanosed with a ‘silent chest’ and tachycardia.
11. Consider ventricular fibrillation or other arrhythmia foremost in an adult with sudden collapse or dizziness.
12. The sudden onset of severe headache adds up to subarachnoid haemorrhage.

IMPORTANT MEDICAL EMERGENCIES IN ADULTS

This section includes summarised protocols for management of emergencies.

Acute anaphylaxis and anaphylactic reactions

Common causes: bee stings, wasp stings, other bites, e.g. Red Imported Fire Ants, parenteral antibiotics, especially penicillin, food reactions e.g. peanuts, fish.

Other causes: allergic extracts, blood products, antivenom, radiological contact ma-terials, anaesthetic agents.

Note: The early danger symptom is itching of the palms of hands and soles of the feet.

Differential diagnosis: syncope

Refer to the adult anaphylaxis kit on page 1244.

First-line treatment (adults)

- Oxygen 6 - 8 L/min (by face mask)
- Adrenaline 0.3 - 0.5 mg (1:1000) IM (if more severe) best given in upper body (e.g. deltoid) (mg = mL of 1:1000 adrenaline)
- If no rapid improvement
    - adrenaline 1 in 10 000: (dilute 1 mL 1:1000 in 9 mL saline)
    - 5 - 10 mL IV over 2 - 5 minutes
- Repeat adrenaline every 5 - 10 minutes as necessary (IM or IV)
- Set up additional IV line (preferably two ‘wide bore’ lines) and infuse colloid solution, e.g. Haemaccel (500 mL → 1 L) or (preferably) crystalloid solution, e.g. N saline (1.5 L → 3 L) 1 part colloid = 3 parts crystalloid (by volume)
- Salbutamol aerosol inhalation (or nebulisation if severe), especially if wheeze/stridor
- Promethazine 10 mg IV slowly (or 25 mg IM) *plus* cimetidine 300 mg IV or ranitidine 50 mg IV
- If very severe add glucagon 1 mg IM or IV
- Admit to hospital (observe at least 4 hours)
- Discharge on diphenhydramine 25 mg qid or promethazine 25 mg tds + prednisolone 50 mg 1 day for 3 days



Figure 130.1 Acute anaphylaxis with respiratory difficulty caused by a European wasp sting

If not responding

- Continue adrenaline every 5 - 10 minutes
- Hydrocortisone 500 mg IV (takes 3 hours to be effective)
- Establish airway (oral airway or endotracheal intubation) if required

Angioedema and acute urticaria

Acute urticaria and angioedema are essentially anaphylaxis limited to the skin, subcutaneous tissues and other specific organs. They can occur together.

Treatment

- Uncomplicated cutaneous swelling - antihistamines e.g. diphenhydramine or promethazine 50 mg (o) or 25 mg IM if more severe
- Upper respiratory involvement
  - adrenaline 0.3 mg SC or IM
  - antihistamine IM

Acute pulmonary oedema

- Keep the patient propped up in bed
- Oxygen by mask or intranasally (8 L/min)
- Insert IV line (large bore canula)
- Frusemide 40 mg IV, increasing to 80 mg IV as necessary (or twice normal oral dose)
- Morphine 1 mg/min IV (slowly up to 5 - 10 mg) + metoclopramide 10 mg IV - usually reserved for chest pain
- Glyceryl trinitrate (nitroglycerin) 300 - 600 ?g sublingual; can use IV nitrates in preference to morphine (if BP > 100 mm Hg)
- CPAP (continuous positive airway pressure) - for unresponsive cases or BiPAP (if available)
- Venesection (if desperate)

Give digoxin if rapid atrial fibrillation and patient not already taking it.

An intravenous infusion of glyceryl trinitrate is superseding morphine use in hospitals.

Note: Keep in mind underlying cause:

e.g. myocardial infarction (? silent)

arrhythmia

cardiomyopathy

anaemia

Severe asthma

Severe asthma is a life-threatening condition that is resistant to standard treatment. It requires intensive medication because of marked obstruction to the air passages, due to severe smooth muscle spasm and in-flammation, producing mucosal oedema and mucous impaction.

Initial treatment

- Oxygen 8 L/min by mask
- Salbutamol 12 puffs (adult) in a spacer or
- Continuous nebulised 0.5% salbutamol by face mask, using compressed (8 L/min) oxygen for nebulisation

Note: Can use salbutamol with ipratropium (0.025%) - 2 mL of each with 4 mL saline, usually for second nebulisation.

- Insert IV line
- Hydrocortisone 4 mg/kg (e.g. 200 - 250 mg) IV statim or prednisolone 1 mg 1 kg(0) then daily
- If still severe, adrenaline 0.3 - 0.5 mg IM

If no response in 30 minutes (or deter-ioration):

- chest X-ray to exclude complications
- arterial blood gases/pulse oximetry
- IV administration of adrenaline 1:10 000 IV (1 mL over 30 seconds) if on monitor then IV infusion of salbutamol 7.5 ?g/kg/h hydrocortisone

If not responding, exhausted and moribund:

- intubation with intermittent positive pressure ventilation (IPPV)
- hydration with IV fluids

Consider isoflurane or halothane inhalation to ‘break’ bronchospasm.

Opioid respiratory depression

- Attend to airway (e.g. pocket mask and bag)
- naloxone 0.2 - 0.4 mg IV and IM

Beware of recurrence of respiratory depression or neurogenic pulmonary oedema from excess naloxone.

Hypoglycaemia

- 50% dextrose 20 - 50 mL IV (if IV line difficult, administer rectally by pressing the nozzle of a large syringe into the anus and injecting slowly) or
- glucagon 1 mL IM or IV (most practical option) then oral glucose

Myocardial infarction/unstable angina

Refer to page 426.

First-line management

- arrange ambulance and hospitalisation
- oxygen (face mask) 6 L/min
- insert IV line
- glyceryl trinitrate (nitroglycerin) 300 ?g ($^{1/2}$ tab) SL, or spray if BP >100 mm Hg
- aspirin 300 mg ($^{1/2 \text{ or}}$1 tab)
- morphine 1 mg per minute IV until pain relief (up to 15 mg) - usually 2 - 10 mg ± metoclopramide 10 mg IV for vomiting

Note: Avoid IM injections.

- ECG

Hyperventilation

Rebreathe slowly from a paper (not plastic) bag

or

Into cupped hands

Status epilepticus and serial seizures

Status epilepticus = repeated convulsions without regaining consciousness after initial tonic - clonic seizure

Serial seizures = repeated convulsions after regaining consciousness

Management

- lie patient on side
- Ensure adequate oxygenation: attend to airway (e.g. Guedel tube); give oxygen 8 L/min (check blood sugar)

If persisting > 5 mmules.

- diazepam 0.05 mg/kg/minute IV until the seizures cease or respiratory depression begins (beware of respiratory depression and other vital parameters): usually 10 - 20 mg bolus in adult or
- clonazepam 1 - 2 mg IV statim then 0.5 - 1 mg/min IV until seizures cease or respiratory depression begins or
- midazolam 0.05 - 0.1 mg/kg IV (max. 10 - 15 mg) or 0.15 mg/kg IM (can be given intranasally or orally by drops into the side of the moull in children 0.3 - 0.5 mg/kg/dose)

Followed by

- phenytoin 1000 mg IV over 20 - 30 min

Other drugs to consider

- Phenobarbitone, thiopentone, paraldehyde, valproate (can use rectally)

BITES AND STINGS

Bites and stings from animals, spiders and insects are commonplace in Australia and the United States but fatal bites are uncommon.

Snake bites

Snake bites are more common and severe in those handling snakes and in those trying to kill the snake. Snakes are more aggressive when mating or sloughing their skin (about four times a year). They strike for one-third of their length at 3.5 metres/second. Over 70% of bites are on the legs. Sea snakes are not a major problem in Australian waters.

First aid

1. Keep the patient as still as possible.
2. Do not wash, cut, manipulate the wound, apply ice or use a tourniquet.
3. Immediately bandage the bite site very firmly (not too tight). A 15 cm crepe bandage is ideal: it should extend above the bite site for 15 cm (e.g. if bitten around the ankle the bandage should cover the leg to the knee).
4. Splint the limb to immobilise it: a firm stick or slab of wood would be ideal.
5. Transport to a medical facility for definitive treatment. Do not give alcoholic beverages or stimulants.
6. If possible, the dead snake should be brought along.

Note: A venom detection kit is used to examine a swab of the bitten area or a fresh urine specimen (the best) or blood.

The bandage can be removed when the patient is safely under medical observation. Observe for symptoms and signs of envenomation.

Envenomation

Not all patients become envenomated and the antivenom should not be given unless there is evidence of this. Envenomation is more likely when the snake has a clear bite, such as in snake handlers or barefooted people or hands placed in burrows.

Important early symptoms of snake bite envenomation include:

- nausea and vomiting (a reliable early symptom)
- abdominal pain
- excessive perspiration
- severe headache
- blurred vision
- difficulty speaking or swallowing
- coagulation defects (e.g. haematuria)
- tender lymphadenopathy

The greatest danger is respiratory obstruction and failure.

Refer to Figure 130.2 for detailed effects.


Symptoms of envenomation
• vomiting/nausea
• haematemesis
• headache
• coughing or spitting blood
• sweating
• muscle pain
abdominal pain
pain in groin
tender lymphadenopathy
haematuria
CNS
• stuporose
• loss of consciousness
or
• neurotoxic effects (esp. cranial nerves)
– ptosis
– blurred, double vision
– difficulty swallowing and speaking
– central voluntary muscle weakness
Bite site
• one or paired puncture mark
(may not be visible)
• ± pain, swelling
or local reaction
• continued bleeding

Figure 130.2 How to recognise snake bite envenomation

Investigations and observation

- Venom detection kit: wound site (best) or urine
- Careful observations (e.g. vital signs, conscious state)
- Test all urine for blood and protein
- Watch for coagulopathy (e.g. spitting and coughing blood, bleeding from wounds/IV site, haematuria)
- Serial whole blood clotting time (a plain glass tube): normal < 5 - 8 min, > 15 min significant
- Coagulation screen

Treatment of envenomation

- Reassure patient at all times
- Set up a slow IV infusion of N saline
- Give SC antihistamine cover (5 min beforehand): keep adrenaline on standby
- Dilute the specific antivenom (1 in 10 in N saline) and infuse slowly over 30 min via the tubing of the saline solution
- Have adrenaline, oxygen, antihistamine and steroids on standby
- Monitor vital signs
- Provide basic life support as necessary

Note 1: The use of prophylactic adrenaline is controversial and some authorities reserve it for a reaction to the antivenom. It is best avoided with brown snake envenomation and with coagulopathy.

Note 2: Do not give antivenom unless clinical signs of envenomation or biochemical signs (e.g. positive urine, or abnormal clotting profile).

Note 3: 1 ampoule may be sufficient but 3 or more may be needed, especially if coagulopathy.

Spider bites

The toxin of most species of spider causes only localised pain, redness and swelling, but the toxin of some, notably the deadly Sydney funnel-web spider (Atrax robustus), can be rapidly fatal.

First aid

- Sydney funnel-web: as for snake bites.
- Other spiders: apply an ice pack, do not bandage.

Treatment

1. Sydney funnel-web

Signs of envenomation (in order):

- muscle fasciculation - limb ? tongue/lip
- marked salivation or lacrimation
- piloerection
- dyspnoea
- neurological symptoms (e.g. disorientation, coma)

Treatment:

- specific antivenom (usually 4 - 8 vials)
- resuscitation and other supportive measures

2. Other spider bites

The toxins of most species of spiders cause only localised symptoms but the venom of a selected few, namely the red-back spider of Australia (Latrodectus mactans hasseltii) and its related black widow spider (Latrodectus mactus), can cause envenomation. This is rarely fatal but is more serious in the young, the frail and the elderly. The bite wounds are prone to infection.

Treatment of envenomation (rarely needed):

- give antihistamine (e.g. IM)
- antivenom IM injection 15 minutes later

Bee stings

First aid

1. Scrape the sting off sideways with a fingernail or knife blade. Do not squeeze it with the fingertips.
2. Apply 20% aluminium sulphate solution (Stingose) or methylated spirits.
3. Apply ice to the site.
4. Rest and elevate the limb that has been stung.

If anaphylaxis, treat as outlined earlier.

Preventive measures (if hypersensitive)

- Avoid bees (and wasps) if possible.
- Immunotherapy to honey bee (or wasp) venom. There is no cross-allergy between the honey bee, the ‘yellow jacket wasp’ and the paper wasp. Specific hyposensitisation against the Vespula species is required. For the bee use pure venom antigen.
- Immunotherapy should be offered to those:
  - with a history of asthma who have had a single severe reaction to a bee sting
  - who have had a minimum of three stings with serial crescendo reactions

  - occupationally exposed who manifest severe reactions
  - with elevated venom-specific IgE (RAST) antibodies, or positive venom prick tests

Centipede and scorpion bites

The main symptom is pain, which can be very severe and prolonged.

First aid

1. Apply local heat (e.g. hot water with ammonia - household bleach)
2. Clean site.
3. Local anaesthetic (e.g. 1 - 2 mL of 1% lignocaine) infiltrated around the site.
4. Consider opioids for pain
5. Consider promethazine injection
6. Check tetanus immunisation status.

Box jellyfish or sea wasp (Chironex fleckeri)

This is the most dangerous jellyfish in Australian waters and has been responsible for at least 80 extremely painful and sudden deaths.[3] Death can occur in minutes due to cardiopulmonary failure. There can be up to 180 metres of stinging tentacles. The sting gives a 'frosted ladder' appearance. The jellyfish is limited to tropical waters north of the Tropic of Capricorn (Exmouth in west to Gladstone in east) and is found in coastal waters during the summer.

Prevention

- Avoid swimming, paddling and wading in 'jellyfish alert' areas in unsafe months.
- Otherwise, use a 'stinger suit'.

Treatment[4]

- The victim should be removed from the water to prevent drowning.
- Inactivate the tentacles by pouring vinegar over them for 30 seconds (do not use alcohol) - use up to 2 L of vinegar at a time.
- Use a cold pack for small stings and ice massage for large areas.
- Check respiration and the pulse.
- Start immediate cardiopulmonary resuscitation if necessary.
- Gain IV access and use colloid; give oxygen and inotropes if necessary.
- Give box jellyfish antivenom IM or by IV injection for major stings (may need several ounpoules).
- Provide pain relief if required (ice, lignocaine and analgesics).

Note: A delayed reaction can occur - the stings can cause pain after weeks (oral steroids are used).

Irukandji syndrome[4]

This is caused by Carukia barnesi, a tiny box jellyfish that can penetrate safety nets for Chironex, and possibly other species of box sellfish. Initially a mild sting with a delayed severe syndrome (usually after 30 minutes):

- severe low back pain and muscle cramps
- chest pain, sweating and anxiety
- anxiety, restlessness, 'impending doom' feeling
- intense headache, vomiting

There is no specific first aid or antivenom but resuscitation measures may be needed as death can occur from pulmonary oedema or cardiac arrest. Measures include morphine 5 mg IV every 5 minutes as required, hypertensive control (e.g. phentolamine, CPAP and oxygen for pulmonary oedema) and possibly IV magnesium.

Stinging fish

The sharp spines of the stinging fish have venom glands which can produce severe pain if they spike or even graze the skin. The best known of these is the stonefish. Others include bullrout, catfish, sea-urchins and crown of thorns (may need an X-ray). The toxin is usually heat-sensitive. Stingrays cause a gash wound with possible superinfection.

Envenomation

- Intense pain
- Localised swelling
- Bluish discolouration

Treatment

- Clean the wound - consider exploration.
- Bathe or immerse the affected part in very warm to hot (not scalding, 43?) water - this may give instant relief.[3]
- Give simple analgesics.
- If pain persists, give a local injection/infiltration of lignocaine 1% or even a regional block. If still persisting, try pyridoxine 50 mg intralesional injection.
- A specific antivenom is available for the sting of the stonefish. Can be given IM or IV.

Mollusc bite (blue-ringed octopus, cone shell)

Mollusc venoms can be rapidly fatal because of prolonged muscular weakness leading to respiratory paralysis.

Treatment

- Compression bandage to bite site (usually hand/arm)
- Immobilise the limb
- Arrange transport (preferably by ambulance) to a medical facility
- Observe (and manage) for respiratory paralysis - ensure adequate ABC

Sandfly bites

For some reason, possibly the nature of body odour, the use of oral thiamine may prevent sandfly bites.

Dose: thiamine 100 mg orally, daily

Other bites and stings

This includes bites from ants, wasps and some jellyfish.

First aid

1. Wash the site with large quantities of cool water.
2. Apply vinegar (liberal amount) or aluminium sulphate 20% solution (Stingose) to the wound for about 30 seconds.
3. Apply ice for several minutes.
4. Use soothing anti-itch cream or 5% lignocaine cream or ointment if very painful.

Medication is not usually necessary unless an acute allergic reaction develops.

The embedded tick

Some species of ticks can be very dangerous to human beings, especially to children. Be careful in children 1 - 5 years - they are usually found in the scalp behind the ears. If they attach themselves to the head or neck, usually behind the ear, a serious problem exists. As it is impossible to distinguish between dangerous and non-dangerous ticks, early removal is mandatory. The tick should be totally removed - and the mouthparts of the tick must not be left behind. Do not attempt to grab the tick by its body and tug. This is rarely successful in dislodging the tick, and more toxin is thereby injected into the host.

As an office procedure, many practitioners grasp the tick's head as close to the skin as possible with fine forceps or tweezers, and pull the tick out sideways with a sharp rotatory action. This is acceptable, but not as effective as the methods described here.

First aid outdoor removal method

- Saturate the tick with petrol or kerosene and leave for 3 minutes.
- Loop a strong thin thread around the tick's head as close to the skin as possible, and pull sharply with a twisting motion.
- A pyrethrin-based spray is often used.

Office procedure

- Infiltrate a small amount of local anaesthetic into the skin around the site of embedment.
- With a number 11 or 15 scalpel blade, make the necessary very small excision, including the mouthparts of the tick, to ensure total removal (Figure 130.3).



Figure 130.3 Removing the embedded tick

- The small defect can usually be closed with a bandaid (or Steri-strips).
- Careful observation is needed after removal.

Human bites and animal bites

These bites can cause problems of suppurative infection and management in general. They are outlined in Chapter 134.

A POT POURRI OF EMERGENCY CALLS

Electric shock

Useful facts

- Direct current (DC) from welding machines or lightning produces more electrolyte tissue damage and burns than AC (domestic supply).
- Injuries occur at sites distant from entry or exit.
- Severe muscle contractions can cause bone fracture.
- Household shocks tend to cause cardiac arrest (ventricular fibrillation) and myocardial damage is common.

- Ischaemic necrosis of a limb is a common effect.
- Apparently minor initial injuries may be very misleading (Figure 130.4).



Figure 130.4 Effect of electric shock passing through the body

- Neurological deficits and psychoneurotic sequelae are common in survivors.

Principles of management

- Make the site safe: switch off the electricity. Use dry wool to insulate rescuers.
- 'Treat the clinically dead'.
- Attend to ABC.
- Give a precordial thump in a witnessed arrest.
- Consider a cervical collar (? cervical fracture).
- Provide basic cardiopulmonary resuscitation, including defibrillation (as required).
- Give a lignocaine infusion (100 mg IV) after cardiac arrest.[5]
- Investigate and consider:
    - careful examination of all limbs
    - X-ray of limbs or spine as appropriate
    - check for myoglobinuria and renal failure
    - give tetanus and clostridial prophylaxis
- Get expert help - intensive care unit, burns unit.

Lightning strike

Prevention (during an electrical storm):

- Don't shelter under trees or tall object (splash phenomenon - Figure 130.5).[6]



Figure 130.5 ‘Splash effect’ where current is reflected from tree

- Stay indoors - shelter in a building or closed car.
- Avoid using telephone.
- Avoid holding metal objects (e.g. golf clubs).
- Keep as low to ground as possible (e.g. curl up in a ditch).
- Avoid being in a group.

Clinical effects:

- Burn injury (90%): the ‘flashover’ phenomenon - clothing disintegrates
- Blast injury (e.g. ruptured spleen, subdural, ruptured eardrum)
- Electrical injury as for houshold shock (uncommon)

Petrol and solvent sniffing

Substances commonly sniffed include petrol, glues, spirit-based paints, paint spray cons and other aerosols.

The three main acute problems:

1. Shaking and fitting: give diazepam IV (as for convulsions). May require paraldehyde IM.
2. Agitation/aggressive behaviour self-haven: try to calm patient in a well-lit room. Give sedation with diazepam. Give haloperidol for hallucinations or delusions.
3. General debilitation: this may include acute infections (e.g. chest infection) or anaemia. Investigation and referral for breaking the habit is necessary.

Near drowning

Near drowning is survival after asphyxia from submersion in a liquid medium.

The rule to remember is that victims can respond to resuscitation after considerable im-mersion time (up to 30 minutes) and that mouth-to-mouth resuscitation should always be attempted even if pulseless or with fixed dilated pupils. The usual routine of basic life support and CPR apply. All symptomatic patients should receive high-flow oxygen and ideally CPAP or BiPAP. Those requiring intubation should receive positive end expiratory pressure (PEEP).

Artificial surfactant given via an endo-tracheal tube has been used successfully in the United Kingdom.

There is no significant difference in outcome and management between salt water and fresh water drowning. Hypothermia should be attended to with warming, such as a hot-air blanket if available and warm fluids.

Epistaxis

Refer to pp. 636 - 7.

Opioid (heroin) overdose

Refer to pp. 1344.

Migraine

Refer to pp. 605 - 6.

VITAL EMERGENCY SKILLS

Cardiopulmonary resuscitation

Cardiopulmonary arrest (CPA). It is essential that all doctors are familiar with the protocol for instituting basic life support in such an eventuality. Sick patients do visit our offices daily and the potential for sudden collapse, including a cardiac arrest, is ever present. About 75% of arrests are due to ventricular fibrillation and more than 75% of victims have severe coronary artery disease.[7] After 3 minutes of CPA (unconsciousness, no pulse, no respiration) there is an increasing risk of permanent cerebral dysfunction.

Important causes of sudden death are outlined in Table 130.1.[8]

Table 130.1 Causes of sudden death

Cardiac arrhythmias
ventricular fibrillation (75%)
ventricular tachycardia
torsade de pointes VT (? drugs)
sick sinus syndrome
severe bradycardia
Sudden pump failure
acute myocardial infarction
cardiomyopathy
Cardiovascular rupture
myocardial rupture
dissecting aneurysm aorta
subarachnoid haemorrhage
Acute circulatory obstruction
pulmonary embolism
Others include
pulmonary hypertension

mitral valve prolapse

electrolyte abnormalities

glue sniffing

The ABC basic life support for cardiac arrest should be followed, but ideally DRABC is best (defibrillation first if a defibrillator is available - the outcome appears to be directly related to the speed of defibrillation).

Basic life support

The following represents a logical ABC plan for the adult patient who collapses or is found apparently unconscious.

1. Shake and shout at the patient.
2. Check breathing.
3. Check pulse (feel carotid adjacent to thyroid cartilage).
4. Call for help (if no pulse).
5. Finger sweep oropharynx (clear it).
6. Place victim on back on firm surface.
7. Thump precordium (if arrest witnessed).
8. Tilt head back (to maximum).
9. Lift chin (use airway if available).
10. Commence basic life support:
    - rescue breaths (RBs) - 2 strong breaths
    - external chest compression (no pause) - 30 compressions
    - continue alternadug 30 compressions (at rate if 100 beats/min) with 2 strong breaths

Note: The ratio of 30:2 with one or two rescuers, especially with the inexperienced, is throw favoured.

The basic schedule for cardiopulmonary resuscitation is presented in Table 130.2 and a flow chart for basic life support in Figure 130.6.

Table 130.2 Basic schedule for CPR

	Infant < 1 year	**Child**	**Adult**
Compression (rate per min)	100	80 - 100 (large) (small)	80 - 100
Depth of compression (cm)	2 - 3	3 - 4	4 - 5
Position of compression	Centre sternum	Centre sternum	4 cm above xiphisternum
Method	2 fingers	1 hand	2 hands
Ventilation (rate per minute)	20	16	12
Head tilt	Nil	Mid	Full


Collapse
Conscious
Unconscious
• make comfortable
• observe
• organise ECG
– blood sugar
– other tests
• lateral position
• clear airway
• check breathing
yes
no
• supine position
• head tilt/jaw lift
2 expired air inflations
(check pulse)
• continue RB
yes
no
• check breathing
and pulse for
5 seconds every
2 minutes
RB
ECC
30 compressions
2 breaths
(check breathing and pulse
for 5 seconds every 2 minutes)

Figure 130.6 Algorithm for basic life support: CPR in the collapsed patient

Method of rescue breathing/ventilation

With the victim's head in the ‘sniffing the morning air’ position (head totally tilted back and chin pulled forward) the rescuer takes a deep breath and seals his or her lips around the mouth or nose of the victim. Pinch the victim's nose if using mouth-to-mouth resuscitation. Two full puffs are given within a few seconds (Figs 130.7 and Figs 130.8). Rescue breathing/ventilation is the only method of artificial respiration that successfully ventilates the patient.[7] If the chest does not move easily, obstruction is present. If available a sucker should be used to clear the oropharynx. Firmly fitting dentures should be left in place as they make artificial respiration easier. A resuscitube or Laerdal pocket mask (which should be in the doctor's bag) is ideal for EAR and saves mouth-to-mouth contact.



For optimal airway patency:

1 Clear foreign matter — from mouth: use finger sweep; from airway: blow between shoulder blades; consider a Heimlich manoeuvre

2 Lay patient supine on flat, firm surface (A). Note how the soft tissue of the pharynx can obstruct the airway by falling backwards.

3 In order to overcome this, apply a head lift (B) plus a chin lift (C) or jaw thrust manoeuvre. (Note: avoid excessive movement of the neck if spinal injury is suspected, but clearing the airway has first priority.)

Slight flexion of the neck with small cushion

Figure 130.7 Basic life support: A = Airway

Rescue breathing:

1 Five full breaths within 10 seconds.
2 Observe rise of chest, not of abdomen.
3 Look, listen and feel for exhalation.
4 Check the carotid pulse.
5 If no pulse, commence full cardiopulmonary resuscitation



Figure 130.8 Basic life support: B = Breathing

External chest compression

Evidence supports the need to maintain cardiac compression following a cardiac arrest.

Compressions are safely performed by finding the xiphisternal notch then placing the broad heel of one hand over the lower half of the sternum (in adults) with the heel of the second hand placed over the first with the fingers interlocked. Remember to keep the arms and elbows straight as the sternum is rhythmically depressed for 4 - 5 cm. Try to keep to this position as ‘wandering’ causes fractured ribs or worse. The fingers must be kept off the chest.

Maintain the compression for 0.5 second, then relax - compressions should be smooth, regular and uninterrupted.

Check pupil size and reaction to light and the carotid or femoral pulse. The compressions should ideally produce an impulse in the femoral pulse. Another person can check for carotid or femoral pulsation during CPR (Figure 130.9).



1 If carotid or femoral pulse is absent, immediately commence external cardiac compression
2 See Figure 130.6 for rhythm.

External cardiac compression with fingers locked (A), and with fingers extended (B). Heel of one hand placed on lower sternum 2 finger breadths above the xiphoid sternal junction. Heel of second hand placed on first. Ensure fingers don't exert pressure. The patient should be lying on a firm surface, the operator level with shoulder.

Figure 130.9 Basic life support: C = Circulation

Maintenance of CPR

Consider ceasing CPR if there is no im-provement in 30 minutes. Exceptions where prolonged CPR can be successful are cold water drowning, marine envenomation, snake bite, and certain poisonings (e.g. cyanide and organophosphate).

Note: Ventilation of the patient in the first minutes is vital to success, especially if a bag and mask are available to perform this. It must not be neglected during basic or advanced cardiorespiratory life support.

Advanced cardiac life support

Advanced life support depends on the availability of skilled personnel and appropriate equipment.

Optimal initial support involves:

- endotracheal intubation (otherwise bag and oxygen)
- ECG monitoring
- intravenous access (large peripheral or central vein)

Optimal initial therapy involves:

- defibrillation
- oxygen
- cardioactive drugs, especially adrenaline

If an ECG recording is unavailable the best course of action is:

- defibrillation if unsuccessful
- adrenaline IV

Advanced life support: Defibrillation

- Minimise interruptions to chest compressions.
- Give a ingle shock instead of stacked shocks (single shock strategy) for ventricular fibrillation/pulseless ventricular

tachycardia.

- Where the arrest is witnessed by a health care professional and a manual defibrillator is available, then up to three shocks may be given (stacked shock strategy) at the first defibrillation attempt.
- Monophasic defibrillation - all shocks 360 joules.
- Bipasic defibrillation - where specific devices have been identified to be efficacious at other energy levels these should be used if known. However, where the health care professional is unsure of the energy level recommended for a specific device, a default energy level of 200#J should be used without delay.
- After each defibrillation attempt give 2 minutes of CPR before checking rhythm and pulse.

For defibrillation, two paddles should be placed correctly on the chest wall, using one of two positions:

- one to right of upper sternum and the other over the apex of the heart (Figure 130.10)



Figure 130.10 Standard position of two paddles for defibrillation

- one over anterior wall of chest and the other under tip of left scapula

Hairs on the chest should be shaved to accommodate the paddles.

A protocol for advanced cardiac life support is presented in Figure 130.10.[10]

Urgent intravenous access

It is preferable to aim for transcutaneous cannulation of veins initially so a peripheral line should be introduced into a vein in the cubital fossa. Several lines may be required with massive blood loss.

Alternative routes:

- central venous cannulation: most doctors should be able to cannulate the external jugular vein with a standard cannula
- peripheral venous cutdown
- intraosseous infusion (see pp. 940 - 1)

Urgent intravenous cutdown

In emergencies, especially those due to acute blood loss, transcutaneous cannulation for the infusion of fluids or transfusion of blood can be difficult. For the short-term situation, a surgical cutdown into the long saphenous vein at the ankle or the cephalic vein at the wrist is life-saving. Ideally, the long saphenous vein should be used in children.

Surface anatomy

Long saphenous vein. The vein lies at the anterior tip of the medial malleolus. The best site for incision is centred about 2 cm above and 2 cm anterior to the most prominent medial bony eminence (Figure 130.11a).


(a)
site of incision
long saphenous vein
2 cm
2 cm
prominence of medial malleous
(b)
cephalic vein
site of incision
styloid process of distal radius
2-3 cm
(c)
vein elevator
distal ligature
proximal ligature
catheter
vein

Figure 130.11 Urgent intravenous cutdown: (a) site of incision over long saphenous vein (medial perspective); (b) site of incision over cephalic vein at wrist (radial or lateral perspective); (c) method of introduction of catheter into vein

Cephalic vein. The cephalic vein 'bisects' the bony eminences of the distal end of the radius as it winds around the radius from the dorsum of the hand to the anterior surface of the forearm. The incision site is 2 - 3 cm above the tip of the radial styloid (Figure 130.11b).

Method of cutdown

After fitting gloves and using a skin preparation:

1. Make a 1.5 - 2 cm transverse skin incision over the vein.
2. Locate the vein by blunt dissection (do not confuse the vein with the pearly white tendons).
3. Loop an aneurysm needle or fine curved artery forceps under and around the vein.
4. Place the ligature around the distal vein and use this to steady the vein.
5. Place a loose knotted ligature over the proximal end of the vein.
6. Incise the vein transversely with a small lancet or scissors or by a carefully controlled stab with a scalpel.
7. Use a vein elevator (if available) for the best possible access to the vein.
8. Insert a catheter (Figure 130.11c); use 14 or 16 gauge.
9. Gently tie the proximal vein to the catheter.

10. After connecting to the intravenous set and checking the flow of fluid, close the wound with a suitable suture material.

REFERENCES

1. Murtagh J. The anatomy of a rural practice. Aust Fam Physician, 1981; 10: 564 - 7.
2. Hogan C. The management of emergencies in general practice. Aust Fam Physician, 1989; 19: 1211 - 19.
3. Sutherland SK. Venomous bites and stings. In: MIMS Disease Index (2nd edn). Sydney: IMS Publishing, 1996: 563 - 7.
4. Fenner P. Marine bites and stings. Today, 2002; 3(1): 26 - 31.
5. Walpole BG. Electric shock. Aust Fam Physician, 1989; 18: 1252 - 8.
6. Crocker B, Thomson S. Lightning injuries. Patient Management, 1991; November: 51 - 5.
7. Kumar PJ, Clarke MC. Clinical Medicine (5th edn). London: Bailliere Tindall, 2000: 746.
8. Tierney LM et al. Current Medical Diagnosis and Treatment (45th edn). New York: McGraw Hill and Lange, 2006: 380 - 1.
9. Jacobs IG (Chairman). # New changes to resuscitation guidelines. Australian Resusation Cornal 2006 < http://www.resus.org.au >
10. American Heart Association in collaboration with International Liaison Committee on Resuscitation. Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care: International Consensus on Science, Part 3: adult basic life support. Resuscitation, 2000; 46(1 - 3): 29 - 71.

CHAPTER 131. THE DOCTOR'S BAG AND OTHER EMERGENCY EQUIPMENT

Almost everyone who goes to bed counts upon a full night's rest: like a picket at the outposts, the doctor must be ever on call.

KARL F. MARX (1796 - 1877)

THE DOCTOR'S BAG AND OTHER EMERGENCY EQUIPMENT

General practitioners who perform home visits and nursing home visits require the traditional doctor's bag that includes the basic tools of trade, drugs (including those for emergency use), stationery and various miscellaneous items. Country doctors will by necessity use their bag for more emergency home and roadside calls.1 These recommended contents are simply a guide for cross-checking.

ESSENTIAL REQUIREMENTS FOR THE BAG

- Sturdiness
- Lockable (e.g. combination lock)
- Ready interior access
- Uncluttered
- Disposable single-use items
- Light, portable equipment
- Regular checks to ensure non-expired drugs
- Storage in a cool place (not boot of car)

Stationery (checklist)

- Practice letterhead and envelopes
- Prescription pads
- Hospital admission forms
- Sickness/off -work certificates
- X-ray, pathology referral forms
- Accounting forms
- Dangerous drugs record books
- Continuation notes
- Large adhesive labels to record visit (attach later to patient's history)
- Tie-on labels for emergencies
- Recommendation forms (to psychiatric/mental hospitals)
- Pens

Miscellaneous items

- Quick reference cards
  - the doctor's bag checklist[1, 2, 3]
  - dosage details of drugs, all age groups
  - important telephone numbers
- Local map
- Phonecard or coins for public telephone use
- Handbook of emergency medicine

Equipment

- Sphygmomanometer (aneroid)
- Stethoscope
- Diagnostic set (auriscope + ophthalmoscope)
- Tongue depressors
- Tourniquet
- Small needle disposal bottle
- Scissors
- Syringes 2, 5, 10 mL
- Needles 19, 21, 23, 25 gauge
- Scalp veins (butterfly) needles
- IV cannulae 16, 18, 20 gauge
- Alcohol swabs
- Micropore tape
- Thermometer
- ‘Spacer’ (e.g. Volumatic, for asthma)
- Artery forceps
- Urine testing sticks
- Pathology specimen bottles
- Skin swabs, throat swabs
- Torch
- Patellar hammer
- Oral airway (e.g. Revivatube, Resuscitube - Figure 131.1, Guedel)
- Scalpel (disposable)
- File (for glass ampoules)
- Examination glove



Figure 131.1 The two-way Resuscitube

Drugs

Drugs (oral)

- Samples of commonly used:
    - analgesics
    - antibiotics
    - antidiarrhoeal agents
    - antiemetics
    - antihistamines
    - sedatives
- Glyceryl trinitrate (nitroglycerin)
- Soluble aspirin (for myocardial infarction)
- Sumatriptan (Imigran)

Drugs (sprays)

- Glyceryl trinitrate
- Salbutamol aerosol (see Spacer)

Drugs (topical)

- Anaesthetic eyedrops

Drugs (injectable)

Refer to Tables 131.1 and 131.2

Table 131.1 Doctor's bag - ideal drugs[5]

Drug	Presentation	Indications
Adrenaline *	1 mg/mL (1:1000)	hypersensitivity reactions and anaphylactic shock; bronchial asthma†; ventricular asystole; croup; ventricular fibrillation (to assist CPR)
Atropine sulphate	0.6 mg/1 mL	bradycardia (after myocardial infarction); 2nd or 3rd degree heart block; ureteric colic†; organophosphate poisoning
Benztropine mesylate *	Cogentin 2 mg/2 mL	acute dystonic reactions
Benzylpenicillin *	3 g with 10 mL	water meningococcaemia, pneumonia (adults)
Dexamethasone	4 mg/1 mL	severe asthma (esp. elderly); moderate to severe croup, anaphylaxis, acute Addisonian crisis
Diazepam *	10 mg/2 mL	status epilepticus and other convulsions such as eclampsia; sedation in acute anxiety and severe tension headache; psychiatric emergencies including acute alcohol withdrawal
Dihydroergotamine	1 mg/mL	severe migraine; cluster headache
Ergometrine maleate	0.25 mg/1 mL	uterine bleeding; abortion or postpartum haemorrhage
Frusemide *	Lasix 20 mg/2 mL	left ventricular failure; acute pulmonary oedema
Glucagon *	1 mg + 1 mL	solvent hypoglycaemia (insulin or oral therapy)
Haloperidol *	Serenace 5 mg/1 mL	psychiatric emergencies such as severe agitation, psychoses; migraine
Hydrocortisone sodium succinate *	Solu-cortef 100 mg/2 mL 250 mg/2 L	anaphylaxis; severe asthma; Addisonian crisis; thyrotoxic crisis; acute allergies

Metoclopramide [*] or Prochlorperazine	Maxolon 10 mg/2 mL Stemetil 12.5 mg/mL	severe vomiting (e.g. Meniere's syndrome, gastritis); acute labyrinthitis; migraine
Morphine sulphate [*]	15 mg/1 mL	acute pulmonary oedema; relief of severe pain (not due to muscular spasm) such as myocardial infarction
Naloxone (more than one ampoule) [*]	Narcan 0.4 mg/mL	opiate respiratory depression
Promethazine [*]	Phenergan 50 mg/2 mL	acute allergic conditions, antiemetic †
Salbutamol [*]	inhaler and/or nebuliser solution	bronchial asthma, other bronchospasm
Tramadol	100 mg/2 mL	moderate to severe pain
Water	5 mL	diluent

[*] essential drugs † May be useful as an alternative drug

All drugs listed are injectable except salbutamol inhaler

Table 131.2 Additional cardiopulmonary drugs (optional)[4, 5]

Injectable drugs	Presentation	Indications
Adenosine	Adenocor 6 mg/2 mL	supraventricular tachycardia
Isoprenaline	Isuprel 1 mg/mL	bradycardia unresponsive to atropine
Metaraminol bitartrate	Aramine 10 mg/mL	non-hypovolaemic shock; anaphylaxis drug-induced, associated with spinal anaesthesia, ? cardiogenic
Terbutaline	Bricanyl 0.5 mg/mL	bronchospasm; asthma; bronchitis; smoke inhalation
Verapamil	Isoptin 5 mg/2 mL	supraventricular tachycardia (with adequate blood pressure)
Heparin	5000 U/mL	thromboembolism, myocardial infarction

Note: The author recommends the MIN-I-JET syringe packs for ideal emergency use. The range includes naloxone 5 mL, aminophylline, atropine, adrenaline, dextrose, lignocaine, isoprenaline, sodium bicarbonate.

A study conducted by Johnston et al.4 of 512 Queensland general practitioners revealed the range of emergency equipment and essential drugs carried with indications of that considered essential. The most common emergencies were acute asthma, psychiatric emergencies, seizures, hypoglycaemia, anaphyl-axis, impaired consciousness, shock, poisoning and overdose. Most GPs stocked 15 or more of the 16 emergency doctor's bag drugs. The drugs used most were adrenaline, benztropine, diazepam, glucagon, haloperidol, hydro-cortisone, naloxone and salbutamol (inhaler).

The issue of what drugs to carry is always an interesting intellectual exercise. It could reduce to just a few items such as adrenaline, diazepam, naloxone and morphine. It is advisable not to carry pethidine as morphine can be used for kidney and biliary colic since it does not aggravate spasm. An alternative injectable analgesic to the traditional opioids is tramadol (available as 100 mg/2 mL injection).

Drugs (suppositories)

- Indomethacin

THE COUNTRY DOCTOR'S BAG

Country doctors, especially in isolated areas, usually carry additional equipment in their motor vehicles when called to the scene of an accident or other emergency. The equipment will vary according to geographic factors, the ambulance service and the special interests and enthusiasm of the practitioner.

Accident kit

The following list represents the contents of an isolated country doctor's kit; it will occupy a standard briefcase only.

- Flashlight and spare batteries
- Sterile compression bandages
- Steri-strips (large and small)
- Plastic container of antiseptic
- Wide-bore needle
- Sterile suture set
- Suture material
- Gillies' forceps
- Small scissors
- Large scissors (for cutting clothing)
- Artery forceps × 2
- Laerdal pocket mask (Figure 131.2)
- Medium-sized torch
- Rigid cervical collar
- Triangular bandages × 2
- Crepe bandages × 2
- Air splints × 2



Figure 131.2 The Laerdal pocket mask

- Disposable scalpel with blade

- Safety pins
- Sterile gauze
- Urinary catheter
- Makeshift hook to suspend IV fluid pack
- IV infusion tubing
- Bottle of Haemaccel × 2
- Fluid pack
- 1 L IV fluid (N saline)

The contents of this bag provide the equipment to cope with common accidents, including:

- bleeding wounds (e.g. arterial bleeders)
- tension pneumothorax
- fractured cervical spine
- fractured limbs
- fractures of the shoulder girdle
- snake bites

In addition, some country doctors carry a trephine to cope with the extradural haematoma.

Resuscitation kit

The country doctor can carry an oxyresuscitator unit with the following standard items:

- oxygen
- suction
- laryngoscopes
- endotracheal tubes
- oropharyngeal airways (e.g. Guedel)
- endotracheal adaptor
- face mask (e.g. Laerdal pocket mask, with one-way valve, or Concorde mask)

The base of the unit contains:

- paediatric tracheostomy tube
- self-retaining tourniquet
- intravenous infusion needle
- intravenous infusion tubing
- haemaccel 500 mL
- disposable scalpel (for intravenous cutdown)
- chromic catgut (for intravenous cutdown)
- mosquito forceps (for intravenous cutdown)
- Magill's forceps

Other equipment that could be carried:

- a balloon resuscitator and sucker + oxygen cylinder (instead of the oxyresuscitator)
- normal saline or Hartman's solution
- sodium bicarbonate (100 mL)
- portable ECG and defibrillator (e.g. Heartstart)

PRECAUTIONS AT THE SCENE OF AN ACCIDENT[5, 6]

- Don't become a casualty yourself.

- Do not speed to the accident.

Medical emergencies in general practice - a management proposal

Note: Using basic equipment and doctor's bag (mainly adult dosages)

Don't forget:

- **Secure IV line**
- **Oxygen**

Acute pulmonary oedema	• frusemide 40 - 80 mg IV (or twice usual dose)
Acute anaphylaxis	• glyceryl trinitrate 1 dose (spray) • consider (especially if chest pain) - morphine 5 mg IV + metoclopramide 10 - 20 mg IV • adrenaline 0.3 - 0.5 mg (1:1000) IM, repeat every 5 minutes as necessary
Angio-oedema and acute urticaria	If no rapid improvement: • salbutamol inhalation • promethazine 10 mg IV • VI fluids • ? hydrocortisone/glucagon • promethazine 25 mg IM
Asthma	• salbutamol 6 (< 6 years) - 12 (adults) puff s by spacer
Croup (severe)	• hydrocortisone 200 mg IV If severe: • adrenaline 0.3 - 0.5 mg 1:1000 IM or SC • dexamethasone 0.15 mg/kg IM
Epilepsy (seizure)	• diazepam 5 - 20 mg IV (rate not exceeding 2 mg/min)
Opiate respiratory depression	• naloxone HCl 0.4 (or 0.2) mg IV + 0.4 mg IM
Myocardial infarction	• aspirin 300 mg soluble tab
Hypoglycaemia	• glyceryl trinitrate spray or tablets (max. 3) • morphine sulphate 2.5 - 5 mg IV + metoclopramide • glucagon 1 mg/mL SC, IM or IV, then sweet drink
Migraine (severe)	• prochlorperazine 12.5 mg IV *or*
Cluster headache	• metoclopramide 10 mg IV ± dihydroergotamine IV or IM *or* • haloperidol 5 mg IM or 1V • 100% oxygen 6 L/min for 15 minutes
Movement disorders (from antipsychotic medication)	• metoclopramide 10 mg IV + dihydroergotamine 0.5 mg IV slowly • benztropine 1 - 2 mg IV or IM
Meningococcaemia	• benzylpenicillin 60 mg/kg IV
Ureteric colic	• morphine 10 - 15 mg IM or IV ± metoclopramide
Vertigo (acute)	• indomethacin suppository • prochlorperazine 12.5 - 25 mg IM or promethazine 25 mg IM
Vomiting	• prochlorperazine 12.5 mg IM or IV or metoclopramide 10 mg

Practice tips

- Check your doctor's bag every month for drugs that may be expired, damaged or in short supply (your practice nurse can do this).
- Replace any used drugs or materials the day after use.
- Always have your bag handy but don't carry it in the car in hot weather. It is best to be able to grab it from a safe accessible spot when you leave for an emergency.
- Security is an issue. Drugs of addiction (pethidine and morphine) may be kept separate and then taken from their secure place when their use is anticipated (e.g. myocardial infarction, severe biliary or kidney colic). Tramadol appears to be a satisfactory alternative.
- Keep a spare kit in your surgery if you or your assistants or locums perform a lot of emergency work.
- Familiarise yourself with the layout of your bag (including ampoule files) so that using it in urgent circumstances is efficient.
- Use a large intravenous cannula wherever possible if rapid infusion is required.

- Be alert for other traffic, hazardous material, HIV contamination, power lines, petrol and other inflammable material, jagged edges.
- Turn ignition off as first measure.
- Ensure proper triage: check airways of all victims first, attend to cervical spine injuries and arrest bleeding.
- Recruit bystanders for simple tasks.
- Control the accident scene: stop traffic with proper lights and signs.

REFERENCES

1. Murtagh J. Drugs for the doctor's bag. Aust Prescriber, 1996; 19(4): 89 - 92.
2. Hogan C. Home visits, emergencies and use of the doctor's bag. Med Observer, 2000: 28 April; 73 - 5.
3. Troller J. The doctor's bag. Essential requirements. Cur Ther, 1990; August: 64 - 5.
4. Johnston CL et al. Medical emergencies in general practice in south-east Queensland: prevalence and practice preparedness. Med J Aust, 2001; 175: 99 - 103.
5. Murtagh J. The doctor's bag. What do you really need? Aust Fam Physician, 2000; 29(1): 25 - 8.
6. Walpole BG. Roadside emergencies. Aust Fam Physician, 1984; 13(4): 249 - 53.

CHAPTER 132. STROKE AND TRANSIENT ISCHAEMIC ATTACKS

A mild attack of apoplexy may be called death's retaining fee.

GILES MENAGE (1613 - 92)

STROKE AND TRANSIENT ISCHAEMIC ATTACKS

Definitions

Stroke A focal neurological deficit lasting longer than 24 hours caused by intracerebral haemorrhage or infarction.

Transient cerebral ischaemic attack (TIA) A focal neurological deficit due to cerebral ischaemia, lasting less than 24 hours.

Stroke in evolution An enlarging neurological deficit, presumably due to infarction, which increases over 24 - 48 hours.

KEY FACTS AND CHECKPOINTS

- A stroke or TIA must be considered a medical emergency.
- One in 10 patients with a TIA are likely to have a stroke shortly afterwards - most within 48 hours. The risk is greatest if older than 60, symptoms last more than 10 minutes and there is weakness or a speech impediment with the TIA.[1]
- Clinical assessment, investigations and treatment should be commenced quickly.
- The best approach to stroke management is aggressive attention to primary and secondary prevention.
- The main risk factors for stroke are atrial fibrillation, hypertension, smoking, age and diabetes.
- Cardiac disease is now a more recognised source of emboli.
- Most patients with a stroke or TIA require urgent imaging to find the cause and guide treatment.
- Ideally patients should be referred to a stroke unit ASAP - within 3 hours.
- Consider the possibility of a cryptogenic stroke, especially from a patent foramen ovale PFO (in 20 - 25% of population and responsible for 50% of these strokes) in relatively young people presenting with a stroke: this leads to paradoxical emboli (from veins to the brain). PFOs may be detected by echocardiography and sealed with a percutaneous closure device.
- Consider the possibility of endocarditis if there is a heart murmur.
- Order a CT or MRI scan on all patients with suspected TIAs and strokes (if not referring to a stroke unit): if normal repeat in 7 days (CT scans unreliable after 7 days).
- Keep in mind atherosclerotic disease of aortic arch as a source of cerebral embolism.
- The place for carotid endarterectomy for asymptomatic carotid stenosis remains controversial. It should be seriously considered if the stenosis is severe, the risk of surgery is low (3% risk of major stroke), the team has proven expertise and the patient is medically fit with a good life expectancy.[2]
- Carotid artery stenting for the treatment and prevention of stroke is an evolving procedure.

MODIFIABLE RISK FACTORS FOR CEREBROVASCULAR DISEASE[2]

Major: hypertension, smoking, cardiovascular disease, atrial fibrillation (especially valvular), diabetes

Others: cardiac failure, dyslipidaemia, obesity, alcohol excess, oral contraception, migraine, stress

Control of risk factors is the key approach to management. Control of hypertension, including systolic hypertension in the elderly, and smoking cessation are vital factors for reduction of the incidence of stroke. A meta-analysis of 14 randomised trials has shown that a reduction of blood pressure of 5 - 6 mmHg is associated with about 40% reduction in stroke incidence.[2] There is a 3 hour time window of opportunity for optimal management.

STROKE

Interesting facts[3]

- Stroke is the third most common cause of death in Australia.
- About one-third of victims will die within 1 month.
- About 50% of ischaemic strokes are preceded by TIAs.
- Thromboembolism from vascular disease outside the brain causes 70% of strokes and 90% of TIAs.
- Such sources are atheromatous plaques within the carotid or vertebral systems or cardiac causes (e.g. postmyocardial infarction).
- Echocardiography is an important investigation with TIAs since LV dysfunction and the size of the left atrium are the strongest independent predictors of thromboembolism.

Pathophysiological groups of cerebral infarction

The three main groups are:

- single penetrator or small vessel disease (lacunar syndrome) - probably due to in situ small vessel disease
- cardioemboli
- large vessel artery-to-artery embolic infarcts (Table 132.1)

Table 132.1 Types and incidence of stroke[3]

Stroke subtype	Frequency (%)
Haemorrhage	
• primary intracerebral	10
• subarachnoid	5
Ischaemic stroke	85
• single penetrator (lacunar)	20
• cardioembolic	25
• large vessel (artery-to-artery)	35
• uncertain/others	20

Differential diagnosis ‘stroke mimics’

- Syncope
- Seizure (and subsequent Todd's paresis)
- Migraine
- Cerebral tumour and other spaceoccupying lesions
- Hypoglycaemia
- Hyponatraemia
- Delirium
- Head injury
- Medically unexplained (e.g. somatisation)

Diagnostic guidelines

- Sudden stroke is typical of embolism.
- The clinical picture depends on the vessel involved.
- In young people < 50 years consider patent foramen ovale.
- With cerebral haemorrhage the stroke evolves steadily, often over hours: the putamen (50%) is the commonest site.
- Lacunar CVAs
    - small deep infarcts
    - pure motor hemiplegia most common effect
    - lack of cortical signs
    - the neurological deficit may progress over 24 - 36 hours
    - outcome usually good
- Investigate all, including SAH, with CT scans or MRI (may need a lumbar puncture for SAH diagnosis).
- Carotid duplex Doppler ultrasound scan can accurately determine atherosclerotic narrowing of the extracranial carotid circulation.

Pitfalls

- Mistaking visual or sensory migraine equivalents in young adults for TIAs
- Mistaking a CVA for labyrinthitis (rare over 50 years)
- Failure to perform carotid duplex Doppler US before starting aspirin for TIA or small stroke (because of missing small haemorrhage, unsuspected tumour or a subdural)
- Diagnosing small stroke as a lacuna (may be a stroke in evolution)

Management

The development of an acute stroke or TIA is a medical emergency and admission to a stroke unit (if available) as soon as possible is advised. The evidence that care in a stroke unit gives the patient the greatest chance of independent survival is proven.[4]

Immediate

- Stabilise ventilation - consider intubation and oxygen.
- Exclude head trauma.
- Obtain urgent CT or MRI scan.
- Treat any seizures.
- Treat any hypoglycaemia.

General

- Carry out investigations, including carotid duplex Doppler US.
- Treat hypertension (systolic > 140) vigorously - it carries a 6 times increased risk of stroke.
- Give IV fluid, electrolyte and nutritional support (nil orally until swallowing has been assessed).
- Good nursing care is the cornerstone of management.
- Physiotherapy and speech therapy.
- Vigorous rehabilitation.
- Intracerebral haemorrhage: considerurgent surgical evacuation for haematomas of the posterior fossa (cerebellum) and cerebral white matter. Shunt insertion may be needed. No medical therapy is of proven value.
- SAH: requires urgent referral (vasospasm and rebleeding are the main causes of morbidity and mortality): nimodipine ± surgery
- Ischaemic stroke: give aspirin 150 - 300 mg (o) daily within 48 hours if CT scan has excluded cerebral haemorrhage or other blood-thinning agents are not used. Early intervention within 3 hours (ideally) of onset with tissue plasminogen activator (rtPA) is currently under investigation, as are cytoprotective agents. Recent trials indicate improved outcomes with rtPA (but not with streptokinase). However, there appears to be a 5 - 7% risk of intracerebral haemorrhage.[2]

Note: Avoid steroids, mannitol, haemodilution and anticoagulation in acute stroke.[5]

TRANSIENT ISCHAEMIC ATTACKS

Features

- Sudden onset
- Complete clinical recovery in less than 24 hours
- Average duration is 5 minutes
- Consciousness usually preserved
- 90% usually in anterior circulation
- Carotid TIAs - unilateral features
- Vertebrobasilar TIAs - often have bilateral or crossed features

A comparison of the main clinical features of carotid (anterior circulation) ischaemia and vertebrobasilar (posterior circulation) ischaemia is presented in Figure 132.1. The carotid circulation accounts for 80% of TIAs.


Carotid artery symptoms
• amaurosis fugax (sudden blindness in one eye)
• dysphasia
• unilateral weakness, paralysis or numbness of face and arm (especially) and leg
internal carotid
external carotid
common carotid
vertebral arteries
right
left
Vertebrobasilar artery symptoms
• homonymous hemianopia
• nausea ± vomiting
• bilateral blurring or blindness
• diplopia
• vertigo
• dysarthria
• hemiplegia
• ataxia ± bilateral weakness or numbness
• drop attacks

Figure 132.1 Cerebral arterial circulation with some important clinical features of carotid and vertebrobasilar ischaemia. Reproduced from C. Kenna and J. Murtagh, *Back Pain and Spinal Manipulation*, Sydney: Butterworths, 1989, with permission

Differential diagnoses of TIAs are presented in Table 132.2.

Table 132.2 Differential diagnosis of TIAs

• Classic migraine (with aura)
• Unusual migraine variants
- hemiplegic
- ophthalmoplegic
- retinal
• Focal epileptic seizures
- complex partial
- simple partial
• Multiple sclerosis
• Transient global amnesia
• Intracranial structural lesions
- arteriovenous malformation
- tumour

- Vestibular disorders
    - acute labyrinthitis
- Benign paroxysmal positional vertigo
- Meniere's syndrome
- Hypoglycaemia
- Adverse drug reactions
- Toxic reactions
- Peripheral nerve lesions
    - carpal tunnel syndrome
    - Bell's palsy
- Psychological conditions
    - hyperventilation
    - panic attacks
    - somatisation

Some ischaemic syndromes

- Transient monocular blindness (amaurosis fugax)
- Transient hemisphere attacks
- The 'locked in' syndrome
- Vertebrobasilar
    - bilateral motor loss
    - crossed sensory and motor loss
    - diplopia
    - bilateral blurring or blindness

Amaurosis fugax

This is the sudden transient loss of vision in one eye (like a 'curtain or shade' coming from above or below) due to the passage of an embolus through the retinal vessels from ipsilateral carotid vessel disease. It is a feature of a TIA in the carotid artery circulation and is often the first clinical evidence of carotid stenosis.[6] About 20% of all TIAs present as amaurosis fugax.[7] Amaurosis fugax may forewarn of the development of hemiparesis or blindness and should be considered a matter for urgent attention and rectification. Carotid endarterectomy may be required for high grade stenosis.

Transient hemisphere attack (usually middle cerebral artery)

- Affects motor or sensory or both
- Usually face and arm (more than leg)
- Dysphasia common

The 'locked in' syndrome

In this syndrome, which may be transient or persistent, patients remain conscious and aware of their dilemma but are unable to speak or move the limbs, particularly the arms. It may be possible to communicate with patients using eye responses to commands. The cause is invariably a lesion in the brain stem.

Significance of TIAs

- Five years after a TIA, 22 - 51% (average 1 in 3) patients (without treatment) will have a stroke.[6, 8] This figure may be higher for those with ipsilateral high-grade (70% or more) carotid stenosis.

- The highest risk is in the first 6 months.
- A carotid artery TIA has more serious prognostic significance. Such patients are at high risk of developing a stroke that is potentially preventable.
- Referral for investigation is appropriate.
- All patients should have a carotid duplex Doppler US and CT scan at presentation.
- Cardiac status should be addressed because of an association with myocardial infarction.

Investigations

- Full blood count
- Blood glucose, creatinine and cholesterol levels
- Thyroid function tests
- Carotid duplex Doppler US (the investigation of choice)
- ECG
- transoesophageal echocardiography
- CT scan (with contrast)

Treatment

- Aim to minimise the risk of a major stroke.
- Determine cause and correct (if possible).
- Early neurorehabilitation.
- Advise cessation of smoking and treat hypertension (if applicable).
- Anti-platelet therapy (especially for carotid ischaemia): aspirin 50 - 300 mg (o) daily (gives 30% protection from stroke or death after TIA)[9] or clopidogrel 75 mg (o) daily or dipyridamole + aspirin 200 mg/25 mg (o) bd (proven better outcomes than aspirin alone)[10] or (especially if aspirin contraindicated) ticlopidine 250 mg (o) bd after food (concern about neutropenia so monitor blood).
- Anticoagulation therapy: warfarin
  - for vertebrobasilar ischaemia (with increasing frequency of TIAs)
  - for failed anti-platelet therapy
  - atrial fibrillation (selected cases) > 65 years of age.
- Carotid endarterectomy has been proven to have a place in the management of carotid artery stenosis and the decision depends on the expertise of the unit.[11] There is no evidence that surgery is appropriate for the asymptomatic patient with a stenosis < 60% (antiplatelet therapy proved to be as effective) or the symptomatic patient with a stenosis less than 30%, but there is a significant benefit for a stenosis greater than 70% (and possibly > 60% in asymptomatic patients). If the stenosis is > 90% refer immediately. Greater than 75% stenosis is associated with a 2% per annum rate of ipsilateral ischaemic stroke.[2]
- Percutaneous transluminal angioplasty (stenting). Carotid artery stenting for the treatment and prevention of stroke is now employed in stroke units as an alternative to surgery.

ATRIAL FIBRILLATION[2, 3]

- The main source of cardioembolic infarction
- Increased with risk factors - hypertension, previous embolism and recent CHF(previous 3 months)
- With non-valvular AF, annual risk of CVA is 2.5% (no risk factors) to 17.6% (2+ risk factors)
- Intermittent AF can also be a risk

Management

- Valvular disease: warfarin - target INR 2 - 3

- Non-valvular AF
  - no risk factors: aspirin 100 - 300 mg/day
  - risk factors: warfarin: INR 2 - 3
  - if warfarin contraindicated: aspirin

CEREBRAL VENOUS THROMBOSIS[2]

This rare cause of stroke may present as an acute or chronic cerebrovascular disorder. It should be suspected particularly in women in the post-partum period preventing with severe headache or focal neurological defects. Diagnosis is by MRI. In the acute phase treatment is by anti-coagulation with heparin followed by warfarin for approximately 6 months.

INDICATIONS FOR CAROTID DUPLEX DOPPLER ULTRASOUND

- Bruit in neck
- TIAs
- Crescendo TIAs (more frequent and longer lasting)
- Vertebrobasilar insufficiency symptoms
- Hemispheric stroke
- Prior to major vascular surgery (e.g. CABG)

WHEN TO REFER

- Refer immediately to a stroke unit
- Suspicion of SAH
- Carotid artery stenosis on carotid duplex Doppler ultrasound scan
- Cerebellar haemorrhage on CT scan
- Stroke in a young patient < 50 years (consider patent foramen ovale and other less common causes)

KEY CHECKPOINTS

Proven strategies

3 proven strategies to improve outcome of acute stroke (level 1 evidence):

- management in a stroke unit
- giving IV tPA within 3 hours of ischaemic stroke
- giving antiplatelet agents within 48 hours of ischaemic stroke

Predicators (ABCD) of a high risk of stroke within the first 7 days of TIA12

- Age > 60
- Blood pressure > 140 systolic and/or 90 diastolic
- Clinical features: unilateral limb weakness or speech disturbance
- Duration of symptoms > 10 minutes

First aid for stroke

- Is it a stroke?
- Think FAST

- ↓
- Face (ask person to smile)
- Arms (raise both arms)
- Speech (speak a simple sentence)
- Time (within 3 hours) then if yes
- Refer to stroke unit ASAP
- Don't give aspirin

REFERENCES

1. Claiborne Johnston S et al. Short-term prognosis after emergency department diagnosis of TIA. JAMA, 2000; 284: 2901 - 6.
2. Tiller J (Chair). *Therapeutic Guidelines*: Neurology (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2002: 79 - 89.
3. Donnan G. How to treat: TIA and stroke. Aust Doctor, 1994; 16 September: 1 - 4.
4. Stroke Unit Trialists' Collaboration. Organised inpatient (stroke unit) care for stroke (Cochrane Review). In: The Cochrane Library, Issue 3, 2004. Chichester, UK: John Wiley & Sons Ltd.
5. Lindley RI, Landau P. Early management of acute stroke Aust Prescr, 2004; 27: 120 - 3.
6. Kumar PJ, Clark ML. *Clinical Medicine* (2nd edn). London: Bailliere Tindall, 1990: 907 - 13.
7. Crimmins DS. How to investigate the patient with TIA. Mod Med Aust, 1995; 38(12): 71 - 4.
8. JoffR. Diagnosis and treatment of transient ischaemic attacks. Mod Med Aust, 1994; 37(8): 18 - 23.
9. Leicester J. Stroke and transient cerebral ischaemic attacks. In: *MIMS Disease Index* (2nd edn). Sydney: IMS Publishing, 1996: 487 - 9.
10. Halkes PH et al. ESPRIT study group. Aspirin + dipyrid amole versus aspirin alone after cerebral ischaemia of arterial origin: a RCT, Lancet, 2006; 367: 1665 - 73.
11. Donnan G (Chair). NH&MRC Working Party. Clinical practice guidelines: Prevention of stroke - the role of anticoagulants, antiplatelets and carotid endarterectomy. Consultation document draft. Canberra, NH&MRC, 1996.
12. Rothwell PM et al. A simple score (ABCD) to identify individuals at high early risk of stroke after transient ischaemic attack. Lancet, 2005; 366: 29 - 36.

CHAPTER 133. THROMBOSIS AND THROMBOEMBOLISM

Thrombosis is an ugly thought; Dicumarol and heparin, Statistics show, prevent most ills Like thrombus of the coronary, infarction of the heart And tests are made so many times That arms turn blue and blood runs thin

DAVID LITTMAN NEJM, 1965, 'THE GOOD OLD DAYS'

THROMBOSIS AND THROMBOEMBOLISM

A thrombus is defined as a clot formed in the circulation from the constituents of the blood, whereas emboli are fragments of thrombus that break off and block vessels downstream. Almost half of adult deaths in Western countries are due to thrombosis of coronary arteries or cerebral arteries and to pulmonary embolism.[1] A thrombus is the result of a cascade of events involving platelets, RBCs, coagulation factors and the vessel wall.

Conditions predisposing to thrombosis:

- thrombophilia
- thrombocytosis (↑ platelets)
- polycythemia rubra vera

THROMBOPHILIA[2]

Thrombophilia refers to a disorder of haemostasis in the form of a primary coagulopathy leading to a tendency to thrombosis. This should be considered in patients with major unprecipitated venous thromboembolism with or without a strong family history of venous thrombosis. There are several causes that can be tested. The factors are both inherited and acquired:

- inherited:
  - factor V Leiden gene mutation (activated protein C resistance)
  - prothrombin gene mutation
  - protein C deficiency
  - protein S deficiency
  - antithrombin deficiency
- acquired:
  - antiphospholipid antibodies (anticardiolipin or anti-$ß_2$ GPI)
  - elevated homocysteine level
  - lupus anticoagulant

The above factors can all be measured in the laboratory with specific genetic tests, coagulation or antibody-based tests. Other acquired causes (regarded as risk factors) include malignancy, OCP, HRT, immobilisation, pregnancy and major surgery. Consideration should be given to screening patients with a DVT, especially in travellers at possible risk. Referral to a haematologist if thrombophilia is proved or suspected is advisable. Of particular interest is factor V Leiden, which occurs in about 5% of Caucasians, and the risk of venous thrombosis is increased three to seven times in heterozygotes and 80 times in homozygotes.[2]

Indications for investigation

- Recurrent or unusual thrombosis
- Venous thromboembolism < 40 years
- Arterial thrombosis < 30 years
- Skin necrosis, especially on warfarin
- Recurrent foetal loss
- Familial thromboembolism

VENOUS THROMBOEMBOLISM

A feature of venous thrombosis is that it develops in normal vessels, with key factors being stasis and increased coagulability including thrombophilia. The classic example is deep venous thrombosis of the leg veins. Another poorly recognised thrombosis is axillosubclavian venous thrombosis, which is associated with pulmonary embolism in 30% of cases.[3]

DEEP VENOUS THROMBOSIS

Past history is very important.

Risk factors

- Family history
- Thrombophilia
- History of previous thromboembolism
- Drugs (e.g. OCP, tamoxifen, HRT)
- Malignancy (watch idiopathic DVT)
- Increasing age; age > 40
- Varicose veins
- Significant illness, esp. heart failure and cancer
- Other chronic illness
- Recent surgery
- Major/orthopaedic surgery
- Immobility
- Long flights
- Pregnancy/puerperium
- Obesity
- Dehydration

There is an up to 20% association with pulmonary emboli, of which 30% may be fatal. DVT may be asymptomatic, but usually causes tenderness in the calf. DVT may present with painless unilateral leg swelling. Because of the potentially serious consequences from untreated thromboembolism, it is essential to objectively confirm or exclude clinically suspected disease.

Investigations

- Duplex ultrasound: accurate for above-knee thrombosis; improving for distal calf
- Should be repeated in 1 week if initial test normal
- Contrast venography: reserved if ultrasound doubtful

Note:

- MRI appears to be very accurate for DVT, but is not yet available generally.
- The plasma D-dimer can be helpful. Where the clinical probability of venous thrombosis is low, a normal D-dimer effectively excludes the diagnosis. Where clinical probability is high, appropriate imaging with Doppler ultrasound or lung scan should be performed. A raised D-dimer is non-specific and does not help with diagnosis.

Treatment[4]

Drugs used in the treatment of thromboembolism are summarised in general format in Table 133.1.

Table 133.1 Drugs employed in thrombotic disorders

Antiplatelet
Aspirin
Clopidogrel
Dipyridamine
Ticlopidine
Glycoprotein IIb/IIIa inhibitors, e.g. abciximab
Anticoagulants
Heparin
• unfractionated/standard
• low molecular weight
- dalteparin
- enoxaparin
Vitamin K antagonists
• phenindione
• warfarin
Hirudins
• bivalirudin
• fondaparinux
• lepirudin
Thrombolytics
Alteplase
Reteplase
Streptokinase
Tenecteplase
Urokinase

- Collect blood for APTT, INR, and platelet count.
- LMW heparin (e.g. enoxaparin 1.5 mg/kg subcutaneously daily or enoxaparin 1 mg/kg sc bd).
- Oral anticoagulant (warfarin) for 3 to 6 months
- Do not give aspirin
- Mobilisation within the limits of pain, tenderness and swelling
- Class II graded compression stocking to affected leg in proximal DVT associated with significant swelling. The stocking may be above or below the knee depending on the extent of the swelling.

Prevention of deep venous thrombosis

Surgery

- Early ambulation
- Unfractionated heparin 5000 U (sc) bd (intermediate risk), LMW heparin for orthopaedic or other high-risk surgery
- Graded pressure elastic stockings
- Physiotherapy
- Pneumatic compression (especially in high-risk patients where heparin is contraindicated)

- Electrical calf muscle stimulation during surgery

Prolonged travel/immobilisation

- Keep hydrated - ample water.
- Avoid or restrict alcohol and coffee.
- Exercises - 3 - 4 minutes per hour, e.g. walking, calf contraction (e.g. foot pumps, Figure 133.1), ankle circles, knee lifts



Figure 133.1 Foot pump exercises to prevent DVT during prolonged air travel.

Figure 133.1 1 Start with the feet flat and both heels on the floor, lift the feet (toes up) as high as you can. 2 Return the feet to the flat position, pressing them firmly on the floor. 3 Lift the heels high, keeping the balls of the feet on the floor. Continue this up-and-down movement for at least 30 seconds.

- Injections LMWH just prior to flying and on arrival for those at high risk. Use a prophylactic dose, e.g. enoxaparin 40 mg or dalteparin 5000 U

Pulmonary embolism[5]

Refer to pages XXX - XXX for more detail on the clinical features and management of pulmonary embolism. CT pulmonary angiography is very specific and appears to be as sensitive as Y/Q scanning for embolism, and is currently the preferred first-line investigation. The basis of treatment is LMW heparin and warfarin, as for DVT.

ARTERIAL THROMBOEMBOLISM

The common serious manifestations of this are myocardial infarction, stroke and occlusion of the arterial system of the lower limbs. Emboli originate from the left side of the heart, the carotid arteries or the iliac arteries. Atrial fibrillation (AF) is an important predisposing factor.

Preventing systemic embolism in atrial fibrillation[5]

AF accounts for about 15% of ischaemic strokes caused by systemic embolism of cardiac thrombi. This risk gradually increases with age. Use of warfarin reduces the annual incidence of stroke during AF from 4.5% to 1.4% - a risk reduction of almost 70%. The decision to use it or an anti-platelet agent is a difficult one and should be made in consultation with a cardiologist. Although aspirin reduces stroke rate by about 20% in patients with AF when compared with no treatment, it has half the effectiveness of warfarin and is less able to prevent severe strokes. As a general rule all patients with AF should start on warfarin unless < 65 years old or have a major contraindication to its use. It is not indicated in patients with lone AF who are less than 60 years of age with no risk factors. If using warfarin, start with a low dose (e.g. 2 - 4 mg) and maintain an INR of 2 - 3 with regular checks. Anticoagulant therapy is also required to prevent embolism after cardioversion.

DRUG TREATMENT OF THROMBOTIC DISORDERS

Warfarin

Warfarin is the key oral agent for the treatment and prevention of venous thromboembolism, unlike the antiplatelet drugs which have little or no benefit and are recommended in arterial disease. Before prescribing warfarin, the risk of bleeding should be evaluated and discussed with each patient. Indications for warfarin treatment are outlined in Table 133.2.

Table 133.2 Warfarin anticoagulation

Indications
• Prosthetic cardiac valves
• Deep venous thrombosis, pulmonary thromboembolism
• Atrial fibrillation (selected cases)
• Postoperatively in lower limb orthopaedic surgery (low dose)
• Postcoronary bypass surgery (selected cases)
• Thrombosis in antiphospholipid antibody syndrome
Contraindications
• Active bleeding
• History of intracranial haemorrhage
• Uncontrolled hypertension
• Liver disease with impaired synthetic function - based on INR
• Pregnancy

Actions[6]

- Antagonises vitamin K.
- Depresses factors VII, IX and X (half-life of 30 - 40 hours) and prothrombin.
- Achieves full anticoagulation effect after 5 - 7 days.
- Prothrombin time (INR ratio) of 2 - 3 times normal control indicates therapeutic effect.
- The INR (international normalized ratio) is a good indicator of effectiveness and risk of bleeding.
- Duration of effect is 4 - 5 days after cessation.
- Antidote is vitamin K ± plasma or prothrombin complex concentrate.

Initiation of warfarin treatment[7, 8]

An estimate of the patient's final steady dose is made. The patient is commenced on this dose and the INR monitored daily and the dose altered accordingly.

- Measure INR first to establish baseline.
- Generally warfarin is commenced on same day or day after heparin is commenced.
- Heparin can be ceased when INR > 2 for 2 consecutive days.
- Typical loading dose is 5 - 10 mg (o) daily for 2 days (avoid dose > 30 mg over 3 days without INR).
- Adjust the dosage according to the INR table (Table 133.3) from the third day.
- Establish the INR in the therapeutic range, usually 2 - 3 (average 2.5).

Table 133.3 Warfarin dosages adjustment[*10]

Day	INR	Dose
1	-	5 - 10 mg[**]
2	< 1.8	5 mg[**]
	1.8 to 2.0	1 mg

	> 2.0	Hold
3	< 2	5 mg
	2.0 to 2.5	4 mg
	2.6 to 2.9	3 mg
	3.0 to 3.2	2 mg
	3.3 to 3.5	1 mg
	> 3.5	Hold
4, and until stabilised	> 1.4	10 mg
	1.4 to 1.5	7 mg
	1.6 to 1.7	6 mg
	1.8 to 1.9	5 mg
	2.0 to 2.3	4 mg
	2.4 to 3.0	3 mg
	3.1 to 3.2	2 mg
	3.3 to 3.5	1 mg
	> 3.5	Hold

[*] This table should be used only if the pretreatment INR is normal.

[**] 5 mg of warfarin should be given to patients who are more likely to be sensitive to warfarin. This includes the elderly, the very ill, the malnourished and patients with abnormal liver function or significant chronic renal failure.

- Maintenance dose is usually reached by day 5.
- The INR reflects the warfarin dose given 48 hours earlier.
- It is best taken in the evening and the INR measured in the morning.

INR measurement schedule

- before treatment on third day
- ↓
- daily for 1 week
- ↓
- 2 times daily for 2 weeks
- ↓
- weekly for 4 weeks
- ↓
- monthly

Note:

- Warfarin should be continued for 3 to 6 months and longer if major risk factors are present.
- Watch for potential drug interactions.

Recommended target INR values are given in the box.

Recommended INR target values

- Prevention of DVT 2.0 - 3.0
- Treatment of DVT or PE 2.0 - 3.0
- Preventing systemic embolism 2.0 - 3.0

  - atrial fibrillation
  - post myocardial infarction
  - tissue heart valve
  - valvular heart disease
- Mechanical prosthetic heart valve 2.5 - 3.5
- Prevent recurrence of MI 2.0 - 3.0
- Antiphospholipid antibody syndrome thrombosis 2.0 - 3.0

Overdosage of warfarin

Signs of warfarin overdosage include:

- unexpected bleeding after minor trauma
- epistaxis
- spontaneous bruising
- unusually heavy menstrual bleeding
- gastrointestinal bleeding

Management of overdosage[9]

1. Urgent measurement of INR is required.
2. If the only evidence of overdosage is a small increase of the INR above the therapeutic range, cessation of warfarin for 1 to 2 days followed by a continuation at a lower dose is appropriate.
3. If the INR is markedly elevated (> 5.0) consider giving a small dose of oral vitamin K (1 - 2 mg).
4. If bleeding is minor, transient action as in point 2 is still appropriate.
5. If bleeding is persistent, or severe, or involves closed body cavities (such as pericardium, intracranial, fascial compartment), urgent admission to hospital is essential. The anticoagulation may need to be reversed by administering oral or parenteral vitamin K. Infusion of fresh frozen plasma and/or prothrombin complex concentrate may also be necessary.

Drug interactions

There are so many potential interactions between warfarin and other drugs that the following general principles should be applied:

1. Maintain the simplest possible drug regimens. Avoid polypharmacy.
2. Aspirin is contraindicated while the patient is on warfarin because of the combined antiplatelet and anticoagulation effects. The risk of gastrointestinal bleeding is also increased. Other NSAIDs should also be avoided (Table 133.4).
3. If the patient's drug regimen must be altered during warfarin therapy, then the INR should be followed closely until stable.

Table 133.4 Some important drug interactions with warfarin

Effects on warfarin activity	Drug
↑ Increased	Allopurinol
	Amiodarone
	Anabolic steroids
	Antibiotics (broad spectrum)
	Antifungals
	Aspirin - salicylates (high doses)

	Chloral hydrate
	Cimetidine
	Clofibrate
	Gemfibrozil
	Metronidazole
	Miconazole
	NSAIDs, including COX-2 inhibitors
	Paracetamol (large doses)
	Phenytoin
	Proton-pump blockers
	Quinidine/quinine
	Ranitidine
	SSRIs
	Sulphonamides
	Tamoxifen
	Thyroxine
	Herbal medicines:
	• dong quai
	• papaya
	•St John's wort
↑ Decreased	Antacids
	Antihistamines
	Barbiturates
	Antiepileptics (e.g. carbamazepine)
	Cholestyramine (reduced absorption)
	Griseofulvin
	Haloperidol
	Oestrogen/oral contraceptives
	Rifampicin
	Vitamin C
Increased or decreased	Alcohol
	Chloral hydrate
	Diuretics
	Ranitidine

Advice to the patient

- Keep to a consistent diet.
- Do not take aspirin or liquid paraffin.

- Always mention that you take warfarin to any doctor, dentist or chemist you are consulting.
- Remember to take tablets strictly as directed and have your blood tests.
- Report signs of bleeding, such as black motions, blood in urine, easy bruising, unusual nose bleeds, heavy periods, ‘purple toes’.

Note: Give the patient the information sheet about risks, especially if ceasing warfarin.

Practice tips[10]

- Consider avoiding use if patient compliance is likely to be poor.
- The INR result reflects the warfarin dose administered 48 - 72 hours earlier.
- Advise and encourage patients to keep a record in an ‘anticoagulant diary’ of drug dosage and INR results.
- An unacceptable INR is > 5.0.
- Discontinue therapy if skin necrosis or ‘purple toes syndrome’ occurs.

REFERENCES

1. Kumar P, Clark M. *Clinical Medicine*. London: Saunders, 2002: 465.
2. Joseph J. *Thrombophilia. Common sense pathology*. RCPA, 2004.
3. Smith A (Chair). *Therapeutic Guidelines: Cardiovascular* (Version 4). Melbourne. Therapeutic Guidelines Ltd, 2003: 179.
4. ibid.: 177 - 9.
5. Gallus AS. Anticoagulation: how to treat. Australian Doctor, 4 March 2005: 29 - 36.
6. Bochner F. *Australian Medicines Handbook*. Adelaide: Australian Medicines Handbook Pty Ltd, 2006: 291 - 2.
7. Walker ID et al. Guidelines on oral anticoagulation (3rd edn). Br J Haematol, 1998; 101: 374 - 87.
8. Gallus AS. Consensus guidelines for warfarin therapy - recommendations from the Australasian Society of Thrombosis and Haemostasis. Med J Aust, 2000; 172: 600 - 5.
9. Baker R et al. Australian and NZ Consensus Guidelines for Warfarin Reversal. MJA, 2004; 181(9): 492 - 7.
10. Campbell P, Roberts R, Eaton V et al. Managing warfarin therapy in the community. Aust Prescr, 2001; 24: 86 - 9.

CHAPTER 134. COMMON SKIN WOUNDS AND FOREIGN BODIES

The variety of foreign bodies which have found their way into the rectum is hardly less remarkable than the ingenuity displayed in their removal.

A turnip has been delivered PR by the use of obstetric forceps.

A stick firmly impacted has been withdrawn by inserting a gimlet into its lower end.

A tumbler, mouth downwards, has several times been extracted by filling the interior with a wet plaster of Paris bandage, leaving the end of the bandage extruding, and allowing the plaster to set.

BAILEY AND LOVE 1943, A SHORT PRACTICE OF SURGERY

COMMON SKIN WOUNDS AND FOREIGN BODIES

Injuries to the skin, including simple lacer-ations, abrasions, contusions and foreign bodies, are among the commonest problems encountered in general practice. To manage these cosmetically important injuries well is one of the really basic and enjoyable skills of our profession.

KEY FACTS AND CHECKPOINTS

- A well-prepared treatment room with good sterilisation facilities, instruments, sterile dressings and an assistant facilitates management.
- With lacerations check carefully for nerve damage, tendon damage and arterial damage.
- Beware of slivers of glass in wounds caused by glass - explore carefully and X-ray (especially with high-resolution ultrasound) if in doubt.
- Beware of electrical or thermal wounds because marked tissue necrosis can be hidden by slightly injured skin.
- Beware also of roller injuries such as car wheels.
- Beware of pressure gun injuries such as oil and paint. The consequences can be disastrous.
- Avoid suturing the tongue, and animal and human bites, unless absolutely necessary.
- Keep drawings or photographs of wounds in your medical records.
- Have a management plan for puncture wounds, including medical needlestick injuries.
- Gravel rash wounds are a special problem because retained fragments of dirt and metal can leave a ‘dirty’ tattoo-like effect in the healed wound.

CONTUSIONS AND HAEMATOMAS

A contusion (bruise or ecchymosis) is the consequence of injury causing bleeding in subcutaneous or deeper tissue while leaving the skin basically intact. It might take weeks to resolve, especially if extensive.

A haematoma is a large collection of extravasated blood that produces an obvious and tender swelling or deformity. The blood usually clots and becomes firm, warm and red; later (about 10 days) it begins to liquify and becomes fluctuant.

Principles of management

- Explanation and reassurance
- RICE (for larger bruises/haematomas) for 48 hours
  - R: rest
  - I: ice (for 20 minutes every 2 waking hours)
  - C: compression (firm elastic bandage)
  - E: elevation (if a limb)
- Analgesics: paracetamol/acetaminophen

- Avoid aspiration (some exceptions)
- Avoid massage
- Heat may be applied after 72 hours as local heat or whirlpool baths
- Consider possibility of bleeding disorder if bleeding out of proportion to injury

PROBLEMATIC HAEMATOMAS

Some haematomas in certain locations can cause deformity and other problems.

Haematoma of nasal septum[1]

Refer p. 638.

Haematoma of the pinna[1]

When trauma to the pinna causes a haematoma between the epidermis and the cartilage, a permanent deformity known as 'cauliflower ear' may result. The haematoma, if left, becomes organised and the normal contour of the ear is lost.

The aim is to evacuate the haematoma as soon as practical and then to prevent it reforming. One can achieve a fair degree of success even on haematomas that have been present for several days.

Method

Under aseptic conditions insert a 25 gauge needle into the haematoma at its lowest point and aspirate the extravasated blood (Figure 134.1a). Apply a padded test tube clamp to the haematoma site and leave on for 30 - 40 minutes (Figure 134.1b). Generally, daily aspiration and clamping are sufficient to eradicate the haematoma completely.



Figure 134.1 Treatment of haematoma of the pinna

Pretibial haematoma

A haematoma over the tibia (shin bone) can be persistently painful and slow to resolve. An efficient method is, under very strict asepsis, to inject 1 mL lignocaine 1% and 1 mL hyalur-onidase and follow with immediate ultrasound. This may disperse or require drainage.

Subungual haematomas

These important haematomas are discussed in Chapter 119.

ABRASIONS

Abrasions vary considerably in degree and potential contamination. They are common with bicycle or motorcycle accidents and skateboard accidents. Special care is needed over joints such as the knee or elbow.

Rules of management

- Clean meticulously, remove all ground-in dirt, metal, clothing and other material.
- Scrub out dirt with sterile normal saline under anaesthesia (local infiltration or general anaesthesia for deep wounds).
- Treat the injury as a burn.
- When clean, apply a protective dressing (some wounds may be left open).
- Use paraffin gauze and non-adhesive absorbent pads such as Melolin.
- Ensure adequate follow-up.
- Immobilise a joint that may be affected by a deep wound.

LACERATIONS

Lacerations vary enormously in complexity and repairability. Very complex lacerations and those involving nerves or other structures should be referred to an expert.

Principles of repair

- Good approximation of wound edges minimises scar formation and healing time.
- Pay special attention to debridement.
- Avoid deep layers of suture material in a contaminated wound - consider drainage.
- Inspect all wounds carefully for damage to major structures such as nerves and tendons and for foreign material:
    - shattered glass wounds require careful inspection and perhaps X-ray.
    - high-energy wounds (e.g. motor mowers) are prone to have metallic foreign bodies and associated fractures.
- Be ready to take X-rays of wounds to look for foreign objects or fractures (compound fractures).
- Trim jagged or crushed wound edges, especially on the face.
- All wounds should be closed in layers.
- Avoid leaving dead space.
- Do not suture an ‘old’ wound (greater than 8 hours) if it is contaminated with primary closure: leave 4 days before suturing if not infected.
- Take care in poor healing areas, such as backs, necks, calves and knees; and in areas prone to hypertrophic scarring, such as over the sternum of the chest and the shoulder.
- Use atraumatic tissue-handling techniques.
- Everted edges heal better than inverted edges.
- Practise minimal handling of wound edges.
- A suture is too tight when it blanches the skin between the thread - it should be loosened.
- Avoid tension on the wound, especially in fingers, lower leg, foot or palm.
- The finest scar and best result is obtained by using a large number of fine sutures rather than a fewer thicker sutures more widely spread.
- Avoid haematoma.

- Apply a firm pressure dressing when appropriate, especially with swollen skin flaps.
- Consider appropriate immobilisation for wounds. Many wound failures are due to lack of immobilisation from a volar slab on the hand or a back slab on the leg.

Practical aspects

Suture material (Table 134.1)

Table 134.1 Selection of suture material (guidelines)

Skin	Nylon 6/0	Face
	Nylon 3/0	Back, scalp
	Nylon 5/0	Elsewhere
Deeper tissue (dead space)	Catgut 4/0	Face
	Dexon/Vicryl 3/0 or 4/0	Elsewhere
Subcuticular	Catgut 4/0	
Small vessel ties	Plain catgut 4/0	
Large vessel ties	Chromic catgut (CCG) 4/0	

- Monofilament nylon sutures are generally preferred for skin repair.
- Use the smallest calibre compatible with required strains.
- The synthetic, absorbable polyglycolic acid or polyglactin sutures (Dexon, Vicryl) are stronger than catgut of the same gauge, but do not use these (use catgut instead) on the face or subcuticularly.

Instruments

Examples of good quality instruments:

- locking needle holder (e.g. Crile-Wood 12 cm)
- skin hooks
- iris scissors
- toothed forceps

Holding the needle

The needle should be held in its middle; this will help to avoid breakage and distortion, which tend to occur if the needle is held near its end (Figure 134.2).



Figure 134.2 Correct and incorrect methods of holding the needle

Dead space

Dead space should be eliminated to reduce tension on skin sutures. Use buried absorbable sutures to approximate underlying tissue. This is done by starting suture insertion from the fat to pick up the fat/dermis interface so as to bury the knot (Figure 134.3).



Figure 134.3 Eliminating dead space

Everted wounds

Eversion is achieved by making the 'bite' in the dermis wider than the bite in the epidermis (skin surface) and making the suture deeper than it is wide. Shown are:

- simple suture (Figure 134.4a)
- vertical mattress suture (Figure 134.4b)



Figure 134.4 Everted wounds: (a) correct and incorrect methods of making a simple suture; (b) making a vertical mattress suture

The mattress suture is the ideal way to evert a wound.

Number of sutures

One should aim to use a minimum number of sutures to achieve closure without gaps but sufficient sutures to avoid tension. Place the sutures as close to the wound edge as possible.

SPECIAL TECHNIQUES FOR VARIOUS WOUNDS

The three-point suture

In wounds with a triangular flap component, it is often difficult to place the apex of the flap accurately. The three-point suture is the best way to achieve this while minimising the chance of strangulation necrosis at the tip of the flap.

Method

1. Pass the needle through the skin of the non-flap side of the wound.
2. Pass it then through the subcuticular layer of the flap tip at exactly the same level as the reception side.
3. Finally, pass the needle back through the reception side so that it emerges well back from the V flap (Figure 134.5).



Figure 134.5 The three-point suture

Triangular flap wounds on the lower leg

Triangular flap wounds below the knee are a common injury and are often treated incorrectly. Similar wounds in the upper limb heal rapidly when sutured properly, but lower limb injury will not usually heal at first intention unless the apex of the flap is given special attention.

Proximally based flap

A fall through a gap in flooring boards will produce a proximally based flap; a heavy object (such as the tailboard of a trailer) striking the shin will result in a distally based flap.

Often the apex of the flap is crushed and poorly vascularised; it will not survive to heal after suture.

Treatment methods (under infiltration with LA)

1. Preferred method: To attempt to salvage the distal flap, scrape away the subcutaneous tissue on the flap and use it as a full-thickness graft.
2. An alternative is to excise the apex of the flap, loosely suture the remaining flap and place a small split-thickness graft on the raw area (Figure 134.6).


donor site (within anaesthetised area)
excised apex of skin flap
and graft site

Figure 134.6 Triangular flap wound repair: proximally based flap

For both methods apply a suitable dressing and strap firmly with a crepe bandage. The patient should rest with the leg elevated for 3 days.

Distally based flap (Figure 134.7)



Figure 134.7 Triangular flap wound repair: distally based flap

This flap, which is quite avascular, has a poorer prognosis. The same methods as for the proximally based flap can be used. Trimming the flap and using it as a full thickness graft has a good chance of repair in a younger person but a poor chance in the elderly.

Repair of cut lip

While small lacerations of the buccal mucosa of the lip can be left safely, more extensive cuts require careful repair. Local anaesthetic infiltration may be adequate, although a mental nerve block is ideal for larger lacerations of the lower lip.

For wounds that cross the vermilion border, meticulous alignment is essential. It may be advisable to premark the vermilion border with gentian violet or a marker pen. It is desirable to have an assistant.

Method

1. Close the deeper muscular layer of the wound using 4/0 CCG. The first suture should carefully appose the mucosal area of the lip, followed by one or two sutures in the remaining layer.
2. Next, insert a 6/0 monofilament nylon suture to bring both ends of the vermilion border together. The slightest step is unacceptable (Figure 134.8). This is the key to the procedure.



Figure 134.8 The lacerated lip: ensuring meticulous suture of the vermilion border

3. Close the inner buccal mucosa with interrupted 4/0 plain catgut sutures.
4. The outer skin of the lip (above and below the vermilion border) is closed with interrupted nylon sutures.

Post repair

1. Apply a moisturising lotion along the lines of the wound.
2. Remove nylon sutures in 3 - 4 days (in a young person) and 5 - 6 days (in an older person).

Repair of lacerated eyelid

General points

- Preserve as much tissue as possible.
- Do not shave the eyebrow.
- Do not invert hair-bearing skin into the wound.
- Ensure precise alignment of wound margins.
- Tie suture knots away from the eyeball.

Method

1. Place an intermarginal suture behind the eyelashes if the margin is involved.
2. Repair conjunctiva and tarsus with 6/0 catgut.
3. Then repair skin and muscle (orbicularis oculi) with 6/0 nylon (Figure 134.9).



Figure 134.9 The lacerated eyelid

Repair of tongue wound

Wherever possible, it is best to avoid repair to wounds of the tongue because these heal rapidly. However, large flap wounds to the tongue on the dorsum or the lateral border may require suturing. The best method is to use buried catgut sutures.

Method

1. Get patient to suck ice for a few minutes, then infiltrate with 1% lignocaine and leave 5 - 10 minutes.
2. Use 4/0 or 3/0 catgut sutures to suture the flap to its bed, and bury the sutures (Figure 134.10).



Figure 134.10 Repair of tongue wound

It should not be necessary to use surface sutures. If it is, 4/0 silk sutures will suffice.

The patient should be instructed to rinse the mouth regularly with salt water until healing is satisfactory.

The amputated finger

In this emergency situation, instruct the patient to place the severed finger directly into a fluid-tight sterile container, such as a plastic bag or sterile specimen jar. Then place this ‘unit’ in a bag containing iced water with crushed ice.

Note: Never place the amputated finger directly in ice or in fluid such as saline. Fluid makes the tissue soggy, rendering microsurgical repair difficult.

Care of the finger stump

Apply a simple, sterile, loose, non-sticky dressing and keep the hand elevated.

Bite wounds

Human bites and clenched fist injuries

Human bites and clenched fist injuries can present a serious problem of infection. β-lactamase producing anaerobic organisms in the oral cavity (e.g. Vincent's) can penetrate the damaged tissue and form a deep-seated infection. Streptococcus species, staphylococcal organisms and Eikenella corrodens are common pathogens. Complications of the infected wounds include cellulitis, wound abscess and lymphangitis. A coctirane review of antibiotic prophylaxis concluded that it significantly reduces the risk of infection.[2]

Principles of treatment

- Clean and debride the wound carefully (e.g. aqueous antiseptic solution or hydrogen peroxide).
- Give prophylactic penicillin if a severe or deep bite.
- Avoid suturing if possible.
- Tetanus toxoid (although minimum risk).
- Consider rare possibility of HIV and hepatitis B or C infections.
- For high-risk wounds, give procaine penicillin 1.5 g IM statim and/or and amoxycillin/clavulanate 875/125 mg bd for 5 days.[3]
- If established infection in a deep wound, take a swab and give metronidazole 400 mg (o) bd for 5 - 10 days plus either cefotaxime 1 g IV 8 hourly or ceftriaxone 1 g IV daily for 5 - 10 days.

Dog bites

Non-rabid

Dog bites typically have poor healing and carry a risk of infection with anaerobic organisms, including tetanus, staphylococci and strep-tococci. Puncture and crush wounds are more prone to infection than laceration. Upto 25% of dog bited wounds become infected with the first signs appearing in about 24 hours.[4]

Principles of treatment:(Figure 134.11)



Figure 134.11 Dog bite treated by antibiotics, sterile dressing and anti-tetanus vaccination

- Clean and debride the wound with aqueous antiseptic, allowing it to soak for 10 - 20 minutes.
- Aim for open healing - avoid suturing if possible (except in 'privileged' sites with an excellent blood supply such as the face and scalp).
- Apply non-adherent, absorbent dressings (paraffin gauze and Melolin) to absorb the discharge from the wound.
- Tetanus prophylaxis: immunoglobulin or tetanus toxoid.
- Give prophylactic penicillin for a severe or deep bite: 1.5 million units procaine penicillin IM statim, then orally for 5 - 10 days. An alternative is amoxycillin/clavulanate for 5-7 days. Use this antibiotic for 7-10 days for an established infection (depending on swalx)[4]
- Inform the patient that slow healing and scarring are likely.

Rabid or possibly rabid dog (or other animal)

(not currently applicable in Australia)

- Wash the site immediately with detergent or saline (preferable) or hydrogen peroxide or soap (if no other option).
- Do not suture.
- If rabid:
  - human rabies immune globin (passive)
  - antirabies vaccine (active)
- Uncertain: capture and observe animal consider vaccination

Cat bites

Cat bites have the greatest potential for suppurative infection with *Pasteurella multocida* being the most common organism. The same principles apply as for the management of human or dog bites. Use amoxycillint davulanate for prophylaxis for 5 days. For infection swab the wound but start with metronidazole + doxycydine or ciprofloxacin.[3] It is important to clean a deep and penetrating wound. Another problem is cat-scratch disorder, presumably caused by a Gram-negative bacterium, Bartonella henselae.

Clinical features of cat-scratch disorder

- An infected ulcer or papule pustule at bite site (30% of cases) after 3 days or so[5]

- 1 - 3 weeks later: fever, headache, malaise regional lymphadenopathy (may suppurate)
- Intradermal skin test positive
- Benign self-limiting course
- Sometimes severe symptoms for weeks, especially in immunocompromised
- Treat with erythromycin or roxithromycin for 10 days.[3]

Coral cuts

Wounds from coral cuts are at risk of serious infection with *Vibrio* organisms (marine pathogens) or *Streptococcus pyogenes*. Such wounds require cleaning with antiseptics, debridement, dressing and antibiotic cover with doxycycline 100 mg bd or cephalexin 500 mg bd for 7 days.

Scalp lacerations in children

If lacerations are small but gaping use the child's hair for the suture, provided it is long enough.

Method

Make a twisted bunch of the child's own hair on each side of the wound. Tie a reef knot and then an extra holding knot to minimise slipping. Ask an assistant to drop compound benzoin tincture solution (Friar's balsam) on the hair knot. Leave the hair suture long and get the parent to cut the knot in 5 days.

Forehead and other lacerations in children

Despite the temptation, avoid using reinforced paper adhesive strips (Steri-strips) for children with open wounds. They will merely close the dermis and cause a thin, stretched scar. They can be used only for very superficial epidermal wounds in conjunction with sutures.

Adhesive glue for wound adhesion

A tissue adhesive glue can be used successfully to close superficial smooth and clean skin wounds, particularly in children. An expensive commercial preparation Histoacryl (active ingredient enbucrilate) is available, but Super-glue also serves the purpose although sterility and toxicity have to be considered. The glue should be used only for superficial, dry, clean and fresh wounds. No gaps are per-missible with this method. Avoid glues if possible.

Wound anaesthesia in children

New topical preparations which provide surface anaesthesia are being used for wound repair in children. They include lignocaine and prilo-caine mixture (EMLA cream) and adrenaline and cocaine (AC) liquid. Use the latter with caution.

Some practitioners use an ice block to freeze the lacerated site. The child is asked to hold the ice while a suture is rapidly inserted.

Removal of skin sutures

Suture marks are related to the time of retention of the suture, its tension and position. The objective is to remove the sutures as early as possible, as soon as their purpose is achieved. The timing of removal is based on common-sense and individual cases. Nylon sutures are less reactive and can be left for longer periods. After suture removal it is advisable to support the wound with micropore skin tape (e.g. Steri-strips) for 1 - 2 weeks, especially in areas of skin tension.

Method

1. Use good light and have the patient lying comfortably.
2. Use fine, sharp scissors which cut to the point or the tip of a scalpel blade, and a pair of fine, non-toothed dissecting forceps that grip firmly.
3. Cut the suture close to the skin below the knot with scissors or a scalpel tip (Figure 134.12a).



Figure 134.12 Removal of skin sutures: (a) cutting the suture; (b) removal by pulling towards wound

4. Gently pull the suture out towards the side on which it was divided - that is, always towards the wound (Figure 134.12b).

When to remove non-absorbable sutures

For removal of sutures after non-complicated wound closure in adults, see Table 134.2.

Table 134.2 Time after insertion for removal of sutures

Area	Days later
Scalp	6
Face	3 (or alternate at 2, rest 3 - 4)
Ear	5
Neck	4 (or alternate at 3, rest 4)
Chest	8
Arm (including hand and fingers)	8 - 10
Abdomen	8 - 10 (tension 12 - 14)
Back	12
Inguinal and scrotal	7
Perineum	2
Legs	10
Knees and calf	12
Foot (including toes)	10 - 12

Note: Decisions need to be individualised according to the nature of the wound and health of the patient and healing. In general, take sutures out as soon as possible. One way of achieving this is to remove alternate sutures a day or two earlier and remove the rest at the usual time. Steri-strips can then be used to maintain closure and healing.

Additional aspects

In children, tend to remove 1 - 2 days earlier. Allow additional time for backs and legs, especially the calf. Nylon sutures can be left longer because they are less reactive. Alternate sutures may be removed earlier (e.g. face in women).

BURNS[5,6]

Management depends on extent and depth (burns are classified as superficial or deep - otherwise first, second or third degree).

First-degree burns are superficial and involve only the epidermis, causing pain, redness and swelling. A scald, which is a burn caused by moist heat, is an example. Healing proceeds quickly.

Second-degree or partial skin thickness burns cause the epidermis to blister and become necrotic with subsequent serous ooze.

In *third-degree* or full thickness burns, there is deep necrosis and perhaps anaesthesia from destroyed nerve endings. If extensive (> 9% of body surface area) and deep there is a possibility of hypovolaemia and shock.

First aid

The immediate treatment of burns, especially for smaller areas, is immersion in cold running water such as tap water, for a minimum of 10 - 15 minutes (ideally 20 minutes). Do not disturb charred adherent clothing but remove wet clothing

Chemical burns should be liberally irrigated with water. Apply 1 in 10 diluted vinegar to alkali burns and sodium bicarbonate solution for acid burns.

Refer the following burns to hospital:

- > 9% surface area, especially in a child (see rule of 9s, p. 1149).
- > 5% in an infant
- all deep burns
- burns of difficult or vital areas (e.g. face, hands, perineum/genitalia, feet)
- burns with potential problems (e.g. electrical, chemical, circumferential)
- suspicion of inhalational injury

Always give adequate pain relief. During transport, continue cooling by using a fine mist water spray.

Treatment

1. *Very superficial - intact skin.* Can be left with application of a mild antiseptic only. e.g. aqueous chlorhexidine. Review if blistering.
2. *Superficial - blistered skin.* Apply a dressing to promote epithelialisation (e.g. hydrocolloid sheets, hydrogel sheets) covered by an absorbent dressing or (best option) a retention adhesive material (e.g. Fixomull) Mefix, Hyperfix with daily or twice daily cleaning of the serous ooze and reapplication of outer stretch bandage. Fixomull can be left in place for up to 2 weeks.

Guidelines to patient for retention dressings

- First 24 hours: keep dry. If there is any ooze coming through the dressing, pat dry with a clean tissue.
- First day 2: wash over dressing twice daily. Use gentle soap and water, rinse then pat dry. Do not soak. Rinse only. Do not remove the dressing as it may cause pain and damage to the wound. If the wound becomes red, hot or swollen or if pain increases return to the clinic.
- From day 7: return to the clinic for removal of the dressing. Two hours prior to coming into the clinic, soak the

dressing with olive oil then coven with Glad Wrap.

Note: Dressing must be soaked off with oil (e.g. olive, baby, citrus or peanut). Debride ‘popped blisters’. Only pop blisters that interfere with dermal circulation.

1. *Deep burns*. If considerable ooze, apply the following in order
    - Solosite gel
    - non-adherent neutral dressing (e.g. Melolin)
    - layer of absorbent gauze or cotton wool (larger burns)

Change every 2 - 4 days with analgesic cover. Surgical treatment, including skin grafting, may be necessary.

Exposure (open method)

- Keep open without dressings (good for face, perineum or single surface burns)
- Renew coating of antiseptic cream every 24 hours

Dressings (closed method)

- Suitable for circumferential wounds
- Cover area with non-adherent tulle (e.g. paraffin gauze)
- Dress with an absorbent bulky layer of gauze and wool
- Use a plaster splint if necessary

Burns to hands

For superficial blistered burn to the hand or similar complex' shaped parts of the body apply strips of the retention stretch adhesive dressings as described above. They conform well to digits. Apply an outer bandage. At 7 days soak the dressings in oil for 2 hours prior to coming into the clinic.

FOREIGN BODIES

Penetrating gun injuries

Injuries to the body from various types of guns present decision dilemmas for the treating doctor. The following information represents guidelines, including special sources of danger to tissues from various foreign materials discharged by guns.

Gunshot wounds

Airgun

The rule is to remove subcutaneous slugs but to leave deeper slugs unless they lie within and around vital structures (e.g. the wrist). A special common problem is that of slugs in the orbit. These often do little damage and can be left alone, but referral to an ophthalmologist would be appropriate.

0.22 rifle (the pea rifle)

The same principles of management apply but the bullet must be localised precisely by X-ray. Of particular interest are abdominal wounds, which should be observed carefully, as visceral perforations can occur with minimal initial symptoms and signs.

0.410 shotgun

The pellets from this shotgun are usually only dangerous when penetrating from a close range. Again, the rule is not to remove deep-lying pellets - perhaps only those superficial pellets that can be palpated.

Pressure gun injuries

Injection of grease, oil, paint and similar substances from pressure guns (Figure 134.13) can cause very serious injuries, requiring de-compression and removal of the substances.


oil injections into digital pulp
high-pressure grease or paint gun injections

Figure 134.13 Dangerous accidential injections into the hand

Grease gun and paint gun

High-pressure injection of paint or grease into the hand requires urgent surgery if amputation is to be avoided. There is a deceptively minor wound to show for this injury, and after a while the hand feels comfortable. However, ischaemia,[1] chemical irritation and infection can follow, with gangrene of the digits, resulting in, at best, a claw hand due to sclerosis. Treatment is by immediate decompression and meticulous removal of all foreign material and necrotic tissue.

Oil injection

Accidental injection of an inoculum in an oily vehicle into the hand also creates a serious problem with local tissue necrosis. If injected into the digital pulp, this may necessitate amputation. Such injections are common on poultry farms, where many fowl-pest injections are administered.

Splinters under the skin

The splinter under the skin is a common and difficult procedural problem. Instead of using forceps or making a wider excision, use a disposable hypodermic needle to 'spear' the splinter (Figure 134.14) and then use it as a lever to ease the splinter out through the skin.



Figure 134.14 Removal of splinters in the skin

A ‘buried’ wooden foreign body can be detected by ultrasound.

Embedded fish hooks

Two methods of removing fish hooks are presented here, both requiring removal in the reverse direction, against the barb. Method 2 is recommended as first-line management.

Method 1

1. Inject 1 - 2 mL of LA around the fish hook.
2. Grasp the shank of the hook with strong artery forceps.
3. Slide a D11 scalpel blade in along the hook, sharp edge away from the hook, to cut the tissue and free the barb (Figure 134.15).



Figure 134.15 Removal of fish hooks by cutting a path in the skin

4. Withdraw the hook with the forceps.

Method 2

This method, used by some fishers, relies on a loop of cord or fishing line to forcibly disengage and extract the hook intact. It requires no anaesthesia and no instruments - only nerves of steel, especially for the first attempt.

1. Take a piece of string about 10 - 12 cm long and make a loop. One end slips around the hook, the other hooking around one finger of the operator.
2. Depress the shank with the other hand in the direction that tends to disengage the barb.
3. At this point give a very swift, sharp tug along the cord.
4. The hook flies out painlessly in the direction of the tug (Figure 134.16).



Figure 134.16 Fisherman's method of removing a fish hook intact

Note: You must be bold, decisive, confident and quick - half-hearted attempts do not work.

For difficult cases, some local anaesthetic infiltration may be appropriate. Instead of a short loop of cord, a long piece of fishing line double looped around the hook and tugged by the hand, or flicked with a thin ruler in the loop, will work.

NEEDLESTICKS AND SHARPS INJURIES

Accidental skin puncture by contaminated ‘sharps’, including needles (with blood or bloodstained body fluids), is of great concern to all health care workers. Another problem that occurs occasionally is the deliberate inoculation of people such as police by angry sociopathic individuals. A needlestick accident is the commonest incident with the potential to transmit infections such as HIV and hepatitis B, C or D. The part of the venipuncture that is most likely to cause the accident is the recapping or resheathing of the needle. This practice should be discouraged.

Infections transmitted by needlestick accidents are summarised in Table 134.3. The risk from a contaminated patient is greatest with hepatitis (10 - 30%), while the risk of seroconversion or clinical infection after a needlestick injury with HIV-positive blood is very low (probably about 1 in 300).[7]

Table 134.3 Infections transmitted by needlestick accidents

Viruses
HIV
Hepatitis B, C, D
Herpes simplex
Herpes varicella zoster
Bacteria
Streptococcal infections
Staphylococcal infections
Syphilis
Tuberculosis
Other
Malaria

Prevention

- Avoid physically struggling with overdose victims or high-risk patients for lavage or venipuncture.
- Needles should not be recapped.
- Dispose of needles immediately and directly into a leak-proof, puncture-proof sharps container.
- Avoid contact with blood.
- Wear protective gloves (does not prevent sharps injury).

Management

- Squeeze out and wash under running tap water with soap and/or dilute sodium hypochlorite solution (e.g. Milton).
- Encourage bleeding.
- Obtain information about and blood from the sharps' victim and the source person (source of body fluid). A known carrier of hepatitis B surface antigen or an HIV-positive source person will facilitate early decision making.

Note: It takes 3 months to seroconvert with HIV so the patient may be infected but negative on initial tests.

Known hepatitis B carrier source person

- If injured person immune - no further action
- If non-vaccinated and non-immune
  - give hyperimmune hepatitis B gammaglobulin (within 48 hours)
  - commence course of hepatitis B vaccination, within 24 hours.

Known hepatitis C carrier source person[8]

- The recipient needs to have follow up HCV antibody tests at 1 and 6 weeks and ALT levels at 4-6 months.
- There is no effective immunoprophylaxis available. Consider early therapy should seroconversion occur.

Known HIV-positive source person[8]

Refer to consultant about relative merits of drug prophylaxis and serological monitoring.

Options

- low risk zidovudine + lamivudine for 4 weeks
- high risk zidovudine (AZT) + lamivudine + indinavir or nelfinavir prophylaxis within 8 hours, preferably within 1 - 2 hours[9], for 6 weeks *or* serological monitoring 0, 4, 6, 12, 24 and 52 weeks[10] (check with consultant and refer to national guidelines)

Unknown risk source person

Take source person's blood (if consent is given) and sharps' victim's blood for hepatitis B (HBsAg and anti-HBs) and hepatitis C and (if high risk for HIV) HIV status tests. Commence hepatitis B vaccination if not vaccinated.

Note: Informed consent for testing and dis-closure of test results for involved person should be obtained.

TETANUS PROPHYLAXIS

Tetanus is a very serious disease but completely preventable by active immunisation. Protection should be universal, especially if the childhood immunisation program is followed. However, all patients with wounds should be assessed for their tetanus status and managed on their merits. For severe wounds the possibility of gas gangrene should also be considered.

Tetanus-prone wounds:

- compound fractures
- penetrating injuries
- foreign bodies
- extensive crushing
- delayed debridement
- severe burns
- pyogenic infection

For the primary immunisation of adults, tetanus toxoid (singly or combined with diphtheria if primary childhood course not given) is given as two doses 6 weeks apart with a third dose 6 months later. Booster doses of tetanus toxoid are given every 10 years or at the time of major injury occurring 5 years after previous dose.

Practice tips

- Have the patient lying down for suturing and parents of children sitting down.
- Avoid using antibiotic sprays and powders in simple wounds - resistant organisms can develop.
- Consider tetanus and gas gangrene prophylaxis in contaminated and deep necrotic wounds.
- Give a tetanus booster if patient has not had one within 5 years for dirty wounds or within 10 years with clean wounds.

- Give tetanus immunoglobulin if patient is not immunised or if the wound is grossly contaminated.
- Never send patients home before thoroughly washing their hair and carefully examining for other lacerations.
- Any laceration in the cheek, mandible or lower eyelid may damage the facial nerve, parotid duct or lacrimal duct respectively.
- When a patient falls onto glass it takes bone to halt its cutting path. Assume all structures between skin and bone are severed.

Passive immunisation

Passive immunisation, in the form of tetanus immunoglobulin 250 units by IM injection, is reserved for non-immunised individuals or those of uncertain immunity wherever the wound is contaminated or has devitalised tissue. Wounds at risk include those con-taminated with dirt, faeces/manure, soil, saliva or other foreign material; puncture wounds; wounds from missiles, crushes and burns.

The guide is outlined in Table 134.4.

Table 134.4 Guide to tetanus prophylaxis in wound management[3]

Time since vaccination	Type of wound	Tetanus toxoid	Tetanus immunoglobulin
History of 3 or more doses of tetanus toxoid			
< 5 years	All wounds	No	No
5 to 10 years	Clean minor wounds	No	No
	All other wounds	Yes	No
> 10 years	All wounds	Yes	No
Uncertain vaccination history or less than three doses of tetanus toxoid			
	Clean minor wounds	Yes	No
	All other wounds	Yes	Yes

REFERENCES

1. Hansen G. *Practice Tips*. Aust Fam Physician, 1982; 11: 867.
2. Medeiros/, Saconato H. Antibiotic prophylaxis for mammalian bites. The Co chrane Database of Systematic Reviews 2001 Issue 2. Art. No: CD 001738. DO1: 10.1002/14651858 CD 001738.
3. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 12). Melbourne: Therapeutic Guidelines Ltd, 2006; 271 - 2.
4. Brown J, Woods ML. Management of bite injuries, Australian Prescriber 2006;29 6 - 8.
5. Tierney LM et al. *Current Medical Diagnosis and Treatment* (41st edn). New York: McGraw Hill and Lange, 2006: 1433.
6. Mashford ML (Chair). *Therapeutic Guidelines: Dermatology* (Version 2). Melbourne: Therapeutic Guidelines Ltd, 2004: 71 - 3
7. Spelman D. Transmission of infection by needlesticks, sharps and splashes. Aust Fam Physician, 1988; 17: 681.
8. Spicer J (Chair). *Therapeutic Guidelines: Antibiotic* (Version 12). Melbourne: Therapeutic Guidelines Ltd, 2006: 183.
9. Gerberding JL. Drug therapy: management of occupational exposures to blood-borne viruses. N Engl J Med, 1995; 332: 444.
10. Hammond L. AIDS and hepatitis B protection strategies. Aust Fam Physician, 1990; 19: 657 - 61.

CHAPTER 135. COMMON FRACTURES AND DISLOCATIONS

The broken bone, once set together, is stronger than ever.

JOHN LYLY (1554 - 1606)

COMMON FRACTURES AND DISLOCATIONS

Common fractures and dislocations usually apply to the limbs, the shoulder girdle and the pelvic girdle and their management requires an early diagnosis to ensure optimum treatment and to prevent complications. Early diagnosis depends on the physician being vigilant and on having knowledge of the less common conditions so that a careful search for the diagnosis can be made.

The diagnosis is dependent on a good history followed by a careful examination, good-quality X-rays appropriate to the injury (e.g. stress view) and, if necessary, special investigations. The golden rule is: if in doubt-X-ray. The family doctor should develop the habit of looking at a patient's X-rays. Such a back-up to the radiologist's report can help avoid missed diagnoses.

There are many pitfalls involved in managing fractures and dislocations. Many injuries, such as fractures of the arm and hand, seem trivial but they can lead to long-term disability. This chapter presents guidelines to help avoid these pitfalls.

KEY FACTS AND CHECKPOINTS

- A fracture usually causes deformity but may cause nothing more than local tenderness over the bone (e.g. scaphoid fracture, impacted fractured neck of femur).
- The classic signs of fracture are:
  - pain
  - tenderness
  - loss of function
  - deformity
  - swelling/bruising
  - crepitus
- X-ray examination of the affected area of the upper limb should include views of joints proximal or distal to the site of the injury, and X-rays in both AP and lateral planes.
- If an X-ray is reported as normal but a fracture is strongly suspected, an option is to splint the affected limb for about 10 days and then repeat the X-ray.
- As a rule, displaced fractures must be reduced whereby bone ends should be placed in proper alignment and then immobilised until union occurs.
- Fractures should be monitored radiologically for loss of position, particularly in the first 1 - 2 weeks following reduction.
- Bone union is assessed clinically by reduced pain at the fracture site and reduced fracture mobility. It is assessed radiologically by X-ray features such as trabecular continuity across the fracture site and bridging callus.
- Non-union is caused by such factors as inadequate immobilisation, excessive distraction, loss of healing callus, infection or avascular necrosis.
- Stiffness of joints is a common problem with immobilisation in plaster casts and slings so the joints must be moved as early as possible. Early use is possible if the fracture is stable.
- A dislocation is a complete disruption of one bone relative to another at a joint.
- A subluxation is a partial displacement such that the joint surfaces are still in partial contact.
- A sprain is a partial disruption of a ligament or capsule of a joint.
- Always consider associated soft-tissue injuries such as neuropraxia to adjacent nerves, vascular injuries and muscle compartment syndromes.
- A stress fracture is an incomplete fracture resulting from repeated small episodes of trauma, which individually would

be insufficient to damage the bone. Stress fractures, especially in the foot, are most likely to result from sport, ballet, gymnastics and aerobics. Their incidence rises sharply at times of increased activity.[1]

- Typical stress fractures (with their usual cause) include:
  - navicular (sprinting sports, football)
  - metatarsal neck (running, walking, basketball, jumping)
  - base of fifth metatarsal (dancing)
  - femur-neck or shaft (distance running)
  - ulna (weight-lifting)
  - distal radial and ulnar epiphyses (gymnastics)
  - talus (running)
  - proximal tibia (running, football)
  - medial tibia (running, football)
  - distal phalanges (guitar playing)
  - cervical spinous process (gardening)
  - lumbar vertebrae-pars interarticularis (fast bowling)
  - spiral humerus (throwing sports)
  - rib fractures-1st (weight-lifting)
  - rib fractures-8th (tennis)
- The key strategy of most reduction manoeuvres is tradtion especialy for dislocations. This may be supplemented with translation or leverage.

TESTING FOR FRACTURES[2]

This method describes the simple principle of applying axial compression for the clinical diagnosis of fractures of bones of the forearm and hand, but also applies to bones of the limbs.

Many fractures are obvious when applying the classic methods of diagnosis but it is sometimes more difficult if there is associated soft-tissue injury from a blow or if there is only a minor fracture such as a greenstick fracture of the distal radius.

If the bone suspected of fracture is compressed gently from end to end, the patient will feel pain. A soft-tissue injury of the forearm will show pain, tenderness, swelling and possibly loss of function. It will, however, not be painful if the bone is compressed axially-that is, in its long axis.

Walking is another method of applying axial compression, and this is very difficult (because of pain) in the presence of a fracture in the weight-bearing axis or pelvis.

Method

1. Grasp the affected area both distally and proximally with your hands.
2. Compress along the long axis of the bones by pushing in both directions, so that the forces focus on the affected area (fracture site; Figure 135.1a). Alternatively, compression can be applied from the distal end with stabilising counterpressure applied proximally (Figure 135.1b).



Figure 135.1 Testing for fractures: (a) axial compression to detect a fracture of the radius or ulna bones; (b) axial compression to detect a fracture of the metacarpal

3. The patient will accurately localise the pain at the fracture site.

PRINCIPLES OF TREATMENT OF LIMB FRACTURES

To reduce any fracture properly, the following steps must be taken (Figure 135.2a).[3]



Figure 135.2 (a) Principles of reduction of fractured bones **(b)** Principles of moulding to maintain reduction: the arrows indicate the three point pressure areas required to maintain reduction

1. Disimpact the fragments, usually by increasing the deformity.
2. Re-establish the correct length of the bone.
3. Re-establish the correct alignment by proper reduction of the fracture.

The above steps will only be achieved with adequate anaesthesia, analgesia and relaxation. Maintenance of the reduction depends upon the moulding which utilises the intact periosteal bridge to hold the fracture fragments in a reduced position. Figure 128.2b illustrates the principle of moulding to maintain reduction.[3]

FRACTURES AND DISLOCATIONS OF THE FACE AND SKULL

Skull fractures

Closed fractures without any neurological symptoms do not require active intervention. Depressed fractures may require elevation of the depressed fragment. Compound fractures of the vault require careful evaluation and referral. Special care is required over the midline as manipulation (usually by elevation) of any depressed fragment can tear the sagittal sinus, causing profuse and fatal bleeding.[4]

Base of skull fractures

These fractures are difficult to diagnose on radiography but their presence is indicated by bleeding from the nose, throat or ears. CSF may be observed escaping, especially through the nose, if the dura is also torn.

Treatment of basal fractures is based on prevention of intracranial infection and avoidance of excessive interference with the nose or ear, such as with packing and naso-gastric tubes. An appropriate antibiotic is cotrimoxazole.[4]

'Malar' fracture

A fractured zygomaticomaxillary complex (malar) is a common body contact sports injury or injury resulting from a fight.

Clinical features:

- swelling of cheek
- circumocular haematoma
- subconjunctival haemorrhage
- palpable step in infraorbital margin
- flat malar eminence when viewed from above
- paraesthesia due to infraorbital nerve injury
- loss of function (i.e. difficulty opening mouth)

Management

- Head injury assessment
- Exclude 'blow-out' fracture of the orbit
- Exclude ocular trauma
  - remove contact lenses if worn
  - check visual acuity
  - check for diplopia
  - check for hyphaema
  - check for retinal haemorrhage
- Persuade patient not to blow nose (can cause surgical emphysema)
- If fracture displaced, refer for reduction under general anaesthesia

Reduction methods

- Elevation by temporal or intraoral approach-healing can be expected in 3 - 4 weeks
- Some require interosseous wiring or plating or pinning

Fracture of mandible

A fracture of the mandible follows a blow to the jaw. The patient may have swelling (which can vary from virtually none to severe), pain, deformity, inability to chew, malalignment of the jaw and teeth and drooling of saliva. Intra-oral examination is important as submucosal ecchymosis in the floor of the mouth is a patho-gnomonic sign.

A simple office test for a suspected fractured mandible is to ask patients to bite on a wooden tongue depressor (or similar firm object). Ask them to maintain the bite as you twist the spatula. If they have a fracture they cannot hang onto the spatula because of pain.[5]

X-rays:

- AP views and lateral obliques
- an orthopantomogram provides a global view

First aid management

- Check the patient's bite and airway
- Remove any free-floating tooth fragments and retain them
- Replace any avulsed or subluxed teeth in their sockets

Note: Never discard teeth.

- First aid immobilisation with a four-tailed bandage (Figure 135.3)



Figure 135.3 Immobilisation of a fractured mandible in a four-tailed bandage

Treatment

Refer for possible internal fixation.

A fracture of the body of the mandible will usually heal in 6 - 12 weeks (depending on the nature of the fracture and fitness of the patient).

Dislocated jaw

The patient may present with unilateral or bilateral dislocation. The jaw will be 'locked' and the patient unable to articulate or close the mouth.

Method of reduction

- Get the patient to sit upright with the head against the wall.
- Wrap a handkerchief around both thumbs and place the thumbs over the lower molar teeth, with the fingers firmly grasping the mandible on the outside.
- Firmly thrusting with the thumbs, push downwards towards the floor (Figure 135.4).



Figure 135.4 Method of reduction of a dislocated jaw by downward traction on the mandible

This action invariably reduces the dislocation, but the reduction can be reinforced by the fingers rotating the mandible upward as the thumbs thrust downwards.

FRACTURED VERTEBRAE/SPINAL COLUMN

Cervical fractures, especially atlas (C1), axis (C2) and odontoid process, require early referral with the neck immobilised in a cervical collar, in a supine position. A hard collar is preferred but a soft collar with sandbags either side of the head to prevent movement will suffice.

Thoracolumbar fractures

Fractures or fracture dislocations of the thoracic and lumbar vertebrae, without neurological deficit, are classified as either stable or unstable.

Stable fractures

- Compression fractures of vertebral body with < 50% loss of vertical height
- Minor fractures
- Laminar fractures

Treatment: rest on firm-to-hard bed for 10 - 28 days depending on symptoms, followed by a brace.

Special problems:

- retroperitoneal haematoma
- paralytic ileus
- associated kidney rupture with L1 fractures
- underlying vertebral body pathology in the elderly (e.g. myeloma or metastases)

Unstable fractures

Burst fractures and shearing fractures are usually unstable. They are often associated with partial or complete paraplegia and require urgent referral.

Fractures of sacrum and coccyx

No treatment apart from symptomatic treatment is required. Manual reduction per rectum can be attempted for significant forward displacement of the coccyx. Advise the use of a rubber ring or special cushion (such as a Sorbo cushion) when sitting. Excision of the coccyx may be considered for persistent discomfort.

FRACTURED RIB

Features:

- pain over the fracture site, especially with deep inspiration and coughing
- localised tenderness and swelling
- pain in the site upon whole chest compression
- X-ray confirms diagnosis and excludes underlying lung damage (e.g. pneumothorax). There is a high incidence of false negative fractures on X-ray, so caution is necessary
- suspect splenic, hepatic and renal trauma with lower rib fractures

Treatment

A simple rib fracture can be extremely painful. The first treatment strategy is to prescribe analgesics, such as paracetamol, and encourage breathing within the limits of pain. If pain persists in cases of single or double rib fracture with no complication, application of a rib support is most helpful.

The universal rib belt

A special elastic rib belt can provide thoracic support and mild compression for fractured ribs (Figure 135.5). Despite its flexibility it gives excellent support and symptom relief while permitting adequate lung expansion. The elastic belt is 15 cm wide and has a Velcro grip fastening, so it can be applied to a variety of chest sizes.



Figure 135.5 Method of application of a universal rib belt

Healing time

3 - 6 weeks; local discomfort may persist much longer.

FRACTURES OF THE CLAVICLE

There is a history of a fall onto the outstretched hand or elbow although this fracture may also occur with a direct blow to the clavicle or the point of the shoulder. The patient has pain aggravated by shoulder movement and usually supports the arm at the elbow and clasped to the chest. The most common fracture site is at the junction of the outer and middle thirds, or in the middle third. Consider the possibility of neurovascular injury.

Treatment

- St John's elevated sling to support arm-for 3 weeks
- Figure of eight bandage (used mainly for severe discomfort)
- Early active exercises to elbow, wrist and fingers
- Active shoulder movements as early as possible

Special problem

Fracture at the lateral end of the bone: this fracture is often subject to delayed or non-union. Consider referral for open reduction.

Healing time

4 - 8 weeks.

The appropriate use of slings for fracture-dislocations is presented in Table 135.1.

Table 135.1 Appropriate use of slings for fracture-dislocations

Collar and cuff	Fractured shaft of humerus
Broad arm sling	Fractured forearm
	Fractured scapula
St John's high sling	Fractured clavicle
	Fractured neck of humerus
	Subluxed acromioclavicular joint
	Dislocated acromioclavicular joint
	Subluxed sternoclavicular joint

FRACTURES OF THE SCAPULA

Fractures of the scapula may include:

- body of scapula: due to a crushing force, considerable blood loss, may be rib fractures
- neck of scapula (may involve joint)
- acromion process (due to a blow or fall on the shoulder)
- coracoid process (due to a blow or fall on the shoulder)

Treatment

- Broad-based triangular sling for comfort
- Early active exercises for shoulder, elbow and fingers as soon as tolerable

- A large glenoid fragment usually requires surgical reduction because of potential glenohumeral joint instability

Healing time

Several weeks to months.

FRACTURES OF THE STERNUM

These are treated symptomatically with analgesics but careful evaluation of thoracic injuries, including cardiac tamponade or myo-cardial contusion, is essential. A significantly depressed fracture should be referred. An ECG is advisable.

ACROMIOCLAVICULAR JOINT DISLOCATION/SUBLUXATION

A fall on the shoulder, elbow or outstretched arm can cause varying degrees of separation of the acromioclavicular (AC) joint, causing the lateral end of the clavicle to be displaced upwards.

- Grades I, II partial separation, involving tearing of the AC capsule and ligaments
- Grade III complete tearing, also affecting the coracoclavicular ligaments

Treatment

- Analgesics
- St John's arm sling (suitable for all injuries)
- Mobilisation exercises as soon as possible and (for grade III)



Figure 135.6 Subluxation of the acromioclavicular joint. Typical appearance

- A compression bandage (or long straps of adhesive low-stretch strapping) with padding at pressure points-elbows, clavicle and coracoid. The clavicle should be manipulated into its correct position and the forearm elevated: applying pressure from above (clavicle) and below (elbow) to achieve compression, apply a bandage over the outer end of the clavicle and round the elbow joint which is flexed to 90°. The bandage or strapping is worn for 2 - 3 weeks.[6]
- Many patients are unable to tolerate this method of treatment. Skin irritation or blisters are common. This occurs

particularly with adhesive strapping and the deformity commonly requires correction after the removal of bandage or strapping. The same effect may be achieved with an orthotic device known as a Kenny-Howard sling or brace.

- The issue of internal fixation versus conservative treatment for a complete dislocation is controversial in that the bulk of patients treated conservatively have minimal residual symptoms. However, a significant proportion have residual symptoms in the form of AC joint pain and traction effects on the brachial plexus due to loss of scapular suspension. The patients most likely to have these symptoms are those with high grades of separation, involvement of the dominant shoulder and participation in employment or sports that place heavy physical demands on the shoulder girdle. If in doubt, referral within the first few weeks of injury for consideration of the pros and cons of conservative versus surgical treatment would be appropriate.

STERNOCLAVICULAR JOINT DISLOCATION/SUBLUXATION

This uncommon injury is caused by a fall or very heavy impact on the shoulder, causing the medial end of the clavicle to move forwards or anterior (making it prominent) or backwards. Plain X-rays are difficult to interpret but a CT scan is the ideal diagnostic method.

Special problem

Backward (inward) displacement of the clavicular end with danger to major blood vessels and trachea. This is one of the few potentially life-threatening orthopaedic injuries. Urgent referral for reduction is essential especially if stridor or venous obstruction is present. A first aid measure is to place a sand bag between the shoulders with the patient supine and extending the abducted arm on the affected side.[7] Closed reduction can usually be achieved under anaesthesia. The reduction is nearly always stable.

Treatment

Forward subluxation or disclocation, unlike posterior dislocation, is nearly always unstable and resists attempts at maintaining closed reduction. Despite the persistence of a medial clavicular swelling, most patients need a sling for only 1 - 2 weeks and the bulk of their pain settles over the following months. Surgery is generally only indicated for an unusually painful and chronic anterior sternoclavicular dislocation.

DISLOCATION OF THE SHOULDER

Dislocations of the shoulder joint can be caused by an impact on the arm by falling directly on the outer aspect of the shoulder, or by a direct violent impact, or by a forceful wrenching of the arm outwards and backwards.

Types of dislocation:

- anterior (forward and downward)-95% of dislocations
- posterior (backward)-diagnosis often overlooked
- recurrent anterior dislocation (recurrent posterior dislocation extremely rare)

Anterior dislocation of the shoulder

Management

AP and lateral X-rays should be undertaken to check the position and exclude an associated fracture. The arm should be assessed for the presence of neurological injury before reduction. Reduction can be achieved under general anaesthesia (easier and more comfortable) or with intravenous pethidine ± diazepam. The following methods can be used for anterior dislocation. Satisfactory analgesia and patient relaxation are vital to the success of any of the methods.

Kocher method

- Elbow flexed to 90° and held close to the body
- Slowly rotate arm laterally (externally)
- Adduct humerus across the body by carrying point of elbow while simultaneously applying longitudinal traction along the line of the humerus
- Rotate arm medially (internally)

Hippocratic method

Apply traction to the outstretched arm by a hold on the hand with countertraction from stockinged foot in the medial wall of the axilla. This levers the head of the humerus back. It is a good method if there is an associated avulsion fracture of the greater tuberosity.

Milch method (does not require anaesthesia or sedation)

- Patient reclines at 30° and with guidance slowly bends the elbow to 90°.
- Operator braces thumb against humeral head.
- The patient is asked to lift the arm (abduct) slowly with the elbow bent so that they can pat the back of their head (requires considerable reassurance and encouragement).
- At this position of external rotation, traction along the line of the humerus (with countertraction) achieves reduction.

Dependent arm in prone position method

- The patient is placed in prone position on bed or trolley with arm hanging limply over the edge and elbow fully extended.
- The shoulder may reduce spontaneously, especially with adequate analgesia.
- The technique may be enhanced by longitudinal traction applied to the arm.

Postreduction

- Reduction is complete if the hand can rest comfortably on the opposite shoulder.
- Confirm reduction by X-ray in two planes and again assess for unsuspected fractures (e.g. glenoid rim or greater tuberosity fractures).
- Keep the arm in a sling for 2 weeks.
- Apply a swathe bandage to the chest wall.
- After immobilisation, begin pendulum and circumduction exercises.
- Combined abduction and lateral rotation should be avoided for 3 weeks.

Posterior dislocation of the shoulder

This is the most commonly misdiagnosed major joint dislocation.[8] Posterior dislocation most often follows an epileptic seizure or electrical shock. The post-ictal patient with a painful shoulder has a posterior dislocation of the shoulder until proven otherwise. Less often this injury is caused by a fall onto the outstretched hand with the arm internally rotated or by a direct blow to the front of the shoulder. If any doubt persists about the diagnosis, a CT scan is appropriate.

The shoulder contour may look normal but the major clinical sign is painful restriction of external rotation, which is usually completely blocked. Beware of the problem of pain in the shoulder after a convulsion. An ‘axillary shoot through’ X-ray view should be routinely ordered following shoulder trauma (Figure 135.7).



Figure 135.7 X-ray (axillary view) illustrating posterior dislocation of the shoulder; the humoral head is pushed backwards with an impaction fracture anteriorly. If the patient prevents satisfactory positioning for X-ray, consider CT scan

Reduction of posterior dislocation

Using appropriate analgesia or anaesthesia, apply traction to the shoulder in 90° of abduction (with the elbow at right angles) and laterally (externally) rotate the limb. Referral for reduction is advisable.

Recurrent anterior dislocation

Acute anterior shoulder dislocation may tear or stretch the anterior capsular ligaments from their bony origin. This may predispose to recurrent anterior dislocation or subluxation. (Recurrent posterior instability is rare.)

A simple procedure for reducing recurrent anterior dislocation:

- Get the patient to sit comfortably on a chair with legs crossed.
- The patient then interlocks hands and elevates the upper knee so that the hands grip the knee.
- The knee is gradually lowered until its full weight is taken by the hands. At the same time the patient has to concentrate on relaxing the muscles of the shoulder girdle. This method usually effects reduction without the use of force.

Recurrent dislocation often requires definitive surgery, depending on the frequency of dislocations and the degree of apprehension between episodes.

The Bankart lesion

Teenagers and young adults who sustain a traumatic dislocation of the shoulders tend to hare the Bankart lesion, which, is avulsion of the anteroinferior capsulolabral complex, leading to a high rate of recurrment dislocation. This should be considered for an arthroscopic anterior stabilisation.

Pitfalls

- Nerve injury, especially axillary (circumflex) nerve.
- A fractured neck of the humerus, especially in the elderly, may mimic a dislocation.
- Associated fractures (greater tuberosity, head of radius, glenoid) may require internal fixation.
- Great difficulty with some reductions (this is often related to inadequate analgesia; the use of excessive force may result in fracture).
- Failing to X-ray all suspected dislocations before and after reduction; failing to obtain an axillary view to show posterior displacement or fractures of the humerus or glenoid.

Orthopaedic problems that cause difficulties in diagnosis and management are outlined in Table 135.2.

Table 135.2 Important orthopaedic problems that cause difficulties in diagnosis and management[8]

Shoulders

• Posterior dislocation of the shoulder
• Recurrent subluxations
• Unstable surgical neck fractures of humerus
• The avascular humeral head
Elbow
• Supracondylar fractures with forearm ischaemia
• Fracture of the lateral humeral condyle in children
• Fractured neck of radius in children
• The Monteggia fracture with dislocation of radial head
Wrist
• Scaphoid fractures
• Scapholunate dislocation
• The unstable Colles' fracture
Fingers
• Phalangeal fractures
• Intra-articular fractures
• Penetrating injuries of the MCP joint
• Gamekeeper's thumb (MCP joint)
The hip
• Developmental dysplasia of the hip
• Septic arthritis
• Slipped capital femoral epiphysis
• Subcapital fractures
• Stress fractures of the femoral neck in athletes
• Impacted subcapital femoral neck fracture in the elderly
Foot and ankle • Talar dome lesions • Stress fractures of the navicular • Intra-articular fractures

FRACTURED GREATER TUBEROSITY OF HUMERUS

Treat with a combination of immediate mobilisation and rest in a sling unless grossly displaced, when surgical reduction is advisable. Shoulder stiffness can be a disabling problem, so early movement is encouraged with review in 7 days. This fracture should be monitored by X-ray within 2 weeks after injury. Undetected displacement may lead to mechanical im-pingement against the acromion. This fracture may also be an indication of the patient having had a transient glenohumeral dislocation.

FRACTURED SURGICAL NECK OF HUMERUS

This usually occurs in the elderly due to a fall onto the outstretched hand. The fragments may be impacted. The greater tuberosity may also be fractured. Watch out for associated dislocation. In adolescents, fracture-separation of the upper humeral epiphysis occurs.

Treatment (no displacement or impaction)

- Triangular sling
- When pain subsides (10 - 14 days) encourage pendulum exercises in the sling
- Aim at full activity within 8 - 12 weeks post injury

Displaced fractures may require internal fixation. Severely comminuted fractures may predispose to post-traumatic osteoarthritis or humeral head avascular necrosis. Referral with a view to prosthetic hemiarthroplasty should be considered.

Healing

Union usually occurs in 4 weeks and consolidation at 6 weeks.

Pitfalls with fractures of the surgical neck

Minimally displaced fractures of the surgical neck of the humerus are usually managed conservatively, but overzealous early mo-bilisation can lead to non-union.[7] If there is a communication of this fracture with joint fluid, movement washes away the fracture haematoma and leads to the development of true pseudo-arthrosis. Judicious early immobilisation will avert this complication.

Always remember the cardinal fracture management rule: ‘First ensure that stability of the fracture is sufficient to allow healing before prescribing rehabilitation exercises or early use of the extremity’.[8]

The management of various humeral fractures is summarised in Figure 135.8.


Management
fracture of greater tuberosity
undisplaced:
• early mobilisation/sling
marked displacement:
• open reduction
fracture of surgical neck
triangular sling
fracture of shaft of humerus
undisplaced:
• collar and cuff
• elbow flexed to 60–70
displaced:
• closed reduction
• then immobilisation
transverse
spiral
intercondylar fractures
refer for reduction (open or closed)

Figure 135.8 Various fractures of the humerus in adults

FRACTURE OF SHAFT OF HUMERUS

Humeral shaft fractures may be:

- spiral-due to a fall on the hand
- transverse or slightly oblique-fall on elbow with arm abducted
- comminuted-heavy blow

Caution: watch for radial nerve palsy.

Treatment

- Perfect bony opposition is not necessary; some overriding is acceptable but distraction of the fragments is not.
- Undisplaced fracture: collar and cuff with elbow flexed to 60 - 70°.
- Significantly displaced humeral shaft fractures may require manipulation under anaesthestic. However, the vast majority of shaft fractures realign to a satisfactory extent under gravitational effects in a sling once muscle spasm and oedema have subsided. A U-shaped hanging cast or slab enhances the gravitational effect and assists splintage.

INTERCONDYLAR FRACTURES IN ADULTS

Intercondylar fractures, which may be T-shaped or Y-shaped, are usually caused by a fall on the point of the elbow which drives the olecranon process upwards, splitting the condyles apart. Fractures involving the joint can cause long-term problems of post-traumatic osteoarthritis and joint stiffness. Referral for reduction (closed or open) is appropriate.

FRACTURES AND AVULSION INJURIES AROUND THE ELBOW JOINT IN CHILDREN

Potentially severe deforming injuries include:

- supracondylar fractures
- fracture of the lateral humeral condyle
- fracture of medial humeral epicondyle (Figure 135.9)



Figure 135.9 Fractures and avulsion injuries around the elbow joint in children

- fracture of neck of radius

Fractures around the elbow in children require referral to consultants experienced in radiology and fracture management.

Supracondylar fractures with forearm ischaemia

Supracondylar fractures represent about half of all elbow fractures in children and most are extension fractures following falls onto the outstretched arm.

Pressure of the displaced bony fragments causing impingement on the brachial artery leads to impending forearm flexor com-partment ischaemia and muscle death. Severe forearm pain is the most significant and important sign of ischaemia. Neuropraxia of the median, radial or ulnar nerves is common. These injuries almost invariably recover.

This diagnosis must always be assumed in displaced supracondylar fractures in children. Thus, it is the general practitioner's re-sponsibility to ensure treatment is expedited.

The fracture is reduced by hyperflexion of the elbow during traction (after lateral displacement has been corrected) and then immobilised in collar and cuff and stockinet vest. The fully flexed elbow with the usually intact posterior periosteal hinge provides fracture stability. Plaster casting is unnecessary and some would suggest contraindicated because of the significant risk of ischaemic contracture. Circulatory status requires monitoring in the first 24 hours following injury. The collar and cuff should be used for 6 weeks. The invariably stiffelbow quickly resolves without a need for formal therapy.

Fracture of the lateral humeral condyle

Fractures of the lateral humeral condyle also result from a fall onto the outstretched arm in children (Figure 135.9). The fracture line passes vertically or obliquely through the lateral condyle and thus crosses the distal humeral growth plate. It occurs in an age group prior to the appearance of the epiphysis of the lateral epicondyle. Pain and swelling over the lateral elbow, but without the gross deformity of a supracondylar fracture of the humerus, could make one suspect this injury. The fracture is commonly overlooked on X-ray. Comparison views of the opposite elbow are particularly helpful in diagnosing this injury.

Recognition of the fracture and early open reduction and internal fixation with wires is vital to reduce the risk of premature plate closure. Such growth plate disturbance may result in a progressive valgus deformity of the elbow and the late development of an ulnar nerve palsy.

Fracture of the medial humeral epicondyle

This problem occurs typically in adolescents following a fall onto the outstretched hand. The medial epicondyle may be avulsed by massive flexor pronator muscle contraction together with abduction stresses on the forearm. Avulsion of the epicondyle occurs in the young patient before the epiphysis is united. If displaced this fracture is best treated by open reduction and internal fixation. Untreated injuries commonly result in non-union, elbow pain and restricted elbow extension.

Fractured neck of radius

This fracture is caused by a child falling on to the outstretched hand. The fracture line is transverse and is situated immediately distal to the epiphysis.

The degree of tilt is critical. Up to 15° of tilt is acceptable but, beyond that, reduction (preferably closed) will be necessary. The head of the radius must never be excised in children.

DISLOCATED ELBOW

A dislocated elbow is caused by a fall on to the outstretched hand, forcing the forearm backwards to result in posterior and lateral displacement (Figures 135.10, 135.11). The peripheral pulses and sensation in the hand must be assessed carefully. It may result in vascular injury to the brachial artery or injury to the median and ulnar news Check the function of the ulnar nerve before and after reduction.



Figure 135.10 Dislocated elbow: uncomplicated posterior dislocation



Figure 135.11 Dislocated elbow showing posterior displacement of the ulna and radius, wih the typical deformity

Treatment

Attempt reduction with patient fully relaxed under anesthesia. It is important to apply traction to the flexed elbow but allowing it to extend approximately 20 - 30° to enable correction of the lateral displacement and then the posterior displacement.

Follow-up

Encourage early mobilisation with gentle exercises in between resting the elbow for 2 - 3 weeks in a collar and cuff with the elbow flexed above 90°, avoiding passive movements. A plaster cast should not be used because of the risk of ischaemic necrosis of muscle. This will minimise the possibility of myositis ossificans. Recurrent dislocation of the elbow is uncommon.

A simple method of reduction

This method reduces an uncomplicated posterior dislocation of the elbow without the need for anaesthesia or an assistant. The manipulation must be gentle and without sudden movement.

Method

1. The patient lies prone on a stretcher or couch, with the forearm dangling towards the floor.
2. Grasp the wrist and slowly apply traction in the direction of the long axis of the forearm (Figure 135.12).



Figure 135.12 Dislocated elbow: method of reduction by traction on the dependent arm

3. When the muscles feel relaxed (this might take several minutes), use the thumb and index finger of the other hand to grasp the olecranon and guide it to a reduced position, correcting any lateral shift.

Pitfalls

- Incomplete reduction: ulna articulates with capitellum and not the trochlea
- Injury to ulnar nerve (spontaneous recovery usually occurs after 6 - 8 weeks)
- Associated fractures (e.g. coronoid process), which may cause instability

FRACTURED HEAD OF RADIUS (ADULTS)

If the fracture is very slight and undisplaced, treat conservatively with the elbow at right angles in a collar and cuff until pain subsides sufficiently to allow flexion/extension and pronation/supination exercises.

Elbow stiffness is a major problem even after apparently trivial radial head fractures. Early mobilisation is vital. Excision of the radial head should be considered for highly comminuted fractures which limit the ability to mobilise the elbow early or predispose to post-traumatic osteoarthritis. Associated distal radioulnar joint or wrist injuries are often overlooked.

FRACTURED OLECRANON

- Comminuted fracture (with little displacement): sling for 3 weeks and active movements.
- Transverse (gap) fracture: open reduction with screw or wire.

MONTEGGIA FRACTURE-DISLOCATION OF THE RADIAL HEAD

Fractures of the proximal third of the ulna with dislocation of the radial head (Monteggia fracture-dislocation) (Figure 135.13) has a history of mismanagement during treatment. The radial head dislocation is easily overlooked.



Figure 135.13 Monteggia fracture-dislocation of the radial head; it is important not to miss a dislocated head of radius with a fracture of the proximal third of the ulna

Redislocation or subluxation of the radial head is common.

Since surgical intervention is advisable, referral of displaced forearm fractures for early surgery is recommended. Surgical plating of the ulnar shaft maintains the radial head in a reduced position. Follow-up X-rays are mandatory to ensure that there has not been a late redislocation of the radial head.

FRACTURE-DISLOCATION IN THE LOWER FOREARM (GALEAZZI INJURY)

This injury is usually caused by a fall on to the hand and is a combination of a fractured radius (at the junction of its middle and distal thirds) and subluxation of the distal radioulnar joint. The patient should be referred, as open reduction is often required.

FRACTURES OF THE RADIUS AND ULNA SHAFTS

General features

It is more common to have both bones broken. Displaced fractures of both forearm bones in the adult require perfect reduction, which can generally only be achieved by surgical reduction and plating. Less than satisfactory reduction interferes with normal pronation and supination. A fracture of one bone alone is uncommon and usually caused by a direct blow. For a fracture of one bone alone look for evidence of an associated dislocation of the other forearm bone. In children, greenstick fractures are common. Fractured radial shafts tend to slip and ulna fractures heal slowly. Dislocation of the head of the radius or inferior radioulnar joint can be missed if X-rays do not include the elbows and wrist joints.

Reduction

- A greenstick fracture is readily straightened by firm pressure.
- A complete fracture (spiral or transverse) is reduced by traction and rotation.
- A slight overlap and angulation is permissible in children but perfect reduction is essential in adults.
- A plaster cast should include both the elbow and the wrist joints.

Healing time: (adults) spiral fracture-6 weeks; transverse fracture-12 weeks

COLLES' FRACTURE OF LOWER END OF RADIUS

A Colles' fracture, probably the most common of all fractures, is a supination fracture of the distal 3 cm of the radius, caused by a fall onto the outstretched hand.

Clinical features

- Usually an elderly woman
- Osteoporosis is common
- Fall on dorsiflexed hand
- Fracture features
  - impaction
  - posterior displacement and angulation
  - lateral displacement and angulation
  - supination
  - dinner fork deformity (Figure 135.14)



Figure 135.14 Dinner fork deformity of Colles' fracture: a fracture of the distal head of the radius showing impaction and posterior displacement and angulation

Treatment

- If minimal displacement-below-elbow plaster for 4 weeks, then a crepe bandage
- If displaced: meticulous reduction under anaesthesia
  - set in flexion 10°, ulnar deviation 10° and pronation (Figure 135.15)



Figure 135.15 Ideal position of the forearm in a Colles' plaster. Note ulnar deviation, slight flexion and pronation

  - below-elbow plaster 4 - 6 weeks (6 weeks maximum time)
  - unstable fractures may require an above-elbow cast initially with the forearm in pronation
  - check X-ray at 10 - 14 days; position may be lost as swelling subsides and plaster becomes loose

Problems associated with Colles' fracture:

- watch for ruptured extensor pollicus longus tendon
- stiffness of the elbow, MCP joints and IP joints
- discomfort at inferior radioulnar joint due to disruption
- regional pain syndrome

Pitfall: the unstable Colles' fracture[8]

With the advent of modern imaging techniques and power equipment it has become a simple procedure to pin unstable Colles' fractures percutaneously, even in the elderly. Thus, severe deformities are now unacceptable. An early percutaneous pin is much simpler than a late osteotomy. Colles' fractures deserve more respect than they received in the past.

Remember the basic classification into intra-articular and extra-articular fractures. Restoring reasonable joint surface alignment is an important part of the treatment and fortunately is usually relieved with simple traction under local or general anaesthesia.

SMITH'S FRACTURE OF LOWER END OF RADIUS

This is often referred to as a ‘reverse Colles' ’. It is caused by a fall on to the back of the hand. The lower fragment is flexed and impacted on the upper fragment. It is reduced and immobilised for 6 weeks in a cast as for Colles' fracture but with the wrist extended. Unstable fractures may require an above-elbow cast initially with the forearm in supination.

ULNA STYLOID FRACTURE

Treat symptomatically. Delayed union or non-union is common, but rarely symptomatic.

RADIAL STYLOID FRACTURE

Undisplaced:	plaster slab for 3 weeks
Displaced:	closed reduction and plaster slab for 6 weeks if this fails-open reduction

SCAPHOID FRACTURES

Scaphoid fractures account for almost 75% of all carpal injuries (Figure 135.16), but are rare in children and the elderly.[9] It is caused typically by a fall on to the outstretched wrist.



Figure 135.16 Typical appearance of a fractured scaphoid

If a scaphoid fracture is highly suspected in the presence of a normal X-ray of wrist, immobilise the wrist in a schaphoid plaster for 10 days, remove it and then re-X-ray. Isotopic bone scan may be indicated in cases where suspicion of fracture is high despite normal X-rays. For undisplaced fractures, 8 weeks in a below-elbow scaphoid cast usually suffices (Figure 135.17). Displaced fractures of the scaphoid require reduction (either open or closed) and, if unstable, internal fixation.



Figure 135.17 Appearance of the scaphoid plaster

All scaphoid fractures require late X-ray evaluation of treatment to diagnose non-union before they become symptomatic from late degenerative changes. Early bone grafting of a non-union can prevent fragment collapse and radioscaphoid degenerative changes.

Pitfall

The fracture may not be apparent on routine wrist X-rays. Specific scaphoid views should be requested.

SCAPHOLUNATE DISSOCIATION

This not uncommon carpal injury results from disruption of the scapholunate interosseous ligament and palmar radiocarpal ligaments. It results in a gap appearing between the scaphoid and lunate bones (the so-called ‘Terry-Thomas’ sign on plain AP X-rays of the wrist) and the scaphoid rotating into a vertical position on lateral X-rays. It is associated with pain in the wrist on dorsiflexion. Median nerve compression may occur after wrist or carpal dislocations.

Early diagnosis with referral simplifies treatment. This injury has only been recognised in recent times.

FRACTURES OF THE HANDS AND FINGERS

Thumb fractures

The thumb's special function renders injuries more difficult than other digits. Fractures well clear of the joints in the proximal and distal phalanges are treated in a similar way to other digits. However, intra-articular injuries are more common and internal fixation is more likely on the thumb than other digits.[10]

Bennett's fracture

This is a fracture-dislocation of the first carpo-metacarpal joint. The larger fragment of the first metacarpal dislocates proximally and laterally (Figure 135.18).



Figure 135.18 Method of reduction of a Bennett's fracture-dislocation of the first carpometacarpal joint

Treatment

Under anaesthesia the thumb is reduced using the forces indicated (Figure 135.18). A scaphoid plaster is applied with the thumb in the open grasp position. If anatomical reduction cannot be achieved by closed means, then open reduction and internal fixation is indicated. Percutaneous pinning with wires under X-ray control is also commonly used to hold an anatomical reduction.

Gamekeeper's thumb

This problematic injury of the metacarpo-phalangeal joint is presented in more detail on page 1380.

METACARPAL FRACTURES

Metacarpal fractures can be stable or unstable, intra-articular or extra-articular, and closed or open. They include the ‘knuckle’ injuries resulting from a punch which is prone to cause a fracture of the neck of the fifth metacarpal. As a general rule, most metacarpal (shaft and neck) fractures are treated by correcting marked displacements with manipulation (under anaesthesia) and splinting with a below-elbow, padded posterior plaster slab that extends up to the dorsum of the proximal phalanx and holds the metacarpophalangeal joints in a position of function (Figure 135.19).[4]



Figure 135.19 Fracture of the metacarpal: showing position of function with posterior plaster slab and the hand gripping a roll of felt padding

There is often a tendency for metacarpal fractures to rotate and this must be prevented. This is best achieved by splinting the MCP joints at 90°, which corrects any tendency to malrotation. If there is gross displacement, shortening or rotation then surgical intervention is indicated. A felt pad acts as a suitable grip. The patient should exercise free fingers vigorously. Remove the splint after 3 weeks and start active mobilisation.

PHALANGEAL FRACTURES

These fractures result from either direct trauma causing a transverse or a comminuted fracture or a torsional force causing an oblique fracture. The tendency to regard fractures of phalanges (especially middle and proximal phalanges) as minor injuries (with scant attention paid to management and particularly to follow-up care) is worth highlighting. These fractures require as near perfect reduction as possible, careful splintage and, above all, early mobilisation once the fracture is stable-usually in 2 - 3 weeks.

Nevertheless, overzealous mobilisation can be as dangerous as prolonged immobilisation. Early operative intervention should be considered if the fracture is unstable.

Angulation is usually obvious but it is most important to check for rotational mal-alignment, especially with torsional fracture. A simple method is to get the patient to make a fist of the hand and check the direction in which the nails are facing. Furthermore each finger can be flexed in turn and checked to see if the fingertips point towards the tubercule of the scaphoid (palpable halfway along the base of the thenar eminence and 1.5 cm distal to the distal wrist crease).

The phalanges

- *Distal phalanges*: usually crush fractures; generally heal simply unless intra-articular. Disturbance of nail growth is common.

- *Middle phalanges*: tend to be displaced and unstable-beware of rotation.
- *Proximal phalanges*: of the greatest concern, especially of the little finger; intra-articular fractures usually need internal fixation.

Treatment

Non-displaced phalanges with no rotational malalignment. Strap the injured finger to the adjacent normal finger with an elastic garter or adhesive tape for 2 - 3 weeks (i.e. 'buddy strapping') (Figure 135.20). Start the patient on active exercises.



Figure 135.20 Treatment of non-displaced phalanges by 'buddy strappping': the fractured finger is strapped to an adjacent healthy finger

or

If pain and swelling is a problem, splint the finger with a narrow dorsal or anterior slab (a felt-lined strip of malleable aluminium can be used) (Figure 135.21).



Figure 135.21 Method of splinting a phalangeal fracture of the index finger by a posterior plaster slab 4

An alternative is to bandage the hand while the patient holds a tennis ball or appropriate roll of bandage in order to maintain appropriate flexion of all interphalangeal joints.

Displaced phalangeal fractures (usually proximal and middle). With suitable anaesthesia correct the deformity by traction and direct digital pressure. Maintain correction by splintage for 2 - 3 weeks. Ensure flexion at the interphalangeal joints with a dorsal padded plaster slab from above the wrist to the base of the fingernail (Figure 135.21).

Intra-articular phalangeal fractures

Intra-articular phalangeal fractures are a great problem in management as subsequent stiffness of even a single interphalangeal joint can be a significant disability. Subsequent degenerative changes are common.

These fractures often occur in association with subluxation or dislocation of the joint. Reduction and fixation of the fracture may be an integral part of restoring joint stability. Displaced intra-articular phalangeal fractures, especially with joint instability, require referral.

Mallet finger

Refer to page 1378.

PENETRATING INJURIES TO THE HAND

Assessing these injuries requires a careful history and examination. The pugilist who sustains a seemingly minor cut over a ‘knuckle’ may have a tooth-penetrating injury to the metacarpophalangeal joint. In the flexed position the dorsal hood is drawn over the joint. The point of penetration of the hood retracts as the finger extends and ‘locks’ saliva into the joint. This injury invariably results in a severe septic arthritis unless aggressively treated with surgical debridement and high-dose antibiotics. Given the common occurrence of oral pathogens, antibiotic cover should include anaerobic organisms.

Refer to pp. 1378 - 9.

DISLOCATED FINGERS

In most cases the distal part dislocates dorsally.

For dislocated fingers immediate reduction is advisable. Test for an associated fracture and X-ray if appropriate. General anaesthesia may be necessary for reduction of a dislocated thumb.

Simple reduction of a dislocated interphalangeal joint

This method employs the principles of using the patient's body weight as the distracting force to achieve reduction of the dislocation. It is relatively painless and very effective.

Method

1. Face the patient, both in standing positions.
2. Firmly grasp the distal part of the dislocated finger. A better grip is achieved by wrapping simple adhesive tape around the end of the finger.
3. Request the patient to lean backward, while maintaining the finger in the fixed position (Figure 135.22).



Figure 135.22 Reduction of a dislocated finger

4. As the patient leans back, sudden, painless reduction should spontaneously occur.

Otherwise under a ring block or sedation apply traction and push the proximal phalangeal head dorsally. Splint the joint for 3 weeks to allow soft-tissue healing.

Pitfalls

- Instability-torn collateral ligaments: unstable in lateral direction
- Interposed volar plate-postreduction full flexion absent
- Fractures of base of phalanx
- Extensor mechanism rupture (e.g. buttonhole deformity at PIP joint or mallet finger deformity at DIP joint)

These problems may need surgical reduction.

FRACTURES OF THE PELVIS

Fractures of the pelvic ring are either:

1. stable: a single fracture
2. unstable: a break at two sites or association with disruption of the symphysis pubis or sacroiliac articulation

Treatment

Stable pelvic fracture:

- symptomatic, especially analgesics
- bed rest as pain symptoms dictate
- attempt walking with an aid as soon as comfortable

Unstable fractures: these are usually serious with possible associated visceral damage or blood loss. Patients should be referred for expert help.

FEMORAL FRACTURES

Femoral neck fractures include:

- subcapital fractures
- intertrochanteric fractures
- stress fractures in the young

Subcapital fractures are usually treated by pinning. Greatly displaced subcapital fractures in the elderly have a high risk of femoral head vascular necrosis. Thus, prosthetic replacement of the femoral head may be considered as a primary option.

A trap can be the impacted subcapital fracture that may allow partial weight-bearing, thus making radiological investigation essential in elderly patients complaining of hip pain. The fracture may not be evident on plain X-rays. If suspicion of fracture is still high, a bone scan should be performed.

Beware of the teenage athlete who complains of hip pain after running. Exclude a slipped upper femoral epiphysis and then a stress fracture. A technetium-99m bone scan will detect the fracture. A stress fracture may displace without warning, posing a serious risk of femoral head avascular necrosis. Thus, stress fractures must be considered for prophylactic pinning.

A summary of the management of other femoral fractures is presented with Figure 135.23.


Fracture
Usual management
subcapital fracture
reduction and pin or prosthesis (older patient)
stress fracture (usually young athlete)
internal fixation
intertrochanteric fracture
pin and plate
femoral shaft
intramedullary nail
supracondylar fracture
internal fixation
condyle:
undisplaced fracture: full limb plaster cylinder or internal fixation
displaced: internal fixation

Figure 135.23 Management of basic fractures of the femur

DISLOCATIONS OF THE LOWER LIMBS

Posterior dislocation of the hip

This causes a very painful shortened leg which is held adducted, medially rotated and slightly flexed. Be careful of sciatic nerve damage. Early reduction within hours minimises the risk of avascular necrosis of the femoral head.

Management

- Adequate analgesia (e.g. IM pethidine)
- X-rays to confirm diagnosis and exclude associated fracture
- Reduction of the dislocated hip under relaxant anaesthesia
- Follow-up X-ray to confirm reduction and exclude any fracture not visible on the first X-ray
- Intra-articular bone fragments need to be excluded by CT scanning

Note: Femoral neurovascular injury may occur in the rare cases of awterior dislocation of the hep.

Patella dislocation and subluxation

An acute dislocation needs to be reduced as a medical emergency.

The dislocated patella, which occurs mainly in children and young adults, especially girls, is always displaced laterally (Figure 135.24), often following a rotational and valgus force. The patient may feel the patella dislocate and it may sometimes reduce spontaneously. It may be associated with an osteochondral injury on the medial facet of the patella or the lateral femoral condyle. There is often a tense joint effusion, especially in the presence of a fracture. Predisposing factors include valgus knees, a small module patella, a laterally placed tibial tuberosity, a shallow patella-femoral groan and ligamentous laxity. Immediate reduction can be attempted by gently flexing the hip to relax the quadnceps, placing the thumb under the lateral edge of the patella and pushing it medially as the knee is extended. This may be attempted without anaesthesia or by using pethidine and intravenous diazepam as a relaxant.



Figure 135.24 Dislocated patella showing lateral displacement

X-rays with anteroposterior, lateral, skyline and intracondylar views should be taken to exclude an associated osteochondral fracture.

The usual RICE treatment should be given initially and crutches provided. Rest of the injured knee is achieved using a knee splint with the knee held in extension and crutches for 4 weeks.

Weight-bearing is permitted when the swelling has subsided and the patient is gradually taken off the crutches. Introduce quadriceps exercises with the knee in extension.

Patellar subluxation is when the patella is mobile and dues not actually dislocate, but results in episodic pain and feelings of instability. Physiotherapy and appropriate splinting for sporting activities is helpful before surgical stabilisation is considers.

Recurrent dislocation/subluxations in young females (14 - 18 years) requires surgery-combined tibial tubercle transfer with lateral release of the capsule. Immediate surgery in the acute phase is undertaken only in the presence of haemarthrosis with an osteochondral fracture.

FRACTURES OF THE PATELLA

- Fractures without displacement: walking plaster cylinder 4 weeks.
- Displaced single transverse fracture: surgical reduction with Kirschner wires.
- Displaced and comminuted fracture: refer for patellectomy.

FRACTURES OF BOTH TIBIA AND FIBULA

The nature and management of these fractures vary considerably. Some fractures are caused by blunt injuries, such as a blow from a motor car bumper, while twisting forces cause a spiral fracture of both bones at different levels. As a general rule, referral of patients to a specialist is necessary, especially where soft-tissue damage is significant. Management of fractures with minimal soft-tissue damage can be summarised thus:

- No or minimal displacement: full-length cast as for isolated fracture of tibia.
- Displacement: reduction under general anaesthesia, then application of cast as above (meticulous alignment essential).
- Period of immobilisation: adults 16 weeks, children 8 weeks.

FRACTURE OF FIBULA[9]

An isolated fracture of the fibula is usually due to stress or to a direct blow. The patient is generally able to stand and move the knee and ankle joints. However, most spiral fractures are associated with injuries of the ankle or knee. The ankle in particular should be examined and X-rayed.

Treatment is usually with analgesics to control the pain and no more than a crepe bandage or a walking stick is necessary. A below-knee walking plaster for about 3 weeks will help those with severe discomfort.

FRACTURE OF THE TIBIAL SHAFT

A fracture of the tibia alone is uncommon in adults but more common in children, due to a twisting injury. Reduction may not be necessary in some patients. Many can be reduced to a satisfactory position in the anaesthetised patient by letting the fractured leg hang over the edge of the table with the knee at a right angle.

A padded cast from the groin to the metatarsal necks is applied with the knee joint at 10° of flexion, and the ankle at a right angle. This should be maintained for 3 - 4 months.

FRACTURE AROUND THE ANKLE

The ankle is one of the areas liable to fractures. The commonest mechanism is forceful inversion of the foot, which can cause fracture of the fibula on a level with the joint line and tearing of the lateral collateral ligament. Other injuries can also occur, such as fracture of the medial malleolus and tearing of the tibiofibular syndesmosis. At least three views on X-ray are needed: AP, lateral and a half oblique 'mortise' view.

Undisplaced, uncomplicated fractures are treated with a plaster cast from just below the knee to the toes for 6 - 8 weeks. The foot must be plantigrade (i.e. with the foot at 90° to the leg and neither in varus nor in valgus).[9] Fractures treated in plaster need X-ray monitoring. Unsuspected displacement may occur as swelling subsides and the plaster loosens. Occult displacement of the fracture leading to mal-union will predispose to ankle osteo-arthritis. Fractures that are displaced or cause instability of the ankle joint require surgery to achieve stability followed by a longer period of immobilisation.

ANKLE/TALUS/SUBTALAR JOINT DISLOCATIONS

These dislokations may result in vascular compromise. The strctched overlying skins may rapidly necrose. Refer early.[7]

STRESS FRACTURES OF THE FOOT

Stress fractures of the navicular, calcaneus and metatarsal bones can be found in otherwise healthy people from the age of 7 onwards. Long-distance runners and high-performance athletes are also susceptible.

Clinical features

- Localised pain during weight-bearing activity
- Localised tenderness and swelling (not inevitable)
- Plain X-rays are necessary but show no fracture in about 50% of cases;[6] X-rays can be repeated in 2 - 3 weeks if a fracture is suspected
- A nuclear bone scan may confirm the diagnosis

Navicular

This hitherto unrecognised stress fracture has become apparent with the advent of CT scanning, which shows up the fracture better than nuclear scanning. It is seen in athletes involved with running sports and presents as poorly localised mid-foot pain. Plain X-ray is usually normal. The fracture, like the scaphoid fracture, is difficult to manage since delayed union and non-union are common. Cast immobilisation for 8 weeks may avoid the need for an operation.

Metatarsal bones

The second metatarsal is probably the most common site of all for stress fracture because it is invariably the largest metatarsal and absorbs a greater load than the others.[1]

Treatment

- Rest is the basis of treatment
- Resting the foot with crutches for 6 weeks provides optimal healing
- Healing usually takes 6 - 8 weeks

- Gradual slow resumption of activity

FRACTURES OF THE TOES

Most toe injuries are easy to treat but, like the fingers, the great and little toes demand special attention. Intra-articular injuries of the great toe (unless undisplaced) should be treated by internal fixation.

'Buddy strapping' can be used for many uncomplicated fractured phalanges of the toes, which tend to angulate and rotate more readily and are often harder to control than finger fractures. Strapping them to their adjacent toes on both sides simultaneously tends to counteract this problem.

Like the little finger, the little toe is injured by forceful abduction and if allowed to heal in that position may leave difficulties in wearing shoes.[9]

Approximate average immobilisation times for various fractures are given in Table 135.3.

Table 135.3 Healing of uncomplicated fractures (adults)

Fracture	(Approximate) average immobilisation time (weeks)
Rib	3 - 6 (healing time)
Clavicle	4 - 8 (2 weeks in sling)
Scapula	weeks to months
Humerus	
• neck	3 - 6
• shaft	8
• condyles	3 - 4
Radius	
• head of radius	3
• shaft	6
• Colles'	4 - 6
Radius and ulna (shafts)	6 - 12
Ulna-shaft	8
Scaphoid	8 - 12
Metacarpals	
• Bennett's fracture	6 - 8
• Other MCs	3 - 4
Phalanges (hand)	
• proximal	3
• middle	2 - 3
• distal	2 - 3
Pelvis	rest in bed 2 - 6
Femur	

• femoral neck	according to surgery
• shaft	12 - 16
• distal	8 - 12
Patella	3 - 4
Tibia	12 - 16
Fibula	0 - 6
Both tibia and fibula	16
Pott's fracture	6 - 8
Lateral malleolus avulsion	3
Calcaneus	
• minor	4 - 6
• compression	14 - 16
Talus	12
Tarsal bones (stress fracture)	8
Metatarsals	4
Phalanges (toes)	0 - 3

Important principles:

- Children under 8 years usually take half the time to heal.
- Have a check X-ray in 1 week (for most fractures).
- Radiological union lags behind clinical union.

Note: There can be considerable variation in immobilisation times depending on factors such as trauma degree and soft-tissue injuries.

DISLOCATION OF TOES

Dislocations occur mainly at the meta-tarsophalangeal joint and are rare; they require special care because of the strong tendons crossing the joint. Perfect reduction of the dislocated great toe is essential and it should be supported by a below-knee plaster cast extending beyond the toes. Temporary internal fixation with a Kirschner wire or open liga-mentous repair may be required.[10]

ANALGESIA AND RELAXATION

For the reduction of dislocations, analgesia and relaxation are appropriate. Resuscitation facilities and an experienced practitioner are required to handle this procedure. All drugs should be given intravenously and titrated to achieve the desired effect. Adverse effects include respiratory depression and hypo-tension.[7] The choice of agents is presented in Table 135.4.

Table 135.4 Analgesic and relaxant/sedative agents

	IV dose	**Antidote**
Relaxant/sedative agent		

Diazepam	0.1 - 0.2 mg/kg (5 - 10 mg)	Flumazenil
Midazolam	0.05 - 0.1 mg/kg (2 - 5 mg)	Flumazenil
Analgesic agent		
Fentanyl	1 - 2 μg/kg (50 - 100 μg)	Naloxone
Morphine	0.1 - 0.2 mg/kg (5 - 15 mg)	Naloxone

PLASTERING TIPS

Plaster of Paris

The bucket of water

- Line the bucket with a plastic bag for easy cleaning.
- The water should be deep enough to allow complete vertical immersion.
- Use cold water for slow setting.
- Use tepid water for faster setting.
- Do not use hot water: it produces rapid setting and brittle plaster.

The plaster rolls

- Do not use plaster rolls if water has been splashed on them.
- Hold the roll loosely but with the free end firm and secure (Figure 135.25).



Figure 135.25 Holding the plaster roll

- Ensure that the centre of the plaster is fully wet.
- Drain surface water after removal from the bucket.
- Gently squeeze the roll in the middle: do not indent.

Padding

- Use Velband or stockinet under the plaster.
- With Velband, moisten the end of the roll in water to allow it to adhere to the limb.
- For legs, make extra padding around the ankle and heel.
- Avoid multiple layers of padding.

Method

- Use an assistant to support the limb where possible (e.g. hold the arm up with fingers of stockinet).
- Lay the bandage on firmly but do not pull tight.
- Lay it on quickly.
- Overlap the bandage by about 25% of its width.

Note: It is good practice to review the patient the next day.

FURTHER READING

Apley AG, Solomon L. *Apley's System of Orthopaedics and Fractures* (9th edn). Oxford: Butterworth-Heinemann, 1993.

REFERENCES

1. Quirk R. Stress fractures. Aust Fam Physician, 1993; 22: 300 - 7.
2. Brentnall E. Diagnosing a fracture. Aust Fam Physician, 1990; 19: 948.
3. McMenimen PJ. Management of common fractures of the upper limb. Aust Fam Physician, 1987; 16: 783 - 91.
4. Cook J, Sankaran B, Wasunna A. *Surgery at the District Hospital: Obstetrics, Gynaecology, Orthopaedics and Traumatology*. Geneva: World Health Organization, 199: 75 - 162.
5. Brentnall E. Spatula test for fracture of mandible. Aust Fam Physician, 1992; 21: 1007.
6. Peterson L, Renström P. *Sports Injuries: their Prevention and Treatment*. Sydney: Methuen, 1986: 179 - 81.
7. Mohammed KD, Sonnabend DH. A GP's guide to the reduction of dislocations. Mod Med Aust, 1996; 39(2): 100 - 8.
8. Young D, Murtagh J. Pitfalls in orthopaedics. Aust Fam Physician, 1989; 18: 645 - 60.
9. Apley AG, Solomon L. *Apley's System of Orthopaedics and Fractures (9th edn)*. Oxford: Butterworth-Heinemann, 1993: 601 - 4.
10. Carter G. Fractures and dislocations of fingers and toes. Aust Fam Physician, 1993; 22: 310 - 17.

CHAPTER 136. COMMON SPORTING INJURIES

Exercise and temperance can preserve something of our early strength even in old age.

CICERO (106 - 43 BC)

COMMON SPORTING INJURIES

Although there is considerable overlap between injuries occurring during everyday activities and those of sporting and recreational activities, there are many injuries that are characteristic to sportspeople. Many of these injuries are the result of trauma of various degrees and include the many varieties of fractures, dislocations and soft-tissue injuries.

On the other hand ‘runner's anaemia’ - an iron deficiency - is considered to be multifactorial, including mild haemolysis and blood loss from the bladder, kidney and gastrointestinal tract (refer p. 829).

INJURIES TO THE EYE

Blunt injuries to the eye are common in sport. Examples include tennis and squash balls, cricket balls and baseballs, and fists and fingers associated with body contact sports. Haemorrhage is the most common problem and occurs throughout the eye: subconjunctivally, in the anterior chamber (hyphaema), into the vitreous, and underneath the retina or choroid.

Another common problem is a corneal abrasion where a small wound can be caused by a foreign body, a fingernail or a contact lens. It needs to be treated with great respect.

Hyphaema

With hyphaema, bleeding from the iris collects in the anterior chamber of the eye (Figure 136.1). The danger is that, with exertion, a secondary bleed from the ruptured vessel could fill the anterior chamber with blood, blocking the escape of aqueous humour and causing a severe secondary glaucoma. Loss of the eye can occur with a severe haemorrhage. It is likely to happen between the second and fourth day after the injury.



Figure 136.1 Hyphaema of the eye showing blood in the anterior chamber. This occurred in a 29-year-old man who was struck in the eye by a squash ball

Management

- First, exclude a penetrating injury.
- Avoid unnecessary movement: vibration will aggravate bleeding. (For this reason, do not use a helicopter if evacuation is necessary.)

- Avoid smoking and alcohol.
- Do not give aspirin (can induce bleeding).
- Prescribe complete bed rest for 5 days and
- review the patient daily.
- Apply padding over the injured eye for 4 days.
- Administer sedatives as required.
- Beware of ‘floaters’, ‘flashes’ and field defects.

Arrange ophthalmic consultation after 1 month to exclude glaucoma and retinal detachment. No sport before this time.

Generally, recovery runs an uneventful course. If secondary bleeding occurs (usually the second, third or fourth day) the patient should be transported immediately to the nearest eye hospital. Evacuate by air (not by helicopter) only if the cabin altitude can be kept below 1300 metres (4000 feet). It is important to prevent vomiting and expansion of air within the eye.

Protective spectacles should always be worn when playing squash. People with monocular vision should be advised not to participate in this sport.

KNOCKED OUT OR BROKEN TEETH

If a permanent (second) tooth is knocked out it can be saved by immediate proper care. Likewise, a broken tooth should be saved and urgent dental attention sought.

The knocked-out tooth

- Place the tooth in its original position, preferably immediately (Figure 136.2): if dirty, put it in milk before replacement or, better still, place it under the tongue and ‘wash it’ in saliva. Do not use water, and do not wipe or touch the root.



Figure 136.2 Replacement of a knocked-out tooth

- Fix the tooth by moulding strong silver foil (e.g. cooking foil) over it and the adjacent teeth.
- Refer the patient to his or her dentist or dental hospital as soon as possible.

Note: Teeth replaced within half an hour have a 90% chance of successful reimplantation.

INJURIES TO THE NOSE

Common injuries to the nose include epistaxis and fractures of the nasal bones.

Epistaxis

First aid is simple tamponade, which is invariably effective. The soft cartilaginous part of the nose should be pinched between the finger and thumb for 5-10 minutes. The head should be kept bent slightly forward. Packing of the nose may be required (pp. 636 - 7).

Fracture of the nose

If deformity is present the patient should be referred for reduction within 7 days. Refer to page 1391.

Septal haematoma

Special care has to be taken of a septal haematoma, which has a tendency to become infected (p. 1390).

SHOULDER INJURIES

Common shoulder injuries acquired in sporting activities include:

- dislocated or subluxed acromioclavicular joint (p. 1393)
- fractured clavicle (p. 672)
- dislocated shoulder (pp. 682 - 4)
- supraspinatus tendonitis (p. 673)

Swimmer's shoulder

Painful shoulders occur in about 60% of elite level swimmers during their career. The basic disorder is rotator cuff tendonitis, particularly supraspinatus tendonitis, which is considered to be associated with abnormals apular positioning and cervicothoracic dysfunction. The best treatment is prevention, which aims at rotator cuff strengthening exercises, better scapulothoracic control, including correction of thoracic extension if it is decreased, and scapular stabilisation exercises.[1].

ELBOW INJURIES

Softtissue disorders of the elbow are extremely common. Two types of tennis elbow are identifiable. ‘Backhand’ tennis elbow or lateral epicondylitis (p. 1396) and ‘forehand’ tennis elbow or medial epicondylitis, which is also known as golfer's elbow or baseball pitcher's elbow. These common problems, often unrelated to sporting activity, are presented in more detail in Chapter 64.

HAND INJURIES

Hand and finger injuries are very important in sporting activities and include fractures and dislocations of phalanges and metacarpals. A mallet finger is a common injury and can result from overuse.

Ligamentous disruption of finger joints can cause instability and require early referral. An example is gamekeeper's thumb, often encountered in skiers, where there is complete tearing of the medial ligament of the metacarpophalangeal joint.

Mallet finger

A mallet finger is a common sports injury caused by the ball (football, cricket ball or baseball) unexpectedly hitting the finger tip and forcing the finger to flex. Such a forced hyperflexion injury to the distal phalanx can rupture or avulse the extensor insertion into its dorsal base. The characteristic swan neck deformity (Figure 136.3) is due to retraction of the lateral bands and hyperextension of the proximal interphalangeal joint.



Figure 136.3 Mallet finger with the 'swan-neck deformity' following rupture of the extensor tendon to the distal phalanx

The 45° guideline

Without treatment, the eventual disability will be minimal if the extensor lag at the distal joint is less than 45°; a greater lag will result in functional difficulty and cosmetic deformity.

Treatment

Maintain hyperextension of the distal interphalangeal joint for 6 weeks, leaving the proximal interphalangeal joint free to flex.

Equipment

- Friar's balsam (will permit greater adhesion of tape)
- Non-stretch adhesive tape, 1 cm wide: two strips approximately 10 cm in length

Method

1. Paint finger with Friar's balsam (compound benzoin tincture).
2. Apply the first strip of tape in a figure of eight configuration. The centre of the tape must engage and support the pulp of the finger. The tapes must cross dorsally at the level of the distal interphalangeal joint and extend to the volar aspect of the proximal interphalangeal joint without inhibiting its movement (Figure 136.4a).



Figure 136.4 Mallet finger: (a) application of first tape; b) application of 'stay' tape

3. Apply the second piece of tape as a 'stay' around the midshaft of the middle phalanx (Figure 136.4b).

Reapply the tape wherever extension of the distal interphalangeal joint drops below the neutral position (usually daily, depending on the patient's occupation). Maintain extension for 6 weeks.

Surgery

Open reduction and internal fixation are reserved for those cases where the avulsed bony fragment is large enough to cause instability, leading to volar subluxation of the distal interphalangeal joint.

Tenpin bowler's thumb

Tenpin bowler's thumb is a common stress syndrome in players. It usually presents as a soft-tissue swelling at the base of the thumb web, with associated pain and stiffness of the digits used for bowling. It may cause a traumatic neuroma of the digital nerve at this site with associated hyperaesthesia.

Management

- Rest
- Massage
- Bevel the bowling ball holes to reduce friction
- An intralesional injection (Figure 136.5) of 0.25 mL of long-acting corticosteroid mixed with local anaesthetic (resistant cases)



Figure 136.5 Tenpin bowler's thumb

SNOW SKIING INJURIES

The most common injuries encountered in snow skiing are soft-tissue injuries and fractures and dislocations.

A study by Robinson showed that the six most common skiing injuries were strains to the medial collateral ligament of the knee 24.3%; contusions of soft tissue (excluding head and neck) 17.6%; lacerations 15.5%; neck and back injuries 7.8%; fractures 7.6%; and dislocations.[2]

There has been a large decrease in injuries relative to participation in the past decade because of improved equipment and attention to safety. The most common fractures in skiers are those involving the tibia and fibula, especially spiral fractures. Other common fractures are of the clavicle, wrist and humerus. Dislocation of the shoulder region (glenohumeral joint and acromioclavicular joint) are due to falls on hard impacted snow.

Ulnar collateral ligament injury ('skier's thumb')[3, 4]

A special injury is skier's thumb (also known as gamekeeper's thumb) in which there is ligamentous disruption of the metacarpophalangeal joint with or without an avulsion fracture of the base of the proximal phalanx at the point of ligamentous attachment. This injury is caused by the thumb being forced into abduction and hyperextension by the ski pole as the skier pitches into the snow.

Diagnosis is made by X-ray with stress views of the thumb. Incomplete tears are immobilised in a scaphoid type of plaster for 3 weeks, while complete tears (the Stener lesion) and avulsion fractures should be referred for surgical repair.

SPINAL PROBLEMS

Spinal dysfunction, particularly of the neck and low back, are very common problems in sport, as for the general population.

Serious problems include pars interarticularis fractures, spondylolisthesis, disc disruptions with prolapse and, rarely, vertebral body fracture. The common problems are the various facet joint syndromes and musculoskeletal strains, which are managed conservatively as outlined in Chapters 37 and 62. The key to management is a conservative approach with a back education and exercise program.

INJURIES TO THE LOWER LIMBS

Injuries due to trauma and overuse of the lower limbs comprise the most frequent group of sports-related disorders requiring medical attention.

The three main causes of overuse trauma are:

- friction (e.g. peritendonitis)
- stress or overload (e.g. hamstring tear, tibialstress fracture)
- ischaemia (e.g. anterior compartment syndrome)

Overuse leg syndromes

Increased community participation in physical activity, including running and jogging, has resulted in a concomitant increase in overuse leg injuries, especially in the lower leg with its weight-bearing load. The common cause is repetitive trauma where the forces involved overwhelm the tissue's ability to repair adequately. Common causes of chronic leg pain include hamstring injuries and injuries to the lower leg.

Principles of management

Prevention:

- maintain ideal weight
- good nutrition
- adequate preparation
- warm-up exercises for the legs
- proper footwear
- proper activity planning

Treatment of injury

- Rest, or relative rest: the patient is allowed to perform activities that do not aggravate the injury.
- Ice: apply an ice pack for 20-30 minutes every 2 hours while awake during the first 48-72 hours post injury.
- Compression: keep the injured muscle or tissue firmly bandaged for at least 48 hours.
- Elevation: rest the leg on a stool or chair until the swelling subsides.
- Correction of predisposing factors (intrinsic or extrinsic) - orthotics for malalignment, correction of training errors.
- NSAIDs for painful inflammatory response.
- Physical therapy (e.g. stretching, mobilisation when acute phase settled).

GROIN PAIN

Groin pain is a particularly common condition among athletes.

Acute groin pain

Acute conditions such as muscle and musculotendinous strains,[5] and overuse injuries such as tendonitis and tendoperiostitis, are generally readily diagnosed and treated.[6] Diagnostic difficulties can arise because of referred pain from the lumbosacral spine, hip and pelvis, including hip labral injuries and a stress fracture of the femoral neck. More common acute groin injuries include injuries to the following muscles and their tendons (Figure 136.6):



Figure 136.6 Muscles in the groin region subject to musculotendinous injuries in the athlete

- adductor longus (e.g. musculotendinous strains)
- rectus femoris
- sartorius
- iliopsoas

Adductor muscle or tendon injury

Features: upper inner thigh pain, local tenderness, resisted hip adduction and a history of twisting, catching or slipping. Initial management is with RICE therapy followed by an adductor strengthening program. Persistent adductor pain will require investigation with ultrasound or MRI to confirm the diagnosis and management (possible steroid injection).

Iliopsoas problems

The typical feature is pain on stretching the hip flexor or with resisted hip flexion. Bursitis may be present. Treatment includes avoidance of aggravating activity and stretching, followed by a graduated re-strengthening program.

Other injuries include:

- SCFE in adolescents
- avulsion fractures in adolescents (e.g. rectus femoris and sartorius on the iliac spines)

Chronic groin pain

There are many causes of chronic groin pain, with bone and joint abnormalities being more likely causes. Important causes include:

- muscle and musculotendinous lesions (e.g. adductor longus tendoperiostitis)
- bursitis (e.g. iliopsoas bursitis)
- osteitis pubis (pubic symphysis)
- stress fractures (e.g. femoral neck and pubic rami)
- sacroiliac and hip joint disorders (e.g. osteoarthritis hip/tumour)
- lumbar spine - L1-2 or L2-3 disc

- ‘occult’ inguinal or femoral hernia

Investigations

- X-ray of pelvis (AP, lateral, oblique)
- Tomography of pubic symphysis (to detect osteitis pubis and pubic instability)
- Bone scan to detect stress fractures or osteitis pubis
- Herniography
- CT scan or MRI or ultrasound (increasing potential)

Osteitis pubis

This is inflammation on each side of the periosteal bone of the symphysis pubis that usually develops insidiously in sportspeople and is associated with pubic bone stress. It often presents in a football player or other athlete involved in sports that involve sudden twisting or turning movements.[5] It may follow injuries of the adductor longus muscle origin.

Clinical features

- Pain in groin (i.e. low anterior pelvis)
- Radiation of pain into adductor muscles of both thighs
- Tendency to ‘duck waddling’ gait
- Pain aggravated by exercise, especially twisting and turning and kicking
- Local ‘pinpoint’ tenderness to palpation over the symphysis
- Positive squeeze test:[5] the patient lies supine with the hips flexed to 45°, the knee flexed to 90° and the feet on the bed. The examiner places their fist between the medial aspects of the patient's knees and the patient is asked to adduct the hips bilaterally against the fist
- MRI scanning is the most sensitive imaging technique[5]

Management

- NSAIDs
- Relative rest from provoking activity
- Conservative rehabilitation program
- Activity modification
- Change to a low impact sport for 3-6 months (e.g. cycling)
- Regular stretching - physiotherapy supervision
- Graduated strengthening program of the adductors and lower abdominal musculature over 10-12 weeks
- Surgery has a place for specific cases

Jock itch

Jock itch, or tinea cruris, is a common infection in the groin area of young men, especially athletes, who are subjected to chaffing in the groins from tight shorts and nylon ‘jock straps’. The feet should be inspected for evidence of tinea pedis. The dermatophyte is transmitted by towels and other objects, particularly in change rooms and communal showers (p. 1154).

HAMSTRING INJURIES

Hamstring strains are common in athletes. The short head of biceps femoris is the most commonly strained component of the hamstring group.

Clinical features:

- a history of a ‘pull’, ‘twinge’, ‘tear’ or ‘twang’ in the back of the thigh

- a soreness and lump develops (with a severe tear a person can collapse)
- localised tenderness
- limitation of straight leg raising
- pain on resisted or active knee flexion or hip extension
- bruising (usually in popliteal fossa) may be present

Management

The immediate goals of treatment of the acute injury are to relieve pain and minimise swelling.

- RICE for 72 hours
- NSAIDs (e.g. aspirin or indomethacin)
- stretching exercises
    - passive stretching after ice treatment
    - then active stretching
    - then isometric contraction exercises

HAEMATOMAS IN MUSCLE (‘CORKED THIGH’)

Haematomas can be intramuscular, intermuscular or interstitial. They usually result from a sharp blow (e.g. knee to the thigh or kick in the anterior compartment of the leg).

An intramuscular haematoma can cause an acute compartment syndrome which may require urgent decompression. One objective of treatment is to prevent excessive scarring. Other complications include infection, cyst formation, thrombophlebitis and myositis ossificans.

Management

- RICE treatment with emphasis on cooling
- Non-weight-bearing, using crutches initially
- Consider admission to hospital or a day surgical unit to check progress
- Referral for expert advice may be appropriate because of the potentially
- serious nature of the injury

INJURIES TO THE KNEE

Knee injuries are common, have multiple clinical disorders and are potentially disastrous to the athlete. The various injuries and overuse syndromes are presented in considerable detail in Chapter 67, on the painful knee. Any twisting or pivoting knee injury resulting in pain or swelling should be considered an ACL disruption until proved otherwise. Often the person is involved in a ball sport and describes a loud ‘pop’ or ‘crack’ at the time of injury.

Acute injuries

Acute injuries (pp. 739 - 41) include:

- meniscal tears
- ligamentous tears and strains (of varying degrees)
    - anterior cruciate ligament (ACL)
    - posterior cruciate ligament (PCL)
    - medial collateral ligament
    - lateral collateral ligament

Overuse syndromes

The knee is very prone to overuse disorders (p. 739). The pain develops gradually without swelling, is aggravated by activity and relieved with rest. It can usually be traced back to a change in the sportsperson's training schedule, footwear, technique or related factors. It may also be related to biomechanical abnormalities, ranging from hip disorders to disorders of the feet.

Overuse injuries include:

- patellofemoral pain syndrome (jogger's knee/runner's knee)
- patellar tendonitis (jumper's knee)
- synovial plica syndrome
- infrapatellar fat-pad inflammation
- anserinus bursitis/tendonitis
- biceps femoris tendonitis
- semimembranous bursitis/tendonitis
- quadriceps tendonitis/rupture
- popliteus tendonitis
- iliotibial band friction syndrome (runner's knee)
- the hamstrung knee

A careful history followed by systematic anatomical palpation around the knee joint will pinpoint the specific overuse syndrome.

OVERUSE INJURIES TO LOWER LEG

A summary of the clinical and management aspects of various injuries is presented in Table 136.1 and in Figure 136.7.

Table 136.1 Clinical comparisons of overuse syndromes in lower leg

Syndrome	Symptoms	Cause	Treatment
Anterior compartment syndrome	Pain in the anterolateral muscular compartment of the leg, increasing with activity. Difficult dorsiflexion of foot, which may feel floppy.	Persistent fast running (e.g. squash, football, middle-distance running).	Modify activities. Surgical fasciotomy is the only effective treatment.
Iliotibial band tendonitis	Deep aching along lateral aspect of knee or lateral thigh. Worse running downhill, eased by rest. Pain appears after 3-4 km running.	Running up hills by longdistance runners and increasing distance too quickly.	Rest from running for 6 weeks. Special stretching exercises. Correct training faults and footwear. ? injection of LA and corticosteroids deep into tender areas.
Tibial stress syndrome or shin splints	Pain and localised tenderness over the distal posteromedial border of the tibia. Bone scan for diagnosis.	Running or jumping on hard surfaces.	Relative rest for 6 weeks. Ice massage. Calf (soleus stretching). NSAIDs. Correct training faults and footwear.
Tibial stress fracture	Pain, in a similar site to shin splints, noted after running. Usually relieved by rest. Bone scan for diagnosis.	Overtraining on hard (often bitumen) surfaces. Faulty footwear.	Rest for 6-10 weeks. Casting not recommended. Graduated training after healing.
Tibialis anterior tenosynovitis	Pain, over anterior distal third of leg and ankle. Pain at beginning and after exercise ± swelling, crepitus. Pain on active or resisted ankle dorsiflexion.	Overuse - excessive downhill running.	Rest, even from walking. Injection of LA and corticosteroid within tendon sheath.

Achilles tendonitis	Pain in the Achilles tendon aggravated by walking on the toes. Stiff and sore in the morning after rising but improving after activity.	Repeated toe running in sprinters or running uphill in distance runners.	Relative rest. Ice at first and then heat. 10 mm heel wedge. Correct training faults and footwear. NSAIDs.


iliotibial band tendonitis
anterior compartment syndrome
Achilles tendonitis
plantar fasciitis
shin splint
tibial stress fracture

Figure 136.7 Common sites of overuse injuries in the lower leg

Common causes of chronic lower leg pain in sportspeople include:[7]

- medial tibial stress syndrome (previously called shin splints)
- stress fractures (Figure 136.8)



Figure 136.8 Typical sites of stress fractures in athletes in the tibia and fibula

- exertional compartment syndrome, especially anterior compartment
- tibialis anterior tenosynovitis (Figure 136.9)



Figure 136.9 Site of tibialis anterior tenosynovitis

- chronic muscle strains

These problems are invariably due to excessive physical demands in athletes striving for the ultimate performance or in the occasional athletes who have made inadequate preparation for their activity. Training errors contribute to a large proportion (60%) of overuse injuries.[7]

Principles of treatment

- Rest or relative rest
- Exercise program (where appropriate)
- Correction of predisposing factors
  - training errors
  - unsuitable footwear
  - inadequate warm-up
  - malalignment
- Analgesics: use NSAIDs only if it is true inflammatory pain (pain at rest)

STRESS FRACTURES

Stress fractures are an important cause of lower leg pain and foot pain in sport, accounting for 5-15% of injuries.[7] Stress fractures occur in the tibia and fibula and in the foot (navicular, calcaneus and metatarsals). The important clinical factor is to keep stress fractures in mind and X-ray the tender area. If the X-ray is negative and there is a high index of suspicion, a radionuclide scan should be ordered.

In the tibia, stress fractures occur mainly in the proximal metaphysis and the junction of the middle and distal thirds of the shaft (Figure 136.10). In the fibula they usually occur 5-7 cm above the tip of the lateral malleolus (Figure 136.8).



Figure 136.10 Site of pain in tibial stress syndrome

These stress fractures usually occur after prolonged and repeated heavy loading such as long-distance running or repeated jumping.

TORN 'MONKEY MUSCLE'

The so-called torn 'monkey muscle', or 'tennis leg', is actually a rupture of the medial head of gastrocnemius at the musculoskeletal junction where the Achilles tendon merges with the muscle (Figure 136.11).



Figure 136.11 ‘Tennis leg’ or ‘monkey muscle’ - illustrating typical site of rupture of the medial head of gastrocnemius at the junction of muscle and tendon left leg)

It is not a torn plantaris muscle as commonly believed. This painful injury is common in middleaged tennis and squash players who play infrequently and are unfit.

Clinical features:

- a sudden sharp pain in the calf (the person thinks he/she has been struck from behind, e.g. a thrown stone)
- unable to put heel to ground
- walks on tiptoes
- localised tenderness and hardness
- dorsiflexion of ankle painful
- bruising over site of rupture

Management

- RICE treatment for 48 hours
- Ice packs immediately for 20 minutes and then every 2 hours when awake (can be placed over the bandage)
- A firm elastic bandage from toes to below the knee
- Crutches can be used if severe
- A raised heel on the shoe aids mobility. Women should wear high heels
- Commence mobilisation after 48 hours rest, with active exercises
- Physiotherapist supervision for gentle stretching massage and then restricted exercise

SPRAINED ANKLE

There are two main ligaments that are subject to heavy inversion or eversion stresses, namely the lateral ligaments and the medial ligaments respectively. Most of the ankle ‘sprains’ or tears involve the lateral ligaments (up to 90%), while the stronger tauter (deltoid) ligament is less prone to injury.

The lateral ligament complex involves three main bands: the anterior talofibular (ATFL), the calcaneo-fibular (CFL) and the posterior talofibular ligament (PTFL) (Figure 136.13).



Figure 136.13 Lateral ligaments of the ankle

Mechanism of injury to lateral ligaments [8]

Forced inversion causes about 90% of all ankle injuries.

Most sprains occur when the ankle is plantar-flexed and inverted, such as when landing awkwardly after jumping or stepping on uneven ground.

Inversion

- Foot in plantarflexion: ATFLinjury likely 50-60%)
- Foot in neutral: CFLinjury likely (10%)
- Foot in dorsiflexion: PTFLinjury likely (5%)

Note: Combined ATFLand CFLinjury (15-25%).

Eversion

- Foot in plantarflexion or neutral: medial ligament (mainly anterior part)

The classification of ankle injuries is presented in Table 136.2.

Table 136.2 Classification of injuries to ankle ligaments (adapted from Litt)[8]

Grade	Functional/clinical	Ligamentous stability	Stress X-rays
I (mild)	minimal pain and swelling minimal bleeding full range of motion heel and toe walking	minor ligamentous injury with only a partial tear of the ligament stable ankle joint	normal
II (moderate)	moderate to severe pain and swelling	similar to grade I only more severe	anterior draw 4 - 14 mm

	considerable bleeding decreased range of motion difficulty in weight-bearing and ambulation	partially unstable joint	talar tilt 5 - 10°
III (severe)	minimal to severe pain and swelling pronounced bleeding minimal range of motion unable to weight-bear	complete ligamentous rupture with unstable joint	anterior draw > 15 mm talar tilt > 20°

Clinical features of sprained lateral ligaments

Common features

- Ankle 'gives way'
- Difficulty in weight-bearing
- Discomfort varies from mild to severe
- Bruising (may take 12-24 hours) indicates more severe injury (Figure 136.12)



Figure 136.12 Sprained ankle with tearing of the lateral ligament complex with obvious haematoma

- May have functional instability: ankle gives way on uneven ground

Physical examination (perform as soon as possible)

- Note swelling and bruising
- Palpate over bony landmarks and three lateral ligaments
- Test general joint laxity and range of motion
- A common finding is rounded swelling in front of lateral malleolus (the 'signe de la coquille d'oeuf ')
- Test stability in AP plane (anterior draw sign)
- Talar tilt test (inversion stress test)

Is there an underlying fracture?[9]

For a severe injury the possibility of a fracture - usually of the lateral malleolus or base of fifth metatarsal - must be considered. If the patient is able to walk without much discomfort straight after the injury, a fracture is unlikely.

Indications for X-ray include:[9]

- inability to weight-bear immediately after injury
- marked swelling and bruising soon after injury
- marked tenderness over the bony landmarks
- marked pain on movement of the ankle
- crepitus on palpation or movement
- point tenderness over the base of the fifth metatarsal
- special circumstances (e.g. litigation potential)

Ottawa rules for ankle and foot X-ray[10]

These rules are a quick and reliable method of selecting which patients with ankle and foot injuries need X-rays to exclude a fracture.

Ankle injury

An X-ray of the ankle is necessary when the patient has pain over the medial or lateral malleolar zone and any one of the following findings:

- there is bone tenderness on palpation of the distal 6 cm of the fibula (posterior tip of lateral malleolus)
- there is bone tenderness on palpation of the distal 6 cm of the tibia (posterior tip of medial malleolus)
- there was an inability to bear weight (walk four steps) *both* immediately after injury and during the clinical examination

Foot injury

Refer for a foot X-ray (suspected mid-foot fracture) if there is pain in the mid-foot and any one of:

- bone tenderness at fifth metatarsal base
- bone tenderness at the navicular bone
- inability to weight bear immediately after injury and when seen

Management

The treatment of ankle ligament sprains depends on the severity of the sprain. Most grade I and II sprains respond well to standard conservative measures and regain full, pain-free movement in 1-6 weeks, but controversy surrounds the most appropriate management of grade III sprains.

Grade I sprain

R rest the injured part for 48 hours, depending on disability

I ice pack for 20 minutes every 3-4 hours when awake for the first 48 hours

C compression bandage (e.g. crepe bandage)

E elevate to hip level to minimise swelling

A analgesics (e.g. paracetamol ± codeine)

R review in 48 hours, then 7 days

S special strapping

Use partial weight-bearing with crutches for the first 48 hours or until standing is no longer painful, then encourage early full weightbearing and a full range of movement with isometric exercises.[7] Use warm soaks, dispense with ice packs after 48 hours. Walking in sand (e.g. along the beach) is excellent rehabilitation. Aim towards full activity by 2 weeks.

Special strapping

A firm support for partial tears in the absence of gross swelling provides excellent symptomatic relief and early mobilisation.

Method

- Maintain the foot in a neutral position (right angles to leg) by getting patient to hold the foot in that position by a long strap or sling.
- Apply small protective pads over pressure points.
- Apply one or two stirrups of adhesive lowstretch 6-8 cm strapping from halfway up medial side, around the heel and then halfway up the lateral side to hold foot in slight eversion (Figure 136.14).



Figure 136.14 Supportive strapping for a sprained ankle: (a) apply protective pads and stay tape; (b) apply stirrups to hold foot in slight eversion and (c) apply an ankle lock tape

- Apply an adhesive bandage (e.g. Acrylastic, 6-8 cm) which can be rerolled and reused.
- Reapply in 3-4 days.
- After 7 days, remove the bandage and use a non-adhesive tubular elasticised support until full pain-free movement is achieved.

Grade II sprain

RICE treatment (as above) for 48 hours but ice (e.g. ACE wrap), should be used every 2-3 hours and no weightbearing (use crutches) for 48 hours. Then permit partial weightbearing with crutches and begin the active exercise program. Follow-up and supportive strapping as for grade I. Note that the ice packs can be placed over the strapping.

Grade III sprain

It would be appropriate to refer this patient with a complete tear. Initial management includes RICE and analgesics and an X-ray to exclude an associated fracture. The three main treatment approaches appear to be equally satisfactory.

Surgical repair

Some specialists prefer this treatment but it is usually reserved for the competitive athlete who demands absolute stability of the ankle.

Plaster immobilisation

This is usually reserved for patients who are unable actively to dorsiflex their foot to a right angle and those who need to be mobile and protected in order to work. The plaster is maintained until the ligament repairs, usually 4-6 weeks. The patient can walk normally when comfortable with a rockered sole or open cast walking shoe.

Strapping and physiotherapy

This approach is generally recommended. After the usual treatment for a grade II repair, including the strapping as described, a heel lock (Figure 136.14c) should be used.

The patient continues on crutches and appropriate physiotherapy is given with care so that the torn ends are not distracted. Strengthened balance is achieved by the use of elastic bands, swimming and cycling.

Non-response to treatment

There are some patients who, despite an apparently straightforward ankle sprain, do not respond to therapy and do not regain a full range of movement. In such patients alternative diagnoses in addition to ligament tearing must be considered (Table 136.3).

Table 136.3 Unstable ankle injuries to be considered in delayed healing (after Brukner)[9]

Osteochondral fracture of the talar dome
Dislocation of the peroneal tendons
Sinus tarsi syndrome
Anteroinferior tibiofibular ligament (syndesmosis) injury
Post-traumatic synovitis
Anterior impingement syndrome
Posterior impingement syndrome
Anterior lateral impingement
Rupture of posterior tibial tendon
Regional pain syndrome
Other fractures
• base fifth metatarsal (avulsion)
• lateral process of talus
• anterior process of the calcaneus
• tibial plafond
• stress fracture navicular

These require careful clinical assessment and further investigation such as bone scans.

Wobbleboard ‘aeroplane’ technique for ankle dysfunction

This involves proprioception exercises for injured ankle ligaments. The ligaments and leg muscles can be strengthened by the use of a wobble board. An improvised wobble board can be constructed by attaching a small piece of wood (10 cm × 5 cm) to the centre of a 30 cm piece of plywood or simply placing a slab of wood on a dome-shaped mound of earth. The patient stands on the board and shifts his or her weight from side to side in neutral (first), forward (after 2-3 days) or extended (later) body positions (Figure 136.15) to improve proprioception and balance.



Figure 136.15 Wobble board technique for ankle dysfunction

TIBIOFIBULAR SYNDESMOSIS INJURY[5]

The syndesmosis of the ankle joint comprises the anterior and posterior tibiofibular ligaments and the interosseous membrane. It is injured commonly in football codes by a dorsiflexion-eversion mechanism. The injury is not commonly recognised and can present as an ankle sprain that is slow to recover. Tenderness is found anteriorly over the ligament and pain (which can radiate proximally between the tibia and fibula) can be produced by forced external rotation in dorsiflexion. An X-ray may determine a widened ankle mortise. More severe cases leading to tibiofibular diastasis require an orthopaedic opinion.

TALAR DOME INJURY[12]

Talar dome lesions represent chip fractures of cartilage or bone from the articular surface of the talus. These occur in about 4-5% of ankle sprains and are usually detected by MRI and/or bone scan. It should be suspected with an inability to weight bear after an ankle sprain and persistent severe pain. Symptomatic lesions are usually treated best with arthroscopic surgery.

HEEL DISORDERS

Important causes of heel pain and other disorders resulting from overuse sporting activities include:

- Achilles tendon disorders
    - tendonitis/peritendonitis
    - tear: partial or complete
- bruised heel
- pump bumps'/bursitis
- calcaneal apophysitis
- plantar fasciitis (p. 752)

- talon noir
- blisters

Achilles tendonitis/peritendonitis[11]

The inflammation that occurs as a combination of degenerative and inflammatory changes due to overuse may appear either in the tendon itself or in the surrounding paratendon. The latter is called peritendonitis rather than tenosynovitis because there is no synovial sheath.

Clinical features:

- history of unaccustomed running or long walk
- common in runners who change routine
- usually young to middle-aged males
- aching pain on using tendon
- tendon feels stiff, especially on rising
- tender thickened tendon
- palpable crepitus on movement of tendon

Ultrasound examination

This is very useful in differentiating between tendonitis, peritendonitis, focal degeneration and a partial tear.

Preventive measures

- Warm-up and stretching exercises in athletes
- Good quality shoes
- 1 cm heel raise

Treatment

- Rest: ? crutches in acute phase, plaster cast if severe
- Cool with ice in acute stage, then heat NSAIDs
- 1-2 cm heel raise under the shoe
- Ultrasound and deep friction massage
- Mobilisation, then graduated stretching exercises

Note: Ensure adequate rest and early resolution because chronic tendonitis is persistent and very difficult to treat.

Avoid corticosteroid injection in acute stages and never give into tendon. Can be injected around the tendon if localised and very tender.

Partial rupture of Achilles tendon

Clinical features

- a sudden sharp pain at the time of injury
- sharp pain when stepping off affected leg
- usually males > 30 years sporadically engaged in sport
- history of running, jumping or hurrying up stairs
- a tender swelling palpable about 2.5 cm above the insertion
- may be a very tender defect about the size of the tip of little finger

Treatment

If palpable gap - early surgical exploration with repair.

If no gap, use conservative treatment:

- initial rest (with ice) and crutches
- 1-2 cm heel raise inside shoe
- ultrasound and deep friction massage
- graduated stretching exercises

Convalescence is usually 10-12 weeks. A poor response to healing manifests as recurrent pain and disability, indicates surgical exploration and possible repair.

Complete rupture of Achilles tendon

This common problem in athletes occurs in a possibly degenerated tendon subjected to a sudden increased load (e.g. a skier with foot anchored and ankle dorsiflexed).

Clinical features

- sudden onset of intense pain
- patient usually falls over
- feels more comfortable when acute phase passes
- development of swelling and bruising
- some difficulty walking, especially on tiptoe

Diagnosis

- Palpation of gap (best to test in first 2-3 hours as haematoma can fill gap)
- Positive Thompson's test (Figure 136.16 and 136.17)



Figure 136.16 Rupture of the right Achilles tendon. This 31-year-old woman injured her heel snow skiing. Examination by firmly compressing the gastroeneminus soleus complex of both legs shows an absent plantar reflex on the left side (positive Thompson's test). Photo courtesy *Bryan Walpole*



Figure 136.17 Thompson's calf squeeze test for ruptured Achilles tendon: (a) intact tendon, normal plantarflexion; (b) ruptured tendon, foot remains stationary

Note: The injury may be missed because the patient is able to plantarflex the foot actively by means of the deep long flexors to the foot.

Treatment

Early surgical repair (within 3 weeks).

'Pump bumps'

A 'pump bump' is a tender bursa over a bony prominence lateral to the attachment of the Achilles tendon. This is caused by inflammation related to poorly fitting footwear irritating a pre-existing enlargement of the calcaneus. Treatment is symptomatic and attention to footwear.

Talon noir

Talon noir or 'black heel', which has a black spotted appearance on the posterior end of the heel, is common in sportspeople, especially squash players. It tends to be bilateral and is caused by the shearing stresses of the sharp turns required in sport. The diagnosis is confirmed by gentle paring away of the hard skin containing old blood.

DISORDERS OF THE FEET AND TOES

Common problems include:

- fractures of toes
- foot strain
- ingrowing toenails
- 'black' nails
- bony outgrowth under the nail (subungual exostosis)
- calluses

- athlete's foot (tinea pedis)
- plantar warts

Black nails (‘soccer toe’)

Black or ‘bruised’ nails are due to subungual haematoma caused by trauma (Figure 136.18). The problem can be acute or chronic and is seen in the great toes. Acute cases are usually the result of the toe being trodden on, while chronic cases are the result of wearing ill-fitting shoes (too narrow or loose) or the toenails being left too long.



Figure 136.18 Black nails (‘soccer toe’), due to subungual haematoma caused by chronic trauma to the great toes. In this case a netball player wearing new ill-fitting shoes

The problem is encountered commonly in sports that involve deceleration forces and include running especially cross-country with downhill running), netball, basketball, tennis, football and skiing.

Treatment

An acute subungual haematoma should be decompressed with a hot needle or other procedure through the nail. A chronic non-painful problem should be left to heal. The toenails will become dystrophic and be replaced by ‘new’ nails.

Attention should be paid to the footwear either by changing it or by placing protective padding in the toes of the running shoes or boots.

INJURIES IN ADOLESCENTS

If an adolescent engaged in sport presents with pain in the leg it is important to consider the following problems.

- SCFE (p. 702)
- avulsion of epiphyses (e.g. ischial tuberosity - hamstring)
- stress fracture
- Osgood-Schlatter disorder
- Scheuermann's disorder
- idiopathic scoliosis

REFERENCES

1. Fitzpatrick J. Shoulder pain a real wet blanket. Aust Dr Weekly, 1993; 5 February: 56.
2. Robinson M. Hazards of alpine sport. Aust Fam Physician, 1991; 20: 961 - 70.
3. Elliott B, Sherry E. Common snow skiing injuries. Aust Fam Physician, 1984; 13: 570 - 4.
4. Brukner P, Khan K. *Clinical Sports Medicine* (2nd edn). Sydney: McGraw-Hill, 2000: 380 - 4.

5. Mashford L (Chair). Therapeutic Guidelines: Rheumatology (Version I) Melbourne: *Therapeutic Guidelines Ltd*, 2006: 168 - 9.
6. Zimmermann G. Groin pain in athletes. Aust Fam Physician, 1988; 17: 1046 - 52.
7. James T. Chronic lower leg pain in sport. Aust Fam Physician, 1988; 17: 1041 - 5.
8. Litt J. The sprained ankle. Aust Fam Physician, 1992; 21: 447 - 56.
9. Brukner P. The difficult ankle. Aust Fam Physician, 1991; 20: 919 - 30.
10. Stiell I. Ottawa ankle rules. Can Fam Physician, 1996; 42: 478 - 80.
11. Brukner P, Khan K. *Clinical Sports Medicine* (2nd edn). Sydney: McGraw-Hill, 2000: 540 - 2.
12. Paoloni J. Acute ankle injuries in sports. Medical observer, 2005; 20 May: 29 - 31.

CHAPTER 137. THE HEALTH OF INDIGENOUS PEOPLES

Health does not just mean the physical well being of the individual but refers to the social, emotional and cultural well being of the whole community. This is a whole of life view and includes the cyclical concept of life-death-life. Health services should strive to achieve the state where every individual can achieve their full potential as human beings (Aborigines) and thus bring about the total well being of their communities. This is an evolving definition (process).

NATIONAL ABORIGINAL COMMUNITY CONTROLLED HEALTH ORGANISATION, SEPTEMBER 1993

THE HEALTH OF INDIGENOUS PEOPLES

The major health challenge in Australia (and several other developed countries) is the health status of Indigenous peoples, which continues to be dramatically worse than that of other people. In some cases, it appears that the gap may be widening, especially for women.[1]

Life expectancies in 1997 for Aboriginal and Torpes Straits Islander men and women were 15 - 20 years below those of other Australians. At the end of 1997 their average life expectancy was 54.1 years for men and 61.2 years for women (compared with others - 75.7 years and 81.4 years, respectively).[2]

The commonest cause of death is cardiovascular disease especially ischaemic heart diseases which causes about 57% of these deaths.[3] The contrast with other Australians is most marked at 25 - 54 years. Diseases of the circulatory system, injury and poisoning, respiratory illness and neoplasms continue to be important causes of death. Deaths from infectious diseases and genitourinary disorders continue to occur at much higher rates than among other Australians.

The increasing incidence of diabetes is of great concern, especially in those changing from a traditional diet to inappropriate Westernised diets. Death rates from diabetes in 1992 - 94 are 12 times higher for men and nearly 17 times higher for women than the rates for other Australians.[1]

The estimated life span prior to white settlement was 40 years with the commonest cause of death being injury, particularly from warfare and murder. Thirteen percent of all children died within the first year of life and 25% by the end of the fifth year.[4] Records written by early settlers indicated that the Indigenous people appeared to be in good health and free from disease. It is estimated that the total Indigenous population was 750000 in 1788. It had fallen to about 70000 in the 1930s after 150 years of exposure to white civilisation. Significant causes were killings by the settlers (recorded as approximately 20000) and disease predominantly.

The main diseases that decimated the population were smallpox (two severe epidemics - 1789 and 1829-30), influenza, TB (very severe), pneumonia, measles, varicella, whooping cough, typhoid and diphtheria. The indigenous population is now approximately 480000.

The level of infant and maternal mortality continues to be a concern. After great reductions in infant mortality rates in the 1970s there has been a levelling off, with rates remaining 3 - 5 times higher than those of other Australians. About 30% of maternal deaths occur in Indigenous women, who contribute only about 3% of confinements.[1]

It is important to understand that Indigenous people and culture must not be seen as homogeneous but rather as diverse, with each group needing a special understanding of its cultural issues.

Practitioners working in primary health care in rural and remote areas in central and Northern Australia are advised to use the CARPA standard treatment Manual.[5]

KEY FACTS AND CHECKPOINTS

- Consider the importance of cultural issues in a consultation with an Indigenous person.
- If assistance is required with a cultural issue, consider involving an Aboriginal Indigenous health worker in the consultation. Such health workers are a vital part of the team.

- Always consider multiple medical conditions in the sick person.
- Remember the importance of opportunistic screening in Indigenous patients for relatively common conditions:
    - type 2 diabetes (20 - 50% incidence)
    - hypertension
    - other cardiovascular risk factors, eg. hyperchofesterdaemia
    - hepatitis B
    - STI urine screening (men and women)
    - Pap smears
    - anaemia in children
    - hearing in children
- Screening investigations to consider include:
    - blood sugar
    - serum lipids
    - urea and electrolytes
    - hepatitis B serology
    - BMI
    - urinalysis
- Cervical cancer is 6 - 8 times more common in Indigenous Australian women.
- Other common cancers are lung and liver.
- Approximately 50% of Indigenous children have chronic tympanic membrane perforations, with very significant consequences for language development and school achievement.
- The Indigenous Australian population has an incidence of end-stage kidney failure 10 times greater than that of other Australians.
- In some regions BCG vaccination is recommended for newborn Indigenous children.
- Influenza and pneumococcal vaccines are recommended for adults over 50 years.
- Depressive illness, like in the general community, is a significant concern.
- Alcohol use is a serious health and community problem. Another drinking problem is Kava, a drink made from a plant nature to the Pacific Islands. Its effects are similar to alcohol and benzodiazepines with marked muscle relaxation.[5] Excessive use causes acute and long-term problems.

NATIONAL SURVEY

The first national survey of the health of Indigenous Australians, completed in 1994, highlighted considerable differences in reported health status according to place of residence.[1] Interestingly, most survey participants (88%) considered themselves to be in good to excellent health.

The survey highlighted the following disorders:

- asthma
- ear and hearing problems
- diabetes
- hypertension
- kidney disorders
- heart disease
- skin disorders
- eye problems, including trachoma
- nutritional status (especially obesity)
- substance abuse (e.g. alcohol, marijuana, petrol sniffing)
- dental problems (a reversed trend of dental caries)
- pneumococcal respiratory disease

The many reasons for the lower health status of Indigenous Australians include poverty, dispossession, geographical isolation, high popu-lation mobility, unemployment, poor housing, low education attainment, temperature extremes in central Australia, increased exposure to infectious diseases, especially in subtropical areas, and lack of appropriate service deliveries.

Poor living conditions contribute to poor health outcomes, such as substance abuse, domestic violence, other social dysfunction and child malnutrition. Other environmental health issues, such as lack of adequate shelter, lack of basic amenities such as clean running water and adequate sewage disposal facilities, and often lack of refrigeration in hot climates all impact on Aboriginal health.

Associated co-morbidities of children admitted to the infectious diseases ward of the Royal Darwin Hospital[6] included dehydration (50%), malnutrition (60%), hypokalaemia (70%), iron deficiency (90%), anaemia (25%), pneumonia (24 - 32%), chronic suppurative otitis media (37%), urinary tract infection (10%) and scabies (25%).

Priority Health problems have been identified by the National Aboriginal Health Strategy and are summarised in Table 137.1.

Table 137.1 Priority Aboriginal health problems

A. Clinical
Diabetes
Cardiovascular disease
Injury (and youth suicide)
Kidney disease
STIs
Mental health
Poor nutrition
Ear infections
Women's problems
B. Socioeconomic
Education of Aboriginal children (particularly in rural and remote areas)
Housing
Water supply
Alcohol and substance misuse
Domestic violence and sexual abuse
Child abuse
Gambling
Unemployment

INDIGENOUS CULTURE AND THE DOCTOR - PATIENT RELATIONSHIP

An understanding of Indigenous cultural issues is fundamental to successful management outcomes.

Doctors should realise that, while examining Indigenous patients, they are themselves being examined. When Indigenous Australians visit the doctor they bring their own cultural expectations with them. ‘A poster or coffee table book on Indigenous culture in the waiting room may form a simple bridge to acceptance.’ [7] Indigenous people are often rather shy and communicate more indirectly than Europeans on important sensitive issues. They may use silence while waiting for answers and this could be a cue for the doctor to use a new line of approach.[8] It is important for doctors working in Indigenous communities to have an appreciation of and respect for Indigenous culture and be aware of its significance in health and behaviour.

An Indigenous patient may feel more relaxed if accompanied by a relative, who can witness what the doctor said and reinforce it later.

Women's business

The Indigenous concept of ‘women's business’; can be defined as the range of experience and knowledge that is the exclusive preserve and domain of Indigenous women. It encompasses issues about menstruation, pregnancy, childbirth and contraception.[9] Such matters are traditionally not discussed directly but are conveyed indirectly through stories, ceremonies and songs. For more traditional women it is taboo to talk about women's health issues to male doctors or male health workers, or to be physically examined by them. Apart from the sense of shame and embarrassment, it represents a transgression of their natural law.[10]

Men's business

Similarly, the cultural issue of men's business needs to be understood and respected. This applies to manhood initiation rites, cir-cumcision, sexuality and sexually transmitted infections.

Sorry business

Sorry business is the process of grieving, and this needs to be clearly understood. There is a cultural obligation for mourners to grieve the death of a relative in a special way. This involves changing their appearance and a deliberate avoidance of any mention of the deceased person's name or any portrayal of his or her likeness.[9] The place where the person died is deserted for a certain time and then smoked out.

COMMON PROBLEMS IN CHILDREN[10]

Indigenous children suffer the same spectrum of health problems as children in developing countries and communities throughout the world, and the infant mortality rate remains high. The major problems are malnutrition, diarrhoeal disease, skin infections and respiratory tract infection. Common problems are presented in Table 137.2.

Table 137.2 Common clinical problems in children

Perinatal
Low birthweight
Asphyxia
Infections
Preschool
Failure to thrive
Malnutrition
Anaemia
Respiratory infection
Diarrhoeal disease
Hepatitis B
Skin infection/infestation
Urinary tract infection
Meningitis

Joint and bone infection
Chronic suppurative otitis media
Later childhood and adolescence
Bacterial and viral infections
Parasitic infestation
Streptococcal infection
• rheumatic fever
glomerulonephritis
Trauma
Substance abuse
Chronic suppurative otitis media

Acute respiratory tract infections are a common reason for admission to hospital. Bacterial pneumonias occur more commonly than in non-Indigenous children and usually present late. Chronic upper respiratory tract disease is typical in younger children and mucopurulent nasal discharge is present in most preschool children.[11] Inappropriate treatment of respiratory tract infection will predispose to a high incidence of low-grade lower respiratory tract disease in later childhood and classic bronchiectasis.

Chronic suppurative otitis media, which is almost universal in preschool children, is often refractory to treatment and can lead to significant hearing impairment in many children. It can develop without apparent preceding acute otitis media and may be related to poor nutrition and anaemia. The basic treatment is ear toilet with povidone - iodine solution, followed by dry mopping with rolled toilet paper/tissue ‘spears’, or initial use of the ‘spears’ followed by acetic acid drops.

Skin infection and infestation are almost as prevalent as respiratory tract disease. Scabies is endemic and occasionally reaches epidemic proportions. It can be a very debilitating problem and can present in very young infants, in the first few weeks of life.[11]

Anaemia, usually iron deficiency, is found in at least 25% of children. Apart from reduced intake, intestinal loss from hookworm and other parasitic infection is an important factor. Treatment includes deworming in addition to iron supplements.

Diarrhoeal disease is a very common reason for hospital admission. Causes of infective gastroenteritis include rotavirus, bacteria in-cluding *Shigella, Salmonella and Campylobacter*, and parasites such as *Giardia lamblia, Strongyloides and Cryptosporidium*.

Other important and serious problems encountered more frequently in Indigenous children include bacterial meningitis (especially due to *Haemophilus influenzae*), septic arthritis and osteomyelitis, pyomyositis, *Streptococcus pyogenes* infections with associated glomerulonephritis and rheumatic fever, urinary tract calculi, urinary tract infection, especially 6-18 months of age) hepatitis B and petrol sniffing. (refer page 1350.)

The ability to achieve appropriate levels of immunisation in these communities will have enormous health benefits. Poliomyelitis, diphtheria, pertussis and tuberculosis are now rare and it is hoped that hepatitis B and H. *influenzae* infections will be drastically reduced.

SPECIFIC DISORDERS REQUIRING ATTENTION

The general practitioner attending Indigenous patients has to develop special skills in the diagnosis and management of the following health concerns:

- diabetes mellitus, frequently with associated hypertension and renal disease

- trauma
- substance abuse, including alcohol and smoking
- ear and eye infections
- respiratory disorders - URTIs and LRTIs, asthma
- skin disorders (e.g. fungal infections, impetigo, leg ulcers, cellulitis, boils)
- parasitic infestations (e.g. scabies, lice)
- gastrointestinal infections (e.g. *Campylobacter* enteritis, giardiasis, *Shigella*)
- sexually transmitted infections
- psychosocial dysfunction
- bites and stings
- severe infections (e.g. meningitis, rheumatic fever, septicaemia)
- hepatitis B
- tropical diseases (where applicable)
- worm infestation

However, the general management of medical disorders follows the principles and treatment guidelines outlined in this book. Antibiotic guidelines for use by Indigenous health workers in rural Indigenous communities are available.[12]

CARDIO VASCULAR DISEASE [3, 13]

Cardiovascular disease, especially ischaemic heart disease, is a major cause of continuing high rates of mortality and morbidity and includes ischaemic and rheumatic heart disease and stroke. Mortality from ischaemic heart disease is almost twice that in the non Indigenous population overall and 6-8 times higher in those aged 25-64 years. Reasons for thus include high smoking rates (twice the rate for others), type 2 diabetics (two to four times higher), obesity and low rates of plupical activity. Rheumatic heart disease is 11 times more common in the Indigenous population.

Target for secondary prevention of cardiovascular disease and diabetes mellitus are as presented in page 1321 (‘Management of diabetes’).

EAR INFECTIONS[5, 12]

Otitis externa and otitis media with its acute and chronic complications are major health problems in rural Indigenous children. Acute otitis media should be treated early and aggressively with antibiotics to prevent chronic suppurative otitis media, which is very difficult to cure once established. Check carefully for a perforation, which may affect management.

Treatment guidelines

- Acute otitis media
  Amoxycillin (o) or cotrimazole (o) or procaine penicillin (IM) for 5 days; if no response, consider amoxycillin/clavulanate or cefaclor
- Acute suppurative otitis media
  Antibiotics (as above) + dry mop ear
- Chronic suppurative otitis media
  Wash with povidone - iodine 5% solution using a 20 mL syringe with plastic tubing 1, 2 or 3 times daily, then dry mop with rolled toilet paper ‘spears’. Teach this method to family members. If available, suction kits are useful. Then instil ciprofloxacillin with hydrocortisone drops 12 hourly, especially in the presence of a perforation of the tympanic membrane.
- Otitis externa
  Gently clean out debris with toilet paper ‘spears‘ followed by acetic acid 0.25%; insert Kenacomb or Sofradex

drops or ointment 12 hourly on a gauze wick (if no perforation), otherwise ciprofloxacillin with hydrocortisone drops 12 hourly

- Acute mastoiditis
  Parenteral IM or IV flucloxacillin/dicloxacillin and hospitalisation

EYE INFECTIONS[5, 12]

Treatment guidelines

- *Periorbital cellulitis and penetrating eye trauma*. Arrange evacuation to hospital; if critically ill or delay in transfer give empirical treatment with ceftriaxone IM or IV once daily. Add gentamicin IM or IV as single dose for a child < 3 months or patients with other risk factors such as diabetes.
- *Conjunctivitis*. Take two swabs (one for microculture and one for *Chlamydia*). Apply topical chloramphenicol eyedrops plus ointment.
- *Neonatal gonococcal ophthalmia and chlamydia infection*. Refer to pp. 552 - 1.
- *Gonococcal conjunctivitis*. Procaine penicillin (IM) statim or single dose oral therapy with amoxycillin plus probenecid (e.g. 3 g + 1 g in adult). Use ceftriaxone IM if penicillin-resistant.
- *Trachoma*. These patients have ‘scratchy’ eyes and watery discharge ± red eye.
    - if over 6 kg and not pregnant: azithromycin (o) as single dose
    - if under 6 kg or pregnant: erythromycin or roxithromycin (o) for 14 days *or* oily tetracycline eyedrops 1 bd for 3 - 6 weeks
    - check and treat household contacts
    - check routinely for ‘follicles’ of trachoma

SKIN AND SOFT-TISSUE INFECTIONS

Skin infections are the commonest presenting problem in many clinics.[16] These include a high incidence of scabies and tinea corporis (ringworm), boils and carbuncles, infected wounds, impetigo and cellulitis. The most serious complication of skin infections is post-streptococcal glomerulonephritis secondary to *S. pyogenes* infection. Scabies is the most common skin infestation and commonly starts as an itchy rash with pinhead papules in the web spaces of the fingers.

Recommended treatment (in summary)[5, 12]

Impetigo and other skin sores

- Soak and remove crusts with povidone - iodine or cetrimide/chlorhexadine
- Antibiotic treatment (if required)
  Bicillin All-Purpose IM, statim dose *or*
  erythromycin (o) 12 hourly or roxithromycin (o) daily for 10 days

Cellulitis (mild - moderate) and erysipelas

Bicillin All-Purpose IM on days 1 and 3 or daily for 3 - 5 days *or*

procaine penicillin IM daily for 5 days

If no improvement: flucloxacillin/dicloxacillin plus probenecid (as below)

Boils, carbuncles, abscesses, bullous impetigo

Flucloxacillin/dicloxacillin (o) + probenecid (o) 12 hourly for 5 - 10 days

Suppurative wound infections

- Use local measures such as aseptic dressings and topical antiseptics
- If necessary, add flucloxacillin/dicloxacillin (as above); consider clindamycin

Tinea corporis (ringworm)

- Use benzoic acid ointment, Whitfield's ointment or one of the imidazole preparations: apply 1 - 3 times daily for 4 - 6 weeks
- Continue topical therapy for 2 weeks after resolution
- Systemic agents may be necessary

Scabies

- Apply permethrin 5% cream or benzyl benzoate 25% emulsion (pp. 1152 and 1134)
- For children less than 2 months use sulphur 5% cream for 2 - 3 days *or*
  crotamiton 10% cream daily for 3 - 5 days
- For infected scabies use flucloxacillin/dicloxacillin or erythromycin

Pityriasis versicolor (white spot)

Refer to p. 1175.

GABHS infections

Group A beta haemolytic streytococci (GABHS) infections are a significant problem, causing diseases such as phanynotonsillitis, impetigo, cellulitis, otitis media and scarlet fever. Two important immunological reactions to the streptococcus toxin are acute rheumatic fever and (post streptococcus) glomerulonephritis (PSGN).[3]

Acute rheumatic fever

Rheumatic fever and its effect are a serious cause of cardiovascular morbidity and mortality in Indigenous people, It is a classic hallmark of overcrowding, poverty and lack of hygiene, all of which facilitates *Streptococcal* Infection. The Australian Indigenous population has the highest role of acute rheumatic fever of any racial group in the world with an incidence of 250-300 per 100000 children[15] (refer to p. 250 - 300 for clinical features).

Treatment is with benzathine penicillin.

Acute glomerulonephritis

There is an association between PSGN (pp. 829 - 30) and *Streptococcal* skin and throat infections. Impetigo is the more frequent antecedent to PSGN. A study based on the fact that the Indigenous population has an incidence of end stage kidney failure 10 times greater than the general population, found that people with a history of PSGN in childhood had a risk of overt albuminuria more than six times that in a control group.[16]. There is no simple treatment of PSGN and preventing *streptococcal* infection remains the most important control strategy. Penicillin is beneficial in preventing spread during epidemics.

Tuberculosis

The rate of TB in Indigenous Australians is 10 - 15 times higher than other people in the Northern Territory.[3] It is related to poverty, overcrowding. Malnutrition and homelessness. Early diagnosis and treatment is the Ley to its control. This includes detection of latent TB infection in those at risk and strategies to prevent transmission in acute cases, BCG immunisation in higher risk groups is a key preventive strategy and is recommended for newborn from high-risk communities.

Loprosy (Hansen's disease)

Leprosy (p. 244) has been endemic in Northern Australia for over 100 years, with the Indigenous population being mainly affected, although the incidence is reducing. Control strategies, include early diagnosis of new cases, treatment with multi, drug therapy, monitoring of treatment to ensure its completion and prevention of nerve function impairment (NFI). NFI is monitored by a brief voluntary muscle and sensory test. It defected early it responds to anti-inflammatory medication such as prednisolone.

BCG vaccination, used to prevent TB, has probably had a protective efficacy against leprosy of approximately 50%.[3]

Worm infestations

Intestinal helminths are common in tropical northern Australia. Symptoms may include diarrhoea and abdominal pain with or without distension. Anaemia is common with hookworm infestation. Refer to Chapter 83 (pp. 971 - 2) for more information.

Treatment [5, 12]

- *Hookworm, roundworm, threadworm* - pyrantel embonate or imbendazole or albendazole
- *Whipworm* - mebendazole or albendazole
- *Strongyloidiasis* - albendazole or thiabendazole
- *Cutaneous larva migrans* - albendazole or thiabendazole

Community worm program. In selected communities a worm eradication program is recommended for children between the ages of 6 months and 12 years with either pyrantel embonate or albendazole.

Sexually transmitted infections

In the management of STIs it is important to be aware of the significance of men's and women's ‘business’; that is, the cultural sensitivities regarding men's and women's special gender feelings and issues, which need to be observed when discussing STIs with Indigenous patients. For some women it is inappropriate for a male doctor to discuss such issues with them but appropriate for a female doctor, nurse or health worker to do so. Similarly, it may be inappropriate for a female health worker to discuss STIs with a male patient.

Always undertake a full STI screen in patients presenting with an STI. Screening consists of counselling, taking blood for RPR, hepatitis B, hepatitis C and HIV, urethral and cervical swabs for gonococcus and *Chlamydia*, ulcer swabs, viral media swab for possible herpes, ‘snip’ biopsy (where appropriate) for Donovanosis or malignancy, and urine for MCU.[12] First voided urine for PCR for gonococcus and *Chlamydia* is an important investigation, especially for those who refuse swabs. Women can take their own tampon swabs for investigation. All sexually active females under 25 years should be screened opportunistically for *Chlamydia* (in particular) and griorhrea. Follow-up after therapy and contact tracing, screening, treatment and notification are all necessary.

Specific treatment

Refer also to Chapter 107.

Urethritis and cervicitis

amoxycillin 3 g (o) + probenecid 1 g (o) - single dose (use ceftriaxone in penicillin-resistant areas)

plus

azithromycin 1 g (o) as single dose

Genital sores

Herpes simplex, syphilis and Donovanosis are much more common than chancroid and lymphogranuloma venereum. Serology for syphilis is essential. Advise to avoid sex and for males to use condoms until treatment is completed and lesions well healed.

Syphilis

benzathine penicillin 1.8 g IM as single dose

Donovanosis (granuloma inguinale)

azithromycin 0.5 - 1 g (o) once daily for 7 days[11] *or*

1 g (o) once weekly for 4 weeks

(if not pregnant or breastfeeding); if so, give erythromycin or roxithromycin[12]

Herpes simplex

See pp. 1130 - 2.

Genital warts

See p. 1133.

Pelvic inflammatory disease

If sexually acquired, it is usually due to *Neisseria gonorrhoeae or Chlamydia trachomatis* (less likely). For treatment, see pp. 1001 - 3.

Vaginal infections

Refer to Chapter 110 for treatment. *Trichomonas vaginalis* is usually sexually transmitted, while *Candida albicans* and bacterial vaginosis is not.

COMMUNICATION TIPS[19]

- Don't assume English is a first language, particularly in remote areas.
- Don't assume a nod means understanding and/or agreement to treatment.
- Check hearing as chronic ear infections can impair it.
- Appreciate the different family network, particularly the tendency of grandmothers and aunts to care for children.
- Don't assume a broken appointment means the patient will not return for treatment. Often family and cultural duties take precedence.
- Be aware of cultural sensitivity.
- Don't touch a patient, particularly of the opposite sex, without seeking permission and explaining what you are doing.
- Be aware that patients may not be comfortable with direct questions about their family and health.
- Don't be too stern or authoritative during a consultation.
- Ensure receptionists and other staff understand the cultural sensitivities of Indigenous patients.
- Be accepting, respectful and non-judgmental.

Practice tips

- Anaemia is common in Indigenous children - be on the lookout for it. Consider giving pyrantel embonate or mebendazole in a hookworm endemic area.
- Asthma is common - consider it in coughing children.
- In children with failure to thrive consider insufficient food, urinary tract infection, GIT infection or parasites and recurrent illness.
- Beware of diarrhoea in children - attend to fluid and electrolyte replacement.
- Think pelvic inflammatory disease in the woman of child-bearing age presenting with abdominal pain. Be watchful for penicillinase-producing *Neisseria gonorrhoeae* (although it remains uncommon in Indigenous communities).
- Consider the possibility of rheumatic fever or glomerulonephritis with S. pyogenes throat infection and treat

with an optimal course of antibiotics (e.g. single injection of benzathine penicillin).

- In tropical areas consider diseases such as melioidosis, dengue and Ross River infection.
- Promote immunisation programs.
- Measles can be a very serious disease in Indigenous children and is highly contagious.
- In the fitting or aggressive patient, alcohol withdrawal is the commonest cause but the possibility of petrol sniffing should also be considered.
- Kidney failure is more common in Indigenous people: look for it if proteinuria, diabetes, hypertension, general debility or recurrent infections.
- Serum creatinine measures are also useful. A level of < 150 μmol/L is regarded as a safe limit for renal function.
- Consider urine albumin/creatinine ratio testing in adults to detect and monitor early renal disease. As renal disease has a rapidly progressive nature regular monitoring is important.[17] Renal function must be monitored at least annually.
- Donovanosis (granuloma inguinale), which can be chronic and progressive, presents initially as raised beefy nodules or sores. Microscopy of scrapings or snip or pinch biopsy will confirm diagnosis.[17]
- If routine swabs for gonococcus and *Chlamydia* infection are unobtainable for cultural reasons, use urine PCR testing.
- Medroxyprogesterone acetate (Depo-Provera) and Implanon are very useful contraceptive agents but always adhere to guidelines for informed consent.
- Because compliance with medication may be a problem with some patients, once-a-day therapy is recommended where possible.
- Point of Care laboratory equipment is very helpful for monitoring diabetic patients as they can be used in remote communities by trained health workers.

REFERENCES

1. Reid J (Chair). *Australia's Health 1996*. Canberra: Australian Institute of Health & Welfare, 1996, 21 - 30.
2. *General Practice in Australia: 2000*. Canberra: Department of Health & Aged Care, 2000: 90 - 2.
3. Couzos 5, Murray R, Aboriginal Primary Health Care. An evidence based Approach (2nd edn) Melbourne: Oxford University Press 2003.
4. Reid J, Trompf P. *The Health of Aboriginal Australia*, Sydney; Itarcourt Brace Jovanavidi, 1991.
5. Central Australian Rural Practitioners Association CARPA Standard Treatment Manual (4th edn.). Alice Springs; ARPA 2003. (Available from CARPA, PO Box 8143, Alice Springs, NT 0871.)
6. Ruben AR, Walker A. Malnutrition among rural Aboriginal children in the Top End of the Northern Territory. Med J Aust, 1995; 162: 400 - 3.
7. Hill P. Aboriginal culture and the doctor - patient relationship. Aust Fam Physician, 1994; 23: 29 - 32.
8. Eades D. They don't speak an Aboriginal language, or do they? In: Keen I ed. *Being Black: Aboriginal Cultures in 'Settled' Australia*. Canberra: Aboriginal Studies Press, 1991: 97 - 115.
9. O'Connor M. Women's business. Aust Fam Physician, 1994; 23: 40 - 4.
10. National Aboriginal Health Strategy Working Party. *A National Aboriginal Health Strategy*. Canberra: Department of Aboriginal Affairs, 1989: 193.
11. Walker A. Common health problems in Northern Territory Aboriginal children. Aust Fam Physician, 1994; 23: 55 - 62.
12. Mashford ML (Chair). Antibiotic Guidelines for Central and Northern Australia and other remote areas. In: *Antibiotic Guidelines* (9th edn). Melbourne: Therapeutic Guidelines Ltd, 1997: 175 - 202.
13. Walsh WF. Editorial: Cardiovascular health in Indigenous Australians: a call for action. MJA, 2001; 175: 351-2.
14. Mashford L (Chair) *Therapeutic Guidelines: Rheumatology* (Version 1) Melbourne. Therapeutic Guidelines Ltd, 2006: 201 - 7.
15. Atkins RC, Editorial: How bright is their future? MSA, 2001; 174:489-90.
16. Crowe C. Common illnesses (Aboriginal health). Aust Fam Physician, 1995; 24: 1469 - 74.

17. Bell D. Chronic disease in Indigenous Australians. Aust Doctor, 2002; 14 June: I - VIII.
18. Ryan K. Skill with indigenous patients: cultural issues. Aust Doctor, 2001; 9 March: 66 - 7.

APPENDIX I. PERCENTILE CHARTS: INFANT GIRLS



INFANT GIRLS

NAME BIRTH DATE NO.

WEIGHT
The child is weighed without clothing except light undergarments

LENGTH
The child lies relaxed on a firm surface. The soles of the feet are held against a firm upright at the zero mark on a parallel rule and a movable square is brought against the vertex. The head is held so the eyes face the ceiling.

HEAD CIRCUMFERENCE

APPENDIX II. PERCENTILE CHARTS: INFANT BOYS



INFANT BOYS

NAME

BIRTH DATE

NO.

WEIGHT

The child is weighed without clothing except light undergarments

LENGTH

The child lies relaxed on a firm surface. The soles of the feet are held against a firm upright at the zero mark on a parallel rule and a movable square is brought against the vertex. The head is held so the eyes face the ceiling.

APPENDIX III. PERCENTILE CHARTS: GIRLS



LENGTH
The child lies relaxed on a firm surface. The soles of the feet are held against a firm upright at the zero mark on a parallel rule and a movable square is brought against the vertex. The head is held so the eyes face the ceiling.

HEIGHT
The child's heels should be together. Heels buttocks and occiput should be against a vertical upright.

WEIGHT
The child is weighed without clothing except light undergarments.

APPENDIX IV. PERCENTILE CHARTS: BOYS



LENGTH
The child lies relaxed on a firm surface. The soles of the feet are held against a firm upright at the zero mark on a parallel rule and a movable square is brought against the vertex. The head is held so the eyes face the ceiling.

HEIGHT
The child's heels should be together. Heels buttocks and occiput should be against a vertical upright.

WEIGHT
The child is weighed without clothing except light undergarments.

APPENDIX V. RESPIRATORY FUNCTION TABLES

Predicted mean values for healthy Australian adults

The mean predicted normal values for Caucasian males and females between 10 and 80 years of age are given in the following tables.

FEV$_1$ (Litres)												**Male**
Height		**145**	**150**	**155**	**160**	**165**	**170**	**175**	**180**	**185**	**190**	**195**
Age	10	2.31	2.54	2.77	3.00	3.23	3.46	3.69	3.92	4.15	4.38	4.61
	12	2.40	2.63	2.86	3.09	3.32	3.55	3.78	4.01	4.24	4.47	4.70
	14	2.49	2.72	2.95	3.18	3.41	3.64	3.87	4.10	4.33	4.56	4.79
	16	2.58	2.81	3.04	3.27	3.50	3.73	3.96	4.19	4.42	4.65	4.88
	18	2.67	2.90	3.13	3.36	3.59	3.82	4.05	4.28	4.51	4.74	4.97
	20	2.67	2.99	3.22	3.45	3.68	3.91	4.14	4.37	4.60	4.83	5.06
	25	2.66	2.92	3.18	3.44	3.70	3.96	4.22	4.48	4.47	5.00	5.26
	30	2.53	2.79	3.05	3.31	3.57	3.83	4.09	4.35	4.61	4.87	5.13
	40	2.26	2.52	2.78	3.04	3.30	3.56	3.82	4.08	4.34	4.60	4.86
	50	1.99	2.25	2.51	2.77	3.03	3.29	3.55	3.81	4.07	4.33	4.59
	60	1.72	1.98	2.24	2.50	2.76	3.02	3.28	3.54	3.80	4.06	4.32
	70	1.45	1.71	1.97	2.23	2.49	2.75	3.01	3.27	3.53	3.79	4.05
	80	1.18	1.44	1.70	1.96	2.22	2.48	2.74	3.00	3.26	3.52	3.78

FVC (Litres)												**Male**
Height		**145**	**150**	**155**	**160**	**165**	**170**	**175**	**180**	**185**	**190**	**195**
Age	10	2.52	2.77	3.02	3.27	3.52	3.77	4.02	4.27	4.52	4.77	5.02
	12	2.68	2.93	3.18	3.43	3.68	3.93	4.18	4.43	4.68	4.93	5.18
	14	2.83	3.08	3.33	3.58	3.83	4.08	4.33	4.58	4.83	5.08	5.33
	16	2.99	3.24	3.49	3.74	3.99	4.24	4.49	4.74	4.99	5.24	5.49
	18	3.15	3.40	3.65	3.90	4.15	4.40	4.65	4.90	5.15	5.40	5.65
	20	3.30	3.55	3.80	4.05	4.30	4.55	4.80	5.05	5.30	5.55	5.80
	25	3.24	3.57	3.89	4.22	4.54	4.87	5.19	5.52	5.84	6.17	6.49
	30	3.10	3.42	3.75	4.07	4.40	4.72	5.05	5.37	5.70	6.02	6.35
	40	3.81	3.13	3.46	3.78	4.11	4.43	4.76	5.08	5.41	5.73	6.06
	50	2.52	2.84	3.17	3.49	3.82	4.14	4.47	4.79	5.12	5.44	5.77
	60	2.23	2.55	2.88	3.20	3.53	3.85	4.18	4.50	4.83	5.15	5.48
	70	1.94	2.26	2.59	2.91	3.24	3.56	3.89	4.21	4.54	4.86	5.19
	80	1.65	1.97	2.30	2.62	2.95	3.27	3.60	3.92	4.25	4.57	4.90

These tables are based on: Knudson RJ, Slatin RC, Lewowitz MD and Burrows B. The maximal expiratory flow volume curve. Normal standards variability, and effect of age. *American Review of Respiratory Disease* 1976, 113:587-600, as used in *Spirometry: The Measurement and Interpretation of Ventilatory Function* , Dr Rob Pierce and Mr David P Johns. National Asthma Campaign, Melbourne, 1975. Reprinted with permission

FEV$_l$ (Litres)												**Female**
Height		**145**	**150**	**155**	**160**	**165**	**170**	**175**	**180**	**185**	**190**	**195**
Age	10	2.06	2.20	2.33	2.47	2.60	2.47	2.87	3.01	3.14	3.28	3.41
	12	2.23	2.37	2.50	2.64	2.77	2.91	3.04	3.18	3.31	3.45	3.58
	14	2.40	2.54	2.67	2.81	2.94	3.08	3.21	3.35	3.48	3.62	3.75
	16	2.57	2.71	2.84	2.98	3.11	3.25	3.38	3.52	3.65	3.79	3.92
	18	2.74	2.88	3.01	3.15	3.28	3.42	3.55	3.69	3.82	3.96	4.09
	20	2.70	2.84	2.97	3.11	3.24	3.38	3.51	3.65	3.78	3.92	4.05
	25	2.60	2.73	2.87	3.00	3.14	3.27	3.41	3.54	3.68	3.81	3.95
	30	2.49	2.63	2.76	2.90	3.03	3.17	3.30	3.44	3.57	3.71	3.84
	40	2.28	2.42	2.55	2.69	2.82	2.96	3.09	3.23	3.36	3.50	3.63

	50	1.07	2.21	2.34	2.48	2.61	2.75	2.88	3.02	3.15	3.29	3.42
	60	1.86	2.00	2.13	2.27	2.40	2.54	2.67	2.81	2.94	3.08	3.21
	70	1.65	1.79	1.92	2.06	2.19	2.33	2.46	2.60	2.73	2.87	3.00
	80	1.44	1.58	1.71	1.85	1.98	2.12	2.25	2.39	2.52	2.66	2.79

FVC (Litres)												**Female**
Height		**145**	**150**	**155**	**160**	**165**	**170**	**175**	**180**	**185**	**190**	**195**
Age	10	2.24	2.40	2.57	2.73	2.90	3.06	3.23	3.39	3.56	3.72	3.89
	12	2.42	2.59	2.75	2.92	3.08	3.25	3.41	3.58	3.74	3.91	4.07
	14	2.60	2.77	2.93	3.10	3.26	3.43	3.59	3.76	3.92	4.09	4.25
	16	2.79	2.95	3.12	3.28	3.45	3.61	3.78	3.94	4.11	4.27	4.44
	18	2.97	3.14	3.30	3.47	3.63	3.80	3.96	4.13	4.29	4.46	4.62
	20	3.15	3.34	3.52	3.71	3.89	4.08	4.26	4.45	4.63	4.82	5.00
	25	3.04	3.23	3.41	3.60	3.78	3.97	4.15	4.34	4.52	4.71	4.89
	30	2.93	3.12	3.30	3.49	3.67	3.86	4.04	4.23	4.41	4.60	4.78
	40	2.71	2.90	3.08	3.27	3.45	3.64	3.82	4.01	4.19	4.38	4.56
	50	2.49	2.68	2.86	3.05	3.23	3.42	3.60	3.79	3.97	4.16	4.34
	60	2.27	2.46	2.64	2.83	3.01	3.20	3.38	3.57	3.75	3.94	4.12
	70	2.05	2.24	2.42	2.61	2.79	2.98	3.16	3.35	3.53	3.72	3.90
	80	1.83	2.02	2.20	2.39	2.57	2.76	2.94	3.13	3.31	3.50	3.68

APPENDIX VI. THE AUSTRALIAN NUTRITION FOUNDATION: WEIGHT FOR HEIGHT CHARTS